TITLE:

Inferring the role of habitat dynamics in driving diversification: evidence for a species pump

in Lake Tanganyika cichlids

AUTHORS: Thijs Janzen[1,2,3*], Rampal S. Etienne[2]

AUTHOR AFFILIATION

1. Max Planck Institute for Evolutionary Biology, Department of Evolutionary Theory,
August-Thienemann-Straße 2, 24306, Plön, Germany.
2. University of Groningen, Groningen Institute for Evolutionary Life Sciences, Box
11103, 9700 CC Groningen, The Netherlands
3. Carl von Ossietzky University of Oldenburg, Institute for Biology, Department of
Molecular Ecology, Carl-von-Ossietzky Straße 9, 26111, Oldenburg, Germany

4. *Corresponding author: Thijs Janzen, Carl von Ossietzky University of Oldenburg,
Institute for Biology, Department of Molecular Ecology, Carl-von-Ossietzky Straße 9,
26111, Oldenburg, Germany
e-mail: thijs.janzen@uni-oldenburg.de

ABSTRACT

Geographic isolation that drives speciation is often assumed to slowly increase over time, for

instance through the formation of rivers, the formation of mountains or the movement of

tectonic plates. Cyclic changes in connectivity between areas may occur with the

advancement and retraction of glaciers, with water level fluctuations in seas between islands

or in lakes that have an uneven bathymetry. These habitat dynamics may act as a driver of

allopatric speciation and propel local diversity. Here we present a parsimonious model of the

interaction between cyclical (but not necessarily periodic) changes in the environment and

speciation, and provide an ABC-SMC method to infer the rates of allopatric and sympatric

speciation from a phylogenetic tree. We apply our approach to the posterior sample of an

updated phylogeny of the *Lamprologini*, a tribe of cichlid fish from Lake Tanganyika where

such cyclic changes in water level have occurred. We find that water level changes play a

crucial role in driving diversity in Lake Tanganyika. We note that if we apply our analysis to

the Most Credible Consensus (MCC) tree, we do not find evidence for water level changes

influencing diversity in the *Lamprologini*, suggesting that the MCC tree is a misleading

representation of the true species tree. Furthermore, we note that the signature of habitat

dynamics is found in the posterior sample despite the fact that this sample was constructed

using a species tree prior that ignores habitat dynamics. However, in other cases this species

tree prior might erase this signature. Hence we argue that in order to improve inference of the

effect of habitat dynamics on biodiversity, phylogenetic reconstruction methods should

include tree priors that explicitly take into account such dynamics.

INTRODUCTION

Environmental changes such as the formation of mountain ridges, the formation of rivers and

the movement of tectonic plates have long been known to be important drivers of speciation

(Coyne and Orr 2004). Repeated environmental changes may thus lead to diversification

patterns. Cyclic changes in the environment can cause populations to continuously switch

between an allopatric and sympatric stage, providing a continuously renewed potential for

speciation. And these cyclic changes can in turn drive diversity towards levels unexpected

given the current geography, sometimes referred to as a "species pump" (Heaney 1985;

Rossiter 1995). Examples of species pumps include environmental fluctuations fragmenting

habitats on the slopes of mountains (Weir 2006; Sedano and Burns 2010; Hutter et al. 2013),

glaciations and postglacial secondary contacts (Barnosky 2005), sea level changes causing

the fusion and fragmentation of islands (Glor et al. 2004; Thorpe et al. 2008, but see

Papadopoulou and Knowles 2015), and fluctuations in water level causing fragmentation and

fusion of lakes with uneven bathymetry, as in the African Rift Lakes (Cohen et al. 1997b;

Alin et al. 1999; McGlue et al. 2008; Ivory et al. 2016).

The African Rift Lakes provide a good starting point in studying the interplay between cyclic

habitat dynamics and speciation, because they have been subject to frequent water level

changes (Cohen et al. 1997b; Alin and Cohen 2003; Ivory et al. 2016), and are well known

for their tremendous biodiversity (Seehausen 2000, 2006; Turner et al. 2001; Wagner et al.

2012, 2014; Brawand et al. 2014). An estimated number of 2000 cichlid fish species (Turner

et al. 2001) have evolved in the African Rift Lakes over the past 10 million years (Genner et

al. 2007; Meyer et al. 2016), and comprise one of the most spectacular adaptive radiations

(Seehausen 2006). The most prominent water level changes took place in Lake Tanganyika,

where the water level has dropped substantially on multiple occasions over the past million

years, sometimes splitting the lake into multiple smaller lakes (Lezzar et al. 1996; Cohen et

al. 1997a, 2007). Being the oldest lake of the three large rift lakes (Cohen et al. 1993), Lake

Tanganyika contains the highest behavioral diversity (Konings 2007) and is the only lake

with a highly resolved phylogeny for cichlid fish. Evidence for the influence of changing

water levels comes from analysis of mitochondrial DNA, which shows that for *Tropheus*

species, some populations have experienced secondary contact upon changes in water level,

potentially increasing genetic diversity and driving speciation (Sturmbauer et al. 2001;

Koblmüller et al. 2011; Sefc et al. 2017). Similar patterns were found for *Variabilichromis*

moorii and *Ophthalmotilapia nasuta* (Sturmbauer et al. 2001), *Telmatochromis temporalis*

(Winkelmann et al. 2016), and *Altolamprologus* (Koblmüller et al. 2016). Comparison of

mitochondrial DNA between populations from deep and shallow areas emphasizes that the

deep areas are habitats that are more persistent over time, with lower genetic variation

(Nevado et al. 2013). Furthermore, *Eretmodus* lineages identified using mitochondrial DNA

are strongly associated with the bathymetric basins of Lake Tanganyika (Verheyen et al.

1996), suggesting that they have independently diversified at low water level.

Aguilée et al. (2013) developed a model for the African Rift Lakes in which populations at

different locations diverge from each other depending on the local habitat, and at the same

time allowed for sympatric speciation by implementing assortative mating that allows for a

single branching point in trait values. Over time the different locations become separated or

are reconnected, and this may drive the formation of new species. The authors conclude that

stable numbers of diversity are best obtained by a fragmented habitat with recurrent merged

states and rapid fluctuations. However, Aguilée et al. (2013) do not compare their results to

empirical data. By contrast, Pigot et al (2010) used a spatially explicit model of landscape

fragmentation, where consecutive splitting of species' geographic ranges drives speciation,

and compared phylogenies generated with their model, with known bird phylogenies. They

found that including this geographical context of speciation explains a large part of the

features exhibited by the reconstructed avian trees. Hence, including a geographical context

of speciation seems a promising research avenue.

Here, we provide a method to infer whether or how cyclic changes in the environment

influence both the generation and the maintenance of biodiversity. We use an extension of the

standard constant-rates birth-death model. Because deriving an expression for the likelihood

of this model for a given set of phylogenetic branching times is difficult, but simulation of

phylogenies under the model is easy, we used approximate Bayesian computation (ABC)

based on sequential Monte Carlo sampling (SMC) to estimate parameters from phylogenies.

We applied our approach to an updated phylogeny of the *Lamprologini*, a tribe of cichlid fish

from Lake Tanganyika in order to assess the importance of these habitat dynamics in shaping

the current biodiversity of cichlids in Lake Tanganyika.

METHODS

Model

To model the interaction between environmental change and speciation, we envisage a lake that consists of a single pocket at high water level, but that splits into two pockets when the water level drops. When the water level drops, we assume that all species distribute themselves equally over the two pockets; similarly, when the water level rises, all species previously contained in the two pockets are combined into the single pocket. Allopatric speciation can only occur when the water level is low. We assume a constant probability rate for allopatric speciation, and hence the waiting time until the next speciation event is exponentially distributed. After this waiting time, one of the two incipient species in either pocket can speciate into a new species. If this allopatric speciation does not occur before the water level rises again, i.e. reflecting that there has not been enough genetic divergence, the two incipient species in the two pockets merge back into one species. This is conceptually similar to the idea of protracted speciation (Etienne and Rosindell 2012): the water level drop initiates the speciation process whereas the allopatric speciation event is the completion of speciation under the protracted speciation model. Sympatric speciation can always occur in our model, either at high water level in the lake, or in both pockets when the water level is low. Extinction is considered to be a background process that occurs locally, i.e. within a pocket. If the water level is high, this causes extinction of a species, if the water level is low, this causes local extinction in one of the pockets.

We implemented our model using a Gillespie algorithm, where the time steps are chosen depending on the rate of possible events. In the model there are five possible events:

    1) A water level change event, inducing incipient species or merger of incipient species.

    2) Sympatric speciation event at high water level, with rate λ_s^h

3) Sympatric speciation event at low water level, with rate λ_s^l

4) Allopatric speciation(-completion) event, at low water level, with rate λ_a^l

5) Extinction event, with rate μ

When the water level drops, all species distribute themselves over both pockets. Thus,

immediately after a water level drop, the number of incipient species is equal to twice the

number of species. When the water level rises, all incipient species that belong to the same

species merge into a single species. During a sympatric speciation event, a single species

splits into two new species, and the original (incipient) species is consumed in the process.

Here we assume that local disruptive selection causes divergence, similar to the

implementation of speciation by Aguilee (Aguilée et al. 2011, 2013). If sympatric speciation

occurs when the water level is low, the species in the other pocket is retained, and thus three

new lineages arise: the first branching point occurs at the water level drop while the second

occurs at the sympatric speciation event (Figure 1).

Figure 1. Schematic representation of the consequences of the three different types of
speciation. Time proceeds from left to right. The dotted blue lines indicate water level
changes. (a) During a sympatric speciation at high water level event, diversification is not
aligned with any associated water level change. (b) During allopatric speciation at low water
level, speciation initiation (incipient species are indicated with a dotted line) coincides with
the water level drop, causing branching events (if speciation-completion occurs before water
level rise) to line up in time. Branching events are conditionally independent of the time of
speciation completion, hence, even when the actual speciation completion events occur at
different time points, branching events in the species tree are identical. (c) During a sympatric

speciation event at low water, the speciation event is independent of the water level changes.
Because the original species is consumed in the process, a new branching event is also added
at the water level change event. Hence, both speciation-completion (b) and sympatric
speciation at low water level (c) cause branching times to line up at the time of water level
drop. Please note that (a) and (c) represent the reconstructed species tree, but (b) does not; the
reconstructed species tree for (b) would not show the branching event in the uppermost part
of the tree.

During an extinction event, one (possibly incipient) species is removed from the simulation.

If the water level is low, this need not lead to the extinction of a species, because the sister

incipient species might remain in the other pocket, ensuring survival of the species.

Maximum Likelihood

Without water level changes, our model reduces to the constant rates birth-death model (Nee

et al. 1994). As a reference therefore, we estimated parameters of the standard birth-death

model using Maximum Likelihood. The likelihood of the birth-death model was calculated

using the function "bd_ML" from the R package DDD. (Etienne et al. 2012).

Fitting the model to empirical data

We performed two different fitting procedures: firstly, we performed a model selection

procedure, where three different water level scenarios were fitted simultaneously to the data

(more information about the chosen scenarios can be found in the next section). The model

selection procedure simultaneously estimates parameter estimates and assesses the fit of the

models. However, because the model selection procedure primarily samples the best fitting

model (by design), it does not allow for the comparison of parameter estimates across

different models. Therefore, we also fitted the three different water level scenarios

independently to the empirical data, and obtained posterior distributions for the parameters

relevant to these scenarios.

We fitted our model to 100 trees randomly sampled from the MCMC chain obtained from the

*BEAST analysis (see below), and to the Most Credible Consensus (MCC) tree.

Water level scenarios

The main focus of our approach is to assess the impact of water level changes on the

diversification rate. Lake Tanganyika experienced low water level stands 35 - 40 k years ago

(kya) (-160 meter), 169 - 193 kya (-250 meter), 262 - 295 kya (-350 meter) , 363 - 393 kya (-

350 meter) and 550 - 1100 kya (-650 – 700 meter) (Lezzar et al. 1996; Cohen et al. 1997a).

The southern and northern basin of Lake Tanganyika are separated from each other by a ridge

at a depth of 500 meter below present level. Although some of these water level changes may

not have split up the lake completely, we assume here that these water level changes still

caused sufficient disruption of migration between the northern and southern basin, to be

equivalent to physical separation. Consequently, high water levels occurred between 0 – 35

kya, 40 – 169 kya, 193 – 262 kya, 295 - 363 kya and 393 – 550 kya. Unfortunately the

geological record does not reveal whether any low water level stands occurred beyond 1.1

million years ago (Ma). This leaves us with two alternative scenarios: either no low water

level stands occurred beyond 1.1 Ma, or these low water level stands have not been preserved

accurately in the geological record.

In order to capture these two scenarios we performed inference using two alternative water

level implementations. Firstly we used the exact literature values, assuming a high water

level stand until 1.1 Ma. We refer to this scenario as LW (Literature Waterlevels). Secondly

we assumed that before 1.1. Ma, water level changes occurred at the same *average* rate of

water level change in the most recent 1.1 million years. In the recent 1.1 million years, the

lake experienced 5 high water level stands, and 5 low water level stands, which amounts to

10 water level changes in total. To extrapolate water level changes to more than 1.1 Ma, we

drew waiting times until the next water level change from an exponential distribution with

rate 10. We refer to this scenario as EW (Extrapolated Water levels). Thirdly we also tested

the null expectation: no effect of water level changes on speciation, we refer to this scenario

as NW (No Water levels). Without water level changes, the model reduces to the constant-

rates birth death model.

Parameter estimation

To fit the model to empirical data we used Approximate Bayesian Computation, in

combination with a Sequential Monte Carlo scheme (ABC-SMC) (Toni et al. 2009).

As summary statistics for the ABC analysis we chose the normalized Lineages Through Time

statistic (Janzen et al. 2015), tree size, Phylogenetic Diversity (AvPD, Schweiger et al. 2008)

and the γ statistic (Pybus and Harvey 2000). On all parameters $(\lambda_s^h, \lambda_s^l, \lambda_a^l, \mu)$ we chose

uniform priors U(-3, 2), on a ^{10}log scale, such that the eventual prior distribution spans $(10^{-3},$

$10^2)$. A ^{10}log scale was chosen to explore parameter space uniformly, and put extra emphasis

on low values. The standard deviation of the normal distribution used to perturb the

parameters was chosen to have a mean of 0, and a standard deviation of 0.05 (on the ^{10}log

transformed parameter), and we updated one parameter each time (e.g. jumps were only made

in one dimension, to avoid extremely low acceptance rates). The number of particles used per

SMC step was 10,000, where a particle is a data structure containing the model choice and

the parameter estimates. To assess the fit of the model to the data we calculated the Euclidian

distance between the summary statistic of the simulated data and the empirical data. To

ensure that the differences in summary statistics were on the same scale, we normalized the

differences. Differences were normalized by dividing each difference by the standard

deviation of that summary statistic of 1,000,000 trees simulated using parameter values

sampled from the prior.

 Model selection

To identify which model best explains the data, we performed ABC model selection, as

described in Toni et al. (2009; 2010). The main difference between standard ABC-SMC and

ABC-SMC including model selection is that the latter adds one parameter, which keeps track

of the model. As jumping kernel between models we assumed a 50% probability of staying at

the same model, and a 25% probability of jumping to either other model. We assumed a

uniform prior across all three models; this translates to a probability of 1/3 for each model in

the first iteration of the ABC-SMC procedure, and hence an expected number of 3333

particles assigned to each model in the first iteration. This reversible jump ABC-SMC model

selection procedure results in a posterior distribution over the three models, where the model

with most support is the model selected most across all particles. We can calculate the Bayes

factor by taking the ratio of the number of particles assigned to the respective models (Toni et

al. 2009). For example, the Bayes factor of LW/EW is the number of particles assigned to the

model with literature water level changes divided by the number of particles assigned to the

model with extrapolated water level changes. Because a model can receive zero particles, we

set the Bayes factor for each model compared to the model with zero particles to the

maximum support possible, which is the total number of particles: 10,000. To calculate the

posterior support for a model, we calculate 2 ln (Bayes factor), following Kass and Raftery

(1995). A transformed Bayes factor over a value of 2 then corresponds to substantial support

for the considered model (Kass and Raftery 1995).

Model selection validation

To assess whether our ABC-SMC method can accurately infer the correct model, we

simulated 100 datasets for each model (NW, LW & EW), with parameter values drawn from

the prior. We report the median Bayes factor across the 100 replicates. If our method can

accurately infer the correct model, we expect the median Bayes Factor (after 2 ln

transformation) to be above 2 when comparing posterior support for the model with which

the data was simulated to the other two models.

Measurement uncertainty

A phylogeny generated with a high rate of allopatric speciation and a high rate of water level

changes tends to have multiple speciation events that are aligned in time (Figure 1, b). This is

due to the fact that the onset of speciation is given by the time of water level change.

Phylogenetic reconstruction methods such as BEAST (Bouckaert et al. 2014) currently do not

allow for simultaneous branching events. Hence, when fitting the model, trees are generated

that are by definition dissimilar from the empirical tree constructed using BEAST, even if

underlying events are close to the original events. To circumvent this we perturbed the

branching time of each node in the trees simulated using our model. In this way speciation

events that were previously aligned in time now occur on slightly different time points, as in

a tree from a BEAST analysis. We perturbed branching times by adding a random number

drawn from a truncated normal distribution with mean 0, standard deviation σ, truncated by

the minimum distance to either the daughter or the parent species. If there were no daughter

lineages present, and the node gave rise to an extant species, the normal distribution was

truncated to the minimum distance to the parent or the present time. Nodes were perturbed

from past to present (leaving the crown in place, to ensure a phylogenetic tree with the same

age as the empirical tree). The standard deviation of the perturbation kernel was included as

an extra parameter to be inferred, with a uniform prior on $(10^{-3}, 10^0)$.

 Empirical data

We fitted our model to the phylogenetic tree of the tribe of *Lamprologini*, the most diverse

tribe within Lake Tanganyika, containing 79 species of cichlids in Lake Tanganyika (Day et

al. 2007; Koblmüller et al. 2007; Sturmbauer et al. 2010). The *Lamprologini* are endemic to

Lake Tanganyika and its surrounding rivers and all species are substrate brooders with shared

paternal and maternal care. In contrast to the mouthbrooding species from the

Haplochromini, the *Lamprologini* show little sexual dimorphism and dichromatism, which

are well-known indicators for sexual selection (Kraaijeveld et al. 2011). We therefore expect

that the *Lamprologini* is a good candidate for picking up signals from water level changes.

We reconstructed a new *Lamprologini* tree following the workflow of the most complete

Lamprologini tree to date, which is a consensus tree based on the mitochondrial ND2 gene

(Sturmbauer et al. 2010), but we added three newly described species (*Lepidiolamprologus*

mimicus (Schelly et al. 2007)*, Neolamprologus timidus* (Kullander et al. 2014b) and

Chalinochromis cyanophleps (Kullander et al. 2014a)). Using phyloGenerator (Pearse and

Purvis 2013), we downloaded sequences from GenBank for nine genes (GenBank access

numbers can be found in the Supplementary Information). Genes were selected on the basis

of species coverage (at least 25% of the 79 Lamprologini species for which molecular data is

available), and whether or not the gene was crucial for inclusion of a species (e.g. for a

number of species, the only available gene was ND2). After selection, our full dataset

consisted of three mitochondrial genes: the NADH dehydrogenase subunit 2 (ND2 gene,

sequences from Kocher et al. 1995; Klett and Meyer 2002; Clabaut et al. 2005; Duftner et al.

2005; Schelly et al. 2006; Day et al. 2007; Koblmüller et al. 2007, 2016; Schwarzer et al.

2009; Wagner et al. 2009; Sturmbauer et al. 2010; O'Quin et al. 2010; Kullander et al. 2014b;

Weiss et al. 2015). The cytochrome b (cytb) gene (sequences from Salzburger et al. 2002;

Nevado et al. 2009; Wagner et al. 2009; O'Quin et al. 2010; Matschiner et al. 2011, 2016;

Kullander et al. 2014b; Shirai et al. 2014) and the cytochrome c oxidase subunit I (COI gene,

sequences from Sparks and Smith 2004; Nevado et al. 2013; Kullander et al. 2014a, 2014b;

Breman et al. 2016; Matschiner et al. 2016) and six nuclear genes: the nuclear locus 38A

(38A, sequences from Muschick et al. 2012; Meyer et al. 2016), the 18S ribosomal RNA

internal-transcribed spacer 1–2 with 5.8S and 28S ribosomal RNA partial sequences (18S,

sequences from Nevado et al. 2009; Koblmüller et al. 2016), the recombinase activating

protein 1 (rag1, sequences from Clabaut et al. 2005; Nevado et al. 2009; Kullander et al.

2014b; Shirai et al. 2014; Koblmüller et al. 2016; Meyer et al. 2016), the endothelin receptor

B1 gene (ednrb1, sequences from Muschick et al. 2012; Santos et al. 2014), the ribosomal

protein S7 (rps7, sequences from Schelly et al. 2006; Meyer et al. 2016)) gene and the rod

opsin gene (RH1, sequences from Sugawara et al. 2002; Spady et al. 2005; Nagai et al. 2011;

Meyer et al. 2015). GenBank access numbers for the used sequences can be found in the

supplementary material.

Sequences were aligned using MAFFT (setting: --auto) (Katoh and Standley 2013), and

subsequently, sequences were cleaned using trimAl (sites with >80% data missing were

removed, e.g. setting –gt 0.2) (Capella-Gutiérrez et al. 2009). Rather than concatenating the

alignments, we partitioned the data into subsets with independent sequence evolution models,

which is more suitable for a dataset which is expected to show incomplete lineage sorting or

hybridization (Meyer et al. 2016). To prepare alignments for use with partitionFinder,

alignments were combined using SequenceMatrix 1.8 (Vaidya et al. 2011). The best

partitioning found by partitionFinder 2.1.1 (Lanfear et al. 2012, 2016), partitioned the data

into 5 subsets (unlinked branches, AICc selection criterion), with all nuclear genes into one

subset (Rps7, ednrb1, 38A, 18S, RAG1 and RH1), with substitution model HKY+I+Γ. The

remaining three mitochondrial genes (ND2, COI and cytb) were placed in separate subsets,

each with a GTR+I+ Γ substitution model.

Using *BEAST (Heled and Drummond 2010) within the BEAST 2 package (Bouckaert et al.

2014), we inferred the time-calibrated species tree. We used an uncorrelated log-normal

relaxed clock and applied two calibration points. Firstly, we calibrated the crown of the

Lamprologini to be 4 million years old (log-normal prior, mean of 4 Myr 95% conf interval:

[3, 5]), based on the results from Meyer et al. (2016). Secondly, we included two riverine

Lamprologini species (*L. congoensis* and *L. teugelsi*), and calibrated the onset of their

branching event at 1.7 Ma (offset 1.1, log normal distribution with mean 1.7, 95% conf

interval [1.15, 3.47], "use originate = true"), following Koblmuller (2010). We applied 1/X

priors on the clock rates, and log-normal priors on the substitution rates. All other priors were

left at their default setting. As tree model we used the birth death model. The used BEAST

configuration file (the Beauti xml) can be found in the supplementary material.

We ran 10 independent STARBEAST MCMC chains, of 750M trees each. Each chain was

verified to have ESS values of at least 100 for all parameters. The first 10M trees were

pruned from these chains as burn-in and then they were combined (we used the species tree,

rather than the individual gene trees) into one large chain (of 7400M trees). Chains were

thinned by taking only each 5,000th tree. Using TreeAnnotator (from the BEAST 2 suite) we

constructed a Maximum Clade Credibility tree (using all 1.48M trees after thinning), storing

the mean heights.

We then pruned the tree from riverine species to obtain the pure *Lamprologini* tree on which

we fitted our model. Instead of performing one ABC-SMC inference on the obtained MCC

tree using a huge number of particles, which would be more accurate but computationally

extremely demanding, we performed 100 parallel inferences using 10,000 particles each. We

report the mean Bayes factor across these replicates.

Branching time uncertainty in the empirical tree

To account for uncertainty in the estimates of branching times in the *Lamprologini* tree we

sampled 100 trees from the posterior distribution obtained by *BEAST. Sampling was

performed at random, irrespective of the likelihood of the trees. In the Supplementary

material we show that the distribution of summary statistics of the subset of 100 trees is

similar to the distribution of the thinned chain. The 100 sampled trees were, like the

Maximum Clade Credibility tree, also pruned to remove the riverine taxa and stored

separately. For all 100 trees we performed both the ABC-SMC model selection algorithm and

the ABC-SMC parameter estimation algorithm, to determine the impact of different

branching times on the inferred water level model and associated parameters, and to

determine whether the MCC tree is a good representation of the underlying variability.

RESULTS

Lamprologini *phylogeny*

The onset of diversification within the *Lamprologini* is estimated to be around 3.96 Ma (95%

Highest Posterior Density interval (HPD): [3.09, 4.91]), which is very close to the prior we

put on the node age, based on previous estimates (Meyer et al. 2016). Furthermore, we

estimate the branching off of the Congo species (*N. congoensis* and *N. teugelsi*) from the

Lamprologini in Lake Tanganyika to have occurred around 1.35 Ma (HPD: [1.12, 1.68]),

which is a bit younger than previously obtained estimates (1.70 Ma, (Sturmbauer et al.

2010)). The topology of the Maximum Clade Credibility tree is largely consistent with

previous findings (Sturmbauer et al. 2010) (Figure 1). Placement of *Neolamprologus*

fasciatus as a close relative to *N. wauthioni* seems to re-iterate previously published evidence

for introgressive hybridization (Koblmüller et al. 2007). For the three species not previously

included in the *Lamprologini* phylogeny, *Lepidiolamprologus mimicus* was placed as a close

relative to the other species within the genus *Lepidiolamprologus*, *Chalinochromis*

cyanophleps was placed as a sister species to *Chalinochromis brichardi*, within the group of

Chalinochromis and *Julidochromis* species, and in agreement with previous analysis

(Kullander et al. 2014b). In contrast to previous findings (Kullander et al. 2014b),

Neolamprologus timidus is not placed as a sister species to *Neolamprologus furcifer*, but

rather associates with *N. mondabu* and *N. falcicula*. Again, in contrast to other previous

findings (Gante et al. 2016), we place *N. olivaceous* outside the *brichardi* complex, which

includes the model system species *N. brichardi* and *N. pulcher*. (but see the DensiTree

representation, which shows that this is not true for all trees). Interestingly, we also do not

infer *N. savory*i to be phylogenetically clustered within the *brichardi* complex (the 'Princess

cichlids' (Gante et al. 2016)), in contrast to Gante *et al.* (2016). We should take into account

however that the analysis by Gante *et al.* is based on on full genome sequences from only a

small group of species, in contrast to the limited number of markers from a large number of

species that we used.

As a reference we inferred speciation and extinction using the constant-rates birth-death

model (Nee et al. 1994). Using Maximum Likelihood (the function bd_ML in the DDD

package (Etienne et al. 2012)), we obtained an estimate of 1.871 myr^{-1} for the speciation rate,

and an estimate of 0.993 myr^{-1} for the extinction rate, for the MCC tree. We find an estimate

of 0.87 myr^{-1} for the diversification rate (speciation - extinction) and an estimate of 0.531

myr^{-1} for the turnover rate (extinction / speciation). For the 100 trees sampled from the full

chain, we obtain estimates of 3.02 myr^{-1} (95% HPD: [1.608, 4.947]) for the speciation rate

and 2.409 myr^{-1} (95% HPD: [0.884, 4.530]) for the extinction rate. This translates into

estimates of 0.61 myr^{-1} (95% HPD: [0.313, 0.985]) for the diversification rate, and 0.765 myr^{-}

^{1} (95% HPD: [0.518, 0.930]) for the turnover rate. Estimates for the birth-death model

obtained during reconstruction of the tree using BEAST indicate an estimate of 0.864 myr^{-1}

(95% HPD: [0.287, 1.459]) for the speciation rate and an estimate of 0.613 myr^{-1} (95% HPD:

[0.181, 0.953]) for the relative death rate, which translates into an estimate for the extinction

rate of 0.52 myr^{-1} (95% HPD: [0.156, 0.823]) per million years. This yields estimates of

0.334 myr^{-1} and 0.613 myr^{-1}for the diversification and turnover rate respectively. The BEAST

inferences include the riverine species, so speciation and extinction rates are expected to be a

bit different.

Figure 2. Phylogenetic hypothesis for the *Lamprologini* and outgroups, based on 3
mitochondrial and 6 nuclear genes, and two calibrations: 4 million years for the root of the
Lamprologini clade, and 1.1 - 3.5 million years for the Congo *Lamprologini* species. Left
panel: DensiTree (Bouckaert and Heled 2014) representation of the MCMC chain obtained
using *BEAST. Shown are trees from a thinned posterior chain, after selecting every
100,000[th] tree. Riverine species are indicated in grey. Right panel: Maximum Clade
Credibility tree. Bars around the node span the 95% HPD for each node. Please note that for
the dual display of both the densitree representation and the MCC phylogeny, some tips of
the MCC phylogeny might appear slightly misaligned. A high resolution version of both the
Densitree representation and the MCC phylogeny can be found in the supplementary
information.

Parameter estimation

We estimated parameter values for the three models for all 100 trees sampled from the

posterior. We report the parameter values across the combined posterior across all 100 trees.

Note that variation in the parameter estimates results from two sources of variation:

branching time variation across the 100 trees, and secondly variation in the parameter

estimate within each ABC-SMC inference.

The model without water level changes is identical to the constant-rates birth-death model,

and we find that our ABC-SMC estimates for sympatric speciation at high water level (λ_s^h)

are slightly lower than the Maximum Likelihood estimate of the birth rate under the constant

rates birth-death model (2.644 Myr^{-1}(95% HPD: [1.208, 4.633], see also Table 1) versus 3.02,

see also Table 1). Similarly, we infer the extinction rate (μ) to also be slightly lower (1.950

Myr^{-1} (95% HPD: [0.188, 4.101]) versus 2.409, see also Table 1). We obtain estimates of

0.694 and 0.738 for diversification and turnover respectively, which are close to the estimates

obtained using Maximum Likelihood (a diversification rate of 0.610 and a turnover rate of

0.765 respectively). Taking into account the 95% confidence intervals on the obtained

parameter estimates and the fact that the ABC-SMC estimates are potentially affected by the

prior while the ML estimates are not, we are confident that estimates obtained using our

ABC-SMC method for the model without water level changes are consistent with the

maximum likelihood estimates under the constant-rates birth-death model.

Using the LW model, which implements water level changes following the literature (e.g.

high water level until ~1.1 Ma, after which a series of water level changes took place), we

infer a lower rate of sympatric speciation at high water level (0.871 Myr^{-1} (95% HPD: [0.227,

3.642])), which is compensated with a high rate of allopatric speciation (6.412 Myr^{-1} (95%

HPD: [0.001, 14.195])) but not with a high rate of sympatric speciation at low water level

(0.028 Myr^{-1} (95% HPD: [0.001, 0.651])), suggesting that water level dynamics are important

drivers of biodiversity, but only through allopatric speciation. Extinction is inferred to be low

(0.037 Myr^{-1} (95% HPD: [0.001, 2.133])). Because of the non-trivial relationship between

speciation at high and low water level, we can no longer calculate diversification and

turnover rates.

Using the EW model, where water level changes are extrapolated beyond 1.1 Ma, we observe

that the rate of sympatric speciation at high water level is inferred to be similar to without

water level changes (2.753 Myr^{-1} (95% HPD: [1.347, 4.383])). Extinction, however, is lower

(0.111 Myr^{-1} (95% HPD: [0.001, 1.627])), and allopatric speciation and sympatric speciation

at low water level are both inferred to be much lower than for the literature water scenario

(0.022 Myr^{-1} (95% HPD: [0.001, 0.466]) and 0.033 Myr^{-1} (95% HPD: [0.001, 0.504])

respectively).

Across the three water level models we observe that the distribution of the post-hoc

perturbations σ does not differ substantially from the prior for the NW and EW water models,

with low estimates (0.036 (95% HPD: [0.001, 0.569]) and 0.030 (95% HPD: [0.001, 0.484])

for the NW and EW model respectively, Table 1). We notice a much higher value of σ

associated with LW (0.174, (95% HPD: [0.001, 0.680])), which also has a much higher

estimate for allopatric speciation at low water level. Allopatric speciation at low water level

potentially causes temporal alignment of branching times and we introduced the parameter σ

to correct simulated phylogenies for this, to allow comparison with phylogenies generated by

*BEAST, which does not allow for temporally aligned branching times. Hence, the higher

inferred value of σ for the LW model confirms the validity of the application of our post-hoc

perturbation.

Table 1. Median posterior density estimate, for sympatric speciation at high water (λ_s^h), extinction (μ), perturbation (σ), sympatric speciation at low water (λ_s^l) and allopatric speciation (λ_a^l). Shown are results for the model with no water level changes (NW), literature values for water level changes (LW) and water level changes extrapolated beyond the literature range (EW). The 95% credibility interval is shown between square brackets. All values are rates per million years.

	λ_s^h	μ	σ	λ_a^l	λ_s^l
NW	2.644 [1.208, 4.633]	1.950 [0.188, 4.101]	0.036 [0.001, 0.569]		
LW	0.871 [0.227, 3.642]	0.037 [0.001, 2.133]	0.174 [0.001, 0.68]	6.412 [0.001, 14.195]	0.028 [0.001, 0.651]
EW	2.753 [1.347, 4.383]	0.111 [0.001, 1.627]	0.030 [0.001, 0.484]	0.022 [0.001, 0.466]	0.033 [0.001, 0.504]

Figure 3. Posterior densities of the pooled posterior distribution across 100 randomly drawn trees from the posterior MCMC chain. Shown are estimates for the three water level scenarios (no water level changes (NW), literature values for water level changes (LW) and extrapolated values for water level changes (EW)). Shown are the posterior density (black line) and the 95% credibility interval (shaded area, blue for NW, gold for LW and green for EW. X-axes are on a ^{10}log scale. The first column shows a sample water level profile, with the water level on the y-axis, and the time before present (in million years) on the x-axis. Note that for the EW model, for each simulation a new profile was generated, and that the shown profile is only one example of such a profile. Because without water level changes, λ_a^l and λ_s^l have no meaning, their posterior distribution is not shown for the NW scenario.

Model fitting

Figure 4. Model selection results on 100 trees randomly drawn from the *BEAST posterior of the *Lamprologini* tree. The top row shows 2 ln (Bayes factors) comparing posterior support of the LW (literature water changes) model with the NW (no water level changes) model, the bottom row shows 2 ln (Bayes factors) of the comparison between the posterior support for the LW model with the EW (extrapolated water level changes) model. A 2ln(Bayes factor) higher than 2 is generally considered to provide substantial evidence in favor of the respective model (Kass and Raftery 1995), which is indicated by the solid lines. The thin dotted line indicates the median 2 ln (Bayes factor) obtained for the MCC tree, for which we do not find substantial support for any of the three models. The thick dotted line indicates the median 2 ln (Bayes factor) for the trees drawn from the *BEAST posterior (e.g. the median of the distribution shown), which is in both cases above 2, indicating substantial support for the LW model compared to the other two models. 2 ln(Bayes factors) higher than 10 are grouped together into one category.

Model selection

When we apply the model selection algorithm to the MCC tree, we find median Bayes factors

(we report here not the raw numbers, but 2 ln(Bayes factor), but for brevity refer to them as

Bayes factors) of 1.64 and 0.9 when comparing the LW model with the NW and EW model

respectively. We thus find no convincing evidence for any of the three models, when fitting

our model to the MCC tree. Alternatively, when we fit to 100 trees randomly sampled from

the *BEAST posterior, we find Bayes factors of 3.60 and 3.65 when comparing LW model

with the NW and EW model respectively. Furthermore, in 77 out of the 100 trees we select

the LW model as the most likely model (based on the Bayes factor), in 17 out of 100 trees we

select the EW model, and only in 6 out of 100 trees we select the model without any water

level changes.

Figure 5. Validation of the ability of our ABC-SMC algorithm to infer the correct model. 100 replicate datasets were generated for each water level model (no water level changes NW, water level changes from the literature LW, or water level changes extrapolated beyond the literature range, EW). The plots show the distribution of the 2 ln(Bayes factor) across all 100 replicate inferences. The dotted line indicates the median 2 ln(Bayes factor). A 2ln(Bayes factor) higher than 2 is generally considered to provide substantial evidence in favor of the respective model (Kass and Raftery 1995). 2 ln(Bayes factors) higher than 10 are grouped together into one category.

Model validation shows that when we simulated data using the NW model, the NW model was selected using our model validation algorithm more than the other two models (59 out of 100 replicates). Median Bayes factors are both higher than 2, with a median of 5.28 and 3.83 versus the LW and EW model respectively, supporting considerable support for the NW

model over the other two models. When data was simulated with the LW model, we selected

the correct model in the majority of 100 replicates (84 out of 100 replicates). The Bayes

factors reflect this, with medians of 18.4 (this is the maximum score) versus both the NW and

EW model. Lastly, when we simulated data using the EW model, we selected the correct

model more than the other two models, in 65 out of 100 replicates. This was reflected by the

Bayes factors as well, as the median Bayes factor versus the NW model was 18.4, and the

median Bayes factor versus the LW model was 7.47.

More interesting is the correct detection rate of a model, which is given by the number of

trees simulated by the model that is selected for that tree. This is equal to asking whether,

given posterior support for a respective model, we also find that the tree for which we find

this support was simulated with the respective model. If our model selection procedure can

not detect models accurately, we expect a detection rate of around 50%, as support is always

divided between two (not three) models. Detection rates larger than 50% support the

conclusion that our model selection procedure can adequately infer the correct model.

We find that across the 300 simulated trees, 120 trees received considerable support for the

LW model over the NW model (e.g. 2 ln (BF LW/NW) > 2), of these 120 trees, 92 trees were

simulated with the LW model, which leads to a correct detection rate of 77% (See Figure 6).

Furthermore, out of 107 trees that received considerable support for the LW model over the

EW model, we find that 83 trees were simulated using the LW model, which translates to a

detection rate of 78%. We find similar detection rates for the NW model: 90% against the

LW model (62 out of 69 detected trees) and 83% against the EW model (54 out of 65

detected trees). Lastly, detection rates for the EW model mirror these findings: a detection

rate of 79% against the NW model (79 out of 100 detected trees), and of 86% against the LW

model (68 out of 79 detected trees).

Posterior support for

Figure 6. Accuracy of assignment of models depending on their posterior support. **A**: The relative fraction of trees simulated with either LW (dark) or NW (light), receiving support for LW (2ln(Bayes Factor LW/NW) > 2), support for NW (2ln(Bayes Factor LW/NW) < -2), or receiving no support for either model. **B**: The relative fraction of trees simulated with either

LW (dark) or EW (light), receiving support for LW (2ln(Bayes Factor LW/EW) > 2), support
for EW (2ln(Bayes Factor LW/EW) < -2), or receiving no support for either model. **C**: The
relative fraction of trees simulated with either NW (dark) or EW (light), receiving support for
NW (2ln(Bayes Factor NW/EW) > 2), support for EW (2ln(Bayes Factor NW/EW) < -2), or
receiving no support for either model.

DISCUSSION

We have presented a model that infers past speciation and extinction rates, and their

interactions with changes in the environment, from a given phylogeny. We have shown that

our model is able to accurately select between different scenarios, including or excluding

environmental change. We applied our model to an updated phylogeny of the cichlid fish

tribe of *Lamprologini* and found evidence that past water level changes have shaped current

cichlid diversity in Lake Tanganyika, when we applied our model to a sample from the

posterior distribution of trees of the *Lamprologini*, as inferred by *BEAST. We asked the

model to select the best fitting of three scenarios: a scenario without any water level changes,

a scenario using the values found in the literature, and a scenario using the mean rate of water

level change found in the literature to extrapolate water level changes beyond the range of

literature values available. We found that the model following literature water levels received

most support, which suggests that water level changes have been an important driver of

diversity in the *Lamprologini*. We note that a model without effect of water level changes on

diversification (NW) can sometimes generate patterns that resemble the predictions of the

preferred model (LW). Yet, we find when fitting our model to trees drawn from the *BEAST

posterior that the distribution of Bayes Factors is skewed towards the model following

literature water levels (LW) and we find support for the model without an effect of water

level changes on diversification (NW) only for a small number of trees, suggesting that this

effect is relatively small.

When we applied our model selection algorithm on the Most Credible Consensus (MCC)

tree, we found contrasting results. Support for both models including water level changes

diminished, and posterior support for the model without any water level changes increased.

Nevertheless, no single model could yield enough support to convincingly reject the other

two. Moreover, results using the MCC tree are markedly different from those using trees

sampled from the posterior. We conclude therefore that the MCC tree, at least for the

Lamprologini, but most likely more generally, provides a poor summary of the true species

tree and of the underlying variation in branching patterns. Hence, we suggest to avoid

reporting MCC trees, and instead to provide the reader with the full posterior distribution, for

instance through a DensiTree plot (Bouckaert and Heled 2014). Posterior inference, for

instance of speciation and extinction rates should preferentially also be performed on multiple

independent samples from the posterior, rather than on the MCC tree, as the underlying

variation might lead to very different results, as we have shown here.

Discrepancies between the MCC tree and the posterior distribution of trees could also

potentially clarify previously recovered inconsistencies when studying diversification, for

example in shrews in the Philippines. The Philippines have been subject to strong sea level

fluctuations, causing the fission and fusion of several islands, primarily during the

Pleistocene (Brown et al. 2013). Population genetic evidence has convincingly shown that the

location of such fused islands correlates strongly with genetic divergence between

populations in many different species (Evans et al. 2003; Linkem et al. 2010; Siler et al.

2010; Oaks et al. 2013). Phylogenetic analysis however, has failed to show any evidence of

diversification associated with Pleistocene water level changes (Esselstyn and Brown 2009).

The basis for this phylogenetic analysis however, was an MCC tree. Repeating the analysis

on the posterior distribution underlying the MCC tree could mitigate these problems, and

could clarify the impact of Pleistocene water level changes on diversification in the

Philippines archipelago.

When allopatric speciation rates are high, the resulting phylogenetic trees have internal nodes that have synchronized branching times, e.g. branching times that align with episodes of water level change. Although Phylogenetic reconstruction software is able to infer simultaneous branching events, it typically uses only two parameters (birth and death) to infer all branching events of the tree. Therefore, if it can accommodate the simultaneous events, it is unlikely to fit well to the non-simultaneous events, and vice-versa. Our finding of evidence for a substantial role of habitat dynamics in diversification can therefore be regarded as conservative. To improve the fit of trees generated by our model with trees generated by *BEAST we included an *a posteriori* perturbation parameter in our model. This parameter determines the standard deviation of a Gaussian perturbation kernel that is applied to each node after the simulation has completed. By perturbing each node, we minimized the probability that branching times align in time. We found that standard deviation increased in size with an increase in allopatric speciation, as expected. A less *ad hoc* solution to deal with the alignment of branching times in the tree would be to incorporate the model presented here as a tree prior in phylogenetic reconstruction software. Although this need not introduce any significant differences in the tree topology, the distribution of branching times could be substantially influenced, and any subsequent inference focusing on such patterns could be very different. Including such models in tree reconstruction software may require incorporation of ABC methods, and will be extremely computationally demanding, but our results justify such an endeavor.

Given that water level changes are only prevalent during the last million years before present, we cannot exclude the possibility that increased diversification due to reasons other than

changing water levels has driven diversification during this period. On average, the LW

model could be represented by a simple birth-death model with a rate shift around one

million years ago. We expect however that although such a model could accommodate the

increased average diversification, it cannot replicate temporal alignment in branching events

due to water level changes. To examine this in more detail, we fitted a simple birth-death

model with a rate shift around one million years ago to the trees obtained from the posterior

(see Supplementary Information).. In the absence of a likelihood for the LW model, we

compared the nLTT statistic for the rate-shift model with that of the LW model, as the nLTT

statistic should be sensitive to detecting temporal alignment of branching events, We find that

our model is much closer to the empirical data than the rate shift model. We attempted to

improve the fit of the rate-shift model by allowing the speciation rate in the model to shift up

and down in line with the literature values of the water level changes. The two rates inferred

by the model then represent speciation at low, and at high water level respectively. Although

we do find an increase in the rate of speciation at low water level, the fit of this rate-shift

model is still worse than that of the LW model. This supports our conclusion that water level

changes influence the phylogeny not only through an increased speciation rate, but also

through temporal alignment of branching times.

Although we refer in our model to the different implementations of speciation as sympatric

and allopatric, care should be taken in interpreting these forms of speciation. We consider

here allopatric speciation only on a large scale, where populations become allopatric over

stretches of hundreds of kilometers (Sturmbauer et al. 2001). Large-scale isolation might not

be necessary for cichlids, as some species can already be limited in gene flow by a sand

stretch of 50 meters separating populations (Rico and Turner 2002). Such micro-allopatric

speciation events are not captured by the allopatric speciation rate in our model. Rather, these

local scale events are captured in our model by sympatric speciation. Hence, sympatric

speciation in our model covers all degrees of speciation ranging from full sympatry to

allopatry, providing that geographical isolation is smaller than that imposed by a water level

change. Allopatric speciation in our model then solely refers to speciation events caused by

geographical isolation over a large distance, driven by changes in water level, and inducing

simultaneous branching events.

In our model we have assumed that when the water level drops, species distribute themselves

equally over the two pockets of water that survive the water level drop. A more realistic

model would allow for a skew towards one of the pockets, either dependent on the respective

sizes of the pockets, the distribution of the species over the lake at high water level, or both.

We have here refrained from including a parameter that regulates the distribution of species

over the two pockets in order to avoid over fitting. Another possible extension of our model

would lie into extending the approach towards three or more pockets, possibly combined with

a parameter governing the distribution of species across these three pockets during a water

level drop. Bathymetric maps of Lake Tanganyika suggest that for some water level changes

it might split into three lakes (Coulter 1991). How a split of a species into three populations,

and associated allopatric divergence and speciation, affects phylogenetic structure and affects

temporal alignment in branching times remains currently unexplored and would be an

interesting avenue for future work.

Our results are strongly in line with population genomic analyses in a number of cichlid

species including *Eretmodus cyanostictus* (Verheyen et al. 1996), *Tropheus moorii*

(Koblmüller et al. 2011; Nevado et al. 2013; Sefc et al. 2017)*, Variabilichromis moorii*

(Nevado et al. 2013)*, Altolamprologus* (Koblmüller et al. 2016) and *Telmatochromis*
temporalis (Winkelmann et al. 2016)*, and resonate with population genomic findings across
the three African Rift Lakes (Sturmbauer et al. 2001). Furthermore, population genetic
studies have shown that water level fluctuations in Lake Malawi have been associated with
population expansion in cichlid species (Arnegard et al. 1999; Sturmbauer et al. 2001;
Genner et al. 2010), suggesting a potential role for water level changes in Lake Malawi as
well. Phylogenetic reconstruction for Malawi cichlid species is problematic however,
partially due to the young age of the species. However, considering that the geological record
of Lake Malawi spans a much larger part of the total lifespan of the lake (Delvaux 1995;
Lyons et al. 2015; Ivory et al. 2016) and thus provides a much better record about water level
fluctuations since the colonization of the lake by cichlids, we expect that modern genetic
developments will soon allow for a thorough understanding of the impact of water level
changes on cichlids in Lake Malawi as well.

Conclusion

Our model integrates standard constant-rate birth-death mechanics with environmental
change and with speciation induced by geographical isolation. We analyzed the phylogeny of
the tribe of *Lamprologini* to see whether past water level changes in Lake Tanganyika have
contributed to the current diversity of cichlid fish in Lake Tanganyika. We find an important
role for environmental changes in driving diversity, and find evidence that past water level
changes have shaped current standing diversity in the tribe of *Lamprologini*. However, we
found that inference of past environmental changes from a single phylogeny, and more
specifically, from the MCC tree, tends to lead to unreliable results. We therefore advocate
caution when using the MCC tree as a basis for further analysis. Furthermore, we argue for

the inclusion of more detailed branching models in phylogenetic reconstruction software,

which allow for the inclusion of an interaction between the environment, and speciation rates.

Acknowledgements

We thank Lucas Molleman for useful discussions. We thank the Netherlands Organisation for

Scientific Research (NWO) for financial support through VIDI and VICI grants awarded to

RSE. We thank the Donald Smits Center for information Technology of the University of

Groningen for their support and providing access to the Millipede and Peregrine high-

performance computing cluster. We thank the Max Planck Institute for Evolutionary Theory

for their support and providing access to their high-performance computing cluster. We thank

the Carl von Ossietzky Universität Oldenburg for their support and providing access to the

CARL computing cluster.

REFERENCES

Aguilée R., Claessen D., Lambert A. 2013. Adaptive radiation driven by the interplay of eco-
evolutionary and landscape dynamics. Evolution (N. Y). 67:1291–1306.

Aguilée R., Lambert A., Claessen D. 2011. Ecological speciation in dynamic landscapes. J.
Evol. Biol. 24:2663–77.

Alin S., Cohen A., Bills R. 1999. Effects of landscape disturbance on animal communities in
Lake Tanganyika, East Africa. Conservation. 13:1017–1033.

Alin S., Cohen A.S. 2003. Lake-level history of Lake Tanganyika, East Africa, for the past
2500 years based on ostracode-inferred water-depth reconstruction. Palaeogeogr.
Palaeoclimatol. Palaeoecol. 199:31–49.

Arnegard M.E., Markert J.A., Danley P.D., Stauffer J.R., Ambali A.J., Kocher T.D. 1999.
Population structure and colour variation of the cichlid fishes Labeotropheus fuelleborni
Ahl along a recently formed archipelago of rocky habitat patches in southern Lake
Malawi. Proc. R. Soc. Lond. B. 266:119–130.

Barnosky A. 2005. Effects of Quaternary climatic change on speciation in mammals. J.
Mamm. Evol. 12:247–264.

Bouckaert R., Heled J. 2014. DensiTree 2: Seeing trees through the forest. bioRxiv.:1–11.

Bouckaert R., Heled J., Kühnert D., Vaughan T., Wu C.H., Xie D., Suchard M.A., Rambaut
A., Drummond A.J. 2014. BEAST 2: A Software Platform for Bayesian Evolutionary
Analysis. PLoS Comput. Biol. 10:1–6.

Brawand D., Wagner C.E., Li Y.I., Malinsky M., Keller I., Fan S., Simakov O., Ng A.Y.,
Lim Z.W., Bezault E., Turner-Maier, J. Johnson J., Alcazar R., Noh H.J., Russell P.,
Aken B., Alföldi J., Amemiya C., Azzouzi N., Baroiller J.-F., Barloy-Hubler F., Berlin
A., Bloomquist R., Carleton K.L., Conte M.A., D'Cotta H., Eshel O., Gaffney L.,
Galibert F., Gante H.F., Gnerre S., Greuter L., Guyon R., Haddad N.S., Haerty W.,
Harris R.M., Hofmann H. a., Hourlier T., Hulata G., Jaffe D.B., Lara M., A.P. L.,
MacCallum I., Mwaiko S., Nikaido M., Nishihara H., Ozouf-Costaz C., Penman D.J.,
Przybylski D., Rakotomanga M., Renn S.C.P., Ribeiro F.J., Ron M., Salzburger W.,
Sanchez-Pulido L., Santos M.E., Searle S., Sharpe T., Swofford R., Tan F.J., Williams
L., Young S., Yin S., Okada N., Kocher T.D., Miska E. a., Lander E.S., Venkatesh B.,
Fernald R.D., Meyer A., Ponting C.P., Streelman J.T., Lindblad-Toh K., Seehausen O.,
Di Palma F. 2014. The genomic substrate for adaptive radiation in African cichlid fish.
Nature. 513:375–381.

Breman F.C., Loix S., Jordaens K., Snoeks J., Van Steenberge M. 2016. Testing the potential
of DNA barcoding in vertebrate radiations: the case of the littoral cichlids (Pisces,
Perciformes, Cichlidae) from Lake Tanganyika. Mol. Ecol. Resour. 16:1455–1464.

Brown R.M., Siler C.D., Oliveros C.H., Esselstyn J. a., Diesmos A.C., Hosner P. a., Linkem
C.W., Barley A.J., Oaks J.R., Sanguila M.B., Welton L.J., Blackburn D.C., Moyle R.G.,
Townsend Peterson a., Alcala A.C. 2013. Evolutionary processes of diversification in a
model island archipelago. Annu. Rev. Ecol. Evol. Syst. 44:411–435.

Capella-Gutiérrez S., Silla-Martínez J.M., Gabaldón T. 2009. trimAl: A tool for automated
alignment trimming in large-scale phylogenetic analyses. Bioinformatics. 25:1972–
1973.

Clabaut C., Salzburger W., Meyer A. 2005. Comparative phylogenetic analyses of the
adaptive radiation of Lake Tanganyika cichlid fish: nuclear sequences are less
homoplasious but also less informative than mitochondrial DNA. J. Mol. Evol. 61:666–
81.

Cohen A.S., Lezzar K.E., Tiercelin J.J., Soreghan M. 1997a. New palaeogeographic and lake-
level reconstructions of Lake Tanganyika: implications for tectonic, climatic and
biological evolution in a rift lake. Basin Res. 9:107–132.

Cohen A.S., Soreghan M., Scholz C.A. 1993. Estimating the age of formation of lakes: an
example from Lake Tanganyika, East African Rift system. Geology. 21:511.

Cohen A.S., Stone J.R., Beuning K.R.M., Park L.E., Reinthal P.N., Dettman D., Scholz C.A.,
Johnson T.C., King J.W., Talbot M.R., Brown E.T., Ivory S.J. 2007. Ecological
consequences of early Late Pleistocene megadroughts in tropical Africa. Proc. Natl.
Acad. Sci. 104:16422–7.

Cohen A.S., Talbot M.R., Awramik S.M., Dettman D.L., Abell P. 1997b. Lake level and
paleoenvironmental history of Lake Tanganyika, Africa, as inferred from late Holocene
and modern stromatolites. Geol. Soc. Am. Bull. 109:444–460.

Coulter G. 1991. Lake Tanganyika and its life. Oxford University Press.

Coyne J., Orr H. 2004. Speciation. Sunderland, Massachusets U.S.A.: Sinauer Associates.

Day J.J., Santini S., Garcia-Moreno J. 2007. Phylogenetic relationships of the Lake
Tanganyika cichlid tribe Lamprologini: the story from mitochondrial DNA. Mol.
Phylogenet. Evol. 45:629–42.

Delvaux D. 1995. Age of Lake Malawi (Nyasa) and water level fluctuations. Mus. roy. Afrr.
centr., Tervuren (Belg.), Dept. Geol. Min., Rapp. ann. 1993 1994. 108:99–108.

Duftner N., Koblmüller S., Sturmbauer C. 2005. Evolutionary relationships of the
Limnochromini, a tribe of benthic deepwater cichlid fish endemic to Lake Tanganyika,
East Africa. J. Mol. Evol. 60:277–289.

Esselstyn J.A., Brown R.M. 2009. The role of repeated sea-level fluctuations in the
generation of shrew (Soricidae: Crocidura) diversity in the Philippine Archipelago. Mol.
Phylogenet. Evol. 53:171–181.

Etienne R.S., Haegeman B., Stadler T., Aze T., Pearson P.N., Purvis A., Phillimore A.B.
2012. Diversity-dependence brings molecular phylogenies closer to agreement with the
fossil record. Proc. R. Soc. B Biol. Sci. 279:1300–1309.

Etienne R.S., Rosindell J. 2012. Prolonging the past counteracts the pull of the present:
protracted speciation can explain observed slowdowns in diversification. Syst. Biol.
61:204–13.

Evans B.J., Brown R.M., McGuire J.A., Supriatna J., Andayani N., Diesmos A., Iskandar D.,
Melnick D.J., Cannatella D.C. 2003. Phylogenetics of fanged frogs: testing

biogeographical hypotheses at the interface of the asian and Australian faunal zones.
Syst. Biol. 52:794–819.

Gante H.F., Matschiner M., Malmstrøm M., Jakobsen K.S., Jentoft S., Salzburger W. 2016.
Genomics of speciation and introgression in Princess cichlid fishes from Lake
Tanganyika. Mol. Ecol.

Genner M.J., Knight M.E., Haesler M.P., Turner G.F. 2010. Establishment and expansion of
Lake Malawi rock fish populations after a dramatic Late Pleistocene lake level rise. Mol.
Ecol. 19:170–82.

Genner M.J., Seehausen O., Lunt D.H., Joyce D.A., Shaw P.W., Carvalho G.R., Turner G.F.
2007. Age of cichlids: new dates for ancient lake fish radiations. Mol. Biol. Evol.
24:1269–82.

Glor R.E., Gifford M.E., Larson A., Losos J.B., Schettino L.R., Chamizo Lara A.R., Jackman
T.R. 2004. Partial island submergence and speciation in an adaptive radiation: a
multilocus analysis of the Cuban green anoles. Proc. R. Soc. B Biol. Sci. 271:2257–65.

Heaney L.R. 1985. Zoogeographic Evidence for Middle and Late Pleistocene Land Bridges
to the Philippine Islands. Mod. Quat. Res. Southeast Asia. 9:127–143.

Heled J., Drummond A.J. 2010. Bayesian Inference of Species Trees from Multilocus Data.
Mol. Biol. Evol. 27:570–580.

Hutter C.R., Guayasamin J.M., Wiens J.J. 2013. Explaining Andean megadiversity: The
evolutionary and ecological causes of glassfrog elevational richness patterns. Ecol. Lett.
16:1135–1144.

Ivory S.J., Blome M.W., King J.W., McGlue M.M., Cole J.E., Cohen A.S. 2016.
Environmental change explains cichlid adaptive radiation at Lake Malawi over the past
1.2 million years. Proc. Natl. Acad. Sci.:201611028.

Janzen T., Höhna S., Etienne R.S.R.S. 2015. Approximate Bayesian Computation of
diversification rates from molecular phylogenies: introducing a new efficient summary
statistic, the nLTT. Methods Ecol. Evol. 6:566–575.

Kass R., Raftery A. 1995. Bayes Factors. J. Amer. Stat. Assoc. 90:773–795.

Katoh K., Standley D.M. 2013. MAFFT multiple sequence alignment software version 7:
Improvements in performance and usability. Mol. Biol. Evol. 30:772–780.

Klett V., Meyer A. 2002. What, if anything, is a Tilapia?-mitochondrial ND2 phylogeny of
tilapiines and the evolution of parental care systems in the African cichlid fishes. Mol.
Biol. Evol. 19:865–883.

Koblmüller S., Duftner N., Sefc K.M., Aibara M., Stipacek M., Blanc M., Egger B.,
Sturmbauer C. 2007. Reticulate phylogeny of gastropod-shell-breeding cichlids from
Lake Tanganyika--the result of repeated introgressive hybridization. BMC Evol. Biol.
7:7.

Koblmüller S., Nevado B., Makasa L., van Steenberge M., Vanhove M.P.M., Verheyen E.,
Sturmbauer C., Sefc K.M. 2016. Phylogeny and phylogeography of Altolamprologus:
ancient introgression and recent divergence in a rock-dwelling Lake Tanganyika cichlid

genus. Hydrobiologia.:1–16.

Koblmüller S., Salzburger W., Obermüller B., Eigner E., Sturmbauer C., Sefc K.M. 2011.
Separated by sand, fused by dropping water: habitat barriers and fluctuating water levels
steer the evolution of rock-dwelling cichlid populations in Lake Tanganyika. Mol. Ecol.
20:2272–90.

Kocher T.D., Conroy J. a, McKaye K.R., Stauffer J.R., Lockwood S.F. 1995. Evolution of
NADH dehydrogenase subunit 2 in east African cichlid fish. Mol. Phylogenet. Evol.
4:420–432.

Kraaijeveld K., Kraaijeveld-Smit F.J.L., Maan M.E. 2011. Sexual selection and speciation:
the comparative evidence revisited. Biol. Rev. Camb. Philos. Soc. 86:367–77.

Kullander S.O., Karlsson M., Norén M. 2014a. Chalinochromis cyanophleps, a new species
of cichlid fish (Teleostei: Cichlidae) from Lake Tanganyika. Zootaxa. 3790:425–438.

Kullander S.O., Norén M., Karlsson M., Karlsson M. 2014b. Description of Neolamprologus
timidus, new species, and review of N . furcifer from Lake Tanganyika (Teleostei:
Cichlidae). Ichthyol.Explor.Freshwaters. 24:301–328.

Lanfear R., Calcott B., Ho S.Y.W., Guindon S. 2012. PartitionFinder: Combined selection of
partitioning schemes and substitution models for phylogenetic analyses. Mol. Biol. Evol.
29:1695–1701.

Lanfear R., Frandsen P.B., Wright A.M., Senfeld T., Calcott B. 2016. PartitionFinder 2: New
Methods for Selecting Partitioned Models of Evolution for Molecular and
Morphological Phylogenetic Analyses. Mol. Biol. Evol.:msw260.

Lezzar K.E., Tiercelin J.J., Batist M., Cohen A.S., Bandora T., Rensbergen P., Turdu C.,
Mifundu W., Klerkx J. 1996. New seismic stratigraphy and Late Tertiary history of the
North Tanganyika Basin, East African Rift system, deduced from multichannel and
high-resolution reflection seismic data and piston core evidence. Basin Res. 8:1–28.

Linkem C.W., Hesed K.M., Diesmos A.C., Brown R.M. 2010. Species boundaries and
cryptic lineage diversity in a Philippine forest skink complex (Reptilia; Squamata;
Scincidae: Lygosominae). Mol. Phylogenet. Evol. 56:572–585.

Lyons R.P., Scholz C.A., Cohen A.S., King J.W., Brown E.T., Ivory S.J., Johnson T.C.,
Deino A.L., Reinthal P.N., Mcglue M.M., Blome M.W. 2015. Continuous 1.3-million-
year record of East African hydroclimate, and implications for patterns of evolution and
biodiversity. PNAS.:2–7.

Matschiner M., Hanel R., Salzburger W. 2011. On the origin and trigger of the notothenioid
adaptive radiation. PLoS One. 6.

Matschiner M., Musilová Z., Barth J.M.I., Starostova Z., Salzburger W., Steel M., Bouckaert
R. 2016. Bayesian Phylogenetic Estimation of Clade Ages Supports Trans-Atlantic
Dispersal of Cichlid Fishes. Syst. Biol.

McGlue M.M., Lezzar K.E., Cohen A.S., Russell J.M., Tiercelin J.-J., Felton A. a., Mbede E.,
Nkotagu H.H. 2008. Seismic records of late Pleistocene aridity in Lake Tanganyika,
tropical East Africa. J. Paleolimnol. 40:635–653.

Meyer B.S., Matschiner M., Salzburger W. 2015. A tribal level phylogeny of Lake
Tanganyika cichlid fishes based on a genomic multi-marker approach. Mol. Phylogenet.
Evol. 83:56–71.

Meyer B.S., Matschiner M., Salzburger W. 2016. Disentangling incomplete lineage sorting
and introgression to refine species-tree estimates for Lake Tanganyika cichlid fishes.
Syst. Biol. syw069:1–20.

Muschick M., Indermaur A., Salzburger W. 2012. Convergent Evolution within an Adaptive
Radiation of Cichlid Fishes. Curr. Biol.:1–7.

Nagai H., Terai Y., Sugawara T., Imai H., Nishihara H., Hori M., Okada N. 2011. Reverse
evolution in RH1 for adaptation of cichlids to water depth in Lake Tanganyika. Mol.
Biol. Evol. 28:1769–1776.

Nee S., May R.M., Harvey P.H. 1994. The reconstructed evolutionary process. Philos. Trans.
R. Soc. Lond. B. Biol. Sci. 344:305–11.

Nevado B., KoblmÜller S., Sturmbauer C., Snoeks J., Usano-Alemany J., Verheyen E. 2009.
Complete mitochondrial DNA replacement in a Lake Tanganyika cichlid fish. Mol.
Ecol. 18:4240–4255.

Nevado B., Mautner S., Sturmbauer C., Verheyen E. 2013. Water-level fluctuations and
metapopulation dynamics as drivers of genetic diversity in populations of three
Tanganyikan cichlid fish species. Mol. Ecol. 22:3933–48.

O'Quin K.E., Hofmann C.M., Hofmann H.A., Carleton K.L. 2010. Parallel Evolution of
opsin gene expression in African cichlid fishes. Mol. Biol. Evol. 27:2839–2854.

Oaks J.R., Sukumaran J., Esselstyn J. a, Linkem C.W., Siler C.D., Holder M.T., Brown R.M.
2013. Evidence for climate-driven diversification? A caution for interpreting ABC
inferences of simultaneous historical events. Evolution (N. Y). 67:991–1010.

Papadopoulou A., Knowles L.L. 2015. Genomic tests of the species-pump hypothesis: Recent
island connectivity cycles drive population divergence but not speciation in Caribbean
crickets across the Virgin Islands. Evolution (N. Y). 69:1501–1517.

Pearse W.D., Purvis A. 2013. phyloGenerator: An automated phylogeny generation tool for
ecologists. Methods Ecol. Evol. 4:692–698.

Pybus O., Harvey P. 2000. Testing macro–evolutionary models using incomplete molecular
phylogenies. Proc. R. Soc. B Biol. Sci. 267:2267–72.

Rico C., Turner G.F. 2002. Extreme microallopatric divergence in a cichlid species from
Lake Malawi. Mol. Ecol. 11:1585–90.

Rossiter A. 1995. The cichlid fish assemblages of Lake Tanganyika: ecology, behaviour and
evolution of its species flocks. Adv. Ecol. Res.

Salzburger W., Baric S., Sturmbauer C. 2002. Speciation via introgressive hybridization in
East African cichlids? Mol. Ecol. 11:619–25.

Santos M.E., Braasch I., Boileau N., Meyer B.S., Sauteur L., Böhne A., Belting H.-G.,
Affolter M., Salzburger W. 2014. The evolution of cichlid fish egg-spots is linked with a

cis-regulatory change. Nat. Commun. 5:5149.

Schelly R., Salzburger W., Koblmüller S., Duftner N., Sturmbauer C. 2006. Phylogenetic
relationships of the lamprologine cichlid genus Lepidiolamprologus (Teleostei:
Perciformes) based on mitochondrial and nuclear sequences, suggesting introgressive
hybridization. Mol. Phylogenet. Evol. 38:426–438.

Schelly R., Takahashi T., Bills R., Hori M. 2007. The first case of aggressive mimicry among
lamprologines in a new species of Lepidiolamprologus (Perciformes: Cichlidae) from
Lake Tanganyika. Zootaxa. 49:39–49.

Schwarzer J., Misof B., Tautz D., Schliewen U.K. 2009. The root of the East African cichlid
radiations. BMC Evol. Biol. 9:186.

Schweiger O., Klotz S., Durka W., Kühn I. 2008. A comparative test of phylogenetic
diversity indices. Oecologia. 157:485–95.

Sedano R.E., Burns K.J. 2010. Are the Northern Andes a species pump for Neotropical birds?
Phylogenetics and biogeography of a clade of Neotropical tanagers (Aves: Thraupini). J.
Biogeogr. 37:325–343.

Seehausen O. 2000. Explosive speciation rates and unusual species richness in
haplochromine cichlid fishes: effects of sexual selection. Adv. Ecol. Res. 31:237–274.

Seehausen O. 2006. African cichlid fish: a model system in adaptive radiation research. Proc.
R. Soc. B Biol. Sci. 273:1987–98.

Sefc K.M., Mattersdorfer K., Ziegelbecker A., Neuhüttler N., Steiner O., Goessler W.,
Koblmüller S. 2017. Shifting barriers and phenotypic diversification by hybridisation.
Ecol. Lett.

Shirai K., Inomata N., Mizoiri S., Aibara M., Terai Y., Okada N., Tachida H. 2014. High
prevalence of non-synonymous substitutions in mtDNA of cichlid fishes from Lake
Victoria. Gene. 552:239–245.

Siler C.D., Oaks J.R., Esselstyn J.A., Diesmos A.C., Brown R.M. 2010. Phylogeny and
biogeography of Philippine bent-toed geckos (Gekkonidae: Cyrtodactylus) contradict a
prevailing model of Pleistocene diversification. Mol. Phylogenet. Evol. 55:699–710.

Spady T.C., Seehausen O., Loew E.R., Jordan R.C., Kocher T.D., Carleton K.L. 2005.
Adaptive molecular evolution in the opsin genes of rapidly speciating cichlid species.
Mol. Biol. Evol. 22:1412–1422.

Sparks J.S., Smith W.L.W.L. 2004. Phylogeny and biogeography of cichlid fishes (Teleostei:
Perciformes: Cichlidae). Cladistics. 20:501–517.

Sturmbauer C., Baric S., Salzburger W., Rüber L., Verheyen E. 2001. Lake level fluctuations
synchronize genetic divergences of cichlid fishes in African lakes. Mol. Biol. Evol.
18:144–54.

Sturmbauer C., Salzburger W., Duftner N., Schelly R., Koblmüller S. 2010. Evolutionary
history of the Lake Tanganyika cichlid tribe Lamprologini (Teleostei: Perciformes)
derived from mitochondrial and nuclear DNA data. Mol. Phylogenet. Evol. 57:266–84.

Sugawara T., Terai Y., Okada N. 2002. Natural Selection of the Rhodopsin Gene During the
Adaptive Radiation of East African Great Lakes Cichlid Fishes. Mol. Biol. Evol.
19:1807–1811.

Thorpe R.S., Surget-Groba Y., Johansson H. 2008. The relative importance of ecology and
geographic isolation for speciation in anoles. Philos. Trans. R. Soc. Lond. B. Biol. Sci.
363:3071–81.

Toni T., Stumpf M.P.H. 2010. Simulation-based model selection for dynamical systems in
systems and population biology. Bioinformatics. 26:104–110.

Toni T., Welch D., Strelkowa N., Ipsen A., Stumpf M.P.H. 2009. Approximate Bayesian
computation scheme for parameter inference and model selection in dynamical systems.
J. R. Soc. Interface. 6:187–202.

Turner G.F., Seehausen O., Knight M.E., Allender C.J., Robinson R. 2001. How many
species of cichlid fishes are there in African lakes? Mol. Ecol. 10:793–806.

Vaidya G., Lohman D.J., Meier R. 2011. SequenceMatrix: Concatenation software for the
fast assembly of multi-gene datasets with character set and codon information.
Cladistics. 27:171–180.

Verheyen E., Rüber L., Snoeks J., Meyer A. 1996. Mitochondrial phylogeography of rock-
dwelling cichlid fishes reveals evolutionary influence of historical lake level fluctuations
of Lake Tanganyika, Africa. Philos. Trans. R. Soc. Lond. B. Biol. Sci. 351:797–805.

Wagner C.E., Harmon L.J., Seehausen O. 2012. Ecological opportunity and sexual selection
together predict adaptive radiation. Nature. 487:366–9.

Wagner C.E., Harmon L.J., Seehausen O. 2014. Cichlid species-area relationships are shaped
by adaptive radiations that scale with area. Ecol. Lett. 17:583–592.

Wagner C.E., McIntyre P.B., Buels K.S., Gilbert D.M., Michel E. 2009. Diet predicts
intestine length in Lake Tanganyika's cichlid fishes. Funct. Ecol. 23:1122–1131.

Weir J.T. 2006. Divergent Timing and Patterns of Species Accumulation in Lowland and
Highland Neotropical Birds. Evolution (N. Y). 60:842–855.

Weiss J.D., Cotterill F.P.D., Schliewen U.K. 2015. Lake Tanganyika — A ' Melting Pot ' of
Ancient and Young Cichlid Lineages (Teleostei : Cichlidae)? :1–29.

Winkelmann K., Rüber L., Genner M.J. 2016. Lake level fluctuations and divergence of
cichlid fish ecomorphs in Lake Tanganyika. Hydrobiologia.:1–14.

1001

1002