Constitutive Immune Activity Promotes Tumorigenesis in *Drosophila* Intestinal Progenitor

Cells.

Kristina Petkau, Meghan Ferguson, Silvia Guntermann, Edan Foley*.

Department of Medical Microbiology and Immunology

Institute of Virology

University of Alberta

Edmonton, Alberta

Canada.

*Corresponding author.

Email: efoley@ualberta.ca

SUMMARY

Gut innate immune defenses contain bacterial populations and protect the host interior from

invasive microbes. Although excess intestinal immune activity frequently promotes inflammatory

illnesses, we know little about the consequences of chronic innate immune activity exclusively in

endodermal gut cells of an otherwise normal animal. To address this question, we generated a

transgenic line that allows us to activate inflammatory signals in adult fly intestinal progenitor

cells. We found that constitutive immune activity in intestinal progenitors disrupts expression of

homeostatic regulators such as Notch signal transduction pathway components and induces

hyperplasia throughout the gut. Consistent with these observations, we found that persistent

immune signaling interferes with progenitor cell differentiation and exacerbates the formation

of Notch-dependent intestinal tumors. These findings uncover a novel link between constitutive

immune activity and tumorigenesis in intestinal stem cells.

INTRODUCTION

The endoderm is at the heart of an ancient, intimate, and continuous relationship between multicellular organisms and the microbial world. Gut commensals negotiate a semi-stable existence within host niches, partially through provision of factors that influence nutrition, development, and immunity in the host (Cho and Blaser, 2012). For their part, hosts invest energy and resources in the containment of intestinal microbes (Hooper et al., 2012). Physical barriers such as the peritrophic matrix of insects, or dense mucosal layers of mammals, keep microbes at an adequate distance from gut cells. Additionally, host-derived reactive oxygen and nitrogen species destroy invading microbes, while antimicrobial effectors block microbial dissemination. The consequences of failed antibacterial defenses are often extreme, and occasionally deadly for the host. For example, dysbiotic microbial communities contribute to traumatic inflammatory bowel diseases in humans (Wlodarska et al., 2015). In recent years, we have made considerable advances in understanding relationships between host immune responses and gut microbiota (Honda and Littman, 2016). Germline-encoded innate immune systems provide critical defenses in the endoderm, a primary site of contact between multicellular organisms and their immediate microbial environment. Thus, characterization of innate immune regulation in endodermal tissues is essential for a full appreciation of the mechanistic basis for the restraint of gut microbes.

The fruit fly, *Drosophila melanogaster*, is widely used for the characterization of intestinal development and function (Buchon et al., 2013, Jiang and Edgar, 2012, Lemaitre and Miguel-Aliaga, 2013). For example, *Drosophila* is an excellent model for the exploration of Notch-mediated regulation of gut development (Guo and Ohlstein, 2015, Micchelli and Perrimon, 2006,

Ohlstein and Spradling, 2006, Ohlstein and Spradling, 2007), and intestinal tumorigenesis (Patel

and Edgar, 2014, Biteau and Jasper, 2011, Marianes and Spradling, 2013, Apidianakis et al., 2009).

In the fly midgut, basal intestinal stem cells (ISC) divide to generate a bipotent transient cell type,

the enteroblast (EB). Delta-Notch signals between ISC-EB progenitor pairs determine the

developmental fate of EBs (Takashima et al., 2011, Perdigoto et al., 2011). High levels of Notch

activity in enteroblasts lead to their differentiation as large, polyploid, absorptive enterocytes

(EC). Lower levels of Notch activity result in the differentiation of enteroblasts as smaller, diploid,

secretory enteroendocrine cells (EE). Notch-dependent control of progenitor cell differentiation

is conserved across vast evolutionary distances, with similar requirements in species as diverse

as fish and rodents (Fre et al., 2005, Stanger et al., 2005, van Es et al., 2005, Crosnier et al., 2005),

and interruptions to Notch signaling lead to intestinal tumor formation in several models

(Kazanjian and Shroyer, 2011, Peignon et al., 2011).

Current data implicate the immune deficiency (IMD) pathway as a central regulator of

antibacterial defenses in the fly gut (Buchon et al., 2009, Lhocine et al., 2008). IMD shares

numerous features with the mammalian tumor necrosis factor (TNF) cascade (Buchon et al.,

2014). In the fly, detection of bacterial peptidoglycan results in the proteolytic removal of thirty

N-terminal amino acids from the Imd adaptor protein by the caspase Dredd (Paquette et al.,

2010). Cleaved Imd associates with the Inhibitor of Apoptosis 2 (IAP2) and Fas-Associated Death

Domain (FADD) orthologs (Guntermann and Foley, 2011, Paquette et al., 2010) to activate the

NF-kB transcription factor Relish (Dushay et al., 1996). Active Relish relocates to the nucleus and

initiates transcription of a broad spectrum of antimicrobial peptides. More recent work showed

that intestinal IMD activity also controls the expression of genes involved in developmental and

metabolic processes (Broderick et al., 2014, Erkosar et al., 2014). These studies hint at novel,

unstudied biological roles for IMD that extend beyond the extermination of unwanted bacteria.

Here, we characterized the consequences of persistent IMD activity in midgut progenitors.

Midgut progenitor cells extend projections into the intestinal lumen (Ohlstein and Spradling,

2006), and express IMD pathway genes at levels comparable to mature epithelial cells (Dutta et

al., 2015). In addition, enteric challenges with bacterial pathogens induce classic IMD-response

genes in progenitor cells (Dutta et al., 2015, Fink et al., 2016), confirming progenitor cells as an

immune-competent compartment. However, the effects of IMD activity on progenitor cells is

unclear. To address this, we engineered a novel IMD variant that permits temporal and cell-

specific activation of IMD. We showed that constitutive immune signaling altered the expression

of Notch pathway regulators, disturbed EE differentiation, and caused hyperplasia throughout

the posterior midgut. Consistent with interactions between IMD and Notch, we found that

constitutive IMD activity greatly exacerbated the formation of Notch-dependent tumors. Given

the involvement of inflammation and Notch signaling in vertebrate intestinal tumor

development, we believe our findings may be relevant to studies that explore relationships

between intestinal inflammation and tumorigenesis.

RESULTS

Constitutive IMD Pathway Activation in Intestinal Stem Cells.

To characterize the effects of constitutive immune signaling in the intestine, we generated a

transgenic *Drosophila* line that allows cell-restricted activation of the IMD pathway. Specifically,

we engineered a truncated IMD protein that uses an internal ATG (at residue 78) as a start codon.

We refer to this truncated protein as ImdCA. ImdCA lacks inhibitory N-terminal amino acids

(Figure 1A), but retains the ability to interact with the FADD adaptor protein (Figure 1B). We

generated transgenic fly lines that permit temperature-dependent expression of ImdCA in fat

tissue (*cgGAL4* ; *GAL80^{ts}*/*UASimdCA* (abbreviated as *cg^{ts}>CA*)), or ISC/EB progenitor cell pairs

(*esgGAL4*, *GAL80^{ts}*, *UASGFP* ; *UASimdCA* (abbreviated as *esg^{ts}>CA*)). Incubation of either

genotype at 29°C, the restrictive temperature for *GAL80^{ts}*, results in tissue-specific expression of

ImdCA in fat body or intestinal progenitor cells, respectively. In the fat body, the principle site of

humoral immunity, ImdCA caused expression of the IMD-responsive antimicrobial peptides *dpt*

and *att* (Figure 1C). The induction of *dpt* and *att* proceeded via a classical IMD response, as null

mutations in *dredd* prevented ImdCA-mediated antimicrobial peptide expression (Figure 1D).

ImdCA is equally effective at inducing antimicrobial expression in midgut progenitor cells (Figure

1E), and expression of ImdCA provided protective benefits against lethal challenges with the

gastrointestinal pathogen *Vibrio cholerae* (Figure 1F). These data show that we have established

a novel fly line that permits tissue-specific activation of IMD responses in *Drosophila*.

FIGURE 1

Figure 1. A truncated Imd protein activates innate immune responses. (A) Schematic

representation of C-terminally myc-tagged Imd and ImdCA. Iap Binding Motif (IBM) and death

domain (DD) are indicated. Western blots show lysates of untransfected S2 cells (lane 1) S2 cells

transfected with expression constructs for Imd-6myc (lane 2), and ImdCA-6myc (lane 3). Numbers

indicated molecular weights in kDa. The membrane was probed with Tubulin as a loading control.

(B) S2 cells were transfected with the indicated expression constructs and input samples (blots

labeled INPUT) were probed for the indicated antigens. Anti-HA immunoprecipitates (blots

labeled IP:HA) were probed with the indicated antibodies. **(C-E)** Quantification of the relative

expression levels of *dpt* and *att* in flies raised at 29°C for the indicated time. Survival curves of

esg^{ts}>CA and *esg^{ts}* flies raised at 29°C for two days, and then challenged with *Vibrio cholerae*.

A Balance of Microbial Cues and Host Immune Status Determines Gene Expression Patterns in the

Adult Midgut.

Intestinal progenitor cells express IMD pathway components at similar levels to epithelial cell

(Figure 5A), and display classical IMD-responses to microbial challenges (Dutta et al., 2015). To

characterize the effects of ImdCA expression in intestinal progenitors, we prepared

transcriptional profiles of the intestines of *esg^{ts}* and *esg^{ts}>CA* adult flies that we raised at 18°C for

ten days, and shifted to 29°C for two days. We examined *esg^{ts}* and *esg^{ts}>CA* flies raised under

conventional or germ-free conditions in these studies. This strategy allowed us to make pairwise

comparisons of the effects of commensal bacteria and ImdCA on the control of host transcription

(Figure 2A-D).

FIGURE 2

 Figure 2. ImdCA and the microbiome interact to control gut transcription. A-B: Graphic

representation of biological functions dysregulated in $esg^{ts}>CA$ flies relative to esg^{ts} flies raised

under conventional (A) or germ-free conditions (B). **C-D:** Graphic representation of biological

functions dysregulated in conventional flies relative to germ-free flies in the absence (C) or

presence (D) of ImdCA expression. For A-D, numbers indicated total numbers of dysregulated

genes. Genes with unknown function were not included in pie charts. **(E-F)** Heatmap of

significance scores for gene ontology terms (minimum 5 genes in each term) that were

significantly dysregulated in $esg^{ts}>CA$ flies relative to esg^{ts} flies (E), or conventionally-reared

relative to germ-free flies (F). In E, GO terms are further subdivided into terms that were

dysregulated in conventionally-reared or germ-free flies. In F, GO terms are subdivided into terms

that were dysregulated in esg^{ts} flies and $esg^{ts}>CA$ flies.

Similar to earlier studies (Broderick et al., 2014, Erkosar et al., 2014), we found that the

microbiome regulates the expression a large cohort of host genes, with a particularly notable

effect on genes involved in metabolic processes (Figure 2A-D, Figure 3). As expected, ImdCA

altered the expression of genes with known roles in innate defenses. However, we also noticed

effects of ImdCA on pathways as diverse as metabolism, transportation and stress responses

(Figures 2, 3).

Comparisons between the individual expression profiles uncovered substantial effects of the

microbiome on ImdCA-regulated transcription. For example, only 35 host genes are induced in

both conventionally-reared and germ-free animals in response to ImdCA (Figure 3E). In contrast,

146 transcripts are only induced in the presence of an intact microbiome, and an additional 21

transcripts are only induced in the absence of a microbiome (Figure 3E). Likewise, microbiome-

response genes are heavily influenced by the extent of immune activity in the gut. For example,

a unique set of 129 genes responds to presence of the microbiome only upon induction of ImdCA

(Figure 3F). GO term analysis showed that the microbiome and immune activity synergistically

influence functional outputs in the gut. For example, ImdCA only promotes protein glycosylation

in the presence of a microbiome (Figure 2E), and the gut microbiome induces tricarboxylic acid

cycle genes only when accompanied by intestinal inflammation (Figure 2F). Combined, these data

show that gut transcriptional outputs reflect the extent of immune activity and microbial

presence in the gut.

Figure 3: A-D: Relative expression of totM (A), pirk (B), pgrp-sc1 (C) and cec (D) in the dissected

intestines of conventionally-reared or germ-free *esg^ts* flies or *esg^ts>CA* flies. For each assay,

expression levels were reported relative to those observed in germ-free *esg^ts* flies. Comparisons

were performed with a Sidak's multiple comparison test. Values show the Sidak's t value for the

respective tests. **E:** Number of genes and GO terms that were up or downregulated in *esg^ts>CA*

flies relative to *esg^ts* flies raised under conventional (CR) or germ-free (GF) conditions. **F:** Number

of genes and GO terms that were up or downregulated in conventionally-reared flies relative to

germ-free flies of the indicated genotypes.

Given the effects of ImdCA on antimicrobial peptide expression, we assumed that ImdCA will

affect composition of the host microbiome. To test this, we treated freshly eclosed *esg^ts* and

esg^ts>CA flies with a regime of antibiotics for six days to eliminate the endogenous microbiome.

We then fed all flies a homogenate prepared from our lab wild-type *Drosophila* strain, and

passaged flies to fresh food twice over seven days to facilitate stable association between

microbes in the homogenate and the recipient flies. After seven days, we shifted recipients to

29°C for an additional seven days to activate the IMD pathway in *esg^ts>CA* flies. We dissected the

intestines of the different populations and used deep-sequencing of 16S DNA to identify the

bacterial communities present in the respective guts (Figure 4). As expected, the bacterial

microbiome of *esg^ts* flies was barely distinguishable from the homogenate. Both microbiomes

contained a limited number of OTUs characterized primarily by *Acetobacter* and *Lactobacilli*

(Figure 4). Contrary to our initial expectation, we did not observe a significant impact of ImdCA

on microbiome composition. In fact, the bacterial communities of the original homogenate, the

esg^{ts} flies, and the *esg^{ts}>CA* flies were barely distinguishable (Figure 4), suggesting that acute

activation of IMD does not significantly alter microbial populations in the gut.

Kruskall-Wallis test: H=0.1 df=2 P=0.9512

Figure 4. Induced expression of ImdCA in intestinal progenitor cells does not have a substantial

impact on gut microbiome composition. Heat map analysis of microbial original taxonomic unit

representation in dissected midguts of the indicated genotypes. All flies were raised under

identical conditions, and fed the same homogenate at the same time. Kruskall-Wallis test results

for degree of difference between the different samples.

Constitutive IMD Pathway Activation Causes Dysplasia in Intestinal Progenitor Cells.

In contrast to the minimal effects on bacterial microbiome composition, we noticed that

ImdCA affected expression of regulators of intestinal homeostasis. This included elements of

Wnt, Ras, Insulin, JNK and JAK/STAT pathways (Figure 5B). This led us to ask if ImdCA impacts

intestinal morphology. To address this question, we examined the posterior midguts of two,

eleven and twenty-six day old *esg^{ts}>CA* and control *esg^{ts}* flies. In these lines, the *esg^{ts}* regulatory

system marks all progenitors with GFP. We found that acute activation of the IMD pathway had

no effect on midgut architecture. The posterior midguts of two-day old *esg^{ts}* and *esg^{ts}>CA* flies

raised at 29°C were essentially indistinguishable (Figure 6A). In both cases, anti-Armadillo (beta-

catenin ortholog) immunofluorescence showed a regular arrangement of large, polyploid ECs,

that were interspersed with small, Prospero positive EE cells. Likewise, the posterior midguts of

both genotypes contained regularly spaced nuclei and evenly distributed GFP positive progenitor

cells (Figure 6A).

A

GENE	ISC	EB	EC	EE	VM
PGRP-LC	6.13	7.96	6.432	1.365	4.263
IMD	3.534	4.249	7.41	7.473	7.55
IAP2	10.171	16.595	13.329	15.8	12.292
DREDD	1.684	1.712	1.694	1.506	1.747
FADD	6.649	11.527	11.19	11.997	3.282
TAB2	2.413	2.222	4.253	6.007	3.373
TAK1	4.121	7.004	7.89	5.929	9.594
KEY	19.999	52.248	42.098	19.39	39.636
IRD5	0.97	0.765	1.076	0.782	0.906
REL	40.397	52.647	39.985	14.71	42.481

Expression value (rpkm) per cell type

B

Gene	Function	Fold Change (*esgCA* V *esg*)	ANOVA p-value
MESK2	Ras signaling	-2.15	0.000349
gfzg	Ras signaling	-1.84	0.000116
cdi	Ras signaling	1.59	0.001404
Plzf	Ras signaling	-1.66	0.025787
CG42684	Ras signaling	2	0.000475
dock	Insulin signaling	-1.56	0.016287
srl	Insulin signaling	1.59	0.012109
sug	Insulin signaling	-1.68	0.025726
egr	JNK signaling	1.68	0.001723
Traf4	JNK signaling	1.64	0.000984
nmo	Wnt signaling	-1.98	0.000068
ebd1	Wnt signaling	1.53	0.008942
sfl	Wnt signaling	1.96	0.002488
Prosap	Wnt signaling	-1.53	0.002859
SOCS36E	JAK/STAT	-1.84	0.038907

C

Gene	Function	Fold Change (*esgCA* V *esg*)	ANOVA p-value
mahe	Notch signaling	1.54	0.000941
neur	Notch signaling	-1.55	0.001857
Nle	Notch signaling	1.63	0.010107
Rala	Notch signaling	2.07	0.000403
stv	Lateral Inhibition	-1.61	0.005874
Cam	Lateral Inhibition	-1.54	0.007552
CG13827	Lateral Inhibition	-1.74	0.003491
CG16713	Lateral Inhibition	-1.86	0.000181
CG5953	Lateral Inhibition	-1.81	0.002646
CG7631	Lateral Inhibition	-2.19	0.000129
CG9194	Lateral Inhibition	-1.89	0.001998

Figure 5. A: Expression of IMD pathway components in intestinal cell types. Expression data taken

from (Dutta et al., 2015). **B-C:** Homeostatic regulators (B) and Notch pathway components (C)

that were dysregulated in esg^{ts}>CA flies relative to esg^{ts} flies.

By day 11, the posterior midguts of esg^{ts}>CA flies appeared moderately dysplastic relative to

esg^{ts} flies. By 26 days, the posterior midguts of control esg^{ts} flies displayed classical hallmarks of

age-dependent dysplasia such as a disorganized epithelium, uneven arrangements of nuclei, and

irregular patterns of GFP expression (Figure 6A). We noticed that midgut architecture was further

disrupted in 26 day-old esg^{ts}>CA flies. In this case, we detected a near complete breakdown of

epithelial organization (Figure 6A), an increase in cell density, and an increase in GFP positive

cells. The hyperplastic phenotype of 26 day old esg^{ts}>CA flies is not a consequence of accelerated

physiological aging, as esg^{ts} and esg^{ts}>CA flies have indistinguishable lifespans (Figure 6I).

Given the hyperplasia and altered expression of homeostatic regulators in older esg^{ts}>CA

flies, we reasoned that ImdCA affects proliferation rates in intestinal progenitor cells. To test this

hypothesis, we used anti-phosphorylated histone H3 immunofluorescence to count mitotic cells

in the intestines of fourteen and twenty-four day-old esg^{ts} and esg^{ts}>CA files. At both ages, we

detected a significant increase in mitotic events in esg^{ts}>CA intestines relative to age-matched

esg^{ts} controls (Figure 6D). These data show that persistent expression of ImdCA in adult midgut

progenitor cells disrupts expression of homeostatic regulators, increases cell proliferation, and

promotes tissue hyperplasia.

Figure 6. Intestinal hyperplasia in the posterior midguts of *esg^ts^>CA* flies. (A-C)

Immunofluorescence imaging of the posterior midguts of *esg^ts^>CA* and *esg^ts^* flies (A); *Dl^ts^>CA* and

*Dl^ts^* flies (B); and *myo1a^ts^>CA* and *myo1a^ts^* flies (C) raised at 29°C for 2, 11 or 26 days as indicated.

Each midgut was stained with Armadillo (to indicate cell borders), and Prospero to reveal

enteroendocrine cells; Hoechst to reveal nuclei; and GFP was visualized to reveal progenitor cells

(A), stem cells (B) and enterocytes (C). All images were taken at 40X magnification and scale bars

show 25 μm. **(D-F)** Number of mitotic cells in the intestines of flies of the indicated genotypes

raised at 29°C for 14 and 24 days. **G:** Visualization of mitotic clones in wild type flies (upper row)

or mitotic clones that express *imdCA* (lower row). All images were taken at 60X magnification and

scale bars show 15 μm **H:** Percentage of large, enterocyte-type cells in wild type clones and *imdCA*

positive clones. Comparisons were performed with a student's t test. **(I)** Survival curves for 150

esg^{ts}>CA and 150 esg^{ts} flies raised at 29°C.

We did not see a similar phenotype when we used a Dl^{ts} line (*UASGFP ; DlGAL4, GAL80^{ts}*) to

express ImdCA in GFP-marked stem cells. In this case, midgut morphology and proliferation rates

were indistinguishable between Dl^{ts}>CA and control Dl^{ts}/+ flies at all times (Figure 6B and E).

Expression of ImdCA in GFP-marked intestinal enterocytes with the $Myo1A^{ts}$ (*UASGFP ;*

Myo1aGAL4, GAL80^{ts}) line led to a mild increase in midgut proliferation at d14 that did not persist

at later times (Figure 6C and F). Thus, expression of ImdCA in progenitor cells led to a marked

hyperplasia throught the posterior midgut. As the Dl^{ts} line, fails to induce hyperplasia our data

suggest that activation of IMD either in progenitor pairs, or enteroblasts alone leads to midgut

hyperplasia.

Persistent IMD Signals Disrupt Notch Activity.

Among homeostatic regulators, ImdCA had a pronounced effect on expression of Notch

pathway components (Figure 5B). This observation matches an earlier report of deregulated

expression of Notch pathway genes in *imd* mutant flies (Broderick et al., 2014), and raises the

possibility that the IMD pathway influences Notch activity in the adult. To test if ImdCA affects

progenitor cell differentiation, we used MARCM analysis to identify individual mitoses and

subsequent differentiation events in adult posterior midguts. As expected, mitotic clones in

conventional flies contained a mix of smaller cells (most likely progenitors and EE), and larger,

polyploid ECs (Figure 6G). In contrast, we rarely detected large, polyploid cells in mitotic clones

that express ImdCA. Instead, ImdCA clones typically contained aggregates of small cells (Figure

6G). These observations suggest that persistent expression of ImdCA in progenitor cells affects

the differentiation of EBs into ECs.

To directly test the possibility that ImdCA disturbs EB differentiation, we determined the

percentage of EE cells in the intestines of twenty-four days old esg^{ts} and esg^{ts}>CA flies. As

expected, the posterior midguts of esg^{ts}>CA flies contained greater numbers of cells per area

than those observed in age-matched, control esg^{ts} flies (Figure 7A-C). However, we also found

that a significantly greater percentage of midgut cells expressed the EE marker Prospero in

esg^{ts}>CA than in esg^{ts} midguts (Figure 7A, B, D). These data show that persistent expression of

ImdCA disrupts the developmental trajectory of progenitor cells, and leads to the differentiation

of supernumerary EE.

IMD Pathway Activity Promotes the Development of Notch-Mediated Intestinal Tumors.

As bacterial challenges enhance *Notch*-deficient tumor formation (Apidianakis et al., 2009,

Jiang et al., 2009), and ImdCA influences *Notch*-dependent events, we asked what effects

persistent immune signals have on models of *Notch*-deficient tumorigenesis in the posterior

midgut. To answer this question, we used a validated, inducible RNAi line that blocks *Notch*

signaling in GFP-marked adult progenitors (esg^{ts}>N^{RNAi}). As anticipated, the posterior midguts of

esg^{ts}>N^{RNAi} flies contained GFP-positive tumors composed of small, irregularly spaced nuclei

within eight days of Notch depletion (Figure 7E, F). Many tumors also contained patches of

Prospero-positive enteroendocrine cells (Figure 7F). The ability of esg^{ts}>N^{RNAi} to promote

tumorigenesis appears to require extrinsic inputs from the microbiome, as removal of the gut

microbiome greatly reduced the incidence of tumor formation in esg^{ts}>N^{RNAi} posterior midguts

(Figure 7G, H). This observation matches an earlier report that growth of *Notch* loss-of-function

tumors requires stress-induced mitogenic cues (Patel et al., 2015).

Figure 7. ImdCA promotes tumorigenesis in Notch-deficient progenitor cells. (A-B) Midguts of 20d old esg^{ts} (A) and esg^{ts}>CA (B) flies stained for DNA and Prospero as indicated. Images were false-colored and merged with DNA in blue and Prospero in red. Images were taken at 40X magnification and scale bars show 25 μm. **C:** Automated quantification of total nuclei in a given field for flies of both genotypes. **D:** Total percentage of Prospero positive cells in the same fields. For C and D comparisons were performed with a student's t test. **E-L:** Visualization of tumorigenesis in 8d old esg^{ts}>N^{RNAi} and esg^{ts}>N^{RNAi}; CA flies. Posterior midguts were stained for Prospero and DNA in conventionally-reared esg^{ts}>N^{RNAi}; flies (E-F); germ-free esg^{ts}>N^{RNAi} flies (G-H); conventionally-reared esg^{ts}>N^{RNAi}; CA flies (I-J); and germ-free esg^{ts}>N^{RNAi}; CA flies (K-L). In all cases, progenitor cells were visualized through the expression of GFP. Scale bars show 50μm for 20X magnification images, and 15μm for 60X images. A representative grade 2 tumor is indicated with an arrowhead in F, and representative grade 3 tumors are shown with an asterisk. **M:** Tumor scoring in the posterior midguts of 8d old flies with the indicated genotypes and culture conditions (n=26) for each condition. Tumor grade was compared between individual treatments using one-way ANOVA, with Tukey's correction. *** indicates P<0.0001.

Whereas elimination of the microbiome blocked tumorigenesis in esg^{ts}>N^{RNAi} flies, co-expression of ImdCA with N^{RNAi} (esg^{ts}>N^{RNAi}; CA) exacerbated the formation of midgut tumors. We invariably detected large patches of GFP and Prospero positive cells in esg^{ts}>N^{RNAi}; CA posterior midguts, with few large cells characteristic of mature ECs (Figure 7 I, J). To determine if ImdCA alone provides sufficient stress signals to promote tumorigenesis in this *Notch* loss-of-function model, we examined the posterior midguts of germ-free esg^{ts}>N^{RNAi}; CA flies. We found

that nearly all midguts of germ-free esg^{ts}>CA, N^{RNAi} flies had accumulations of small, GFP-positive

cells, large patches of Prospero positive tumors, and few large polyploid ECs (Figure 7K, L). To

quantify the effects of microbiome elimination on tumorigenesis, we developed a grading system

for tumor scale in the posterior midgut. We defined tumor-free intestines as Grade 1 (e.g. Figure

7G, H), tumors that only contained GFP-positive clusters of cells as Grade 2 (Figure 7F, asterisks),

and tumors that contained clusters of GFP-positive and Prospero-positive cells as Grade 3 (Figure

7F, arrowhead). We used this system to quantify tumor grade in 26 intestines from each

experimental condition. This scale confirmed a significant reduction of tumor formation in germ-

free esg^{ts}>N^{RNAi} flies (Figure 7M), and a significant increase of tumorigenesis in germ-free

esg^{ts}>N^{RNA} ; CA flies (Figure 7M).This phenotype is remarkably similar to one described earlier for

bacterial infection of esg^{ts}>N^{RNAi} flies (Apidianakis et al., 2009, Jiang et al., 2009), and shows that

immune signaling in progenitor cells alone is sufficient to promote *Notch*-dependent tumor

growth in adult midguts.

337 **DISCUSSION**

338

339   Animal genomes encode evolutionarily conserved systems that mitigate the threats posed by

340 potentially harmful microbes. For example, homeostatic resistance systems clear invading

341 microbes, while tolerance mechanisms limit the damage caused by invaders (Medzhitov et al.,

342 2012, Schneider and Ayres, 2008) . In this study, we used the genetically accessible fruit fly to

343 activate immune responses exclusively in intestinal progenitor cells. We focused on progenitors,

344 as they are immune-competent, but comparatively understudied from the perspective of IMD.

345 We showed that persistent immune activity causes intestinal hyperplasia, disrupts progenitor cell

346 differentiation, and fuels Notch-dependent tumorigenesis. These phenotypes occur without

347 significant shift in bacterial populations, consistent with altered disease tolerance in the host.

348 Our studies match reports of relationships between the microbiota, inflammation and colorectal

349 cancer (Belkaid and Hand, 2014, Irrazabal et al., 2014), and uncover a novel requirement for

350 regulated innate immune activity in the overall maintenance of progenitor cell homeostasis in

351 the fly.

352   In *Drosophila*, several homeostatic devices control progenitor cell proliferation and

353 differentiation, often in response to bacterial products (Amcheslavsky et al., 2009, Buchon et al.,

354 2009, Jiang et al., 2009). For example, flies raised in the absence of a microbiome undergo fewer

355 mitoses in the midgut (Buchon et al., 2009). In addition, flies raised in a germ-free environment

356 have atypical numbers of prospero-positive EE cells, a lower density of ECs, and deregulated

357 expression of Notch pathway components (Broderick et al., 2014). It is unclear how much of this

358 response requires an intact IMD pathway, although recent transcriptional work established that

mutations of the IMD pathway impact numerous aspects of intestinal transcription (Broderick et

al., 2014, Erkosar et al., 2014), including Notch pathway components (Broderick et al., 2014). Our

work shows that expression of ImdCA in progenitor cells alone recapitulates key features of those

host responses to gut microbes. For example, of the 253 genes differentially expressed in the

intestines of conventionally-reared flies compared to germ-free flies, 87 are affected by

expression of ImdCA. Additionally, expression of ImdCA in progenitor cells boosts intestinal

mitoses, affects EE cell differentiation, and increases EC density. Combined, these data implicate

the IMD pathway as a novel regulator of progenitor cell proliferation.

As the microbiome induces progenitor cell proliferation, we initially assumed that ImdCA-

dependent hyperplasia was the result of a dysbiosis in the gut caused by immune activity.

However, expression of ImdCA does not influence microbiome composition. These data raise the

possibility that common intestinal microbes are insensitive to IMD activity in the gut, or that

elevated immune signaling activates counter-defenses in commensal populations, thereby

allowing persistence of commensal bacteria. This matches experimental data that commensal

bacteria effectively overcome host innate defenses in other models. For example, many gut-

resident *Firmicutes* and *Bacteroidetes* are not susceptible to antimicrobial peptides (Cullen et al.,

2015), and NOD deficient mice do not display dysbiosis (Robertson et al., 2013, Shanahan et al.,

2014). These observations lead us propose that constitutive activation of IMD acts directly on the

host to induces intestinal hyperplasia. Although out data are more consistent with IMD acting in

enteroblasts to promote hyperplasia, additional experiments are required to rule out a need for

IMD activity in progenitor pairs. We believe that the ability of ImdCA to promote Notch-

dependent tumors is of particular significance in the context of aging. As flies age, clones of *Notch*

mutant cells spontaneously develop in the intestine (Siudeja et al., 2015), and host immune

activity gradually rises (Buchon et al., 2009). Our data suggest that age-dependent increases in

immune activity may contribute to the formation, or growth, of Notch-dependent tumors in the

fly. Indeed, we found that elimination of the microbiome was sufficient to arrest the appearance

of Notch-dependent tumors.

The role of Notch in tumorigenesis is not restricted to flies. Several studies established clear

links between deregulated Notch activity and carcinogenesis in vertebrates (Rizzo et al., 2008,

Shih Ie and Wang, 2007). Colorectal cancer lines and adenocarcinomas express elevated levels of

Notch1 and Notch 2 (Fre et al., 2009, Guilmeau et al., 2010, Reedijk et al., 2008, Sikandar et al.,

2010). In addition, manipulations of Notch pathway activity modify tumorigenesis in mouse

models of colorectal cancer (Fre et al., 2009, Rodilla et al., 2009). The mechanisms that activate

Notch in colon cancer are unclear at present. Given the broad conservation of immune activity

and Notch signaling in the guts of flies and mammals, our work raises the possibility that chronic

inflammation is a long-term activator of Notch-dependent tumorigenesis. The data in this study

present a simple model to precisely describe the relationships between intestinal immune

activity and the formation of Notch-dependent tumors.

EXPERIMENTAL PROCEDURES

***Drosophila* husbandry.**

All flies were raised on standard corn meal medium (Nutri-Fly Bloomington Formulation, Genesse Scientific). Germ-free husbandry conditions are described in the Supplementary methods. For 16S deep-sequencing, we raised freshly eclosed virgin females flies on antibiotic medium for 5 d at 18°C, then switched to sterile antibiotic-free food for 1 d. Next, flies were fed a homogenate prepared from w^{1118} flies for 16 h. Afterwards, flies were raised on sterile antibiotic-free food at 18°C for 7 d. Flies were moved to 29°C for another 7 days, and 10 guts/sample were dissected. We extracted microbial DNA with the UltraClean® Microbial DNA Isolation Kit (MoBio Laboratories Inc.), and amplified 16S DNA with Platinum® PFX Taq (Invitrogen), followed by purification with the QIAquick® PCR Purification Kit (Qiagen). Concentration was measured on the Qubit® 2.0 (Invitrogen) and 1 ng was used for library prep. Libraries were prepared using the Nextera XT DNA Library Preparation Kit (Illumina). We purified libraries using Ampure Beads (Qiagen) and quantified using the Qubit® 2.0 (Invitrogen) and Bioanalyzer 2100 (Agilent). Pooled libraries were loaded on the Miseq (illumina) using the MiSeq Reagent Kit v3 (600-cycle) for sequencing. 16S sequences were assembled using DNASTAR Navigator, and annotated with the greengenes database. For infection studies, freshly eclosed flies were raised for 5 d at 18°C, then switched to 29°C for 2 d. 24 h before infection 100 µl of a *V. cholerae* C6706 glycerol stock was spread on LB agar plates and grown at 29°C. Flies were starved for 2 h before infection. The bacterial lawn was scraped from the plate and mixed into LB broth, then diluted to an OD600 of 0.125, flies were fed 3ml bacterial supernatant on a cotton plug, and dead flies counted every 2-

4 h. To generate mitotic clones, flies were raised at 18°C for 5-6 d, incubated at 37°C for 2 h, and

raised for an additional 8-10 days at 25°C.

Cell culture and molecular biology.

The Imd expression construct used in this study has been described elsewhere (Guntermann and

Foley, 2011). To generate ImdCA, we used pENTR/D-TOPO ImdCA forward and reverse primers

(supplemental material) to amplify a truncated Imd. We then cloned truncated Imd with into the

pENTR/D-TOPO plasmid according to manufacturer's instructions (Invitrogen). We recombined

TOPO-ImdCA with pAWM, or pTW (LR recombination, Invitrogen) to generate ImdCA-6myc and

UAS-ImdCA expression plasmids. Transgenic lines were generated by Bestgene Inc.

Immunoprecipitation and Western blot protocols have been described elsewhere (Guntermann

and Foley, 2011). For immunoprecipitation assays, we incubated cell lysate with mouse anti-HA

(Sigma, 1:500) at 4°C overnight, added protein G-Sepharose beads and incubated for an

additional hour at 4°C. Beads were pelleted by centrifugation at 300 X g for 30 s and washed in

lysis buffer three times. After discarding the supernatant, beads were re-suspended in 2X sample

buffer, and analyzed by Western blot.

Gene expression analysis.

We used TRIZOL to prepare RNA from dissected adult intestines (Guntermann and Foley, 2011).

Microarray studies were performed in triplicate on virgin flies that we raised on regular or

antibiotic-treated food food for 10-11 days at 29°C. We then shifted flies to 29°C for another 2

days, after which guts were dissected for RNA extraction (5 females and 5 males per sample). We

used 100 ng purified RNA to make labeled cRNA for microarrays using the GeneChip® 3' IVT Plus

Reagent Kit (Affymetrix), then fragmented cRNA and then hybridized to the

GeneChip®Drosophila Genome 2.0 Array (Affymetrix). Preliminary analysis was done in the

Transciptome Analysis Console (TAC) software (Affymetrix). We analyzed gene expression data

using FlyMINE (Lyne et al., 2007) and Panther (Thomas et al., 2003). Array data has been

submitted to the NCBI GEO database (accession ID: GSE89445). For qPCR studies, we prepared 1

µg cDNA from purified RNA using qScript cDNA Supermix (Quanta Biosciences, Inc.) and then

performed qPCR using the PerfeCTa® SYBR®Green FastMix (Quanta Biosciences, Inc.). All qPCR

studies were performed in triplicate and relative expression values were calculated using delta

delta Ct calculations. In each case, expression levels were normalized to *actin*.

Immunofluorescence.

The immunofluorescence protocol used in this study has been described elsewhere (Petkau et

al., 2014). All immunofluorescence images were prepared from the posterior midgut of adult flies

at a distance of approximately 50 µM from the hindgut transition. To quantify phospho-histone

H3 positive cells, guts were visualized under the microscope and scanned from posterior midgut

(hindgut transition area) to anterior midgut (crop). To quantify Prospero positive cells, we used

the Columbus software to identify the percentage of Hoechst positive cells that were also

prospero positive. 3-D reconstructions were created with Volocity® 6.3 (Perkin Elmer).

Germ-Free Adult Fly Husbandry.

Larval growth and development is supported by commensal microbes. As our primary goal was to examine

the effects of immune signaling in an adult model of intestinal tumorigenesis, we were concerned that

elimination of the larval microbiome would influence gut development, and possibly confound

interpretation of our data. For that reason, we chose to raise larvae under conventional conditions, and

exclusively remove the bacterial microbiome from adults. To generate germ-free animals, we raised

freshly eclosed adult flies on sterile food containing an antibiotic cocktail (100 µg/ml Ampicillin, 100 µg/ml

Neomycin, 100 µg/ml Metronidazole and 50 µg/ml Vancomycin). This protocol allowed us to raise larvae

in the presence of a conventional microbiome, and restrict our examination of germ-free phenotypes to

the adult life stage. We confirmed bacterial elimination by plating fly homogenates on agar plates

permissive for the growth of *Lactobacilli* and *Actetobacter*.

Primers used in this study.

Gene	Forward Primer	Reverse Primer
pENTR/D-TOPO ImdCA	CACCGCAGCTCCCGTGGACGACAACG	CTAGCTGTTTGTCTTGCGC
16S	AGAGTTTGATCCTGGCTCAG	GGCTACCTTGTTACGACTT
att	AGTCACAACTGGCGGAAC	TGTTGAATAAATTGGCATGG
dpt	ACCGCAGTACCCACTCAATC	ACTTTCCAGCTCGGTTCTGA
actin	TGCCTCATCGCCGACATAA	CACGTCACCAGGGCGTAAT
totM	ACCGGAACATCGACAGCC	CCAGAATCCGCCTTGTGC
pirk	AGAGCACGAGCAGGGTAAATC	TGTTGTTCTCAATGCGGTACTC

pgrp-sc1	AAGCGATCGTCAACTATT	GAGAGCCACTTTGGAAACCA
cec	TGTAAGCTAGTTTATTTCTATGG	GATGAGCCTTTAATGTCC

476 **Antibodies used in this study.**

Antigen	Use	Source
Tubulin.	Western blot (1:1000)	Developmental Studies Hybridoma Bank E7
cMyc	Western blot (1:5000)	Sigma
HA	Western blot (1:4000)	Sigma
JNK	Western blot (1:4000)	Santa Cruz sc-571
Armadillo	Immunofluorescence (1:100)	Developmental Studies Hybridoma Bank N2 7A1
Prospero	Immunofluorescence (1:100)	Developmental Studies Hybridoma Bank MR1A
phospho- H3(Ser10)	Immunofluorescence (1:1000)	Millipore
pgrp-sc1	AAGCGATCGTCAACTATT	GAGAGCCACTTTGGAAACCA
cec	TGTAAGCTAGTTTATTTCTATGG	GATGAGCCTTTAATGTCC

Fly Genotypes

Clonal analysis was performed with flies of the following genotypes: *hsFLP, UASGFP/X* ;

tubGAL80, neoFRT40A/neoFRT40A ; *tubGAL4/+* (wild-type), and *hsFLP, UASGFP/X* ; *tubGAL80,*

neoFRT40A/neoFRT40A ; *tubGAL4/UASimdCA* (ImdCA positive clones).

AUTHOR CONTRIBUTIONS

K.P., M.F., S.G., and E.F. conceived and designed experiments; K.P., M.F., and S.G. performed the

experiments; K.P., M.F., and E.F performed data analysis and wrote the paper.

ACKNOWLEDGEMENTS

Transgenic flies were provided by Bruce Edgar (esg^{ts}), Bruno Lemaitre (*dredd*), the Bloomington

Drosophila stock Center (*cgGAL4* and *GAL80^{ts}*), and the Vienna *Drosophila* Resource Center

(N^{RNAi}). The research was funded by a grant from the Canadian Institutes of Health Research to

EF (MOP 77746). We acknowledge the microscopy support from Dr. Stephen Ogg and the Faculty

of Medicine and Dentistry core imaging service, the Cell Imaging Centre, University of Alberta.

Microarrays were processed at the Alberta Transplant Applied Genomics Center. We are grateful

to Juan Jovel for bioinformatics advice.

REFERENCES

AMCHESLAVSKY, A., JIANG, J. & IP, Y. T. 2009. Tissue damage-induced intestinal stem cell
division in Drosophila. *Cell Stem Cell,* 4**,** 49-61.
APIDIANAKIS, Y., PITSOULI, C., PERRIMON, N. & RAHME, L. 2009. Synergy between bacterial
infection and genetic predisposition in intestinal dysplasia. *Proc Natl Acad Sci U S A,* 106**,**
20883-8.
BELKAID, Y. & HAND, T. W. 2014. Role of the microbiota in immunity and inflammation. *Cell,*
157**,** 121-41.
BITEAU, B. & JASPER, H. 2011. EGF signaling regulates the proliferation of intestinal stem cells in
Drosophila. *Development,* 138**,** 1045-55.
BRODERICK, N. A., BUCHON, N. & LEMAITRE, B. 2014. Microbiota-induced changes in drosophila
melanogaster host gene expression and gut morphology. *MBio,* 5**,** e01117-14.
BUCHON, N., BRODERICK, N. A., CHAKRABARTI, S. & LEMAITRE, B. 2009. Invasive and
indigenous microbiota impact intestinal stem cell activity through multiple pathways in
Drosophila. *Genes Dev,* 23**,** 2333-44.
BUCHON, N., BRODERICK, N. A. & LEMAITRE, B. 2013. Gut homeostasis in a microbial world:
insights from Drosophila melanogaster. *Nat Rev Microbiol,* 11**,** 615-26.
BUCHON, N., SILVERMAN, N. & CHERRY, S. 2014. Immunity in Drosophila melanogaster--from
microbial recognition to whole-organism physiology. *Nat Rev Immunol,* 14**,** 796-810.
CHO, I. & BLASER, M. J. 2012. The human microbiome: at the interface of health and disease.
Nat Rev Genet, 13**,** 260-70.
CROSNIER, C., VARGESSON, N., GSCHMEISSNER, S., ARIZA-MCNAUGHTON, L., MORRISON, A. &
LEWIS, J. 2005. Delta-Notch signalling controls commitment to a secretory fate in the
zebrafish intestine. *Development,* 132**,** 1093-104.
CULLEN, T. W., SCHOFIELD, W. B., BARRY, N. A., PUTNAM, E. E., RUNDELL, E. A., TRENT, M. S.,
DEGNAN, P. H., BOOTH, C. J., YU, H. & GOODMAN, A. L. 2015. Gut microbiota.
Antimicrobial peptide resistance mediates resilience of prominent gut commensals
during inflammation. *Science,* 347**,** 170-5.
DUSHAY, M. S., ASLING, B. & HULTMARK, D. 1996. Origins of immunity: Relish, a compound Rel-
like gene in the antibacterial defense of Drosophila. *Proc Natl Acad Sci U S A,* 93**,** 10343-
7.
DUTTA, D., DOBSON, A. J., HOUTZ, P. L., GLASSER, C., REVAH, J., KORZELIUS, J., PATEL, P. H.,
EDGAR, B. A. & BUCHON, N. 2015. Regional Cell-Specific Transcriptome Mapping Reveals
Regulatory Complexity in the Adult Drosophila Midgut. *Cell Rep,* 12**,** 346-58.
ERKOSAR, B., DEFAYE, A., BOZONNET, N., PUTHIER, D., ROYET, J. & LEULIER, F. 2014. Drosophila
microbiota modulates host metabolic gene expression via IMD/NF-kappaB signaling.
PLoS One, 9**,** e94729.
FINK, C., HOFFMANN, J., KNOP, M., LI, Y., ISERMANN, K. & ROEDER, T. 2016. Intestinal FoxO
signaling is required to survive oral infection in Drosophila. *Mucosal Immunol,* 9**,** 927-36.
FRE, S., HUYGHE, M., MOURIKIS, P., ROBINE, S., LOUVARD, D. & ARTAVANIS-TSAKONAS, S. 2005.
Notch signals control the fate of immature progenitor cells in the intestine. *Nature,* 435**,**
964-8.

FRE, S., PALLAVI, S. K., HUYGHE, M., LAE, M., JANSSEN, K. P., ROBINE, S., ARTAVANIS-TSAKONAS,
S. & LOUVARD, D. 2009. Notch and Wnt signals cooperatively control cell proliferation
and tumorigenesis in the intestine. *Proc Natl Acad Sci U S A,* 106**,** 6309-14.
GUILMEAU, S., FLANDEZ, M., MARIADASON, J. M. & AUGENLICHT, L. H. 2010. Heterogeneity of
Jagged1 expression in human and mouse intestinal tumors: implications for targeting
Notch signaling. *Oncogene,* 29**,** 992-1002.
GUNTERMANN, S. & FOLEY, E. 2011. The protein Dredd is an essential component of the c-Jun
N-terminal kinase pathway in the Drosophila immune response. *J Biol Chem,* 286**,**
30284-94.
GUO, Z. & OHLSTEIN, B. 2015. Stem cell regulation. Bidirectional Notch signaling regulates
Drosophila intestinal stem cell multipotency. *Science,* 350.
HONDA, K. & LITTMAN, D. R. 2016. The microbiota in adaptive immune homeostasis and
disease. *Nature,* 535**,** 75-84.
HOOPER, L. V., LITTMAN, D. R. & MACPHERSON, A. J. 2012. Interactions between the
microbiota and the immune system. *Science,* 336**,** 1268-73.
IRRAZABAL, T., BELCHEVA, A., GIRARDIN, S. E., MARTIN, A. & PHILPOTT, D. J. 2014. The
multifaceted role of the intestinal microbiota in colon cancer. *Mol Cell,* 54**,** 309-20.
JIANG, H. & EDGAR, B. A. 2012. Intestinal stem cell function in Drosophila and mice. *Curr Opin*
Genet Dev, 22**,** 354-60.
JIANG, H., PATEL, P. H., KOHLMAIER, A., GRENLEY, M. O., MCEWEN, D. G. & EDGAR, B. A. 2009.
Cytokine/Jak/Stat signaling mediates regeneration and homeostasis in the Drosophila
midgut. *Cell,* 137**,** 1343-55.
KAZANJIAN, A. & SHROYER, N. F. 2011. NOTCH Signaling and ATOH1 in Colorectal Cancers. *Curr*
Colorectal Cancer Rep, 7**,** 121-127.
LEMAITRE, B. & MIGUEL-ALIAGA, I. 2013. The digestive tract of Drosophila melanogaster. *Annu*
Rev Genet, 47**,** 377-404.
LHOCINE, N., RIBEIRO, P. S., BUCHON, N., WEPF, A., WILSON, R., TENEV, T., LEMAITRE, B.,
GSTAIGER, M., MEIER, P. & LEULIER, F. 2008. PIMS modulates immune tolerance by
negatively regulating Drosophila innate immune signaling. *Cell Host Microbe,* 4**,** 147-58.
LYNE, R., SMITH, R., RUTHERFORD, K., WAKELING, M., VARLEY, A., GUILLIER, F., JANSSENS, H., JI,
W., MCLAREN, P., NORTH, P., RANA, D., RILEY, T., SULLIVAN, J., WATKINS, X.,
WOODBRIDGE, M., LILLEY, K., RUSSELL, S., ASHBURNER, M., MIZUGUCHI, K. & MICKLEM,
G. 2007. FlyMine: an integrated database for Drosophila and Anopheles genomics.
Genome Biol, 8**,** R129.
MARIANES, A. & SPRADLING, A. C. 2013. Physiological and stem cell compartmentalization
within the Drosophila midgut. *Elife,* 2**,** e00886.
MEDZHITOV, R., SCHNEIDER, D. S. & SOARES, M. P. 2012. Disease tolerance as a defense
strategy. *Science,* 335**,** 936-41.
MICCHELLI, C. A. & PERRIMON, N. 2006. Evidence that stem cells reside in the adult Drosophila
midgut epithelium. *Nature,* 439**,** 475-9.
OHLSTEIN, B. & SPRADLING, A. 2006. The adult Drosophila posterior midgut is maintained by
pluripotent stem cells. *Nature,* 439**,** 470-4.
OHLSTEIN, B. & SPRADLING, A. 2007. Multipotent Drosophila intestinal stem cells specify
daughter cell fates by differential notch signaling. *Science,* 315**,** 988-92.

PAQUETTE, N., BROEMER, M., AGGARWAL, K., CHEN, L., HUSSON, M., ERTURK-HASDEMIR, D.,
REICHHART, J. M., MEIER, P. & SILVERMAN, N. 2010. Caspase-mediated cleavage, IAP
binding, and ubiquitination: linking three mechanisms crucial for Drosophila NF-kappaB
signaling. *Mol Cell,* 37**,** 172-82.
PATEL, P. H., DUTTA, D. & EDGAR, B. A. 2015. Niche appropriation by Drosophila intestinal stem
cell tumours. *Nat Cell Biol,* 17**,** 1182-92.
PATEL, P. H. & EDGAR, B. A. 2014. Tissue design: how Drosophila tumors remodel their
neighborhood. *Semin Cell Dev Biol,* 28**,** 86-95.
PEIGNON, G., DURAND, A., CACHEUX, W., AYRAULT, O., TERRIS, B., LAURENT-PUIG, P.,
SHROYER, N. F., VAN SEUNINGEN, I., HONJO, T., PERRET, C. & ROMAGNOLO, B. 2011.
Complex interplay between beta-catenin signalling and Notch effectors in intestinal
tumorigenesis. *Gut,* 60**,** 166-76.
PERDIGOTO, C. N., SCHWEISGUTH, F. & BARDIN, A. J. 2011. Distinct levels of Notch activity for
commitment and terminal differentiation of stem cells in the adult fly intestine.
Development, 138**,** 4585-95.
PETKAU, K., PARSONS, B. D., DUGGAL, A. & FOLEY, E. 2014. A deregulated intestinal cell cycle
program disrupts tissue homeostasis without affecting longevity in Drosophila. *J Biol*
Chem, 289**,** 28719-29.
REEDIJK, M., ODORCIC, S., ZHANG, H., CHETTY, R., TENNERT, C., DICKSON, B. C., LOCKWOOD, G.,
GALLINGER, S. & EGAN, S. E. 2008. Activation of Notch signaling in human colon
adenocarcinoma. *Int J Oncol,* 33**,** 1223-9.
RIZZO, P., OSIPO, C., FOREMAN, K., GOLDE, T., OSBORNE, B. & MIELE, L. 2008. Rational targeting
of Notch signaling in cancer. *Oncogene,* 27**,** 5124-31.
ROBERTSON, S. J., ZHOU, J. Y., GEDDES, K., RUBINO, S. J., CHO, J. H., GIRARDIN, S. E. &
PHILPOTT, D. J. 2013. Nod1 and Nod2 signaling does not alter the composition of
intestinal bacterial communities at homeostasis. *Gut Microbes,* 4**,** 222-31.
RODILLA, V., VILLANUEVA, A., OBRADOR-HEVIA, A., ROBERT-MORENO, A., FERNANDEZ-
MAJADA, V., GRILLI, A., LOPEZ-BIGAS, N., BELLORA, N., ALBA, M. M., TORRES, F.,
DUNACH, M., SANJUAN, X., GONZALEZ, S., GRIDLEY, T., CAPELLA, G., BIGAS, A. &
ESPINOSA, L. 2009. Jagged1 is the pathological link between Wnt and Notch pathways in
colorectal cancer. *Proc Natl Acad Sci U S A,* 106**,** 6315-20.
SCHNEIDER, D. S. & AYRES, J. S. 2008. Two ways to survive infection: what resistance and
tolerance can teach us about treating infectious diseases. *Nat Rev Immunol,* 8**,** 889-95.
SHANAHAN, M. T., CARROLL, I. M., GROSSNIKLAUS, E., WHITE, A., VON FURSTENBERG, R. J.,
BARNER, R., FODOR, A. A., HENNING, S. J., SARTOR, R. B. & GULATI, A. S. 2014. Mouse
Paneth cell antimicrobial function is independent of Nod2. *Gut,* 63**,** 903-10.
SHIH IE, M. & WANG, T. L. 2007. Notch signaling, gamma-secretase inhibitors, and cancer
therapy. *Cancer Res,* 67**,** 1879-82.
SIKANDAR, S. S., PATE, K. T., ANDERSON, S., DIZON, D., EDWARDS, R. A., WATERMAN, M. L. &
LIPKIN, S. M. 2010. NOTCH signaling is required for formation and self-renewal of tumor-
initiating cells and for repression of secretory cell differentiation in colon cancer. *Cancer*
Res, 70**,** 1469-78.
SIUDEJA, K., NASSARI, S., GERVAIS, L., SKORSKI, P., LAMEIRAS, S., STOLFA, D., ZANDE, M.,
BERNARD, V., RIO FRIO, T. & BARDIN, A. J. 2015. Frequent Somatic Mutation in Adult

Intestinal Stem Cells Drives Neoplasia and Genetic Mosaicism during Aging. *Cell Stem*
Cell, 17**,** 663-74.
STANGER, B. Z., DATAR, R., MURTAUGH, L. C. & MELTON, D. A. 2005. Direct regulation of
intestinal fate by Notch. *Proc Natl Acad Sci U S A,* 102**,** 12443-8.
TAKASHIMA, S., ADAMS, K. L., ORTIZ, P. A., YING, C. T., MORIDZADEH, R., YOUNOSSI-
HARTENSTEIN, A. & HARTENSTEIN, V. 2011. Development of the Drosophila entero-
endocrine lineage and its specification by the Notch signaling pathway. *Dev Biol,* 353**,**
161-72.
THOMAS, P. D., CAMPBELL, M. J., KEJARIWAL, A., MI, H., KARLAK, B., DAVERMAN, R., DIEMER,
K., MURUGANUJAN, A. & NARECHANIA, A. 2003. PANTHER: a library of protein families
and subfamilies indexed by function. *Genome Res,* 13**,** 2129-41.
VAN ES, J. H., VAN GIJN, M. E., RICCIO, O., VAN DEN BORN, M., VOOIJS, M., BEGTHEL, H.,
COZIJNSEN, M., ROBINE, S., WINTON, D. J., RADTKE, F. & CLEVERS, H. 2005.
Notch/gamma-secretase inhibition turns proliferative cells in intestinal crypts and
adenomas into goblet cells. *Nature,* 435**,** 959-63.
WLODARSKA, M., KOSTIC, A. D. & XAVIER, R. J. 2015. An integrative view of microbiome-host
interactions in inflammatory bowel diseases. *Cell Host Microbe,* 17**,** 577-91.
642