Characteristics of functional enrichment and gene expression level of

human putative transcriptional target genes

Naoki Osato[1*]

[1]Department of Bioinformatic Engineering, Graduate School of Information Science

and Technology, Osaka University, Osaka 565-0871, Japan

[*]Corresponding author

E-mail address: naokiosato11@gmail.com, nosato@ist.osaka-u.ac.jp

Abstract

Background: Transcriptional target genes show functional enrichment of genes. However, how many and how significantly transcriptional target genes include functional enrichments are still unclear. To address these issues, I predicted human transcriptional target genes using open chromatin regions, ChIP-seq data and DNA binding sequences of transcription factors in databases, and examined functional enrichment and gene expression level of putative transcriptional target genes.

Results: Gene Ontology annotations showed four times larger numbers of functional enrichments in putative transcriptional target genes than gene expression information alone, independent of transcriptional target genes. To compare the number of functional enrichments of putative transcriptional target genes between cells or search conditions, I normalized the number of functional enrichment by calculating its ratios in the total number of transcriptional target genes. With this analysis, native putative transcriptional target genes showed the largest normalized number of functional enrichments, compared with target genes including 5 – 60% of randomly selected genes. The normalized number of functional enrichments was changed according to the criteria of enhancer-promoter interactions such as distance from transcriptional start sites and orientation of CTCF-binding sites. Forward-reverse orientation of CTCF-binding sites showed significantly higher normalized number of functional enrichments than the other orientations. Journal papers showed that the top five frequent functional enrichments were related to the cellular functions in the three cell types. The median

expression level of transcriptional target genes changed according to the criteria of

enhancer-promoter assignments (i.e. interactions) and was correlated with the changes

of the normalized number of functional enrichments of transcriptional target genes.

Conclusions: Human putative transcriptional target genes showed significant functional

enrichments. Functional enrichments were related to the cellular functions. The

normalized number of functional enrichments of human putative transcriptional target

genes changed according to the criteria of enhancer-promoter assignments and

correlated with the median expression level of the target genes. These analyses and

characters of human putative transcriptional target genes would be useful to examine

the criteria of enhancer-promoter assignments and to predict the novel mechanisms and

factors such as DNA binding proteins and DNA sequences of enhancer-promoter

interactions.

Keywords: transcriptional target genes, functional enrichment, gene expression,

transcription factors, enhancer, open chromatin regions, ChIP-seq, CTCF

Background

More than 400 types of cells have been found in the human body. Human

development is accompanied by the differentiation of stem cells into various cell types,

leading to a diversification of their phenotypes and functions. For example, the

development of the immune system involves differentiation and diversification of stem

cells into various types of mature immune cells. The functions of monocytes include

phagocytosis and antigen presentation. $CD4^+$ T cells, however, play a central role in

cell-mediated immunity and are involved in the activation of phagocytes and

antigen-specific cytotoxic T-lymphocytes, and the release of various cytokines in

response to an antigen. The $CD20^+$ B cells are involved in the production of antibodies

against antigens.

Differentiation of cells is often triggered by the expression of transcription factors

(TF) followed by the expression of their target genes, which results in the

transformation of cells into other cell types. For example, the transcription factors PU.1

and CCAAT enhancer-binding protein α (C/EBPα) play a critical role in the expression

of myeloid-specific genes and the generation of monocytes and macrophages [1, 2]. The

transcription factor GATA-3 is essential for early T cell development and the

differentiation of naive $CD4^+$ T cells into Th2 effector cells [3]. E2A, EBF1, PAX5, and

Ikaros are among the most important transcription factors that control early

development in mice, thereby conditioning homeostatic B cell lymphopoiesis [4].

We previously examined the differentiation of monocytes and macrophages in

mice, and discovered that the transcription factor IRF8 was essential for cellular

differentiation [5]. An analysis of transcription factor-binding sites (TFBS) revealed that

IRF8 regulated the expression of KLF4 through the IRF8 transcriptional cascade.

Functional enrichment analyses revealed that the target genes of IRF8 showed

functional enrichment for antigen presentation, whereas those of KLF4 showed

functional enrichments for phagocytosis and locomotion. These results suggested that

the transcriptional cascades of IRF8 and KLF4 included different functional modules of

target genes.

Functional enrichments of transcriptional cascades of IRF8 and KLF4 appeared to

be related to the cellular functions of monocytes and macrophages. Although several

transcription factors were expressed in monocytes and macrophages, the number of

these transcriptional target genes that resulted in functional enrichments remains

unknown. Whether transcriptional target genes in other human cells show functional

enrichments remain unclear. If the transcriptional target genes showed significant

functional enrichment, analyzing transcriptional target genes would be useful in

identifying genes involved in a specific cellular function. Using the budding yeast,

previous studies examined the functional enrichments on a genome-scale genetic

interaction map using the GeneMANIA algorithm [6-8]. Using bacterial systems, the

analyses of functional enrichments of predicted regulatory networks were performed

using Gene Ontology annotations [9]. Various databases of functional annotations of

genes and pathways exist. Analysis of functional enrichments is expected to be useful

for understanding the association of genes involved in similar functions and same

pathways, and for predicting unknown gene functions such as non-protein-coding

RNAs. In addition, the extent of enhancer contribution to functional enrichments of

transcriptional target genes remains unknown.

In this study, transcriptional target genes were predicted using public databases of

open chromatin regions of human monocytes, naive CD4$^+$ T, CD20$^+$ B cells, HUVEC,

IMR90, MCF-7, HMEC, H1-hESC, iPSC, and ChIP-seq data of human H1-hESC cells

and known transcription factor binding sequences. Functional enrichment analyses of

putative transcriptional target genes were conducted using 10 different annotation

databases of functional annotations and pathways. The gene expression level of

transcriptional target genes was examined in the cells.

Results

Prediction of transcriptional target genes

To examine functional enrichments of transcriptional target genes in a genome

scale, transcriptional target genes were predicted in human monocytes, CD4$^+$ T cells,

and CD20$^+$ B cells. Searches for known transcription factor binding sequences, which

were collected from various databases and papers, were conducted in open chromatin

regions of the promoter sequences of RefSeq transcripts (Figure 1, see Methods).

Among 6,277 transcription factor binding sequences derived from vertebrates, 4,373

were linked to 1,018 TF transcripts computationally (see Methods). To maintain the

sensitivity of the searches for transcription factor binding sites and as some transcription

factors will recognize multiple distinctly different sequence motifs, transcription factor

binding sequences that targeted the same genes were recognized as redundant, and one

of the sequences was used [10] (see Methods). In total, 3,337 transcription factor

binding sequences in human monocytes, 3,652 in CD4$^+$ T cells, and 3,187 in CD20$^+$ B

cells were identified with their target genes, which were selected from highly expressed

genes in a cell (top 30% expression level, see Methods).

The total numbers of unique highly expressed target genes of transcription factor

binding sequences were 4,481, 7,558, and 4,753 in monocytes, CD4$^+$ T cells, and

CD20$^+$ B cells respectively using promoters. The mean target genes of a transcription

factor were 124, 164, and 144 in monocytes, CD4$^+$ T cells, and CD20$^+$ B cells,

respectively, with the corresponding medians being 24, 33, and 24, respectively. With

regard to the genomic localizations of TFBS, 51%, 65%, and 61% of TFBS were

located within promoters (±5 kb of TSS) of target genes in monocytes, CD4$^+$ T cells,

and CD20$^+$ B cells, respectively (according to association rule 1, see Methods).

Functional enrichments of putative transcriptional target genes

Functional enrichments of the putative target genes were examined. The

distribution of functional enrichments in transcriptional target genes was predicted

using genome sequences of promoters in the three cell types (Figure 1 and Table 1, see

Methods). Furthermore, the effect of transcriptional target genes including randomly

selected genes on functional enrichments was investigated using DNase-DGF data of

monocytes, CD4$^+$ T and CD20$^+$ B cells, HUVEC, IMR90, MCF-7, HMEC, and

ChIP-seq data of H1-hESC (Figure 2A and B, and Figure S1 and S2, see Methods). The

native putative transcriptional target genes not including randomly selected genes

showed the highest functional enrichments using Gene Ontology, GO Slim, KEGG,

Pathway Commons, WikiPathways, InterPro and UniProt functional regions (Domains)

in both DNase-DGF and ChIP-seq data of the five types of cells. Of the 10 databases

used in this analysis, the Gene Ontology database consists of three types of functional

annotations, i.e., 20,836 biological processes, 9,020 molecular functions, and 2,847

cellular components. The numbers of functional enrichments of Gene Ontology

annotations in target genes of a transcription factor were 2,902, 4,077, and 2,778 in

monocytes, CD4$^+$ T cells, and CD20$^+$ B cells, respectively. An examination of

functional enrichments of highly expressed genes (top 30% expression level),

independent of transcriptional target genes, revealed 237, 301, and 239 'unique' Gene

Ontology annotations in monocytes, CD4$^+$ T cells, and CD20$^+$ B cells, respectively

(Table 1). Further, the examination of functional enrichments of highly expressed target

genes (top 30% expression level) in target genes revealed 1,271, 1,654, and 1,192

'unique' Gene Ontology annotations in monocytes, CD4$^+$ T cells, and CD20$^+$ B cells,

respectively i.e., These numbers were four times larger than functional enrichments

identified by gene expression information alone, independent of transcriptional target

genes, suggesting that transcriptional target genes were frequently associated with

similar functions or pathways (Table S3 and S4).

Functional enrichments of transcriptional target genes from other databases were

also examined (Table 1). KEGG, Target genes of transcription factors, Disease Ontology,

GO Slim, Pathway Commons, Cellular biomarkers, Target genes of microRNAs,

Protein domains, and WikiPathways had 95, 16, 127, 12, 242, 17, 97, 303, and 105

unique functional annotations, respectively. The numbers of functional enrichments of

transcriptional target genes in the other annotation databases except for microRNAs and

Protein domains were significantly higher than gene expression information alone,

independent of transcriptional target genes, as well as Gene Ontology annotations

(Table 1). The functional enrichments of transcriptional target genes from Pathway

Commons for monocytes, CD4$^+$ T cells, and CD20$^+$ B cells are shown in Table 2 and

Table S5. Functional enrichments were found to be related to cellular functions, e.g.,

interferon signaling, GMCSF (Granulocyte-macrophage colony-stimulating factor, a

kind of cytokine)-mediated signaling events, antigen processing-cross presentation in

monocytes; TCR (T-cell receptor) signaling in naive CD4$^+$ T cells, IL-12 (Interleukin-12,

a kind of cytokine)-mediated signaling events, and downstream signaling in naive CD8$^+$

T cells in CD4$^+$ T cells; interferon alpha/beta signaling, IL8- and CXCR2 (Chemokine

receptor type 2, a kind of cytokine)-mediated signaling events, and BCR (B cell antigen

receptor) signaling pathway in CD20$^+$ B cells. WikiPathways, KEGG and GO also

revealed that functional enrichments were associated with cellular functions (Table S6,

S7 and S8).

Effect of enhancer-promoter association rules on functional enrichments

To understand the effect of 'promoter and extended regions for

enhancer-promoter association (EPA)' on the functional enrichments of target genes, the

rule of extended regions was modified according to four criteria (Figure 3A and see

Methods) [11], and functional enrichments were investigated.

According to the association rule (1), the means of target genes were 177, 217,

and 175 in monocytes, CD4$^+$ T cells, and CD20$^+$ B cells, respectively, whereas the

corresponding medians were 55, 58, and 37, respectively (Table S9). The numbers of

functional enrichments of Pathway Commons annotations using promoter regions were

1,005, 1,806, and 821 in monocytes, CD4$^+$ T cells, and CD20$^+$ B cells, respectively

(Table S10). With the use of EPA (association rule 1), the numbers of functional

enrichments of Pathway Commons annotations were 3,087, 7,216, and 3,900,

representing 3.07-, 4.00-, and 4.75-fold increases, respectively, in the three cells types.

Additionally, the numbers of 'unique' Pathway Commons annotations with promoter

regions were 321, 415, and 329 in monocytes, $CD4^+$ T cells, and $CD20^+$ B cells,

respectively; the corresponding numbers with the use of EPA (association rule 1) were

364, 437, and 364, representing 1.13-, 1.05-, and 1.11-fold increases, respectively, in the

three cell types. The normalized numbers of functional enrichments of Pathway

Commons annotations were 44.75, 84.51, and 59.32, representing 1.84-, 2.80-, and

3.32-fold increases, respectively, in the three cell types (association rule 1, Table 3).

Other cell types also showed the same tendencies (Table 3).

The normalized numbers of the functional enrichments of transcriptional target

genes showed association rule (4) as the highest number, followed by association rule

(1) and (2) in the three cell types. Although association rule (3) was the longest among

the four criteria, it showed the lowest number of functional enrichments in the three cell

types (Figure 3A and Table 3). ChIP-seq data of 19 TF in H1-hESC (Human embryonic

stem cells) also showed almost the same tendency (difference between association rule

(4) and (1) was not statistically significant, probably due to a large number of

transcriptional target genes predicted using 19 TF ChIP-seq data. Several thousands of

target genes of each TF were predicted. Some of them would be indirect interactions

between TF and genome DNA, which were identified by ChIP-seq experiments. (Table

S11, see Additional file).

Differences in functional enrichments using Pathway Commons were examined

between promoters *versus* EPA (association rule 1) (Table 4 and Table S12). A

comparison of 321 and 364 functional enrichments using the promoters and EPA,

respectively, in monocytes revealed that 152 (47% in promoters, 42% in extended

regions) of them were common. For example, IFN-gamma (Interferon gamma) pathway,

GMCSF (Granulocyte-macrophage colony-stimulating factor, a kind of

cytokine)-mediated signaling events, and PDGF (Platelet-derived growth factor)

receptor signaling network were enriched using extended regions (association rule 1) as

opposed to promoters (Table S12). The comparison of 415 (promoters) and 437

(extended regions) functional enrichments in $CD4^+$ T cells revealed that 163 of them

(39% in promoters, 37% in extended regions) were common. IFN-gamma pathway,

TCR (T-cell receptor) signaling in naive $CD4^+$ T cells, and IL3 (Interleukin-3, a kind of

cytokine)-mediated signaling events were enriched using extended regions. The

comparison of 329 (promoters) and 364 (extended regions) functional enrichments in

$CD20^+$ B cells revealed that 171 of them (52% in promoters, 47% in extended regions)

were common. IL5-mediated signaling events, IL4-mediated signaling events, and

cytokine signaling in immune system were enriched in $CD20^+$ B cells using extended

regions. Only about 40% of functional enrichments of Pathway Commons annotations

were unchanged between promoters and EPA. EPA significantly affected the functional

enrichments of transcriptional target genes. Journal papers showed that frequent

functional enrichments were related to the cellular functions in the three cell types

(Table 4). These results showed that new functional enrichments related to cellular

functions were identified using extended regions for enhancer-promoter association.

Effect of CTCF-binding sites on functional enrichments

CTCF have the activity of insulators to block the interaction between enhancers

and promoters [12]. Recent studies identified a correlation between the orientation of

CTCF-binding sites and chromatin loops (Figure 3B) [13]. Forward–reverse (FR)

orientation of CTCF-binding sites are frequently found in chromatin loops. To examine

the effect of forward–reverse orientation of CTCF-binding sites on functional

enrichments of target genes, 'promoter and extended regions for enhancer-promoter

association (EPA)' were shortened at the genomic locations of forward–reverse

orientation of CTCF-binding sites, and transcriptional target genes were predicted from

the shortened regions using TFBS (see Methods). The numbers of functional

enrichments of target genes were investigated. According to EPA (association rule 4)

that were shortened at genomic locations of forward–reverse orientation of

CTCF-binding sites, the means of target genes were 67, 64, and 77 in monocytes, $CD4^+$

T cells, and $CD20^+$ B cells, respectively, whereas the corresponding medians were 23,

21, and 20, respectively (Table S13). The normalized numbers of functional

enrichments of Pathway Commons annotations using EPA were 71.42, 108.08, and

90.99 in monocytes, $CD4^+$ T cells, and $CD20^+$ B cells, respectively (Table 5). With the

use of EPA shortened at forward–reverse orientation of CTCF-binding sites, the

normalized numbers of functional enrichments of Pathway Commons annotations were

196.58, 220.54, and 220.77, representing 2.75-, 2.04-, and 2.43-fold increases,

respectively, in the three cells types. Additionally, the normalized numbers of functional

enrichments of 'unique' Pathway Commons annotations with EPA were 5.09, 5.34, and

6.00 in monocytes, CD4$^+$ T cells, and CD20$^+$ B cells, respectively; the corresponding

normalized numbers with the use of EPA shortened at forward–reverse orientation of

CTCF-binding sites were 9.88, 10.72, and 9.10, representing 1.94-, 2.01-, and 1.52-fold

increases, respectively, in the three cell types (Table S14). Other cell types also showed

the same tendencies (Table 5). The normalized numbers of functional enrichments were

significantly increased between EPA and EPA shortened at forward–reverse orientation

of CTCF-binding sites in Gene Ontology, Disease Ontology, Pathway Commons, GO

Slim, WikiPathways, KEGG, InterPro and UniProt functional regions (Domains)

annotations. These increases were also significant, compared with EPA shortened at

CTCF-binding sites without the consideration of their orientation. Transcriptional target

genes predicted from EPA shortened at forward–reverse orientation of CTCF-binding

sites tend to include similar function of genes significantly.

Differences in functional enrichments obtained using EPA *versus* EPA shortened

at forward–reverse orientation of CTCF-binding sites were examined using the

functional enrichments of Pathway Commons (Table 6 and Results in Additional file).

Transcriptional target genes predicted from EPA shortened at the CTCF-binding sites

tended to include the similar function of genes. About 40 – 80% of functional

enrichments were unchanged between promoters and EPA shortened at forward–reverse

orientation of CTCF-binding sites, and the functional enrichments observed in EPA

shortened at forward–reverse orientation of CTCF-binding sites as opposed to

promoters included various immunological terms. Journal papers showed that the top

five frequent functional enrichments were related to the cellular functions in the three

cell types (Table 6). These results showed that new functional enrichments related to

cellular functions were identified using forward–reverse orientation of CTCF-binding

sites.

Comparison of expression levels of putative transcriptional target genes

To examine the relationship between functional enrichments and expression

levels of target genes, the expression levels of target genes predicted from promoters

and three types of 'promoter and extended regions for enhancer-promoter assignment

(EPA)' were investigated in monocytes, CD4$^+$ T, H1-hESC and iPSC (Figure 4). Median

expression levels of the target genes of the same transcription factor binding sequences

were compared between promoters and three types of EPA. Red and blue dots in Figure

4 show statistically significant difference of the distribution of expression levels of

target genes between promoters and EPA. Additionally, "red dots" show the median

expression level of target genes of a TFBS was 'higher' in EPA than promoters, and

"blue dots" show the median expression level of target genes of a TFBS was 'lower' in

EPA than promoters. The ratios of red dots were higher in EPA (association rule 4) that

were shortened at forward–reverse orientation of CTCF-binding sites *versus* promoters

(left graph in Figure 4) than EPA (association rule 4) *versus* promoters (right graph) in

monocytes and CD4$^+$ T cells. The ratios of blue dots were higher in EPA (association

rule 4) that were shortened at forward–reverse orientation of CTCF-binding sites *versus*

promoters (left graph) than EPA (association rule 4) *versus* promoters (right graph) in

H1-hESC and iPSC. Moreover, the ratio of the sum of median expression levels

between the three types of EPA and promoters in monocytes and $CD4^+$ T cells was the

highest in EPA shortened at forward–reverse orientation of CTCF-binding sites (Table

S16). Conversely, the ratio of the sum of median expression levels between the three

types of EPA and promoters in H1-hESC and iPSC was the lowest in EPA shortened at

forward–reverse orientation of CTCF-binding sites.

EPA shortened at forward–reverse orientation of CTCF-binding sites changed (i.e.

increased or decreased) the expression levels of target genes more than the other types

of EPA. This implied that gene expression tended to be activated in monocytes and

$CD4^+$ T cells, but repressed in H1-hESC and iPSC by enhancers. EPA shortened at

forward–reverse orientation of CTCF-binding sites also showed the highest normalized

number of functional enrichments of transcriptional target genes, as shown in the

previous paragraphs.

Discussion

Genome-wide functional enrichments and gene expression levels of putative

target genes of human transcription factors were investigated. Human putative

transcriptional target genes showed significantly larger numbers of functional

enrichments than gene expression information alone, independent of transcriptional

target genes. Moreover, when the number of functional enrichments of human putative

transcriptional target genes was normalized by the total number of transcriptional target

genes, native putative transcriptional target genes showed the highest ratio of functional

enrichments, compared with target genes partially including randomly selected genes.

The ratio of functional enrichments was decreased according to the increase of the ratio

of randomly selected genes in target genes. These tendencies were observed in putative

transcriptional target genes predicted from both open chromatin regions and ChIP-seq

data of transcription factors. Prediction of transcriptional target genes from open

chromatin regions includes false positives, since DNase I cleavage bias affects the

computational analysis of DNase-seq experiments [14]. However, the detection of

ChIP-seq peaks is also changed depending on the methods to identify them and the

depth of DNA sequencing of ChIP-seq experiments [15]. Though human putative

transcriptional target genes include false positives, they showed significantly the largest

number of functional enrichments, compared with target genes including 5 – 60% of

randomly selected genes (Figure 2).

The median expression level of human putative transcriptional target genes was

changed according to the criteria of enhancer-promoter assignments, and was correlated

with the normalized number of functional enrichments. The median expression level of

transcriptional target genes was 'decreased' significantly in transcriptional target genes

predicted using enhancers, compared with those predicted using promoters in H1-hESC

and iPSC, and the median expression level was 'increased' significantly in target genes

predicted using enhancers, compared with those predicted using promoters in

monocytes and $CD4^+$ T cells. These results implied that transcription factors bound in

enhancers act as repressors in H1-hESC (ES) and iPSC, but those act as activators in

monocytes and CD4$^+$ T cells. The change of functional roles of transcription factors

depending on the cell types would be analyzed and reported elsewhere.

The median expression level was increased significantly in target genes predicted

using enhancers, compared with those predicted from promoters in immune cells using

gene expression data (Blueprint RNA-seq RPKM data; GSE58310), but smaller number

of target genes showed the increase of median expression level using gene expression

data (ENCODE; GSM984609). The results of the analyses may be slightly different

depending on gene expression data. H1-hESC (ES) and iPSC showed a strong tendency

of decrease of median expression levels of transcriptional target genes between

enhancers and promoters.

The gene symbols of transcription factors were sometimes different among

databases, because more than one gene symbol are assigned to some transcription

factors and some gene symbols are spelled in several different ways. These differences

need to be identified with manual curations. This analysis will be required to predict

transcriptional cascades by associating transcription factors with transcriptional target

genes consisting of transcription factors. In the analyses of transcriptional cascades, to

reduce false positive predictions of enhancer-promoter associations from open

chromatin regions, the identification of DNase peaks will be modified using a new tool

such as HINT [16].

In this study, I focused on three types of immune cells and stem cells such as

H1-hESC and iPSC to examine transcriptional target genes in a genome scale, since in

my previous study, I examined transcriptional cascades involved in the differentiation of

immune cells as introduced in Background [5]. Furthermore, I confirmed the features of

functional enrichments of putative transcriptional target genes are commonly found in

other four types of normal and disease cells (HUVEC, IMR90, MCF-7 and HMEF).

It is difficult to predict enhancer-promoter associations using a single parameter,

so that machine learning methods to combine several parameters have been proposed

[17-19]. These methods showed high accuracy in predicting enhancer-promoter

associations (I tried to use some of the tools, but they did not work properly. I am

waiting for the authors to update the tools). However, molecular mechanisms of

enhancer-promoter interactions are not clearly understood. CTCF has been found to

bind at chromatin interaction anchors and form chromatin interactions [12]. About

20-40% of chromatin interaction anchors included DNA binding sequences of CTCF,

when I examined public Hi-C experimental data [20, 21]. Among 33,939 RefSeq

transcripts, 7,202 (21%), 4,404 (13%), and 6,921 (20%) (p-value $< 10^{-5}$ in the search for

CTCF-binding motifs using FIMO) to 9,608 (28%), 5,806 (17%), and 9,137 (27%)

(p-value $< 10^{-4}$) of transcripts had forward–reverse orientation of CTCF-binding sites

within 1 Mb from transcriptional start sites in the three immune cell types, respectively.

These analyses implied that other factors might be involved in chromatin interactions.

ZNF143 has been reported to locate at promoter regions of chromatin interaction

anchors [22]. To predict the other factors and molecular mechanisms, the analyses in

this study would be useful to examine further the criteria in predicting

enhancer-promoter associations. Machine learning methods need the information what

parameters should be used for prediction, so it would be better to choose parameters

involved in predicting enhancer-promoter associations. To improve the prediction and

understand the molecular mechanisms of enhancer-promoter interactions, I am

promoting the analyses of chromatin interaction anchors, and the results of the analyses

will be reported elsewhere.

Conclusion

In this study, human transcriptional target genes were predicted using open

chromatin regions, ChIP-seq data, and DNA binding sequences of transcription factors

in databases. Human putative transcriptional target genes showed significant functional

enrichments. Journal papers showed that frequent functional enrichments were related

to the cellular functions. The normalized number of functional enrichments was the

highest in native putative transcriptional target genes, compared with target genes

partially replaced with randomly selected genes. The normalized number of functional

enrichments of human putative transcriptional target genes changed according to the

criteria of enhancer-promoter assignments and correlated with the median expression

level of the target genes. The normalized numbers of functional enrichments of

transcriptional target genes did not show the highest number in the criterion of

enhancer-promoter assignments covering the longest distance from transcriptional start

site among four criteria. This suggested that there is a criterion of enhancer-promoter

assignments that shows the highest normalized number of functional enrichments. The

median expression level of transcriptional target genes was 'decreased' significantly in

transcriptional target genes predicted using enhancers, compared with those predicted

using promoters in H1-hESC and iPSC, and the median expression level was 'increased'

significantly in target genes predicted using enhancers, compared with those predicted

using promoters in immune cells. These results implied that transcription factors bound

in enhancers act as repressors in H1-hESC (ES) and iPSC, but those act as activators in

immune cells. These analyses and characters of human putative transcriptional target

genes would be useful to examine the criteria of enhancer-promoter assignments and to

predict the novel mechanisms and factors such as DNA binding proteins and DNA

sequences of enhancer-promoter interactions.

Methods

Searches for transcription factor binding sequences from open chromatin regions

To examine transcriptional regulatory target genes, bed files of hg19 narrow

peaks of ENCODE DNase-DGF and DNase data for Monocytes-CD14$^+$_RO01746

(GSM1024791; UCSC Accession: wgEncodeEH001196), CD4$^+$_Naive_Wb11970640

(GSM1014537; UCSC Accession: wgEncodeEH003156), CD20$^+$_RO01778

(GSM1014525; UCSC Accession: wgEncodeEH002442), H1-hESC (GSM816632;

UCSC Accession: wgEncodeEH000556), iPSC (GSM816642; UCSC Accession:

wgEncodeEH001110), HUVEC (GSM1014528; UCSC Accession:

wgEncodeEH002460), IMR90 (GSM1008586; UCSC Accession: wgEncodeEH003482),

MCF-7 (GSM816627; UCSC Accession: wgEncodeEH000579), and HMEC

(GSM816669; UCSC Accession: wgEncodeEH001101) from the ENCODE website

(http://hgdownload.cse.ucsc.edu/goldenPath/hg19/encodeDCC/wgEncodeUwDgf/)

were used. For comparison with transcriptional target genes predicted using ChIP-seq

data, bed files of hg19 narrow peaks of ENCODE ChIP-seq data for 19 transcription

factors (TF) (BACH1, BRCA1, C/EBPbeta, CHD2, c-JUN, c-MYC, GTF2I, JUND,

MAFK, MAX, MXI1, NRF1, RAD21, RFX5, SIN3A, SUZ12, TBP, USF2, ZNF143) in

H1-hESC   from   the   ENCODE   website

(https://genome.ucsc.edu/cgi-bin/hgFileUi?db=hg19&g=wgEncodeAwgTfbsUniform)

were utilized.

To identify transcription factor binding sites (TFBS) from the DNase-DGF data,

TRANSFAC (2013.2), JASPAR (2010), UniPROBE, BEEML-PBM, high-throughput

SELEX, Human Protein-DNA Interactome, and transcription factor binding sequences

of ENCODE ChIP-seq data were used [23-29]. Position weight matrices of transcription

factor binding sequences were transformed into TRANSFAC matrices and then into

MEME matrices using in-house Perl scripts and transfac2meme in MEME suite [30].

Transcription factor binding sequences of transcription factors derived from vertebrates

were used for further analyses. Searches were conducted for transcription factor binding

sequences from the central 50-bp regions of each narrow peak using FIMO with p-value

threshold of 10^{-5} [31]. Transcription factors corresponding to transcription factor

binding sequences were searched computationally by comparing their names and gene

symbols of HGNC (HUGO Gene Nomenclature Committee) -approved gene

nomenclature   and   31,848   UCSC   known   canonical   transcripts

(http://hgdownload.cse.ucsc.edu/goldenpath/hg19/database/knownCanonical.txt.gz),   as

transcription factor binding sequences were not linked to transcript IDs such as UCSC,

RefSeq, and Ensembl transcripts.

Prediction of transcriptional target genes

Target genes of a transcription factor were assigned when its TFBS was found in

DNase-DGF narrow peaks in promoter or extended regions for enhancer-promoter

association of genes (EPA). Promoter and extended regions were defined as follows:

promoter regions were those that were within distances of ±5 kb from transcriptional

start sites (TSS). Promoter and extended regions were defined as per the following four

association rules, which are similar or same as those defined in a previous study [11]:

(1) the basal plus extension association rule assigns a basal regulatory domain to each

gene regardless of other nearby genes. The domain is then extended to the basal

regulatory domain of the nearest upstream and downstream genes, and includes a 5 kb +

5 kb basal region and an extension up to 300 kb or the midpoint between the TSS of the

gene and that of the nearest gene upstream and downstream; (2) 5 kb + 1 kb basal

region and an extension up to 1 Mb; (3) the two nearest genes association rule, which

extends the regulatory domain to the TSS of the nearest upstream and downstream

genes without the limitation of extension length; and (4) the single nearest gene

association rule, which extends the regulatory domain to the midpoint between the TSS

of the gene and that of the nearest gene upstream and downstream without the limitation

of extension length. Association rule (1) was used in our previous study [5]. Association

rule (2), (3), and (4) were the same as those in Figure 3A of the previous study [11],

however, association rules (3) and (4) did not have the limitation of extension length in

this study. The genomic positions of genes were identified using 'knownGene.txt.gz'

file in UCSC bioinformatics sites [32]. The file 'knownCanonical.txt.gz' was also

utilized for choosing representative transcripts among various alternate forms for

assigning promoter and extended regions for enhancer-promoter association of the

genes. From the list of transcription factor binding sequences and transcriptional target

genes, redundant transcription factor binding sequences were removed by comparing

the target genes of a transcription factor binding sequence and its corresponding

transcription factor; if identical, one of the transcription factor binding sequences was

used. When the number of transcriptional target genes predicted from a transcription

factor binding sequence was less than five, the transcription factor binding sequence

was omitted.

Gene expression analyses

For gene expression data, RNA-seq reads mapped onto human hg19 genome

sequences were obtained, including ENCODE long RNA-seq reads with poly-A of

monocytes CD14$^+$ cells, CD20$^+$ B cells, H1-hESC, iPSC, HUVEC, IMR90, MCF-7, and

HMEC (GSM984609, GSM981256, GSE26284, GSM958733, GSM2344099,

GSM2344100, GSM958734, GSM2400222, GSM765388, and GSM758571), and

UCSF-UBC human reference epigenome mapping project RNA-seq reads with poly-A

of naive CD4$^+$ T cells (GSM669617). Two replicates were present for monocytes

CD14$^+$ cells, CD20$^+$ B cells, H1-hESC, iPSC, HUVEC, IMR90, MCF-7, and HMEC

and a single one for CD4$^+$ T cells. RPKMs of the RNA-seq data were calculated using

RSeQC [33]. For monocytes, Blueprint RNA-seq RPKM data (GSE58310,

GSE58310_GeneExpression.csv.gz, Monocytes_Day0_RPMI) was also used [34].

Based on RPKM, UCSC transcripts with expression levels among top 30% of all the

transcripts were selected in each cell type.

Functional enrichment analyses

The functional enrichments of target genes of a TFBS and its corresponding

transcription factor were examined using GO-Elite v1.2.5 with *p*-value threshold at 1,

and after GO-Elite analyses a false discovery rate (FDR) test was performed with

q-value threshold at 10^{-3} to correct for multiple comparisons of thousands of groups of

transcriptional target genes in each cell type and condition [35]. For examining

functional enrichments of high or low expressed genes independent of transcriptional

target genes, the *p*-value threshold was set to 0.01 or 0.05 to confirm that the results

were not significantly changed. UCSC gene IDs were transformed into RefSeq IDs prior

to GO-Elite analyses. GO-Elite uses 10 databases for identifying functional

enrichments: (1) Gene Ontology, (2) Disease Ontology, (3) Pathway Commons, (4) GO

Slim, (5) WikiPathways, (6) KEGG, (7) Transcription factor to target genes, (8)

microRNA to target genes, (9) InterPro and UniProt functional regions (Domains), and

(10) Cellular biomarkers (BioMarkers). To calculate the normalized numbers of

functional enrichments of target genes, the numbers of functional enrichments were

divided by the total number of target genes in each cell type and condition, and were

multiplied by 10^5. In tables showing the numbers of functional enrichments in 10

databases, heat maps were plotted according to *Z*-scores calculated from the numbers of

functional enrichments of each database using in-house Excel VBA scripts. In the

comparisons of the normalized numbers of functional enrichments of target genes in

cell types and conditions, if the number of a functional annotation in a cell type or

condition was two times larger than that in the other cell type or condition, the

functional annotation was recognized as more enriched than the other cell type or

condition.

To investigate whether the normalized numbers of functional enrichments of

transcriptional target genes correlate with the prediction of target genes, a part of target

genes were changed with randomly selected genes with high expression level (top 30%

expression level), and functional enrichments of the target genes were examined. First,

5%, 10%, 20%, 40%, and 60% of target genes were changed with randomly selected

genes with high expression level in monocytes, CD4$^+$ T cells, and CD20$^+$ B cells.

Second, as another randomization of target genes, the same number of 5%, 10%, 20%,

40%, and 60% of target genes were selected randomly from highly expressed genes,

then added them to the original target genes, and functional enrichments of the target

genes were examined. All analyses were repeated three times to estimate standard errors

(Figure 2A and B, Figure S1 and S2, and Table S1). The same analysis was performed

using DNase-DGF data and ChIP-seq data of 19 TF in H1-hESC. Transcriptional target

genes were predicted from promoter (Tables S2).

CTCF-binding sites

CTCF   ChIP-seq   data   for   monocytes   CD14$^+$ cells

(GSM1003508_hg19_wgEncodeBroadHistoneMonocd14ro1746CtcfPk.broadPeak.gz),

CD4$^+$ T   cells   (SRR001460.bam),   CD20$^+$ B   cells

(GSM1003474_hg19_wgEncodeBroadHistoneCd20CtcfPk.broadPeak.gz),   H1-hESC

(wgEncodeAwgTfbsUtaH1hescCtcfUniPk.narrowPeak.gz), iPSC (GSE96477), HUVEC

(wgEncodeAwgTfbsUwHuvecCtcfUniPk.narrowPeak.gz),   IMR90

(wgEncodeAwgTfbsSydhImr90CtcfbIggrabUniPktfbsf.narrowPeak.gz),   MCF-7

(wgEncodeAwgTfbsUwMcf7CtcfUniPktfbsf.narrowPeak.gz),   and   HMEC

(wgEncodeAwgTfbsUwHmecCtcfUniPktfbsf.narrowPeak.gz)   were   used.

SRR001460.bam was sorted and indexed by SAMtools and transformed into a bed file

using bamToBed of BEDTools [36, 37]. ChIP-seq peaks were predicted by SICER-rb.sh

of SICER with optional parameters 'hg19 1 200 150 0.74 200 100' [38]. Extended

regions for enhancer-promoter association (association rule 4) were shortened at the

genomic locations of CTCF-binding sites that were the closest to a transcriptional start

site, and transcriptional target genes were predicted from the shortened enhancer

regions   using   TFBS.   Furthermore,   promoter   and   extended   regions   for

enhancer-promoter association (association rule 4) were shortened at the genomic

locations of forward–reverse orientation of CTCF-binding sites. When forward or

reverse orientation of CTCF-binding sites were continuously located in genome

sequences   several   times,   the   most   external   forward–reverse   orientation   of

CTCF-binding sites were selected.

Abbreviations

TF: transcription factors

TFBS: transcription factor binding sites

TSS: transcriptional start sites

ENCODE: encyclopedia of DNA elements

ChIP-seq: ChIP-sequencing, chromatin immunoprecipitation followed by massively

parallel DNA sequencing

RNA-seq: RNA-sequencing

EPA: promoter and extended regions for enhancer-promoter association

FR: forward-reverse

RF: reverse-forward

FF: forward-forward

RR: reverse-reverse

Declarations

Ethics approval and consent to participate

577   I used experimental data in public databases, so this is not applicable.

Consent for publication

580   Not applicable

Availability of data and material

I used experimental data in public and commercial databases, so this is not applicable.

TRANSFAC, which is a database of transcription factors, is commercial database.

Competing interests

The authors declare that they have no competing interests

Funding

Publication charges for this article have been funded by JSPS KAKENHI Grant

Number 16K00387. This research was partially supported by the Platform Project for

Supporting in Drug Discovery and Life Science Research（Platform for Dynamic

Approaches to Living System）from Japan Agency for Medical Research and

Development (AMED). This research was partially supported by Development of

Fundamental Technologies for Diagnosis and Therapy Based upon Epigenome Analysis

from Japan Agency for Medical Research and Development (AMED). This work was

partially supported by JST CREST Grant Number JPMJCR15G1, Japan.

Authors' contributions

NO designed and performed the research

Acknowledgements

The supercomputing resource was provided by Human Genome Center of the Institute

of Medical Science at the University of Tokyo. Computations were partially performed

on the NIG supercomputer at ROIS National Institute of Genetics.

Authors' information

References used in Table 4.

Monocytes: IFN-gamma pathway[39], IL3-mediated signaling events[40],

GMCSF-mediated signaling events[41], PDGF receptor signaling network[42], VEGF

and VEGFR signaling network[43]

CD4+ T cell: Integrin family cell surface interactions[44], IFN-gamma pathway[45],

LKB1 signaling events[46], TCR signaling in naive CD4+ T cells[47], IL3-mediated

signaling events[48]

CD20+ B cell: Integrin family cell surface interactions[49], IL5-mediated signaling

events[50], Insulin Pathway[51], mTOR signaling pathway[52], Sphingosine

1-phosphate (S1P) pathway[53]

References used in Table 6.

Monocytes: CXCR4-mediated signaling events[54], GPVI-mediated activation

cascade[55], Signaling events mediated by TCPTP[56], p38 MAPK signaling

pathway[57], IL4-mediated signaling events[58]

CD4+ T cell: CXCR4-mediated signaling events[59], Fc-epsilon receptor I signaling in

mast cells[60], JNK signaling in the CD4+ TCR pathway[61], Reelin signaling

pathway[62], Semaphorin interactions[63]

CD20+ B cell: Syndecan-4-mediated signaling events[64], CXCR4-mediated signaling

events[65], EPO signaling pathway[66], Stabilization and expansion of the E-cadherin

adherens junction[67], Interleukin-1 signaling[68]

Endnotes

References

Uncategorized References

1.   Valledor AF, Borras FE, Cullell-Young M, Celada A: **Transcription factors**
that regulate monocyte/macrophage differentiation. *Journal of leukocyte*
biology 1998, **63**(4):405-417.

2.   Nagamura-Inoue T, Tamura T, Ozato K: **Transcription factors that regulate**
growth and differentiation of myeloid cells. *International reviews of*
immunology 2001, **20**(1):83-105.

3.   Ho IC, Tai TS, Pai SY: **GATA3 and the T-cell lineage: essential functions**
before and after T-helper-2-cell differentiation. *Nature reviews Immunology*
2009, **9**(2):125-135.

4.   Rothenberg EV: **B cell specification from the genome up**. *Nature immunology*
2010, **11**(7):572-574.

5.   Kurotaki D, Osato N, Nishiyama A, Yamamoto M, Ban T, Sato H, Nakabayashi
J, Umehara M, Miyake N, Matsumoto N *et al*: **Essential role of the**
IRF8-KLF4 transcription factor cascade in murine monocyte
differentiation. *Blood* 2013, **121**(10):1839-1849.

6.   Costanzo M, Baryshnikova A, Bellay J, Kim Y, Spear ED, Sevier CS, Ding H,
Koh JL, Toufighi K, Mostafavi S *et al*: **The genetic landscape of a cell**. *Science*
(New York, NY) 2010, **327**(5964):425-431.

7.   Warde-Farley D, Donaldson SL, Comes O, Zuberi K, Badrawi R, Chao P, Franz
M, Grouios C, Kazi F, Lopes CT *et al*: **The GeneMANIA prediction server:**

biological network integration for gene prioritization and predicting gene
function. *Nucleic acids research* 2010, **38**(Web Server issue):W214-220.

8.   Costanzo M, VanderSluis B, Koch EN, Baryshnikova A, Pons C, Tan G, Wang
W, Usaj M, Hanchard J, Lee SD *et al*: **A global genetic interaction network**
maps a wiring diagram of cellular function. *Science (New York, N Y)* 2016,
353(6306).

9.   Marbach D, Costello JC, Kuffner R, Vega NM, Prill RJ, Camacho DM, Allison
KR, Kellis M, Collins JJ, Stolovitzky G: **Wisdom of crowds for robust gene**
network inference. *Nature methods* 2012, **9**(8):796-804.

10.   Badis G, Berger MF, Philippakis AA, Talukder S, Gehrke AR, Jaeger SA, Chan
ET, Metzler G, Vedenko A, Chen X *et al*: **Diversity and complexity in DNA**
recognition by transcription factors. *Science (New York, NY)* 2009,
324(5935):1720-1723.

11.   McLean CY, Bristor D, Hiller M, Clarke SL, Schaar BT, Lowe CB, Wenger AM,
Bejerano G: **GREAT improves functional interpretation of cis-regulatory**
regions. *Nature biotechnology* 2010, **28**(5):495-501.

12.   Wendt KS, Yoshida K, Itoh T, Bando M, Koch B, Schirghuber E, Tsutsumi S,
Nagae G, Ishihara K, Mishiro T *et al*: **Cohesin mediates transcriptional**
insulation by CCCTC-binding factor. *Nature* 2008, **451**(7180):796-801.

13.   Guo Y, Xu Q, Canzio D, Shou J, Li J, Gorkin DU, Jung I, Wu H, Zhai Y, Tang Y
et al: **CRISPR Inversion of CTCF Sites Alters Genome Topology and**
Enhancer/Promoter Function. *Cell* 2015, **162**(4):900-910.

14.   Gusmao EG, Allhoff M, Zenke M, Costa IG: **Analysis of computational**
footprinting methods for DNase sequencing experiments. *Nature methods*
2016, **13**(4):303-309.

15.   Thomas R, Thomas S, Holloway AK, Pollard KS: **Features that define the best**
ChIP-seq peak calling algorithms. *Brief Bioinform* 2016.

16.   Gusmao EG, Dieterich C, Zenke M, Costa IG: **Detection of active**
transcription factor binding sites with the combination of DNase
hypersensitivity and histone modifications. *Bioinformatics (Oxford, England)*
2014, **30**(22):3143-3151.

17.   He B, Chen C, Teng L, Tan K: **Global view of enhancer-promoter**
interactome in human cells. *Proc Natl Acad Sci U S A* 2014,

111(21):E2191-2199.

18.   Wang D, Rendon A, Wernisch L: **Transcription factor and chromatin features**
predict genes associated with eQTLs. *Nucleic acids research* 2013,
41(3):1450-1463.

19.   Whalen S, Truty RM, Pollard KS: **Enhancer-promoter interactions are**
encoded by complex genomic signatures on looping chromatin. *Nature*
genetics 2016, **48**(5):488-496.

20.   Mifsud B, Tavares-Cadete F, Young AN, Sugar R, Schoenfelder S, Ferreira L,
Wingett SW, Andrews S, Grey W, Ewels PA *et al*: **Mapping long-range**
promoter contacts in human cells with high-resolution capture Hi-C. *Nature*
genetics 2015, **47**(6):598-606.

21.   Javierre BM, Burren OS, Wilder SP, Kreuzhuber R, Hill SM, Sewitz S, Cairns J,
Wingett SW, Varnai C, Thiecke MJ *et al*: **Lineage-Specific Genome**
Architecture Links Enhancers and Non-coding Disease Variants to Target
Gene Promoters. *Cell* 2016, **167**(5):1369-1384 e1319.

22.   Bailey SD, Zhang X, Desai K, Aid M, Corradin O, Cowper-Sal Lari R,
Akhtar-Zaidi B, Scacheri PC, Haibe-Kains B, Lupien M: **ZNF143 provides**
sequence specificity to secure chromatin interactions at gene promoters. *Nat*
Commun 2015, **2**:6186.

23.   Wingender E, Dietze P, Karas H, Knuppel R: **TRANSFAC: a database on**
transcription factors and their DNA binding sites. *Nucleic acids research*
1996, **24**(1):238-241.

24.   Portales-Casamar E, Thongjuea S, Kwon AT, Arenillas D, Zhao X, Valen E,
Yusuf D, Lenhard B, Wasserman WW, Sandelin A: **JASPAR 2010: the greatly**
expanded open-access database of transcription factor binding profiles.
Nucleic acids research 2010, **38**(Database issue):D105-110.

25.   Newburger DE, Bulyk ML: **UniPROBE: an online database of protein**
binding microarray data on protein-DNA interactions. *Nucleic acids*
research 2009, **37**(Database issue):D77-82.

26.   Zhao Y, Stormo GD: **Quantitative analysis demonstrates most transcription**
factors require only simple models of specificity. *Nature biotechnology* 2011,
29(6):480-483.

27.   Jolma A, Yan J, Whitington T, Toivonen J, Nitta KR, Rastas P, Morgunova E,

Enge M, Taipale M, Wei G *et al*: **DNA-binding specificities of human**
transcription factors. *Cell* 2013, **152**(1-2):327-339.

28.   Xie Z, Hu S, Blackshaw S, Zhu H, Qian J: **hPDI: a database of experimental**
human protein-DNA interactions. *Bioinformatics (Oxford, England)* 2010,
26(2):287-289.

29.   Kheradpour P, Kellis M: **Systematic discovery and characterization of**
regulatory motifs in ENCODE TF binding experiments. *Nucleic acids*
research 2014, **42**(5):2976-2987.

30.   Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, Ren J, Li WW,
Noble WS: **MEME SUITE: tools for motif discovery and searching**. *Nucleic*
acids research 2009, **37**(Web Server issue):W202-208.

31.   Grant CE, Bailey TL, Noble WS: **FIMO: scanning for occurrences of a given**
motif. *Bioinformatics (Oxford, England)* 2011, **27**(7):1017-1018.

32.   Karolchik D, Barber GP, Casper J, Clawson H, Cline MS, Diekhans M, Dreszer
TR, Fujita PA, Guruvadoo L, Haeussler M *et al*: **The UCSC Genome Browser**
database: 2014 update. *Nucleic acids research* 2014, **42**(Database
issue):D764-770.

33.   Wang L, Wang S, Li W: **RSeQC: quality control of RNA-seq experiments**.
Bioinformatics (Oxford, England) 2012, **28**(16):2184-2185.

34.   Saeed S, Quintin J, Kerstens HH, Rao NA, Aghajanirefah A, Matarese F, Cheng
SC, Ratter J, Berentsen K, van der Ent MA *et al*: **Epigenetic programming of**
monocyte-to-macrophage differentiation and trained innate immunity.
Science (New York, NY) 2014, **345**(6204):1251086.

35.   Zambon AC, Gaj S, Ho I, Hanspers K, Vranizan K, Evelo CT, Conklin BR, Pico
AR, Salomonis N: **GO-Elite: a flexible solution for pathway and ontology**
over-representation. *Bioinformatics (Oxford, England)* 2012,
28(16):2209-2210.

36.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis
G, Durbin R, Genome Project Data Processing S: **The Sequence**
Alignment/Map format and SAMtools. *Bioinformatics (Oxford, England)*
2009, **25**(16):2078-2079.

37.   Quinlan AR, Hall IM: **BEDTools: a flexible suite of utilities for comparing**
genomic features. *Bioinformatics (Oxford, England)* 2010, **26**(6):841-842.

38.   Zang C, Schones DE, Zeng C, Cui K, Zhao K, Peng W: **A clustering approach**
for identification of enriched domains from histone modification ChIP-Seq
data. *Bioinformatics (Oxford, England)* 2009, **25**(15):1952-1958.

39.   Kraaij MD, Vereyken EJ, Leenen PJ, van den Bosch TP, Rezaee F, Betjes MG,
Baan CC, Rowshani AT: **Human monocytes produce interferon-gamma upon**
stimulation with LPS. *Cytokine* 2014, **67**(1):7-12.

40.   Ebner S, Hofer S, Nguyen VA, Furhapter C, Herold M, Fritsch P, Heufler C,
Romani N: **A novel role for IL-3: human monocytes cultured in the presence**
of IL-3 and IL-4 differentiate into dendritic cells that produce less IL-12
and shift Th cell responses toward a Th2 cytokine pattern. *J Immunol* 2002,
168(12):6199-6207.

41.   Francisco-Cruz A, Aguilar-Santelises M, Ramos-Espinosa O, Mata-Espinosa D,
Marquina-Castillo B, Barrios-Payan J, Hernandez-Pando R:
Granulocyte-macrophage colony-stimulating factor: not just another
haematopoietic growth factor. *Med Oncol* 2014, **31**(1):774.

42.   Shaw RJ, Doherty DE, Ritter AG, Benedict SH, Clark RA:
Adherence-dependent increase in human monocyte PDGF(B) mRNA is
associated with increases in c-fos, c-jun, and EGR2 mRNA. *J Cell Biol* 1990,
111(5 Pt 1):2139-2148.

43.   Heil M, Clauss M, Suzuki K, Buschmann IR, Willuweit A, Fischer S, Schaper
W: **Vascular endothelial growth factor (VEGF) stimulates monocyte**
migration through endothelial monolayers via increased integrin expression.
Eur J Cell Biol 2000, **79**(11):850-857.

44.   Hogg N, Laschinger M, Giles K, McDowall A: **T-cell integrins: more than just**
sticking points. *J Cell Sci* 2003, **116**(Pt 23):4695-4705.

45.   Ngai P, McCormick S, Small C, Zhang X, Zganiacz A, Aoki N, Xing Z: **Gamma**
interferon responses of CD4 and CD8 T-cell subsets are quantitatively
different and independent of each other during pulmonary Mycobacterium
bovis BCG infection. *Infect Immun* 2007, **75**(5):2244-2252.

46.   MacIver NJ, Blagih J, Saucillo DC, Tonelli L, Griss T, Rathmell JC, Jones RG:
The liver kinase B1 is a central regulator of T cell development, activation,
and metabolism. *J Immunol* 2011, **187**(8):4187-4198.

47.   Gallegos AM, Xiong H, Leiner IM, Susac B, Glickman MS, Pamer EG, van

Heijst JW: **Control of T cell antigen reactivity via programmed TCR**
downregulation. *Nature immunology* 2016, **17**(4):379-386.

48.   Kronin V, Hochrein H, Shortman K, Kelso A: **Regulation of T cell cytokine**
production by dendritic cells. *Immunol Cell Biol* 2000, **78**(3):214-223.

49.   Lu TT, Cyster JG: **Integrin-mediated long-term B cell retention in the splenic**
marginal zone. *Science (New York, NY)* 2002, **297**(5580):409-412.

50.   Baumann MA, Paul CC: **Interleukin-5 and human B lymphocytes**. *Methods*
1997, **11**(1):88-97.

51.   Winer DA, Winer S, Shen L, Wadia PP, Yantha J, Paltser G, Tsui H, Wu P,
Davidson MG, Alonso MN *et al*: **B cells promote insulin resistance through**
modulation of T cells and production of pathogenic IgG antibodies. *Nat Med*
2011, **17**(5):610-617.

52.   Limon JJ, Fruman DA: **Akt and mTOR in B Cell Activation and**
Differentiation. *Front Immunol* 2012, **3**:228.

53.   Sorrentino R, Bertolino A, Terlizzi M, Iacono VM, Maiolino P, Cirino G,
Roviezzo F, Pinto A: **B cell depletion increases**
sphingosine-1-phosphate-dependent airway inflammation in mice. *Am J*
Respir Cell Mol Biol 2015, **52**(5):571-583.

54.   Caulfield J, Fernandez M, Snetkov V, Lee T, Hawrylowicz C: **CXCR4**
expression on monocytes is up-regulated by dexamethasone and is
modulated by autologous CD3+ T cells. *Immunology* 2002, **105**(2):155-162.

55.   Kral JB, Schrottmaier WC, Salzmann M, Assinger A: **Platelet Interaction with**
Innate Immune Cells. *Transfus Med Hemother* 2016, **43**(2):78-88.

56.   Heinonen KM, Bourdeau A, Doody KM, Tremblay ML: **Protein tyrosine**
phosphatases PTP-1B and TC-PTP play nonredundant roles in macrophage
development and IFN-gamma signaling. *Proc Natl Acad Sci U S A* 2009,
106(23):9368-9372.

57.   Liu HS, Pan CE, Liu QG, Yang W, Liu XM: **Effect of NF-kappaB and p38**
MAPK in activated monocytes/macrophages on pro-inflammatory
cytokines of rats with acute pancreatitis. *World J Gastroenterol* 2003,
9(11):2513-2518.

58.   Bonder CS, Finlay-Jones JJ, Hart PH: **Interleukin-4 regulation of human**
monocyte and macrophage interleukin-10 and interleukin-12 production.

Role of a functional interleukin-2 receptor gamma-chain. *Immunology* 1999,
96(4):529-536.

59. Murakami T, Nakajima T, Koyanagi Y, Tachibana K, Fujii N, Tamamura H,
Yoshida N, Waki M, Matsumoto A, Yoshie O *et al*: **A small molecule CXCR4**
inhibitor that blocks T cell line-tropic HIV-1 infection. *J Exp Med* 1997,
186(8):1389-1393.

60. Koyasu S, D'Adamio L, Arulanandam AR, Abraham S, Clayton LK, Reinherz
EL: **T cell receptor complexes containing Fc epsilon RI gamma homodimers**
in lieu of CD3 zeta and CD3 eta components: a novel isoform expressed on
large granular lymphocytes. *J Exp Med* 1992, **175**(1):203-209.

61. Dong C, Yang DD, Wysk M, Whitmarsh AJ, Davis RJ, Flavell RA: **Defective T**
cell differentiation in the absence of Jnk1. *Science (New York, NY)* 1998,
282(5396):2092-2095.

62. Huang Y, Clarke F, Karimi M, Roy NH, Williamson EK, Okumura M,
Mochizuki K, Chen EJ, Park TJ, Debes GF *et al*: **CRK proteins selectively**
regulate T cell migration into inflamed tissues. *J Clin Invest* 2015,
125(3):1019-1032.

63. Kumanogoh A, Marukawa S, Suzuki K, Takegahara N, Watanabe C, Ch'ng E,
Ishida I, Fujimura H, Sakoda S, Yoshida K *et al*: **Class IV semaphorin Sema4A**
enhances T-cell activation and interacts with Tim-2. *Nature* 2002,
419(6907):629-633.

64. Endo T, Ito K, Morimoto J, Kanayama M, Ota D, Ikesue M, Kon S, Takahashi D,
Onodera T, Iwasaki N *et al*: **Syndecan 4 Regulation of the Development of**
Autoimmune Arthritis in Mice by Modulating B Cell Migration and
Germinal Center Formation. *Arthritis Rheumatol* 2015, **67**(9):2512-2522.

65. Beider K, Ribakovsky E, Abraham M, Wald H, Weiss L, Rosenberg E, Galun E,
Avigdor A, Eizenberg O, Peled A *et al*: **Targeting the CD20 and CXCR4**
pathways in non-hodgkin lymphoma with rituximab and high-affinity
CXCR4 antagonist BKT140. *Clin Cancer Res* 2013, **19**(13):3495-3507.

66. Kimata H, Yoshida A, Ishioka C, Masuda S, Sasaki R, Mikawa H: **Human**
recombinant erythropoietin directly stimulates B cell immunoglobulin
production and proliferation in serum-free medium. *Clin Exp Immunol* 1991,
85(1):151-156.

67. Koizumi M, Hiasa Y, Kumagi T, Yamanishi H, Azemoto N, Kobata T, Matsuura
B, Abe M, Onji M: **Increased B cell-activating factor promotes tumor**
invasion and metastasis in human pancreatic cancer. *PLoS One* 2013,
8(8):e71367.

68. Rosser EC, Oleinika K, Tonon S, Doyle R, Bosma A, Carter NA, Harris KA,
Jones SA, Klein N, Mauri C: **Regulatory B cells are induced by gut**
microbiota-driven interleukin-1beta and interleukin-6 production. *Nat Med*
2014, **20**(11):1334-1339.

69. de Wit E, Vos ES, Holwerda SJ, Valdes-Quezada C, Verstegen MJ, Teunissen H,
Splinter E, Wijchers PJ, Krijger PH, de Laat W: **CTCF Binding Polarity**
Determines Chromatin Looping. *Molecular cell* 2015, **60**(4):676-684.

Figure

868
869

Figure 1. Analyses of functional enrichments of putative transcriptional target

genes. Transcriptional target genes were predicted using open chromatin regions

(DNase-DGF) and known transcription factor binding sequences. Functional

enrichments of target genes were analyzed using 10 annotation databases, and were

changed based on the criteria of promoter and extended regions for enhancer-promoter

association (EPA). To compare with the tendency of the normalized numbers of

functional enrichments, the median expression levels of target genes were examined

using promoter and EPA.

a

GO (Gene Ontology)

* Wilcoxon signed-rank test $p < 10^{-2}$

Legend: Monocyte, CD4+ T cell, CD20+ B cell, H1-hESC, iPSC

Ratio of putative transcriptional target genes replaced with randomly selected genes

Ratio of randomly selected genes added to putative transcriptional target genes

b

Wilcoxon signed-rank test $p < 10^{-2}$

Legend: Pathway Commons, KEGG, GO, TF Targets, CTD Ontology, GO Slim, BioMarkers, Domains, WikiPathways

Normalized number of functional enrichments (Annotation databases except for GO)

Normalized number of functional enrichments (GO)

Ratio of putative transcriptional target genes replaced with randomly selected genes

Ratio of randomly selected genes added to putative transcriptional target genes

Transcription factors (TF)

Putative transcriptional target genes

Randomly selected genes

Figure 2. Effect of randomly selected genes on functional enrichments. (a) Effect of

randomly selected genes on functional enrichments using DNase-DGF data.

Transcriptional target genes were predicted using DNase-DGF data in human

monocytes, $CD4^+$ T, $CD20^+$ B, other four somatic and two stem cells (H1-hESC and

iPSC) (see also Figure S1). To test whether slight changes of transcriptional target genes

were reflected in the normalized number of their functional enrichments, the ratio of

randomly selected genes in the target genes of each TF was changed between 5% and

60%. In the left part of the graphs, randomly selected genes were replaced with the

target genes where the total number of target genes was unchanged. In the right part of

the graphs, randomly selected genes were added to the target genes where the total

number of target genes was increased. The result of Gene Ontology annotation was

shown. The results of Pathway Commons and KEGG were shown in Figure S1. Native

target genes showed the most functional enrichments in most cell types. **(b)** Effect of

randomly selected genes on functional enrichments using ChIP-seq data. Transcriptional

target genes were predicted using ChIP-seq data of 19 TF in H1-hESC. The results of

nine functional annotation databases were shown and the result of target genes of

microRNAs was shown in Figure S2. Native target genes showed the most functional

enrichments using most annotation databases except for low frequent functional

annotations. Putative transcriptional target genes tend to include similar function of

genes.

Figure 3. Criteria of promoter and extended regions for enhancer-promoter

association and features of chromatin interactions. (a) Computationally-defined

regulatory domains [11]. The transcription start site (TSS) of each gene is indicated as

an arrow. The corresponding regulatory domain for each gene is shown in a matching

color as a bracketed line. The *basal plus extension* association rule assigns a basal

regulatory domain to each gene regardless of genes nearby (thick line, Association rule

1 and 2) (see Methods). The domain is then extended to the basal regulatory domain of

the nearest upstream and downstream genes. The *two nearest genes* association rule

extends the regulatory domain to the TSS of the nearest upstream and downstream

genes (Association rule 3). The *single nearest gene* association rule extends the

regulatory domain to the midpoint between this gene's TSS and the nearest gene's TSS

both upstream and downstream (Association rule 4). (**b**) Forward–reverse orientation of

CTCF-binding sites are frequently found in chromatin interactions. CTCF can block the

interaction between enhancers and promoters limiting the activity of enhancers to

certain functional domains. Figures adapted from [13, 69] and the bottom figure in

panel b was in the book 'THE CELL: A molecular approach 6e, Figure 7.21, 2013

Sinauer   Associates,   Inc'   and   Image   68   insulators   in

'http://slideplayer.com/slide/3836520'.

Figure 4. Comparison of the median expression levels of transcriptional target

genes. The median expression levels of the target genes of the same transcription factor

binding sequences were compared between promoters and three types of promoter and

extended regions for enhancer-promoter association (EPA). Red and blue dots show

statistically significant difference of the distribution of expression levels of target genes

between promoters and EPA. Red dots show the median expression level of target genes

was higher in EPA than promoters, and blue dots show the median expression level of

target genes was lower in EPA than promoters. The median expression levels of putative

transcriptional target genes were significantly lower using TF bound in enhancers than

using TF bound in promoters in stem cells such as H1-hESC and iPSC, and higher by

TF bound in enhancers in immune cells. These results implied that enhancers

significantly affect the expression of target genes.

Tables

Tables are attached in separate PDF files (Table 1 – 6) in Attachment folder, because the

tables are colored to understand them easily and statistically different numbers are

shown with lines and asterisks.

Table 1.

Table 1. Number of functional enrichments and unique functional enrichemnts of putative transcriptional target genes.

Number of functional enrichments of putative trancriptional target genes

	KEGG	TF Targets	CTD Ontology	GO Slim	GO	Pathway Commons	BioMarkers	MicroRNA	Domains	WikiPathways
Monocyte	349	107	209	114	2,902	1,005	42	451	1,202	242
CD4+ T cell	317	135	278	77	4,077	1,806	47	754	1,401	405
CD20+ B cell	323	103	170	88	2,778	821	39	948	950	288

Number of unique functional enrichments of gene expression information alone and putative transcriptional target genes.
* Wilcoxon signed-rank test *p* < 0.01 (except for MicroRNA and Domains)

Gene expression information alone										
Monocyte	43	0	35	11	237	101	7	314	404	58
CD4+ T cell	47	0	19	9	301	165	9	136	397	81
CD20+ B cell	42	0	27	12	239	247	6	370	409	65
Putative transcriptional target genes										
Monocyte	95	16	127	12	1,271	242	17	97	303	105
CD4+ T cell	105	26	146	23	1,654	415	24	224	585	133
CD20+ B cell	93	23	96	23	1,192	329	16	231	397	106

Table 2

Table 2. Functional enrichments of putative transcriptional target genes using Pathway Commons

Monocyte - Pathway Commons	No. of TFs
Proteoglycan syndecan-mediated signaling events	18
Regulation of CDC42 activity	15
LKB1 signaling events	14
Glypican pathway	13
Interferon Signaling	13
Sphingosine 1-phosphate (S1P) pathway	13
IL5-mediated signaling events	12
Syndecan-1-mediated signaling events	12
IL3-mediated signaling events	11
Mitotic Prophase	11
Golgi Cisternae Pericentriolar Stack Reorganization	11
Interferon alpha/beta signaling	11
IFN-gamma pathway	10
Signaling events mediated by Hepatocyte Growth Factor Receptor (c-Met)	10
Recruitment of mitotic centrosome proteins and complexes	10

CD4+ T cell - Pathway Commons	No. of TFs
TCR signaling in naive CD8+ T cells	36
IL12-mediated signaling events	24
Downstream signaling in naive CD8+ T cells	21
TCR signaling in naive CD4+ T cells	21
IL12 signaling mediated by STAT4	20
Validated transcriptional targets of AP1 family members Fra1 and Fra2	19
CXCR4-mediated signaling events	17
ATF-2 transcription factor network	15
Thrombin/protease-activated receptor (PAR) pathway	14
TCR signaling	14
PAR1-mediated thrombin signaling events	14
Downstream TCR signaling	14
Internalization of ErbB1	13
Urokinase-type plasminogen activator (uPA) and uPAR-mediated signaling	13
ErbB receptor signaling network	13

CD20+ B cell - Pathway Commons	No. of TFs
Interferon alpha/beta signaling	12
Alpha6Beta4Integrin	11
Validated targets of C-MYC transcriptional activation	11
IL8- and CXCR2-mediated signaling events	10
Antigen processing-Cross presentation	9
BCR signaling pathway	9
IL6-mediated signaling events	9
Cell junction organization	9
ER-Phagosome pathway	8
Regulation of CDC42 activity	8
CXCR4-mediated signaling events	8
CDC42 signaling events	8
Syndecan-4-mediated signaling events	8
Noncanonical Wnt signaling pathway	7
Class I MHC mediated antigen processing & presentation	7

Table 3. Normalized number of functional enrichments of putative transcriptional target genes using promoter and extended regions for enhancer-promoter association.
* Wilcoxon signed-rank test *p* < 0.05

	KEGG	CTD Ontology	GO Slim	GO	Pathway Commons	Domains	Wiki Pathways
Monocyte							
Promoter	8.46	5.07	2.76	70.34	24.36	29.13	5.87
Association rule 1	10.44	6.48	2.01	133.54	44.75	42.19	11.10
Association rule 2	9.03	6.29	1.45	125.13	38.69	40.17	8.30
Association rule 3	8.06	5.29	1.37	106.60	24.25	38.96	7.62
Association rule 4	11.47	8.22	2.46	164.18	71.42	47.85	12.78
CD4⁺ T cell							
Promoter	5.30	4.64	1.29	68.11	30.17	23.41	6.77
Association rule 1	13.60	7.07	2.74	142.40	84.51	43.78	13.65
Association rule 2	13.57	6.69	3.05	141.15	86.36	46.33	12.02
Association rule 3	12.40	5.89	2.50	115.76	68.84	41.85	10.00
Association rule 4	16.40	7.86	4.03	177.55	108.08	53.86	16.72
CD20⁺ B cell							
Promoter	7.02	3.70	1.91	60.39	17.85	20.65	6.26
Association rule 1	8.88	6.21	2.59	104.55	59.32	34.34	8.29
Association rule 2	8.60	5.32	1.55	105.34	57.31	38.05	9.95
Association rule 3	9.01	5.28	1.42	88.85	26.49	35.17	8.26
Association rule 4	9.95	6.62	3.07	134.46	90.99	41.30	10.67
HUVEC							
Promoter	7.50	5.74	1.34	79.71	22.21	22.40	4.59
Association rule 1	11.62	8.95	3.74	160.52	37.11	65.15	8.88
Association rule 2	11.87	8.77	4.76	176.26	36.86	74.71	8.66
Association rule 3	9.84	7.70	4.55	154.06	34.29	73.98	7.74
Association rule 4	13.46	8.99	5.08	191.73	59.10	82.65	10.13
IMR90							
Promoter	6.63	5.21	2.43	68.31	11.64	25.50	5.47
Association rule 1	9.63	6.78	2.85	122.38	24.89	65.39	8.03
Association rule 2	9.32	6.56	1.43	115.31	16.36	60.36	7.47
Association rule 3	9.59	5.63	1.55	99.10	15.39	54.82	5.82
Association rule 4	10.55	7.68	3.76	142.10	30.05	74.86	8.78
MCF-7							
Promoter	6.97	5.43	1.69	77.88	24.71	25.99	8.77
Association rule 1	9.45	6.26	2.03	121.84	22.38	55.19	9.86
Association rule 2	9.44	6.18	0.89	111.77	21.59	53.90	7.97
Association rule 3	9.83	5.13	1.12	100.35	18.96	55.30	7.37
Association rule 4	10.80	7.38	2.77	149.64	31.20	62.40	11.07
HMEC							
Promoter	8.76	5.03	1.53	65.80	14.07	26.52	6.70
Association rule 1	10.88	7.04	1.96	138.27	18.28	72.55	8.08
Association rule 2	9.22	6.31	1.63	141.57	18.84	73.09	7.99
Association rule 3	9.30	6.50	1.68	123.46	17.27	64.67	7.75
Association rule 4	11.78	7.53	3.48	166.71	23.02	90.11	10.10
H1-hESC							
Promoter	8.45	3.47	2.49	73.33	14.12	24.35	4.84
Association rule 1	9.54	4.82	1.97	113.37	21.86	37.66	6.30
Association rule 2	10.76	5.82	1.69	123.45	21.27	38.26	5.35
Association rule 3	8.00	4.84	1.43	101.12	18.00	33.59	5.62
Association rule 4	10.56	5.37	3.27	141.38	35.12	46.16	6.73
iPSC							
Promoter	8.64	2.05	1.58	67.32	22.75	12.36	6.42
Association rule 1	10.25	5.64	1.66	107.50	18.35	26.77	8.59
Association rule 2	11.70	6.20	1.01	101.76	17.00	28.13	8.16
Association rule 3	11.69	6.89	1.01	90.85	16.60	24.89	7.76
Association rule 4	11.39	7.21	2.53	136.46	23.60	31.49	9.27

Z-Score scale: −2, −1, 0, 1, 2

954   Table 4

Table 4. Differences in functional enrichments using Pathway Commons. Immune cell-related functional annotations were enriched more in 'promoter and extended regions for enhancer-promoter association (EPA)' than promoters. Five annotations are shown in each cell type. These functions are confirmed to be related to the cellular functions by reference journal papers.

Cell type	Term of functional annotation	No. of TF EPA (Association rule 4)	No. of TF Promoter	P-value	Experimental support
Monocytes	IFN-gamma pathway	51	10	6.89×10^{-4}	Ref. 39
	IL3-mediated signaling events	46	11	4.69×10^{-3}	Ref. 40
	GMCSF-mediated signaling events	43	10	5.25×10^{-3}	Ref. 41
	PDGF receptor signaling network	43	8	1.34×10^{-2}	Ref. 42
	VEGF and VEGFR signaling network	44	9	2.10×10^{-3}	Ref. 43
CD4[+] T cell	Integrin family cell surface interactions	112	13	2.59×10^{-12}	Ref. 44
	IFN-gamma pathway	107	12	5.26×10^{-12}	Ref. 45
	LKB1 signaling events	107	11	1.96×10^{-12}	Ref. 46
	TCR signaling in naive CD4[+] T cells	103	21	3.98×10^{-8}	Ref. 47
	IL3-mediated signaling events	100	8	9.54×10^{-13}	Ref. 48
CD20[+] B cell	Integrin family cell surface interactions	69	2	5.17×10^{-11}	Ref. 49
	IL5-mediated signaling events	64	0	2.21×10^{-11}	Ref. 50
	Insulin Pathway	63	0	3.15×10^{-11}	Ref. 51
	mTOR signaling pathway	63	0	3.15×10^{-11}	Ref. 52
	Sphingosine 1-phosphate (S1P) pathway	63	0	3.15×10^{-11}	Ref. 53

955   Chi-square tests were conducted using the total number of putative transcriptional target genes (Table S9).

Table 5

Table. 5 Normalized number of functional enrichments of putative transcriptional target genes using CTCF binding sites. * Wilcoxon signed-rank test $p < 0.05$

	KEGG	CTD Ontology	GO Slim	GO	Pathway Commons	Domains	Wiki Pathways
Monocytes							
Association rule 4	11.47	8.22	2.46	164.18	71.42	47.85	12.78
CTCF (FR+RF+FF+RR)	13.19	10.39	2.74	134.26	34.37	44.46	12.96
CTCF (FR)	42.92	19.53	5.66	509.86	196.58	112.14	35.11
CD4⁺ T cells							
Association rule 4	16.40	7.86	4.03	177.55	108.08	53.86	16.72
CTCF (FR+RF+FF+RR)	26.33	8.73	5.11	206.05	130.71	57.53	23.56
CTCF (FR)	69.39	14.66	24.91	560.44	220.54	133.26	46.54
CD20⁺ B cells							
Association rule 4	9.95	6.62	3.07	134.46	90.99	41.30	10.67
CTCF (FR+RF+FF+RR)	8.78	4.22	2.78	94.13	27.70	28.55	7.08
CTCF (FR)	28.86	9.72	6.61	304.01	220.77	99.89	22.68
HUVEC							
Association rule 4	13.46	8.99	5.08	191.73	59.10	82.65	10.13
CTCF (FR+RF+FF+RR)	19.35	10.59	3.85	155.11	28.36	57.87	13.58
CTCF (FR)	33.71	14.82	16.33	454.47	61.85	163.38	29.38
IMR90							
Association rule 4	10.55	7.68	3.76	142.10	30.05	74.86	8.78
CTCF (FR+RF+FF+RR)	13.88	17.55	9.22	252.24	39.92	123.18	22.45
CTCF (FR)	30.22	15.89	11.34	402.40	53.56	180.71	37.42
MCF-7							
Association rule 4	10.80	7.38	2.77	149.64	31.20	62.40	11.07
CTCF (FR+RF+FF+RR)	11.39	8.30	4.92	234.48	30.40	126.70	13.99
CTCF (FR)	22.34	11.48	6.09	397.69	51.39	145.43	37.58
HMEC							
Association rule 4	11.78	7.53	3.48	166.71	23.02	90.11	10.10
CTCF (FR+RF+FF+RR)	19.68	13.54	4.29	232.47	27.97	106.24	9.84
CTCF (FR)	29.59	14.46	6.97	390.45	43.90	124.81	30.78
H1-hESC							
Association rule 4	10.56	5.37	3.27	141.38	35.12	46.16	6.73
CTCF (FR+RF+FF+RR)	12.95	5.37	2.45	131.46	29.57	44.15	11.08
CTCF (FR)	28.21	10.55	9.02	303.41	56.95	139.44	19.72
iPSC							
Association rule 4	11.39	7.21	2.53	136.46	23.60	31.49	9.27
CTCF (FR+RF+FF+RR)	17.17	2.33	1.69	120.87	25.60	21.06	6.42
CTCF (FR)	36.94	6.09	6.86	274.30	31.37	83.40	16.80

Z-Score legend: -2 -1 0 1 2

960   Table 6

Table 6. Differences in functional enrichments using Pathway Commons. Immune cell-related functional annotations were enriched more in EPA shortened at forward–reverse orientation of CTCF-binding sites than EPA without CTCF. Five annotations are shown in each cell type. These functions are confirmed to be related to the cellular functions by reference journal papers.

Cell type	Term of functional annotation	No. of TF EPA (Association rule 4) shortened by FR CTCF	No. of TF EPA (Association rule 4) without CTCF	P-value	Experimental support
Monocytes	CXCR4-mediated signaling events	33	14	7.81×10^{-12}	Ref. 54
	GPVI-mediated activation cascade	26	3	8.97×10^{-15}	Ref. 55
	Signaling events mediated by TCPTP	23	10	1.46×10^{-8}	Ref. 56
	p38 MAPK signaling pathway	16	8	6.97×10^{-6}	Ref. 57
	IL4-mediated signaling events	11	3	8.80×10^{-6}	Ref. 58
CD4[+] T cell	CXCR4-mediated signaling events	181	82	$< 2.20 \times 10^{-16}$	Ref. 59
	Fc-epsilon receptor I signaling in mast cells	85	25	$< 2.20 \times 10^{-16}$	Ref. 60
	JNK signaling in the CD4+ TCR pathway	65	18	$< 2.20 \times 10^{-16}$	Ref. 61
	Reelin signaling pathway	62	26	$< 2.20 \times 10^{-16}$	Ref. 62
	Semaphorin interactions	51	14	$< 2.20 \times 10^{-16}$	Ref. 63
CD20[+] B cell	Syndecan-4-mediated signaling events	85	40	$< 2.20 \times 10^{-16}$	Ref. 64
	CXCR4-mediated signaling events	77	15	$< 2.20 \times 10^{-16}$	Ref. 65
	EPO signaling pathway	46	19	4.50×10^{-15}	Ref. 66
	Stabilization and expansion of the E-cadherin adherens junction	32	16	1.06×10^{-9}	Ref. 67
	Interleukin-1 signaling	13	1	3.98×10^{-8}	Ref. 68

961   Chi-square tests were conducted using the total number of putative transcriptional target genes (Table S12).