Neuronal components of evaluating the human origin of abstract shapes

Hagar Goldberg[1], Yuval Hart[2], Avraham E Mayo[3,4], Uri Alon[3,4] and Rafael Malach[1]

• Department of Neurobiology, Weizmann Institute of Science, Rehovot, Israel

• School of Engineering and Applied Sciences, Harvard University, Cambridge, MA USA

• Department of Molecular Cell Biology, Weizmann Institute of Science, Rehovot, Israel

• The Theatre Lab, Weizmann Institute of Science, Rehovot, Israel

Corresponding author. Fax: +972 8 9344131. E-mail addresses: rafi.malach@weizmann.ac.il, rafi.malach@gmail.com (R. Malach).

Abstract

Communication through visual symbols is a key aspect of human culture. However, to what

extent can people distinguish between human-origin and artificial symbols, and the neuronal

mechanisms underlying this process are not clear. Using fMRI we contrasted brain activity

during presentation of human-created abstract shapes and random-algorithm created shapes,

both sharing similar low level features.

We found that participants correctly identified most shapes as *human* or *random*. The lateral

occipital complex (LOC) was the main brain region showing preference to human-made shapes,

independently of task. Furthermore, LOC activity was parametrically correlated to beauty and

familiarity scores of the shapes (rated following the scan). Finally, a model classifier based only

on LOC activity showed human level accuracy at discriminating between human-made and

randomly-made shapes.

Our results highlight the sensitivity of the human brain to social and cultural cues, and point to

high-order object areas as central nodes underlying this capacity.

Introduction:

Humans are social and tuned to social cues. They therefore need to distinguish true social cues

from other stimuli. Previous studies have shown that human observers can distinguish biological

motion from random motion, even in impoverished stimuli such as point-light displays

(Johansson, 1973). This is suggested to be an intrinsic ability of the visual system, which has

evolved to preferentially attend to other humans, as shown in newborn babies (Simion et al.,

2008). These findings suggest that essential information such as social recognition can be

derived from minimalistic dynamic displays. In addition to direct communication (through verbal

and body language) human culture has ways of indirect communication through visual symbols -

a conventional representation of concepts through scripts and art; starting from cave paintings

(Chauvet et al., 1996), and continuing nowadays with symbols such as emoticons.

Similarly to the way humans can distinguish human walk from random motion generated by spot

light displays, we hypothesize that humans can successfully recognize abstract shapes that have

been generated by other humans compared to similar shapes created randomly. We further

hypothesize that the neuronal mechanisms that underlie this capacity are likely to be ingrained in

core systems and hence independent of the task performed.

What could be the aspects of the shapes that underlie the ability of participants to categorize

them into *human* vs. *random*? The symbolic meaning of a shape and its familiarity can serve to

assess the shape's origin. Another relevant feature is the shape's aesthetic value, if indeed people

tend to create more beautiful shapes. Thus, in the present study we focused on beauty and

familiarity- i.e. iconicity - the sense that a shape represents a familiar icon. We also examined the

inverse of familiarity- i.e. "weirdness"- a subjective sense that a figure is strange and unfamiliar.

Both beauty and weirdness are intuitive and powerful yet subjective impressions which are

difficult to define. Substantial work has suggested beauty and weirdness carry evolutionary

advantages. For example, neuro-imaging study showed overlap in brain regions which function

both during processing of aesthetic artworks and during appraisal of evolutionary important

objects (e.g. attractive potential mates, or desirable food) (Brown et al., 2011). Visual artistic

representations of beauty (Kawabata and Zeki, 2004, Di Dio et al., 2007, Ishizu and Zeki, 2011,

Cattaneo et al., 2015, Vartanian and Goel, 2004, Lacey et al., 2011) or naturalistic stimuli

beyond the arts domain (Brown et al., 2011, Chatterjee et al., 2009, Kirk, 2008, Lacey et al.,

2011) were showed to be linked to experience of reward, pleasure, and attitudes to external

information (approach/withdrawal). Similarly, the concept of weirdness was attributed to signals

of danger and risk (Rotshtein et al., 2001). Sensing beauty and weirdness thus seem important for

human behavior and survival, but we have little understanding of their neuronal correlates.

Another line of work considered the contrast between artificial and familiar images in a number

of studies of the human visual cortex (e.g. Fourier descriptors, and scrambled images; Lerner et

al., 2002, Tsao et al., 2003, Aalto et al., 2002, Murray et al., 2002, Malach et al., 1995). Visual

objects such as face images were rendered bizarre by inverting internal face features (Rotshtein

et al., 2001). These studies showed an increased activity of high order object areas (lateral

occipital complex - LOC) to coherent objects whether they are familiar or unfamiliar (Malach et

al., 1995, Kanwisher et al., 1997, Kanwisher et al., 1996, Grill-Spector et al., 2001, Rotshtein et
al., 2001).

However, in previous work a control over low-level features was lacking. In fact, we are not
aware of systematic study that directly compared brain activation to human-generated vs.
randomly generated shapes constructed of similar low-level components. Furthermore the
contribution of beauty and familiarity to the ability to evaluate human origin of shapes has not
been explored directly in previous research.

To address these issues, we examined, using fMRI, human perception of simple shapes made by
other humans, compared to similar shapes generated by an artificial algorithm. We specifically
examined to what extent brain regions respond differentially to these two categories - and to
what extent the beauty, iconicity and weirdness dimensions contribute to this differentiation.
To this end we employed a novel design in which a large group of 101 people were asked to
create simple shapes and to rank them according to how appealing they were (interesting and
beautiful) (Noy et al., 2012). Additional shapes were generated by an artificial, random walk
algorithm sampling from the space of all possible shapes and excluding shapes that were
generated by human observers. The generated shape ensemble included a wide spectrum of
beauty and familiarity levels.

We found that participants which were unfamiliar with the shapes were able to successfully
distinguish between human and random origin shapes. Our brain imaging results show that LOC
activity was significantly higher for the human-made shapes compared to the *random* ones.
Furthermore, this differential activity was a combined result of a positive correlation to beauty
and iconicity (familiarity) and a negative correlation to weirdness (unfamiliarity) of the abstract
shapes regardless of task. Our results point to the high-order object related complex (LOC -
Malach et al., 1995) as a pivotal node in endowing human observers with the ability to recognize
shared symbolic meaning and distinguish human from artificially created shapes.

Results:
Here we aimed to study the behavioral and neuronal mechanisms of distinguishing whether an
abstract shape was created by a human from a given space of shapes or by an algorithm that
makes a random choice from the same space of shapes. We used a rich yet fully determined
space of shapes, made of ten contiguous squares. Shapes were built by either 101 human players
in a computer-shape-generation game (see methods and Noy et al., 2012) or a random choice

95 algorithm from the same space of shapes. Thus, we contrasted algorithm-created with human-

96 created abstract shapes having similar low-level features. The human-created shapes were further

97 subdivided into three groups of different appeal ratings (by their human creators, see Methods).

98 Subjects unfamiliar with the shapes underwent fMRI scanning while watching the shapes in a

99 block design. The shapes were presented in two different experiments. In the first, subjects

100 performed a color discrimination task (experiment and task 1), and in the second, a human vs.

101 random algorithm origin discrimination task (experiment and task 2). Following the scan,

102 subjects gave their subjective evaluations of beauty, weirdness and iconicity of the shapes (see

103 Methods for details).

105 <u>Behavioral results:</u> Subjects successfully distinguished between most human and random

106 creation shapes (Fig. 2a). Most categories were successfully classified by the subjects (85-90%

107 accuracy), except *not chosen* category which was correctly classified (as *human*) in only 30% of

108 the cases (chance level = 50%). This difference in success rate between *not chosen* category and

109 the other categories was significantly lower (Mann-Whitney U = 0, $n_1 = n_2 = 7$ p < 0.005, two-

110 tailed). The difference between all other categories was not significant. A bias-free signal

111 detection analysis indicated that subjects were able to reliably distinguish between human and

112 random made shapes (N = 13, mean d' = 2.03, SE = ±0.34). No difference in reaction times was

113 found between the four categories in both experiments.

114 Three subjective aspects of the shapes were examined; beauty, weirdness and iconicity.

115 Weirdness and iconicity scores complemented each other – while weirdness focused mainly on

116 the level of unfamiliarity of a shape, iconicity reflected the level of familiarity and distinct

117 meaning. The classification to human origin was best modeled by the interaction of the beauty

118 and weirdness scores rather than the two scores separated

119 ($P(human) = (1 + 13.7\,e^{-6.6\,WB})^{-1}$) (see Methods and SI). Beauty scores were lowest for

120 *random* and *not chosen* shapes, and were significantly higher for *chosen* and even higher for *top*

121 *rate* (Mann-Whitney between *random* and *not chosen*: U = 20, $n_1 = n_2 = 7$, p = 0.521, two-tailed;

122 between all other categories: U = 0, $n_1 = n_2 = 7$, p < 0.005, two-tailed. see Fig. 2b). Thus

123 indicating cross population consistency between the creators of the shapes and the scanned

124 participants.

Weirdness scores on the other hand distinguished between *human* and *random* categories; the
random category received the highest score, *not chosen* received a significantly lower score,
chosen and *top rated* received the lowest scores, significantly different from the first two (Mann-
Whitney between *chosen* and *top rated*: U = 0.531, $n_1 = n_2 = 7$, p = 0.08, two-tailed; between all
other categories: U = 1, $n_1 = n_2 = 7$, p < 0.005, two-tailed. see Fig. 2c). Thus, subjective beauty
was positively correlated, and weirdness was negatively correlated to classification as *human*.
Furthermore each parameter separated between different shape categories.
132
<u>Brain imaging results:</u> In order to examine a possible implicit differentiation between human-
made shapes and randomly made shapes, a direct contrast of BOLD activity (task 1) between
human blocks and *random* blocks was conducted. The contrast map (Fig. 3) revealed highly
localized preferential activations to *human* vs. *random* in the lateral occipital complex (LOC,
Malach et al., 1995). Preferential activation to *random* shapes was spread over a wider range of
the cortex, particularly in parietal and frontal regions. Its most significant activation was located
in the inferior parietal gyrus (IFG).
Cross-task ROI analysis demonstrated a consistently stronger response to *human* blocks relative
to *random* blocks in bilateral LOC in both tasks (paired t test; (experiment 1) left LOC, N = 17, t
= 2.74, p < 0.05; right LOC, N = 17, t = 3.92, p < 0.005; (experiment 2) left LOC, N = 15, t =
3,93, p < 0.005; right LOC, N = 15, t = 3.29, p < 0.05). ROI analysis inspecting the averaged
LOC activity (beta weight) per category revealed a gradual positive response in bilateral LOC
along the "appeal" axis (Fig 3c). *Random* shapes showed the lowest response in LOC, top rated
showed the highest response (followed by not chosen and chosen respectively) (One way
repeated measures ANOVA; (experiment 1) left LOC, N = 17, F = 2.751 not significant, Linear
trend: F = 5.174, p < 0.05; right LOC, N = 17, F = 4.35, p < 0.05, Linear trend: F = 7.974, p <
0.05; (experiment 2) left LOC, N = 15, F = 6.145, p < 0.05, Linear trend: F = 28.721, p < 0.0005;
right LOC, N = 15, F = 8.052, p < 0.0005, Linear trend: F = 26.281, p < 0.0005).
Since beauty, weirdness and iconicity scores were predictive for blocks classification
(*human/random*) we wanted to study their neuronal correlates. A whole brain parametric GLM
analysis was therefore conducted (Fig. 4). The results revealed a consistent parametric
relationship between LOC activity to beauty, weirdness and iconicity measures regardless of task
(color/ shape origin discrimination). Parametric brain maps for beauty scores show focused

activation in the LOC. However in frontal areas, weirdness and iconicity showed a task-related

selectivity: in task 1 frontal region such as dorsal premotor cortex and inferior frontal gyrus

(IFG) were parametrically correlated to weirdness (positive) and iconicity (negative). In task 2

lateral frontal cortex and superior parietal lobe showed parametric correlation to both weirdness

(negative) and iconicity (positive) measures (Fig. 4). Finding a common neuronal network to

both weirdness and iconicity with inverse correlation, supports the idea that these parameters

reflect two opposite aspects of shapes familiarity and symbolic meaning.

In order to disentangle the coupling between the subjective parameter of beauty and the objective

parameter of symmetry (Spearman correlation, $r = 0.92$, $p < 5*10^{-11}$), we calculated a subjective

beauty score – blocks were ranked by their beauty score per subject, and then beauty scores of

blocks with same ranking were averaged across subjects. The result is a perceptual beauty score

which is independent of physical attributes such as symmetry. ROI analysis demonstrated a

significant positive correlation between both symmetry/beauty and LOC activity (averaged

normalized beta weights). However, the subjective beauty scores showed a significantly higher

correlation compared to the symmetry scores (Spearman correlation; (experiment 1) left LOC,

Symmetry: $r = 0.4$, ns; Beauty: $r = 0.5$, $p < 0.005$; right LOC, Symmetry: $r = 0.47$, $p < 0.05$;

Beauty: $r = 0.72$, $p < 5*10^{-5}$; (experiment 2) left LOC, Symmetry: $r = 0.57$, $p < 0.005$; Beauty: r

$= 0.69$, $p < 5*10^{-5}$; right LOC, Symmetry: $r = 0.5$, $p < 0.05$; Beauty: $r = 0.81$, $p < 5*10^{-7}$, see Fig.

5). The correlation scores were further compared using bootstrapping method, where data points

were randomly chosen (with replacements) from each dataset and the correlation was calculated.

Each process was repeated 1000 times. (Spearman correlation; (experiment 1) left LOC,

Symmetry: $r = 0.28 \pm 0.17$, Beauty: $r = 0.54 \pm 0.13$, d'=1.2 ; right LOC, Symmetry: $r = 0.18 \pm 0.17$,

Beauty: $r = 0.72 \pm 0.18$, d'=2.9; (experiment 2) left LOC, Symmetry: $r = 0.35 \pm 0.15$, Beauty: r =

0.69 ± 0.06, d'=2.1; right LOC, Symmetry: $r = 0.28 \pm 0.16$, Beauty: $r = 0.81 \pm 0.05$, d'=3.1. All

values are Mean $\pm$STD, $N_{(symmetry)} = 16$, $N_{(beauty)} = 13$). According to these results perceptual

beauty showed a graded response across the entire dynamic range enhancing correlation and

significance while symmetry correlation highly depended on the points chosen as evident from

its large correlation variation (Fig. 5).

Iconicity group scores were also correlated to LOC activity; Spearman correlation; (experiment

1) left LOC, $r = 0.51$, $p < 0.01$; right LOC, $r = 0.65$, $p < 0.0005$. (experiment 2) left LOC, r =

0.65, $p < 0.0005$; right LOC, $r = 0.57$, $p < 0.005$.

Lastly, we tested whether one could infer a shape's origin by the beta activity of LOC. We
compared the average LOC activity for each block with its probability to be classified as human.
We fitted a non-linear hyperbolic tangent classifier with a random partial sample of data points
(23/28) and iterated the process 1000 times. We found that the averaged fitted function predicted
78% of the shapes accurately by using the average beta activity of the block (compared to 50%
for chance performance). Interestingly, the classifier's ability resembled the subject's behavioral
results with 86% correct in *random*, *chosen* and *top-rated* categories, and only 57% in the *not-*
chosen category. Thus, LOC activity might serve as a predictor to the shape's origin.

Discussion*:*

The human brain is skilled in distinguishing between the familiar and the strange, between the
natural and the artificial, and here we examined its ability to distinguish between human and
random creations. We showed here a tight connection between behavior and brain function
related to the process of identifying the origin of human-generated versus random abstract
shapes. Our findings revealed that human subjects could correctly identify shapes as human
made or randomly made. Subjects showed a general agreement that *random* shapes appeared
weirder. Moreover their aesthetic evaluation of the shapes (Fig. 2b) was very similar to the
evaluation made by the shapes' creators (the three categories of human made shapes - see
methods). These findings suggest a general agreement among individuals about what is
considered human, meaningful and beautiful.

While previous studies used complex naturalistic images or objects, in the current study we used
relatively simple and well-controlled shapes. Although similar in low level features, some of
them (*random* shapes) were out of the common human scheme as manifested by the players
playing the game. Indeed they were never created by human players (although probabilistically
they should have been created), and they were perceived differently by the human brain.

Brain activity - specifically, the LOC, showed preferential activation to *human* relative to
random blocks even in task 1 (color discrimination) in which attention was targeted to color
rather than shapes. This supports our hypothesis that the human brain is capable of recognizing
human creation even when not explicitly instructed to do so.

The LOC, well established as a hub of visual object recognition (Malach et al., 1995, Grill-
Spector et al., 2001) - was the central region showing a preferred activation to human-made

shapes compared to *random* shapes (task independent). The complementary, preferred activation
to *random* shapes was found in parietal regions, motor cortex and most significantly in the
inferior frontal gyrus (IFG, see figure 3). These findings are compatible with previous studies
which reported that IFG is responsive to unexpected stimuli (Huettel and McCarthy, 2004), and
to incongruent stimuli specifically within a social context (Shibata et al., 2011).
We examined 3 subjective shape characteristics with predictive value to classify as
human/random; beauty, weirdness and iconicity. Using a whole brain analysis exploratory
approach - we searched brain networks which were parametrically connected to each external
measure. These analyses (Fig. 4) showed that LOC was a central node - parametrically correlated
to shapes' beauty, weirdness (inversely correlated) and iconicity. Moreover, using LOC activity
might allow predicting shape's origin with human level accuracy (Fig. 6). Thus, our findings
point to the LOC as a central node for human vs. random origin, with shape's beauty and
symbolic meaning playing a role in that evaluation.
231
Brain and beauty:
Previous studies of beauty, focusing on response to visual arts or objects perception, found
activations in reward related areas, such as the orbito-frontal cortex (Kawabata and Zeki, 2004,
Brown et al., 2011, Ishizu and Zeki, 2011, Kirk, 2008, Lacey et al., 2011, Cela-Conde et al.,
2004), in emotion related areas like the amygdala (Brown et al., 2011, Di Dio et al., 2007, Ishizu
and Zeki, 2011) and insula (Brown et al., 2011, Di Dio et al., 2007) and also in motor areas
(Kawabata and Zeki, 2004, Ishizu and Zeki, 2011), pointing to a link between the experience of
beauty to reward or information gathering and response upon it.
Recent studies also point to high order visual areas, and specifically the LOC, as an aesthetic
center in the brain (Cattaneo et al., 2015, Chatterjee et al., 2009, Kirk, 2008, Lacey et al., 2011,
Vartanian and Goel, 2004). Some reported that high order visual areas show preferential activity
for aesthetic value only for representational art and not for abstract art (Cattaneo et al., 2015,
Vartanian and Goel, 2004).
In the current study we explored which brain regions were sensitive to the beauty of abstract
shapes. Our results revealed a consistent (experiments 1 and 2) positive parametric correlation
between subjective beauty scores and bilateral LOC (Fig. 4a). The consistency and specificity of

LOC in beauty analyses, suggests that beauty evaluation of abstract shapes is an automatic -

bottom up process which is less dependent on the attention to the shape.

Previous studies pointed to the LOC as a central player in symmetry evaluation (Hodgson, 2009,

Chatterjee et al., 2009, Beck et al., 2005, Sasaki et al., 2005). In the present study symmetry

indeed showed a significant correlation to beauty ratings. Importantly, the correlation between

LOC activity and symmetry was significantly weaker and less reliable compared to the beauty

score correlations (Fig. 5). Thus, our paradigm of abstract shapes allowed the decoupling of the

symmetry measure from that of the beauty scores. Our results show that subjective beauty

evaluation is better correlated to LOC activity compared to shapes' symmetry.

Brain and familiarity:

According to our results - both weirdness and iconicity, measures of familiarity and symbolic

meaning, were correlated with LOC activation. Previous works showed increased activation in

LOC which was similar for familiar and unfamiliar objects as long as the object is a coherent one

(Malach et al., 1995, Kanwisher et al., 1997, Kanwisher et al., 1996, Grill-Spector et al., 2001),

tracing a difference in activation between coherent objects response to non-coherent ones (using

Fourier descriptors, and scrambled images; Lerner et al., 2002, Tsao et al., 2003, Aalto et al.,

2002, Murray et al., 2002, Malach et al., 1995). This is compatible with the earlier finding of an

enhanced activity in mid-fusiform gyrus in a response to abstract shapes, associated with long-

term familiarization (Gauthier et al., 1999). Similarly, right LOC showed enhanced activity to

abstract object structures following a short-term learning (familiarization) (de Beeck et al.,

2006). Furthermore, LOC preferential activity to familiar (iconic) objects was shown in a visual

imagery with haptic perception task (Deshpande et al., 2010, Lacey et al., 2010).

Our abstract shapes paradigm introduced a broad range of familiarity levels, which allowed to

explore the effect of familiarity on LOC activity on a graded scale instead of a binary approach

(familiar/unfamiliar) and without distortion of object's features (Lerner et al., 2002, Tsao et al.,

2003, Aalto et al., 2002, Murray et al., 2002, Malach et al., 1995, Rotshtein et al., 2001). Shapes

with different familiarity levels manifested a difference in LOC activation although composed of

similar low level features.

Interestingly, weirdness and iconicity showed inversed correlation patterns within very similar

neuronal networks, supporting the idea that they reflect the two ends of shapes familiarity and

meaning (not only theoretically but also neuronaly). While the beauty parametric analysis

showed a consistent and local activity of LOC, both weirdness and iconicity parametric mapping

showed a task dependent activation, and a broader network involvement, including frontal and

parietal regions in addition to LOC (Fig. 4b,c).

It could have been argued that the identification of human origin shapes was mainly due to

highly recognizable symbols (e.g. letters and digits). However- such iconic shapes were only a

small portion (~10%) of the entire ensemble (sup Fig. 1). In fact, the shape ensemble introduced

a gradient of iconicity as was shown by both ratings and brain responses. Furthermore, it has

been well established that letters and digit representations are left- lateralized (Hasson et al.,

2002, Fiez and Petersen, 1998, McCandliss et al., 2003) while we consistently found a slight

right-hemisphere bias of the *human* vs. *random* contrast (sup Fig. 2).

De Beeck et al have showed an overall increased response to trained abstract objects compared

to untrained abstract objects. Right LOC in particular showed the strongest increased response as

a result to the training (de Beeck et al., 2006). In our study there was no formal training and all

the shapes were novel, however we suggest that the human schema is an expression of an

inherent training for social representations in the human mind. This schema led human players to

explore certain shapes, and avoid other shapes (unlike the random walk algorithm), similarly this

schema guided participants to successfully distinguish between human made and randomly made

shapes. It is an open question whether this schema was developed as a result or is the source of

human graphic communication. An interesting future avenue related to this question will be to

investigate whether these findings are reproducible across cultures, within cultures, and in young

infants. It will also be of interest to study whether humans on the autistic spectrum experience

and neuronaly process these shapes similarly to typical individuals.

Conclusion:

Our study examined the behavioral and neuronal components of evaluating human origin of

abstract shapes. Our results point to high order object areas (LOC) as a central node in beauty

representation, and in symbolic meaning attribution of abstract shapes. Both aspects had a

significant contribution to the classification of shape origin as *human* or *random*, and may have a

key role in visual human communication (such as visual art).

Materials and methods:

participants:

Seventeen healthy right handed subjects (ages 28 ± 3.8, 10 females) participated in the fMRI experiments. Fifteen of them participated in both experiments 1 and 2. Fourteen filled a beauty, iconicity and weirdness evaluation questionnaire post the fMRI scan.

Task and stimuli:

Shape stimuli. Shapes of ten contiguous identical green squares were created in a shape-search computer game, by either 101 human players or a random walk algorithm (Noy et al., 2012). In each shape, squares were connected by an edge. Players moved one square at a time to create new shapes. They were instructed to place beautiful and interesting shapes into a gallery by pressing a button. There was no limit to the gallery. Players played for 15 min and created about 310 shapes, of which they chose 46 shapes on average to the gallery. At the end of the game, players chose the 5 most creative shapes from their own gallery. The shape space includes 36,446 possible shapes.

Shapes were classified into four categories based on their origin (human/algorithm), and their appeal ratings (by their human creators). *Not chosen shapes* were created by human players but never chosen as beautiful shapes (by the players). *Chosen shapes* were created by human players, chosen as beautiful and interesting to the gallery but never rated as most creative shapes. *Top rated shapes* were created by human players, chosen as beautiful and interesting shapes and were rated by players as most creative shapes. *Random* (*Never human-created shapes)* were created only by a random walk algorithm (and never by human players) on the space of shapes, where the next shape is chosen randomly from neighboring shapes that are one move of a square away from the current shape. Length of walks was sampled from the distribution of walk lengths of the human players. For each category, the 20 most frequent shapes, i.e. the shapes that were the most common to many players (or random walks in the random category), were chosen for the fMRI experiment (see the shapes in sup. Fig. 1).

Experimental design

During the fMRI scan the created shapes were presented in homogeneous-category blocks lasting 9 sec, followed by a 9 sec fixation screen. Each block consisted of 9 images (one second each);

eight images in light green and one image in dark green. Each category was presented in seven
different blocks. To reduce scan novelty effect, an extra block (which was not analyzed) was
added to the beginning of each experiment, 29 blocks were presented in total.
Each subject watched the same sequence of blocks twice (once for each task). In the first
experiment (task 1) subjects were required to classify the stimuli according to color; light green
(press 1) or dark green (press 2). In the second experiment (task 2) following each block,
subjects were required to classify the shapes of the preceding block as human creation (press 1)
or random algorithm creation (press 2).
349

Subjective and objective shape evaluations
Following the scan, subjects evaluated each shape's beauty level on a 1-4 scale (4 being most
beautiful), chose the 20 weirdest shapes, and also chose the most iconic shapes in blocks of 20
shapes.
Block's beauty score was calculated as the summation of beauty scores of all the shapes in the
block. Weirdness and iconicity scores were calculated independently, in the same manner; A
shape's weirdness/iconicity score was equal to the number of subjects which rate the shape as
weird/iconic. A block's weirdness/iconicity score was calculated as the summation of
weirdness/iconicity   scores   of   all   shapes   in   the   block.
In addition- the shapes were analyzed according to symmetry as an objective parameter. Block's
symmetry was calculated as the summation of all rotation and reflection symmetry groups of all
the shapes in the block.
362

MRI Data Acquisition and Preprocessing
The data were acquired on a 3 Tesla Trio Magnetom Siemens scanner at the Weizmann Institute
of Science. Functional images of blood oxygenation level dependent (BOLD) contrast
comprising of 46 axial slices were obtained with a T2*-weighted gradient echo planar imaging
(EPI) sequence ($3 \times 3 \times 3$ mm voxel , TR = 3000 ms, TE = 30, flip angle = 75°, FOV 240 mm)
covering the whole brain. Anatomical images for each subject were acquired in order to
incorporate the functional data into the 3D Talairach space (Talairach and Tournoux, 1988) using
3-D T1- weighted images with high resolution ($1 \times 1 \times 1$ mm voxel, MPRAGE sequence, TR=
2300 ms, TE= 2.98 ms).

The first 7 images of each functional scan (including the extra initial block and rest) were

discarded. Functional scan preprocessing included 3D motion correction and filtering out of low

frequency noise (slow drift), and spatial smoothing using an isotropic Gaussian kernel of 6 mm

full-width-half-maximum (FWHM). The functional images were superimposed on 2D anatomic

images and incorporated into the 3D data sets through trilinear interpolation. Statistical analysis

was based on a general linear model in which all stimuli conditions were defined as predictors,

and convolved with the hemodynamic response function (HRF).

Data analysis:

In order to learn about the connection between shapes' characteristics, perception and brain

activity several measurements were examined; Reaction times (both tasks), response accuracy

(*human*/*random*, task2), subjective evaluations post-scan (beauty, weirdness and iconicity), and

symmetry   score.

To investigate a possible difference in reaction times between shape categories (*random, not*

chosen, chosen, top rated), Mann-Whitney tests were calculated for experiment 1 and 2 within

subject and between subjects. Aesthetic (beauty) ratings for each shape were collected by each

subject post the scan (on a 1-4 scale). To control for difference in rating patterns each subject's

ratings were Z normalized. Five GLM analyses were conducted; the first included four

predictors: *random, not chosen, chosen, top rated*. A second GLM analysis with two predictors

based on the category's creator: *human* or *random*. In addition, in order to relate subjective

blocks' characteristics to brain activity, three parametric GLM analyses were conducted for

beauty, weirdness and iconicity. In these multi-subject, random effect analyses each block of

shapes received a weight according to its score (beauty/weirdness/iconicity) which was

represented in the model as differential amplitude of the BOLD signal. Beauty, being an

individual score calculated per subject separately, was z normalized between subjects.

Model selection of beauty and weirdness as well as beauty and iconicity fit to the probability to

be classified as human was done based on the spearman correlation between model predictions

and the Akaike information criterion (AIC). Models were generated as all possible combinations

of the three parameters, either alone or coupled together. In order to have the monotonicity of the

two scores increase in the same direction, weirdness score was represented as $e^{-Weirdness\ score}$

(see SI for more details).

In all GLM analyses beta coefficients were calculated for the regressors, and a Student's t-test

was performed. Multi-subject analysis was based on a random-effect GLM. Multi-subject

contrast maps (*human* vs. *random*, or category vs. category) were projected on an unfolded,

inflated Talairach-normalized brain. Significance levels were calculated, taking into account the

minimum cluster size and the probability threshold of a false detection of any given cluster. This

was accomplished by a Monte Carlo simulation (cluster-level statistical threshold estimator in

"Brain Voyager" software).

In the first experiment (task1) for *human* vs. *random* contrast (Fig. 3) a minimum cluster size of

103 voxels was significant. A minimum cluster size of 71 voxels was significant for *top rated* vs.

chosen, 78 voxels for both *top rated* vs. *not chosen*, and *chosen* vs. *not chosen* (sup Fig. 2a). The

minimum significant cluster size for each *human* category vs. *random* (sup Fig. 2b) was 103

voxels (*top rated*), 91 voxels (*chosen*) and 85 voxels (*not chosen*). For the parametric maps (Fig.

4) a minimum cluster size of 84 voxels was significant for beauty in task 1, and 90 voxels in task

2. For weirdness a minimum cluster size of 98 voxels was significant in task 1, and 105 voxels in

task 2. For iconicity a minimum cluster size of 90 voxels was significant in task 1, and 104

voxels in task 2.

ROI definition and analysis

Analysis of LOC-relevant voxels was conducted by defining a group bilateral ROI within the

LOC using the contrast *human > random* in one task, and sampled in the other task. Note that the

inverted contrast i.e. *random> human* failed to reveal any voxels in the LOC region (see Fig. 3).

The ROI's averaged beta weight (across voxels) was calculated per subject, for each predictor.

Two-tailed paired t-tests (within subjects) were conducted between *human* and *random* beta

weights for unaware (task 1) and aware (task 2) stimuli. A beta weight was extracted for each

block and was plotted as a function of subjective beauty and symmetry (Fig. 5). Spearman

correlation was calculated between participant's average beta for each block (averaged over each

ROI) and their aforementioned features.

Acknowledgements

We thank Nahum Stern, Fanny Atar and Dr. Edna Haran-Furman for their assistance in the imaging setup

and fMRI data collection. We thank Dr. Lior Noy for developing the shapes-creation computer game.

This study was supported by the EU FP7 VERE, EU HBP Flagship (R.M.), ICORE ISF grants (R.M.), the

Helen and Martin Kimmel award (R.M.), the Braginsky Center for the Interface Between Science and

Humanities (U.A). U.A. is the incumbent of the Abisch-Frenkel Professorial Chair.

References

AALTO, S., N T NEN, P., WALLIUS, E., METS HONKALA, L., STENMAN, H., NIEMI, P. &
KARLSSON, H. 2002. Neuroanatomical substrata of amusement and sadness: a PET
activation study using film stimuli. *Neuroreport,* 13**,** 67.
BECK, D. M., PINSK, M. A. & KASTNER, S. 2005. Symmetry perception in humans and
macaques. *Trends in cognitive sciences,* 9**,** 405-406.
BROWN, S., GAO, X., TISDELLE, L., EICKHOFF, S. B. & LIOTTI, M. 2011. Naturalizing
aesthetics: brain areas for aesthetic appraisal across sensory modalities. *Neuroimage,* 58**,**
250-258.
CATTANEO, Z., LEGA, C., FERRARI, C., VECCHI, T., CELA-CONDE, C. J., SILVANTO, J.
& NADAL, M. 2015. The role of the lateral occipital cortex in aesthetic appreciation of
representational and abstract paintings: A TMS study. *Brain and cognition,* 95**,** 44-53.
CELA-CONDE, C. J., MARTY, G., MAESTÚ, F., ORTIZ, T., MUNAR, E., FERNÁNDEZ, A.,
ROCA, M., ROSSELLÓ, J. & QUESNEY, F. 2004. Activation of the prefrontal cortex in
the human visual aesthetic perception. *Proceedings of the National Academy of Sciences*
of the United States of America, 101**,** 6321-6325.
CHATTERJEE, A., THOMAS, A., SMITH, S. E. & AGUIRRE, G. K. 2009. The neural
response to facial attractiveness. *Neuropsychology,* 23**,** 135.
CHAUVET, J.-M., BRUNEL DESCHAMPS, E. & HILLAIRE, C. 1996. *Dawn of art: the*
Chauvet Cave: the oldest known paintings in the world, HN Abrams.
DE BEECK, H. P. O., BAKER, C. I., DICARLO, J. J. & KANWISHER, N. G. 2006.
Discrimination training alters object representations in human extrastriate cortex. *The*
Journal of Neuroscience, 26**,** 13025-13036.
DESHPANDE, G., HU, X., LACEY, S., STILLA, R. & SATHIAN, K. 2010. Object familiarity
modulates effective connectivity during haptic shape perception. *Neuroimage,* 49**,** 1991-
2000.
DI DIO, C., MACALUSO, E. & RIZZOLATTI, G. 2007. The golden beauty: brain response to
classical and renaissance sculptures. *PloS one,* 2**,** e1201.
FIEZ, J. A. & PETERSEN, S. E. 1998. Neuroimaging studies of word reading. *Proceedings of*
the National Academy of Sciences, 95**,** 914-921.
GAUTHIER, I., TARR, M. J., ANDERSON, A. W., SKUDLARSKI, P. & GORE, J. C. 1999.
Activation of the middle fusiform'face area'increases with expertise in recognizing novel
objects. *Nature neuroscience,* 2**,** 568-573.
GRILL-SPECTOR, K., KOURTZI, Z. & KANWISHER, N. 2001. The lateral occipital complex
and its role in object recognition. *Vision research,* 41**,** 1409-1422.
HASSON, U., LEVY, I., BEHRMANN, M., HENDLER, T. & MALACH, R. 2002. Eccentricity
bias as an organizing principle for human high-order object areas. *Neuron,* 34**,** 479-490.

HODGSON, D. 2009. Evolution of the visual cortex and the emergence of symmetry in the
Acheulean techno-complex. *Comptes Rendus Palevol,* 8**,** 93-97.
HUETTEL, S. A. & MCCARTHY, G. 2004. What is odd in the oddball task?: Prefrontal cortex
is activated by dynamic changes in response strategy. *Neuropsychologia,* 42**,** 379-386.
ISHIZU, T. & ZEKI, S. 2011. Toward a brain-based theory of beauty. *PloS one,* 6**,** e21852.
JOHANSSON, G. 1973. Visual perception of biological motion and a model for its analysis.
Attention, Perception, & Psychophysics, 14**,** 201-211.
KANWISHER, N., CHUN, M. M., MCDERMOTT, J. & LEDDEN, P. J. 1996. Functional
imaging of human visual recognition. *Cognitive Brain Research,* 5**,** 55-67.
KANWISHER, N., WOODS, R. P., IACOBONI, M. & MAZZIOTTA, J. C. 1997. A locus in
human extrastriate cortex for visual shape analysis. *Cognitive Neuroscience, Journal of,*
9**,** 133-142.
KAWABATA, H. & ZEKI, S. 2004. Neural correlates of beauty. *Journal of Neurophysiology,*
91**,** 1699-1705.
KIRK, U. 2008. The neural basis of object-context relationships on aesthetic judgment. *PloS one,*
3**,** e3754.
LACEY, S., FLUECKIGER, P., STILLA, R., LAVA, M. & SATHIAN, K. 2010. Object
familiarity modulates the relationship between visual object imagery and haptic shape
perception. *Neuroimage,* 49**,** 1977-1990.
LACEY, S., HAGTVEDT, H., PATRICK, V. M., ANDERSON, A., STILLA, R.,
DESHPANDE, G., HU, X., SATO, J. R., REDDY, S. & SATHIAN, K. 2011. Art for
reward's sake: Visual art recruits the ventral striatum. *Neuroimage,* 55**,** 420-433.
LERNER, Y., HENDLER, T. & MALACH, R. 2002. Object-completion effects in the human
lateral occipital complex. *Cerebral Cortex,* 12**,** 163-177.
MALACH, R., REPPAS, J., BENSON, R., KWONG, K., JIANG, H., KENNEDY, W.,
LEDDEN, P., BRADY, T., ROSEN, B. & TOOTELL, R. 1995. Object-related activity
revealed by functional magnetic resonance imaging in human occipital cortex.
Proceedings of the National Academy of Sciences, 92**,** 8135-8139.
MCCANDLISS, B. D., COHEN, L. & DEHAENE, S. 2003. The visual word form area:
expertise for reading in the fusiform gyrus. *Trends in cognitive sciences,* 7**,** 293-299.
MURRAY, S. O., KERSTEN, D., OLSHAUSEN, B. A., SCHRATER, P. & WOODS, D. L.
2002. Shape perception reduces activity in human primary visual cortex. *Proceedings of*
the National Academy of Sciences, 99**,** 15164-15169.
NOY, L., HART, Y., ANDREW, N., RAMOTE, O., MAYO, A. & ALON, U. Year. A
quantitative study of creative leaps. *In:* Proc. Third Int. Conf. Comput. Creativity,
Dublin, 2012. Citeseer, 72-76.
ROTSHTEIN, P., MALACH, R., HADAR, U., GRAIF, M. & HENDLER, T. 2001. Feeling or
Features:: Different Sensitivity to Emotion in High-Order Visual Cortex and Amygdala.
Neuron, 32**,** 747-757.
SASAKI, Y., VANDUFFEL, W., KNUTSEN, T., TYLER, C. & TOOTELL, R. 2005. Symmetry
activates extrastriate visual cortex in human and nonhuman primates. *Proceedings of the*
National Academy of Sciences of the United States of America, 102**,** 3159-3163.
SHIBATA, H., INUI, T. & OGAWA, K. 2011. Understanding interpersonal action coordination:
an fMRI study. *Experimental Brain Research,* 211**,** 569-579.
SIMION, F., REGOLIN, L. & BULF, H. 2008. A predisposition for biological motion in the
newborn baby. *Proceedings of the National Academy of Sciences,* 105**,** 809-813.

TALAIRACH, J. & TOURNOUX, P. 1988. Co-planar stereotaxic atlas of the human brain. 3-
Dimensional proportional system: an approach to cerebral imaging.
TSAO, D. Y., FREIWALD, W. A., KNUTSEN, T. A., MANDEVILLE, J. B. & TOOTELL, R.
B. 2003. Faces and objects in macaque cerebral cortex. *Nature neuroscience,* 6**,** 989-995.
VARTANIAN, O. & GOEL, V. 2004. Neuroanatomical correlates of aesthetic preference for
paintings. *Neuroreport,* 15**,** 893-897.

569
Figure 1. Experimental design .(a) Experimental protocol: shape images from four categories were presented in a
block design with 9 second blocks, followed by 9 second fixation screen. Shape categorization was based on its
creation process during a computer game in a previous work (Noy et al., 2012, see methods); *Top rated shapes* were
created by human players and rated as most creative shapes. *Chosen shapes* were created by human players, chosen
as beautiful shapes but never rated as most creative shapes. N*ot chosen shapes* were created by human players but
never chosen as beautiful shapes or rated (by the players). *Never created shapes* were created by a random
algorithm, and never by human players. (b) Example for a block of each category. Each block included 9 images
(one second each) of the same category; 8 images in light green and one in dark green.

Figure 2. Behavioral measurements (a) Group's average percent success rate (human/random) in each category; *Human- top rated, Human- chosen, Human- not chosen, Algorithm-random*. Two-tailed, within subjects Mann-Whitney between categories, n1 = n2 = 7, p < 0.005**, error bars indicate the groups' standard error. (b) Group's average beauty scores per category (z-score normalized). (c) Group's average weirdness scores per category.

603
604
Figure 3. Comparison of *Human* and *Random* cortical activations. All human generated shapes *(Human)* vs. computer generated shapes *(Random)* multi subjects activity map (experiment 1, N = 17, corrected p < 0.05) is presented on an inflated cortex, in a lateral view (a) and unfolded cortex (b). Color scale indicates t values. Yellow-orange scale represents regions which were more activated while watching human generated blocks compared to random shapes (blue-green scale). Most significant activated regions are marked on the inflated map; Lateral Occipital cortex (LOC), Inferior frontal gyrus (IFG).
(c) LOC ROI analysis: repeated measures ANOVA between averaged beta values of each category; *Human- top rated, Human- chosen, Human- not chosen, Algorithm-random*. Left side - experiment 1 (N = 17, p < 0.05*), right side - experiment 2 (N = 15, p < 0.005**).

615
616
Figure 4. Parametric mapping. Cortical activity maps of multi subjects, random effect, parametric GLM analysis
of beauty (a), weirdness (b), and iconicity (c) presented on an unfolded cortex, right side experiment 1 (N = 13, 13,
14), bottom panel experiment 2 (N = 12, 12, 13), both maps are corrected for multiple comparisons, p < 0.05. Color
scale indicates t values. Yellow- orange scale represents regions which showed positive parametric relation with
beauty/weirdness/iconicity scores. Blue-green scale represents regions which showed negative parametric relation
with beauty/weirdness/ iconicity scores.

630
Figure 5. Correlations of LOC activity and symmetry/beauty. (a) Scatter plots present the relation between
averaged blocks' symmetry (x axis) and group's averaged brain activity (normalized beta weight, y axis) in bilateral
LOC. Each dot represents one block, and color indicates the blocks' category (*Top rated* in red, *chosen* in orange,
not chosen in green, *random* in blue) N = 13. Spearman correlation; (experiment 1) left LOC, r = 0.4, ns ; right
LOC, r = 0.5, p < 0.05. (experiment 2) left LOC, r = 0.6, p < 0.005; right LOC, r = 0.5, p < 0.05. (b) The relation
between group's averaged beauty ratings (x axis), and the group's averaged brain activity (normalized beta weight, y
axis) in LOC. Each dot represents mixed blocks with similar beauty ranking, color indicates the blocks' category. N=
16. Spearman correlation; (experiment 1) left LOC, r = 0.52, p < 0.005 ; right LOC, r = 0.69, p < 0.0005.
(experiment 2) left LOC, r = 0.69, p < 0.0005; right LOC, r = 0.83, p < 0.0005.

640
Figure 6. Classification of blocks' origin based on their average LOC activity results in human-level
accuracy. (a) We fitted a hyperbolic tangent classifier, $P(human) = a + d\,Tanh[b\,x + d]$ (where x is taken to be
the averaged LOC activity of the block) to the data points by a bootstrapping method. Each iteration, 23/28 points
were chosen randomly and the best fit parameters were extracted. We repeated this process 1000 times and averaged
the parameters of all the runs. Shown is the averaged model. Model parameters are (mean±ste): a=0.48±0.03,
b=45±9, c=5±1, d=0.33±0.04. (b) Classification accuracy of the classifier on the different categories (top rated,
chosen, not-chosen, random). Error bars are STE and are calculated by 100 random sampling with replacements of
the real data points per each category.

a.

c.

b.

d.

Supplementary Figure 1. The entire stimuli ensemble. (a) *Top rated* shapes, (b) *Chosen* shapes, (c) *Random - Never created* shapes, (d) *Not chosen* shapes.

652
Supplementary Figure 2. Comparison cortical activations between categories (experiment 1). Multi subjects
activity maps (N = 17, corrected p < 0.05) are presented on an unfolded cortex. Color scale indicates t values. (a)
Contrast maps between all the human generated shapes categories *(Human)*. Yellow- orange scale represents regions
which were more activated while watching blocks from the category in the left side of the contrast. Blue-green scale
represents regions which were more activated while watching blocks from the category in the right side of the
contrast. (b) Contrast maps between the *Human* categories and *Random* category (computer generated shapes).
Color scale indicates t values. Yellow- orange scale represents regions which were more activated while watching
shapes of *human* categories, compared to *random* shapes (blue-green scale).

Supplementary Information

<u>Modelling shapes' probability to be classified as human created using the behavioral scores of</u>

<u>subjects</u>

Here we describe the model selection process of finding the best model for the probabilities to be

classified as human-created vs. random created shapes. The probability function was fit using a

Logit function, $P(x,y) = \frac{1}{1+e^{f(x,y)}}$. The variables of the model were the three subjective scores in

our experiment – beauty (denoted as b), weirdness (denoted as w) and iconicity (denoted as i).

Since weirdness and iconicity carry inverse correlations to the behavioral data (iconicity

increases the probability to be classified as human, while weirdness decreases that same

probability), we chose to map the weirdness score in the following way - $w' \to e^{-w}$.

We tested all possible linear combinations of the individual scores, their pair and triplet

interactions, yielding:

(S1) $f(b, w', i) = m_0 + m_1 b + m_2 w' + m_3 i + m_{12}b * w' + m_{13}b * i + m_{23} w' * i +$

$m_{123} b * w' * i$

We assessed each model's correlation with the data as well as its Akaike Information Criterion

(AIC) score to attain the most accurate and simplest best fit model. The most accurate model

with least number of parameters is one containing the interaction term between beauty and

weirdness as a sole parameter, suggesting that it is the combination of beauty and familiarity that

is the dominant component of classifying a shape as having human origin. In Table S1 we list the

highest correlation and AIC scores of the different models for both beauty and weirdness and

beauty and iconicity.

Table S1: Pearson correlation, p-values and AIC scores of Logit models with beauty,
weirdness and iconicity scores.

	Model	Model parameters	Spearman correlation	P-value	AIC scores
1	$f(b, w', i) = b * w'$	$\dfrac{1}{1 + e^{2.6 - 6.6\,b\,w'}}$	0.96	$4*10^{-21}$	5.84
2	$f(b, w', i) = i$	$\dfrac{1}{1 + e^{1.5 - 7\,i}}$	0.94	$5*10^{-19}$	6.32
3	$f(b, w', i) = i * w'$	$\dfrac{1}{1 + e^{1.4 - 7\,i\,w'}}$	0.94	$1*10^{-18}$	6.4
4	$f(b, w', i) = b$	$\dfrac{1}{1 + e^{3.8 - 8\,b}}$	0.93	$5*10^{-17}$	6.54

5	$f(b, w', i) = b * i$	$\dfrac{1}{1 + e^{1.3 - 9.2\, b\, i}}$	0.92	$5*10^{-16}$	7.04
6	$f(b, w', i) = b * w' * i$	$\dfrac{1}{1 + e^{1.2 - 9.3\, b\, i\, w'}}$	0.92	$1*10^{-15}$	7.13
7	$f(b, w', i) = w'$	$\dfrac{1}{1 + e^{7.1 - 9\, w'}}$	0.89	$2*10^{-12}$	5.36