1

Social ecology of a woodland songbird community: from individual

movements to the emergence of population social structure

Damien R. Farine[a,b,c], Ben C. Sheldon*[a]

[a] Edward Grey Institute of Field Ornithology, Department of Zoology, University of Oxford,

South Parks Road, Oxford OX1 3PS, United Kingdom.

[b] Department of Collective Behaviour, Max Planck Institute for Ornithology,

Universitätsstrasse 10, 78457 Konstanz, Germany.

[c] Department of Biology, University of Konstanz, Universitätsstrasse 10, 78457 Konstanz,

Germany.

* Corresponding author:

damien.farine@orn.mpg.de

ABSTRACT

Social structure can have profound evolutionary and ecological implications for
animal populations. Structure can arise and be maintained via social preferences or
be indirectly shaped by habitat structure. Quantifying the drivers of social structure
is important to understand how social networks can shape evolutionary landscapes.
Here, we study a large community of wild birds fitted with uniquely-coded passive
integrated transponder (PIT) tags and recorded on a grid of automated feeders
fitted with radio frequency identification (RFID) antennae. These data reveal that
preferred movement pathways between sites that are consistent between years and
not predicted by habitat features alone drive between-year consistent multi-level
community structure in the social network. Our study highlights how ecological
factors can shape social structure at the population scale, which has widespread
implications for understanding eco-evolutionary dynamics.

INTRODUCTION

The social environment can fundamentally shape the life histories of animals. Who
individuals associate with can determine the information they have access to [1, 2, 3],
affect how well they can exploit resources [4, 5, 6], and impact their ability to
successfully reproduce [7, 8, 9, 10]. Social structure can also influence population
processes, such as the spread of information within [11, 12, 13] and between [14] species,
and the spread of disease among individuals [15, 16]. Finally, individuals can experience
selection arising from properties of their social groups [7, 8] or their communities [17].
Quantifying the factors that shape individuals' social environments is therefore a
key step in determining the evolutionary drivers of sociality [18, 19] as the mechanisms
that determine where, and with whom, individuals live their lives are what selection
arising from population processes (e.g. disease burden or information use) can act
upon.

Population social structure (the patterns of connections that emerges from
interactions among individuals, often represented using social networks) is

generally considered to arise from individuals making decisions about whom to associate with [20]. However, because spatial proximity is a key requirement for associations or interactions (i.e. connections) among animals [21], patterns of connections among individuals are bound to be shaped by a number of extrinsic factors, such as geometry and structure of the habitat that promote or restrict individual movements. At a local scale, habitat structure, such as understorey density, can channel individual movements, thus influencing the propensity for contact among individuals. At a broader scale, movement 'highways' can significantly reduce the social distance among individuals despite a large spatial separation, and therefore facilitate the flow of information or disease [22]. For example, sleepy lizards (*Tiliqua rugosa*) living in open habitats had fewer contacts with conspecifics than those living in more structured habitats [23]. Whilst habitat features can promote movements, the geometry of habitat (such as habitat fragmentation) can introduce barriers that constrain individual movements. Such barriers can result in a structure where the social distance between two individuals (i.e. their degrees of separation in a social network) on either side of this barrier could be much greater than their actual spatial distance. Habitat geometry and features can therefore impose structure in a social network that could easily be interpreted as arising socially. Given that local heterogeneity in gene flow can lead to rapid evolutionary differentiation [24], integrating knowledge about fine-scale environmental heterogeneity into studies of social structure could fundamentally alter our understanding of adaptation and the ability for animals to respond to selective pressures.

A major challenge in identifying extrinsic factors that drive population social structure is the need to have information about the movement of individuals in space, the patterns of social connections among the majority of individuals in a population, and information about the habitat in which they live. Recent technological advances have eased the logistical constraints of sampling many individuals moving across large areas [25, 26, 27]. In particular, passive integrated transponder (PIT) tags are cheap electromagnetic tags that can be fitted to many

individuals at once [28], thus overcoming the challenges associated with studying

entire animal communities. Because PIT tags do not rely on battery power to emit a

signal, but instead are detected by affecting the magnetic field in radio frequency

identification (RFID) antennas, they provide the unprecedented ability to track

individuals across years and life stages. Further, advances in the analytical tools,

such as social network analysis [29] and community detection algorithms [30], are

facilitating greater insight into patterns of social structure. However, although

technological advances have underpinned a boom in the study of animals' social

networks [31], much less effort has focused on identifying key mechanisms that form

communities and shape social structure at the scale of populations.

In this paper, we investigate the patterns of individual movement and the resulting

community structure in a large contiguous mixed population comprising 5 species

of wild songbird over 4 winters [Farine et al *in prep.*]. We first quantify movements

across the landscape across all four years to determine if individuals move evenly

through space, or if there are consistent movement corridors. We determine

whether preferred movement pathways are maintained across generations, and

whether they are associated with understorey habitat density and the geometry of

the woodland. Second, we use patterns of connections among individuals to

construct social networks, and use these social networks to determine at what

scales population social structure is most consistent across years. Doing so helps us

to elucidate the relative importance of different potential drivers (both social and

habitat) that underpin population social structure.

RESULTS

Quantifying individual movements

To determine whether individuals moved freely through Wytham Woods, or if

movement patterns are shaped by features of the habitat, we first quantified the

daily movement rates between different feeding stations. Our population consisted

of a total of 5163 unique PIT-tagged individuals (see Methods), of which 3051 were

great tits (*Parus major*), 3960 were blue tits (*Cyanistes caeruleus*), 403 were marsh

tits (*Poecile palustris*), 232 were coal tits (*Periparus ater*), and 110 were nuthatches

(*Sitta europaea*). We detected the presence and group membership of birds using a

grid of 65 bird feeders filled with unhusked sunflower seed, and studied the

population over 4 winters.

We first created a movement network combining the data across all years (Figure 1,

see methods). We found that the majority of movements (n=83071) occurred

between feeding stations where more birds were present. However, not all birds

followed the same patterns. First, we found that birds in their first winter made

many more moves than adult birds: first year birds accounted for 61% of all

movements despite making up only made up 38% of the population on average.

Further, their movement networks included many more long-distance movements

than those of adults (Figure 1), resulting in a significantly higher average movement

distance (Figure S1). Second, we found that although great and blue tits exhibited

similar movement patterns, marsh tit movements were typically much more

localized (Figures S2, S3). Thus, age and species both affected the distribution of

overall movements. These patterns suggest that many social processes, such as

disease spread or information transmission (both within and between species) will

be linked to demographic processes.

Figure 1: Total movements of (A) all birds, (B) adults, and (C) juveniles from all

species over 4 winters of data. The thickness of each line represents the number of

observations of a bird moving between the two feeding stations (black points) in the

same day. The size of the points represents the average number of individuals

observed at each feeding station. The green outline represents the outline of

Wytham Woods and four small external woodplots.

Testing the predictors of movement propensity

Habitat structure is likely to shape movement patterns by birds during the winter in

several ways. First, for arboreal species, individuals are more likely to move through

the forest than over open group, and thus the distance between foraging patches

needs to be calculated through the forest and thus the presence of open areas is

likely to have some impact on the spatial patterns of movements. Second, within the

forest, the understory structure of the habitat could create preferred areas of

movement by providing shelter from predators. Finally, additional factors such as

local traditions could maintain consistent movement pathways, when generations

mix and where social learning occurs. We compared the patterns of movement

observed across years, and relate these to spatial distance and habitat features. To

avoid the potential confounding effect of population density in our statistical

analyses, we re-defined edges in the movement network as the probability that an

individual detected at either site moves between them on a given day (see

methods). Edge weights ranged from 0 representing no movement between

adjacent sites to 0.83 representing that each individual detected had an 83% chance

of moving between the two sites on a given day (Figure S4). For each pair of sites,

we also estimated the density of understorey vegetation between them using data

collected at 100m intervals throughout Wytham Woods [32] (see methods). Because

Wytham Woods is unevenly shaped, we also calculated distance between feeding

sites as *forest distance*, which is the shortest line without leaving boundaries of the

forest. To enable direct comparison between each predictor variable, we scaled each

predictor to zero mean and unit variance.

Weighted multiple regression quadratic assignment procedure (MRQAP) [33] revealed

that the movements by birds between feeding stations were significantly more

similar from year to year than expected by chance, even when accounting for forest

distance and habitat structure (see Table 1, Figure 2). In all years, birds were

significantly more likely to move between 'close' feeding stations than distant ones,

and were also more likely to move between feeders that were separated by higher

understory habitat density. However, the propensity to move between feeders

observed in previous years was consistently the strongest predictor of future

movements. The coefficient values for movements predicted by the previous year

are typically an order of magnitude larger than those of other predictor variables,

suggesting that some undetected factors are driving patterns of movements by birds

across this woodland. Because for many feeding stations the distance and forest

distance are very similar (see Figure 2), we also tested whether birds were less

likely to move between feeding sites that were separated non-forest by calculating

the difference between forest distance and Euclidian distance. We found that birds

moved less between feeders with a larger difference in distance (i.e. the path

through the forest was much longer than the straight-line path), suggesting that

birds are avoiding crossing open fields (Table 2).

Table 1: Results of multiple regression quadratic assignment procedure used to test

whether previously observed patterns of movement (probability of moving between

sites per capita), distance between sites (distance through the forest), and habitat

density (percentage cover) between sites explain the observed patterns of

movement. Bold values represent significant coefficients, * represents significance

at P < 0.01, ** represents significance at P < 0.001. All variables are scaled to 0 mean

and unit variance to enable comparison between effect sizes.

Year	Previous Year	Forest Distance	Habitat Density
2013	**0.903****	**-0.028***	-0.010
2014	**0.877****	**-0.060****	-0.010
2015	**0.787****	**-0.091****	-0.019

Figure 2: Total movement of all birds (from Figure 1A) overlaid on the shrub layer

density (0.5m to 2.5m above ground). Areas with a white background have no

habitat data available.

Table 2: Results of multiple regression quadratic assignment procedure used to test

whether birds moved less between feeding sites separated by open space. For each

pair of sites, we calculated the difference between the forest distance and the

Euclidian distance. Bold values represent significant coefficients, * represents

significance at P < 0.01, ** represents significance at P < 0.001. All variables are

scaled to 0 mean and unit variance to enable comparison between effect sizes.

Year	Previous Year	Relative Distance
2013	**0.919****	-0.001
2014	**0.906****	**-0.019***
2015	**0.828****	**-0.029***

Quantifying community structure

Population-level social structure can have significant implications for population

processes. For example, high levels of clustering can reduce the spread of disease

within populations [34]. Thus far, we have shown that both social and habitat features

are likely to contribute to where animals move. Here we investigate how these

patterns contribute to the emergent structure of the population. From the

detections of individuals at feeding stations, we inferred flock membership and

social network structure (see methods) using well established approaches

developed for this population [7, 12, 13, 14, 29, 35, 36, 37, 38]. We then inferred communities in

the social networks using the edge betweenness community detection algorithm in

the R package *igraph* [39], and explored the between-year stability of individual co-

membership to the same community (see Methods).

To investigate population social structure at different scales, we cut the social

network into k communities, where ranged from 2 to 65 (the latter representing the

total number of locations with a feeding station). If communities are structured

exclusively by extrinsic fctors, we expected a drop in the stability of co-membership

by individuals as we created more fine-scale communities. For example, if a

population is spread across three isolated patches of woodland, then we expect that

birds will always occur within the same three communities (one for each woodland)

each year. By contrast, if communities are structured socially, then we expected

smaller communities (local cliques) to be more stable. For example, for a territorial

pair-living species living in a lattice-like uniform environment, an algorithm will be

able to isolate each pair when identifying k=N/2 communities, whereas the

communities detected for smaller k values will be essentially random. When applied

to our social network data, we uncovered two scales that maximized the propensity

for pairs of individuals that were observed in the same community to be re-

observed in the same community in the following year. When social networks were

partitioned into 2 or 3 communities (Figure 3), individuals observed in the same

community in one year and observed again in the following year re-occurred in the

same community approximately 90% of the time. These communities largely

represent the geometry of Wytham Woods, with two core habitats (north-west and

the east), and a small wood to the south that is only attached by a narrow neck of

vegetation representing a third community.

Figure 3: The probability that two birds observed in the same community in one

year remain in the same community in the following year, given that both are

observed. The probability is calculated with the social network partitioned into the

same specified number of communities (2 to 65) in all years.

Specifying the algorithm to detect 4 communities significantly decreased the

probability of individuals being re-observed in the same community (no overlap in

the 95% confidence intervals in Figure 4), suggesting that there is no stable 4[th]

community. Partitioning the network further by specifying the algorithm to detect

more than 4 communities then increased probability that two individuals occurring

in the same community in one year and both observed in the following year again

re-occurred in the same community. Thus, the patterns of social organization at

both the population scale (2-3 communities) and at a local scale (>50 communities)

were extremely stable year-to-year. This result suggest that multiple levels of

community structure exist in this population. Partitioning the network into larger

number of communities did not result in one community per feeding station, instead

several large communities were maintained and many small ones were created

(Figure 5). Finally, we found no evidence that the composition of species in

communities changed based on how many communities were created (Figure S5).

Thus, the partitioning of the network into more (smaller) communities did not

segregate individuals into species-specific clusters, and so stable community

structure at a local scale was not explained by simple species-level processes.

Figure 4: The social network for each year of the study partitioned into 3

communities. Each point represents one individual, and its colour represents the

community it is assigned into. The size of each point represents its weighted degree

(larger points have more and/or stronger connections to other individuals). Points

are drawn at the average location that the individual was observed, with a small

amount of jittering added to reduce the overlap between individuals observed in the

same location.

Figure 5: The social network for each year of the study partitioned into 65

communities. Each point represents one individual, and its colour represents the

community it is assigned into. The size of each point represents its weighted degree

(larger points have more and/or stronger connections to other individuals). Points

are drawn at the average location that the individual was observed, with a small

amount of jittering added to reduce the overlap between individuals observed in the

same location.

Linking movement patterns to community structure

What drives the consistent patterns of community structure in the social network of

this population? Understanding how communities emerge in the social networks is

important for determining which behaviours (i.e. mechanisms) group-living could

act on as agents of selection. Uncovering the processes that drive community

structure becomes notably more complex when studying species where individuals

form fission-fusion groups, as no single individual is likely to be responsible for the

global patterns of connection. Instead social structure arises from the interactions

among all individuals taken together. One plausible mechanism that is likely to

underpin some aspects of community structure is the patterns of movements.

Individuals living at two locations with frequent movements of individuals between

them will be more likely to be connected in the social network, and therefore more

likely to be in the same community and share similar social environments.

We investigate whether the regular movements of individuals between particular

feeding stations are responsible for global community structure (see Methods). We

found that when we partitioned the network into few communities, the individuals

at feeding stations connected by many movements occurring were very likely to be

allocated to the same community (Figure 6). This supports our hypothesis that the

extrinsic habitat features shape the broad patterning of the community (i.e. the

presence of 2-3 distinct clusters of individuals, see Figure 3). However, at more local

scale, we found that the assortativity coefficient decreased (Figure 6). Thus, as the

social network is partitioned into more communities, movements between sites

explained less of the community structure, despite the fact that individuals become

more likely to re-occur in the same communities across years (see Figure 4).

Figure 6: The correlation of movements between feeding stations and community

structure decreases with increasing numbers of communities. Each feeding station

is allocated to the community in which the majority of individuals are members and

used as a trait value to calculate assortment using the raw movement networks

(Figure 1A) and the per capita movement network (Figure S2A). High values

represent stronger connections between feeding stations in the same community.

DISCUSSION

Our study revealed two levels of social structure in a large community of wild birds

containing several thousands of individuals from five species. At a broad scale, the

social network contained two to three communities that were easily predicted by

the regular movement paths used by birds. The movement of birds through the

woodland were repeatable each year, but the similarity in movements across years

was only partly by the geometry of the study area and fine-scale variation in habitat

structure. Our results suggest that some other processes, potentially social

processes such as local traditions [12, 40], may also be involved. If that is the case, then

broad-scale social structure could be, in part, the result of a socially-transmitted

inter-generational effect. At a more local scale, we found highly stable social

structure, with local clusters of individuals from all five species re-associating each

year to maintain consistent communities. Our study highlights how multiple factors

shape the social ecology in a population of wild birds.

The link between extrinsic habitat factors and community structure in animal

populations has been investigated before. For example, community and sub-

community structure in Galapagos sealions *Zalophus wollebaeki* are largely driven

by the structure of male territories [41]. However, territorial behaviours are unlikely

to play a major role in structuring the winter population of birds in Wytham Woods

because the majority of individuals were great tits and blue tits, which are both non-

territorial during the winter. In non-territorial wintering golden-crowned sparrows

(*Zonotrichia atricapilla*), social network communities were also found to be more

structured than expected given the spatial overlap in the home-range of birds [42]. The

surprising aspect from that study is that golden-crowned sparrows maintain a

stable community structure in their wintering range despite having migrated a long

distance from their breeding areas. The phenomenon observed in both tits and

sparrows suggests that winter sociality is likely to play an important role that goes

beyond simple group size effects, and thus could have carry-over effects into

breeding performance in the following spring (e.g. [7]).

A potentially important feature that we extracted in our study was differential

movement patterns between classes of individuals. We found that juveniles typically

made more long-distance movements than adults. This pattern, which is likely to

arise from juvenile dispersal behaviour, has a number of implications for social

processes. To overcome strong seasonal changes in the environment, juvenile tits

rely on learning from adults [43]. As they move through the landscape, juveniles copy

the adults in their local environment, and a recent study demonstrated that tits

exhibit a strong conformist (copy the local majority) social learning strategy [12].

However, stability in local community structure means that local traditions

(behaviours that differ among different social groups in a population) can easily

become entrenched [12]. Juveniles could therefore play an important role in

facilitating transfer of new information into social groups. At the same time,

juveniles are likely to come into contact with a greater number of individuals, and

thus could play an important role in spreading diseases or pathogens across

communities (as suggested in humans [44]).

By investigating the stability of community structure at different scales, we found

evidence that tits in Wytham Woods live in a multi-level community structure.

Multi-level community structure occurs when animals form small groups, or

clusters, of individuals with whom they associate most strongly, and a larger groups

composed of individuals with whom they associate loosely or indirectly. There is

increasing interest in multi-level community structure as it can have major

implications for how social processes occur [45, 46, 47, 48]. Multiple factors can shape the

movement (or not) of individuals among social units. These factors can be broadly

split into two categories: social factors (such as relatedness [49, 50, 51], cultural

similarity [52], or species identity [45]) and habitat factors (features of the environment

that modulate how individuals move, and thus whom they encounter). Although

recent studies have begun teasing apart social versus habitat factors that determine

the patterns of contact among individuals with different phenotypic characteristics

[37], little is known what drives the emergence of global population-level structure.

Cantor et al [52] recently used simulations to suggest that multi-level communities can

emerge when individual segregate into clans formed around similar cultural

behaviours. In our study, we found evidence that both habitat and social factors

contribute to multi-level community structure. The general geometry of Wytham

Woods introduced a repeatable set of large-scale communities (Figure 3).

Environmental features (notably habitat density) that promote or hinder movement

by individuals (Figure 2) can shape whom individuals encounter. Together, these

features are likely to play a major role in how the population is broadly structured.

In addition, by studying birds that form mixed-species communities, our study

highlights that complex interaction patterns, such as associations among kin or

cooperation, are not pre-requisites for multi-level structure to emerge. Instead,

social mechanisms, such as social preference[37] and phenotypic drivers[35], determine

who individuals affiliate with, while the woodland geometry and habitat density

determine who individuals can come into contact with.

The presence of multi-level community structure can have implications for

evolutionary dynamics of populations. First, restricted movement can reduce gene

flow and lead to divergence in the evolutionary trajectories of sub-parts of each

population. Garant et al. [24] demonstrated that differential dispersal reinforces local

variation in selection for nestling body mass. In their study, they found that trends

in genotypic variance for body mass in nestlings were very different in the eastern

sectors of Wytham Woods and the northern sectors. These two areas represent the

two largest population-level communities we found in our study. Second,

individuals in the same community will have more similar social environments than

individuals occurring in different communities. Thus, any social effects arising via

the social environment, such as indirect genetic effects [53], could accelerate patterns

of divergence within single populations. Finally, the social environment itself can act

as an agent of selection [18, 54], and therefore processes that shape social structure are

likely to impact the overall strength and direction of selection experienced by

populations.

Together the findings from our study highlight how stable social structure can be

maintained in populations as a consequence of both social affiliations and features

of the habitat that individuals live in. The combination of strong clustering together

with some random movements in networks can facilitate the spread of disease or

information through the network [34]. This prediction is supported by evidence from

the rapid spread of a recent disease [55] and of novel traditions [12] in this population.

At the same time, consistent population social structure can lead to phenotypic and

genotypic divergence [24], with potential implications for how animals can adapt to

changing environmental conditions. Integrating information about animal social

structure with data on both pulse and long-term selective events could yield novel

insights into the evolution of social behaviour. However, as our study highlights,

determining the capacity for populations to respond to selective pressures will

require an understanding of a range of different drivers shaping their social

structure.

DATA COLLECTION METHODS

Study location, study species, and population dynamics

The study was undertaken in Wytham Woods, Oxfordshire, UK (51° 46' N, 01° 20'

W), a 385ha area of broadleaf deciduous woodland surrounded by open farmland.

Pairs of birds hold territories during the breeding season (April – June), but form

loose fission-fusion groups during the winter, flocking with unrelated individuals

that forage for ephemeral food sources. Flocks often contain multiple species [38], and

our study also includes data on the five most common flocking species: blue tits,

marsh tits, coal tits, and nuthatches. Tits are generally short-lived—great tits have a

mean life span of 1.9 years. This short generation time results in high annual

population turn-over and inter-annual variation in population sizes. Very good

breeding conditions lead to large population sizes, whereas poor breeding

conditions result in fewer juveniles and a much reduced population size.

Understorey habitat density

We used data from Kirby et al. [32] to quantify the habitat structure between each

feeding site. In that study, the authors recorded, among other measures, the shrub

cover density (0.5m to 2.5m above ground) along the diagonal of 164 different 10m

x 10m quadrats equally spaced throughout Wytham Woods. Here we use data from

the 2012 census, which falls roughly in the middle of our study period. To

extrapolate from the 164 sites, we generated a surface plot where we extrapolated

the data to a 10 x 10m grid of points. The resulting figure accurately captures

variation in habitat density based on our knowledge of the study site. To calculate

habitat density between each pair of feeding sites, we calculated the mean habitat

density along a 20m-wide transect connecting the two sites.

PIT-tagging birds

All birds in the study were caught in either a nest box (as parents and as chicks) or a

mist-net (approximately half the population are birds that immigrate). Each bird

was fitted with uniquely numbered British Trust for Ornithology metal leg ring, and

a uniquely-coded passive integrated transponder (PIT) tag (IB Technologies, UK)

that was fully enclosed in a molded plastic ring fitted to the other leg. Each PIT tag

contains a unique code that can be recorded by antennae (see next section), and

these were then matched to the bird's ring number. We ceased fitting PIT tags to

coal tits from October 2012 as the tags were aggravating pox lesions on birds legs

during a naturally-occurring epidemic. For each bird that was caught and tagged, we

recorded the age and sex (where possible following [56]).

Data collection

We placed 65 automated feeding stations in a evenly-spaced grid covering the

entirety of Wytham Woods and small isolated patches of woodland nearby.

Commercial bird feeders (Jacoby Jayne, UK) were fitted with a radio frequency

identification (RFID) antenna on each of the lower two access holes and other access

holes were blocked. The antennae recorded the unique PIT tag code, time and date

for each visit by a marked bird. For the duration of the study, the feeding stations

were scheduled to open and begin logging at 6am on Saturday mornings, and shut

after dusk on Sunday evening. Feeders were in place December to February in the

winter 2011-12 (13 weeks), from December to early March of winters 2012-13 and

2013-14 (14 weeks each), and for January and February in the winter of 2014-15 (8

weeks). This data collection resulted in 49 unique weekends and 98 complete data

logging days over the 4 winters. In latter years, 6 feeders covering two external sites

were replaced for a separate experiment, and thus the data were not included in

these analyses.

Inferring flocks and flock membership

The data logged from the PIT tag detections produces bursts of detections in the

temporal data stream. These vary in length depending on the size of the flocks

present which increases during the course of the day, [57]. We used a recently-

developed statistical method called Gaussian Mixture Models [58] to extract the start

and end times for each distinct flock. This machine-learning method statistically fits

Gaussian curves of varying sizes to each burst in the data and allocates each record

to the distribution, or 'gathering event', into which it falls. Gaussian Mixture Models

provide a more robust estimation of the social network structure than alternative

methods [59].

Constructing movement networks

From the logging data, we recorded every case of movement by a bird from one

feeding station to another within the same day (a total of 83071 movements over 4

winters), and used these to create movement networks that quantify the

connectedness between each pair of feeding stations in the study. We used daily

sequential detections to maximize our chances of correctly inferring direct

movement pathway (e.g. moving between locations A and D via locations B and C)

and minimizing our chances of incorrectly inferring movement pathways (e.g.

estimating only a direct movement between A and D). Because the number of

movements are inherently linked to number of individuals present at a specific pair

of feeders, we also created a network describing the rate of movement between

feeders, where the rate was defined as the probability that an individual at one focal

feeder would be observed moving to the other focal feeder within a day.

Constructing social networks

We defined edges in the social network using the simple ratio index: $E_{AB} =$

$\frac{x}{x+y_{AB}+y_A+y_B}$, where E_{AB} is the edge weight between individuals A and B, x is the

number of times they were detected in the same flock, y_{AB} is the number of

occasions they were both detected at the same time but not in the same flock, y_A is

the number of detections of A where B was not seen, and y_B is the number of

detections of B where A was not seen. The networks for each year were constructed

using R package *asnipe* [60].

Linking movements to community structure

For each year's social network, we first split the network by specifying $k = 2$ to 65

communities (see main text). For each value of k, we allocated individuals to their

most common feeder, and for each of the feeding stations, select the community that

the majority of individuals present were members of. This enabled us to create a

community label for each feeding station (and for each value of k), and link these to

the network formed by the movement of individuals (Figure 1). To test whether the

movement network shaped the community structure in the network for a given

value of k, we quantified the assortativity coefficient of the network using

communities as discrete trait values [61, 62]. Assortativity is the measure of how well

connected alike nodes are compared to how well connected dislike nodes are,

ranging from 1 (all edges connect nodes with the same traits) to -1 (all edges

connect nodes with different traits).

REFERENCES

1.   Dall SRX*, et al.* Information and its use by animals in evolutionary ecology.
Trends in Ecology and Evolution **20**, 187-193 (2005).
542

543   2.   Seppanen JT, Forsman JT, Monkkonen M, Thomson RL. Social information use
544   is a process across time, space, and ecology, reaching heterospecifics. *Ecology*
545   **88**, 1622-1633 (2007).
546

547   3.   Valone TJ. From eavesdropping on performance to copying the behavior of
548   others: a review of public information use. *Behav Ecol Sociobiol* **62**, 1-14
549   (2007).
550

551   4.   Aplin LM, Farine DR, Mann RP, Sheldon BC. Individual-level personality
552   influences social foraging and collective behaviour in wild birds. *Proc R Soc B*
553   **281**, 20141016 (2014).
554

555   5.   Pruitt JN, Keiser CN. The personality types of key catalytic individuals shape
556   colonies' collective behaviour and success. *Anim Behav* **93**, 87-95 (2014).
557

558   6.   Pruitt JN, Riechert SE. How within-group behavioural variation and task
559   efficiency enhance fitness in a social group. *Proc R Soc B* **278**, 1209-1215
560   (2011).
561

562   7.   Farine DR, Sheldon BC. Selection for territory acquisition is modulated by
563   social network structure in a wild songbird. *J Evolution Biol* **28**, 547-556
564   (2015).
565

566   8.   Formica VA*, et al.* Phenotypic Assortment Mediates the Effect of Social
567   Selection in a Wild Beetle Population. *Evolution* **65**, 2771-2781 (2011).
568

569   9.   Formica VA*, et al.* Fitness consequences of social network position in a wild
570   population of forked fungus beetles (Bolitotherus cornutus). *J Evolution Biol*
571   **25**, 130-137 (2012).
572

573   10.   Wey TW, Burger JR, Ebensperger LA, Hayes LD. Reproductive correlates of
574   social network variation in plurally breeding degus (Octodon degus). *Anim*
575   *Behav* **85**, 1407-1414 (2013).
576

577   11.   Allen J, Weinrich M, Hoppitt W, Rendell L. Network-Based Diffusion Analysis
578   Reveals Cultural Transmission of Lobtail Feeding in Humpback Whales.
579   *Science* **340**, 485-488 (2013).
580

581   12.   Aplin LM*, et al.* Experimentally induced innovations lead to persistent culture
582   via conformity in wild birds. *Nature* **518**, 538-541 (2015).
583

584   13.   Aplin LM, Farine DR, Morand-Ferron J, Sheldon BC. Social networks predict
585   patch discovery in a wild population of songbirds. *Proc R Soc B* **279**, 4199-
586   4205 (2012).
587

588   14.   Farine DR, Aplin LM, Sheldon BC, Hoppitt W. Interspecific social networks
589   promote information transmission in wild songbirds. *Proc R Soc B* **282**,
590   20142804 (2015).
591

592   15.   Adelman JS, Moyers SC, Farine DR, Hawley DM. Feeder use predicts both
593   acquisition and transmission of a contagious pathogen in a North American
594   songbird. *Proc R Soc B* **282**, 20151429 (2015).
595

596   16.   VanderWaal KL, Atwill ER, Isbell LA, McCowan B. Linking social and pathogen
597   transmission networks using microbial genetics in giraffe (Giraffa
598   camelopardalis). *J Anim Ecol* **83**, 406-414 (2013).
599

600   17.   Pruitt JN, Goodnight CJ. Site-specific group selection drives locally adapted
601   group compositions. *Nature* **514**, 359–362 (2014).
602

603   18.   Farine DR, Montiglio PO, Spiegel O. From individuals to groups and back: the
604   evolutionary implications of group phenotypic composition. *Trends Ecol Evol*
605   **30**, 609-621 (2015).
606

607   19.   Goodale E*, et al.* Interspecific information transfer influences animal
608   community structure. *Trends in Ecology and Evolution* **25**, 354-361 (2010).
609

610   20.   Kurvers RHJM*, et al.* The evolutionary and ecological consequences of animal
611   social networks: emerging issues. *Trends Ecol Evol* **29**, 326-335 (2014).
612

613   21.   Farine DR. Proximity as a proxy for interactions: issues of scale in social
614   network analysis. *Anim Behav* **104**, e1-e5 (2015).
615

616   22.   Brockmann D, Helbing D. The Hidden Geometry of Complex, Network-Driven
617   Contagion Phenomena. *Science* **342**, 1337-1342 (2013).
618

619   23.   Leu ST*, et al.* Environment modulates population social structure:
620   experimental evidence from replicated social networks of wild lizards. *Anim*
621   *Behav*, (in press).
622

623   24.   Garant D*, et al.* Evolution driven by differential dispersal within a wild bird
624   population. *Nature* **433**, 60-65 (2005).
625

626   25.   Kays R, Crofoot MC, Jetz W, Wikelski M. Terrestrial animal tracking as an eye
627   on life and planet. *Science* **348**, aaa2478 (2015).
628

629   26.   Krause J*, et al.* Reality mining of animal social systems. *Trends Ecol Evol* **28**,
630   541-551 (2013).
631

632   27.   Strandburg-Peshkin A, Farine DR, Couzin ID, Crofoot MC. Shared decision-
633   making drives collective movement in wild baboons. *Science* **348**, 1358-1361
634   (2015).
635

636   28.   Bonter DN, Bridge ES. Applications of radio frequency identification (RFID) in
637   ornithological research: a review. *J Field Ornithol* **82**, 1-10 (2011).
638

639   29.   Farine DR, Whitehead H. Constructing, conducting, and interpreting animal
640   social network analysis. *J Anim Ecol* **84**, 1144-1163 (2015).
641

642   30.   Fortunato S. Community detection in graphs. *Phys Rep* **486**, 75-174 (2010).
643

644   31.   Krause J, James R, Franks DW, Croft DP. *Animal Social Networks*. Oxford
645   University Press (2015).
646

647   32.   Kirby KJ*, et al.* Changes in the tree and shrub layer of Wytham Woods
648   (Southern England) 1974-2012: local and national trends compared. *Forestry*
649   **87**, 663-673 (2014).
650

651   33.   Dekker D, Krackhardt D, Snijders T. Sensitivity of MRQAP tests to collinearity
652   and autocorrelation conditions. *Psychometrika* **72**, 563-581 (2007).
653

654   34.   Eames KTD. Modelling disease spread through random and regular contacts
655   in clustered populations. *Theor Popul Biol* **73**, 104-111 (2008).
656

657   35.   Aplin LM*, et al.* Individual personalities predict social behaviour in wild
658   networks of great tits (Parus major). *Ecol Lett* **16**, 1365–1372 (2013).
659

660   36.   Aplin LM*, et al.* Consistent individual differences in the social phenotypes of
661   wild great tits (Parus major). *Anim Behav* **108**, 117-127 (2015).
662

663   37.   Farine DR*, et al.* The role of social and ecological processes in structuring
664   animal populations: a case study from automated tracking of wild birds. *Roy*
665   *Soc Open Sci* **2**, 150057 (2015).
666

667   38.   Farine DR, Garroway CJ, Sheldon BC. Social network analysis of mixed-
668   species flocks: exploring the structure and evolution of interspecific social
669   behaviour. *Anim Behav* **84**, 1271-1277 (2012).
670
671   39.   Csardi G, Nepusz T. The igraph software package for complex network
672   research. *InterJournal* **Complex Systems**, 1695 (2006).
673
674   40.   Mueller T*, et al.* Social Learning of Migratory Performance. *Science* **341**, 999-
675   1002 (2013).
676
677   41.   Wolf JBW, Mawdsley D, Trillmich F, James R. Social structure in a colonial
678   mammal: unravelling hidden structural layers and their foundations by
679   network analysis. *Anim Behav* **74**, 1293-1302 (2007).
680
681   42.   Shizuka D*, et al.* Across-year social stability shapes network structure in
682   wintering migrant sparrows. *Ecol Lett* **17**, 998-1007 (2014).
683
684   43.   Slagsvold T, Wiebe KL. Learning the ecological niche. *Proc R Soc B* **274**, 19-23
685   (2007).
686
687   44.   Del Valle SY, Hyman JM, Hethcote HW, Eubank SG. Mixing patterns between
688   age groups in social networks. *Soc Networks* **29**, 539-554 (2007).
689
690   45.   Bell HL, Ford HA. A Comparison of the Social-Organization of 3 Syntopic
691   Species of Australian Thornbill, Acanthiza. *Behav Ecol Sociobiol* **19**, 381-392
692   (1986).
693
694   46.   de Silva S, Wittemyer G. A Comparison of Social Organization in Asian
695   Elephants and African Savannah Elephants. *Int J Primatol* **33**, 1125-1141
696   (2012).
697
698   47.   Grueter CC, Matsuda I, Zhang P, Zinner D. Multilevel Societies in Primates and
699   Other Mammals: Introduction to the Special Issue. *Int J Primatol* **33**, 993-
700   1001 (2012).
701
702   48.   Whitehead H*, et al.* Multilevel Societies of Female Sperm Whales (Physeter
703   macrocephalus) in the Atlantic and Pacific: Why Are They So Different? *Int J
704   Primatol* **33**, 1142-1164 (2012).
705
706   49.   Archie EA, Moss CJ, Alberts SC. The ties that bind: genetic relatedness
707   predicts the fission and fusion of social groups in wild African elephants. *Proc
708   R Soc B* **273**, 513-522 (2006).
709
710   50.   Croft DP*, et al.* The role of relatedness in structuring the social network of a
711   wild guppy population. *Oecologia* **170**, 955-963 (2012).
712

713   51.   Godfrey SS*, et al.* A contact based social network of lizards is defined by low
714   genetic relatedness among strongly-connected individuals. *Anim Behav* **97**,
715   35-43 (2014).
716
717   52.   Cantor M*, et al.* Multilevel animal societies can emerge from cultural
718   transmission. *Nature Communications* **6**, (2015).
719
720   53.   Moore AJ, Brodie ED, Wolf JB. Interacting phenotypes and the evolutionary
721   process .1. Direct and indirect genetic effects of social interactions. *Evolution*
722   **51**, 1352-1362 (1997).
723
724   54.   Wolf JB, Moore AJ. Interacting Phenotypes and Indirect Genetic Effects. In:
725   *Evolutionary Behavioral Ecology* (ed^(eds Westneat DF, Fox CW). Oxford
726   University Press (2010).
727
728   55.   Lachish S, Lawson B, Cunningham AA, Sheldon BC. Epidemiology of the
729   Emergent Disease Paridae pox in an Intensively Studied Wild Bird
730   Population. *Plos One* **7**, (2012).
731
732   56.   Svensson L. *Identification Guide to European Passerines*, 4th edition edn. BTO
733   (1992).
734
735   57.   Farine DR, Lang SDJ. The early bird gets the worm: foraging strategies of wild
736   songbirds lead to the early discovery of food sources. *Biol Letters* **9**,
737   20130578 (2013).
738
739   58.   Psorakis I, Roberts SJ, Rezek I, Sheldon BC. Inferring social network structure
740   in ecological systems from spatio-temporal data streams. *J R Soc Interface* **9**,
741   3055-3066 (2012).
742
743   59.   Psorakis I*, et al.* Inferring social structure from temporal data. *Behav Ecol*
744   *Sociobiol* **69**, 857-866 (2015).
745
746   60.   Farine DR. Animal Social Network Inference and Permutations for Ecologists
747   in R using asnipe. *Methods Ecol Evol* **4**, 1187–1194 (2013).
748
749   61.   Farine DR. Measuring phenotypic assortment in animal social networks:
750   weighted associations are more robust than binary edges. *Anim Behav* **89**,
751   141-153 (2014).
752
753   62.   Shizuka D, Farine DR. Measuring the robustness of network community
754   structure using assortativity. *Anim Behav* **122**, 237-246 (2016).
755
756
757

A

B

C

N

0   500

m

●   60

·   20

━━   900

──   300