1

Stable multi-level social structure is maintained by habitat

geometry in a wild bird population

Damien R. Farine*[a,b,c,d], Ben C. Sheldon[a]

[a] Edward Grey Institute of Field Ornithology, Department of Zoology, University of Oxford,

South Parks Road, Oxford OX1 3PS, United Kingdom.

[b] Department of Collective Behaviour, Max Planck Institute for Ornithology,

Universitätsstrasse 10, 78457 Konstanz, Germany.

[c] Chair of Biodiversity and Collective Behaviour, Department of Biology, University of

Konstanz, Universitätsstrasse 10, 78457 Konstanz, Germany.

[d] Center for the Advanced Study of Collective Behaviour, University of Konstanz,

Universitätsstrasse 10, 78457 Konstanz, Germany.

* Corresponding author:

dfarine@orn.mpg.de

ABSTRACT

Social structure can have profound evolutionary and ecological implications for

animal populations. Structure can arise and be maintained via social preferences or

be indirectly shaped by habitat structure. Understanding how social structure

emerges is important for understanding the potential links between social structure

and evolutionary and ecological processes. Here, we study a large community of

wild birds fitted with uniquely-coded passive integrated transponder (PIT) tags and

recorded on a grid of automated feeders fitted with radio frequency identification

(RFID) antennae. We show that both large-scale and fine-scale network

communities are consistent across years in this population, despite high

generational turn-over. Studying the process that generates community structure,

here the movement of individual birds across the woodland, suggests an important

role of habitat geometry in shaping population-level social community structure.

Our study highlights how relatively simple factors can produce apparent emergent

social structure at the population scale, which has widespread implications for

understanding eco-evolutionary dynamics.

INTRODUCTION

The social environment can profoundly shape the life histories of animals. Who

individuals associate with can determine the information they have access to (Dall

et al. 2005; Seppanen *et al.* 2007; Valone 2007), affect how well they can exploit

resources (Pruitt & Riechert 2011; Aplin *et al.* 2014; Pruitt & Keiser 2014), and

impact their ability to successfully reproduce (Formica *et al.* 2011; Formica *et al.*

2012; Wey *et al.* 2013; Farine & Sheldon 2015). Social structure can also influence

population processes, such as the spread of information within (Aplin *et al.* 2012;

Allen *et al.* 2013; Aplin *et al.* 2015a) and between (Farine *et al.* 2015a) species, and

the spread of disease among individuals (VanderWaal *et al.* 2013; Adelman *et al.*

2015). Finally, individuals can experience selection arising from properties of their

social groups (Formica *et al.* 2011; Farine & Sheldon 2015) or their communities

(Pruitt & Goodnight 2014). Quantifying the factors that shape individuals' social

environments is a key step in identifying the evolutionary drivers of sociality

(Goodale *et al.* 2010; Farine, Montiglio & Spiegel 2015), as the mechanisms that

determine where, and with whom, individuals live their lives are what selection

arising from population processes (e.g. via disease burden or information use) can

act upon.

Social structure (the pattern of connections that emerges from interactions among

individuals, often represented using social networks) is generally considered to

arise from individuals making decisions about with whom to associate (Hinde 1976;

Whitehead 2008; Kurvers *et al.* 2014). The overall structural properties of a

population are likely to be emergent outcomes of the combined behaviours and

social decisions of all individuals in the population, which for most animals includes

heterospecifics. A number of recent studies have highlighted potential multi-level

social structure in animal populations (Wittemyer, Douglas-Hamilton & Getz 2005;

Archie, Moss & Alberts 2006; Schreier & Swedell 2009), with smaller 'groups' or

'communities' of individuals being embedded in larger communities containing

multiple groups. The implications of such population structure are profound, for

example individuals within a community would experience more similar modulating

effects of the social environment on selection relative to individuals from different

communities (Montiglio, McGlothlin & Farine 2018). However, we still know

relatively little about how multi-level societies emerge.

Because spatial proximity is a necessary requirement for associations or

interactions (i.e. connections) among individuals in many species of animals (Farine

2015), patterns of connections among individuals are bound to be shaped by a

number of extrinsic factors, such as geometry and structure of habitat features that

promote or restrict individual movements (Wiens 1976; O'Brien *et al.* 2006; Cox *et*

al. 2016; He, Maldonado-Chaparro & Farine 2019). At a local scale, habitat structure,

such as the presence of understorey density, can channel individual movements,

thus increasing the propensity for contact among individuals. At a broader scale,

movement 'highways' can significantly reduce the social distance among individuals

despite a large spatial separation, and therefore facilitate the flow of information or

disease (Brockmann & Helbing 2013). For example, sleepy lizards (*Tiliqua rugosa*)

living in open habitats had fewer contacts with conspecifics than those living in

more structured habitats (Leu *et al.* 2016). Whilst habitat features can promote

movements, the geometry of habitat (such as the boundary of fragmentized habitat)

can also introduce barriers that constrain individual movements. Such barriers can

result in a structure where the social distance between two individuals (i.e. their

degrees of separation in a social network) on either side of this barrier could be

much greater than their actual spatial distance. Habitat geometry and features can

therefore impose structure in a social network that could easily be interpreted as

arising socially. Given that local heterogeneity in gene flow can lead to rapid

evolutionary differentiation (Garant *et al.* 2005), integrating knowledge about fine-

scale environmental heterogeneity into studies of social structure could

fundamentally alter our understanding of adaptation and the ability for animals to

respond to selective pressures.

A major challenge in identifying extrinsic factors that drive population social

structure is the need to track the movement of individuals in space, patterns of

social connections among the majority of individuals in a population, together with

information about the habitat in which they live (Strandburg-Peshkin *et al.* 2017).

Recent technological advances have eased the logistical constraints of sampling

many individuals moving across large areas (Tomkiewicz *et al.* 2010; Rutz *et al.*

2012; Ryder *et al.* 2012; Krause *et al.* 2013; Finn *et al.* 2014; Kays *et al.* 2015; Levin

et al. 2015; Strandburg-Peshkin *et al.* 2015; Jacoby & Freeman 2016). In particular,

passive integrated transponder (PIT) tags are cheap electromagnetic tags that can

be fitted to many individuals at once (Bonter & Bridge 2011), thus overcoming the

challenges associated with studying entire animal communities. Because PIT tags do

not rely on battery power to emit a signal, but instead are detected by affecting the

magnetic field in radio frequency identification (RFID) antennas, they provide the

capability to track individuals across years and life stages. Further, advances in the

analytical tools, such as social network analysis (Farine & Whitehead 2015) and

community detection algorithms (Fortunato 2010), are facilitating greater insight

into patterns of social structure. These technological advances have underpinned a

boom in the study of animals' social networks (Krause *et al.* 2015), and in networks

formed by the movement of individuals across space (Finn *et al.* 2014; Jacoby &

Freeman 2016). However, much less effort has focused on identifying the key

mechanisms that shape overall social structure at the scale of populations.

In this paper, we investigate whether habitat structure affects patterns of

movement, and in turn drive social structure, in a large population comprising 5

species of wild songbird. A number of previous studies in our population have

investigated how genetic traits (Radersma *et al.* 2017), phenotypic traits such as

age, personality, sex, and immigration status (Quinn *et al.* 2011; Aplin *et al.* 2013;

Aplin *et al.* 2014; Farine 2014; Aplin *et al.* 2015b; Farine *et al.* 2015b), species

identity (Farine, Garroway & Sheldon 2012; Farine *et al.* 2014), and diel cycles

(Farine & Lang 2013; Hilleman *et al.* in press) affect individual-level movement and

social behaviour, and that who individuals associate with can impact their fitness

(Farine & Sheldon 2015). Using data from this population collected over 4

consecutive winters, we first quantify overall patterns of movements across the

landscape each winter to determine whether individuals move evenly through

space, or if there are consistent movement corridors. Based on studies showing that

birds are sensitive to open habitat (Quinn *et al.* 2012) (which heighten the risk of

predation), we predict that birds should show a preference for moving along

corridors comprising denser understory habitat. Second, we use patterns of

associations among individuals in each winter to construct social networks, and use

these social networks to determine whether social structure is consistent across

winters. Given that approximately 50% of individuals in the population each winter

are first-year birds, and approximately 50% of first-year birds are immigrants into

the population (born outside the study area), we predict that there would be no

consistent social structure at local scales. However, because the study area

remained identical across winters, we predict that any effects of the habitat on the

social structure of the population would be repeated each winter.

METHODS

Study location, study species, and population dynamics

The study was undertaken in Wytham Woods, Oxfordshire, UK (51° 46' N, 01° 20'

W), a 385ha area of broadleaf deciduous woodland 'island' surrounded by intensive

agriculture. Pairs of birds hold territories during the breeding season (April – June),

but form loose fission-fusion groups during the winter, flocking with unrelated

individuals that forage for ephemeral food sources. Flocks often contain multiple

species (Farine, Garroway & Sheldon 2012), and our study also includes data on the

five most common flocking species: great tits (*Parus major*), blue tits (*Cyanistes*

caeruleus), marsh tits (*Poecile palustris*), coal tits (*Periparus ater*), and nuthatches

(*Sitta europaea*). Tits are generally short-lived—great tits have a mean life span of

1.9 years. This short generation time results in high annual population turn-over

and inter-annual variation in population sizes. Good breeding conditions lead to

large population sizes, whereas poor breeding conditions result in fewer juveniles

and a much reduced population size.

PIT-tagging birds

All birds in the study were caught in either a nest box (as parents and as chicks) or a

mist-net (approximately half the population are birds that immigrate). Each bird

was fitted with uniquely numbered British Trust for Ornithology metal leg ring, and

a uniquely-coded passive integrated transponder (PIT) tag (IB Technologies, UK)

that was fully enclosed in a moulded plastic ring fitted to the other leg. Each PIT tag

contains a unique code that can be recorded by antennae (see next section), and

these were then matched to the bird's ring number. We ceased fitting PIT tags to

coal tits from October 2012 as the tags were aggravating pox lesions on birds legs

during a naturally-occurring epidemic. For each bird that was caught and tagged, we

recorded the age and sex (where possible following Svensson 1992).

Data collection

We placed 65 automated feeding stations in a evenly-spaced grid covering the

entirety of Wytham Woods and small isolated patches of woodland nearby.

Commercial bird feeders (Jacobi Jayne, UK) were fitted with a radio frequency

identification (RFID) antenna on each of the lower two access holes and other access

holes were blocked. The antennae recorded the unique PIT tag code, time and date

for each visit by a marked bird. For the duration of the study, the feeding stations

were scheduled to open and begin logging at 6am on Saturday mornings, and shut

after dusk on Sunday evening. Feeders were in place December to February in the

winter 2011-12 (13 weeks), from December to early March of winters 2012-13 and

2013-14 (14 weeks each), and for January and February in the winter of 2014-15 (8

weeks). This data collection resulted in 49 unique weekends and 98 complete data

logging days over the 4 winters. In later winters, 6 feeders covering two external

sites were replaced for a separate experiment, and thus the data were not included

in these analyses.

Constructing movement networks

From the logging data, we recorded every case of movement by a bird from one

feeding station to another within the same day (a total of 83071 movements over 4

winters), and used these to create movement networks that quantify the

connectedness between each pair of feeding stations in the study. We used daily

sequential detections to maximize our chances of correctly inferring direct

movement pathway (e.g. moving between locations A and D via locations B and C)

and minimizing our chances of incorrectly inferring movement pathways (e.g.

estimating only a direct movement between A and D). Because the number of

movements are inherently linked to the number of individuals present at a specific

pair of feeders, we also created a network describing the rate of movement between

feeders, where the rate was defined as the probability that an individual at one focal

feeder would be observed moving to the other focal feeder within a day.

Understory habitat density

We used data from Kirby et al. (Kirby *et al.* 2014) to quantify the habitat structure

between each feeding site. In that study, the authors recorded, among other

measures, the shrub cover density (0.5m to 2.5m above ground) along the diagonal

of 164 different 10m x 10m quadrats equally spaced throughout Wytham Woods.

Here we use data from the 2012 census, which falls roughly in the middle of our

study period, but note that there were no major changes to the habitat across years.

To extrapolate from the 164 sites, we generated a surface plot where we

extrapolated the data to a 10 x 10m grid of points using the vgm function in the R

package gstat (Gräler, Pebesma & Heuvelink 2016) with a spherical model, omitting

fitting the nugget component. The resulting figure accurately captures variation in

habitat density based on our knowledge of the study site. To calculate habitat

density between each pair of feeding sites, we calculated the mean habitat density

(as estimated by the surface) along a 20m-wide transect connecting the two sites.

Inferring flocks and flock membership

The data logged from the PIT tag detections produces bursts of detections in the

temporal data stream. These vary in length depending on the size of the flocks

present (which increases during the course of the day, Farine & Lang 2013;

Hilleman *et al.* in press). We used a recently-developed statistical tool for analyzing

such data involving Gaussian Mixture Models (Psorakis *et al.* 2012) to extract the

start and end times for each distinct flock. This machine-learning method

statistically fits Gaussian curves of varying sizes to each burst in the data and

allocates each record to the distribution, or 'gathering event', into which it falls. We

have shown that the Gaussian Mixture Model approach provides a more robust

estimation of the social network structure than alternative methods (Psorakis *et al.*

2015).

Constructing social networks

We defined edges in the social network using the simple ratio index (Hoppitt &

Farine 2018): $E_{AB} = \frac{x}{x + y_{AB} + y_A + y_B}$, where E_{AB} is the edge weight between individuals

A and B, x is the number of times they were detected in the same flock, y_{AB} is the

number of occasions they were both detected at the same time but not in the same

flock, y_A is the number of detections of A where B was not seen, and y_B is the

number of detections of B where A was not seen. The networks for each winter

were constructed using R package *asnipe* (Farine 2013).

Detecting consistent social structure across winters

For each winter's social network, inferred structural communities in each social

network using the edge betweenness community detection algorithm in the R

package *igraph* (Csardi & Nepusz 2006). We specified the algorithm to detect k = 2

to 65 communities, and recorded the identities of individuals in each community. If

communities are structured exclusively by extrinsic factors, we expected a drop in

the stability of co-membership by individuals as we created finer grained

communities. For example, if a population is spread across three isolated patches of

woodland, and birds do not move between woodlands, then we expect that birds

will always occur within the same three communities (one for each woodland patch)

each winter. By contrast, if communities are structured socially, then we expected

smaller communities (local cliques) to be more stable, but little stability in global

structure. For example, for a territorial pair-living species living in a lattice-like

uniform environment, an algorithm will be able to isolate each pair when identifying

k=N/2 communities, whereas the communities detected for smaller k values will be

essentially random.

Linking movements to community structure

To test for a link between community structure and movement networks, we

allocated individuals to their most common feeder, and for each of the feeding

stations, selected the community of which the majority of individuals present were

members. This enabled us to create a community label for each feeding station (and

for each value of k), and link these to the network formed by the movement of

individuals (Figure 1). To test whether the movement network shaped the

community structure in the network for a given value of k, we quantified the

assortativity coefficient of the network using communities as discrete trait values

(Farine 2014; Shizuka & Farine 2016). Assortativity is the measure of how well

connected alike nodes are compared to how well connected dislike nodes are,

ranging from 1 (all edges connect nodes with the same traits) to -1 (all edges

connect nodes with different traits).

RESULTS

Movement networks

We first quantified the daily movement rates between different feeding stations. Our

population consisted of a total of 6299 unique PIT-tagged individuals, of which 2230

were great tits, 3304 were blue tits, 237 were marsh tits, 169 were coal tits, and 78

were nuthatches. We detected 83701 movements, detections at two feeders on the

same day, across all winters (Figure 1). Overall, birds moved very little—our

analysis was based on 9612 'bird winters', meaning that each bird moved (on

average) fewer than 10 times over a whole winter. We found that most movements

occurred between feeding stations where more birds were present.

Not all birds followed the same pattern. Birds in their first winter moved more often

than older birds: first-year birds accounted for 61% of all movements, despite

making up only made up 38% of the population on average. Of the juveniles that

moved, each made on average 20.8 moves, which was significantly more than the

average 12.9 moves made by adults detected moving at least once (GLM:

$\beta \pm SE=0.477 \pm 0.007$, $z=67.08$, $P<0.001$, family=Poisson). Further, their movement

networks included more long-distance movements than those of adults, resulting in

a significantly higher average movement distance ($\beta \pm SE=29.5 \pm 1.35$,

$t=21.77, P<0.001$, see also Figures 1 and S1).

We also found species-level differences. Great and blue tits, which comprise the

large majority of individuals in the population, exhibited similar movement patterns

(difference=0.5m, P=0.520, see Figures S2-3). By contrast, marsh tit movements

were typically much more localized (difference blue tit – marsh tit: 48.8m, P<0.001;

difference great tit – marsh tit: 29.3m, P<0.001; see Figures S2-3). Overall, age had

the greatest effect on the distribution of movements, with some differences among

species.

Figure 1: Total movements of (A) all birds, (B) adults, and (C) juveniles (first

winters) from all species over 4 winters of data. The thickness of each line

represents the number of observations of a bird detected at two feeding stations

(black points) in the same day. The size of the points represents the average number

of individuals observed at each feeding station. The green outline represents the

outline of the wooded areas, which are surrounded by open agricultural land. Note

that there is very little movement of birds between Wytham Woods and the four

external woodlots, highlighting how closed this population is during the winter. See

Figure S1 for statistical evidence that juveniles make significantly more long-

distance movements.

Do understory density or habitat geometry predict movement propensity?

We compared the patterns of movement observed across winters, and related these

to spatial distance and habitat features. We first re-defined edges in the movement

network as the probability that an individual detected at either site moves between

them on a given day. Edge weights ranged from 0 representing no movement

between adjacent sites to 0.83 representing that each individual detected had an

83% chance of moving between the two sites on a given day (see Figure S4).

Weighted multiple regression quadratic assignment procedure (MRQAP)(Dekker,

Krackhardt & Snijders 2007) revealed that the movements by birds between feeding

stations were significantly more similar from winter-to-winter than expected by

chance, even when accounting for distance through the forest and habitat structure

(see Table 1, Figure 2). In all winters, birds were significantly more likely to move

between 'close' feeding stations than distant ones. However, contrary to our

predictions, we found no effect of understory habitat density on the propensity for

birds to move between feeders.

Table 1: Results of multiple regression quadratic assignment procedure used to test

whether previously observed patterns of movement (probability of moving between

sites per capita), distance between sites (through the forest), and habitat density

(percentage cover) between sites explain the observed patterns of movement. Bold

values represent significant coefficients, * represents significance at $P < 0.01$, **

represents significance at $P < 0.001$, based on a two-tailed test. All variables are

scaled to 0 mean and unit variance to enable comparison between effect sizes.

Winter	Previous Winter	Forest Distance	Habitat Density
2013	**0.903****	**-0.028***	-0.010
2014	**0.877****	**-0.060****	-0.010
2015	**0.787****	**-0.091****	-0.019

Figure 2: Total movement of all birds (from Figure 1A) overlaid on the understorey habitat density (0.5m to 2.5m above ground) calculated at 164 equally-spaced quadrats (see Methods). Areas within the polygon with a white background have no habitat data available, and any links that included such areas were excluded from the analyses. Areas outside of the black polygons are open agricultural land. Blue areas have a largely open understory, whereas green areas have very thick understory. The vast majority of Wytham Woods is closed canopy.

Because for many pairs of feeding stations the geometric distance and forest distance are very similar see Figure 2), we used this information to also test whether birds were less likely to move between feeding sites that were separated by non-forest. We did this by calculating the difference between forest distance and Euclidian distance. We found that birds moved less between feeders with a larger difference in distance (i.e. the path through the forest was much longer than the straight-line path), suggesting that birds are avoiding crossing open habitats (Table 2). However, the propensity to move between feeders observed in previous winters

was consistently the strongest predictor of future movements. The coefficient values

for movements predicted by the previous winter were typically an order of

magnitude larger than those of other predictor variables, suggesting that additional

undetected factors—potentially social—are driving patterns of movements by birds

across this woodland.

Table 2: Results of multiple regression quadratic assignment procedure used to test

whether birds moved less between feeding sites separated by open space. For each

pair of sites, we calculated the difference between the forest distance and the

Euclidian distance. Bold values represent significant coefficients, * represents

significance at P < 0.01, ** represents significance at P < 0.001, based on a two-tailed

test. All variables are scaled to 0 mean and unit variance to enable comparison

between effect sizes.

Winter	Previous Winter	Relative Distance
2013	**0.919****	-0.001
2014	**0.906****	**-0.019***
2015	**0.828****	**-0.029***

Does the population have consistent social structure?

Population-level social structure can have significant implications for population

processes. For example, high levels of clustering can reduce the spread of disease

within populations (Eames 2008). Thus far, we have shown that habitat geometry,

as well as additional unknown factors, contribute to consistent patterns of animal

movements. We next investigated how these patterns contribute to the emergent

structure of the mixed-species population.

We uncovered two scales that maximized the propensity for pairs of individuals

observed in the same community in winter t to be observed in the same community

in winter $t + 1$. When social networks were partitioned into 2 or 3 communities

(Figure 3), individuals observed in successive winters were seen in the same

community each winter approximately 90% of the time. These macro communities
largely reflect the geometry of the study site, with two core habitats (north-west and
the east), and a smaller patch of forest to the south, that is only attached by a narrow
neck of vegetation, representing a third community (Figure 4). Specifying the
algorithm to detect 4 communities significantly decreased the probability of
individuals being re-observed in the same community (no overlap in the ranges in
Figure 3), suggesting that there is no stable 4th community. Partitioning the network
further by specifying the algorithm to detect more than 4 communities then
increased the probability that two individuals observed in successive winters were
observed in the same community. Thus, the patterns of social organization at both
the population scale (2-3 communities) and at a local scale (>30 communities) were
the most stable winter-to-winter. This result suggest that multiple levels of
community structure exist in this population, with micro communities nested within
macro communities.
406

407
Figure 3: The probability that two birds observed in the same community in one
winter remain in the same community in the following winter, given that both are
observed. Points represent the mean, and lines represent the range from the 3 pairs

of winters. The probability is calculated with the social network partitioned into the

same specified number of communities (2 to 65) in all winters.

Partitioning the network into a larger number of communities did not result in one

community per feeding station, instead several large communities were maintained

and many small, spatially-overlapping, ones were created (Figure 5). Finally, we

found no evidence that the composition of species in communities changed based on

how many communities were created (Figure S5). Thus, the partitioning of the

network into more (micro) communities did not segregate individuals into species-

specific clusters, and so stable community structure at a local scale was not

explained by simple species-level processes. We note here that there is extensive

evidence of social processes driving community structure at these local scales

(Krause *et al.* 2015). Such studies of animal social networks are increasingly

accounting for effects such as individual home-ranges using null model (Farine

2017). However, fewer studies have explored links between home ranges and

community structure (but see Shizuka *et al.* 2014). Our study highlights the

importance of exploring this link.

Figure 4: The social network for each winter of the study when partitioned into 3

communities. Each point represents one individual (N=3019, 2598, 2294, 1701

respectively), and the colours represents the community each node is assigned to.

The size of each point represents its weighted degree (larger points have more

and/or stronger connections to other individuals). Points are drawn at the average

location that the individual was observed, with a small amount of jittering added to

reduce the overlap between individuals observed in the same location. Community

memberships were inferred using the edge betweenness algorithm applied to each

winter independently with the number of communities set to 3.

Figure 5: The social network for each winter of the study partitioned into 65

communities. Each point represents one individual, and colours represents the

community each node is assigned into (per Figure 4). The size of each point

represents its weighted degree (larger points have more and/or stronger

connections to other individuals). Points are drawn at the average location that the

individual was observed, with a small amount of jittering added to reduce the

overlap between individuals observed in the same location. Community

memberships were inferred using the edge betweenness algorithm applied to each

winter independently with the number of communities set to 65.

Linking movement patterns to community structure

We investigated whether the regular movements of individuals between particular

feeding stations were responsible for global community structure. Individuals living

at two locations with frequent movements of individuals between them will be more

likely to be connected in the social network, and therefore more likely to be in the

same community and share similar social environments. We found that when we

partitioned the network into few communities, almost all of the movements were

between feeders in the same community (Figure 6). This supports our hypothesis

that extrinsic large-scale habitat features shape the broad patterning of the

community (i.e. the presence of 2-3 distinct clusters of individuals, see Figure 4) via

individual movement (i.e. by disconnecting feeders not directly connected by

forest). However, at more local scale, we found that the assortativity coefficient

decreased (Figure 6). Thus, as the social network is partitioned into more

communities, movements between sites explained less of the community structure,

despite the fact that individuals become more likely to re-occur in the same

communities across winters (see Figure 3).

Figure 6: The correlation of movements between feeding stations and community

structure decreases with increasing numbers of communities. Each feeding station

is allocated to the community in which the majority of individuals are members and

used as a trait value to calculate assortment using the raw movement networks

(Figure 1A) and the per capita movement network (Figure S2A). High values

represent stronger connections between feeding stations in the same community.

Points represent the mean, and lines represent the range across the 4 winters.

DISCUSSION

Our study revealed two levels of social structure, maintained across winters, in a

large population of wild birds containing thousands of individuals of five species. At

a broad scale, the social network contained two or three communities that were

predicted by the regular movement paths used by birds. The movement of birds

through the woodland were repeatable each winter, but the similarity in

movements across winters was only partly explained by the geometry of the study

area. Our analyses suggest, in fact, that some other processes, potentially social

processes such as the persistence of local traditions (Mueller *et al.* 2013; Aplin *et al.*

2015a; Jesmer *et al.* 2018), may also be involved. If that is the case, then broad-scale

social structure could be, in part, the result of a socially-transmitted inter-

generational effect. At a more local scale, we found highly stable social structure,

with micro communities of individuals comprising all five species re-associating

each winter to maintain consistent communities. Our study thus highlights how

factors operating at different scales can shape the social ecology in a wild bird

population.

The link between extrinsic habitat factors and community structure in animal

populations has been investigated before. For example, community and sub-

community structure in Galapagos sealions *Zalophus wollebaeki* are largely driven

by the structure of male territories (Wolf *et al.* 2007). However, territorial

behaviours are unlikely to play a major role in structuring the winter population of

birds in Wytham Woods because the majority of individuals were great tits and blue

tits, which are both non-territorial during the winter, although there is some

evidence of linkages between summer and winter social structure (Farine & Sheldon

2015; Firth & Sheldon 2016). What is more likely is that individuals have relatively

small homeranges that overlap with many other individuals, and that these are not

uniformly distributed across the habitat. In non-territorial wintering golden-

crowned sparrows (*Zonotrichia atricapilla*), social network communities were also

found to be consistent across winters at very small spatial scales (Shizuka *et al.*

2014). The surprising aspect from that study was that golden-crowned sparrows

exhibit such stability despite having migrated a long distance from their breeding

areas. This aspect is partly replicated in our tit population as approximately 50% of

the individuals in any given winter are first-year birds and most birds leave the

study area during the summer (Matechou *et al.* 2015). The phenomenon observed in

both tits and sparrows suggests that winter sociality is likely to play an important

role that goes beyond simple group size effects, and thus could have carry-over

effects into the territory structure (Firth & Sheldon 2016) and breeding

performance in the following spring (e.g. Farine & Sheldon 2015).

The unexplained consistent structural patterns in both our study and in the golden-

crowned sparrow study (Shizuka *et al.* 2014) could represent local traditions that

are passed on through social learning. In this scenario, juvenile and immigrant birds

copy the movement behaviours of older, resident, birds. This could explain why

patterns remained consistent over 4 winters, well beyond the generation time of tits

(typically <2 years). Such findings would not be unprecedented. For example, one

study in whooping cranes found that juveniles socially learn the migration routes

from older individuals (Mueller *et al.* 2013). A previous study in our population

showed that social learning can easily lead to persistent local cultures, and that the

presence of experienced individuals facilitates the rapid adoption of new behaviours

by the next generation (Aplin *et al.* 2015a). The role of such traditions in shaping

animal movements and subsequent community structure warrant much further

investigation.

Another potentially important feature that we extracted in our study was

differential movement patterns between classes of individuals. We found that

juveniles typically made more long-distance movements than adults (Figures 1 &

S1). This pattern, which is likely to be linked to juvenile dispersal behaviour, has a

number of implications for social processes. To overcome strong seasonal changes

in the environment, juvenile tits rely on learning from adults in their local

environment (Slagsvold & Wiebe 2007). These juveniles therefore play an important

role in shaping the overall structure of the social network, and could play a major

role as transmission vectors. By coming into contact with a greater number of

individuals, they could facilitate the spread diseases or pathogens across

communities (as suggested in humans, Del Valle *et al.* 2007), or even introduce

novel behaviours into populations.

By investigating the stability of community structure at different scales, we found

evidence that tits in Wytham Woods live in a multi-level community structure.

Multi-level community structure occurs when animals form small groups, or

clusters, of individuals with whom they associate most strongly, and larger groups

in which these clusters are embedded. There is increasing interest in multi-level

community structure as it can have major implications for how social processes

occur (Bell & Ford 1986; de Silva & Wittemyer 2012; Grueter *et al.* 2012; Whitehead

et al. 2012). Multiple factors can shape the movement (or not) of individuals among

social units. These factors can be broadly split into two categories: social factors

[such as relatedness (Archie, Moss & Alberts 2006; Croft *et al.* 2012; Godfrey *et al.*

2014), cultural similarity (Cantor *et al.* 2015), or species identity (Bell & Ford

1986)] and habitat factors [features of the environment that modulate where in the

environment individuals are found (Croft *et al.* 2003), how they move, and thus

whom they encounter (He, Maldonado-Chaparro & Farine 2019)]. Although recent

studies have begun to tease apart social versus habitat factors that determine the

patterns of contact among individuals with different phenotypic characteristics

(Farine *et al.* 2015b), little is known what drives the emergence of global

population-level structure. Cantor *et al.* (2015) used simulations to suggest that

multi-level communities can emerge when individual segregate into clans formed

around similar cultural behaviours. In our study, we found evidence that both

environmental and social factors contribute to producing a hierarchical community

structure. The general geometry of Wytham Woods is likely to have introduced a

repeatable set of large-scale communities (Figure 3). Thus, the shape of the forest is

plays a major role in how the population is broadly structured (Figure 4). By

studying birds that form mixed-species communities, our study highlights that

hierarchical community structure can be the by-product of external processes.

Further, the majority of individuals in our study were great tits and blue tits. These

birds moved in similar ways, and we could not decompose our networks into groups

of one versus the other. Thus, while social mechanisms, such as social preference

(Farine *et al.* 2015b) and phenotypic drivers (Croft *et al.* 2003; Croft *et al.* 2009;

Aplin *et al.* 2013), can play a large role in determining who individuals affiliate with,

the woodland geometry and the resulting behaviour of all individuals combined can

generate large-scale static population structures.

The presence of multi-level community structure can have implications for

evolutionary dynamics of populations. First, restricted movement can reduce gene

flow and lead to divergence in the evolutionary trajectories of sub-parts of each

population. Garant *et al.* (2005) demonstrated that differential dispersal reinforces

local variation in selection for nestling body mass. In their study, they found that

trends in phenotypic variance for body mass in nestlings were very different in the

eastern sectors of Wytham Woods and the northern sectors. These two areas

represent the two largest population-level communities we found in our study.

Second, individuals in the same community will have more similar social

environments than individuals occurring in different communities. Thus, any social

effects arising via the social environment, such as indirect genetic effects (Moore,

Brodie & Wolf 1997), could accelerate patterns of divergence within single

populations. Finally, the social environment itself can act as an agent of selection

(Wolf & Moore 2010; Farine, Montiglio & Spiegel 2015), and therefore processes

that shape social structure are likely to impact the overall strength and direction of

selection experienced by populations (Montiglio, McGlothlin & Farine 2018).

One potential limitation of our study is that we employed bird feeders to detect the

presence of birds. This approach is what enabled us to collect information on so

many individuals simultaneously, but in doing so, we could have also influenced the

behaviour of birds. There are three reasons why we do not think that the presence

of bird feeders impacted our results. First, the feeders were all identical and open at

the same time, meaning that there would be little reason for a bird to choose to

relocate to another feeder (and as noted, most birds did not move, and birds moved

on average only once every two study days). Second, feeders were evenly spaced out

on a grid, meaning that, in the absence of habitat or behavioural heterogeneity,

movements between any pair of adjacent feeders should be equally likely. Finally,

feeders were open only two days per week, and remained shut for 5 consecutive

days. This means that birds would have had to maintain their natural foraging

behaviour rather than adapting to a new regime (noting that tits can lose up to 10%

of their body weight in a single winters' night, Owen 1954).

Together the findings from our study highlight several ways in which stable social

structure can be maintained in populations. The combination of strong clustering

together with some random movements in networks can facilitate the spread of

disease or information through the network (Eames 2008). This prediction is

supported by the rapid spread and establishment of novel traditions (Aplin *et al.*

2015a) in this population. At the same time, consistent population social structure

can lead to phenotypic and genotypic divergence (Garant *et al.* 2005), with potential

implications for how animals can adapt to changing environmental conditions.

Integrating information about animal social structure with data on both short-term

and long-term selective events could yield novel insights into the evolution of social

behaviour. As our study highlights, determining the capacity for populations to

respond to selective pressures will require an understanding of a range of different

drivers that could shape their social structure.

ACKNOWLEDGEMENTS

We thank Keith Kirby for discussion on the habitat data he collected, and for making

these data available to us. We also thank the large number of contributors to the

data collection, in particular to Keith McMahon.

REFERENCES

Adelman, J.S., Moyers, S.C., Farine, D.R. & Hawley, D.M. (2015) Feeder use predicts
both acquisition and transmission of a contagious pathogen in a North
American songbird. *Proceedings of the Royal Society B,* **282**, 20151429.
Allen, J., Weinrich, M., Hoppitt, W. & Rendell, L. (2013) Network-Based Diffusion
Analysis Reveals Cultural Transmission of Lobtail Feeding in Humpback
Whales. *Science,* **340**, 485-488.
Aplin, L.M., Farine, D.R., Mann, R.P. & Sheldon, B.C. (2014) Individual-level
personality influences social foraging and collective behaviour in wild birds.
Proceedings of the Royal Society B, **281**, 20141016.
Aplin, L.M., Farine, D.R., Morand-Ferron, J., Cockburn, A., Thornton, A. & Sheldon, B.C.
(2015a) Experimentally induced innovations lead to persistent culture via
conformity in wild birds. *Nature,* **518**, 538-541.
Aplin, L.M., Farine, D.R., Morand-Ferron, J., Cole, E.F., Cockburn, A. & Sheldon, B.C.
(2013) Individual personalities predict social behaviour in wild networks of
great tits (Parus major). *Ecology Letters,* **16**, 1365–1372.
Aplin, L.M., Farine, D.R., Morand-Ferron, J. & Sheldon, B.C. (2012) Social networks
predict patch discovery in a wild population of songbirds. *Proceedings of the*
Royal Society B, **279**, 4199-4205.
Aplin, L.M., Firth, J.A., Farine, D.R., Voelkl, B., Crates, R.A., Culina, A., Garroway, C.J.,
Hinde, C.A., Kidd, L.R., Psorakis, I., Milligan, N.D., Radersma, R., Verhelst, B. &
Sheldon, B.C. (2015b) Consistent individual differences in the social
phenotypes of wild great tits (*Parus major*). *Animal Behaviour,* **108**, 117-127.
Archie, E.A., Moss, C.J. & Alberts, S.C. (2006) The ties that bind: genetic relatedness
predicts the fission and fusion of social groups in wild African elephants.
Proceedings of the Royal Society B, **273**, 513-522.
Bell, H.L. & Ford, H.A. (1986) A Comparison of the Social-Organization of 3 Syntopic
Species of Australian Thornbill, Acanthiza. *Behavioral Ecology and*
Sociobiology, **19**, 381-392.
Bonter, D.N. & Bridge, E.S. (2011) Applications of radio frequency identification
(RFID) in ornithological research: a review. *Journal of Field Ornithology,* **82**,
1-10.
Brockmann, D. & Helbing, D. (2013) The Hidden Geometry of Complex, Network-
Driven Contagion Phenomena. *Science,* **342**, 1337-1342.
Cantor, M., Shoemaker, L.G., Cabral, R.B., Flores, C.O., Varga, M. & Whitehead, H.
(2015) Multilevel animal societies can emerge from cultural transmission.
Nature Communications, **6**.
Cox, D.T.C., Inger, R., Hancock, S., Anderson, K. & Gaston, K.J. (2016) Movement of
feeder-using songbirds: the in uence of urban features. *Scientific Reports,* **6**,
37669.
Croft, D.P., Arrowsmith, B.J., Bielby, J., Skinner, K., White, E., Couzin, I.D., Magurran,
A.E., Ramnarine, I. & Krause, J. (2003) Mechanisms underlying shoal

678   composition in the Trinidadian guppy, Poecilia reticulata. *Oikos,* **100,** 429-
679   438.
680 Croft, D.P., Hamilton, P.B., Darden, S.K., Jacoby, D.M.P., James, R., Bettaney, E.M. &
681   Tyler, C.R. (2012) The role of relatedness in structuring the social network of
682   a wild guppy population. *Oecologia,* **170,** 955-963.
683 Croft, D.P., Krause, J., Darden, S.K., Ramnarine, I.W., Faria, J.J. & James, R. (2009)
684   Behavioural trait assortment in a social network: patterns and implications.
685   *Behavioral Ecology and Sociobiology,* **63,** 1495-1503.
686 Csardi, G. & Nepusz, T. (2006) The igraph software package for complex network
687   research. *InterJournal,* **Complex Systems,** 1695.
688 Dall, S.R.X., Giraldeau, L.A., Olsson, O., McNamara, J.M. & Stephens, D.W. (2005)
689   Information and its use by animals in evolutionary ecology. *Trends in Ecology*
690   *and Evolution,* **20,** 187-193.
691 de Silva, S. & Wittemyer, G. (2012) A Comparison of Social Organization in Asian
692   Elephants and African Savannah Elephants. *International Journal of*
693   *Primatology,* **33,** 1125-1141.
694 Dekker, D., Krackhardt, D. & Snijders, T. (2007) Sensitivity of MRQAP tests to
695   collinearity and autocorrelation conditions. *Psychometrika,* **72,** 563-581.
696 Del Valle, S.Y., Hyman, J.M., Hethcote, H.W. & Eubank, S.G. (2007) Mixing patterns
697   between age groups in social networks. *Social Networks,* **29,** 539-554.
698 Eames, K.T.D. (2008) Modelling disease spread through random and regular
699   contacts in clustered populations. *Theoretical Population Biology,* **73,** 104-
700   111.
701 Farine, D.R. (2013) Animal Social Network Inference and Permutations for
702   Ecologists in R using asnipe. *Methods in Ecology and Evolution,* **4,** 1187–1194.
703 Farine, D.R. (2014) Measuring phenotypic assortment in animal social networks:
704   weighted associations are more robust than binary edges. *Animal Behaviour,*
705   **89,** 141-153.
706 Farine, D.R. (2015) Proximity as a proxy for interactions: issues of scale in social
707   network analysis. *Animal Behaviour,* **104,** e1-e5.
708 Farine, D.R. (2017) A guide to null models for animal social network analysis.
709   *Methods in Ecology and Evolution,* **8,** 1309-1320.
710 Farine, D.R., Aplin, L.M., Garroway, C.J., Mann, R.P. & Sheldon, B.C. (2014) Collective
711   decision-making and social interaction rules in mixed-species flocks of
712   songbirds. *Animal Behaviour,* **95,** 173-182.
713 Farine, D.R., Aplin, L.M., Sheldon, B.C. & Hoppitt, W. (2015a) Interspecific social
714   networks promote information transmission in wild songbirds. *Proceedings*
715   *of the Royal Society B,* **282,** 20142804.
716 Farine, D.R., Firth, J.A., Aplin, L.M., Crates, R.A., Culina, A., Garroway, C.J., Hinde, C.A.,
717   Kidd, L.R., Milligan, N.D., Psorakis, I., Radersma, R., Verhelst, B., Voelkl, B. &
718   Sheldon, B.C. (2015b) The role of social and ecological processes in
719   structuring animal populations: a case study from automated tracking of wild
720   birds. *Royal Society Open Science,* **2,** 150057.
721 Farine, D.R., Garroway, C.J. & Sheldon, B.C. (2012) Social network analysis of mixed-
722   species flocks: exploring the structure and evolution of interspecific social
723   behaviour. *Animal Behaviour,* **84,** 1271-1277.

Farine, D.R. & Lang, S.D.J. (2013) The early bird gets the worm: foraging strategies of
wild songbirds lead to the early discovery of food sources. *Biology Letters,* **9,**
20130578.
Farine, D.R., Montiglio, P.O. & Spiegel, O. (2015) From individuals to groups and
back: the evolutionary implications of group phenotypic composition. *Trends*
in Ecology & Evolution, **30,** 609-621.
Farine, D.R. & Sheldon, B.C. (2015) Selection for territory acquisition is modulated
by social network structure in a wild songbird. *Journal of Evolutionary*
Biology, **28,** 547-556.
Farine, D.R. & Whitehead, H. (2015) Constructing, conducting, and interpreting
animal social network analysis. *Journal of Animal Ecology,* **84,** 1144-1163.
Finn, J.T., Brownscombe, J.W., Haak, C.R., Cooke, S.J., Cormier, R., Gagne, T. &
Danylchuk, A.J. (2014) Applying network methods to acoustic telemetry data:
Modeling the movements of tropical marine fishes. *Ecological Modelling,* **293,**
139-149.
Firth, J.A. & Sheldon, B.C. (2016) Social carry-over effects underpin trans-seasonally
linked structure in a wild bird population. *Ecology Letters,* **19,** 1324-1332.
Formica, V.A., McGlothlin, J.W., Wood, C.W., Augat, M.E., Butterfield, R.E., Barnard,
M.E. & Brodie, E.D. (2011) Phenotypic Assortment Mediates the Effect of
Social Selection in a Wild Beetle Population. *Evolution,* **65,** 2771-2781.
Formica, V.A., Wood, C.W., Larsen, W.B., Butterfield, R.E., Augat, M.E., Hougen, H.Y. &
Brodie, E.D. (2012) Fitness consequences of social network position in a wild
population of forked fungus beetles (*Bolitotherus cornutus*). *Journal of*
Evolutionary Biology, **25,** 130-137.
Fortunato, S. (2010) Community detection in graphs. *Physics Reports-Review Section*
of Physics Letters, **486,** 75-174.
Garant, D., Kruuk, L.E.B., Wilkin, T.A., McCleery, R.H. & Sheldon, B.C. (2005)
Evolution driven by differential dispersal within a wild bird population.
Nature, **433,** 60-65.
Godfrey, S.S., Anasri, T.H., Gardner, M.G., Farine, D.R. & Bull, C.M. (2014) A contact
based social network of lizards is defined by low genetic relatedness among
strongly-connected individuals. *Animal Behaviour,* **97,** 35-43.
Goodale, E., Beauchamp, G., Magrath, R.D., Nieh, J.C. & Ruxton, G.D. (2010)
Interspecific information transfer influences animal community structure.
Trends in Ecology and Evolution, **25,** 354-361.
Gräler, B., Pebesma, E. & Heuvelink, G. (2016) Spatio-Temporal Interpolation using
gstat. *The R Journal,* **8,** 204-218.
Grueter, C.C., Matsuda, I., Zhang, P. & Zinner, D. (2012) Multilevel Societies in
Primates and Other Mammals: Introduction to the Special Issue. *International*
Journal of Primatology, **33,** 993-1001.
He, P., Maldonado-Chaparro, A. & Farine, D.R. (2019) The role of habitat
configuration in shaping social structure: a gap in studies of animal social
complexity. *Behavioral Ecology and Sociobiology*, 9.
Hilleman, F., Cole, E.F., Keen, S.C., Sheldon, B.C. & Farine, D.R. (in press) Diurnal
variation in the production of vocal information about food supports a model

of social adjustment in wild songbirds. *Proceedings of the Royal Society B,*
10.1098/rspb-2018-2740.
Hinde, R.A. (1976) Interactions, Relationships and Social-Structure. *Man,* **11,** 1-17.
Hoppitt, W. & Farine, D.R. (2018) Association indices for quantifying social
relationships: how to deal with missing observations of individuals or
groups. *Animal Behaviour,* **136,** 227-238.
Jacoby, D.M.P. & Freeman, R. (2016) Emerging Network-Based Tools in Movement
Ecology. *Trends in Ecology & Evolution,* **31,** 301-314.
Jesmer, B.R., Merkle, J.A., Goheen, J.R., Aikens, E.O., Beck, J.L., Courtemanch, A.B.,
Hurley, M.A., MeWhicter, D.E., Miyasaki, H.M., Monteith, K.L. & Kauffman, M.J.
(2018) Is ungulate migration culturally transmitted? Evidence of social
learning from translocated animals. *Science,* **361,** 1023-1025.
Kays, R., Crofoot, M.C., Jetz, W. & Wikelski, M. (2015) Terrestrial animal tracking as
an eye on life and planet. *Science,* **348,** aaa2478.
Kirby, K.J., Bazely, D.R., Goldberg, E.A., Hall, J.E., Isted, R., Perry, S.C. & Thomas, R.C.
(2014) Changes in the tree and shrub layer of Wytham Woods (Southern
England) 1974-2012: local and national trends compared. *Forestry,* **87,** 663-
673.
Krause, J., James, R., Franks, D.W. & Croft, D.P. (2015) *Animal Social Networks.* Oxford
University Press, Oxford, UK.
Krause, J., Krause, S., Arlinghaus, R., Psorakis, I., Roberts, S. & Rutz, C. (2013) Reality
mining of animal social systems. *Trends in Ecology & Evolution,* **28,** 541-551.
Kurvers, R.H.J.M., Krause, J., Croft, D.P., Wilson, A.D.M. & Wolf, M. (2014) The
evolutionary and ecological consequences of animal social networks:
emerging issues. *Trends in Ecology & Evolution,* **29,** 326-335.
Leu, S.T., Farine, D.R., Wey, T.W., Sih, A. & Bull, M. (2016) Environment modulates
population social structure: experimental evidence from replicated social
networks of wild lizards. *Animal Behaviour,* **111,** 23-31.
Levin, I.I., Zonana, D.M., Burt, J.M. & Safran, R.J. (2015) Performance of Encounternet
Tags: Field Tests of Miniaturized Proximity Loggers for Use on Small Birds.
Plos One, **10**.
Matechou, E., Cheng, S.C., Kidd, L.R. & Garroway, C.J. (2015) Reproductive
consequences of the timing of seasonal movements in a non-migratory wild
bird population. *Ecology,* **96,** 1641-1649.
Montiglio, P.O., McGlothlin, J.W. & Farine, D.R. (2018) Social structure modulates the
evolutionary consequences of social plasticity: taking a social network
perspective of interacting phenotypes. *Ecology & Evolution,* **8,** 1451-1464.
Moore, A.J., Brodie, E.D. & Wolf, J.B. (1997) Interacting phenotypes and the
evolutionary process .1. Direct and indirect genetic effects of social
interactions. *Evolution,* **51,** 1352-1362.
Mueller, T., O'Hara, R.B., Converse, S.J., Urbanek, R.P. & Fagan, W.F. (2013) Social
Learning of Migratory Performance. *Science,* **341,** 999-1002.
O'Brien, D., Manseau, M., Fall, A. & Fortin, M.J. (2006) Testing the importance of
spatial configuration of winter habitat for woodland caribou: An application
of graph theory. *Biological Conservation,* **130,** 70-83.
Owen, D.F. (1954) The winter weights of titmice. *Ibis,* **96,** 299-309.

Pruitt, J.N. & Goodnight, C.J. (2014) Site-specific group selection drives locally
adapted group compositions. *Nature,* **514,** 359–362.
Pruitt, J.N. & Keiser, C.N. (2014) The personality types of key catalytic individuals
shape colonies' collective behaviour and success. *Animal Behaviour,* **93,** 87-
95.
Pruitt, J.N. & Riechert, S.E. (2011) How within-group behavioural variation and task
efficiency enhance fitness in a social group. *Proceedings of the Royal Society B,*
278, 1209-1215.
Psorakis, I., Roberts, S.J., Rezek, I. & Sheldon, B.C. (2012) Inferring social network
structure in ecological systems from spatio-temporal data streams. *Journal of*
the Royal Society Interface, **9,** 3055-3066.
Psorakis, I., Voelkl, B., Garroway, C.J., Radersma, R., Aplin, L.M., Crates, R.A., Culina,
A., Farine, D.R., Firth, J.A., Hinde, C.A., Kidd, L.R., Milligan, N.D., Roberts, S.J.,
Verhelst, B. & Sheldon, B.C. (2015) Inferring social structure from temporal
data. *Behavioral Ecology and Sociobiology,* **69,** 857-866.
Quinn, J.L., Cole, E.F., Bates, J., Payne, R.W. & Cresswell, W. (2012) Personality
predicts individual responsiveness to the risks of starvation and predation.
Proceedings of the Royal Society B, **279,** 1919-1926.
Quinn, J.L., Cole, E.F., Patrick, S.C. & Sheldon, B.C. (2011) Scale and state dependence
of the relationship between personality and dispersal in a great tit
population. *Journal of Animal Ecology,* **80,** 918-928.
Radersma, R., Garroway, C.J., Santure, A.W., de Cauwer, I., Farine, D.R., Slate, J. &
Sheldon, B.C. (2017) Social and spatial effects on genetic variation between
foraging flocks in a wild bird population. *Molecular Ecology,* **26,** 5807–5819.
Rutz, C., Burns, Z.T., James, R., Ismar, S.M.H., Burt, J., Otis, B., Bowen, J. & St Clair, J.J.H.
(2012) Automated mapping of social networks in wild birds. *Current Biology,*
22, R669-R671.
Ryder, T.B., Horton, B.M., van den Tillaart, M., Morales, J.D. & Moore, I.T. (2012)
Proximity data-loggers increase the quantity and quality of social network
data. *Biology Letters,* **8,** 917-920.
Schreier, A.L. & Swedell, L. (2009) The Fourth Level of Social Structure in a Multi-
Level Society: Ecological and Social Functions of Clans in Hamadryas
Baboons. *American Journal of Primatology,* **71,** 948-955.
Seppanen, J.T., Forsman, J.T., Monkkonen, M. & Thomson, R.L. (2007) Social
information use is a process across time, space, and ecology, reaching
heterospecifics. *Ecology,* **88,** 1622-1633.
Shizuka, D., Chaine, A.S., Anderson, J., Johnson, O., Laursen, I.M. & Lyon, B.E. (2014)
Across-year social stability shapes network structure in wintering migrant
sparrows. *Ecology Letters,* **17,** 998-1007.
Shizuka, D. & Farine, D.R. (2016) Measuring the robustness of network community
structure using assortativity. *Animal Behaviour,* **122,** 237-246.
Slagsvold, T. & Wiebe, K.L. (2007) Learning the ecological niche. *Proceedings of the*
Royal Society B, **274,** 19-23.
Strandburg-Peshkin, A., Farine, D.R., Couzin, I.D. & Crofoot, M.C. (2015) Shared
decision-making drives collective movement in wild baboons. *Science,* **348,**
1358-1361.

Strandburg-Peshkin, A., Farine, D.R., Crofoot, M.C. & Couzin, I.D. (2017) Habitat and
social factors shape individual decisions and emergent group structure
during baboon collective movement. *eLife,* **6,** e19505.
Svensson, L. (1992) *Identification Guide to European Passerines,* 4th edition edn.
BTO, Stockholm.
Tomkiewicz, S.M., Fuller, M.R., Kie, J.G. & Bates, K.K. (2010) Global positioning
system and associated technologies in animal behaviour and ecological
research. *Philosophical Transactions of the Royal Society B-Biological Sciences,*
365, 2163-2176.
Valone, T.J. (2007) From eavesdropping on performance to copying the behavior of
others: a review of public information use. *Behavioral Ecology and*
Sociobiology, **62,** 1-14.
VanderWaal, K.L., Atwill, E.R., Isbell, L.A. & McCowan, B. (2013) Linking social and
pathogen transmission networks using microbial genetics in giraffe (Giraffa
camelopardalis). *Journal of Animal Ecology,* **83,** 406-414.
Wey, T.W., Burger, J.R., Ebensperger, L.A. & Hayes, L.D. (2013) Reproductive
correlates of social network variation in plurally breeding degus (Octodon
degus). *Animal Behaviour,* **85,** 1407-1414.
Whitehead, H. (2008) *Analyzing animal societies.* University of Chicago Press,
Chicago, USA.
Whitehead, H., Antunes, R., Gero, S., Wong, S.N.P., Engelhaupt, D. & Rendell, L. (2012)
Multilevel Societies of Female Sperm Whales (Physeter macrocephalus) in
the Atlantic and Pacific: Why Are They So Different? *International Journal of*
Primatology, **33,** 1142-1164.
Wiens, J.A. (1976) Population Responses to Patchy Environments. *Annual Review of*
Ecology and Systematics, **7,** 81-120.
Wittemyer, G., Douglas-Hamilton, I. & Getz, W.M. (2005) The socioecology of
elephants: analysis of the processes creating multitiered social structures.
Animal Behaviour, **69,** 1357-1371.
Wolf, J.B. & Moore, A.J. (2010) Interacting Phenotypes and Indirect Genetic Effects.
Evolutionary Behavioral Ecology (eds D.F. Westneat & C.W. Fox), pp. 225-245.
Oxford University Press, Oxford.
Wolf, J.B.W., Mawdsley, D., Trillmich, F. & James, R. (2007) Social structure in a
colonial mammal: unravelling hidden structural layers and their foundations
by network analysis. *Animal Behaviour,* **74,** 1293-1302.
896
897