**Title:** Super-resolution microscopy of the ß-carboxysome reveals a homogenous matrix

**Authors:** Matthew J. Niederhuber *,[†], Talley J. Lambert[‡], Clarence Yapp[§], Pamela A.

Silver*,[†], Jessica K. Polka*,[†]

**Affiliations:**

* Department of Systems Biology, Harvard Medical School, 02115, Boston,

Massachusetts, United States.

† Wyss Institute for Biologically Inspired Engineering, Harvard University, 02115,

Boston, Massachusetts, United States.

‡. Department of Cell Biology, Harvard Medical School, 02115, Boston, Massachusetts,

United States.

§. Image and Data Analysis Core, Harvard Medical School, 02115, Boston,

Massachusetts, United States.

**Abstract:**

Carbon fixation in cyanobacteria makes a major contribution to the global carbon

cycle. The cyanobacterial carboxysome is a proteinaceous microcompartment that

protects and concentrates the carbon-fixing enzyme RuBisCO in a paracrystalline lattice,

making it possible for these organisms to fix $CO_2$ from the atmosphere. The protein

responsible for the organization of this lattice in beta-type carboxysomes of the

freshwater cyanobacterium *Synechococcus elongatus*, CcmM, occurs in two isoforms

thought to localize differentially within the carboxysome matrix. Here, we use widefield

timelapse and 3D-structured illumination microscopy (3D-SIM) to study the recruitment

and localization of these two isoforms. We demonstrate that this super-resolution

technique is capable of successfully resolving the outer protein shell of the carboxysome

from its internal cargo. We develop an automated analysis pipeline to analyze and

quantify 3D-SIM images and generate a population level description of carboxysome

shell protein, RuBisCO, and CcmM isoform localization. We find that both CcmM

isoforms colocalize in space and time, prompting a revised model of the internal

arrangement of the beta carboxysome.

**Introduction:**

RuBisCO (ribulose-1,5-bisphosphate carboxylase/oxygenase), the primary

enzyme of carbon fixation, is ubiquitous in autotrophic organisms. However, it is

notoriously inefficient and readily participates in energetically wasteful side-reactions

with $O_2$ (Andersson *et al*., 2008). Many organisms that rely on RuBisCO to acquire

inorganic carbon tune the enzyme's local microenvironment to achieve sufficient levels

of carbon fixation. While plants can do this within membrane barriers, many single-celled

organisms have evolved a strategy of proteinaceous compartmentalization.

The carboxysome is a bacterial microcompartment (BMC) found in cyanobacteria

and prokaryotic chemoautotrophs. Composed of thousands of individual subunits, the

carboxysome encapsulates RuBisCO and carbonic anhydrase (CA) within a selectively

permeable icosahedral shell, making survival possible at atmospheric levels of $CO_2$

(Figure 1A) (Rae *et al*., 2013; Kerfeld *et al*., 2016).

While evolutionarily distinct carboxysomes share many similar structural features,

the RuBisCO in the beta-type carboxysome (ß-carboxysome) of the freshwater

cyanobacterium *Synechococcus elongatus* PCC7942 is organized into a paracrystalline

lattice (Kaneko *et al*., 2006). The structural protein CcmM has been implicated in

connecting individual RuBisCO enzymes into an ordered matrix, resulting in the

observed dense crystalline packing (Long *et al*., 2007; Cot *et al*., 2008, Long *et al.,* 2010;

Cameron *et al*., 2013). The full-length 58kDa CcmM protein of PCC7942 is characterized by an N-terminal carbonic anhydrase (CA) binding domain and three repeating domains toward the C-terminal end that have sequence homology to the small subunit of RuBisCO (Figure 1B) (Price *et al*., 1993). Due to an internal ribosomal entry site (IRES), CcmM is translated as two isoforms, a full-length protein (M58) and a 35kDa truncated version (M35) that consists of only the three small subunit like (SSU-like) domains (Long *et al*., 2007; Long *et al*., 2010).

The current model of ß-carboxysome internal structure suggests that the small subunit like domains (SSU-like) of CcmM displace some of the small subunits of the $L_8S_8$ RuBisCO holoenzyme and tether multiple enzymes together into a dense network (Figure 1C) (Long *et al*., 2007; Long *et al*., 2010; Rae *et al*., 2013; Kerfeld *et al.,* 2016). Co-purification and yeast 2-hybrid experiments have shown that RuBisCO readily complexes with both isoforms of CcmM in PCC7942 (Long *et al*., 2007; Cot *et al*., 2008). In addition, fluorescence tagging experiments have demonstrated that M35 alone is capable of nucleating assembly of tagged RuBisCO if at least two of the SSU-like domains are present. However, M35 is not sufficient to cause encapsulation by recruiting shell components in the absence of M58 (Cameron *et al*., 2013). Densitometry analyses of western blot experiments have found imbalanced ratios of RbcS to RbcL (~5:8) in PCC7942 cell lysates, suggesting that the missing RbcS slots may be occupied by CcmM SSU domains (Long *et al*., 2011). These findings have supported a model of carboxysome architecture and step-wise assembly in which M35 nucleates a core of RuBisCO. In this model, M58 is limited to a layer beneath the shell where it can interact with CcmN, CA, and shell components. While this model predicts that the CcmM isoforms are differentially localized and that the timing of their integration may differ, these predictions have yet to be directly tested.

In order to test this model, we fluorescently tagged CcmM to visualize its

assembly and the localization of CcmM isoforms *in vivo*. Previous work has shown that

completely replacing CcmM with tagged fusions does not disrupt carboxysome function

(Long *et al*., 2007; Cameron *et al*., 2015). Loss of either isoform of CcmM creates

mutants that fail to produce carboxysomes and have a high $CO_2$ requirement (HCR) for

growth (Woodger *et al.,* 2005; Long *et al*., 2010). However, complementation of such

mutants with tagged copies of CcmM has been shown to rescue this phenotype (Long *et*

*al*., 2007; Cameron *et al*., 2015).

Here we show the successful complementation of a Δ*ccmM* mutant with a dually

fluorescently tagged integration of *ccmM*. This dual tagging strategy does not

significantly impair carboxysome formation, morphology, or function. In our timelapse

fluorescence microscopy data we find no evidence of differential CcmM isoform

assembly during carboxysome biogenesis. Using super-resolution 3D Structured

Illumination Microscopy (3D-SIM), we examined the relative localization of carboxysome

shell, CcmM, and RuBisCO proteins. Our 3D-SIM data does not support a model of

carboxysome internal organization in which M58 is limited to a layer just beneath the

external protein shell. These data support a revised model of carboxysome internal

structure in which M35 and M58 isoforms are both integrated deep within the core of the

carboxysome and are both involved with the early steps of carboxysome assembly.

**Results:**

**Fluorescent fusions to CcmM complement a deletion of the wildtype**

We integrated an N- and C-terminal fluorescently-tagged copy of *ccmM* (*mTagBFP2-*

*ccmM-mNeonGreen*) driven by the IPTG inducible promoter P*trc* into the genome of an

*S. elongatus* PCC7942 Δ*ccmM*::*HygR* mutant strain (Figure 2A-B) and verified

expression with western blot analysis (Figure 2C). This strategy allowed us to visually

separate full length CcmM from truncated M35 by blue fluorescence. We found that P*trc*

was sufficiently leaky to produce protein without induction. Consequently we never

induced the Δ*ccmM* + *mTagBFP2-CcmM-mNeonGreen* strain, including for western blot

analysis, unless otherwise noted. We blotted lysates of WT, Δ*ccmM*, and Δ*ccmM* +

*mTagBFP2-CcmM-mNeonGreen* strains for the presence of CcmM (Figure 2C). In WT

7942 cells, we detected two bands at ~60 kDa and ~40 kDa, corresponding to the two

native CcmM isoforms M58 and M35. In the Δ*ccmM* strain, these bands are absent. In

the Δ*ccmM* + *mTagBFP2-CcmM-mNeonGreen* gene complement, we detected two

bands at ~140 kDa and ~70 kDa, corresponding to molecular weight shifts of 60kDa

(both fluorescent tags) and 30kDa (mNeonGreen alone), respectively. The observed

weight shift of the tagged CcmM demonstrates that the CcmM IRES is not impaired by

the presence of N- and C-terminal tags, and that both long and short CcmM isoforms are

translated. We also made a reciprocally tagged version of the CcmM complement

(mNeonGreen-CcmM-mTagBFP2), but did not use it for further analysis because

western blotting showed the presence of protein degradation (Supplemental Figure S1).

Our tagged *ccmM* complement grew at a rate comparable to WT at atmospheric

levels of $CO_2$ (Figure 2D). We grew WT 7942, the Δ*ccmM* mutant, and the Δ*ccmM* +

*mTagBFP2-CcmM-mNeonGreen* complement in ambient air and 2% $CO_2$ conditions,

measured optical density at 750nm ($OD_{750}$), and calculated doubling times between the

18- and 90-hour time points. Both WT and Δ*ccmM* cells cultured with 2% $CO_2$ grew well

with similar doubling times of 17.6 $\pm$ 0.4 hours ($n$ = 5, Error = SD) and 15.8 $\pm$ 2.1 hours

($n$ = 6, Error = SD) respectively. As expected, Δ*ccmM* mutant growth was dependent on

supplemented $CO_2$ (Long *et al*., 2010); these cells showed no growth in ambient air

conditions. In contrast, the Δ*ccmM* + *mTagBFP2-CcmM-mNeonGreen* complement grew

well in air with a doubling time of 23.3 $\pm$ 2.4 hours ($n$ = 6, Error = SD). This growth rate is

similar to that of WT cells in air, which had a doubling time of 20.6 $\pm$ 1.8 hours ($n$ = 6,

Error = SD), indicating that our engineered strain is capable of fixing carbon, and that

carboxysome function is not severely impaired by the dual-fusion *ccmM* complement.

Because both the M58 and M35 isoforms of CcmM are required for growth without $CO_2$

supplementation (Long *et al.*, 2010), our data indicate that both isoforms produced by

our dual fusion are functional.

Carboxysomes formed in the $\Delta ccmM$ + *mTagBFP2-CcmM-mNeonGreen*

complement were structurally similar to those of WT cells. Using transmission electron

microscopy, we imaged ultrathin sections of fixed cells from WT 7942, $\Delta ccmM$, and

$\Delta ccmM$ + *mTagBFP2-CcmM-mNeonGreen* strains (Figure 3). Carboxysomes in WT

PCC7942 cells had an average maximum width of 179 $\pm$ 33 nm ($n$ = 26, Error = SD).

Carboxysomes in $\Delta ccmM$ + *mTagBFP2-CcmM-mNeonGreen* cells generally had a

normal appearance and shape with visibly angled facets and often clear hexagonal

geometry, but were noticeably larger, with an average maximum diameter of 319 $\pm$ 92

nm ($n$ = 41, Error = SD). This is not unexpected; it has been previously reported that His-

tagged *ccmM* complements increase carboxysome size (Long *et al.,* 2007).

Occasionally, dense masses at the cell poles were observed in $\Delta ccmM$ + *mTagBFP2-*

*CcmM-mNeonGreen* micrographs, most likely indicating protein aggregation. Because

we are using this strain without induction and no effective repressors exist, we cannot

know if reducing expression would prevent aggregate formation. However, IPTG-

induced cells observed with 3D-SIM display a dramatic increase in polar protein

aggregation, often filling a large portion of the cell, confirming that overexpression of this

construct leads to aggregation (Supplemental Figure S2).

**Both CcmM isoforms share similar recruitment kinetics and localization**

155
Both CcmM isoforms are recruited to growing carboxysomes with similar timing

and kinetics. Using widefield microscopy, we imaged the CcmM complement

(mTagBFP2-CcmM-mNeonGreen) at 10 minute intervals for 40 hours (Figure 4). Particle

tracking revealed that both the CcmM isoforms assembled slowly, with an average

assembly time of approximately 10.50 $\pm$ 5.39 hours ($n$ = 8, Error = SD) for M58-

mNeonGreen and M35-mNeonGreen, and 12.06 $\pm$ 7.27 hours ($n$ = 8, Error = SD) for

mTagBFP2-M58 (Figure 4C). This is longer than the ~6-7 hour maturation time

previously reported for RuBisCO (Chen *et al.*, 2013), perhaps as a result of temperature

or light levels during imaging. Assembly times were estimated based on the time it took

a particle to go from the intensity required for automatic particle detection (methods) to

the first value $\geq$ 0.8 on the normalized intensity plot.

Next, a four-parameter logistic equation was used to approximately model the

particle tracks. These models highlight similar assembly kinetics for both CcmM isoforms

within the same carboxysome.  Based on these fit curves, half-max values of green and

blue particle assemblies were found to have an absolute average difference of 1.18 $\pm$

0.68 hours ($n$ = 8, Error = SD). If one CcmM isoform precedes the other during

carboxysome biogenesis, the half-max values of each isoform would be expected to be

significantly different. However, a comparison of half-max values from eight measured

assemblies found no significant difference (Wilcoxon matched-pairs signed rank test,

number of pairs = 8, P = 0.945). In order to test if CcmM isoforms differ in rate of

assembly, channel specific assembly times (average reported above) were analyzed

and found not to be significantly different among paired fluorescent channels (Wilcoxon

matched-pairs signed rank test, number of pairs = 8, P = 0.375). This indicates that

CcmM isoforms M35 and M58 assembled at relatively similar rates.

**M58 is not confined to a sub-shell layer in the carboxysome**

Super-resolution microscopy of fluorescently tagged M58 and M35 show indistinguishable distributions in the carboxysome. We used 3D-Structured Illumination Microscopy (3D-SIM) to acquire super-resolution images of the Δ*ccmM* + *mTagBFP2-CcmM-mNeonGreen* strain, as well as fluorescently tagged RuBisCO (RbcL-sfGFP) and the shell protein CcmK4 (CcmK4-mTagBFP2 and CcmK4-sfGFP) in PCC7942. 3D-SIM images of strains with both GFP- and BFP-tagged CcmK4 resolved the external carboxysome shell as a ring structure in the xy-plane (Figure 5A-F), demonstrating that 3D-SIM provides sufficient resolution to identify peripherally localized proteins in the carboxysome. To confirm these rings were not an artifact of SIM reconstruction, we imaged a strain of PCC7942 integrated with a GFP labeled copy of the large subunit of RuBisCO (RbcL-sfGFP). This strain has been previously used successfully to study RuBisCO assembly and spacing (Savage *et al*., 2010; Chen *et al*., 2013). 3D-SIM of RbcL-sfGFP showed no ring-like structures, producing only solid regularly-spaced particles (Figure 5G-I). To test if M58 was in fact limited to a sub-shell layer, we imaged the Δ*ccmM* + *mTagBFP2-CcmM-mNeonGreen* gene complement with 3D-SIM. Both CcmM isoforms produced solid objects similar to those observed with the RbcL-sfGFP strain (Figure 5J-N). These qualitative results suggested that the tagged M58 isoform of CcmM was not excluded from integration into the carboxysome core as would be expected if the protein was only present as a thin layer just beneath the exterior shell. The reconstructed intensity histogram (RIH) score from the SIMcheck plugin (Ball *et al*., 2015) was used to verify reconstruction quality. Reconstruction scores for all conditions were of similar quality (Supplemental Table S1).

We developed an automated image analysis pipeline and a metric to quantify the degree to which fluorescently-labeled proteins were localized at the perimeter of the carboxysome versus uniformly distributed throughout the matrix. The max-to-center

intensity (MTC) ratio is calculated for each particle as the peak intensity of the average

radial profile divided by the intensity at the particle centroid.  A perfectly uniform protein

distribution would result in the peak particle intensity overlapping the particle centroid,

and yield an MTC ratio of 1.00. Higher MTC values result from particles with decreased

intensity at their centers, indicating a more ring-link appearance.

ANOVA testing confirmed that mean MTC values were significantly

heterogeneous between all the labeled carboxysome proteins (Welch's ANOVA, $F_{4,}$

$_{7378}$=160.48, $P$<0.001). A Games-Howell test with $q_{critical}$(k=5, df=∞, α=0.001)=5.484 was

then used to make further pairwise comparisons of labeled carboxysome protein MTC

values (Supplementary Table S2). This automated analysis of RbcL-sfGFP and CcmK4-

sfGFP images confirmed our initial observations that carboxysomes appear

fundamentally different between these strains (Games-Howell test, MD=0.02, $q$=22.26);

similar results were obtained from the comparison of RbcL-sfGFP and CcmK4-

mTagBFP2 (Games-Howell test, MD=0.03, $q$=26.38) (Figure 6A). RbcL-sfGFP particles

had an average diameter of 244 $\pm$ 45 nm and an average MTC value of 1.002 $\pm$ 0.005 ($n$

= 3505), with a maximum intensity ratio of 1.04. Based on this, we calculated an upper

ratio threshold of 1.05, below which objects are likely to be uniform, by adding two

standard deviations to the RbcL-sfGFP max ratio of 1.04. This threshold is denoted on

MTC ratio versus diameter plots as a horizontal dotted line (Figure 6).

In contrast to RbcL, we found that shell proteins displayed high MTC ratios.

CcmK4-sfGFP particles had an average diameter of 253 $\pm$ 34 nm ($n$ = 3459), and in

comparison to RbcL-sfGFP particles, a significantly greater average MTC ratio of 1.023

$\pm$ 0.077 (Games-Howell test, MD=0.02, $q$=22.26) (Figure 6H). Approximately 10% of

CcmK4-sfGFP particles have MTC ratios $\geq$ 1.05, denoting a ring-like appearance (Figure

6B). Analysis of CcmK4-mTagBFP2 particles also revealed a significantly different mean

MTC ratio compared to labeled RbcL-sfGFP (Games-Howell test, MD=0.03, $q$=26.38)

(Figure 6H), with the CcmK4-mTagBFP2 particles having an average size of 244 $\pm$ 32

nm and average MTC ratio of 1.036 $\pm$ 0.093 (Figure 6C). Of note, the percent of ring-like

particles identified in CcmK4-mTagBFP2 cells (18.16% $\geq$ $MTC_{1.05}$) was significantly

greater than the percentage found in CcmK4-sfGFP cells (9.90% $\geq$ $MTC_{1.05}$) (Games-

Howell test, MD=0.01, $q$=8.51) (Figure 6H). This increase in the fraction of rings

identified with BFP labeled CcmK4 over those identified with GFP may be the result of

the improvement in resolution due to the shorter wavelength of blue versus green

fluorophores. Based on the smallest resolvable shell in the CcmK4-mTagBFP2 strain,

which had an apparent diameter of 197nm, and accounting for an expected shell depth

of 5-6nm, we do not expect to resolve by SIM even true ring-like particles smaller than

approximately 220nm in diameter (Kaneko *et al.*, 2006). This lower resolvable limit is

denoted on plots of tagged M58 and of M35 particles by a grey region in Figure 6D-E.

245           Analysis of the $\Delta ccmM$ + *mTagBFP2-CcmM-mNeonGreen* gene complement

identified a small number of tagged M58 particles as ring-like (Figure 6D). Our analysis

found 0.58% (19 of 3303) of the mTagBFP2-M58 particles had MTC ratios of $\geq$ 1.05,

with an average size of 233 $\pm$ 46 nm and average MTC ratio of 1.003 $\pm$ 0.010 (Figure

6F). We found a significant difference in the MTC ratios of mTagBFP2-M58 particles vs

both shell-labeled CcmK4-mTagBFP2 (Games-Howell test, MD=0.03, $q$=25.27) and

CcmK4-sfGFP (Games-Howell test, MD=0.02, $q$=20.69) (Figure 6H). This supports

qualitative observations that particles of the labeled M58 strain did not appear as ring-

like as those in the strains with labeled carboxysome shells. mTagBFP2-M58 particles

also had significantly different MTC ratios compared to RbcL-sfGFP particles (Games-

Howell test, MD=0.001, $q$=10.05), but this difference was produced by only 19 ring-like

particles in the mTagBFP2-M58 strain (Figure 6H). Taken together, these test results

indicate that though mTagBFP2-M58 labeled carboxysomes have a small but statistically

significant number of MTC ratios $\geq$ 1.05, the mean ratio is not similar to that produced by

labeled carboxysome shell proteins, as would have been expected if the M58 isoform

followed a strict sub-shell localization.

The Δ*ccmM* + *mTagBFP2-CcmM-mNeonGreen* complement, which expresses

both full-length M58-mNeonGreen and the short isoform M35-mNeonGreen, produced 2

particles with MTC ratios ≥ 1.05 (Figure 6E). Only 0.06% of particles were found above

that threshold, with an average size of 244 ± 47 nm (*n* = 3437) and average MTC ratio of

1.002 ± 0.005 (Figure 6F). The mean MTC ratio of these particles was significantly

different than those of both labeled CcmK4-sfGFP (Games-Howell test, MD=0.02,

*q*=22.20) and CcmK4-mTagBFP2 (Games-Howell test, MD=0.03, *q*=26.34) (Figure 6H).

Pair-wise comparison of M58-mNeonGreen and M35-mNeonGreen particles to RbcL-

sfGFP found no significant difference in mean MTC ratios (Games-Howell test,

MD=0.00005, *q*=0.57) (Figure 6G), indicating that the population of green fluorescent

particles produced in the Δ*ccmM* + mTagBFP2-CcmM-mNeonGreen strain were solid,

similar to those observed in the strain with labeled RuBisCO.

**Discussion:**

**The carboxysome can support the addition of large foreign proteins via direct**

**fusion to CcmM**

The function of CcmM and other components of the carboxysome are robust to

the addition of large protein fusions, further supporting these fusions as a means to

localize foreign cargo to the carboxysome for engineering applications. Previous work

has shown that tagged CcmM gene complements can rescue the HCR phenotype of

CcmM mutants without significant impairment of carboxysome function compared to WT

cells (Long *et al.*, 2010; Cameron *et al.*, 2015). We find that such complementation is

also possible when fluorescent tags are fused to both ends of CcmM, significantly

increasing the size of both the M58 and M35 isoforms. These results suggest that

heterologous proteins could be simultaneously fused to multiple loci in the matrix,

opening the door for localizing multipart enzymatic pathways inside the carboxysome.

**M35 and M58 assemble simultaneously during carboxysome biogenesis**

M35 has been hypothesized to coordinate the organized paracrystalline cargo

assembly of RuBisCO, with M58 limited to a peripheral sub-shell layer (Long *et al.,* 2007;

Cot *et al.,* 2008; Long *et al*., 2010; Long *et al*., 2011; Rae *et al*., 2013). Based on this

architecture, we expected M35 to assemble prior to M58 in the same manner as shell

recruitment follows RuBisCO (Chen *et al*., 2013). However, we were unable to resolve

any significant temporal difference between the CcmM isoforms. This concurrent timing

suggests that M58 interacts with RuBisCO even at the early stages of carboxysome

assembly, possibly in the internal core of the carboxysome. Assuming that the

carboxysome core grows by addition of material to the outside of a stable paracrystalline

RuBisCO assembly (Chen *et al*., 2013), these results also suggest that M58 does not

assemble as a separate layer around a pre-existing M35/RuBisCO core.

**M58 is not exclusive to a subshell layer**

3D-SIM experiments and quantitative analysis do not support a model of

carboxysome internal structure in which M58 is limited to a sub-shell localization.  The

preexisting model described above proposes that M58 is not present within the primary

M35-Rubisco matrix. This makes logical sense based on the fact that M35, which does

not contain the N-terminal domain, presumably can achieve denser packing of RuBisCO

enzymes.

We expected that if M58 was in fact limited to a sub-shell layer ~5-6nm beneath

the outer surface, it would appear as a ring similar to CcmK4 by 3D-SIM. While SIM

imaging of CcmK4 did reveal the expected ring-like distribution, carboxysomes with

fluorescently-labeled M58 appeared predominantly as solid objects, with only a very a

small population (0.58%) showing dips in fluorescence intensity at their centers.

Consequently, we find no strong evidence by super-resolution microscopy to support a

model of carboxysome structure in which M58 is excluded from the carboxysome core,

suggesting that the internal carboxysome structure contains a mixed population of both

M35 and M58 coordinating with RuBisCO.

317           This finding raises new questions about the biogenesis of carboxysomes. For

example, M58 is thought to play a role in recruitment of shell components (Cot *et al*.,

2008; Long *et al.,* 2011; Kinney *et al*., 2012; Cameron *et al*., 2013). However, shell

proteins are recruited to carboxysomes only late in their maturation process (Chen *et al.,*

2013). Therefore, if M58 is present even at the early stages of carboxysome assembly,

what factors are responsible for delaying shell recruitment? This question is a promising

avenue for future work.

**Materials and Methods:**

**Western blot analysis**

327           For SDS-PAGE, log-phase ($OD_{750}$ = 1.0) cultures of cyanobacterial strains were

pelleted and frozen at -20°C. Frozen pellets were thawed in B-PER Bacterial Protein

Extraction Reagent (ThermoScientific), without DNAse or Lysozyme, and normalized by

wet pellet mass. Resuspended cells were mixed directly with SDS loading buffer and

boiled. Boiled samples were run on a 4-20% Tris-Glycine NuPAGE gel (Invitrogen) at

120V. SDS-PAGE separated proteins were transferred to a nitrocellulose membrane

with an iBlot2 transfer device (ThermoScientific).

334           The membrane was blocked with 5% skim milk in TBST (50mM Tris, 150mM

NaCl, 0.1% Tween 20), then probed with 1:10,000 of anti-CcmM primary antibody (a gift

of Ben Long), followed by a horseradish peroxidase-conjugated goat anti-rabbit

secondary antibody (Agrisera AS09 602). The membrane was incubated with

SuperSignal West Dura Extended Duration chemiluminescent substrate

(Thermoscientific) and imaged with the Bio-Rad ChemiDoc system.

**Transmission Electron Microscopy**

For TEM experiments, 1mL of mid-log cultures were pelleted (6k rpm, 5 min) and

concentrated 2x in sterile $dH_2O$. The cells were then mixed 1:1 with fixative (1.25%

formaldehyde, 2.5% glutaraldehyde, 0.03% picric acid, 0.1M sodium cacodylate buffer,

pH 7.4), incubated at RT for 30 min, then pelleted and stored in fixative at $4^{o}C$ overnight.

Fixed cells were embedded in epon resin and cut into ultra-thin sections (~60-80nm).

Sections were stained with uranyl acetate and lead citrate, then imaged on a Tecnai G2

Spirit Bio TWIN TEM.

**Cell culture**

Wild-type, mutant, and modified strains of *Synechococcus elongatus* PCC7942

were grown in BG-11 media, with 1 g/L HEPES adjusted to pH 8.5. Cultures were grown

at $30\text{-}35^{o}C$ under constant light with an approximate intensity of 1000 $lm/m^2$. Cultures

were grown in 125ml or 250ml flat bottom flasks, shaking at 150 rpm. Wild type 7942

and knockout complements were grown with ambient air conditions.  Mutant

$\Delta ccmM$::*HygR* cultures were supplemented  with 2% $CO_2$ and 25 µg/mL hygromycin B.

Strains with fluorescently labeled carboxysome components, carrying kanamycin or

spectinomycin resistance genes, were grown with 25 µg/mL kanamycin or 25µg/mL

spectinomycin respectively. For the CcmM complement strain, $\Delta ccmM$ + *mTagBFP2-*

*CcmM-mNeonGreen*, cultures were grown with 5 µg/mL kanamycin and 25 µg/mL

hygromycin B. For growth curve experiments, biological replicates were made by

isolating colonies grown on BG11 agar with 1 g/L HEPES, 50mM $HCO_3^-$, and 1mM

$Na_2S_2O_3$. Isolates were screened by PCR to confirm the presence or absence of the WT

*ccmM* gene. Growth curve cultures were prepared by washing starter cultures in fresh

BG11 pH 8.5 media, then diluting to OD750 of ~0.1 in 40 mL BG11 in 250mL flat bottom

flasks.

**Image acquisition**

For both conventional fluorescence and 3D-SIM experiments, 5µL of either mid-

log cells ($OD_{750}$ = 1-2) for timelapse experiments or early-log cells ($OD_{750}$ = 0.5 - 1.2) for

3D-SIM were spotted on 2% agarose BG11 pads. Agarose pads were placed sample-

side down in MatTek 35mm glass bottom microwell dishes with No. 1.5 coverslips.

Sterile water was spotted within the dish to mitigate dehydration of the agarose pad, and

the dish was sealed with parafilm. Though strains expressing labeled CcmM were under

the control of the IPTG inducible promoter P*trc*, no induction was used with these cells

due to adequate basal expression. Cells expressing RbcL-sfGFP were imaged on BG11

agarose pads with the addition of 50µM IPTG following previously reported protocols

(Savage *et al.*, 2010). SIM of uninduced RbcL-sfGFP cells also produced solid particles,

but were not as bright as induced cells and produced several false-positives in our

analysis pipeline (Supplemental figure S3). For timelapse experiments, samples were

recovered for >1h at 34-35°C, under ~1000 $lm/m^2$, and in ambient air, before being

imaged using a Nikon TE2000 and a 100x oil objective. Cells were kept at 30°C under

~800 $lm/m^2$ light intensity during timelapse, and imaged every 10min.

3D-SIM data was collected on a DeltaVision OMX V4 Blaze system (GE

Healthcare) equipped with a 60x / 1.42 N.A. Plan Apo oil immersion objective lens

(Olympus), and three Edge 5.5 sCMOS cameras (PCO).  mTagBFP2 fluorescence was

excited with a 405nm laser and collected with a 477/35 emission filter, mNeonGreen with

a 514nm laser and 541/22 emission filter, and sfGFP with a 488nm laser and a 528/48

emission filter.  Z-stacks of ~2 microns were acquired with a z-step of 125 nm and with

15 raw images per plane (five phases, three angles).  Spherical aberration was

minimized using immersion oil matching (Hiraoka *et al.*, 1990). Super-resolution images

were computationally reconstructed from the raw datasets with a channel-specific

measured optical transfer function (OTF) and a Wiener filter constant of 0.001 using

CUDA-accelerated 3D-SIM reconstruction code based on Gustafsson *et al.*

*(*2008).  Axial chromatic aberration was measured using TetraSpeck beads (Thermo

Fisher) a nano-grid control slide and multi-channel datasets were registered using the

image registration function in softWoRx.

**Image analysis**

SIM datasets were imported into MATLAB (Mathworks) using the bioformats

reader, followed by thresholding and background subtraction. The threshold value was

empirically determined and assessed by visualizing the resultant centroid position and

mask in MATLAB and Imaris (Bitplane AG) respectively. Segmented objects were further

analyzed if they met a cutoff for volume, surface area, sphericity, major-minor axis ratio,

and mean radius.

We calculated the MTC (max to center) ratio for each object. From the centroid,

the radially averaged intensity for each sub-sampled (0.5 pixels) radius was measured.

This was done for each Z-plane and then averaged over the Z-axis. A polynomial curve

was fit to the averaged radial profile of each segmented object in MATLAB. The most

suitable polynomial fit (up to 6th order) was selected based on meeting the Bayesian

Information Criterion.  Then, the 1st and 2nd derivative of each fitted curve were

calculated to find the root with the global maximum. The intensity at this root was

calculated and divided by the intensity at the centroid. This ratio was calculated for each

object and plotted versus its mean radius in MATLAB.

Notably, the resolvable carboxysome ring limit of 220nm that we estimate in our

analysis is well above the ~110nm theoretical resolution limit of 3D-SIM (at 520nm

wavelength; Gustafsson *et al.,* 2000; Gustafsson *et al*., 2008). The difference is likely

due to a combination of practical limitations along with the probable overestimation of

carboxysome diameter in our analysis pipeline due to convolution of the true object size

with the point spread function in the fluorescence images.

Timelapse data of Δ*ccmM* + *mTagBFP2-CcmM-mNeonGreen* were processed

with ImageJ (FIJI) (Schindelin *et al*., 2012). Timelapse data from two sequential

acquisitions of the same field of view were imported into FIJI with the Bio-Formats Import

plugin with autoscaling, concatenated, and separated by channel. Channels were

registered using rigid registration in the StackReg FIJI plugin, then background

subtracted with a 50-pixel radius rolling ball. Particle tracking was performed in FIJI with

the TrackMate plugin using the LoG detector (Laplacian of Gaussian filter) with 0.5μm

estimated blob diameter, 0.5 threshold, and sub-pixel localization. Automatically

generated tracks were manually extended backwards in time 20-30 frames by selecting

0.5μm diameter regions at or near site where the particle was first detected in order to

capture a background fluorescence level. Particle tracks were corrected for

photobleaching by dividing intensities by an average of 7 reference particles, normalized

to the first frame of the time lapse, in each channel. Reference particles were objects

observed to be relatively stable compared to particles that lost or gained fluorescence

due to carboxysome biogenesis. The same reference particles were used for correction

in both channels. To generate the timelapse montage, brightness and contrast values

were manually adjusted separately in each channel to enhance visibility in the figure.

**Cloning and strain construction**

Plasmids used for *S. elongatus* strain construction were made by Gibson

assembly. Tagged CcmM plasmids were cloned by inserting the *ccmM* gene, along with

mNeonGreen and mTagBFP2 coding sequences with additional flexible linker

sequences, into the previously described pDFS21 neutral site II (NS2) integration

plasmid downstream of the IPTG inducible P*trc* promoter (Savage *et al*., 2010). The

*ccmM* gene was obtained directly from the *S. elongatus* PCC7942 genome via PCR with

Phusion polymerase (NEB). The *ccmK4* gene for cloning shell fusion plasmids was

obtained as a synthesized gBlock (IDT) and cloned into the pDFS21 integration plasmid.

*S. elongatus* strains were transformed with integration plasmids by incubating ~5

fold concentrated $OD_{750}$ 1-2 culture with 10-100ng plasmid at 30$^{o}$C in the dark overnight,

followed by plating transformants on selective BG11 agar (Clerico *et al*., 2007). The

mutant Δ*ccmM* strain was made by integrating a hygromycin B resistance gene into the

native *ccmM* ORF, followed by selective plating and screening (Sachdeva G., 2014).

**Acknowledgements**

We thank Ben Long (ANU) for many helpful discussions and the kind gift of anti-

CcmM serum, Jennifer Waters and the Harvard Cell Biology Microscopy Facility for their

guidance on SIM acquisition and analysis, Lin Shao for CUDA-accelerated SIM

reconstruction code, Hunter Elliott (Harvard Medical School) for his great help and

intellectual input on developing the SIM analysis pipeline, David Savage (UC Berkeley)

for collaborating with us on exploring other microscopy methods, the Harvard Medical

School EM facility for all their amazing work preparing samples and helping us with EM

imaging. We would also like to acknowledge Gairik Sachdeva and Simon Kretschmer

who originally constructed the Δ*ccmM::HygB* knockout strain in the Silver lab.

**References**

Andersson I, Backlund A. (2008). Structure and function of Rubisco. Plant Physio
and Biochem. 46:275-291.
468
469 Badger MR, Hanson D, Price GD. (2002). Evolution and diversity of CO2
470 concentrating mechanisms in cyanobacteria. Funct. Plant Biol. 29:161–
471 173.
472
473 Badger MR, Price GD. (2003). CO2 concentrating mechanisms in cyanobacteria:
474 molecular components, their diversity and evolution. J Exp.
475 Bot. 54:609–622.
476
477 Ball G, Demmerle J, Kaufmann R, Davis I, Dobbie IM, Schermelleh L. (2015).
478 SIMcheck: a Toolbox for Successful Super-resolution Structured Illumination
479 Microscopy. Sci Rep. 5:15915.
480
481 Berla BM, Saha R, Immethun CM, Maranas CD, Moon TS, Pakrasi HB. (2013).
482 Synthetic biology of cyanobacteria: unique challenges and opportunities. Front
483 Microbiol. 4(246):1-14.
484
485 Cameron JC, Wilson SC, Bernstein SL, Kerfeld CA. (2013). Biogenesis of a
486 bacterial organelle: the carboxysome assembly pathway. Cell. 155:1131-1140.
487
488 Chen AH, Robinson-Mosher A, Savage DF, Silver PA. Polka JK. (2013). The
489 bacterial carbon-fixing organelle is formed by shell envelopment of preassembled
490 cargo. PLOS One. 8(9):1-13.
491
492 Clerico EM, Ditty JL, Golden SS. (2007). Specialized techniques for site-directed
493 mutagenesis in cyanobacteria. Methods Mol Biol. 362:155-171.
494
495 Cot SS, So AK, Espie GS. (2008). A multiprotein bicarbonate dehydration
496 complex essential to carboxysome function in cyanobacteria. J Bacteriol.
497 190:936-945.
498
499 Gustafsson MGL. (2000). Surpassing the lateral resolution limit by a factor of two
500 using structured illumination microscopy. J of Microsc. 198(2):82-87.
501
502 Gustafsson MGL, Shao L, Carlton PM, Wang CJR, Golubovskaya IN, Cande WZ,
503 Agard DA, Sedat JW. (2008). Three-dimensional resolution doubling in wide-field
504 fluorescence microscopy by structured illumination. Biophys J. 94:4957-4970.
505
506 Hiraoka Y, Sedat JW, Agard DA. (1990). Determination of three-dimensional
507 imaging properties of a light microscope system. Partial confocal behavior in
508 epifluorescence microscopy. Biophys J. 57(2):325-333.
509

510 Kaneko Y, Danev R, Nagayama K, Nakamoto H. (2006). Intact carboxysomes in
511 cyanobacterial cell visualized by Hilbert Differential Contrast Transmission
512 Electron Microscopy. J Bacteriol. 188(2):805-808.
513
514
515 Kerfeld CA, Melnicki MR. (2016). Assembly, function and evolution of
516 cyanobacterial carboxysomes. 32:66-75.
517
518 Kinney JN, Salmeen A, Cai F, Kerfeld CA. (2012). Elucidating essential role of
519 conserved carboxysomal protein CcmN reveals common feature of bacterial
520 microcompartment assembly. 2012. J Biol Chem. 287(21):17729-17736.
521
522 Long BM, Murray BR, Whitney SM, Price GD. (2007). Analysis of carboxysomes
523 from *Synechococcus* PCC7942 reveals multiple rubisco Complexes with
524 carboxysomal proteins CcmM and CcaA. J Biol Chem. 282(40):29323-29335.
525
526 Long BM, Rae BD, Badger MR, Price GD. (2011). Over-expression of the ß-
527 carboxysomal CcmM protein in *Synechococcus* PCC7942 reveals a tight co-
528 regulation of carboxysomal carbonic anhydrase (CcaA) and M58 content.
529 Photosynth Res. 109:33-45.
530
531 Long BM, Tucker L, Badger MR, Price GD. (2010). Functional cyanobacterial ß-
532 carboxysomes have an absolute requirement for both long and short forms of the
533 CcmM Protein. Plant Physiol. 153:285-293.
534
535 Polka JK, Silver PA. (2013). Building synthetic cellular organization. Mol Biol Cell.
536 24:3585-3587
537
538 Price GD, Howitt SM, Harrison K, Badger MR. (1993). Analysis of a genomic
539 DNA region from the cyanobacterium *Synechococcus* sp. Strain PCC7942
540 involved in carboxysome assembly and function. J Bacteriol. 175(10):2871-2879.
541
542 Price GD, Badger MR, Woodger FJ, Long BM. (2007). Advances in
543 understanding the cyanobacterial $CO_2$-concentrating-mechanism (CCM):
544 functional components, Ci transporters, diversity, genetic regulation and
545 prospects for engineering into plants. J Exper Bot. 59(7):1441-1461.
546
547 Rae BD, Long BM, Badger MR, Price GD. (2012). Structural determinants of the
548 outershell of ß-carboxysomes in *Synechococcus elongatus* PCC7942: roles for
549 CcmK2, K3-K4, CcmO, and CcmL. PLOS One. 7(8):1-11.
550
551 Rae BD, Long BM, Badger MR, Price GD. (2013). Functions, compositions, and
552 evolutions of the two types of carboxysomes: polyhedral microcompartments that
553 facilitate $CO_2$ fixation in cyanobacteria and some proteobacteria. Microbiol Mol
554 Biol Rev. 77(3):357-379.
555

556   Rae BD, Long BM, Whitehead LF, Förster B, Badger MR, Price GD. (2013).
557   Cyanobacterial carboxysomes: microcompartments that facilitate $CO_2$ fixation. J
558   Mol Microbiol Biotechnol. 23:300-307.
559

560   Sachdeva G. (2014). Controlling fluxes for microbial metabolic engineering.
561   Doctoral dissertation, Harvard University.
562

563   Savage DF, Afonso B, Chen AH, Silver PA. (2010). Spatially ordered dynamics
564   of the bacterial carbon fixation machinery. Science. 327:1258-1261.
565

566   Schindelin J, Arganda-Carreras I, Frise E. *et al*. (2012). Fiji: an open-source
567   platform for biological-image analysis. Nature Methods. 9(7): 676-682.
568

569   Schindelin J, Rueden CT, Hiner MC, Eliceiri KW. (2015). The ImageJ ecosystem:
570   An open platform for biomedical image analysis. Mol Reprod Dev. 82(7-8):518-
571   29.
572

573   Tanaka S, Sawaya M, Philips M, Yeates T. (2008). Insights from multiple
574   structures of the shell proteins from the ß-carboxysome.  Prot Sci. 18:108-120.
575

576   Woodger FJ, Badger MR, Price GD. (2005). Sensing of inorganic carbon
577   limitation in *Synechococcus* PCC7942 is correlated with the size of the internal
578   inorganic carbon pool and involves oxygen. Plant Physiol. 139(4):1959-1969.
579

580   Yokoo R, Hood RD, Savage DF. (2015). Live-cell imaging of cyanobacteria.
581   Photosynth Res. 126(1):33-46.

596
Figure 1. Current model of the cyanobacterial carboxysome internal structure.
(A) *S. elongatus* utilizes the carboxysome to concentrate and protect its
RuBisCO enzymes. Through a multi-step process, bicarbonate passes into the
carboxysome, is converted to $CO_2$ by carbonic anhydrase, and is then fixed by
RuBisCO to form 3-phosphoglyceric acid (3-PGA). (B) Legend of proteins and
protein complexes of the carboxysomes interior. *CcmM* is an essential
carboxysome gene that is translated as two different isoforms, the full-length
58kDa M58 and a truncated 35kDa form, M35. CcmM has an N-terminal domain
that complexes with CcaA carbonic anhydrase, and three repetitive C-terminal
domains that have homology with the small subunit of RuBisCO. (C) These small
subunit-like domains (SSU-like), which make up the majority of the M35 protein,
are believed to displace a portion of the RuBisCO small subunits and physically
tether the enzymes together within the core of the carboxysome, to the exclusion
of the long M58 isoform which has been hypothesized to remain just beneath the
carboxysome shell.

612
Figure 2. Fluorescently tagged *ccmM* integrated into the genome of a Δ*ccmM*
knockout produces both CcmM isoforms and and restores growth in ambient air
conditions. (A) The Δ*ccmM* knockout was made by integrating a hygromycin
resistance marker into the genomic site of *ccmM*. (B) Diagram of fluorescently
tagged *ccmM*. N- and C-terminal fusions of mTagBFP2 and mNeonGreen
produces short and long isoforms with different fluorescent characteristics. This
construct was integrated at Neutral Site II of the 7942 Δ*ccmM* genome to yield
the Δ*ccmM* + *mTagBFP2-CcmM-mNeonGreen (+B-ccmM-N)* strain. (C) Anti-
CcmM western blot of WT, Δ*ccmM*, and *+B-ccmM-N*. Full length M58 isoform of
CcmM is ~58 kDa, while the truncated M35 is ~35 kDa. The fluorescent fusions
of mTagBFP2-M58-mNeonGreen and M35-mNeonGreen have expected
molecular weights of approximately 113 kDa and 63 kDa respectively. (D)
Growth curves of WT 7942, Δ*ccmM*, and the *+B-ccmM-N* genetic complement.
Dotted lines denote cultures grown with 2% supplemented $CO_2$. Solid lines
denote cultures grown in ambient air conditions. n = 3 biological replicates, error
bars = standard deviation.

A    *S. elongatus* PCC7942

B    *S. elongatus ΔccmM::HygB*

C    *S. elongatus ΔccmM::HygB + mTagBFP2-ccmM-mNeonGreen*

629
Figure 3. Transmission electron micrographs show the formation of
carboxysomes in Δ*ccmM* cells complemented with fluorescently tagged *ccmM*.
(A-D) Carboxysomes (white arrows) in wild type PCC7942 cells show
characteristic hexagonal geometry. Diameters of WT carboxysomes labeled in
micrographs range from 150 - 280 nm with an average of 180 $\pm$ 33nm (N=???).
(E-H) Micrographs of the Δ*ccmM* knockout show the absence of carboxysomes.
(I-L) Integration of the fluorescently tagged *ccmM* construct *B-ccmM-N* into the
Δ*ccmM* knock out rescues the formation of apparently normal carboxysomes.
These carboxysomes had geometry similar to that observed in wild type cells.
Carboxysomes with tagged CcmM appeared to range more widely in size than in
wild type cells with an average diameter of 319 $\pm$ 92nm (N=???). Some (J & L)
were of typical size (200nm in diameter), while others (K) are larger at 300-330
nm. Larger carboxysomes were also observed, like that seen in (I), which
approaches 600nm at its widest point.

Figure 4. CcmM isoforms co-assemble at the site of new carboxysomes. (A) Merged montage of carboxysome biogenesis in the Δ*ccmM* + *mTagBFP2-ccmM-mNeonGreen* gene complement strain shows simultaneous co-localization of both blue (mTagBFP2-M58) and green (M58-mNeonGreen / M35-mNeonGreen) particles. White arrows indicate the new carboxysome. White asterisk indicates the parent carboxysome. (B) Channel specific intensity plots of carboxysome biogenesis depicted in panel A reveal sigmoidal-like dynamics in assembly timing in M35 and M58 CcmM isoforms. Vertical blue and green lines denote point at which automatic particle tracking first identified the new carboxysome. Tracks were extended backwards to measure background by manually measuring spot at or near the new particle (Methods). Four-parameter logistic equations were fit to each track; black solid or dotted vertical lines denote half-max values of each channel. (C) Statistics of sigmoidal curves fit to 8 measured biogenesis events find similar assembly dynamics in both green and blue channels. Assembly time was measured from first time point of automatic tracking to first value $\geq$ 0.8 on normalized intensity plot.

Figure 5. SIM of fluorescently tagged carboxysomes resolves the ring of the outer shell, and an absence of shell-like localization of either RuBisCO or CcmM. (A-C) SIM of CcmK4-mTagBFP2 resolves the carboxysome shell as a hollow ring, while (D-F) SIM of RbcL-sfGFP produces solid particles. (G-K) SIM of ΔccmM + mTagBFP2-ccmM-mNeonGreen produces RuBisCO-like, apparently solid particles. M35/M58-mNeonGreen particles are pseudocolored magenta. Numbers identify individual carboxysomes and correspond to magnified panels on the right.

Figure 6. Analysis of 3D-SIM data. Using radially averaged intensity profiles of fluorescent particles in 3D-SIM data, maximum intensity to center (MTC) intensity ratios can measure the degree to which a fluorescent object is resolved as a ring. (A) Analysis of RbcL-sfGFP 3D-SIM data found no objects with a max-to-center ratio (MTC) greater than 1.04, with an average ratio of $1.002 \pm 0.005$ and average particle size of $244 \pm 45$ nm marked by the vertical line. An upper ratio limit of 1.05, below which objects are likely to be solid, was calculated by adding two standard deviations to the RbcL-sfGFP max ratio. The 1.05 MTC cutoff is marked by a horizontal line on all plots. (B) Size versus intensity ratio plot of CcmK4-sfGFP shows a population of particles (345, 9.97%) with MTC ratios $\geq$ 1.05. (C) Plot of CcmK4-mTagBFP2 shows a two-fold increase in the fraction of ring-like particles compared to the GFP tagged version (474, 18.16%). (D) Analysis of particles in the blue channel of the $\Delta ccmM \pm mTagBFP2\text{-}CcmM\text{-}mNeonGreen$ strain, representing M58, finds only a few particles with noticeable ring-like appearance (19, 0.58%), with the vast majority of particles appearing

solid. Based on the size of smallest detected ring in the tagged shell strains (196nm) and a shell thickness of 5-6nm, the minimum diameter at which ring-like shapes can be detected is calculated as 220nm; this is marked in gray. Ring-like particles within this region are unlikely to be resolved due to their small size. (E) Plot of particles in the green channel of Δ*ccmM* $\pm$ *mTagBFP2-CcmM-mNeonGreen* strain, representing M58-mNeonGreen and M35-mNeonGreen are predominantly solid with only 2 ring-like structures identified (2, 0.06%). (F) Table summarizing analysis statistics. In microscopy panels, scale bars are 200nm, and numbers denote MTC ratio.

726
727 Figure S1. Protein degradation in variant *ccmM* gene complements. Western blot
728 analysis of Δ*ccmM* + *mNeonGreen-ccmM-mTagBFP2*, and Δ*ccmM* + *ccmM-mTagBFP2*
729 revealed prominent unexpected bands (marked by red asterix) that could be indicative of
730 protein degradation in the cell. The Δ*ccmM* + *mTagBFP2-ccmM-mNeonGreen* strain did
731 not show similar unexpected bands. L = cell lysate and P = post-lysis pellet.
732

733
734 Figure S2. Induction of the P*trc*::Δ*ccmM* + *mTagBFP2-ccmM-mNeonGreen* in *S.*
735 *elongatus* leads to protein aggregation. SIM images of Δ*ccmM* + *mTagBFP2-ccmM-*
736 *mNeonGreen* cells induced with 50µM IPTG show large aggregations of fluorescent
737 signal near the cell polls.

738
Figure S3. Uninduced cells expressing RbcL-sfGFP still produce solid particles.
Qualitative data and image analysis results of PCC7942 RbcL-sfGFP cells without any
IPTG induction show similar incidence of low MTC ratio solid particles compared to cells
induced with 50μM IPTG. Three particles were identified as ring-like with MTC ratios >
1.05, but were confirmed to be three particle clusters incorrectly analyzed as single
objects by our analysis program.
745
746

| Strain | Tag | IPTG (μM) | # Imgs | Refractive Index | Wiener Filter | RIH (477) | RIH (528) | RIH (541) | RIH (608) |
|--------|-----|-----------|--------|------------------|---------------|-----------|-----------|-----------|-----------|
| DFS724 | RbcL-sfGFP | 50 | 31 | 1.518 | 0.01 | | 12.74 ± 1.63 | | 2.78 ± 0.66 |
| AHC134 | CcmK4-sfGFP | - | 31 | 1.517 | 0.01 | | 13.15 ± 1.78 | | 3.53 ± 1.08 |
| MJN31 | CcmK4-mTagBFP2 | 0 | 29 | 1.517 | 0.01 | 8.28 ± 1.84 | | | 2.77 ± 0.96 |
| MJN28 | mTagBFP2-CcmM-mNeon | 0 | 47 | 1.517 | 0.01 | 9.45 ± 2.80 | | 12.58 ± 2.51 | 3.10 ± 0.84 |

747
Tables S1. Structured Illumination Microscopy reconstruction scores. Reconstructed
Intensity Histogram scores (RIH) were averaged across all image sets for each strain
used for image analysis. Scores of 6-12 are considered good, and greater than 12
excellent. Columns with RIH scores are channel specific. Channel 608 captured the red
auto-fluorescence of cells which was not used for any analysis. Error = Standard
deviation.
754

**A**

| Welch's ANOVA | Degrees of freedom | $F_s$ | $P$ |
|---|---|---|---|
| Among groups | 4.00 | 160.48 | $6.51 \times 10^{-132}$ |
| Within groups | 7377.44 | | |

**B**

| Games-Howell test | Mean difference | q | $> q_{crit}$ 5.484 ($\alpha=0.001$) |
|---|---|---|---|
| CcmM-mNeon vs CcmK4-sfGFP | 0.020465 | 22.20 | Significant |
| CcmM-mNeon vs CcmK4-mTagBFP2 | 0.033914 | 26.34 | Significant |
| CcmM-mNeon vs mTagBFP2-CcmM | 0.001283 | 9.56 | Significant |
| CcmM-mNeon vs RbcL-sfGFP | 0.000050 | 0.57 | Not significant |
| CcmK4-sfGFP vs CcmK4-mTagBFP2 | 0.013449 | 8.51 | Significant |
| CcmK4-sfGFP vs mTagBFP2-CcmM | 0.019182 | 20.69 | Significant |
| CcmK4-sfGFP vs RbcL-sfGFP | 0.020515 | 22.26 | Significant |
| CcmK4-mTagBFP2 vs mTagBFP2-Ccm | 0.032631 | 25.27 | Significant |
| CcmK4-mTagBFP2 vs RbcL-sfGFP | 0.033964 | 26.38 | Significant |
| mTagBFP2-CcmM vs RbcL-sfGFP | 0.001333 | 10.05 | Significant |

755
756 Table S2. Statistical test results of particle MTC ratio comparison.
757
758

| Strain or Plasmid | Genotype | Resistance | Reference |
|---|---|---|---|
| *E-coli strains* | | | |
| DH5-alpha | Host strain for plasmid construction (PAS644) | | |
| DH10-beta | Host strain for plasmid construction (PAS641, PAS642, PAS643) | | |
| Turbo | Host strain for plasmid construction (PAS645) | | |
| *S. elongatus strains* | | | |
| PCC7942 | Wild-type *Synechococcus elongatus,* ATCC 33912 | | |
| Δ*ccmM* | PCC7942 Δ*ccmM*::*HygR* knock out | HygB | |
| PAS646 | PCC7942 *Ptrc:CcmK4-mNeon* integrated in NS2 | Kan | This work |
| PAS647 | PCC7942 Δ*CcmM::HygR* + *Ptrc:mNeon-CcmM-mTagBFP2* integrated in NS2 | HygB, Kan | This work |
| PAS648 | PCC7942 Δ*CcmM::HygR* + *Ptrc:CcmM-mTagBFP2* integrated in NS2 | HygB, Kan | This work |
| PAS649 | PCC7942 Δ*CcmM::HygR* + *Ptrc:mTagBFP2-CcmM-mNeon* integrated in NS2 | HygB, Kan | This work |
| PAS650 | PCC7942 *Ptrc:CcmK4-mTagBFP2* integrated in NS2 | Kan | This work |
| PAS651 | PCC7942 *Papca:CcmK4-mNeon* integrated in NS1 | Sp | Chen et al., 2013 |
| PAS652 | PCC7942 *Ptrc:RbcL-sfGFP* integrated in NS2 | Kan | Savage et al., 2010 |
| *Plasmids* | | | |
| pAM2314 | *S. elongatus* Neutral Site I integration vector | Sp | Mackey et al., 2007 |
| pAM1579 | *S. elongatus* Neutral Site II integration vector | Kan | Mackey et al., 2007 |
| pDFS21 | derivative of pAM1579 with *trc* promoter from pTRC99 and *lacI* | Kan | Savage et al., 2010 |
| pAHC134 | P*apca*::*ccmK4-sfGFP* | Sp | Chen et al., 2013 |
| pDFS724 | P*trc*::*RbcL-sfGFP* | Kan | Savage et al., 2010 |
| PAS641 | P*trc*::*mNeonGreen-CcmM-mTagBFP2* cloned into pDFS21 at XbaI and NotI sites | Kan | This work |
| PAS642 | P*trc*::*CcmM-mTagBFP2* cloned into pDFS21 at XbaI and NotI sites | Kan | This work |
| PAS643 | P*trc*::*mTagBFP2-CcmM-mNeonGreen* cloned into pDFS21 at XbaI and NotI sites | Kan | This work |
| PAS644 | *Ptrc*::*CcmK4-mNeonGreen* cloned into pDFS21 at XbaI and NotI sites | Kan | This work |
| PAS645 | *Ptrc*::*CcmK4-mTagBFP2, mTagBFP2* cloned in place of *mNeonGreen* in pDFS21 | Kan | This work |

759
760 Table S3. Strains and plasmids

761
762
763