Feedback Determines the Structure

of Correlated Variability in Primary Visual Cortex

Adrian G. Bondy[1,2], Ralf Haefner[3], and Bruce G. Cumming[1]

[1]   Laboratory of Sensorimotor Research, National Eye Institute, NIH

    49 Convent Drive, Rm. 2A50

    Bethesda, MD 20892

[2]   Brown-NIH Neuroscience Graduate Partnership Program

    185 Meeting Street, Box GL-N

    Providence, Rhode Island 02912

[3]   Brain & Cognitive Sciences, University of Rochester

     Meliora Hall

     Rochester, NY 14627

Correspondence:

Adrian Bondy

Princeton Neuroscience Institute

Washington Road

Princeton, New Jersey 08540

adrian.bondy@gmail.com

301-451-4926

The variable spiking discharge of sensory neurons in response to a fixed stimulus tends to be weakly correlated (spike-count correlation, r_{sc}). This is widely thought to reflect stochastic noise in shared sensory afferents, in which case it places strict limits on the fidelity of sensory coding. However, it may also be generated by changes over time in feedback from higher-order brain regions. Currently, the relative contribution of these sources is unknown. We addressed this by measuring spiking activity in populations of primary visual cortical (V1) neurons in rhesus monkeys performing different visual discrimination tasks on the same set of visual inputs. We found that the structure of r_{sc} (the way r_{sc} varied with neuronal stimulus preference) changed dramatically with task instruction, despite identical retinal input. Therefore, even at the earliest stage in the cortical visual hierarchy, r_{sc} structure during task performance primarily reflects feedback dynamics, not noise in sensory afferents. As a consequence, previous analyses of how r_{sc} constrains sensory processing need not apply. Furthermore, these results imply that correlations between the activity of single neurons and choice depend on feedback engaged by the task.

Judgments made about sensory events (i.e. perceptual decisions) rely on the spiking discharge of sensory neurons. For this reason, there has been longstanding interest in the observation that this discharge tends to be variable given a fixed stimulus[1,2]. In principle, this variability could confound perceptual judgments, impairing the fidelity of sensory information in the brain. Even worse, this variability tends to be weakly correlated amongst sensory neurons (spike-count correlation; r_{sc})[3], meaning it cannot trivially be averaged away[4]. For this reason, r_{sc} is widely referred to as "correlated noise"[5–8].

This way of thinking has underlied several influential lines of research in systems neuroscience. One has sought to understand the magnitude of the perceptual impairment introduced by r_{sc} in different behavioral contexts[5,8–15]. When r_{sc} is distributed in such a way that correlated fluctuations mimic the sensory events being detected or discriminated, it could severely impair perceptual accuracy[11,15,16]. A related line of research has sought to understand how correlated variability affects the choices subjects make in perceptual discrimination tasks from trial to trial[17–19]. These studies have shown that r_{sc} structure

can give rise to a weak correlation between variability in single neurons and perceptual reports (Choice

Probability; CP), consistent with the notion that CP observed in real neurons reflects the causal influence

of correlated sensory neuronal variability on perception.

However, we currently know very little about the origin of r_{sc}, making it unclear to what degree

these conclusions are correct. A frequent (although typically unstated) assumption is that r_{sc} in sensory

neurons is generated by shared variability in common afferent inputs. Consistent with this idea, r_{sc}

correlates with the physical proximity and similarity in stimulus preference of neuronal pairs[8,20–23], which

are also predictive of the degree of feedforward input convergence. If this explanation is correct, it

supports the traditional view of r_{sc} as "confounding noise" since it arises from stochastic processes in the

sensory encoding pathway. However, the bulk of synaptic inputs to sensory cortical neurons are not

strictly "feedforward" in nature[24,25]. Consequently, variation over time in shared inputs from downstream

areas (i.e. "top-down"; "feedback"), may make a significant contribution to r_{sc}. These signals may reflect

endogenous processes like attention, arousal, or perceptual state, and could be under voluntary control. In

principle, this source of correlated variability need not confound perceptual judgments, but instead reflect

ongoing neuronal computations.

Several recent studies have shown that r_{sc} does change to some degree with task context[12,26–28],

suggesting a top-down component. These studies have shown that r_{sc} in populations of sensory neurons

can either increase or decrease depending on attentional state or other task demands. However, prior

studies have made only limited measures of r_{sc} structure and how this changes with task, yet these are

critical for understanding how r_{sc} arises and how it relates to task performance. Furthermore, the relative

magnitude of feedforward versus top-down contributions to r_{sc} has not been determined. It also unknown

whether task-dependent changes in r_{sc} reflect an adaptive reduction of sensory noise or whether r_{sc} is, in

the first instance, generated by variability over time in top-down inputs reflecting downstream

computations.

In the present study, we used large-scale neuronal population recordings in behaving macaques, along with careful behavioral control and a novel analytical approach, to significantly advance our understanding of these fundamental questions. Subjects performed different orientation discrimination tasks using the same set of stimuli. The only difference between tasks was the set of orientations being discriminated. If r_{sc} primarily reflects noisy sensory encoding, it should be invariant to changes in the task given fixed retinal input. Alternatively, if r_{sc} structure changes dynamically with the task, this would indicate that it reflects top-down signals. This experimental approach, inspired by a previous study[27], was combined with large-scale population recordings, allowing us to estimate the full r_{sc} matrix – that is, how r_{sc} varies as a function of all possible combinations of pairwise orientation preference. This made it possible to directly infer which components were fixed and which changed with the task. Strikingly, we could not identify a component that remained fixed. Instead we observed a pattern of task-dependent changes that was highly systematic, and could be modeled as the effect of a single modulatory input that targets the two task-relevant subpopulations of V1 neurons in an alternating fashion across trials.

These data give unprecedented insight into the functional role of r_{sc} structure in task performance. First, they show that the task-dependent changes in r_{sc} structure appear to degrade the task performance of an ideal observer of V1 activity alone, because they mimic task-relevant stimulus changes. However, our discovery of the feedback origin of these correlations means that they need not degrade performance, and points to the possibility that they may instead be a signature of ongoing neuronal computations. Indeed, recent circuit models of perceptual inference predict feedback signals whose statistics reflect the subject's prior beliefs about the task, yielding predictions which closely match our obervations[29,30]. Second, we show quantitatively that these feedback dynamics are the primary source of the choice-related activity we observed in V1, clarifying an ongoing debate[31] about the interpretation of choice-related signals in sensory neurons. We conclude that r_{sc} in sensory neurons reveals less than previously thought about the feedforward transmission of sensory information in the brain, but potentially much more about the interareal computations underlying sensory processing.

Results

We trained two rhesus monkeys (*Macaca mulatta*) to perform a two-alternative forced choice

(2AFC) coarse orientation discrimination task (Fig. 1), used previously[32]. On a given trial, the subject was

shown a dynamic, 2D filtered noise stimulus for 2 seconds, after which it reported the stimulus orientation

by making a saccade to one of two choice targets (oriented Gabor patches). Different task contexts were

defined by the pair of discriminandum orientations. The stimuli were bandpass filtered in the Fourier

domain to include only orientations within a predetermined range. The stimulus filter was centered on one

of the two task orientations and its orientation bandwidth was used to control task difficulty. We included

0%-signal trials, for which the stimuli were unfiltered for orientation (and thus the same regardless of

context), to examine the effect of task context on r_{sc} in the presence of a fixed retinal input.

In order to detect any effect of task context on r_{sc} structure, it is critical that subjects based their

choices on the presence of the correct orientation signals. To ensure this, we used psychophysical reverse

correlation[32–34] to directly measure the influence of different stimulus orientations on the subject's choices

(the "psychophysical kernel"). We found that subjects required multiple days of retraining after a change

in the task context to fully update their psychophysical kernel. For this reason, we kept the task context

fixed for the duration of each recording session, and only undertook recordings in a new task context after

subjects had updated their kernel (Supplementary Fig. 1). This is a significant advance over past studies of

the effect of task context on neuronal responses, which typically have not quantified the extent to which

behavioral strategy truly matches task instruction.

We recorded spiking activity in populations of single V1 neurons using multi-electrode arrays

while the subjects performed the task. We determined the preferred orientation of each neuron by

measuring its response to oriented stimuli (see Methods) in separate blocks of trials during which subjects

passively fixated. Neurons were excluded from analysis if they were not well orientation tuned. The final

dataset includes 811 simultaneously recorded pairs from 200 unique cells across 41 recording sessions.

For each pair, we calculated its r_{sc} value as the Pearson correlation between the set of trial-duration spike-

counts across trials of the same stimulus condition. While measuring r_{sc} only across 0%-signal trials

isolated any changes due to the task context, we found similar results within each signal level (Fig. 6).

Therefore, to increase statistical power, we report r_{sc} values measured across all trials, after normalizing

spike counts to remove the effect of stimulus drive on firing rates.

R_{sc} structure changes systematically with task context

Recording large populations gave us the power to measure the full "r_{sc} matrix": that is, how r_{sc}

varied as a function of all possible combinations of orientation preference. This is the first time that such

detailed measures of r_{sc} structure have been made while animals perform a discrimination task. To assess

the presence of task-dependent r_{sc} structure in the data, we we first divided the recording sessions into two

groups based on the task context used (Fig. 2b). We estimated the smoothed, average r_{sc} matrix associated

with each subset (Fig. 2a,c) by pooling r_{sc} values measured across the subset of sessions along with

measures of the neuronal preferred orientation. Across both subsets of sessions, we observed a tendency

towards higher values of r_{sc} for pairs of neurons with more similar orientation preferences (i.e. higher

values closer to the diagonal of the matrix), consistent with numerous prior observations[3] (Fig. 2d).

Traditionally, such observations were presumed to reflect "limited-range correlations" that depend only on

similarity in stimulus preference[5,9,10], equivalent to a rotationally-symmetric (Toeplitz) correlation matrix.

In contrast, in our data this was due to distinct patterns in the two matrices: we observed the highest values

of r_{sc} amongst pairs that shared a preferred orientation close to a discriminandum, and the lowest values of

r_{sc} tended to occur amongst pairs preferring opposite task orientations. Because the task context differed

between the two subsets, this yielded matrices with a lattice-like pattern offset along the diagonal by an

amount reflecting the task context. In other words, r_{sc} structure changed dramatically with task context,

consistent with the presence of task-dependent feedback and inconsistent with a fixed r_{sc} structure

primarily driven by sensory afferent noise.

To summarize this task-dependent structure across the entire dataset (Fig. 2e) we expressed each

neuron's preferred orientation relative to the task orientations on its respective recording session, such that

0º and 90º always indexed the task orientations. This combined matrix clearly illustrates the task-

dependent pattern of r_{sc} structure in the V1 population, a pattern that was consistent across both subjects

(see Supplementary Fig. 2a,b). This finding is consistent with, but expands upon, more limited measures

made in area MT in a direction discrimination task[27]. (For a detailed comparison with those results, see

Supplementary Discussion §4 and Supplementary Fig. 11.) As in previous studies, there was a great deal

of variability between individual r_{sc} values, even amongst pairs with similar orientation preferences and

task (Fig. 2d,f) demonstrating that factors not considered here also contribute to r_{sc}.

Importantly, we observed a different result during separate blocks of trials in the same recording

sessions, during which the subject fixated passively for reward but the same set of stimuli was shown.

During these blocks, the highest values of r_{sc} tended to occur along the diagonal, independent of

orientation preference or task (see Supplementary Fig. 3). This demonstrates that the task-dependent

pattern observed during task performance depends on active task engagement, and cannot be explained,

for instance, simply as an effect of adaptation to task experience. We performed a number of additional

analyses to rule out any possibility that our findings could be explained merely as an effect of changing

retinal input across task contexts, such as effects related to stimulus history or eye movements (see

Supplementary Discussion §1 and Supplementary Figs. 4-7). Taken together, these controls strengthen our

interpretation that centrally-generated signals reflecting task engagement underlie the task-dependent r_{sc}

structure we observed.

Segregating fixed and task-dependent components of r_{sc} structure

Our dataset of r_{sc} measurements made in large, heterogeneous populations across diverse task

contexts allowed us to directly estimate the r_{sc} structure that was fixed versus dynamically changing with

task. To do this, we modeled the raw r_{sc} values using two structured components: 1) a fixed r_{sc} matrix

describing the dependence of r_{sc} on pairwise orientation preference regardless of task, and 2) a task-

dependent r_{sc} matrix capturing the dependence of r_{sc} on pairwise orientation preference *relative* to the task

orientations. We used ridge regression to find the form of these two component matrices that best

predicted the raw r_{sc} measurements. To reduce the number of regressors without constraining the form

these two components could take, we parametrized the matrices as 8x8 grids of basis functions (see

schematic in Fig. 3a and Methods).

This modeling approach allowed us to address two related questions. First, the form of the fitted

components serves to identify the nature of the dynamic and fixed r_{sc} structure in the V1 population.

Second, comparing models that included either or both components provides a quantitative test for the

origin of the r_{sc} structure we observed. When we jointly fit both components to the data, the inferred task-

dependent component (Fig. 3c) recapitulated the lattice-like structure we observed in the smoothed,

average data (Fig. 2e). The fixed component (Fig. 3d) was smaller in amplitude and, interestingly,

appeared also to contain a weak lattice-like structure, offset by approximately 30º. This is likely due to the

fact that we did not uniformly sample across all possible task contexts, with tasks discriminating

orientations near 30º/120º being overrepresented (see Fig. 2b). Next, we compared reduced models in

which only one of the two components was used. Strikingly, cross-validated model accuracy was

increased when we removed the fixed component entirely, but reduced by about half when we removed

the task-dependent component (Fig. 3b). This suggests that the dependence of r_{sc} on orientation preference

in our data can be explained best as a completely dynamic phenomenon, with no detectable dependence

that is invariant to the task. We found that that all of these modeling results could be replicated when the

fixed and task-dependent components were parametrized in a different way (using a variable number of

basis functions with locations fit to the data, instead of a fixed grid of basis functions) suggesting they do

not depend on the particular parametric assumptions that were made (see Supplementary Discussion §3

and Supplementary Fig. 10).

R_{sc} depends on a large number of factors irrelevant to the present study, such as physical proximity

between pairs and similarity in tuning along many stimulus dimensions apart from orientation[3,22]. We

were only interested in the dependence on orientation preference and task, and therefore a model that

describes this correctly will only explain a small fraction of the variance in r_{sc}. (This can be appreciated in

figure 2d and f, where pairs with similar locations on the abscicca have substantial variation in r_{sc}.) To

estimate this fraction, we assessed the accuracy with which we could predict individual r_{sc} values from a

smoothed matrix built with other pairs. This showed that, in principle, 3.6% of the variance is explainable,

of which the majority was explained by the regression model above. We also found that, across cross-

validation folds, the fitted model components were highly consistent (mean correlation of 0.99),

suggesting the inferred structure is robust to noise in the data despite the low absolute value of variance

explained. Additionally, as we will discuss, the task-dependent pattern of r_{sc} we identify is likely to be

critically important during performance of the task being performed by our subjects.

R_{sc} structure during task performance reflects a single mode of variability

In our approach so far, we aimed to describe r_{sc} structure with as few assumptions as possible.

Having established that the observed r_{sc} structure can be best described assuming it is entirely task-

dependent, we next sought to identify a more parsimonious and intuitive description of this task-

dependency. We started with the observation that the pattern we observed – increased correlation between

pairs preferring the same task orientation and decreased correlation for pairs preferring opposing task

orientations – would be consistent with feature-selective feedback which varied in its allocation from trial

to trial between the two task-relevant orientations, as has been shown in recent theoretical studies[30,35]. To

quantify this observation, we performed an eigendecomposition of the smoothed, average r_{sc} matrix (Fig.

4a). We found that it had a single eigenvalue significantly larger than would be predicted by chance,

consistent with the correlation structure being determined largely by a single mode. Moreover, the first

eigenvector contained a peak and trough at the two discriminandum orientations, respectively, suggesting

a mode of variability which increases the firing rate of neurons supporting one choice and decreases the

firing rate of neurons supporting the other choice (Fig. 4b). To model this, we assumed all observed r_{sc}

values could be predicted by a single eigenvector which we constrained to be the difference of two von

Mises functions centered 90° apart with variable amplitude and width (see Fig. 4c). We found that this

simpler model in fact performed better than the more complex regression model in predicting individual

r_{sc} values, capturing about 80% of the explainable variance in r_{sc} (see Fig. 4e). This suggests that the r_{sc}

structure we observed in V1 could indeed be well described as the result of a single source of covariability

that changed dynamically with the task.

We compared the "single eigenvector" model with another simple parametric model that more

closely reflected standard assumptions about r_{sc} structure in sensory brain areas. This model predicted that

r_{sc} depends only on the difference in preferred orientation between pairs of neurons regardless of task[5,9,10]

("limited-range correlations" yielding an r_{sc} matrix with a diagonal ridge) and would be consistent with r_{sc}

structure due to common afferent inputs. We modeled this dependence as a von Mises function of

preferred orientation difference (Fig. 4d). This model performed much worse in predicting the observed

set of r_{sc} values, in fact not exceeding chance performance (Fig. 4e). (This result was qualitatively

replicated in both subjects individually; see Supplementary Fig. 10). Importantly, both of these simple

parametric models predict a dependence of r_{sc} on preferred orientation difference similar to what we found

in the data (Fig. 2d) and has been observed previously[8,20-23] – however, in the case of the "single

eigenvector" model, this is due to task-dependent changes in r_{sc} while for the "diagonal ridge" model,

there is no effect of task context. Notably, we found that during the passive fixation blocks, the "diagonal

ridge" model performed better (Supplementary Fig. 3), quantitively supporting the observation that the

task-dependent correlations we observed require active task engagement.

Effect of task-dependent r_{sc} structure on neural coding

We next sought to address the functional importance of the r_{sc} we observed on sensory coding.

Many studies have shown that r_{sc} in sensory neurons can impair information transmission, particularly

when r_{sc} resembles correlations due to task-related stimulus changes[5,8–15]. We estimated this task-related

stimulus correlation as the product of the slopes of a pair's mean response functions along the task axis

(i.e. as a function of orientation signal strength; Fig. 5a)[16], normalized by the product of the neuronal

variances. When we plotted these values as a smooth, task-aligned matrix (Fig. 5b), we observed a lattice-

like pattern strikingly similar to the observed r_{sc} matrix (Fig. 2e). Confirming this similarity, the task-

dependent component of r_{sc} structure identified by the regression model was highly correlated on a pair-

by-pair basis with the stimulus-induced correlations ($r=0.62$, Fig. 5c). This matches our earlier

observation that r_{sc} structure was consistent with feedback that alternatingly targeted the task-relevant

neuronal pools, which is similar to the effect of varying the stimulus along the axis defining the task.

Thus, the observed r_{sc} structure appears not to improve, but rather to degrade, the sensory

representation. However, our results highlight a problem with this interpretation and any purely

feedforward account of the functional role of r_{sc}. Namely, r_{sc} that is generated endogenously need not be

problematic at all (e.g. if the decoder had access to those endogenous signals). Indeed, the propagation of

feedback signals that are matched to the statistics of the relevant sensory stimuli may be an adaptive

strategy for bringing prior knowledge to bear, as predicted by recent models of probabilistic perceptual

inference[29,30]. R_{sc} resembling stimulus-induced correlations emerge in this model[29] as a consequence of

the subject developing the appropriate priors about the task, yielding predictions that both match our

empirical findings and offer a normative explanation.

Relationship between r_{sc} structure and perceptual choice

Correlations between trial-to-trial variability of single neurons and choice[36,37] have been frequently

observed throughout sensory cortex. Theoretical studies have emphasized that this suggests the presence

of spike-count correlation with a particular structure[17–19,38,39]. After all, if many sensory neurons have

variability that is correlated with choice, this implies that neurons supporting the same choice are

themselves correlated. However, this could be compatible with either or both of two causal mechanisms:

1) correlated fluctuations directly affect the choices a subject makes trial to trial (a feedforward source of

choice-related activity); or 2) the correlated fluctuations reflect variation across trials in a feedback signal

related to the upcoming choice (a feedback source). As we show, our detailed measures of r_{sc} structure

during task performance can address this ongoing debate.

First, we reasoned that a signature of feedback related to the upcoming choice would be r_{sc}

structure in V1 whose magnitude depends systematically on variability in choice. Consistent with this

prediction, we found that the amplitude of the r_{sc} structure was attenuated on high-signal trials relative to

0% signal trials, in a manner which depended systematically on signal strength (Fig. 6a,b). However, this

attenuation was modest, even at the highest signal level we analyzed (11% reduction), despite the highly

uneven distribution of choices. This rules out the extreme scenario in which feedback perfectly reflects

choice. Supporting this conclusion, we found that the r_{sc} structure, when calculated using only spikes from

different 200-ms windows during the trial, showed a stable timecourse (after a precipitous drop at the first

time point) and did not grow in amplitude with decision formation (Fig. 8). Taken together, these

observations support the conclusion that the r_{sc} structure reflects variation in feedback signals only

partially correlated with the subject's final choices. These could include a combination of bias, attention to

orientation, prior beliefs, and/or a decision variable.

Next, we assumed standard feedforward pooling (i.e. linear readout weights applied to the sensory

pool) to determine if the observed r_{sc} structure would be quantitatively consistent with the observed

choice-related activity. To do this, we made use of recent theoretical work which analytically relates r_{sc}

structure, readout weights, and choice-related activity[17]. We calculated Choice Probability (CP), which

quantifies the probability with which an ideal observer could correctly predict the subject's choices using

just that neuron's responses[36,37], for each recorded neuron. We found an average CP of 0.54 for task-

relevant neurons, significantly above chance level (Fig. 7a) and similar in magnitude to another study

using the same task[32]. We found that the r_{sc} structure we observed would be sufficient to produce a pattern

of CP across the population consistent with the data (Fig. 7), across a wide range of possible readout

schemes (Supplementary Fig. 9). Next, we considered the contribution of the different inferred sources of

r_{sc} to CP. (For top-down sources of correlation this is equivalent to assuming that the sensory population is

read out without taking into account the top-down signal.) This allows us to treat all sources of r_{sc}

equivalently, and compare them quantitatively. When we considered a population containing only the

"task-dependent" component of r_{sc} structure identified in the regression model (Fig. 3c), predicted CP was

only slightly reduced. Assuming only the "fixed" component (Fig. 3d), however, drastically reduced

predicted CP below what we observed (Fig. 7b,c). Thus, our data rule out the view that a significant

component of CP merely reflects the feedforward effect of stochastic noise in the afferent sensory

pathway. Instead, the main feedforward source of CP appears to depend on task-dependent changes in r_{sc}

structure that subsequently influence perceptual judgments.

Discussion

Spike-count correlations between sensory neurons have typically been described as reflecting noise

that corrupts sensory encoding[5,8–15]. However, little is known about the origin of r_{sc}, in particular whether

it is due to variability in shared sensory afferents or rather due to changes over time in feedback signals.

We addressed these outstanding questions by recording from populations of V1 neurons using multi-

electrode arrays while macaque subjects performed a set of orientation discrimination tasks. This approach

allowed us to estimate the entire matrix describing the dependence of r_{sc} on pairwise orientation

preference (Fig. 2), providing an unprecedently clear picture of r_{sc} structure in a behaving animal. By

determining to what extent the r_{sc} matrix was fixed, and what extent it changed with task, we could infer

the relative importance of feedforward and feedback pathways in generating it (Fig. 3). We found

systematic and novel structure in the r_{sc} matrix that changed in a predictable manner with the task. Using

multiple modeling approaches, we found that the fixed r_{sc} structure was much smaller than the task-

dependent structure, so much so that we could not estimate a fixed component reliably. Remarkably, a

single source of task-dependent feedback captured the pattern we observed (Fig. 4). This feedback input

increased and decreased the firing rate of neurons tuned for the two task-relevant orientations in a push-

pull manner.

This task-dependent modulation did not appear to be beneficial to task performance (Fig. 5), at

least not in the manner this has typically been examined (i.e. feedforward decoding of the sensory

population alone). Instead, the inferred feedback signals appeared to mimic task-relevant stimulus

changes, confounding the choices of an observer using only the sensory population. However, because the

correlations reflect downstream computations, they need to not be limiting in this way to the subject. Thus

our results highlight the fundamental insufficiency of considering the theoretical implications of r_{sc} in

terms of purely feedforward information transmission, as almost all such studies have done to date.

Consistent with several past studies[31,40,41], we found evidence for choice-related feedback, as

shown by the finding that correlated fluctuations in V1 are more pronounced on trials where the subject's

choices were more variable (Fig. 6). However, this effect was relatively weak, and we observed that task-

dependent r_{sc} structure did not grow in amplitude with decision formation (Fig. 8), suggesting processes

indirectly related to choice may be responsible for the feedback generating the correlations. More

importantly, we found that the standard assumption that correlated fluctuations influence choice through

feedforward pathways[17–19,38,39] predicted CP in the V1 population that matched the data (Fig. 7), the first

empirical test of the theoretical relationship between r_{sc} in sensory neurons, CP, and readout[17]. However,

the r_{sc} structure responsible changed with the task, demonstrating that it does not simply reflect afferent

noise. Taken together, our results instead favor the notion that choice-related activity comes about through

self-reinforcing loops of reciprocal connectivity between cortical areas, as has also been suggested by

other studies[30,41,42].

The inferred source of task-dependent feedback resembles previous reports about the effects of

feature-based attention on visual cortical neurons[35,43]. Feature-based attention enhances the firing rate of

neurons tuned for the attended stimulus feature, and decreases the firing rate of neurons tuned for

unattended stimulus features. One possibility is that our task engages feature-based attention which varies

over time in its allocation between the two task-relevant orientations. However, this does not appear to

provide an adaptive increase in the efficacy of transmission of relevant stimulus information[44,45], contrary

to traditional notions of neuronal attentional effects. However, as discussed above, once a top-down

contribution to correlations is recognized, it is not possible to infer the effectiveness of information

transmission from the activity of a population of sensory neurons alone.

Our findings thus emphasize the need for new theoretical models that predict context-dependent

feedback during perceptual processing. Currently, models based on hierarchical probabilistic

inference[29,30,46,47] do predict such feedback signals, and account for many of our experimental findings.

This work builds on the longstanding idea that the goal of a perceptual system is to generate valid

inferences about the structure of the outside world, rather than to faithfully represent sensory input[48,49].

This requires combining sensory input with prior beliefs. Fluctuations in prior beliefs therefore introduce

correlated variability. As a consequence, the theory provides a normative framework for predicting the

form this covariability will take in a particular behavioral context. During perceptual decision making,

correlations resembling those induced by the stimulus naturally emerge as a consequence of the subject

acquiring the appropriate prior beliefs about the structure of the sensory environment[29]. Further

development of this and other circuit models of perceptual decision making are needed to generate

quantitative predictions which can be further tested empirically.

Figure 1. Orthogonal orientation discrimination task. **a**. Schematic illustration of the task. Two example

task contexts shown (cardinal and oblique discriminations). The task context was fixed in a given recording

session, but varied across sessions. **b**. Psychometric function for monkey *'lem'*, example session. Black

curve is a probit fit, and error bars are 95% confidence intervals. **c**. Histograms showing the distribution of

psychometric thresholds across sessions for the two subjects. Threshold is defined as the signal level

eliciting 75% correct performance. Black triangle indicates the threshold corresponding to the example

session in (b). **d**. Example single stimulus frames corresponding to the two example task contexts in (a).

The stimuli consisted of dynamic, white noise filtered in the Fourier domain for orientation (see Methods).

The filter was centered on one of the two task orientations and its bandwidth determined signal strength.

Figure 2. R_{sc} structure in V1 depends systematically on task context. a,c. Observed r_{sc} matrices for the

two subsets of sessions grouped by task context, as indicated in (b). The matrices were obtained by

pooling the set of r_{sc} measurements made within each subset and applying a von Mises smoothing kernel

(approximating a 2D wrapped Gaussian with 15° s.d.). The colored dots indicate those regions

corresponding to pairs preferring the same or opposing task orientations. In both matrices, there are higher

r_{sc} between pairs preferring the same task orientation and lower r_{sc} for pairs preferring opposing task

orientations. **b**. Polar histogram shows the distribution of task contexts used across sessions, with color

indicating the division into two subsets. Note that the period is 90° because of the orthogonality of the

discriminanda. Colored arrows indicate the mean task context associated with each subset. **d.** Scatter plot

showing a weak, but significant, dependence of r_{sc} on the difference in preferred orientation of neuronal

pairs ($p=9*10^{-4}$, bootstrap test). Black line is (type II) regression line. **e.** Average r_{sc} matrix observed

across all session, shown in a task-aligned coordinate frame. Each pair's preferred orientations are

expressed relative to the task orientations (defined as 0º and 90º). Color scale as in (a). **f.** Scatter plot

showing a dependence of r_{sc} on distance from the peak (0º/0º or 90º/90º) in the matrix in (e). This

dependence was stronger than the dependence on difference in preferred orientation ($r=-0.17$, $p<0.05$,

bootstrap test), suggesting the task-aligned lattice pattern we observed captures a more important feature

of r_{sc} structure. Black line is (type II) regression line.

Figure 3. Segregating fixed and task-dependent components of r_{sc} structure. a. To estimate fixed and

task-dependent components of r_{sc} structure, we used a regression model (see Methods) with two component

matrices composed of a grid of 8x8 von Mises basis functions, whose amplitudes were fit to the data (the

set of raw r_{sc} measurements). **b**. Goodness-of-fit for the model that included both components and for two

reduced models that included only one of the two components. Values are expressed relative to an estimate

of the explainable variance in the data (see Methods). Error bars are +/- 1 SEM obtained from repeated 50-

fold cross-validation. **c,d**. Estimated components from the combined model. The amplitude of the task-

dependent component (c) was considerably larger than the fixed component (d) when jointly estimated, and

closely resembled the lattice-like shape of the task-aligned, average r_{sc} matrix (Fig. 2e). To quantify this

amplitude difference, we compared the relative variance across the fitted parameters for the two

components. This value was greater for the task-dependent component by a factor of 2.08. Note that

orientation preferences for the task-dependent component, but not the fixed component, are expressed

410   relative to the task orientations (i.e. rotated with the task). Mean r_{sc} values are close to 0 due to the inclusion

411   of a model constant.

Figure 4. R_{sc} structure during task performance reflects a single mode of variability. a.
Eigenspectrum for the average, task-aligned r_{sc} matrix in Fig. 2e. The largest eigenvalue was significantly larger than would be predicted by chance (p<0.001, permutation test, one-sided). The chance distribution (mean +/- 1 SEM in blue) was determined by adding a random offset to the preferred orientations of each pair (i.e. permuting each included r_{sc} value along the diagonal). **b.** The eigenvector corresponding to the largest eigenvalue in (a). (We first removed the mean r_{sc} value from the matrices to ignore any flat eigenvectors. Error bar is +/- bootstrap SEM). The dark gray vertical bar indicates the peak in the eigenvector +/- 1 bootstrap SEM. This was not significantly different from 0° (p=0.078, bootstrap test, one-sided), indicating close alignment between the dynamic pattern of r_{sc} in V1 and the subject's task. **c.** Schematic of "single eigenvector" model in which r_{sc} structure is described in terms of a single mode of variability in the population. The eigenvector is parameterized as the difference between two von Mises functions 90° apart. **d.** We compared this model to one in which r_{sc} structure depended only on the difference in preferred orientations, independent of task. This dependence was modeled as a von Mises function centered on zero, yielding an r_{sc} matrix with a diagonal ridge. **e.** Goodness-of-fit for the models in (c) and (d), calculated as % variance explained relative to our estimate of the explainable variance in the data, as in Fig. 3. Error bars obtained through repeated 50-fold cross-validation.

Figure 5. R_{sc} structure matches effect of task-related stimulus variability. **a**. Responses (mean +/- 1 SEM) to the stimuli used in the task at various signal strengths for two example neurons. For the purposes of illustration, the two task orientations are simply labeled positive and negative. This pair was typical in that the response functions (f_1 and f_2) are approximately linear over the range of signal strengths used. For this reason, we calculated the signal-induced correlation as the normalized product of the derivatives $f_1'f_2'$ [16]. **b**. The matrix of $f'f'$ values, as a function of task-aligned pairwise orientation preference, obtained using kernel smoothing as in Fig. 2. We observed a lattice-like pattern that was very similar to the structure of task-dependent r_{sc} we observed during task performance (Fig. 2e), suggesting task-dependent feedback introduces correlations that resemble the effect of stimulus changes relevant for the task. **c**. Scatter plot of the task-dependent (putatively top-down) component of r_{sc} (Fig. 3c) against $f'f'$ values for each recorded neuronal pair. The two were highly correlated across the population (r=0.62, p<0.001, bootstrap test, one-sided).

Figure 6. *R_{sc}* structure depends on variability in choice. a. The average, task-aligned r_{sc} matrix, shown

separately for each stimulus strength. The matrix associated with the 0% signal trials reflects dynamic

changes in r_{sc} structure across task contexts despite identical retinal input. A qualitatively similar structure

was apparent at non-zero signal levels. Spike counts were z-scored to eliminate the effect of stimulus

drive. **b.** We quantified the slope of a regression line comparing the r_{sc} values measured at each signal

level against the r_{sc} values measured at the 0% signal level. The magnitude of the slope indicates the

degree of attenuation of the r_{sc} structure at a given signal level. We observed a weak but significant

negative correlation ($p=0.038$, bootstrap test, one-sided) between this slope value and signal strength

(error bars are +/- 1 bootstrap SEM), suggesting the feedback variability generating the r_{sc} structure is

attenuated on high-signal trials, when there was also less variability in choice. Dotted line is fitted

regression line.

Figure 7. The task-dependent component of r_{sc} structure accounts for choice-related activity. a. Histogram of observed CPs, from the subset of neurons (n=138) significantly preferring one of the two task orientations (d'>0.9 at highest signal level). The mean of 0.54 was significantly above chance (p<0.001, bootstrap test using cell resampling, one-sided). CPs that were individually significant (p<0.05, bootstrap test using trial resampling, one-sided) are shown in black. **b.** To assess the possibility of a feedforward source of CP, we made use of the known analytical relationship between spike-count correlations, readout weights, and CPs, under the assumption of a linear decoder applied to a population of sensory neurons[17] (see Methods). CP is defined as a continuous function of task-aligned preferred orientation, analogous to our description of the r_{sc} matrix in Fig. 2e. The dashed black line shows the profile of CP observed across preferred orientations, after smoothing with a von Mises kernel approximating a wrapped Gaussian with 10° s.d. We applied a fixed sign convention to the CP values across all neurons, equivalent to arbitrarily calling the 0°-choice the preferred one. The predicted CP profiles (solid lines) show the CP elicited by reading out a sensory population with different r_{sc} structures. Readout weights across orientations were unobserved and the profiles shown are averages of a large set generated from different assumed readout weight profiles. **c.** Mean CP (using the traditional sign convention) associated with the profiles in (b), +/- 1 bootstrap SEM obtained by resampling from the data. Note that the mean observed CP is lower here than in (a) because all neurons are included, regardless of their orientation preference.

Figure 8. Temporal dynamics of r_{sc} structure. a. The average, task-aligned r_{sc} matrix obtained using spike

counts from 200-ms windows during the stimulus presentation. The task-aligned lattice structure was present at

all time points (4 examples shown). **b-c**. Plots showing the temporal dynamics of two statistical measures of the

observed r_{sc} structure (mean +/- 1 bootstrap SEM). The colored lines indicate the example time points shown in

(a). The population mean r_{sc} value (b) showed a sharp drop shortly after stimulus onset, as seen in other

studies[50], and then a gradual recovery over the course of the trial. The slope of the regression line of r_{sc}

obtained in each 200-ms window against r_{sc} obtained from trial-length spike counts is in (c). Apart from an

increased slope at the first time point, likely due to the onset of the visual stimulus, this showed no significant

modulation over the course of the trial. Values are all significantly less than 1 because smaller counting

windows introduced a source of uncorrelated noise across trials.

Methods

Electrophysiology

We recorded extracellular spiking activity from populations of V1 neurons in two male, awake, head-fixed rhesus monkeys (*Macaca mulatta*). For the majority of the recordings, monkey '*lem*' was 14 while monkey '*jbe*' was 16 years old, before which time they had each experienced extensive behavioral training, including on other behavioral paradigms for monkey '*lem*'. Monkey *'lem'* could not be pair housed due to antisocial behavior. Both monkeys were implanted with a head post and scleral search coils under general anaesthesia[51]. In monkey '*lem*', a recording chamber was implanted over a craniotomy above the right occipital operculum, as described previously[52], by which we introduced linear microelectrode arrays (U- and V-probes, Plexon; 24-contacts, 50 or 60 µm spacing) at an angle approximately perpendicular to the cortical surface with a custom micro-drive. We positioned the linear arrays so that we roughly spanned the cortical sheet, as confirmed with current-source density analysis, and removed them after each recording session. In monkey '*jbe*', a planar "Utah" array (Blackrock Microsystems; 96 electrodes 1mm in length inserted to target supragranular layers, 400 um spacing) was chronically implanted, also over the right occipital operculum. All procedures were performed in accordance with the U.S. Public Health Service Policy on the humane care and use of laboratory animals and all protocols were approved by the National Eye Institute Animal Care and Use Committee.

Broadband signals were digitized at 30 or 40 kHz and stored to disk. Spike sorting was performed offline using custom software in MATLAB®. First, spikes were detected using a voltage threshold applied to high-pass filtered signals. Next, triggered waveforms were projected into spaces defined either by principal components or similarity to a template. Clusters boundaries were finally estimated with a Gaussian mixture model, and then rigorously verified and adjusted by hand when needed. In the linear array recordings, spike sorting yield and quality was substantially improved by treating sets of three or four neighboring contacts as "n-trodes". As this was not possible with the Utah array due to the greater interelectrode spacing, we excluded pairs of neurons recorded on the same electrode to avoid

contamination by misclassification. Neurons from separate recording sessions were treated as

independent. To reduce the possibility that a single neuron from the Utah array contributed to two

datasets, we included only sessions that were separated by at least 48 hours (with a median separation of 5

days). We excluded from analysis those neurons whose mean evoked firing rate did not exceed 7

spikes/second.

Visual stimuli

All stimuli were presented binocularly on two gamma-corrected cathode ray tube (CRT) monitors

viewed through a mirror haploscope, at 85 or 100Hz. The monitors subtended 24.1° x 19.3° of visual

angle (1280 x 1024 pixels). The stimuli presented during performance of the discrimination task consisted

of bandpass filtered dynamic white noise, as described previously[32]. Briefly, stimuli were filtered in the

Fourier domain with a polar-separable Gaussian. The peak spatial frequency was optimized for the

recorded neuronal population (1 and 4 cpd medians for *'lem'* and '*jbe*', respectively) while the peak

orientation could take one of two orthogonal values the animal had to discriminate in a given session. The

angular s.d. of the filter modulated the orientation bandwidth and was varied trial to trial. A 2D Gaussian

contrast envelope was applied to the stimulus so that its spatial extent was as small as possible while still

covering the minimum response fields of the neuronal populations being recorded. The median envelope

s.d. was 0.6 degrees for both animals. The median stimulus eccentricity was 5.4 degrees for *'lem'* and 0.5

degrees for '*jbe*'. In Fig. 1, we quantify orientation bandwidth as % signal strength. This was calculated

as $100 * R$, where R is the length of the resultant vector associated with the angular component of the

stimulus filter.

We estimated neuronal orientation preferences in separate blocks of trials, using 420-ms

presentations of the following types of stimuli, presented at a range of orientations: 1) an orientation

narrowband version of the stimulus described above (10° angular s.d.); 2) sinusoidal gratings; and 3)

circular patches of dynamic 1D noise patterns (random lines). The preferred orientation of a neuron was

calculated as the circular mean of its orientation tuning curve. For each neuron, from among the set of

tuning curves elicited by the different stimulus types described above, we chose as the final estimate of

preferred orientation the one with the smallest standard error, obtained by resampling trials. We excluded

from further analysis all neurons where this exceeded 5°. On a subset of sessions, we also used these

orientation-tuning blocks to present examples of the 0%-signal orientation-filtered noise stimuli. These

were presented at the same location and size as during task performance, allowing us to calculate r_{sc}

structure in the absence of task engagement but with identical retinal input.

Orthogonal orientation discrimination task

The animals performed a coarse orientation discrimination task using the orientation-filtered noise

stimuli, as described previously[32]. To initiate a trial, the subject had to acquire a central fixation square.

After a delay of 50 ms, the stimulus appeared for a fixed duration of 2 seconds. The trial was aborted if

the subject broke fixation at any point during the stimulus presentation, defined as either 1) making a

microsaccade covering a distance greater than a predefined threshold (typically 0.5°) or 2) a deviation in

mean eye position from the center of the fixation point of more than a predefined threshold, typically 0.7°.

At the end of the stimulus presentation, two choice targets appeared. These were Gabor patches of 2-3° in

spatial extent, oriented at each of the two discriminandum orientations. The locations of the choice targets

depended on the task. For orientation pairs near horizontal and vertical (-22.5° – +22.5° and 67.5° –

112.5°), the choice targets were positioned along the vertical meridian, at an eccentricity of about 3°, with

the more vertically-oriented target appearing always in the upper hemifield. For orientation pairs near the

obliques (22.5° – 67.5° and 112.5° – 157.5°), the choice targets were positioned along the horizontal

meridian, at the same range of eccentricities, with the smaller of the two orientations always appearing in

the left hemifield. (We use the convention that horizontal is 0° and that orientation increases with

clockwise rotation.) To penalize random guessing, the volume of liquid reward delivered after correct

choices was doubled with each consecutive correct choice, up to a maximum of four times the initial

amount. Since we were primarily interested in the effect of task engagement on neuronal activity, we

applied a behavioral criterion to our data, excluding sessions where the subject's psychophysical threshold

(defined as the signal level eliciting 75% correct performance) exceeded 14% signal. A two-pass

presentation procedure was used. Each instance of a stimulus (generated with a given noise seed) was

shown twice per experimental block. This allowed us to account for any effect of fluctuations in the

stimulus on r_{sc} (see Supplementary Discussion §1.1 and Supplementary Fig. 5).

Spike-count correlation measurements

Spike-count correlations were calculated as the Pearson correlation between spike counts, counted

over the entire duration of the stimulus, with a 50-ms delay to account for the typical V1 response latency.

Spike counts were first z-scored separately within each experimental block (typically a set of 100-200

trials lasting about 10 minutes) and each stimulus condition. This removed correlations related to long-

term firing rate nonstationarities and allowed us to combine trials at different signal levels without

introducing correlations related to similarity in stimulus preference. We used a balanced z-scoring method

proposed recently to prevent bias related to differences in choice distributions across signal levels[53]. We

excluded pairs that were not simultaneously isolated for at least 25 trials total. The median number of

trials per pair during task performance was 752.

A main goal of the study was to measure how spike-count correlation varies with pairwise

orientation. We illustrate this dependence in several figures as a smoothed function estimated from

measures of r_{sc} combined across multiple recording sessions, which we then sampled discretely with $1°$

resolution. The smoothed estimates were obtained using a bivariate von Mises smoothing kernel. A point

in the correlation matrix $\mathbf{C}$ was given as:

$$\mathbf{C}(x,y) = \frac{\sum_{i=1}^{n} z_i K(x,y,\theta_i,\phi_i)}{\sum_{i=1}^{n} K(x,y,\theta_i,\phi_i)} \text{, where } K(x,y,\theta_i,\phi_i) = e^{\kappa\,(\cos(\theta_i - x) + \cos(\phi_i - y))}, \tag{1}$$

z_i is the i^{th} r_{sc} measurement, θ_i and ϕ_i are the preferred orientations of the i^{th} pair, and κ is the von Mises

width parameter. We set $\kappa = 1.3\pi$, yielding a smoothing kernel closely approximating a bivariate

wrapped Gaussian with 15° s.d. (Note that this smoothing procedure was only performed to generate

figures in the manuscript, and was not applied as a pre-processing step in any of the quantitative analyses.)

In some cases, we expressed the r_{sc} matrix in a task-aligned coordinate frame (e.g. Fig. 2e), for which the

preferred orientations of the i^{th} pair relative to the task orientations were used for θ_i and ϕ_i. Since there

were always two orthogonal task orientations, we averaged across both possible alignments, such that

$\mathbf{C}(x, y) = \mathbf{C}(x + 90°, y + 90°)$. All angular quantities were doubled for the calculations, as orientation

has a period of 180°. To generate the kernel-smoothed profile of CP (Fig. 7), we used a one-dimensional

equivalent of the procedure above, in which preferred orientations were parameterized only by a single

parameter.

Regression model

We used a multilinear regression model to identify fixed and task-dependent components of the

structured correlations we observed. We describe the set of observations (811 individual pairwise r_{sc}

measurements) in terms of a set of two underlying correlation structures: one defining r_{sc} as a function of

pairwise preferred orientation alone ("fixed") and the other defining r_{sc} as a function of pairwise preferred

orientation relative to the task orientations ("task-dependent"). In order to provide a continuous and

smooth description of the data, each component was parameterized as the sum of an array of $n \ x \ n$ evenly

spaced basis functions. Each observation, y_i, was expressed as:

$$y_i = x_i^{fixed} \cdot \beta^{fixed} + x_i^{task} \cdot \beta^{task} + c + \varepsilon_i \tag{2}$$

x_i^{fixed} and x_i^{task} are length-n^2 vectors of loadings onto the basis functions, which were given by

evaluating the basis functions at the location corresponding to the pairwise orientation preference of the i^{th}

pair. β^{fixed} and β^{task} are the length-n^2 vectors of amplitudes of the basis functions (coefficients to be fit),

c is a model constant, and $\cdot$ is the element-wise product. For the basis functions, we used bivariate von

Mises functions, with no correlation and equal width in both dimensions. Thus the k^{th} loading ($x_i^{fixed}(k)$

or $x_i^{task}(k)$) was given by:

$$x_i(k) = \frac{e^{\kappa\left(\cos\left(\theta_i - \mu_k^1\right) + \cos\left(\phi_i - \mu_k^2\right)\right)}}{Z} \qquad (3)$$

where θ_i and ϕ_i are the preferred orientations of the i^{th} pair (relative to the task orientations in the case of

the task-dependent loadings), μ_k is a pair of orientations defining the location of the k^{th} basis function, Z is

a normalization constant such that the sum of all loadings for observation i ($x_i^{fixed} + x_i^{task}$) is 1, and κ is

the basis function width. Again, angular quantities were doubled. κ acts as a smoothing hyperparameter.

We found that arrays of 8x8 were sufficient to describe the structure of the two components. Because the

observations were pairwise correlations, it was sufficient only to fit the upper triangular portion of the

array of basis functions. Thus, each component was described by 36 parameters (although the effective

number of parameters is significantly less because of the basis function smoothness and the ridge penalty).

We fit the model using ridge regression. The unique optimal solution could therefore be derived

analytically as $\hat{B} = (X^T X + \alpha I)^{-1} X^T Y$, where X is the concatenated design matrix combining x^{fixed} and

x^{task} and α is the ridge parameter, which penalized the squared amplitude of the basis functions. The

optimal values of the hyperparameters α and κ were chosen under 50-fold cross-validation.

Simple Parametric Models

We modeled the observed set of r_{sc} values using two simple parametric models: a "single

eigenvector" model and a "diagonal ridge" model. In the "single eigenvector" model, each observation y_i

was modeled as the outer product of an eigenvector X, evaluated at the relevant pair of orientations. The

eigenvector was parametrized as the difference of two von Mises functions separated by 90°:

$$X(\mu) = a_1 e^{\kappa_1 \cos(\mu+b)} - a_2 e^{\kappa_2 \cos(\mu+b+\pi)} \qquad (4)$$

632 where μ is the difference in preferred orientation and the task orientation (in angle-doubled radians), the

633 $a's$ are the amplitudes to be fit, the $\kappa's$ are the widths to be fit, and b is the offset of the eigenvector peak

634 and trough from the task orientations (allowing a mismatch between the model eigenvector and the task,

635 and also fit to the data). A given r_{sc} value y_i was given by:

$$y_i = X(\theta_i)X(\phi_i) + c + \varepsilon_i \tag{5}$$

where θ_i and ϕ_i are the task-aligned preferred orientation of the pair and c is a model constant. The model

contained six total free parameters which were fit using gradient descent to minimize the squared error in

the r_{sc} predictions.

In the "diagonal ridge" model, r_{sc} values were modeled as a decaying function of the difference in

preferred orientation, independent of task. The dependence was modeled as a von Mises function. A given

r_{sc} value y_i was modeled as:

$$y_i = a \cdot e^{\kappa \, \cos(\theta_i - \phi_i)} + c + \varepsilon_i \tag{6}$$

where θ_i and ϕ_i are the preferred orientation of the pair, c is a model constant, and a and κ parameterize

the von Mises function. The model contained three total free parameters which were fit using gradient

descent to minimize the squared error in the r_{sc} predictions.

Estimating explainable variance

While the above models did not explain more than a small percentage of the variance of the raw

observed r_{sc} values, this is not surprising as the raw correlation data do not vary smoothly with preferred

orientation (reflecting both noise, and the fact that r_{sc} is known to depend on parameters other than

orientation.[3,22,23]). For this reason, we measured goodness-of-fit relative to an estimate of the explainable

variance, which we took as the variance explained simply by a smoothed version of the raw data (sum of

values in fixed and task-aligned matrices was 3.6%). Smoothing was performed with a von Mises kernel,

with width chosen to maximize variance explained.

Choice probability predictions

Choice Probability was calculated in the standard way[36]. We only used 0%-signal trials, as the uneven choice distributions elicited by signal trials yield noisier CP measurements. Assuming feedforward pooling with linear readout weights, the relationship between the covariance matrix for a population of neurons, the readout weight of each neuron, and the Choice Probability (*CP*) of each neuron is:

$$CP_k = \frac{1}{2} + \frac{2}{\pi} \operatorname{sgn}(\xi_k) \arctan \sqrt{2\xi_k^{-2} - 1}^{-1} \quad \text{with} \quad \xi_k = \frac{(\mathbf{C}\beta)_k}{\sqrt{C_{kk}\beta^{\mathrm{T}}\mathbf{C}\beta}} \tag{7}$$

where CP_k is the CP of neuron k with respect to choice 1, β is the vector of readout weights and $\mathbf{C}$ is the covariance matrix[17]. We used this known relationship to quantify the CPs that would be associated with the r_{sc} structure we observed and the fixed and task-dependent components we identified, assuming only a feedforward source of CP (Fig. 7). CPs, r_{sc} structure, and readout weights were described as task-aligned functions of preferred orientation. This is equivalent to assuming a population of infinite size that is homogeneous at a given orientation. For the fixed component of r_{sc}, which was indexed relative to raw orientation preferences, we generated a task-aligned version by substituting the observed r_{sc} values with model fits (using only a fixed component of the model) and then generating a smoothed task-aligned matrix as in Fig. 2e, using these substituted values. To guarantee real-valued CPs on [0,1], we performed the calculations using a symmetric positive definite approximation[54] of the r_{sc} matrices, which introduced negligible error.

Since the readout weights were unknown, we generated a random distribution of 1000 plausible readout weight profiles that could support task performance. To generate a sample from this distribution, we started with a vector of random weights (drawn from a normal distribution) and applied the 90° symmetry inherent in the task, such that $\beta_\theta = -\beta_{\theta+90}$, where β_θ is the weight assigned to neurons with task-aligned preferred orientation θ. Then, we smoothed with a wrapped Gaussian kernel with 15° s.d. and excluded profiles which did not have a circular mean within 22.5° of choice 1 (0°). In practice, we found

the CP predictions to be insensitive to the readout weights (Supplementary Fig. 10), which is not

surprising for a nearly rank-1 matrix (since for exactly rank-1 matrices, the CPs are independent of the

weights)[17].

Estimating mean covariance for a population of neurons is necessarily more error-prone than

estimating mean correlation, as the former is sensitive to sampling error in measurements of average

spike-count variance (and therefore firing rate), so for this reason we chose to perform the calculations

using correlations (see Supplemental Discussion §2). We can use correlations interchangeably with

covariances in Eq. 7, under the simplifying assumption that the variance is uniform as a function of

preferred orientation. If Σ is the correlation matrix for a population with uniform variance α, then it

follows that:

$$\xi_k = \frac{a(\Sigma\beta)_k}{\sqrt{a\Sigma_{kk}\beta^{\mathrm{T}}(a\Sigma)\beta}} = \frac{(\Sigma\beta)_k}{\sqrt{\Sigma_{kk}\beta^{\mathrm{T}}\Sigma\beta}} \tag{8}$$

where $\Sigma_{kk} \equiv 1$ for all k. We felt that spike-count variance that depended systematically on preferred

orientation was unlikely to be a feature of the V1 representation, and thus that the advantages of using

correlations outweighed the cost.

Noise in the decision process after pooling (pooling noise) has the effect of uniformly scaling

down CPs, such that ξ_k in Eq. 7 is substituted with: $\frac{(\mathbf{C}\beta)_k}{\sqrt{C_{kk}(\beta^{\mathrm{T}}\mathbf{C}\beta + \sigma_{pool}^2)}}$, where σ_{pool}^2 is the variance of the

pooling noise[6]. We found that non-zero pooling noise was needed to avoid overestimating the magnitude

of CP from the observed correlation structure. We used a fixed value of pooling noise in our predictions

such that the average squared difference between the CP profile predicted from the observed correlation

matrix and the observed CP profile was minimized. Empirically, we found that pooling noise variance of

0.6 was optimal. Since our spike counts were normalized to have unit variance, this implies pooling noise

whose variance is 60% of the average spike-count variance of single neurons. This should be interpreted

with care, as overestimation of CPs may also be an artefact related to the assumption of a homogeneous

population[17]. Alternatively, the need to invoke pooling noise may be due to nonuniform sensory

integration across the trial, which is distinct but which would also have an attenuating effect on CP when

measured over the entire trial.

Statistics

Statistical tests were performed non-parametrically using bootstrapping or other resampling

methods, as described, with 1,000 resamples. Nonparametric statistical testing is superior when the

underlying distribution of the data cannot be assumed. When p-values of $p<0.001$ are reported, this

indicates the null hypothesis can be ruled out with the most confidence possible given the number of

resamples performed. In most cases, resampling was performed from the set of recorded neuronal pairs

(n=811), and always with replacement. In all figures, one asterisk indicates significant at the $p<0.05$ level,

two indicates $p<0.01$, and three indicates $p<0.001$.

Data availability

All relevant data are available upon reasonable request from the authors.

Code availability

All computer code used to generate the results are available upon request from the authors.

1. Tomko, G. J. & Crapper, D. R. Neuronal variability: non-stationary responses to identical visual
stimuli. *Brain Res.* **79,** 405–418 (1974).

2. Henry, G. H., Bishop, P. O., Tupper, R. M. & Dreher, B. Orientation specificity and response
variability of cells in the striate cortex. *Vision Res.* **13,** 1771–1779 (1973).

3. Cohen, M. R. & Kohn, A. Measuring and interpreting neuronal correlations. *Nat. Neurosci.* **14,**
811–9 (2011).

4. Shadlen, M. N. & Newsome, W. T. The variable discharge of cortical neurons: implications for
connectivity, computation, and information coding. *J. Neurosci.* **18,** 3870–3896 (1998).

5. Sompolinsky, H., Yoon, H., Kang, K. & Shamir, M. Population coding in neuronal systems with
correlated noise. *Phys. Rev. E* **64,** 51904 (2001).

6. Gu, Y., Angelaki, D. E. & DeAngelis, G. C. Contribution of correlated noise and selective
decoding to choice probability measurements in extrastriate visual cortex. *Elife* **3,** 1–19 (2014).

7. Liu, S., Dickman, J. D., Newlands, S. D., DeAngelis, G. C. & Angelaki, D. E. Reduced choice-
related activity and correlated noise accompany perceptual deficits following unilateral vestibular
lesion. *Proc. Natl. Acad. Sci. U. S. A.* **110,** 17999–8004 (2013).

8. Zohary, E., Shadlen, M. N. & Newsome, W. T. Correlated neuronal discharge rate and its
implications for psychophysical performance. *Nature* **370,** 140–143 (1994).

9. Abbott, L. F. & Dayan, P. The effect of correlated variability on the accuracy of a population code.
Neural Comput. **11,** 91–101 (1999).

10. Snippe, H. & Koenderink, J. Information in channel-coded systems: correlated receivers. *Biol.*
Cybern. **190,** 183–190 (1992).

11. Averbeck, B. B., Latham, P. E. & Pouget, A. Neural correlations, population coding and
computation. *Nat. Rev. Neurosci.* **7,** 358–66 (2006).

12. Cohen, M. R. & Maunsell, J. H. R. Attention improves performance primarily by reducing
interneuronal correlations. *Nat. Neurosci.* **12,** 1594–600 (2009).

13. Graf, A. B. a, Kohn, A., Jazayeri, M. & Movshon, J. A. Decoding the activity of neuronal
populations in macaque primary visual cortex. *Nat. Neurosci.* **14,** 239–45 (2011).

14. Mitchell, J. F., Sundberg, K. a & Reynolds, J. H. Spatial attention decorrelates intrinsic activity
fluctuations in macaque area V4. *Neuron* **63,** 879–88 (2009).

15. Johnson, K. O. Sensory discrimination: decision process. *J. Neurophysiol.* **43,** 1771–1792 (1980).

16. Moreno-Bote, R. *et al.* Information-limiting correlations. *Nat. Neurosci.* **17,** 1410–1417 (2014).

17. Haefner, R. M., Gerwinn, S., Macke, J. H. & Bethge, M. Inferring decoding strategies from choice
probabilities in the presence of correlated variability. *Nat. Neurosci.* **16,** 235–42 (2013).

18. Nienborg, H. & Cumming, B. G. Correlations between the activity of sensory neurons and
behavior: how much do they tell us about a neuron's causality? *Curr. Opin. Neurobiol.* **20,** 376–
381 (2010).

19. Shadlen, M. N., Britten, K. H., Newsome, W. T. & Movshon, J. A. A computational analysis of the
relationship between neuronal and behavioral responses to visual motion. *J. Neurosci.* **16,** 1486–
510 (1996).

20.   Bair, W., Zohary, E. & Newsome, W. T. Correlated firing in macaque visual area MT: time scales
757   and relationship to behavior. *J. Neurosci.* **21,** 1676–97 (2001).

758   21.   Lee, D., Port, N. L., Kruse, W. & Georgopoulos, a P. Variability and correlated noise in the
759   discharge of neurons in motor and parietal areas of the primate cortex. *J. Neurosci.* **18,** 1161–70
760   (1998).

761   22.   Smith, M. A. & Kohn, A. Spatial and temporal scales of neuronal correlation in primary visual
762   cortex. *J. Neurosci.* **28,** 12591–603 (2008).

763   23.   Kohn, A. & Smith, M. a. Stimulus dependence of neuronal correlation in primary visual cortex of
764   the macaque. *J. Neurosci.* **25,** 3661–73 (2005).

765   24.   Callaway, E. M. Feedforward, feedback and inhibitory connections in primate visual cortex. *Neural
766   Netw.* **17,** 625–32 (2004).

767   25.   Sillito, A. M., Cudeiro, J. & Jones, H. E. Always returning: feedback and sensory processing in
768   visual cortex and thalamus. *Trends Neurosci.* **29,** 307–16 (2006).

769   26.   Ruff, D. A. & Cohen, M. R. Attention can either increase or decrease spike count correlations in
770   visual cortex. *Nat. Neurosci.* **17,** 1591–7 (2014).

771   27.   Cohen, M. R. & Newsome, W. T. Context-dependent changes in functional circuitry in visual area
772   MT. *Neuron* **60,** 162–73 (2008).

773   28.   Mitchell, J. F., Sundberg, K. a & Reynolds, J. H. Differential attention-dependent response
774   modulation across cell classes in macaque visual area V4. *Neuron* **55,** 131–41 (2007).

775   29.   Lange, R. D. & Haefner, R. M. Inferring the brain's internal model from sensory responses
776   in a probabilistic inference framework. *bioRxiv* (2016).

777   30.   Haefner, R. M., Berkes, P. & Fiser, J. Perceptual Decision-Making as Probabilistic Inference by
778   Neural Sampling. *Neuron* **90,** 649–660 (2016).

779   31.   Cumming, B. G. & Nienborg, H. Feedforward and feedback sources of choice probability in neural
780   population responses. *Curr. Opin. Neurobiol.* **37,** 126–132 (2016).

781   32.   Nienborg, H. & Cumming, B. G. Decision-Related Activity in Sensory Neurons May Depend on
782   the Columnar Architecture of Cerebral Cortex. *J. Neurosci.* **34,** 3579–3585 (2014).

783   33.   Nienborg, H. & Cumming, B. G. Psychophysically measured task strategy for disparity
784   discrimination is reflected in V2 neurons. *Nat. Neurosci.* **10,** 1608–1614 (2007).

785   34.   Ahumada Jr, A. J. Perceptual classification images from Vernier acuity masked by noise. in
786   *Perception ECVP abstract* **25,** 0 (Pion Ltd, 1996).

787   35.   Ecker, A. S., Denfield, G. H., Bethge, M. & Tolias, A. S. On the Structure of Neuronal Population
788   Activity under Fluctuations in Attentional State. *J. Neurosci.* **36,** 1775–1789 (2016).

789   36.   Britten, K. H., Newsome, W. T., Shadlen, M. N., Celebrini, S. & Movshon, J. A. A relationship
790   between behavioral choice and the visual responses of neurons in macaque MT. *Vis. Neurosci.* **13,**
791   87 (1996).

792   37.   Crapse, T. B. & Basso, M. A. Insights into Decision-Making Using Choice Probability. *J.
793   Neurophysiol.* jn.00335.2015 (2015). doi:10.1152/jn.00335.2015

38.   Nienborg, H., Cohen, M. R. & Cumming, B. G. Decision-Related Activity in Sensory Neurons:

Correlations Among Neurons and with Behavior. *Annu. Rev. Neurosci.* **35,** 463–483 (2012).

39.   Crapse, T. B. & Basso, M. A. Insights into decision making using choice probability. *J.*
Neurophysiol. **114,** 3039–3049 (2015).

40.   Nienborg, H. & Cumming, B. G. Decision-related activity in sensory neurons reflects more than a
neuron's causal effect. *Nature* **459,** 89–92 (2009).

41.   Wimmer, K. *et al.* Sensory integration dynamics in a hierarchical network explains choice
probabilities in cortical area MT. *Nat. Commun.* **6,** 6177 (2015).

42.   Kwon, S. E., Yang, H., Minamisawa, G. & O'Connor, D. H. Sensory and decision-related activity
propagate in a cortical feedback loop during touch perception. *Nat. Neurosci.* **19,** (2016).

43.   Treue, S. & Martínez Trujillo, J. C. Feature-based attention influences motion processing gain in
macaque visual cortex. *Nature* **399,** 575–9 (1999).

44.   James, W. *The Principles of Psychology*. (H Holt, 1890).

45.   Posner, M. I. & Petersen, S. E. The attention system of the human brain. *Annu. Rev. Neurosci.* 25–
42 (1990).

46.   Orbán, G., Berkes, P., Fiser, J. & Lengyel, M. Neural Variability and Sampling-Based Probabilistic
Representations in the Visual Cortex. *Neuron* **92,** 530–543 (2016).

47.   Tajima, C. I. *et al.* Population Code Dynamics in Categorical Perception. *Sci. Rep.* **6,** 22536 (2016).

48.   Knill, D. C. & Richards, W. *Perception as Bayesian Inference*. (Cambridge University Press,
1996).

49.   Von Helmholtz, H. *Handbuch der physiologischen Optik*. **9,** (Voss, 1867).

50.   Churchland, M. M. *et al.* Stimulus onset quenches neural variability: a widespread cortical
phenomenon. *Nat. Neurosci.* **13,** 369–78 (2010).

51.   Judge, S. J., Richmond, B. J. & Chu, F. C. Implantation of magnetic search coils for measurement
of eye position: An improved method. *Vision Res.* **20,** 535–538 (1980).

52.   Cumming, B. G. & Parker, A. J. Binocular neurons in V1 of awake monkeys are selective for
absolute, not relative, disparity. *J. Neurosci.* **19,** 5602–18 (1999).

53.   Kang, I. & Maunsell, J. H. R. Potential confounds in estimating trial-to-trial correlations between
neuronal response and behavior using choice probabilities. *J. Neurophysiol.* **108,** 3403–15 (2012).

54.   Higham, N. J. Computing a nearest symmetric positive semidefinite matrix. *Linear Algebra Appl.*
103, 103–118 (1988).

Acknowledgements

We thank Bob Wurtz and James McFarland for useful discussions; Richard Krauzlis, Bevil Conway, and Ali Ghazizadeh for comments on an earlier version of the manuscript; and Beth Nagy, Irina Bunea, and Denise Parker for veterinary care.

Author Contribution

A.G.B. and B.G.C. conceived and designed the experiments. A.G.B. performed the experiments and all aspects of the analysis. A.G.B. and B.G.C. wrote the paper. R.H. advised and assisted with the data analysis and the paper. B.G.C. advised at all stages.

Competing Financial Interests

The authors declare no competing financial interests.