Title

Multi-omic analysis of urothelial cancer patients treated with PD-L1

blockade demonstrates the contribution of both systemic and somatic

factors to the biology of response and resistance

Authors

Alexandra Snyder[1]*, Tavi Nathanson[2]*, Samuel Funt[1]*, Arun Ahuja[2], Jacqueline Buros Novik[2], Matthew D.

Hellmann[1], Eliza Chang[2], Bulent Arman Aksoy[2], Hikmat Al-Ahmadie[3], Erik Yusko[4], Marissa Vignali[4], Sharon

Benzeno[4], Mariel Boyd[1], Meredith Moran[1], Gopa Iyer[1], Harlan S. Robins[4,5], Elaine R. Mardis[6], Taha Merghoub[1],

Jeff Hammerbacher[2]^, Jonathan Rosenberg[1]^, Dean Bajorin[1]^

*These authors contributed equally.

^These authors are co-corresponding.

Affiliations:

[1]Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY; Department of Medicine,

Weill Cornell Medical College, New York, NY 10065

[2]Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY

10029

[3]Department of Pathology, Memorial Sloan Kettering Cancer Center, New York, NY 10065

[4]Adaptive Biotechnologies, Seattle, WA 98102

[5]Fred Hutchinson Cancer Research Center, Seattle, Washington 98109

[6]Institute for Genomic Medicine; The Research Institute at Nationwide Children's Hospital; The Ohio State

University College of Medicine; Columbus, OH 43205

The authors declare no potential conflicts of interest.

Abstract

Background: Inhibition of programmed death-ligand one (PD-L1) with atezolizumab can induce durable clinical benefit (DCB) in patients with metastatic urothelial cancers, including complete remissions in patients with chemotherapy refractory disease. Although mutation load and PD-L1 immune cell (IC) staining have been associated with response, they lack sufficient sensitivity and specificity for clinical use. Thus, there is a need to evaluate the peripheral blood immune environment and to conduct detailed analyses of mutation load, predicted neoantigens and immune cellular infiltration in tumors to enhance our understanding of the biologic underpinnings of response and resistance.

Methods and Findings: We performed whole exome sequencing (WES), RNA sequencing (RNA-seq), and T cell receptor sequencing (TCR-seq) of pre-treatment tumor samples as well as TCR sequencing of matched, serially collected peripheral blood pre- and post-treatment with atezolizumab. These parameters were assessed for correlation with DCB (defined as progression free survival (PFS) > 6 months) and overall survival (OS), both alone and in the context of clinical and intratumoral parameters known to be predictive of survival in this disease state.

Patients with DCB displayed a higher proportion of tumor infiltrating T lymphocytes (TIL) (n=24, Mann-Whitney p=0.047). Pre-treatment peripheral blood TCR clonality below the median was associated with improved PFS (n=29, log-rank p=0.048) and OS (n=29, log-rank p=0.011). Patients with DCB also demonstrated more substantial expansion of tumor-associated TCR clones in the peripheral blood 3 weeks after starting treatment (n=22, Mann-Whitney p=0.022). The combination of high pre-treatment peripheral blood TCR clonality with elevated PD-L1 IC

staining in tumor tissue was strongly associated with poor clinical outcomes (n=10, HR=86.22,

95% CI (2.55, 491.65)). Marked variations in mutation loads were seen with different somatic

variant calling methodologies, which in turn impacted associations with clinical outcomes.

Missense mutation load, predicted neoantigen load and expressed neoantigen load did not

demonstrate significant association with DCB (n=25, Mann-Whitney p=0.22, n=25, Mann-

Whitney p=0.55, and n=25, Mann-Whitney p=0.29 respectively). Instead, we found evidence of

time-varying effects of somatic mutation load on progression-free survival in this cohort (n=25,

p=0.044).

Conclusions: These results demonstrate the complex nature of immune response to checkpoint

blockade and the compelling need for greater interrogation and data integration of both host and

tumor factors. Incorporating these variables in prospective studies will facilitate identification and

treatment of resistant patients.

64   Introduction

Atezolizumab has demonstrated responses in 15-25% of patients with advanced urothelial

carcinoma and improved survival compared to historical expectations (1,2). Similar to predictive

factor analyses in melanoma, colon cancer and non-small cell lung cancer studies with other

checkpoint blockade agents, Rosenberg and colleagues reported a statistically significant

association between mutation load and response to atezolizumab in urothelial cancer patients

(2). However, mutation load in the atezolizumab study was predicted based on an estimate

using a targeted panel and not with WES. Similar to findings from prior studies, the association

between this predicted mutation load and outcomes in patients with urothelial cancer was not

dichotomous; there were tumors from patients with elevated mutation load that did not respond

to therapy, and vice versa. Additionally, positive PD-L1 staining of infiltrating immune cells by

immunohistochemistry was associated with, but poorly predicted, response. A statistical model

suggested that both PD-L1 staining and mutation load impacted the likelihood of response.

However, the authors did not recommend its clinical use.

Collectively, studies to date imply that a combination of immune parameters are

necessary to gain further precision in determining the likelihood of benefit from these

immunotherapies and that a single biologic marker will be insufficient. There have been few

attempts to integrate molecular and immunologic data from patients treated with checkpoint

blockade and their tumors. Consequently we performed whole exome (WES), RNA, and T cell

receptor (TCR) sequencing of tumor samples from patients treated with atezolizumab as well as

TCR sequencing of matched, serially collected peripheral blood.

In this cohort of patients, we illustrate the importance of host immune factors, including

intratumoral and peripheral T cell receptor clonality, infiltration and expansion, to clinical

outcomes. We did not find a significant association between mutation or expressed neoantigen

load and progression free survival or DCB (defined as progression free survival (PFS) > 6

months). However, we did demonstrate a time-dependent relationship between mutation load

and outcome, implyling that patients who experienced rapid progression displayed systemic

indicators of immune deficiency despite elevated mutation load in the tumors. Calculation of the

hazard ratios for each measured biomarker and clinical factor underscores the concept that a

complex interaction of both host and tumor variables determines whether a patient will

experience clinical benefit from anti-PD-L1 therapy.

Methods

<u>Patients and clinical characteristics</u>

All patients had locally advanced or metastatic urothelial carcinoma and were treated at

Memorial Sloan Kettering Cancer Center (n=29) on protocol NCT02108652 (2). All patients

initiated therapy in 2014 and were treated with atezolizumab 1200 mg IV every 3 weeks, and

had consented to Institutional Review Board-approved protocols permitting tissue and blood

collection and sequencing. Patient tumor samples were assessed prospectively and centrally

(by HistoGeneX, Brussels, Belgium) for PD-L1 expression by immunohistochemistry with the

SP142 assay (Ventana, AZ, USA) (1). The PD-L1 tumor-infiltrating immune cell (IC) status was

defined by the percentage of PD-L1-positive immune cells in the tumor microenvironment: IC0

(<1%), IC1 (≥1% but <5%), and IC2/3 (≥5%) as defined in the original study. Smoking status

was evaluated using previously completed self-reported smoking questionnaires or review of

medical records.

<u>Tumor and blood samples</u>

All tumor tissue used for sequencing was obtained prior to dosing with atezolizumab. Tumor

samples used for whole exome sequencing were all formalin-fixed paraffin embedded (FFPE).

The presence of tumor tissue in the sequenced samples was confirmed by examination of a

representative hematoxylin and eosin-stained slide by a genitourinary pathologist (H.A.).

Peripheral blood mononuclear cells (PBMCs) were isolated and stored as previously described

(3). PBMC were collected pre-treatment and during treatment.

Clinical efficacy analysis

Tumor responses to atezolizumab were evaluated by CT scan every 9 weeks for the first 12

months following day 1 of cycle 1. After 12 months, tumor assessments were performed every

12 weeks. The response evaluation criteria in solid tumors (RECIST) version 1.1 was used to

define objective clinical responses by the institutional radiologist.

DNA extraction and high-throughput TCRβ Sequencing

Genomic DNA was purified from total PBMCs and tumor samples using the Qiagen DNeasy

Blood extraction kit. The $TCR\beta$ CDR3 regions were amplified and sequenced using immunoSEQ

(Adaptive Biotechnologies, Seattle, WA) as previously described (4). In brief, bias-controlled V

and J gene primers were used to amplify rearranged V(D)J segments for high-throughput

sequencing at ~20X coverage. After correcting sequencing errors via a clustering algorithm,

CDR3 segments were annotated according to the International ImMunoGeneTics Collaboration

(5) to identify the V, D, and J genes that contributed to each rearrangement. A mixture of

synthetic TCR analogs in each PCR was used to estimate the absolute template abundance

(i.e., the number of cells bearing each unique TCR sequence) from sequencing data, as

previously described (6). The estimated TIL content was calculated as previously described (6–

8). To determine TIL content in FFPE samples as a T cell fraction, we amplified several

housekeeping genes and quantitated their template counts to determine the amount of DNA

usable for TCRB sequencing. ImmunoSEQ then amplifies and sequences the molecules with

rearranged TCRb chains. Because the immunoSEQ assay aligns sequences to the the IMGT

database, sequences are annotated as complete VDJ rearrangements or non-productive

rearrangements (a stop codon or out of frame CDR3 region was generated during VDJ

recombination in one of the alleles; all downstream analysis in this work proceeded with

complete, productive sequences. To estimate the number of starting templates that were in the

sample, the number of sequence reads for each TCRB sequences is measured. Synthetic

control templates were also spiked into each sample, thereby enabling quantitation of input

TCRB templates from the read counts. To determine the proportion of T cells in the FFPE

samples, we amplified several housekeeping genes and quantitated their template counts to

determine the amount of DNA usable for TCRB sequencing. For each sample, Shannon entropy

was also calculated on the clonal abundance of all productive TCR sequences in the data set.

Shannon entropy was normalized to the range by dividing Shannon entropy by the logarithm of

the number of unique productive TCR sequences in the data set. This normalized entropy value

was then inverted (12 normalized entropy) to produce the clonality metric. Those T cell clones

whose frequencies differed between samples from a given subject taken at different time points,

or between cell populations (e.g., between total PBMCs and tumor) were computationally

identified as previously described (9). The input data consisted of the absolute abundance for

each TCR clone in each sample. Fisher's exact test was used to compute a p-value for each

clone across the two samples, against the null hypothesis that the population abundance of the

clone is identical in the two samples. We corrected for multiple testing to control FDR using the

Benjamini-Hochberg procedure and employed a significance threshold of 0.01 on adjusted p-

values.

<u>Whole Exome Sequencing</u>

Twenty-six FFPE-derived tumor and frozen PBMC-derived normal paired samples were

sequenced by exome hybrid capture using the IDT xGen Whole Exome Panel

(https://www.idtdna.com/pages/products/nextgen/target-capture/xgen-lockdown-panels/xgen-

exome-panel) and standard protocols. Briefly, each sample was used to create a barcoded

Illumina library, tumor samples were pooled at optimal multiplex to create an equimolar pool into

the hybrid capture reaction, which was performed according to the manufacturer's suggested

protocol. Similarly, normal samples were pooled and introduced to the hybrid capture reaction.

Following the recovery of captured library fragments, PCR amplification was performed, the

resulting pools of fragments were quantitated using qPCR (Kapa Bio), and sequenced in

separate lanes by paired end 150 bp reads, using the Illumina HiSeq 4000.

Somatic Variant Calling

DNA sequencing data for the tumor and normal samples were aligned to the GRCh37 reference

using bwa-mem (v. 0.7.10) with default settings. The resulting BAMs were processed through

Picard MarkDuplicates and the GATK (v. 3.5-0) pipeline including Base Quality Score

Recalibration and Indel Realignment. Single nucleotide variants were called from Mutect (v.

1.1.6) and Strelka (v. 1.0.14) with default settings. Variants from either call were included and

the variants calls were further filtered to those with depth (in normal and tumor samples) ≥ 7

reads, > 10% tumor VAF and ≤ 3% normal VAF (10). Mutations per megabase was computed

by normalizing the number of mutations by the number of loci with ≥ 7 reads in normal and

tumor samples.

Variants were annotated as missense variants by Varcode (v. 0.5.10,

https://github.com/hammerlab/varcode) and PyEnsembl (v. 1.0.3,

https://github.com/hammerlab/pyensembl) using Ensembl Release 75 and annotated as

deleterious using PolyPhen (v. 2.2.2).

Mutational signatures were inferred from the somatic mutation calls using *deconstructSigs*.

<u>HLA typing</u>

HLA types for each patient were computed from the normal sequencing data using OptiType (v.

1.0.0).

<u>RNA-seq</u>

RNA was extracted from twenty-six FFPE tumor samples and evaluated for quality and quantity

using the Agilent RNA pico chip. Each sample was prepared for sequencing by constructing an

Illumina Tru-Seq Stranded RNA kit, according to the manufacturer's protocol. The resulting

libraries were amplified by PCR, quantitated, pooled and processed through a hybrid capture

intermediate using the IDT xGen Exome reagent (as above). The captured fragments were

quantitated, diluted and were sequenced using 2 x 150 bp paired end reads on the Illumina

HiSeq 4000.

The RNA sequencing data were aligned to the GRCh37 reference an Ensembl Release 75

using STAR (v. 2.4.1d) and transcript quantification was performed using kallisto (v. 0.42.3).

The STAR alignment was only used for identifying variant-supporting reads in the RNA. For

gene-level analysis, the transcript quantifications were aggregated to the gene level using

tximport (http://f1000research.com/articles/4-1521/v1).

Expressed mutations and neoantigens were computed using Isovar (v. 0.0.6,

https://github.com/hammerlab/isovar). This examined the number of RNA reads containing the

mutation and the surrounding nucleotide context. Mutations (and neoantigens) were counted as

expressed when there were at least 3 reads containing the mutated base.

sleuth (v. 0.28.1) was used for differential expression analysis and GSEA was used for pathway

enrichment analysis. ESTIMATE was used to quantify immune and stromal scores from RNA-

seq data.

Neoantigen calling

Neoantigens were computed from all nonsynonymous mutations using Topiary (v. 0.1.0,

https://github.com/hammerlab/topiary) and NetMHCCons (v. 1.1) and the computed HLA alleles

from OptiType. As with expressed mutations, expressed neoantigens were those supported in

the RNA with at least 3 uniquely-mapped reads matching the cDNA sequence.

Statistical analysis

All statistical analysis was performed in Python and R (v. 3.3.1). Cohorts (v. 0.4.0,

https://github.com/hammerlab/cohorts) was used to orchestrate the analysis. The Mann Whitney

and Fisher's Exact test were performed using the Python scientific computing library, SciPy (v.

0.18.1). Kaplan-Meier curves were computed with Lifelines (v. 0.9.1.0). Survival and logistic

regression models were estimated using PyStan (v. 2.12.0.0) and the Stan statistical computing

software (v. 2.12.0). Survival analyses utilized a proportional hazards piecewise exponential

model with a random walk baseline hazard. The analysis for presence of a time-varying

covariate effect was performed in R using the *survival* package (v. 2.39.5), whereas the

estimation of the time-varying covariate effect was performed using Stan. This analysis

estimated the covariate effect at each timepoint with a random-walk prior. In some cases,

alternative specifications of models written in Stan were interrogated as sensitivity analyses; see

the github repo for details.

Results

Patient Characteristics

29 patients with metastatic urothelial cancer from a single institution treated with atezolizumab

as part of a single-arm phase II study (IMvigor 210, NCT 02108652) were included in the

analyses. The patients displayed characteristics typical of the metastatic urothelial cancer

population studied in IMvigor 210: a preponderance of males with urothelial cancers of bladder

origin, greater than half of whom had a reported prior smoking history (Table 1). Patients had an

ECOG performance status of 0 or 1, and had 0 to 3 prior regimens of chemotherapy. Of this

group, 25 patients had sufficient tumor tissue for WES, 26 for RNA-seq and 24 for TCR-seq. 29

had a pre-treatment peripheral blood sample on which TCR-Seq could be performed; 24 had

one pre-treatment and at least one post-treatment peripheral blood collection.

Table 1: Baseline characteristics		
Variable	Durable Clinical Benefit (n=9)	No Durable Benefit (n=20)
Median Age (range)	66 (57-74)	72 (46-81)
Male Sex	9 (100%)	16 (80%)
Site of primary tumor		
Bladder	7 (78%)	19 (95%)
Upper tract	2 (12%)	1 (5%)
ECOG performance status		

0	0 (0%)	1 (5%)
1	9 (100%)	19 (95%)
History of tobacco use		
No	4 (44%)	7 (35%)
Yes	5 (56%)	13 (65%)
Hemoglobin concentration < 10gm/dL	1 (11%)	3 (15%)
Albumin ≤ lower limit of normal	2 (22%)	5 (25%)
Metastatic sites at baseline		
Visceral^	3 (33%)	16 (80%)
Liver	2 (22%)	9 (45%)
Lymph node only	6 (67%)	4 (20%)
Number of previous systemic regimens in the metastatic setting		
0	5 (56%)	3 (20%)
1	2 (22%)	15 (70%)
2	1 (11%)	1 (5%)
≥ 3	1 (11%)	1 (5%)
Previous neoadjuvant or adjuvant chemotherapy, with first progression within ≤12 months	3 (33%)	2 (10%)
Time since previous chemotherapy ≤ 3 months	2 (29%)	5 (26%)
Prognostic risk group for previously treated patients*		
Low	3 (42%)	7 (37%)
Intermediate	2 (29%)	5 (26%)
High	2 (29%)	7 (37%)
Prognostic risk group for previously untreated patients†		

Low	0 (0%)	0 (0%)
Intermediate	2 (100%)	0 (0%)
High	0 (0%)	1 (100%)
Intravesical BCG administered (%)	2 (22%)	10 (50%)
^Visceral metastasis defined as liver, lung, bone, or any non-lymph node or soft tissue metastasis.*Based on (11) †Based on (12)		

Intratumoral and peripheral TCR features associate with durable clinical benefit

The importance of T cells to the anti-tumor response has long been known (13); the relevance of intratumoral and peripheral T cell receptor (TCR) clonality to the anti-tumor response is an area of active study. A single previous study of melanoma patients treated with anti-PD-1 therapy demonstrated that patients whose tumors featured both high levels of tumor infiltrating T lymphocytes (TIL) along with high TIL clonality were more likely to experience radiographic response to therapy (8). A separate study examined the peripheral TCR repertoire in anti-CTLA-4-treated patients with prostate cancer or melanoma, and found that clonotype stability was associated with response (14). To our knowledge, no prior study has reported both intratumoral and peripheral TCR clonality in a single population treated with checkpoint blockade therapy.

We performed TCR sequencing of tumors and peripheral blood mononuclear cells (PBMC) at serial time points in our cohort. Due to limitations in sample availability, this analysis included tumors from 24 patients and peripheral blood from 29 patients, including pre-treatment samples in all patients, and up to 8 total time points. A median of 141,255 (range 43,052-335,089) T cells were analyzed per peripheral blood sample, including 82,636 (range 24,095-

207,860) unique TCRs, with a median clonality of 0.080 (range 0.014-0.37) and a median T cell

proportion of 0.31 (range 0.082-0.64). In the tumors, the corresponding values included 1,402

(range 63-133,167) T cells, 1,086 (range 67-56,273) unique TCRs, clonality of 0.096 (range

0.033-0.34) and T cell proportion of 0.097 (range 0.0098-0.33).

In our patient group, we first asked whether there was an association between outcome

and either tumor TCR clonality or TIL proportion, or with TCR clonality in the peripheral blood.

Consistent with the data from Tumeh and colleagues (8), tumors from patients who experienced

a DCB exhibited a higher TIL proportion and higher TCR clonality than those patients who

experienced progressive disease (Fig 1A, Supplementary Fig S1A). Importantly, TIL proportion

alone was also associated with DCB, with a median of 0.21 (range 0.049-0.33) in tumors from

patients who had PFS greater than 6 months, versus 0.069 (range 0.0098-0.24) in tumors from

patients who did not (n=24, Mann-Whitney p=0.047, Fig 1B); it remains unclear whether TIL

clonality adds to TIL proportion in its association with DCB in this study (TIL proportion and

clonality versus TIL proportion alone, n=24, log-likelihood p=0.100).

Fig 1. T-cell receptor clonality and treatment response.

(A) High-throughput quantitative sequencing of the rearranged TCR β genes using the

immunoSEQ assay was performed. The x-axis represents clonality of the T-cell repertoire.

Absence of DCB (red dots) was associated with lower levels of T-cell infiltration and T-cell

clonality than in tumors from patients with DCB (green dots). TIL proportion and TIL clonality

were not significantly more predictive of DCB than TIL proportion alone (n=24, log-likelihood

p=0.100). (B) TIL proportion alone was associated with DCB, with a median of 0.21 (range

0.049-0.33) in tumors from patients who had DCB, versus 0.069 (range 0.0098-0.24) in tumors

from patients who did not (n=24, Mann-Whitney p=0.047). (C) Patients with a pre-treatment

peripheral TCR clonality less than the median exhibited improved progression free survival

(n=29, log-rank p=0.048). (D) Patients with a pre-treatment, peripheral TCR clonality less than

the median exhibited improved overall survival (n=29, log-rank p=0.011). (E) There was a

significant association between TCR clonality in the peripheral blood prior to initiating treatment

and overall survival greater than 12 months (OS DCB, TCR clonality 0.060 (range 0.022-0.21);

OS less than 12 months: 0.15 (range 0.031-0.35) (n=29, Mann-Whitney p=0.0061). (F) There

was no significant association between pre-treatment peripheral TCR clonality and DCB (DCB,

TCR clonality 0.068 (range 0.027-0.21); no DCB: 0.14 (range 0.022-0.35) (n=29, Mann-Whitney

p=0.25). (G) Expansion of TCR clones found in the tumor infiltrating T lymphocytes (TIL, orange

bars) occurred in the peripheral blood 3 weeks after initiating treatment in all patients. (H) The

number of TCR clones found in TIL that expanded in the peripheral blood 3 weeks after initiating

treatment was 8.00 (range 4.00-12.00) in patients with DCB and 2.50 (range 1.00-18.00) in non-

DCB patients (n=22, Mann-Whitney p=0.022).

We next examined pre-treatment peripheral blood clonality and its relationship to DCB.

Because a diverse TCR repertoire in circulation may increase the likelihood that a tumor specific

T cell population is present, we hypothesized that T cell receptor clonality would be inversely

associated with response. Indeed, low pre-treatment peripheral TCR clonality associated with

improved PFS (n=29, log-rank p=0.048) and OS (n=29, log-rank p=0.011) (Figs 1C, 1D, 1E),

although not with DCB (Fig 1F).

Finally, we explored the relationship between intratumoral and peripheral TCR clonality.

Variations in individual T cell clones present in tumors can be tracked in the peripheral blood

during treatment (examples in Supplementary Fig S1B). Expansion of tumor-associated TCRs

occurred in the peripheral blood in all patients (Fig 1G). However, a more pronounced

expansion of intratumoral TCR clones was observed in DCB patients at three weeks after

initiation of treatment (second dose of therapy) (Fig 1H) that diminished by 6 weeks after

therapy initiation (Supplementary Fig S1C). Interestingly, all patients with low pre-treatment

peripheral TCR clonality and high TIL clonality survived greater than one year (Supplementary

Fig S1D).

Association of tumor genetic features with progression free or overall

survival

To further examine intratumoral factors associated with therapeutic efficacy, we performed WES

on 25 formalin fixed, paraffin-embedded (FFPE) archived tumor samples. Mean target coverage

was 129 (range 44-194) in tumors, and 73 (range 59-91) in normal tissue. Single nucleotide

variants were identified and annotated as silent, missense or nonsense mutations (Fig 2A).

There was no significant association between median missense mutation load and DCB

(median mutations per megabase 3.24 (range 0.038-11.46) in patients with DCB compared to

0.45 (range 0.019-9.90) in those without DCB, n=25, Mann-Whitney p=0.22, Fig 2B). There was

also no significant association between missense mutation load and overall survival greater

than 12 months (n=25, Mann-Whitney p=0.37, Supplementary Fig S2A). In a survival analysis

for time to disease progression or mortality, the estimated hazard ratio associated with increase

in missense SNV count per megabase was 0.92 (95% CI 0.78 - 1.09). These results are not

surprising given that the present sample size (n=25) is underpowered to detect an effect of

magnitude similar to that observed by Rosenberg and colleagues (2) (power=0.2 assuming

median of 12.4 vs 6.4 mutations per megabase among patients with DCB vs non-DCB

response).

Fig 2. Single nucleotide variants and treatment response.

(A) Single nucleotide variants, premature stop codons, transversions, mutations in start or stop

codons and splice site variants as well as transitions and transversions were called for all

samples. (B) Median mutations per megabase of 3.24 (range 0.038-11.46) in tumors from

patients who progressed at or after 6 months, as compared to 0.45 (range 0.019-9.90) in those

who progressed in less than 6 months (n=25, Mann-Whitney p=0.22). (C) Median expressed

neoantigens in tumors from patients who progressed at or after 6 months was 1.32 (range 0.00-

6.06), versus 0.29 (range 0.00-5.70) in those who progressed before 6 months (n=25, Mann-

Whitney p=0.29).

When filtering to expressed mutations, we found a median of 0.79 (range 0.00-3.36)

expressed mutations per megabase for patients with DCB and a median of 0.16 (range 0.00-

3.34) expressed mutations per megabase for patients without DCB (n=25, Mann-Whitney

p=0.26, Supplementary Fig S2B). Consistent with known importance of specific variant calling

pipelines to output (15,16), we found that different filtering techniques impacted the association

with DCB (Supplementary Table S1). Missense mutation load, when counting only mutations

that were removed after post-processing (via Base Quality Score Recalibration (BQSR) and

depth/variant allele frequency (VAF) filtering), was predictive of response (n=25, Mann-Whitney

p=0.0078).

One hypothesis for explaining the association between mutation load and outcome to

treatment with checkpoint blockade is the generation of neoantigens, altered peptides presented

by the major histocompatibility complex that are capable of eliciting an anti-tumor T cell

response and are more common with increased mutation load. After performing *in silico* HLA

typing (Methods), we examined predicted neoantigens that are 8 to 11 amino acids in length

resulting from the nonsynonymous mutations of patients treated with atezolizumab. There was

no significant association between predicted neoantigens per megabase and either DCB or 12

month overall survival. Patients with DCB had a median 4.58 (range 0.037-39.48) predicted

neoantigens per megabase while patients without DCB had 1.35 (range 0.00-20.22) (n=25,

Mann-Whitney p=0.55, Supplementary Figs 2C, 2D). Filtering of predicted neoantigens to focus

only on those expressed in RNA (Methods) also demonstrated no significant association

between expressed predicted neoantigens and clinical benefit with atezolizumab (n=25, Mann-

Whitney p=0.29, Fig 2C and Supplementary Fig S2B). Again, we acknowledge the limitations in

statistical power to detect associations due to the sample size of our study.

Impact of mutation load on response likelihood increases over time

Given that the mutation load and outcomes were weakly associated in the complete IMvigor210

dataset and not statistically significantly associated in this cohort, we embarked upon an

exploration of additional factors, including tumor microenvironmental and systemic measures,

which may modify the importance of this variable or independently impact outcomes.

To this end, we examined the time-varying impact of mutation load on PFS to see if

mutation load had a differential impact on early hazards in contrast to late hazards. We found

evidence of time-varying effects of somatic mutation load on progression-free survival in this

cohort (n=25, p=0.044 for association of scaled Schoenfeld residuals with log(time)). There was

little association of somatic mutation load with mortality or disease progression in the first 3

months (n=25, HR=0.91, 95% CI (0.75, 1.07); Fig 3A). In contrast, there was a notable

association of somatic mutation load with clinical events occurring more than 3 months after

treatment (n=11, HR=0.69, 95% CI (0.38, 0.99)). When a similar analysis was performed for

time-varying association with OS, the evidence in support of time-varying effects was similar

(n=25, p=0.082, Fig 3B). Among patients who survived longer than 3 months, the number of

somatic mutations per megabase was associated with a lower risk of mortality (n=11, HR=0.80,

95% CI (0.60, 1.00)) than during the first three months (n=25, HR=1.02, 95% CI (0.79, 1.22), Fig

3B). There was, however, weak evidence in support of a threshold at 3 months after follow-up.

In a nonparametric analysis, we found that the reduction in risk associated with somatic

mutation load increased steadily over time (Figs 3C, 3D, 3E).

Fig 3. Time dependent relationship between mutation load and treatment response.

(A) There was no significant association between somatic mutation load and PFS for events

occurring in the first 3 months (red box: HR=0.91, 95% CI (0.75, 1.07)), as compared to those

more than 3 months following therapy (blue box: HR=0.69, 95% CI (0.38, 0.99)). (B) There was

no significant association between somatic mutation load and OS for events occurring in the first

3 months (red box: HR=1.02, 95% CI (0.79, 1.22)), as compared to those more than 3 months

following therapy (blue box: HR=0.80, 95% CI (0.60, 1.00)). (C) The hazard ratio for each

mutation per megabase, estimated at each unique failure time. Red box plots summarize 50%

and 95% posterior intervals for each observed failure/censor time, with median values shown in

green. Time (Days) is plotted on a log-scale. Estimates are not independent from one another

since the model utilizes a random-walk parameterization to allow the variance in hazard over

time to be modeled flexibly. (D) Summary of estimated change in hazard for disease

progression or death with each observed mutation per megabase, reported as exp(beta) or

Hazard Ratios. Estimates here are reported per interval of failure time, with failure times

segmented at regular intervals of log(time) since therapy. Boxplots depict 50% and 95%

posterior intervals for hazard ratio at each interval, with median values shown with horizontal

lines. Intervals with no boxplot are those during which no failure or censor events were

observed. Note that these estimates are not independent from one another since the model

utilizes a random-walk parameterization to allow the variance in hazard over time to be modeled

flexibly. (E) Summary of estimated change in hazard for disease progression or death with each

observed mutation per megabase, reported as exp(beta) or Hazard Ratios. Estimates here are

reported per interval of failure time, with failure times segmented at regular intervals of days

since therapy. Boxplots depict 50% and 90% posterior intervals for hazard ratio at each interval,

with median estimates shown as horizontal lines. Intervals with no boxplot are those during

which no failure or censor events were observed. Note that these estimates are not independent

from one another since the model utilizes a random-walk parameterization to allow the variance

in hazard over time to be modeled flexibly.

These data suggest that in patients with rapidly progressive disease, factors other than

mutation load likely determine their outcome. This observation is not surprising in that clinical

factor analysis of this disease state has identified a heterogeneous population of patients, with 5

clinical factors distinguishing those likely to experience a rapid and early death from those more

likely to survive longer (11). We hypothesized that such patients might simply be too clinically

and systemically unwell to mount the necessary immune response, despite some of them

harboring tumor biomarkers thought to confer a likelihood of DCB, including elevated mutation

load. When we examined the 5-factor score in this subset relative to the rest of the dataset, we

found that indeed patients who survived less than or equal to 3 months exhibited a significantly

higher 5-factor score (3.00 (range 2.00-4.00), in contrast to 1.50 (range 0.00-4.00) in patients

who survived longer than 3 months (n=26, Mann-Whitney p=0.018, Supplementary Fig S3A).

Patients surviving less than 3 months were much more likely to have liver metastases: 100% in

patients surviving less than or equal to 3 months and 22% in patients surviving longer than 3

months (n=29, Fisher's Exact p=0.00097, Supplementary Fig S3B). There were no significant

differences in these patients with respect to BCG exposure (n=29, Fisher's Exact p=0.20),

missense SNV load (n=25, Mann-Whitney p=0.26) and pre-treatment peripheral TCR clonality

(n=29, Mann-Whitney p=0.12). These data suggest that there is a subset of nearly end-stage

patients with cancer in whom clinical variables may negate immunological response despite the

presence of one or more favorable tumor-associated biomarkers. The inclusion of these clinical

variables is warranted in future studies.

Examination of the tumor microenvironment shows evidence for adaptive

immunity and suppression in responding tumors

Several studies have suggested that an "inflamed" tumor microenvironment, tumor or immune

cell PD-L1 expression increase the likelihood of response to checkpoint blockade. As seen in

the published IMVigor 210 cohort, PD-L1 IC expression was significantly associated with DCB in

this subset (n=29, Spearman rho=0.48 p=0.0083, Supplementary Fig S4A). We quantified

immune infiltration from RNA-seq using ESTIMATE (17). The immune score, while associated

with the TIL proportion estimated through TCR Seq (Supplementary Fig S4B), was estimated to

be 764.37 (range -1195.08-1509.65) in patients with DCB and 263.49 (range -1100.78-1734.28)

in patients without DCB but was not significantly different (n=26, Mann-Whitney p=0.33,

Supplementary Fig S4C). When we performed gene set enrichment analysis (GSEA) using the

Hallmark Geneset (18), we did not observe any differentially expressed gene sets between

tumors from patients with DCB versus no DCB. Furthermore, RNA expression of *PD-L1* did not

correlate with reported immune cell PD-L1 staining level (n=26, Spearman rho=0.045 p=0.83,

Supplementary Fig S4D). We did not observe a difference in tumor MHC class I expression

according to DCB (Supplementary Fig S4E, *HLA-A*: n=26, Mann-Whitney p=0.26, *HLA-B*: n=26,

Mann-Whitney p=0.36, *HLA-C*: n=26, Mann-Whitney p=0.24).

Given that such agnostic approaches did not reveal a clear association between tumor

microenvironment factors and response, we pursued a hypothesis-driven approach examining

the genes that show upregulation at the cell surface during T cell exhaustion. When categorized

by DCB, there was no significant difference in expression of such genes, including *CTLA-4,*

TIGIT, HAVCR2 (TIM-3) or *LAG-3* (19). When grouped by PD-L1 staining, we found low

expression of all markers in the PD-L1 low group (IC0), as expected. However, in the PD-L1

high group (IC2), *HAVCR* exhibited significantly higher expression in tumors from patients who

experienced DCB than in those who did not (Supplementary Fig S4F). Interestingly, of the three

IC2 tumors, two had missense SNV loads significantly below the median (17 and 57); the third

had 412 SNVs. Additionally, although Rosenberg and colleagues (2) found that among the four

TCGA subtypes of RNA expression, luminal cluster II showed a significantly higher response

rate, no significant association was found here between the four clusters and DCB (n=20,

Fisher's Exact p=0.36) (Supplementary Fig S4G), nor between the luminal/basal sub-

categorization and DCB (n=20, Fisher's Exact p=1.00), possibly due to sample size.

Relative importance of somatic, immune, and clinical factors in resistance

and response to PD-L1 blockade

Unanswered questions that arise from the many studies of biomarker correlates of checkpoint

blockade response are whether measures such as mutation load, PD-L1 staining, and others

reflect the same "tumor state," or if each confers an independent effect on outcome?

When examined in conjunction with mutation load, the greater the expression of PD-L1,

the more negative the association of mutation load with hazard (i.e. higher mutation load was

associated with longer survival). Among patients with tumors showing little-to-no expression of

PD-L1 (IC0 rated), each unit increase in missense SNV count per megabase was associated

with a negligible change in hazard (n=4, HR=1.43, 95% CI (0.75, 2.98)). Among patients with

tumors expressing PD-L1 at moderate or high levels (IC1 or IC2 staining), missense SNV count

per megabase was associated with lower risk for disease progression or mortality (among IC1:

n=11, HR=0.75, 95% CI (0.47, 1.14); among IC2: n=10, HR=0.73, 95% CI (0.48, 1.06)).

Although our limited sample size precludes making an assertion that mutation load is associated

with survival in any particular subgroup (e.g. when looking among IC1 and IC2 tumors alone);

our data do support the presence of an interaction among these variables (p=0.046 for

interaction; Supplementary Fig S5A). Given the plausibility of the finding that somatic mutation

load may correlate better with survival among patients with an inflamed tumor

microenvironment, the addition of somatic mutation load to PD-L1 IC staining warrants further

study.

We found a similar albeit weaker interaction effect when looking at association of

somatic mutation load (missense SNV count per megabase) and progression-free survival

according to the presence/absence of liver metastasis prior to treatment administration (p=0.14

for interaction). Among patients without liver metastasis, somatic mutation load was associated

with a lower risk for disease progression or mortality (n=16, HR=0.73, 95% CI (0.50, 1.02),

Supplementary Fig S5B) than patients with liver metastasis (n=9, HR=0.96, 95% CI (0.66, 1.37),

Supplementary Fig S5B).

To our surprise, although both PD-L1 staining and mutation load were each associated

with response in the original study (2) these variables did not correlate with each other (Fig 4A).

Furthermore, pre-treatment peripheral TCR clonality did not correlate with mutation load (Fig

4B). The lack of association between these variables suggests that each might confer an

independent or semi-independent impact on the likelihood of response to therapy. TCR clonality

and infiltration did, however, correlate with PD-L1 IC score: those tumors with higher clonality or

higher infiltration also featured higher PD-L1 staining (p=0.02 and p=0.01, respectively, Figs 4C,

4D).

Fig 4. Associations between measured somatic, immune and clinical variables.

(A) Although both PD-L1 staining and mutation load each were weakly associated with

response, these variables were not correlated with each other (n=25, Spearman rho=0.14

p=0.51). (B) Pre-treatment peripheral TCR clonality did not correlate with mutation load (n=25,

Pearson r=0.0017 p=0.99). (C) TIL proportion as estimated by TCR sequencing was associated

with PD-L1 IC staining (n=24, Spearman rho=0.51 p=0.010). (D) TIL clonality was associated

with PD-L1 IC staining (n=24, Spearman rho=0.48 p=0.017). (E) Hazard associated with

log(pre-treatment peripheral TCR clonality) by level of immune cell PD-L1 expression (IC0, IC1

or IC2). (F) Multivariate survival analysis of various clinical, peripheral and intratumoral

biomarkers for association with time to disease progression or mortality (PFS), utilizing a

varying-coefficient model which allows the hazard associated with a one-unit increase in a

biomarker's value to vary according to level of intratumoral PD-L1 expression (IC score).

In an analysis to see whether the association between pre-treatment peripheral TCR

clonality and progression-free survival varied by PD-L1 IC score, we found some evidence of an

interaction (p=0.015 for interaction; Fig 4E). Among patients with low levels of PD-L1

expression, there was little association between pre-treatment peripheral TCR clonality and

progression-free survival (among IC0: n=4, HR=1.87, 95% CI (0.49, 4.86); among IC1: n=11,

HR=0.69, 95% CI (0.14, 1.85)). Among patients with high levels of PD-L1 expression, by

comparison, we observed almost complete separation of progression-free survival according to

pre-treatment peripheral TCR clonality (among IC2: n=10, HR=86.22, 95% CI (2.55, 491.65);

Fig 4E). Similar results were seen in analyses with respect to OS, and in a logistic regression

analysis for DCB (Supplementary Figs 5C, 5D, 5E).

To resolve the hypothesis that those patients with low peripheral TCR clonality simply

were healthier, we examined the association between 5-factor score and pre-treatment

peripheral TCR clonality and did not find such an association (n=26, Spearman rho=0.25

p=0.22, Supplementary Fig S5F).

In a multivariate survival model for time to disease progression or mortality, which allows

the effect of each biomarker to vary according to intratumoral PD-L1 IC score, we find that the

correlation of each intratumoral, peripheral or clinical biomarker with disease progression or

mortality is relatively independent of the others (Fig 4F, Supplementary Fig S5G). Perhaps with

the notable exception of the impact of liver metastatic status on time to progression or survival,

the correlation of each intratumoral or peripheral biomarker with outcome is strongest in the

group with the highest levels of immune cell PD-L1 expression (Supplementary Table S2).

551 Discussion

The treatment of previously incurable metastatic solid tumors with checkpoint blockade agents

has led to dramatic success in a minority of patients, a finding that has generated substantial

excitement in the field, with associated correlative studies and drug development. Despite such

studies, a deeper understanding of the biology of response and resistance is often eclipsed by

the search for biomarkers. Of the multitude of studied biomarkers, to date, PD-L1 staining

(either tumor cell or immune cell) and mutation load have emerged as the most consistently

positive predictors of response to checkpoint blockade (10,20,21). However, in all studies the

exceptions are patients who benefit despite having tumors with low mutation load or who lack

high PD-L1 staining, which demonstrates the inadequacy of these biomarkers in excluding and

therefore denying a patient with limited or no treatment options access to potentially life-saving

therapy.

Here, we undertook the in-depth characterization of tumors and peripheral blood from 29

patients treated on IMvigor 210, a Phase II study in which 310 patients were treated with the

anti-PD-L1 agent atezolizumab. Although the overall study found significant associations

between mutation load as measured by the Foundation Medicine targeted sequencing panel

and radiographic response (2), there was no statistically significant association between

mutation load and durable clinical benefit or survival in the patient subset studied here, despite

the similarity of our study population to the parent study. This contrast may be due to a

combination of factors. First, though statistically significant, the association in the overall study

was not categorical: as in other studies of mutation load, this factor alone was not predictive of

response. Second, we have less power to detect this association in our smaller subset

compared with the larger studied cohort. Third, standardized definitions and calculations of

mutation load do not exist as yet; each published study has used differing methodologies

(2,10,20,21). Indeed, in this study, depending on the method used, the association between

mutation load and clinical outcomes varied from $p<0.08$ to $p>0.4$ (AUCs and p-values in

Supplementary Table S1). To illustrate the fickle nature of defining mutation load, counting only

the mutations excluded by BQSR, as opposed to only those remaining after BQSR, showed a

significant association with DCB. Together, these findings underscore the need for improved

and standardized mutation calling methods. The weak association of mutation load with DCB

and the lack of such standardization render this biomarker unfit for application to individual

patients at present. Furthermore, the time-dependent relationship between mutation load and

survival implies that a clinical and immunological state may exist in patients with advanced

cancer, such that patients with very rapidly progressing disease and expected death in <3

months do not respond despite the presence of positive biomarkers.

In an attempt to deepen our understanding of the biology of response and resistance, we

studied additional factors: immune factors, conveyed here as peripheral and intratumoral TCR

clonality and TIL proportion; and systemic factors such as the 5-factor score and presence of

liver metastases. We found that even in this small dataset, TCR clonality below the median in

the peripheral blood prior to treatment, expansion of tumor-associated TCR in the periphery 3

weeks after initiating treatment, and higher TIL proportion all associated with clinical benefit.

These data suggest that TCR sequencing provided additional insights into response and

resistance beyond mutation load and PD-L1 staining. With respect to biomarker development,

our study implies that non-invasive metrics such as pre-treatment peripheral TCR clonality and

known prognostic features such as the presence of liver metastases may be worthy of further

study in urothelial cancer patients treated with PD-L1 blockade.

Finally, though limited in power by the small sample size, we attempted to integrate the

importance of the studied variables. This analysis demonstrated both hypothesized and

unexpected interactions. For example, while mutation burden seemed to associate with

outcome more significantly in PD-L1 IC1 and-2 tumors, high PD-L1 IC staining in the setting of

high peripheral TCR clonality was associated with a substantial hazard for poor outcome. Given

the significance of PD-L1 expression in mediating response to anti-PD-L1 therapy, the presence

of these interactions may argue in their favor as predictive rather than prognostic biomarkers.

Further analysis is required to elucidate the role of these biomarkers in mediating response to

checkpoint blockade.

In conclusion, we have demonstrated that to truly understand and ultimately circumvent

resistance to checkpoint blockade, we must pursue integrated studies of the somatic, immune

and systemic features of each treated patient.

613 Acknowledgements: Carol Aghajanian and Jedd Wolchok for their mentorship of AS.

615 Funding: This work was supported by the Ludwig Center for Cancer Research and the NIH/NCI

616 Cancer Center Support Grant P30 CA008748.

618 Supplementary Information, Captions

619 Supplementary Tables

Supplementary Table S1. Choice of depth and VAF filtering, as well as whether or not to run

Base Quality Score Recalibration (BQSR), resulted in differences in predictive value for

mutation load. We chose to optimize for precision, choosing the highlighted set of filters/BQSR.

Precision is defined as the fraction of filtered missense mutations in IMPACT genes that were

actual IMPACT panel variants. Recall is defined as the fraction of actual IMPACT panel variants

that were found in the filtered missense results.

Supplementary Table S2. Summary of results from multivariate survival analysis of various

clinical, peripheral and intratumoral factors to estimate their independent association with

hazard for disease progression or mortality (PFS) and for mortality (OS) according to level of

intratumoral PD-L1 expression (IC grade). Results are summarized as median and 95%

posterior intervals.

Supplementary Figures

Supplementary Fig S1A. Tumors with less than the median TIL proportion or clonality were less likely to display DCB (25%, versus no DCB: 81%, n=24, Fisher's Exact p=0.021).

Supplementary Fig S1B. TCR overlap between the pre-treatment and 3-week posttreatment peripheral blood in one patient with limited clinical benefit (PFS=37 days) and one patient with durable clinical benefit (CR at 630 days after starting treatment). The association between pre-treatment peripheral blood TCR sequences (x axis) and posttreatment peripheral blood TCR sequences (y axis) is overlaid with the presence of tumor associated T cell clones. Gray indicates TCRs present only in the peripheral blood; blue indicates TCRs present in the tumor and blood; orange indicates TCRs present in the tumor and expanded in the blood with treatment.

Supplementary Fig S1C. There was no significant expansion of TIL-associated TCR clones between pre-treatment (3.00 (range 1.00-9.00)) and 6 weeks post-treatment (2.00 (range 1.00-8.00)), n=20, Mann-Whitney p=0.17.

Supplementary Fig S1D. The combination of high pre-treatment TIL and low pre-treatment peripheral blood TCR clonality were predictive of DCB (n=24, Fisher's Exact p=0.0069) and DCB-OS (n=24, Fisher's Exact p=0.014). For DCB, a logit model combining both was more predictive than peripheral blood (n=24, log-likelihood p=0.00029) or TIL (n=24, log-likelihood p=0.00051) clonality alone. For DCB-OS, both combined were more predictive than TIL (n=24, log-likelihood p=0.0029) clonality alone.

Supplementary Fig S2A. No significant association between the number of missense SNV per

megabase and overall survival, with 2.13 (range 0.038-11.46) in tumors from those patients who

survived greater than 12 months, versus 0.48 (range 0.019-9.90) in those who did not (n=25,

Mann-Whitney p=0.37).

Supplementary Fig S2B. No significant difference between median expressed neoantigens in

tumors from patients who survived greater than or equal to 12 months: was 1.31 (range 0.00-

6.06), versus 0.35 (range 0.00-5.30) in those who survived less than 12 months (n=25, Mann-

Whitney p=0.36).

Supplementary Fig S2C. No significant difference between median predicted neoantigens per

megabase: 4.58 (range 0.037-39.48) in tumors from patients with DCB, as compared to 1.35

(range 0.00-20.22) in those who progressed in less than 6 months (no DCB) (n=25, Mann-

Whitney p=0.55).

Supplementary Fig S2D. No significant difference between median predicted neoantigens per

megabase in tumors from those patients who survived greater than 12 months (here used to

define OS DCB) was 3.56 (range 0.037-39.48) as compared to 1.37 (range 0.00-20.22) in those

who did not (no DCB) (n=25, Mann-Whitney p=0.81).

Supplementary Fig S3A. Patients who survived less than 3 months (red box) exhibited a

significantly higher 5-factor score (3.00 (range 2.00-4.00), as compared to 1.50 (range 0.00-

4.00) in patients who survived >3mo (blue box) (n=26, Mann-Whitney p=0.018).

Supplementary Fig S3B. Patients who survived less than or equal to 3 months (red box) were

more likely to have liver metastases (100% in patients who survived less than or equal to 3

months and 22% in patients who survived longer than 3 months, n=29, Fisher's Exact

p=0.00097).

Supplementary Fig S4A. PD-L1 IC staining as reported by the sponsor in the published study

(2) and outcome in our cohort were significantly associated in this sub-study (n=29, Spearman

rho=0.48 p=0.0083).

Supplementary Fig S4B. ImmuneScore was associated with TIL proportion (n=24, Spearman

rho=0.47 p=0.022).

Supplementary Fig S4C. There was no association between ImmuneScore and DCB (DCB,

764.37 (range -1195.08-1509.65); no DCB 263.49 (range -1100.78-1734.28) (n=26, Mann-

Whitney p=0.33).

Supplementary Fig S4D. PD-L1 expression as measured by RNA-seq was not associated with

PD-L1 IC level (n=26, Spearman rho=0.045 p=0.83). Tumor cell PD-L1 staining was not

available.

Supplementary Fig S4E. HLA Class I expression was not associated with DCB (HLA-A: n=26,

Mann-Whitney p=0.26, HLA-B: n=26, Mann-Whitney p=0.36, HLA-C: n=26, Mann-Whitney

p=0.24).

Supplementary Fig S4F. Expression of other inhibitory markers, in particular *HAVCR2* (also

known as *TIM-3*) was higher in DCB patients in the IC2 group.

Supplementary Fig S4G. No association was found between TCGA RNA Subtype and response

in this sub-study (n=20, Fisher's Exact p=0.36).

Supplementary Fig S5A. Hazard associated with log(missense SNV count per megabase) by

level of immune cell (IC0, IC1 or IC2) PD-L1 expression.

Supplementary Fig S5B. Hazard associated with log(missense SNV count per megabase) by

presence or absence of liver metastasis at enrollment.

Supplementary Fig S5C. Association of peripheral TCR clonality prior to treatment with time to

mortality (OS) varies according to immune cell (IC0, IC1 or IC2) PD-L1 expression.

Supplementary Fig S5D. Association of peripheral TCR clonality prior to treatment with DCB

varies according to immune cell (IC0, IC1 or IC2) PD-L1 expression.

Supplementary Fig S5E. Association of peripheral TCR clonality prior to treatment with DCB

(OS) varies according to immune cell (IC0, IC1 or IC2) PD-L1 expression.

Supplementary Fig S5F. There was no significant relationship between 5-Factor score and pre-

treatment TCR clonality (n=26, Spearman rho=0.25 p=0.22).

Supplementary Fig S5G. Multivariate survival analysis of various clinical, peripheral and

intratumoral biomarkers for association with time to mortality (OS), utilizing a varying-coefficient

model which allows the hazard associated with a one-unit increase in a biomarker's value to

vary according to level of intratumoral PD-L1 expression (IC score). Note that the x-axis has

been truncated at a value of 10 for clarity even though this results in the exclusion of some

estimated HR values (specifically that for pre-treatment peripheral TCR clonality among IC2

patients).

Supplementary Fig S6A. No significant association between the number of missense SNV found

on MSK-IMPACT and DCB (DCB 0.13 (range 0.00-0.31) versus no DCB 0.046 (range 0.00-

0.37) (n=25, Mann-Whitney p=0.42).

Supplementary Fig S6B. No significant association between the number of missense SNV found

on MSK-IMPACT and overall survival (survival greater than 12 months 0.093 (range 0.00-0.31),

versus less than 12 months 0.074 (range 0.00-0.37) (n=25, Mann-Whitney p=0.78).

Supplementary Fig S6C. There was no significant difference in APOBEC signature found in

tumors from patients with PFS DCB (0.19 (range 0.00-0.56)) as compared to no DCB (0.00

(range 0.00-0.46)) (n=25, Mann-Whitney p=0.23).

Supplementary Fig S6D. There was a significant correlation between missense SNV count and

APOBEC signature mutations (n=25, Pearson r=0.40 p=0.048).

Supplementary Fig S6E. There was no significant association between FGFR3 mutations or

expression (n=26, Mann-Whitney p=0.39).

Supplementary Fig S6F. There was no significant association between MYC expression and

outcome measured by PFS DCB (n=26, Mann-Whitney p=0.87).

Supplementary Fig S6G. There was no significant association between DNA damage response

(DDR) mutations rated as "possible" by PolyPhen and DCB (n=25, Mann-Whitney p=0.20).

Supplementary Fig S6H. Univariate association of exonic SNV, missense SNV and neoepitope

load with DCB, with (blue bars) and without (red bars) filtering by expression.

Supplementary Fig S6I. Univariate association of exonic SNV, missense SNV and neoepitope

load with OS greater than 12 months, with (blue bars) and without (red bars) filtering by

expression.

Supplementary Fig S6J. Univariate association of exonic SNV, missense SNV and neoepitope

load with PFS, showing results with (blue bars) and without (red bars) filtering by expression.

Supplementary Fig S6K. Univariate association of exonic SNV, missense SNV and neoepitope

load with OS, showing results with (blue bars) and without (red bars) filtering by expression.

Supplementary Fig S6L. Univariate association of expressed/total ratio for exonic SNV,

missense SNV, and neoantigen loads with PFS.

Supplementary Fig S6M. Univariate association of expressed/total ratio for exonic SNV,

missense SNV, and neoantigen loads with OS.

Supplementary Fig S6N. Univariate association of expressed/total ratio for exonic SNV,

missense SNV, and neoantigen loads with DCB (PFS greater than 12 months).

Supplementary Fig S6O. Univariate association of expressed/total ratio for exonic SNV,

missense SNV, and neoantigen loads with OS greater than 12 months.

Supplementary Fig S6P. There was no significant association between pack years of reported

smoking history and DCB (n=29, Mann-Whitney p=0.87).

Supplementary Fig S6Q. There were significant associations between PFS (n=29, log-rank

p=0.024) and OS (n=29, log-rank p=0.018) and the presence of liver metastasis.

Supplementary Fig S6R. There was a significant association between the presence of visceral

metastases and poor overall survival (n=29, log-rank p=0.020).

Supplementary Fig S6S. There was not a significant association between 5-factor score and OS

(n=26, log-rank p=0.13).

Supplementary Fig S6T. No significant association between the number of expressed missense

SNV per megabase and DCB (DCB 0.79 (range 0.00-3.36), versus no DCB: 0.16 (range 0.00-

3.34)), n=25, Mann-Whitney p=0.26.

References

1. Powles T, Eder JP, Fine GD, Braiteh FS, Loriot Y, Cruz C, et al. MPDL3280A (anti-PD-L1) treatment leads to clinical activity in metastatic bladder cancer. Nature. 2014;515(7528):558–62.

2. Rosenberg JE, Hoffman-Censits J, Powles T, van der Heijden MS, Balar AV, Necchi A, et al. Atezolizumab in patients with locally advanced and metastatic urothelial carcinoma who have progressed following treatment with platinum-based chemotherapy: a single-arm, multicentre, phase 2 trial. Lancet. 2016;387(10031):1909–20.

3. Postow MA, Yuan J, Kitano S, Lesokhin AM, Wolchok JD. Markers for anti-cytotoxic T-lymphocyte antigen 4 (CTLA-4) therapy in melanoma. Methods Mol Biol. 2014;1102:83–95.

4. Robins HS, Campregher PV, Srivastava SK, Wacher A, Turtle CJ, Kahsai O, et al. Comprehensive assessment of T-cell receptor beta-chain diversity in alphabeta T cells. Blood. 2009 Nov 5;114(19):4099–107.

5. Lefranc M-P, Giudicelli V, Duroux P, Jabado-Michaloud J, Folch G, Aouinti S, et al. IMGT®, the international ImMunoGeneTics information system® 25 years on. Nucleic Acids Res. 2015 Jan;43(Database issue):D413–22.

6. Wu D, Emerson RO, Sherwood A, Loh ML, Angiolillo A, Howie B, et al. Detection of minimal residual disease in B lymphoblastic leukemia by high-throughput sequencing of IGH. Clin Cancer Res. 2014 Sep 1;20(17):4540–8.

7. Hsu MS, Sedighim S, Wang T, Antonios JP, Everson RG, Tucker AM, et al. TCR Sequencing Can Identify and Track Glioma-Infiltrating T Cells after DC Vaccination. Cancer Immunol Res. 2016 May;4(5):412–8.

8. Tumeh PC, Harview CL, Yearley JH, Shintaku IP, Taylor EJM, Robert L, et al. PD-1 blockade induces responses by inhibiting adaptive immune resistance. Nature. 2014 Nov 27;515(7528):568–71.

9. DeWitt WS, Emerson RO, Lindau P, Vignali M, Snyder TM, Desmarais C, et al. Dynamics of the cytotoxic T cell response to a model of acute viral infection. J Virol. 2015 Apr;89(8):4517–26.

10. Rizvi, N., Hellmann, M, Snyder, Et al A. Mutational Landscape Determines Sensitivity to Programmed Cell Death-1 Blockade in Non-Small Cell Lung Cancer. Science. 2015;

11. Sonpavde G, Pond GR, Rosenberg JE, Bajorin DF, Choueiri TK, Necchi A, et al. Improved 5-Factor Prognostic Classification of Patients Receiving Salvage Systemic Therapy for Advanced Urothelial Carcinoma. J Urol. 2016;195(2):277–82.

12. Bajorin DF, Dodd PM, Mazumdar M, Fazzari M, McCaffrey JA, Scher HI, et al. Long-term

survival in metastatic transitional-cell carcinoma and prognostic factors predicting outcome
of therapy. J Clin Oncol. 1999 Oct;17(10):3173–81.

13. Dunn GP, Old LJ, Schreiber RD. The immunobiology of cancer immunosurveillance and
immunoediting. Immunity. 2004;21(2):137–48.

14. Cha E, Klinger M, Hou Y, Cummings C, Ribas A, Faham M, et al. Improved survival with T
cell clonotype stability after anti-CTLA-4 treatment in cancer patients. Sci Transl Med.
2014;6(238):238ra70.

15. Alioto TS, Buchhalter I, Derdak S, Hutter B, Eldridge MD, Hovig E, et al. A comprehensive
assessment of somatic mutation detection in cancer using whole-genome sequencing. Nat
Commun. 2015 Dec 9;6:10001.

16. Qiu P, Pang L, Arreaza G, Maguire M, Chang KC, Marton MJ, et al. Data Interoperability of
Whole Exome Sequencing (WES) Based Mutational Burden Estimates from Different
Laboratories. Int J Mol Sci [Internet]. 2016;17(5). Available from:
http://dx.doi.org/10.3390/ijms17050651

17. Bindea G, Mlecnik B, Tosolini M, Kirilovsky A, Waldner M, Obenauf AC, et al.
Spatiotemporal dynamics of intratumoral immune cells reveal the immune landscape in
human cancer. Immunity. 2013;39(4):782–95.

18. Liberzon A, Subramanian A, Pinchback R, Thorvaldsdottir H, Tamayo P, Mesirov JP.
Molecular signatures database (MSigDB) 3.0. Bioinformatics. 2011;27(12):1739–40.

19. Tirosh I, Izar B, Prakadan SM, Wadsworth MH 2nd, Treacy D, Trombetta JJ, et al.
Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq.
Science. 2016;352(6282):189–96.

20. Van Allen EM, Miao D, Schilling B, Shukla SA, Blank C, Zimmer L, et al. Genomic
correlates of response to CTLA-4 blockade in metastatic melanoma. Science.
2015;350(6257):207–11.

21. Snyder A, Makarov V, Merghoub T, Yuan J, Zaretsky JM, Desrichard A, et al. Genetic basis
for clinical response to CTLA-4 blockade in melanoma. N Engl J Med. 2014;371(23):2189–
99.

22. Forbes SA, Beare D, Gunasekaran P, Leung K, Bindal N, Boutselakis H, et al. COSMIC:
exploring the world's knowledge of somatic mutations in human cancer. Nucleic Acids Res.
2015 Jan;43(Database issue):D805–11.

23. Rosenthal R, McGranahan N, Herrero J, Taylor BS, Swanton C. DeconstructSigs:
delineating mutational processes in single tumors distinguishes DNA repair deficiencies
and patterns of carcinoma evolution. Genome Biol. 2016 Feb 22;17:31.