Title: Genetic variation for mitochondrial function in the New Zealand freshwater snail *Potamopyrgus antipodarum*

Authors: Joel Sharbrough[1, 3], Jennifer L. Cruise[2], Megan Beetch[2], Nicole M. Enright[1], & Maurine Neiman[1]

[1] University of Iowa, Department of Biology, 143 Biology Building, Iowa City, IA 52242
[2] University of St. Thomas, Department of Biology, 2115 Summit Ave., Saint Paul, MN 55105
[3] Colorado State University, Department of Biology, Anatomy-Zoology Building E212, Fort Collins, CO 80523

Corresponding Author:

Joel Sharbrough
Colorado State University, Department of Biology
Anatomy-Zoology Building E212
Fort Collins, CO 80523
Email: jsharbro@rams.colostate.edu
Phone: (970) 491-2256

Other Contact Information:
Jennifer L. Cruise: jlcruise@stthomas.edu
Megan Beetch: mbeetch@purdue.edu
Nicole M. Enright: enrightnm@g.cofc.edu
Maurine Neiman: maurine-neiman@uiowa.edu

Running Title: Asexual lineages vary in mitochondrial function

Article Type: Original Article

ABSTRACT

The proteins responsible for mitochondrial function are encoded by two different genomes with

distinct inheritance regimes, rendering rigorous inference of genotype–phenotype connections

intractable for all but a few model systems. Asexual lineages provide a powerful means of

addressing these challenges by enabling comparisons among genetically identical individuals in

which nuclear and mitochondrial genomes are co-inherited. Here, we used *Potamopyrgus*

antipodarum, a New Zealand freshwater snail that is an emerging model system for the evolution

of sexual reproduction, adapting several methods for measuring mitochondrial function in

mollusks at three distinct levels of biological organization: mitogenomic, organellar, and

organismal. We applied these methods to multiple asexual lineages of *P. antipodarum* and

evaluated whether there exists genetic variation (that is, the raw material for adaptive evolution)

for mitochondrial genome copy number, mitochondrial membrane potential, and organismal

oxygen consumption. Our analyses revealed that asexual lineages of *P. antipodarum* differed in

terms of mitochondrial function at all three levels of biological organization. The discovery of

lineage-level variation in terms of mitochondrial function combined with the methods developed

here set the stage to study central evolutionary questions including whether mitochondrial

mutation accumulation influences the maintenance of sexual reproduction in natural populations.

Keywords: Cellular respiration, oxygen consumption, mitochondrial function, asexual

reproduction, mtDNA copy number, mitochondrial membrane potential

INTRODUCTION

Mitochondrial function is of critical importance to eukaryotic health (for example, Chen *et al.*,
2007, Dowling, 2014), and genetic variation for mitochondrial function has been linked to
evolutionary adaptation (for example, Rawson and Burton, 2002) and disease (DiMauro and
Schon, 2001). The role of genetic variation for mitochondrial function is complicated by direct
interaction between nuclear encoded and mitochondrially encoded proteins, particularly with
respect to oxidative phosphorylation (OXPHOS) (reviewed in Rand *et al.,* 2004). Mitonuclear
cooperation is particularly important for proper enzyme function in OXPHOS complexes I, III,
IV, and V because these complexes are comprised of subunits encoded by both genomes.
Accordingly, discordance between mitochondrial and nuclear genomes has been demonstrated to
have negative fitness and/or functional consequences in a variety of animals, including copepods
(Ellison and Burton, 2006), *Drosophila* (Meiklejohn *et al.,* 2013; Pichaud *et al.,* 2013), and
salamanders (Lee-Yaw *et al.,* 2014).

   In sexually reproducing organisms, the maintenance of mitonuclear cooperation is further
complicated by the expectation that the different mechanisms of nuclear *vs*. mitochondrial
genome inheritance will differentially affect the generation, maintenance, and distribution of
genetic variation. In particular, biparental inheritance and meiotic recombination in the nuclear
genome will increase effective population size relative to the (typically) uniparentally inherited
and non-recombinant mitochondrial genome (reviewed in Barr *et al.,* 2005; Neiman and Taylor,
2009). This logic is the basis for the expectation that mitochondrial genomes will, when
compared to the nuclear genome, experience reduced efficacy of selection and suffer an
increased rate of accumulation of mildly deleterious mutations (reviewed in Neiman and Taylor,

2009). This mechanism is also thought to generate selection favoring compensatory changes in

nuclear-encoded mitochondrial subunits (Sloan *et al.,* 2013; Zhang and Broughton, 2013).

Asexual taxa provide a particularly interesting context in which to evaluate mitochondrial

function and evolution because the absence of recombination and segregation in asexual nuclear

genomes means that asexual lineages will transmit their mitochondrial genomes in complete

linkage disequilibrium (LD) with the nuclear genome. The co-transmission of nuclear and

mitochondrial genomes is expected to result in two non-mutually exclusive evolutionary

scenarios: (1) increased LD should decrease the efficacy of selection in both genomes *via*

selective interference (reviewed in Neiman and Taylor, 2009), and (2) tight linkage between

nuclear and mitochondrial genomes may allow epistatic variation to become available to natural

selection (reviewed in Neiman and Linksvayer, 2006). As such, the loss of sex likely has

profound consequences for the evolution of mitochondrial function, most of which remain

unevaluated.

Potamopyrgus antipodarum, a New Zealand freshwater snail, is very well suited for

investigating mitochondrial function in the absence of sex because otherwise similar obligately

sexual and obligately asexual individuals frequently coexist in natural populations. The transition

from sexual ancestor to asexual descendent has occurred multiple separate times within this

species, providing many so-called "natural experiments" into the consequences of asexuality

(Neiman *et al.,* 2011; Paczesniak *et al.,* 2013). Surveys of patterns of molecular evolution in the

mitochondrial genome of *P. antipodarum* have demonstrated that asexual lineages accumulate

nonsynonymous substitutions in their mitochondrial genomes more rapidly than their sexual

counterparts (Neiman *et al.,* 2010; Sharbrough *et al.,* 2016) and retain radical amino acid-

changing polymorphisms longer than sexual lineages (Sharbrough *et al.,* 2016). What remains

unevaluated is whether and how these accumulated changes resulting from the loss of sex have

shaped variation in mitochondrial function. Because natural selection can only operate on genetic

variation that is linked to fitness, the critical first step in evaluating how asexuality impacts

evolution of mitochondrial function in *P. antipodarum* is to evaluate whether asexual lineages

exhibit variation in mitochondrial function. The direct transmission of nuclear and mitochondrial

genetic information from mother to daughter in asexuals means that comparisons of

mitochondrial function across asexual lineages raised under shared environmental conditions will

provide an estimate of heritable variation for mitochondrial function in asexual lineages of *P.*

antipodarum.

Here, we use a common-garden approach to test whether asexual lineages of *P.*

antipodarum vary in mitochondrial function at three distinct levels of biological organization: (1)

mitogenomic (mtDNA copy number), (2) organellar (mitochondrial membrane potential and

electron transport), and (3) organismal (total oxygen (O_2) consumption), three traits that have all

been linked to proper mitochondrial function in other taxa. First, mitochondrial genome copy

number is tightly regulated and is positively correlated with cytochrome C oxidase activity (Van

den Bogert *et al.,* 1993), and low copy number has been implicated in a number of human

diseases (reviewed in Montier *et al*., 2009). Second, elevated mitochondrial membrane potential,

generated by electron transport through OXPHOS, is positively correlated with the ratio of ATP:

ADP, indicating that stronger membrane potentials reflect healthier mitochondria (Nicholls,

2004). Third, because O_2 is the final electron acceptor for the production of ATP (Chance and

Williams, 1955), total O_2 consumption represents an integrated measure of the ATP production

capacity of an organism. We therefore adapted well-established mitochondrial functional assays,

previously employed exclusively in model organisms, to probe the number of mitochondrial

genome copies relative to the nuclear genome, estimate the strength of the electrochemical

gradient, and quantify the flow of electrons through the electron transport chain (ETC) to test for

genetic variation in mitochondrial function in asexual lineages of *P. antipodarum.* Because heat

stress can increase inefficiencies in oxidative phosphorylation (Heise *et al.,* 2003), we also used

heat treatments and aquatic respirometry to quantify and compare total O_2 consumption under

heat stress across asexual *P. antipodarum* lineages. Together, these analyses revealed substantial

variation for mitochondrial function across asexual lineages at all three levels of biological

organization. More broadly, the assays adapted here provide a suite of useful experimental tools

for investigating mitochondrial function in *P. antipodarum* and, potentially, other mollusks.

MATERIALS AND METHODS

Snail husbandry

We compared mitochondrial function across a diverse array of asexual *P. antipodarum* lineages

(Neiman *et al.,* 2011; Paczesniak *et al.,* 2013; Table S1) reared under identical conditions for

multiple generations, enabling us to interpret across-lineage variation in our various measures of

mitochondrial function as representing genetic variation for these traits. Asexuality was

established for each lineage by determining ploidy using flow cytometry (sexual *P. antipodarum*

are diploid, asexuals are polyploid), as described in Neiman *et al.* (2011). Asexual lineages were

chosen for functional assays to represent the range of mitochondrial diversity found in New

Zealand populations (Neiman *et al.,* 2011) and to maximize our ability to compare different

functional assays. Adult female snails were selected arbitrarily from lineage populations for each

assay. Following standard laboratory protocols for *P. antipodarum* (e.g., Zachar and Neiman,

2013), snails were housed at 16ºC on an 18 hr light/6 hr dark schedule and fed *Spirulina* algae 3x

per week.

Mitochondrial function at the genomic level

Mitochondrial genome copy number is known to affect respiratory capacity (Van den Bogart *et*

al., 1993) and to be dynamically regulated in response to various cellular environmental cues

(Hori *et al.,* 2009; Matsushima *et al.,* 2010; Kelly *et al*., 2012). As such, mitochondrial genome

copy number represents a potentially important phenotypic level at which selection could favor a

compensatory mechanism to reduce the potentially negative consequences of mitochondrial

mutation accumulation. To test whether asexual *P. antipodarum* exhibited across-lineage

phenotypic variation for mitochondrial copy number, we estimated mitochondrial genome copy

number relative to a putatively single-copy nuclear gene using quantitative PCR (qPCR) in six

asexual triploid lineages. A putatively single-copy gene from the *P. antipodarum* nuclear

genome (*rad21*; Neiman *et al.,* unpublished) and mitochondrial *cytB* were chosen for analysis.

Primer pairs were designed to generate a 264-bp *rad21* product (F: 5'–

GATTCCAACAACTGATGTTTG –3', R: 5'–CAAAACTTACTCTAAATCTGC–3') and a

194-bp amplicon from *cytB* (F: 5'–TATGAATATTCAGATTTTTTAAATA–3', R: 5'–

CCTTAACTCCTAATCTTGGTAC–3'). For measurement standards, each of these products was

cloned from total DNA from a single *P. antipodarum* individual into the pGEM T-Easy plasmid

vector (Promega Corp., Madison, WI). Linearized plasmids were diluted in the presence of

carrier human genomic DNA to produce samples containing 300 – 300,000 copies of either the

nuclear or mitochondrial amplicon. To evaluate genome copy numbers, total DNA was isolated

from three to five individual snails from each lineage using a DNeasy Plant Mini Kit (Qiagen,

Valencia, CA), and nuclear and mitochondrial targets were amplified in triplicate in separate

reactions on the same plate, together with serial dilutions of the cloned standards, using

quantitative PCR on a StepOne Plus real-time thermal cycler (Applied Biosystems, Foster City,

CA). We converted quantitation cycle values (Cq, the PCR cycle at which amplification products

accumulated above a defined threshold) from snail samples into copy numbers using standard

curves generated from the cloned standards, as in Miller *et al.* (2003), and the ratio of

mitochondrial to haploid nuclear genome copies was determined for each sample; all lineages

examined were triploid, established by flow cytometric data. Because the ratio of *cytB* to *rad21*

fit the assumptions of normality (Shapiro-Wilks $W = 0.956$, $p = 0.228$) and variances between

lineages were not significantly different (Levene's $F = 0.632$, $p = 0.677$), we compared inferred

mitochondrial genome copy number across lineages using a one-way ANOVA and pairwise t-

tests as implemented in the R v 3.2.4 (R Core Team 2016).

Mitochondrial function at the organellar level

JC-1 assay

Mitochondrial membrane potential determines the strength of the electrochemical gradient

mitochondria use to phosphorylate ADP to ATP. Accordingly, the amplitude of membrane

potential is positively correlated with the ratio of ATP to ADP in the cell (Nicholls, 2004).

Variation in this mitochondrial membrane potential has been linked to cellular aging (Nicholls,

2004) and survival (Callegari *et al.,* 2011), making mitochondrial membrane potential a useful

measure of mitochondrial performance. The JC-1 assay (Garner and Thomas, 1999) measures the

strength of the electrochemical gradient in mitochondrial isolates using a fluorescent dye, JC-1.

With UV illumination, JC-1 will fluoresce green when dispersed and red when aggregated. In the

absence of an electrochemical gradient, JC-1 will diffuse randomly across membranes and will

aggregate in proportion to its concentration, such that under our experimental conditions, the

ratio of red: green fluorescence should be approximately 1. If isolated mitochondria establish and

maintain electrochemical gradients, JC-1, a small, positively charged molecule, will diffuse

down the gradient and accumulate inside mitochondrial matrices, yielding an elevated ratio of

red: green. A relatively high ratio of red: green is indicative of increased mitochondrial

membrane potential and thereby suggests increased mitochondrial function (Garner and Thomas,

1999). We used the uncoupler carbonyl cyanide m-chlorophenyl hydrazone (CCCP) to abolish

the electrochemical gradient across the mitochondrial inner membrane in replicate subsamples,

allowing us to control for background levels of fluorescence of JC-1 and of mitochondrial

membranes unrelated to mitochondrial function.

To test for genetic variation in mitochondrial function in *P. antipodarum* in terms of

mitochondrial membrane potential, we assayed 8-10 individual adult female snails from six

distinct asexual lineages representing a diverse subset of the natural mitochondrial haplotype

diversity found in New Zealand (Neiman *et al.,* 2011; Paczesniak *et al.,* 2013; Table S1). For

each snail, we first removed its shell and briefly washed the collected tissues by centrifugation at

600 x g in extraction buffer (10.0 mM HEPES (pH 7.5), 0.2 M mannitol, 70.0 mM sucrose, 1.0

mM EGTA). We rapidly homogenized these tissues on ice in extraction buffer containing 2

mg/ml fatty acid-free bovine serum albumin (fafBSA) using a micropestle and then centrifuged

the homogenate at 4ºC for 5 minutes at 600 x g. The supernatant was recovered and held on ice

separately while the pellet was re-homogenized and centrifuged again, as above. The pooled

mitochondrial-enriched supernatant was centrifuged at 12,000 x g for 10 minutes and the pellet

resuspended in buffer containing 10.0 mM HEPES (pH 7.5), 0.25 M sucrose, 1.0 mM ATP, 0.08

mM ADP, 5.0 mM sodium succinate, 2.0 mM K_2HPO_4, and 1mM DTT. We divided each sample

into three subsamples of 30 µl and added 500 µl assay buffer containing 20.0 mM MOPS (pH

7.5), 110.0 mM KCl, 10mM ATP, 10.0 mM $MgCl_2$, 10.0 mM sodium succinate, and 1.0 mM

EGTA. The first subsample was incubated with buffer alone, to monitor background

fluorescence, the second subsample was incubated with 2 µM JC-1, and the third with 2 µM JC-

1 and 30 µM CCCP. All three subsamples were incubated in the dark at 37ºC for 20 minutes,

after which the ratio of red: green fluorescence was determined using flow cytometry (Becton

Dickinson FacsCalibur, Franklin Lakes, NJ). Ungated data were collected from several hundred

to several thousand mitochondrial particles per sample, on forward scatter (FSC) and side scatter

(SSC), FL1 (green fluorescence, log scale), and FL2 (red fluorescence, log scale). We plotted

FL1 vs. FL2 for each sample, after gating out debris in FlowJo v 10.0.8 (FlowJo, Ashland, OR)

and derived the ratio of red to green for each particle. The median ratio for each sample run was

used for all comparisons.

Because red: green fluorescence ratios were not normally distributed (Shapiro-Wilks $W =$

0.858, $p = 7.131$ x 10^{-6}), we used a log transformation of the data to meet the normality

assumption of parametric analyses (Shapiro-Wilks $W = 0.969$, $p = 0.15$). We also found that

ratios of red: green across lineages did not have equal variances (Levene's $F = 3.093$, $p = 0.016$),

so we used a one-way ANOVA with a White adjustment for unequal variances (McKinnon and

White, 1985) and lineage as a random factor to determine whether lineage affected the dependent

variable of red: green ratio. We performed pairwise comparisons between lineages using

Welch's t-tests (to reflect unequal variances) and the Holm procedure for Bonferroni correction

(Holm 1979) using R v 3.2.4.

MTT assay

Electron flow through the electron transport chain (ETC) provides the energy necessary to

establish a proton gradient, such that increased flow should produce a corresponding increase in

mitochondrial membrane potential. When isolated mitochondria are incubated with the

compound MTT (3-(4,5-dimethylthizol-2-yl) diphenyltetrazolium bromide), MTT accepts

electrons from the ETC, forming a purple formazan product that is proportional to electron flow,

and is the basis of a colorimetric assay of electron transport (Liu *et al.,* 1997). We performed the

MTT assay by resuspending mitochondrial pellets pooled from 3-4 snails (obtained as described

above) in 100 µl buffer (125 mM KCl, 2 mM K_2HPO_4, 1 mM $MgCl_2$, and 20 mM HEPES,

adjusted to pH 7.4 with KOH) with 6 mM succinate as an energy substrate. We then added these

resuspended mitochondria to wells in 96-well plates. The MTT reaction was initiated by adding

10 µl of 2.5 mg/ml MTT to each well, and the plate was incubated for 2 hours at 37°C to allow

electrons from the ETC to reduce MTT. A 20% SDS 50% dimethylformamide solubilization

solution was applied to each well and the plate incubated overnight, after which the reduced

formazan product was measured as A_{570}, in a XL-800 microplate reader (Bio-Tek Instruments,

Winooski, VT). We determined background absorbance from duplicate mitochondrial samples

incubated without an energy substrate and subtracted this background value from all readings.

Phosphate-buffered saline was used as a negative control and 0.1 mM dithiothreitol as a positive

control for MTT reduction. A fraction of each original mitochondrial sample was lysed in SDS

and then used in a bicinchoninic acid assay (Smith *et al.,* 1985) to determine protein

concentration. MTT reduction was expressed as A_{570}/ug protein.

While MTT values did not fit the assumptions of a normal distribution (Shapiro-Wilks W

$= 0.840$, $p = 2.436 \times 10^{-5}$), these values did meet the assumptions of normality following a log

transformation (Shapiro-Wilks $W = 0.961$, $p = 0.140$). Log-transformed MTT values did not

display unequal variances (Levene's $F = 0.964$, $p = 0.452$), allowing us to compare log-

transformed MTT values using a one-way ANOVA with lineage as a random factor to test

whether different asexual lineages exhibited different levels of electron flow through the ETC (as

implemented in R v 3.2.4).

Mitochondrial function at the organismal level

Ectotherms respire at higher rates at elevated temperatures, such that O_2 becomes limiting as

temperature increases (Abele *et al.,* 2007). Accordingly, heat stress can reveal inefficiencies in

ATP production (Heise *et al.,* 2003). By this logic, we hypothesized that measuring O_2

consumption under heat stress would increase our ability to detect existing genetic variation for

mitochondrial function. There is some evidence from earlier studies that *P. antipodarum* are

susceptible to heat stress: at elevated temperature, snails exhibit decreased fecundity (Dybdahl

and Kane, 2005) and tend to consume more O_2 (Hudson, 1983).

To further define the range of heat stress likely to alter mitochondrial function in *P.*

antipodarum, we used a behavioral assay that measures the amount of time that a snail takes to

right itself when placed ventral side-up and compared righting ability in 13 asexual lineages (N =

10 individuals per lineage) across three temperature treatments (16ºC, 22ºC, 30ºC). Righting

ability is a commonly used method to gauge levels of snail stress (for example, Orr *et al.,* 2007).

For example, snails exposed to hypoxic conditions exhibit increased righting time (and elevated

HSP70 gene expression) compared to unexposed snails (Fei *et al.,* 2008). We began by

incubating adult female *P. antipodarum* in carbon-filtered tap water (that is, the water in which

the snails are housed) at the test temperature for 1 hour. Next, we placed snails ventral side-up in

a petri dish and measured the number of seconds elapsed until the snail righted itself, up to 180

seconds. We compared righting time across temperatures and lineages with a Kruskal-Wallis test

(as implemented in R v 3.2.4) and found that both temperature (Kruskal-Wallis, $\chi^2 = 14.218$, df

$= 2, p = 0.00082$) and lineage (Kruskal-Wallis, $\chi^2 = 122.64$, df $= 12, p < 2.2$ x 10^{-16}) significantly

affected righting time. Righting time was ~24% faster at 22ºC (mean (+/- SD) = 48.82 (+/-

56.09) seconds) than at 16ºC (mean (+/- SD) = 62.50 (+/- 63.26) seconds), but this difference

was not significant (Mann-Whitney, $U = 9244.5, p = 0.0754$). Conversely, righting took ~ 37%

longer at 30ºC (mean (+/- SD) = 83.02 (+/- 73.74) seconds) than at 16ºC (Mann-Whitney U =

8192, $p = 0.0254$) and ~85% longer at 30ºC than at 22ºC (Mann-Whitney, $U = 5996.5, p =$

0.00014, Figure S1). These results are consistent with other righting time assays from other snail

species in that some amount of stress decreases righting time (for example, presence of predators,

Orr *et al.,* 2007), and some stimuli increase righting time (for example, hypoxia, Fei *et al.,* 2008),

suggesting that our observed responses indicate stress in *P. antipodarum* exposed to elevated

temperature (here, 30ºC).

After determining that snails do appear to be stressed by elevated temperatures, we tested

whether there exists genetic variation for O_2 consumption under heat stress in *P. antipodarum* by

performing closed-system aquatic respirometry on seven asexual lineages of *P. antipodarum* (N

= 10 per lineage) at three different incubation temperatures (16ºC, 22ºC, 30ºC) using a

Strathkelvin Instruments RC200a respiration chamber, 892 Oxygen Meter, and a 1302 Clark-

type oxygen electrode (Strathkelvin Instruments, Motherwell, UK). We calibrated the electrode

daily using the solubility of O_2 at each respective temperature (16ºC – 309.0 µmol/liter, 22ºC –

279.0 µmol/liter, 30ºC – 236.0 µmol/liter). A high calibration point was obtained by stirring

carbon-filtered water vigorously for 30 minutes prior to calibration, while we used a 2% sodium

sulfite solution as a low calibration standard. We incubated each snail at the prescribed

temperature for one hour prior to measurement, placed snails into the cell chamber, and obtained

O_2 concentration readings for each snail every second for 1 hour. We maintained constant

temperature in the respiration chamber by pumping temperature-controlled water into the cell

chamber's water jacket. Upon completion of the 1-hour test period, we blotted each snail dry and

measured its wet mass on a Denver Instruments Cubis Analytical Balance (Denver Instrument,

Bohemia, NY).

Snail wet mass was significantly and positively correlated with O_2 consumption

(Spearman's rho = 0.17, p = 0.016, Figure S2). Because there is significant variation for snail

wet mass between asexual lineages (Kruskal-Wallis, $\chi^2 = 76.82$, df = 6, $p < 0.0001$), we

calculated the residuals of wet mass *vs*. O_2 consumption using a linear model. Square-root-

transformed O_2 consumption residuals were not significantly different from a normal distribution

(Shapiro-Wilks, $W = 0.9894$, $p = 0.14$), allowing us to implement a linear regression model to

test whether temperature and/or lineage affected O_2 consumption. We then used a two-way

ANOVA to address whether the fixed factor of temperature, the random factor of lineage, and

the interaction between temperature and lineage affected the dependent variable of wet-mass-

corrected O_2 consumption.

Comparison of mitochondrial functional assays

Because each of the three assays that we developed and employed is designed to reflect some

aspect of mitochondrial function, we should in principle expect some correlations between

different assays if the assays in question actually provide meaningful measures of mitochondrial

function. In particular, mitochondrial genome copy number measures overall system capacity for

ATP production, the JC-1 assay measures the strength of the electrochemical gradient, the MTT

assay measures electron flow though the OXPHOS pathway, and O_2 consumption under heat

stress measures the overall electron acceptor turnover in conditions when ATP is in demand. To

test for associations between different mitochondrial functional assays, we performed all 15

possible pairwise comparisons of mitochondrial functional assays by comparing the mean trait

value per lineage across assays using Spearman's rank correlation (as implemented by the *Hmisc*

package for R v 3.2.4), correcting for multiple comparisons using the Holm procedure. For these

analyses, we included two additional sample populations that were not used in any of the tests

for genetic variation: one field-collected sample from a lake with a high frequency of sexual

individuals ("KnSF12") and an inbred sexual lineage that has been maintained in the lab for >20

generations ("Y2", Table S1).

RESULTS

Mitochondrial function at the genomic level

To test whether *P. antipodarum* exhibits heritable variation for mtDNA copy number, we

compared qPCR amplification of the mitochondrially encoded *cytB* locus to amplification of a

putatively single copy nuclear gene, *rad21*, in six asexual lineages of *P. antipodarum* reared in a

common-garden setting (N = 3-8 per lineage). The mean (+/- SD) number of *cytb* copies to the

number of *rad21* copies was 13.715 (+/- 3.571), though we also found significant differences in

this ratio across asexual lineages (one-way ANOVA, $F_{5, 25} = 2.72$, $p < 0.0428$, Figure 1),

indicating that lineages differ in mitochondrial genome copy number. We performed pairwise t-

tests of copy number ratio among each pair of lineages and found that a single lineage (Gr5;

mean copy number ratio = 9.423, SD = 1.483) with very low mitochondrial copy number

appeared to be driving the significant among-lineage variance. To test if the lack of differences

amongst the remaining lineages could be an artifact of low statistical power, we performed

power analyses using the *pwr* package in R v 3.2.4 for each of the 28 pairwise comparisons. We

found mean power (+/- SD) = 0.402 (+/- 0.398), indicating both that power was low to moderate

but also that power varied substantially across comparisons. For all comparisons involving the

Gr5 lineage, mean power (+/- SD) = 0.881 (+/- 0.224), while for all comparisons not involving

Gr5, mean power (+/- SD) was 0.162 (+/- 0.183). This result is likely driven by the low

mitochondrial genome copy number of Gr5 relative to the other lineages and suggests that

sample size was likely too small to detect the more minor differences in copy number, if any,

that might exist between other asexual lineages.

Mitochondrial function at the organellar level

JC-1

Membrane potentials in mitochondria isolated from six asexual lineages of *P. antipodarum* were

compared using the JC-1 assay, in which the ratio of red to green fluorescence is indicative of

membrane potential. We found significant differences in the log-transformed ratios of red: green

across asexual lineages (Welch's one-way ANOVA, $F_{5, 52} = 6.628$, $p = 7.671$ x 10^{-5}, Figure 2a).

Pairwise *posthoc* comparisons revealed three significantly different groups ($p < 0.05$ after

correcting for multiple comparisons) amongst the six lineages (Figure 2a). Unlike mitochondrial

genome copy number, this variation in mitochondrial membrane potential does not appear to be

driven by a single lineage. Because CCCP-treated samples did not vary across lineages (one-way

ANOVA, $F_{5,52} = 1.367$, $p = 0.252$), the variation observed for red: green fluorescence across

lineages likely reflects variation in mitochondrial membrane potential. In particular, these data

suggest that under the current rearing and assay conditions, the Gr5 lineage appears to have the

strongest mitochondrial membrane potential (mean (+/- SD) = 3.859 (+/- 1.546)) and the DenA

lineage (mean (+/- SD) = 1.775 (+/- 0.468)) appears to have the weakest mitochondrial

membrane potential.

MTT

MTT reduction by isolated mitochondria is due to acceptance of electrons from the electron

transport chain of the inner membrane, meaning that the extent of MTT reduction should reflect

electron flow through the OXPHOS pathway. In general, the outcome of the MTT assay would

be expected to be correlated with the strength of the proton gradient measured in the JC-1 assay,

although that gradient can also be affected by the presence of uncoupling proteins, which reduce

the gradient by letting protons leak back into the matrix, dissipating energy as heat. Additionally,

MTT does not accept electrons from all the reactions in the electron transport chain, so that some

complex-specific differences in function may not be represented. Despite clear evidence for

across-lineage variation in JC-1, we did not detect any differences amongst asexual lineages in

MTT reduction (one-way ANOVA, $F_{5, 38} = 0.751$, $p = 0.591$, Figure 2b), even though we had

relatively high power (=0.92, using the pwr.anova.test function implemented in the *pwr* package

in R v 3.2.4).

Mitochondrial function at the organismal level

Organismal O_2 consumption under stressful conditions is expected to reflect electron acceptor

turnover under maximal ATP production (Abele *et al.,* 2007). To test for genetic variation in O_2

consumption in heat-stressed *P. antipodarum*, we performed closed-system aquatic respirometry

for seven asexual lineages (N = 10 per lineage) at 16ºC, 22ºC, and 30ºC. We found that elevated

temperatures significantly affected mass-corrected O_2 consumption residuals (two-way ANOVA,

$F_{1, 191} = 79.118$, $p = 4.44 \times 10^{-16}$) and that there was a significant interaction between lineage and

temperature (Figure 3, two-way ANOVA, $F_{1, 191} = 3.919$, $p = 0.00102$). Using a one-way

ANOVA within each temperature treatment, we found that lineage had a significant effect on O_2

consumption at 22ºC (one-way ANOVA, $F_{6, 58} = 3.002$, $p = 0.0127$) and at 30ºC (one-way

ANOVA, $F_{6, 62} = 3.823$, $p = 0.00263$), but not at 16ºC (one-way ANOVA, $F_{6, 64} = 1.892$, $p =$

0.096). In particular, lineages responded differently to elevated temperature, with some lineages

(for example, Ta10) exhibiting relatively high O_2 consumption at high temperatures and others

maintaining similar levels of O_2 consumption (for example, Gn5) across temperatures. This

result demonstrates that genetically distinct lineages of *P. antipodarum* consume different

amounts of O_2 in response to heat stress.

Comparison of mitochondrial functional assays

To determine the extent to which our different measures of mitochondrial function are associated,

we performed Spearman's rank correlation for each of the 15 possible pairwise comparisons of

the assays for mitochondrial genome copy number, log-transformed JC-1 red: green ratios, log-

transformed MTT values, and square-root-transformed O_2 consumption at each temperature.

None of the correlations remained significant after the Holm correction for multiple comparisons.

The absence of significant relationships between assay results is not surprising in light of the fact

that we had power of ~ 0.35 – 0.65 (using the pwr.r.test function implemented in the *pwr*

package in R v 3.2.4) to detect a relationship between two variables at the $p < 0.003$ level, the

alpha required by the Holm procedure. Power notwithstanding, there were some interesting

results that deserve further scrutiny. First, neither mitochondrial genome copy number nor O_2

consumption values for 16º C or 30º C were correlated with any other phenotypic measure

(Figure 4). Second, while the MTT assay requires pooling of mitochondria from 3-4 individual

snails per measurement and is thus somewhat less sensitive than JC-1, which only requires

mitochondria from one snail, we did find that MTT reduction was positively correlated with JC-1

measures of the proton gradient in the six asexual and two sexual *P. antipodarum* lineages

examined (Spearman's rho = 0.81, $p = 0.0149$, Figure 4f). There was a trend towards a negative

correlation between mean log-transformed JC-1 values and O_2 consumption at 22ºC for the six

asexual lineages (Spearman's rho = -0.77, $p = 0.0724$, Figure 4k), and mean log-transformed

MTT values were negatively correlated with O_2 consumption at 22ºC for the six asexual lineages

(Spearman's rho = -0.89, p = 0.0188, Figure 4k). Together, these tentative relationships between

the MTT assay (reflecting OXPHOS electron flow) and O_2 consumption and between the MTT

assay and the JC-1 assay (reflecting mitochondrial membrane potential) suggest that high

mitochondrial membrane potential is expected to result in low O_2 consumption under O_2-limiting

conditions in *P. antipodarum*. These results also suggest that genome copy number, JC-1, and

organismal O_2 consumption provide orthogonal lines of inquiry into mitochondrial function.

DISCUSSION

Genetic variation for mt function at three different levels of biological organization

We used obligately asexual lineages of *Potamopyrgus antipodarum*, a freshwater New Zealand

snail, to test for genetic variation in mitochondrial function in a common-garden setting. We

found significant levels of variation at all three levels of biological organization that we assayed:

mitochondrial genome copy number, mitochondrial membrane potential, and variation in O_2

consumption in response to heat stress. These results demonstrate that substantial levels of

variation for mitochondrial function exist in asexual lineages of this species and suggest that

mitochondrial function at the organellar level is related to mitochondrial function at the

organismal level. Importantly, the methods described here can be easily adapted to other non-

model organisms, especially mollusks, providing a new means of quantifying genotype-

phenotype relationships.

The extent to which this variation in mitochondrial functional phenotypes contributes to

fitness in *P. antipodarum* remains to be directly evaluated. Even so, the substantial across-

lineage variation that we discovered provides functional evidence of high levels of asexual

phenotypic diversity in *P. antipodarum*, consistent with previous reports that asexual *P.*

antipodarum harbor substantial genetic (Jokela *et al.,* 2003; Paczesniak *et al.*, 2013) and

phenotypic (Kistner and Dybdahl, 2013) diversity. The multiple separate transitions to asexuality

in *P. antipodarum* (Neiman *et al.,* 2011; Paczesniak *et al.,* 2013) may help explain the substantial

levels of variation found here, as asexual lineages represent "snapshots" of local sexual

population diversity. Thus, variation within asexual lineages should be a subset of the variation

found in source sexual populations (for example, Jokela *et al.,* 1997). The mitochondrial

phenotypes we observed in one lineage, Gr5, were particularly distinct: snails from this lineage

exhibited relatively low mitochondrial copy number, high mitochondrial membrane potential and

electron flow, and low O_2 consumption at 22ºC. Together, these phenotypic values suggest that

the Gr5 lineage exhibits relatively high mitochondrial function, indicating that genetic dissection

of its mitochondrial haplotype may prove illuminating. Further comparisons with other *P.*

antipodarum lineages and in other conditions will provide substantial insight into the relative

fitness of this particular mitonuclear combination. Future studies should also focus on a

particular mitochondrial haplotype that appears to be especially common amongst asexual *P.*

antipodarum (Neiman *et al.,* 2011; Paczesniak *et al.,* 2013). Paczesniak *et al.* (2013) showed that

this haplotype is often found in divergent nuclear backgrounds, consistent with a scenario where

this haplotype is spreading into new populations and lineages. Evaluating mitochondrial function

of the common mitochondrial haplotype in *P. antipodarum* with divergent vs. similar nuclear

backgrounds and in various biologically relevant conditions would shed light onto intraspecific

mitonuclear coevolution in the context of asexuality and on whether selection is effective at

favoring epistatic combinations in the absence of sex.

That the phenotypic variation for mitochondrial function shown here resides in traits that

have been demonstrated to influence fitness in other organisms (Chen *et al.,* 2007; Dowling,

2014) means that this variation could have major implications for asexual lineage success. In

particular, asexual *P. antipodarum* are known to harbor mitochondrial genomes with high

mutational loads relative to sexual lineages (Neiman *et al.,* 2010; Sharbrough *et al.,* 2016),

suggesting that accelerated mutation accumulation in asexual lineages likely affects

mitochondrial function. Whether the mutations present in asexual mitochondrial genomes

negatively impact fitness and/or function remains an unanswered question of great interest, and

comparing mitochondrial function in *P. antipodarum* across reproductive modes will provide

critical tests of the hypothesis that mitochondrial mutation accumulation in asexuals is

deleterious.

Relationship between mitochondrial functional assays

All else being equal, stronger membrane potentials, greater electron flow, lower O_2 consumption,

and higher mitochondrial genome copy number should all positively correlate with overall

mitochondrial function. Because all else is almost certainly not equal (for example, variation in

temperature/resources in native range, phylogenetic constraint, *etc*.), it is possible that high

mitochondrial performance in common-garden conditions does not reflect mitochondrial

performance in nature. Despite this caveat, the tools developed here will provide an important

starting point by which to interrogate mitochondrial function in non-model systems.

Although the number of asexual lineages included in the present study was relatively

small, we did observe some interesting relationships between mitochondrial genome copy

number, electron flux through the ETC, mitochondrial membrane potential, and organismal O_2

consumption (Figure 4). In particular, we found an inverse relationship between O_2 consumption

at 22ºC and electron flow through the ETC and a positive relationship between electron flow

through the ETC and mitochondrial membrane potential. The latter relationship is not

particularly surprising in light of the fact that as electrons pass through the ETC, we expect a

corresponding increase in mitochondrial membrane potential as H+ ions are released into the

intermembrane space. The inverse relationship between decreased electron flux and elevated O_2

consumption is more difficult to understand because O_2 consumption is expected to increase with

electron flux. Importantly, correlations between organismal-level O_2 consumption and organelle-

level electron flux are difficult to disentangle because many layers of respiratory regulation can

contribute to increased organismal O_2 consumption beyond ETC inefficiencies (Brand and

Nicholls, 2011). Furthermore, electrons leaked back across the inner membrane though

uncoupling proteins contribute to electron flux but not O_2 consumption, and elevated

temperatures in a marine mollusk have been shown to increase electron leakage of this type, as

well as O_2 consumption in the formation of reactive oxygen species (Abele *et al*., 2002), such

that connecting organismal O_2 consumption to organellar electron flux is a non-trivial exercise.

Understanding the relationship between organellar and organismal variation in mitochondrial

function would provide helpful context for interpreting functional variation in *P. antipodarum*,

and future efforts towards examining these relationships in more detail are necessary.

Implications for the loss of sex

One of the primary hypotheses put forth as an explanation for the prevalence of sexual

reproduction in eukaryotes is the accumulation of slightly deleterious mutations in asexual

lineages, eventually causing mutational meltdown and lineage extinction (Neiman and Taylor,

2009). Empirical evidence in support of this hypothesis remains largely indirect: (1) the

relatively "twiggy" distribution of asexual taxa on the eukaryotic phylogeny compared to sexual

taxa (Williams, 1975), and (2) the accumulation of putatively deleterious mutations in nuclear

(Tucker *et al.,* 2013; Hollister *et al.,* 2015) and organelle genomes of asexual lineages (Neiman

et al., 2010; Henry *et al.,* 2012; Sharbrough *et al.,* 2016). What remains unresolved is whether

the mutations being accumulated by asexual lineages actually decrease function and/or fitness.

Establishing the presence of genetic variation for mitochondrial function represents the first step

in determining whether mitochondrial mutation accumulation in the absence of sex causes such a

decline. More specifically, if mitochondrial haplotype is not linked to mitochondrial phenotype,

mutation accumulation in mitochondrial genomes in asexual lineages would not cause decreases

in function, such that sexual reproduction must be associated with other benefits to persist in the

face of asexual competitors. Here, we show that snails with different mitochondrial genetic

backgrounds have different mitochondrial phenotypes in a common-garden environment,

suggesting that mutations present in certain lineages have some impact on function, thereby

setting the stage for direct comparisons between mitochondrial function in sexual and asexual *P.*

antipodarum.

One alternative possibility to decreased mitochondrial function in asexual lineages is that

asexuals have access to genetic variation that sexuals do not. This scenario is not wholly

unexpected: because the nuclear and mitochondrial genomes are transmitted in complete LD in

asexuals, advantageous combinations of mutations (that is, epistatic variation) can become

visible to selection (Neiman and Linksvayer, 2006). By contrast, in sexual lineages, the

mitochondrial genome is swapped across diverse nuclear backgrounds each generation, such that

selection cannot act effectively on phenotypes linked to epistatic mitonuclear interactions. The

implications are that interacting nuclear and mitochondrial subunits may experience particularly

tight and effective coevolution. Such a phenomenon might exhibit a very similar pattern of

molecular evolution to that of accelerated deleterious mutation accumulation, meaning that

testing the functional consequences of asexual mutation accumulation is critical to determining

whether the absence of sex inevitably leads to mutation meltdown and lineage extinction.

DATA ARCHIVING

All data will be made available on Dryad upon acceptance.

CONFLICT OF INTEREST

The authors declare no conflict of interest for the work described here.

FUNDING

This work was supported by the National Science Foundation (NSF: MCB – 1122176; DEB – 1310825) and the Iowa Academy of Sciences (ISF #13-10).

ACKNOWLEDGEMENTS

We thank JD Woodell for assistance with aquatic respirometry. We thank Claire Tucci, Meagan Luse, and Nikhil Puttagunta for their help measuring snail righting behavior.

TITLES AND LEGENDS TO FIGURES

Figure 1. MtDNA copy number variation in six asexual lineages of *P. antipodarum*. Box-

and-whisker plot depicting quantitative PCR estimates (rank-ordered by median) of *cytB* copy

number relative to a putatively single-copy nuclear gene, *rad21*. MtDNA copy number was

significantly different across lineages (one-way ANOVA, $F = 2.72$, df = 5, $p = 0.043$), which

appears to be driven by lineage Gr5. Boxes represent Inner Quartile Ranges (IQR) with error

bars extending 1.5x beyond IQRs. Data points falling outside error bars are denoted by open

circles. Shared lowercase letters indicate $p > 0.05$ for pairwise t-tests corrected for multiple

comparisons using the Holm procedure for multiple comparisons. N = 10 individuals for all

lineages.

Figure 2. Organellar function of mitochondrial fractions isolated from six lineages of *P.*

***antipodarum*.** a) Estimate of mitochondrial membrane potential using the ratio of red to green

JC-1 dye fluorescence from individual snails (N = 10 individuals for all lineages). Lineages were

rank-ordered by median with boxes representing Inner Quartile Ranges (IQR); error bars extend

1.5x beyond IQRs. Data points falling outside error bars are denoted by open circles. Shared

lowercase letters indicate $p > 0.05$ for pairwise Welch's T-tests, corrected using the Holm

procedure for multiple comparisons. b) Estimate of electron flux through OXPHOS pathway

using the colorimetric MTT reduction assay from pooled mitochondrial extractions (3 snails

pooled per replicate, 3 – 6 replicates per lineage). Lineages were rank-ordered by median value

with boxes representing IQRs and error bars extending 1.5x beyond IQRs. Data points falling

outside error bars are denoted by open circles. Lineages did not appear to differ in MTT

reduction.

Figure 3. Interaction plot between O_2 consumption residuals, temperature, and snail lineage. Lines indicate best-fit linear regression of O_2 consumption across temperature pairs (for example, 16ºC – 22ºC) for seven asexual lineages of *P. antipodarum.* O_2 consumption was measured for 10 individual snails at each temperature for each lineage. O_2 consumption is affected by temperature as well as by the interaction between temperature and lineage. Within temperatures, lineage had a significant effect on O_2 consumption at 22ºC (one-way ANOVA, $F_{6, 58} = 3.002$, $p = 0.0127$) and at 30ºC (one-way ANOVA, $F_{6, 62} = 3.823$, $p = 0.00263$).

Figure 4. Comparisons of mitochondrial functional assays. Spearman's rank correlations for all pairwise comparisons of mitochondrial functional assays. Best-fit linear regression lines (black) are only shown for those comparisons that were significant at the $p < 0.05$ level. a) mtDNA copy number vs. JC-1 (Spearman's rho = -0.1, $p = 0.823$). b) mtDNA copy number vs. MTT reduction (Spearman's rho = -0.33, $p = 0.42$). c) mtDNA copy number vs. O_2 consumption at 16ºC (Spearman's rho = 0.03, $p = 0.957$). d) mtDNA copy number vs. O_2 consumption at 22ºC (Spearman's rho = 0.03, $p = 0.957$). e) mtDNA copy number vs. O_2 consumption at 30ºC (Spearman's rho = 0.43, $p = 0.396$). f) JC-1 vs. MTT reduction (Spearman's rho = 0.81, $p = 0.0149$). g) JC-1 vs. O_2 consumption at 16ºC (Spearman's rho = -0.43, $p = 0.397$). h) JC-1 vs. O_2 consumption at 22ºC (Spearman's rho = -0.77, $p = 0.0724$). i) JC-1 assay vs. O_2 consumption at 30ºC (Spearman's rho = 0.2, $p = 0.704$). j) MTT reduction vs. O_2 consumption at 16ºC (Spearman's rho = 0.14, $p = 0.787$). k) MTT reduction vs. O_2 consumption at 22ºC (Spearman's rho = -0.89, $p = 0.0188$). l) MTT reduction vs. O_2 consumption at 30ºC (Spearman's rho = -0.09, $p = 0.872$). m) O_2 consumption at 16ºC vs. O_2 consumption at 22ºC (Spearman's rho = -0.36, $p =$

600   0.432). n) O_2 consumption at 16ºC vs. O_2 consumption at 30ºC (Spearman's rho = -0.14, p =

601   0.76). o) O_2 consumption at 22ºC vs. O_2 consumption at 30ºC (Spearman's rho = -0.04, p =

602   0.939).

603 **SUPPLEMENTARY MATERIAL TITLES TO FIGURES AND LEGENDS**
604

Figure S1. Variation in righting time in response to heat stress in *P. antipodarum*. Ten

asexual lineages (N = 10 snails per lineage) were incubated at either 16ºC, 22ºC, or 30ºC for 1

hour prior to assaying behavioral responses to elevated temperature. Temperature (Kruskal-

Wallis, χ^2 = 14.218, df = 2, p = 0.00082) and lineage (Kruskal-Wallis, χ^2 = 122.64, df = 12, $p <$

2.2×10^{-16}) both significantly affected righting time, indicating that elevated temperature can be

used to induce stress in *P. antipodarum*.

Figure S2. Relationship between snail wet mass and oxygen consumption in *P. antipodarum*.

There was a significant positive correlation between snail wet mass and O_2 consumption

(Spearman's rho = 0.2, p = 0.0044).

REFERENCES

Abele D, Heise K, Portner HO, Puntarulo S (2002). Temperature-dependence of mitochondrial

function and production of reactive oxygen species in the intertidal mud clam *Mya*

arenaria. *J Exp Biol* **205:** 1831-1841.

Barr CM, Neiman M, Taylor DR (2005). Inheritance and recombination of mitochondrial

genomes in plants, fungi and animals. *New Phytol* **168:** 39-50.

Brand, M. D., & Nicholls, D. G. (2011). Assessing mitochondrial dysfunction in cells.

Biochem J **435**: 297-312.

Callegari S, McKinnon RA, Andrews S, de Barros Lopes MA (2011). The MEF2 gene is

essential for yeast longevity, with a dual role in cell respiration and maintenance of

mitochondrial membrane potential. *FEBS Lett* **585:** 1140-1146.

Chance B, Williams GR (1955). Respiratory enzymes in oxidative phosphorylation. I. Kinetics

of oxygen utilization. *J Biol Chem* **217:** 383-393.

Chen JH, Hales CN, Ozanne SE (2007). DNA damage, cellular senescence and organismal

ageing: causal or correlative? *Nucleic Acids Res* **35:** 7417-7428.

DiMauro S, Schon EA (2001). Mitochondrial DNA mutations in human disease. *Am J Med*

Genet **106:** 18-26.

Dowling DK (2014). Evolutionary perspectives on the links between mitochondrial genotype and

disease phenotype. *Biochim Biophys Acta* **1840:** 1393-1403.

Dybdahl MF, Kane SL (2005). Adaptation vs. phenotypic plasticity in the success of a

clonal invader. *Ecology,* **86:** 1592-1601.

Ellison CK, Burton RS (2006). Disruption of mitochondrial function in interpopulation hybrids

of *Tigriopus californicus*. *Evolution* **60:** 1382-1391.

Fei G, Guo C, Sun HS, Feng ZP (2008). HSP70 reduces chronic hypoxia-induced neural

suppression via regulating expression of syntaxin. In: Poulin MJ, Wilson RJA (eds)

Integration in Respiratory Control, Springer: New York, pp. 35-40.

Garner DL, Thomas CA (1999). Organelle-specific probe JC-1 identifies membrane potential

differences in the mitochondrial function of bovine sperm. *Mol Repro Dev* **53:** 222-229.

Heise K, Puntarulo S, Portner HO, Abele D (2003). Production of reactive oxygen species by

isolated mitochondria of the Antarctic bivalve *Laternula elliptica* (King and Broderip)

under heat stress. *Comp Biochem Physiol C Toxicol Pharmacol* **134:** 79-90.

Henry L, Schwander T, Crespi BJ (2012). Deleterious mutation accumulation in asexual *Timema*

stick insects. *Mol Biol Evol* **29:** 401-408.

Hollister JD, Greiner S, Wang W, Wang J, Zhang Y, Wong GK, *et al.* (2015). Recurrent loss of

sex is associated with accumulation of deleterious mutations in *Oenothera*. *Mol Biol Evol*

32: 896-905.

Hori A, Yoshida M, Shibata T, Ling F (2009). Reactive oxygen species regulate DNA copy

number in isolated yeast mitochondria by triggering recombination-mediated replication.

Nucl Acids Res **37:** 749-761.

Hudson JAL (1983). Oxygen consumption of the mollusc *Potamopyrgus antipodarum*

in relation to habitat. Ph.D. Thesis. University of Canterbury.

Jokela J, Lively CM, Fox JA, Dybdahl MF (1997) Flat reaction norms and "frozen" phenotypic

variation in clonal snails (*Potamopyrgus antipodarum*). *Evolution* **51:** 1120-1129.

Kelly RD, Mahmud A, McKenzie M, Trounce IA, St John JC (2012). Mitochondrial DNA copy

number is regulated in a tissue specific manner by DNA methylation of the nuclear-

encoded DNA polymerase gamma A. *Nucl Acids Res* **40:** 10124-10138.

Lee-Yaw JA, Jacobs CG, Irwin DE (2014). Individual performance in relation to cytonuclear

discordance in a northern contact zone between long-toed salamander (*Ambystoma*

macrodactylum) lineages. *Mol Ecol* **23:** 4590-4602.

Liu Y, Peterson DA, Kimura H, Schubert D (1997) Mechanism of cellular 3-(4,5-

dimethylthiazo-2-yl)-2,5-diphenyltetrazolium bromide (MTT) reduction. *J Neurochem*

69: 581-593.

MacKinnon JG, White H (1985). Some heteroskedasticity-consistent covariance matrix

estimators with improved finite sample properties. *J Econometr* **29:** 305-325.

Matsushima Y, Goto Y, Kaguni LS (2010). Mitochondrial Lon protease regulates mitochondrial

DNA copy number and transcription by selective degradation of mitochondrial

transcription factor A (TFAM). *Proc Natl Acad Sci USA* **107:** 18410-18415.

Meiklejohn CD, Holmbeck MA, Siddiq MA, Abt DN, Rand DM, Montooth KL (2013). An

incompatibility between a mitochondrial tRNA and its nuclear-encoded tRNA synthetase

compromises development and fitness in *Drosophila*. *PLoS Genet* **9:** e1003238.

Miller FJ, Rosenfeldt FL, Zhang C, Linnane AW, Nagley P (2003). Precise determination of

mitochondrial DNA copy number in human skeletal and cardiac muscle by a PCR-based

assay: lack of change of copy number with age. *Nucleic Acids Res* **31:** e61.

Montier, L. L. C., Deng, J. J., & Bai, Y. (2009). Number matters: control of mammalian

mitochondrial DNA copy number. *J Genet Genom*, **36**: 125-131.

Neiman M, Hehman G, Miller JT, Logsdon Jr JM, Taylor DR (2010). Accelerated mutation

accumulation in asexual lineages of a freshwater snail. *Mol Biol Evol* **27:** 954-963.

Neiman M, Linksvayer TA. (2006). The conversion of variance and the evolutionary potential of

restricted recombination. *Heredity* **96:** 111-121.

Neiman M, Paczesniak D, Soper DM, Baldwin AT, Hehman G (2011). Wide variation in ploidy

level and genome size in a New Zealand freshwater snail with coexisting sexual and

asexual lineages. *Evolution* **65:** 3202-3216.

Neiman M, Taylor DR (2009). The causes of mutation accumulation in mitochondrial genomes.

Proc Roy Soc Lond B **276:** 1201-1209.

Nicholls DG (2004). Mitochondrial membrane potential and aging. *Aging Cell* **3:** 35-40.

Paczesniak D, Jokela J, Larkin K, Neiman M (2013). Discordance between nuclear and

mitochondrial genomes in sexual and asexual lineages of the freshwater snail P*otamopyrgus*

antipodarum. *Mol Ecol* **22:** 4695-4710.

Pichaud N, Ballard JW, Tanguay RM, Blier PU (2013). Mitochondrial haplotype divergences

affect specific temperature sensitivity of mitochondrial respiration. *J Bioenerget*

Biomembr **45:** 25-35.

Rand DM, Haney RA, Fry AJ (2004). Cytonuclear coevolution: the genomics of cooperation.

Trends Ecol Evol **19:** 645-653.

Rawson PD, Burton RS (2002). Functional coadaptation between cytochrome c and cytochrome

c oxidase within allopatric populations of a marine copepod. *Proc Natl Acad Sci USA* **99:**

12955-12958.

Sharbrough J, Luse M, Boore JL, Logsdon Jr JM, Neiman M (2016). Radical changes persist

longer in the absence of sex. *bioRχiv.* doi: http://dx.doi.org/10.1101/049924.

Sloan DB, Triant DA, Wu M, Taylor DR (2014). Cytonuclear interactions and relaxed selection

accelerate sequence evolution in organelle ribosomes. *Mol Biol Evol* **31:** 673-682.

Smith PK, Krohn RI, Hermanson GT, Mallia AK, Gartner FH, Provenzano MD, *et al.* (1985).

Measurement of protein using bicinchoninic acid. *Anal Biochem* **150:** 76–85.

Tucker AE, Ackerman MS, Eads BD, Xu S, Lynch M (2013). Population-genomic insights into

the evolutionary origin and fate of obligately asexual *Daphnia pulex*. *Proc Natl Acad Sci*

USA **110:** 15740-15745.

Van den Bogert C, De Vries H, Holtrop M, Muus P, Dekker HL, Van Galen MJ, et al (1993).

Regulation of the expression of mitochondrial proteins: relationship between mtDNA

copy number and cytochrome-c oxidase activity in human cells and tissues. *Biochim*

Biophys Acta **1144:** 177-183.

Williams GC (1975). *Sex and evolution*. Princeton University Press: Princeton.

Zachar, N., and M. Neiman. 2013. Profound effects of population density on fitness-

related traits in an invasive freshwater snail. PLoS One **8**:e80067.

Zhang F, Broughton RE (2013). Mitochondrial-nuclear interactions: compensatory

evolution or variable functional constraint among vertebrate oxidative phosphorylation

genes? *Gen Biol Evol* **5:** 1781-1791.