Striatal action-value neurons reconsidered

Lotem Elber-Dorozko[1] and Yonatan Loewenstein[1,2]

[1]The Edmond & Lily Safra Center for Brain Sciences, The Hebrew University of Jerusalem.

[2]Department of Neurobiology, The Alexander Silberman Institute of Life Sciences and the

Federmann Center for the Study of Rationality, The Hebrew University of Jerusalem.

Corresponding author and lead contact: Lotem.elber@mail.huji.ac.il

Abstract

It is generally believed that during economic decisions, striatal neurons represent the values

associated with different actions. This hypothesis is based on a large number of studies, in

which the neural activity of striatal neurons was measured while the subject was learning to

prefer the more rewarding action. Here we show that these publications are subject to at

least one of two critical confounds and that most are subject to both. First, we show that even

weak temporal correlations in the neuronal data may result in an erroneous identification of

action-value representations. We demonstrate this by erroneously identifying action-value

representations, both in simulations and in the neural activity recorded in unrelated

experiments. Second, we show that the experiments and analyses designed to dissociate

action-value representation from the representation of other decision variables cannot do so.

Specifically, we show that neurons representing policy may be erroneous identified as

representing action-values. We suggest different solutions to identifying action-value

representation that are not subject to these confounds. Applying one of these solutions to

previously identified action-value neurons in the basal ganglia we fail to detect action-value

representations there. Thus, we conclude that the claim that striatal neurons encode action-

values must await new experiments and analyses.

Key words

action-value; striatum; temporal correlations;

There is a long history of operant learning experiments, in which a subject, human or animal,

repeatedly chooses between actions and is rewarded according to its choices. A popular theory

posits that the subject's decisions in these tasks utilize estimates of the different *action-values*.

These action-values correspond to the expected reward associated with each of the actions, and

actions associated with a higher estimated action-value are more likely to be chosen [1]. In recent

years, there is a lot of interest in the neural mechanisms underlying this computation [2,3]. In

particular, based on electrophysiological, fMRI and intervention experiments, it is now widely

accepted that a population of neurons in the striatum represents these action-values, adding sway

to this action-value theory [4–22]. Here we challenge the evidence for action-value representation

in the striatum by describing two major confounds that have been overlooked when analyzing the

data.

To identify neurons that represent the internal values of the different actions, researchers have

searched for neurons whose firing rate is significantly correlated with the average reward

associated with exactly one of the actions. There are several ways of defining the average reward

associated with an action. For example, the average reward can be defined by the reward schedule,

e.g., the probability of a reward associated with the action. Alternatively, one can adopt the

subject's perspective, and use the subject-specific history of rewards and actions in order to

estimate the average reward. In particular, the Rescorla–Wagner model (equivalent to the standard

ones-state Q-learning model) has been used to estimate action-values [4,6]. In this model, the value

associated with an action i in trial t, termed $Q_i(t)$, is an exponentially-weighted average of the

rewards associated with this action in past trials:

$$Q_i(t + 1) = Q_i(t) + \alpha(R(t) - Q_i(t)) \quad \text{if } a(t) = i \qquad (1)$$

$$Q_i(t + 1) = Q_i(t) \quad \text{if } a(t) \neq i$$

where $a(t)$ and $R(t)$ denote the choice and reward in trial t, respectively, and α is the learning

rate.

In a two-alternative task, the probability of choosing an action is a sigmoidal function, typically

softmax, of the difference of the action-values (see also [23]):

$$\Pr(a(t) = 1) = \frac{1}{1 + e^{-\beta(Q_1(t) - Q_2(t))}} \qquad (2)$$

where β is a parameter that determines the bias towards the action associated with the higher

action-value. The parameters of the model, α and β, can be estimated from the behavior, allowing

the researchers to compute Q_1 and Q_2 on a trial-by-trial basis.

In principle, one can identify the neurons that represent action-value by identifying neurons for

which the regression of the trial-by-trial spike count on one of the variables $Q_i(t)$ is statistically

significant. Using this framework, electrophysiological studies have found that the firing rate of a

substantial fraction of striatal neurons (12%-40% for different significance thresholds) is

significantly correlated with an action-value. These and similar results were considered as

evidence that neurons in the striatum represent action-values [4–6,9–13].

In this paper we conduct a systematic literature search and conclude that the literature has, by and

large, ignored two major confounds in this analysis. First, it is well-known that spurious

correlations can emerge in correlation analysis if the variables are temporally correlated [24,25].

Here we show that neurons can be erroneously identified as representing action-values when their

firing rates are weakly temporally correlated. Second, it is also well-known that lack of a statistical

significance in the analysis does not imply lack of correlation. Because in standard analyses

neurons are classified as representing action-values if they have a significant regression coefficient

on *exactly* one action-value and because decision-related variables such as policy are correlated

with both action-values, neurons representing other decision-related variables may be

misclassified as representing action-values. We propose different approaches to address these

issues. Applying one of them to recordings from the basal ganglic, we fail to identify any action-

values representation there. Thus, we conclude that the hypothesis that striatal neurons represent

action-value still remains to be tested by experimental designs and analyses that are not subject to

these confounds. In the Discussion we address additional conceptual issues with such a

representation.

This paper discusses a methodological problem that may also be of relevance in other fields of

biology in general and neuroscience in particular. Nevertheless, the focus of this paper is a single

scientific claim, namely, that action-value representation in the striatum is an established fact. Our

criticism is restricted to the representation of action-values, and we do not make any claims

regarding the possible representations of other decision variables, such as policy, chosen-value or

reward-prediction-error. Moreover, we do not make any claims about the possible representations

of action-values elsewhere in the brain, although our results suggest caution when looking for such

representations.

The paper is organized in the following way. We commence by describing a standard method for

identifying action-value neurons. Next, we show that this method erroneously classifies simulated

neurons whose activity is temporally correlated as representing action-value. We show that this

confound brings into question the conclusion of many existing publications. Then, we propose

different methods for identifying action-value neurons, that overcome this confound. Applying

such a method to basal ganglia recordings, in which action-value neurons were previously

identified, we fail to conclusively detect any action-value representations. We continue by

discussing the second confound: neurons that encode the policy, the probability of choice, may be

erroneously identified as representing action-value, even when the policy is the result of learning

algorithms that are devoid of action-value calculation. Then we discuss a possible solution to this

confound.

Results

Identifying action-value neurons

We commence by examining the standard methods for identifying action-value neurons using a

simulation of an operant learning experiment. We simulated a task, in which the subject repeatedly

chooses between two alternative actions, which yield a binary reward with a probability that

depends on the action. Specifically, each session in the simulation was composed of four blocks

such that the probabilities of rewards were fixed within a block and varied between the blocks.

The probabilities of reward in the blocks were (0.1,0.5), (0.9,0.5), (0.5,0.9) and (0.5,0.1) for actions

1 and 2, respectively (Fig. 1A). The order of blocks was random and a block terminated when the

more rewarding action was chosen more than 14 times within 20 consecutive trials [4,10].

To simulate learning behavior, we used the Q-learning framework (Eqs. (1) and (2) with $\alpha = 0.1$

and $\beta = 2.5$ (taken from distributions reported in [6]) and initial conditions $Q_i(1) = 0.5$). As

demonstrated in Fig. 1A, the model learned, such that the probability of choosing the more

rewarding alternative increased over trials (black line). To model the action-value neurons, we

simulated neurons whose firing rate is a linear function of one of the two Q-values and whose

spike count in a 1 sec trial is randomly drawn from a corresponding Poisson distribution (see

Materials and Methods). The firing rates and spike counts of two such neurons, representing

action-values 1 and 2, are depicted in Fig. 1B in red and blue, respectively.

One standard method for identifying action-value neurons is to compare the firing rates after

learning by comparing the spike counts at the end of the blocks (horizontal bars in Fig. 1B).

Considering the red-labeled Poisson neuron, the spike count in the last 20 trials of the second

block, in which the probability of reward associated with action 1 was 0.9, was significantly higher

than that count in the first block, in which the probability of reward associated with action 1 was

0.1 ($p < 0.01$; rank sum test). By contrast, there was no significant difference in the spike counts

between the third and fourth blocks, in which the probability of reward associated with action 1

was equal ($p = 0.91$; rank sum test; Fig. 1B, red). This is consistent with the fact that the red-

labeled neuron was an action 1-value neuron: its firing rate was a linear function of the value of

action 1. Similarly for the blue labeled neuron, the spike counts in the last 20 trials of the first two

blocks were not significantly different ($p = 0.92$; rank sum test), but there was a significant

difference in the counts between the third and fourth blocks ($p < 0.001$; rank sum test). These

results are consistent with the probabilities of reward associated with action 2 and the fact that in

our simulations, this neuron's firing rate was modulated by the value of action 2 (Fig. 1B, blue).

This approach for identifying action-value neurons is limited, however, for several reasons. First,

it considers only a fraction of the data, the last 20 trials in a block. Second, action-value neurons

are not expected to represent the block average probabilities of reward. Rather, they will represent

a subjective estimate, which is based on the subject-specific history of actions and rewards.

Therefore, it is more common to identify action-value neurons by regressing the spike count on

subjective action-values, estimated from the subject's history of choices and rewards [4–6,10–12].

Note that when studying behavior in experiments, we have no direct access to these estimated

action-values, in particular because the values of the parameters α and β are unknown. Therefore,

following common practice, we estimated the values of α and β from the model's sequence of

choices and rewards using maximum likelihood, and used the estimated learning rate (α) and the

choices and rewards to estimate the action-values (thin lines in Fig. 1C, see Materials and

Methods). These estimates were similar to the true action-value, which underlay the model's

choice behavior (thick lines in Fig. 1C).

Next, we regressed the spike count of each simulated neuron on the two estimated action-values

from its corresponding session. As expected, the t-values of the regression coefficients of the red-

labeled action 1-value neuron was significant for the estimated Q_1 (t_{18} (Q_1) = 4.05) but not for

the estimated Q_2 (t_{18} (Q_2) = −0.27). Similarly, the t-values of the regression coefficients of the

blue-labeled action 2-value neuron was significant for the estimated Q_2 (t_{18} (Q_2) = 3.05) but not

for the estimated Q_1 (t_{18} (Q_1) = 0.78).

A population analysis of the t-values of the two regression coefficients is depicted in Fig. 1D,E.

As expected, a substantial fraction (42%) of the simulated neurons were identified as action-value

neurons. Only 2% of the simulated neurons had significant regression coefficients with both

action-values. Such neurons are typically classified as state (ΣQ) or policy (also known as

preference) (ΔQ) neurons, if the two regression coefficients have the same or different signs,

respectively [10]. Note that despite the fact that by construction, all neurons were action-value

neurons, not all of them were detected as such by this method. This failure occurred for two

reasons. First, the estimated action-values are not identical to the true action-values, which

determine the firing rates. This is because of the finite number of trials and the stochasticity of

choice (note the difference, albeit small, between the thin and thick lines in Fig. 1C). Second and

more importantly, the spike count in a trial is only a noisy estimate of the firing rate because of the

Poisson generation of spikes.

Several prominent studies have implemented the methods we described in this section and reported

that a substantial fraction (10-40% depending on significance threshold) of striatal neurons

represent action-value [4,10,11]. In the next two sections we show that these methods, and similar

methods employed by other studies [5–9,12–14,17–22] are all subject to at least one of two major

confounds.

Confound 1 – Temporal correlations

Simulated random-walk neurons are erroneously identified as action-value neurons

The red and blue-labeled neurons in Fig. 1D were classified as action-value neurons because their

t-values were improbable under the null hypothesis that the firing rate of the neuron is not

modulated by action-values. The significance threshold (t=2) was computed assuming that trials

are independent in time. To see why this assumption is essential, we consider a case in which it is

violated. Fig. 2A depicts the firing rates and spike counts of two simulated Poisson neurons, whose

firing rates follow a bounded Gaussian random-walk process:

$$f(t + 1) = [f(t) + z(t)]_+ \quad (3)$$

where $f(t)$ is the firing rate in trial t (we consider epochs of 1 second as "trials"), $z(t)$ is a diffusion

variable, randomly and independently drawn from a normal distribution with mean 0 and variance

$\sigma^2 = 0.01$ and $[x]_+$ denotes a linear-threshold function, $[x]_+ = x$ if $x \geq 0$ and 0 otherwise.

These random-walk neurons are clearly not action-value neurons. Nevertheless, we tested them

using the analyses depicted in Fig. 1. To that goal, we randomly matched the trials in the simulation

of the random-walk neurons to the trials in the simulation depicted in Fig. 1A. This was done

because the random-walk is completely unrelated to the experimental settings or to any kind of

learning. Then, we considered the spike counts of the random-walk neurons in the last 20 trials of

each of the four blocks in Fig. 1A (block being defined by the simulation of learning and is

unrelated to the neural activity of the random-walk neurons). Surprisingly, when considering the

top neuron in Fig. 2A and utilizing the same analysis as in Fig. 1B, we found that its spike count

differed significantly between the first two blocks ($p < 0.01$, rank sum test) but not between the

last two blocks ($p = 0.28$, rank sum test), similarly to the simulated action 1-value neuron of Fig.

1B (red). Similarly, the spike count of the bottom random-walk neuron matched that of a simulated

action 2-value neuron (compare with the blue-labeled neuron in Fig. 1B; Fig. 2A).

Moreover, we regressed each vector of spike counts for 20,000 random-walk neurons on randomly

matched estimated action-values from Fig. 1E and computed the t-values (Fig. 2B). This analysis

classified 42% of these random-walk neurons as action-value neurons (see Fig. 2C). In particular,

the top and bottom random-walk neurons of Fig. 2A were identified as action-value neurons for

action 1 and 2, respectively (squares in Fig. 2B).

To further quantify this result, we computed the fraction of random-walk neurons erroneously

classified as action-value neurons as a function of the diffusion parameter σ (Fig. 2D). When $\sigma=0$,

the spike counts of the neurons in the different trials are independent and the number of random-

walk neurons classified as action-value neurons is slightly less than 10%, the fraction expected by

chance from a significance criterion of 5% and two statistical tests, corresponding to the two

action-values. The larger the value of σ, the higher the probability that a random-walk neuron will

pass the selection criterion for at least one action-value and thus be erroneously identified as an

action-value, state or policy neuron.

The excess action-value neurons in Fig. 2 emerged because the statistical analysis was based on

the assumption that the different trials are independent from each other. In the case of a regression

of a random-walk process on an action-value related variable, this assumption is violated. The

reason is that in this case, both predictor (action-value) and the dependent variable (spike count)

slowly change over trials, the former because of the learning and the latter because of the random

drift. As a result, the statistic, which relates these two signals, is correlated between temporally-

proximate trials, violating the independence-of-trials assumption of the test. Because of these

dependencies, the expected variance of the statistic (be it average spike count in 20 trials or the

regression coefficient), which is calculated under the independence-of-trials assumption, is an

underestimate of the actual variance. Therefore, the fraction of random-walk neurons classified as

action-value neurons increases with the magnitude of the drift, which is directly related to the

magnitude of correlations between spike counts in proximate trials (Fig. 2D). The phenomenon of

spurious significant correlations in time-series with temporal correlations has been described

previously in the field of econometrics and a formal discussion of this issue can be found in

[24,25].

Is this confound relevant to the question of action-value representation in the striatum?

<u>Is a random-walk process a good description of striatal neurons' activity?</u>

The Gaussian random-walk process is just an example of a temporally correlated firing rate and

we do not argue that the firing rates of striatal neurons follow a random-walk process. However,

any other types of temporal correlations, e.g., oscillations or trends, will violate the independence-

of-trials assumption, and may lead to the erroneous identification of neurons as representing

action-values. Such temporal correlations can also emerge from stochastic learning. For example,

in Fig. S1 we consider a model of operant leaning that is based on covariance synaptic plasticity

[26–30]. Because such plasticity results in slow changes in the firing rates of the neurons, applying

the analysis of Fig. 1E to our simulations results in the erroneous identification of 43% of the

simulated neurons as representing action-values. This is despite the fact that action-values are not

computed, either explicitly or implicitly.

<u>Are temporal correlations in neural recordings sufficiently strong to affect the analysis?</u>

To test the relevance of this confound to experimentally-recorded neural activity, we repeated the

analysis of Fig. 2B,C on neurons recorded in two unrelated experiments: 89 neurons from

extracellular recordings in the motor cortex of an awake monkey (Fig. S2A-B) and 39 auditory

cortex neurons recorded intracellularly in anaesthetized rats (Fig. S2C-D). We regressed the spike

counts on randomly matched estimated action-values from Fig. 1E. In both cases we erroneously

identified action-value representations (36% and 23%, respectively) a fraction comparable to that

reported in the striatum).

<u>Strong temporal correlations in the striatum</u>

To test the relevance of this confound to striatal neurons, we considered previous recordings from

neurons in the nucleus accumbens (NAc) and ventral pallidum (VP) of rats in an operant learning

experiment [7] and regressed their spike counts on the action-values of the simulations in Fig. 1E.

Again, we erroneously identified a substantial fraction of neurons (43%) as representing action-

value, a fraction comparable to that reported in the striatum (Fig. S3).

<u>Haven't previous publications acknowledged this confound and successfully addressed it?</u>

We conducted an extensive literature search to see whether previous studies have identified this

confound and addressed it (see Materials and Methods). Two studies noted that processes such as

slow drift in firing rate may violate the independence-of-trials assumption of the statistical tests

and suggested unique methods to address this problem [6,9]. One method [6] relied on permutation

of the spike count within a block (Fig. S4, see Materials and Methods) and another [9], relied on

using spikes in previous trials as predictors (Fig. S5). However, both approaches fail to account

for all temporal correlations and as a result, they still erroneously identify unrelated and random-

walk neurons as action-value neurons (Figs. S4 and S5). The failure of both these approaches stems

from the fact that we lack a complete model of the learning-independent temporal correlations. As

a result, these methods are unable to remove *all* the temporal correlations from the vector of spike-

counts.

Our literature search yielded four additional methods that have been used to identify action-value

neurons. However, as depicted in Figs. S6 (corresponding to the analyses in [4,7]), S7

(corresponding to the analysis in [10]), S8 (corresponding to the analysis in [8]) and S9

(corresponding to the analysis in [17]), all these additional methods erroneously identify neurons

from unrelated recordings and random-walk neurons as action-value neurons, in numbers

comparable to those reported in the striatum (Figs. S6-S9). The erroneous detection of action-value

representation in S9 is particularly surprising, because this analysis utilizes a trial-design

experiment in which trials with different reward probabilities are randomly intermingled.

However, because in this analysis the regression's predictors are estimated action-values, which

are temporally correlated, even in a trial-design some temporal correlations remain. From this

example it follows that some trial-design experiments are still subject to the temporal correlations

confound.

<u>Shouldn't adding blocks solve the problem?</u>

In Figs. 1 and 2 we considered a learning task composed of four blocks and a mean length of 174

trials (standard deviation 43 trials). It is tempting to believe that experiments with more blocks and

trials (e.g., [7,8]) will be immune to this confound. The intuition is that the larger the number of

trials, the less likely it is that a neuron that is not modulated by action-value (e.g., a random-walk

neuron) will have a large regression coefficient on one of the action-values. However, surprisingly,

this intuition is wrong. In Fig. S10 we show that doubling the number of blocks does not decrease

the fraction of neurons erroneously classified as representing action-values. For the case of

random-walk neurons, it can be shown that, contrary to this intuition, the fraction of erroneously

identified action-value neurons is expected to increase with the number of trials [25]. This is

because the expected variance of regression coefficients under the null hypothesis is proportional

to the inverse of the number of degrees of freedom, which increases with the number of trials. As

a result, the threshold for classifying a regression coefficient as significant decreases with the

number of trials.

Possible solutions to the temporal correlations confound

<u>Permutation analysis</u>

Trivially, an action-value neuron (or any task-related neuron) should be more strongly correlated

with the action-values of the experimental session, in which the neuron was recorded than with

action-values of other sessions (recorded in different days). We propose to use this requirement in

a permutation test, as depicted in Fig. 3. We first consider the two simulated action-value neurons

of Fig. 1B. For each of the two neurons, we computed the t-values of the regression coefficients

of the spike counts on each of the estimated action-values in all possible sessions (see Materials

and Methods). Fig. 3A depicts the two resulting distributions of t-values. As a result of the

temporal correlations, the 5% significance boundaries (vertical dashed lines), which are exceeded

by exactly 5% of t-values in each distribution, are substantially larger (in absolute value) than 2,

the standard significance boundaries. We posit that the neuron is significantly correlated with an

action-value if the t-value of the regression on the action-value (from the corresponding session)

exceeds the significance boundaries derived from the permutation test.

Indeed, when considering the Top (red) simulated action 1-value neuron, we find that its spike

count has a significant regression coefficient on the estimated Q_1 from its session (red arrow) but

not on the estimated Q_2 (blue arrow). Importantly, because the significance boundary exceeds 2,

this approach is less sensitive than the original one (Fig. 1) and indeed, the regression coefficients

of the Bottom simulated neuron (blue) do not exceed the significance level (red and blue arrows)

and thus this analysis fails to identify it as an action-value neuron. Considering the population of

simulated action-value neurons of Fig. 1, this analysis identified 29% of the action-value neurons

of Fig. 1 as such (Fig. 3B, black), demonstrating that this analysis can identify action-value

neurons. When considering the random-walk neurons (Fig. 2), this method classifies only

approximately 10% of the neurons as action-value neurons, as predicted by chance. Similar results

were obtained for the motor cortex and auditory cortex neurons (not shown).

<u>Permutation analysis of basal ganglia neurons</u>

Importantly, this permutation method can also be used to reanalyze the activity of previously-

recorded neurons. To that goal, we considered the recordings reported in [7]. The results of their

model-free method (Fig. S6) imply that approximately 23% of the striatal neurons represent action-

values at different phases of the experiment. As a first step, we estimated the action-values and

regressed the spike counts in the different phases of the experiment on the estimated action-values,

as in Fig. 1 (see Materials and Methods). The results of this analysis implied that 32% of the

neurons represent action values (p<0.01). Next, we applied the permutation analysis. Remarkably,

this analysis yielded that only 3.6% of the neurons have a significantly higher regression

coefficient on an action-value from their session than on other action-values (Fig. 3C). Similar

results were obtained when performing a similar model-free permutation analysis (not shown).

These results raise the possibility that all or much of the apparent action-value representation in

[7] is the result of the temporal correlations confound.

It should be noted that the fraction of action-value neurons reported in [7] is low relative to other

publications, a difference that has been attributed to the location of the recording in the striatum

(ventral as opposed to dorsal). It would be interesting to apply this method to other striatal

recordings [4,8,10].

<u>Trial-design experiments</u>

Another way of overcoming the temporal correlations confound is to use a trial design in the

experiment. The idea is to randomly mix the reward probabilities, rather than use blocks as in Fig.

1. For example, consider the experimental design depicted in Fig. 4A. Each trial is presented in

one of four clearly-marked contexts (color coded). The reward probabilities associated with the

two actions are fixed within a context but differ between the contexts. Within each context the

participant learns to prefer the action associated with a higher probability of reward. If the contexts

are randomly mixed, then by construction, the reward probabilities are temporally independent.

Therefore, in a regression analysis, in which the reward probabilities are predictors, the

independence assumption is not violated.

We simulated learning in a session composed of 280 trials, randomly divided into 4 different

contexts (Fig. 4A). Learning followed the Q-learning equations (Eqs. 1 and 2), independently for

each context. Next, we simulated action-value neurons, as in Fig. 1A, whose firing rate was a linear

function of the action-value in the relevant context (dots in Fig. 4A). Next, we regressed the spike

counts of the neurons in the last 80 trials (approximately 20 trials in each context) on the

corresponding reward probabilities (Fig. 4B). Indeed, 54% of the neurons were classified this way

as action-value neurons (Fig. 4C 10% is chance level). By contrast, considering random-walk

neurons, only 10% were erroneously classified as action-value neurons, a fraction comparable to

that expected by chance.

Although no study has used a design exactly like the one we suggest here, three studies have used

designs in which cues mark reward probabilities, in order to study neural representation in the

striatum [14,17,21] (but see Fig. S9 for [17]). These studies reported significant modulation of

activity by the difference between the two action-values (a measure considered as policy [10]) but

they did not report a representation of exactly one action-value.

Two points are noteworthy. First, in such trial-design experiments the participant learns to

associate cue and reward (cue-values), and not action and reward (action-values) [31]. This may

also be a concern for the analysis in Fig. 4. Second, as discussed in details in the next section,

policy representation can emerge without action-value representation [26–29,32–36]. Therefore,

the results reported in [14,17,21] cannot be taken as evidence for action-value representation in

the striatum.

Confound 2 – correlated decision variables

In the previous sections we demonstrated that irrelevant temporal correlations may lead to the

erroneous identification of neurons as representing action-value, even if their activity is task-

independent. Here we address an unrelated confound. We show that neurons that encode different

decision variables, in particular policy, may be erroneously identified as representing action-

values. For clarity, we will commence by discussing this caveat independently of the temporal

correlations confound. Specifically, we show that neurons whose firing rate encodes the policy

(probability of choice) may be erroneously identified as representing action-values, even when this

policy emerged in the absence of any implicit or explicit action-value representation. We will

conclude by discussing a possible solution that addresses both confounds.

Policy without action-value representation

It is well-known that operant learning can occur in the absence of any value computation, e.g., as

a result of direct-policy learning [30]. Several studies have shown that reward-modulated synaptic

plasticity can implement direct policy reinforcement learning [26–29,32–36].

For concreteness, we consider a particular reinforcement learning algorithm, in which the

probability of choice $\Pr(a(t) = 1)$ is determined by a single parameter W that is learned in

accordance to REINFORCE [37] learning algorithm: $\Pr(a(t) = 1) = \frac{1}{1+e^{-W(t)}}$ where $\Delta W(t) =$

$\alpha(2 \cdot R(t) - 1) \cdot (a(t) - \Pr(a(t) = 1))$, where α is the learning rate, $R(t)$ is the binary reward

in trial t and $a(t)$ is a binary variable indicating whether action 1 was chosen in trial t;

$W(t = 1) = 0$. For biological implementation of this algorithm see [28,32].

We tested this model in the schedule of Fig. 1. (Fig. 5A). As expected, the model learned to prefer

the action associated with a higher probability of reward, completing the four blocks within 228

trials on average (standard deviation 62 trials).

Spike count of neurons representing policy are correlated with estimated ΔQ

Despite the fact that the learning was value-independent, we can still fit a Q-learning model to the behavior, extract best-fit model parameters and compute action-values (see also Fig. S1). The computed action-values are presented in Fig. 5B. Note that according to Eq. (2), the probability of choice is a monotonic function of the difference between Q_1 and Q_2. Therefore, we expected that the probability of choice will be correlated with the computed Q_1 and Q_2, with opposite signs (Fig. 5C).

We simulated policy neurons as Poisson neurons whose firing rate is a linear function of the policy $\Pr(a)$ (Materials and Methods). Next, we regressed the spike count of these neurons on the two action values that were computed from behavior (same as in Figs. 1D,E, 2B,C, S1C,D, S2B,D, S3). Indeed, as expected, 14% of the neurons were significantly correlated with both action values with opposite signs (chance level for each action value is 5%, chance level for both with opposite signs is 0.125%), as depicted in Fig. 5D,E. These results demonstrate that neurons representing value-independent policy can be classified as representing ΔQ.

Neurons representing policy may be erroneously classified as action-value neurons

Surprisingly however, 38% of policy neurons were significantly correlated with *exactly* one estimated action-value, and therefore would have been classified as action-value neurons in the standard method of analysis (10% chance level).

To understand why this erroneous classification emerged, we note that a neuron is classified as representing action-value if its spike count is significantly correlated with one of the action values, but not the other. The confound that led to the classification of policy neurons as representing action-values is that *a lack of statistically significant correlation does not imply lack of correlation.*

All policy neurons are modulated by the probability of choice, a variable that is correlated with

the difference in the two action-values. Therefore, this probability of choice is expected to be

correlated with both action-values, with opposite signs. However, because the neurons are Poisson,

the spike count of the neurons is a noisy estimate of the probability of choice. As a result, in most

cases (86%), the regression coefficients do not cross the significance threshold for *both* action-

values. More often (38%), only one of them crosses the significance threshold, resulting in an

erroneous identification of the neurons as representing action values.

Is this confound relevant to the question of action-value representation in the striatum?

<u>If choice is included as a predictor, is policy representation still a relevant confound?</u>

It is common, (although not ubiquitous) to attempt to differentiate action-value representation from

choice representation by including choice as another regressor in the regression model [5,6,9–

14,17,21]. Such analyses may be expected to exclude policy neurons, whose firing rate is highly

correlated with choice, from being classified as action-value neurons. However, repeating this

analysis (Materials and Methods) for the policy neurons of Fig. 5, we still classify 36% of policy

neurons as action-value neurons (Figure S11A).

An alternative approach has been to consider only those neurons whose spike count is not

significantly correlated with choice [19,22]. Repeating this analysis for Fig. 5 neurons, we still

find that 24% of the neurons are erroneously classified as action-value neurons (8% are classified

as policy neurons).

<u>Is this confound the result of an analysis that is biased against policy representation?</u>

The analysis depicted in Figs. 1D,E, 2B,C, 4B-E, and 5D,E is biased towards classifying neurons

as action-value neurons, at the expense of state or policy neurons, as noted by [8]. This is because

action-value classification is based on a single significant regression coefficient whereas policy or

state classification requires two significant regression coefficients. Therefore, [8] have proposed

an alternative approach. First, compute the statistical significance of the whole regression model

for each neuron (using f-value). Then, classify those significant neurons according to the t-value

of regression coefficients with the two action-values (Fig. S11B). Applying this analysis to the

policy neurons of Fig. 5 with a detection threshold of 5% we find that indeed, this method is useful

in detecting which decision variables are more frequently represented (its major use in [8]): 25%

of the neurons are indeed classified as representing policy (1.25% expected chance). Nevertheless,

12% of the neurons are still classified as action-value neurons (2.5% expected by chance; Fig.

S11B).

<u>Additional issues</u>

In many cases, the term action-value was used, while the reported results were equally consistent

with other decision variables. In some cases, significant correlation with both action-values (with

opposite signs) or significant correlation with the difference between the action-values was used

as evidence for 'action-value representations' [17,18,20–22]. Similarly, other papers did not

distinguish between neurons whose activity is significantly correlated with one action-value and

those whose activity is correlated with both action-values [6,9,12–14]. Finally, in one study the

two action-values were fully correlated [5].

A possible solution to the policy confound

The policy confound emerged because policy and action-values are correlated. To distinguish

between the two possible representations, we should seek a variable that is correlated with the

action-value but uncorrelated with the policy. Consider the sum of the two action-values. It is easy

to see that $\text{Corr}(Q_1 + Q_2, Q_1 - Q_2) \propto \text{Var}(Q_1) - \text{Var}(Q_2)$. Therefore, if the variances of the two

action-values are equal, their sum is uncorrelated with their difference. An action-value neuron is

expected to be correlated with the sum of action-values. By contrast, a policy neuron, modulated

by the difference in action-values is expected to be uncorrelated with this sum.

We repeated the simulations of Fig. 4 (which addresses the temporal correlations confound),

considering three types of neurons: action-value neurons (of Fig. 1), random-walk neurons (of Fig.

2), and policy neurons (of Fig. 5). As in Fig. 4, we regressed the spike counts of the three types

of neurons in the last 80 trials of the session on the sum of reward probabilities (state). We found

that only 5% and 6% of the random-walk and policy neurons, respectively, were significantly

correlated with the sum of reward probabilities (5% chance). By contrast, 38% of the action-value

neurons were significantly correlated with this sum.

This method is able to distinguish between policy and action-value representations. However, it

will fail in the case of state representation because both state and action-values are correlated with

the sum of probabilities of reward. To dissociate between state and action-value representations,

we can consider the difference in reward probabilities as this difference is correlated with the

action-value but is uncorrelated with the state. Regressing the spike count on *both* the sum and

difference in the probabilities of reward, a random-walk neuron is expected to be correlated with

none, a policy neuron is expected to be correlated only with the difference, whereas an action-

value neuron is expected to be correlated with both. This is depicted in Fig. 6A,C,E, where we

report the t-value of these correlations. We now classify a neuron that passes both significance

tests as an action-value neuron. Indeed, for a significance threshold of $p<0.05$ (for each test), Only

0.4% of the random-walk neurons and 4% of the policy neurons were classified as action-value

neurons. By contrast, 22% of the action-value neurons were classified as such. Note that in this

analysis only when more than 5% of the neurons are classified as action-value neurons we have

support for the hypothesis that there is action-value rather than policy or state representation.

To conclude, it is worthwhile repeating the key features of the analysis method proposed in this

section:

1) Trial design is necessary because otherwise temporal correlations in spike count may

inflate the fraction of neurons that pass the significance tests.

2) Regression should be performed on reward probabilities and not on estimated action-

values. The reason is that because the estimated action-values evolve over time, this trial

design does not eliminate all temporal correlations between them (Fig. S9).

3) Reward probabilities associated with the two actions should be chosen such that their

variances should be equal. Otherwise policy or state neurons may be erroneously classified

as action-value neurons.

Discussion

In this paper, we performed a systematic literature search to discern the methods that have been

previously used to infer the representation of action-values in the striatum. We showed that none

of these methods overcome two critical confounds: (1) neurons with temporal correlations in their

firing rates may be erroneously identified as representing action-values and (2) neurons whose

activity co-varies with other decision variables, such as policy, may be erroneously classified as

representing action-values. Finally, we discuss possible experiments and analyses that can address

the question of whether striatal neurons encode action-values.

Temporal correlations and action-value representations

It is well known in statistics that the regression coefficient between two independent, slowly-

changing variables, is on average larger (in absolute value) than this coefficient when the series

are devoid of a temporal structure. Thus, if these temporal correlations are overlooked, the

probability of a false-positive would be underestimated [24]. When searching for action-value

representation in a block design, then by construction, there are positive correlations in the

predictor (action-values). Positive temporal correlations in the dependent variable (neural activity)

will result in an inflation of the false-positive observations, compared with the naïve expectation.

This confound occurs only when there are temporal correlations in both the predictor and the

dependent variable. In a trial design, in which the predictor is randomized and thus has no temporal

structure, we do not expect this confound. However, when studying incremental learning, it is

difficult to randomize the predictor, making it difficult to identify neural correlates of learning.

With respect to the dependent variable (neural activity), temporal correlations in BOLD signal and

their consequences have been discussed [38,39]. Considering electrophysiological recordings,

there have been attempts to remove these correlations, e.g., using previous spike counts as

predictors [9]. However, these are not sufficient because they are unable to remove all task-

independent temporal correlations (see also Figs S4-S10).

One may argue that the fact that action-value representations are reported mostly in a specific brain

area, namely, the striatum, is an indication that their identification there is not a result of the

temporal correlations confound. However, the probability of a false-positive identification of a

neuron as representing action-value depends on the magnitude of temporal correlations in the

neural activity (Fig. 2D). Different brain regions are characterized by different spiking statistics.

Thus, we expect different levels of erroneous identification of action-value neurons in different

parts of the brain and in different experimental settings. Indeed, the fraction of erroneously

identified action-value neurons differed between the auditory and motor cortices (compare Figs.

S2B and S2D). Furthermore, many studies reported action-value representation outside of the

striatum, in brain areas including the supplementary motor area and presupplementary eye fields

[19], the substantia nigra/ventral tegmental area [18] and ventromedial prefrontal cortex, insula

and thalamus [17]. It is also worthwhile noting that [40] have reported that results relating neural

activity with model-based learned value are robust to the parameters fitted in the model, namely

the learning rate. The fact that the correlation of neural activity with value is robust to different

learning rates is in line with the result that temporal correlations in time series yield, on average,

larger regression coefficients.

Considering the ventral striatum, our analysis indicates that the identification of action-value

representations there may have been erroneous, resulting from temporally correlated firing rates

(Figs. 3 and S3). We were unable to directly analyze recordings from the dorsal striatum because

relevant raw data is not publically available. However, previous studies have reported that the

firing rates of dorsal-striatal neurons change slowly over time [41,42]. As a result, identification

of apparent action-value representation in dorsal-striatal neurons may also be the result of this

confound.

Temporal correlations – beyond action-value representation

It should be noted that temporal correlations naturally emerge in experiments composed of

multiple trials. Participants become satiated, bored, tired, etc., which may affect neuronal activity.

In particular, learning in operant tasks is associated, by construction, with temporally correlated

trials, the learning itself. If neural activity is correlated with performance (e.g., accumulated

rewards in the last several trials) then it is expected to have temporal correlations, which may lead

to an erroneous classification of the neurons as representing action-values.

We should be particularly cautious about temporal correlations when searching for the neural

correlates of continuous behaviors. In natural settings, many behavioral variables, such as the

location of the animal or the activation of different muscles change at relatively long time-scales.

Any temporal correlations in the neural recording, be it in fMRI, electrophysiology or calcium

imaging may result in an erroneous identification of correlates of these behavioral variables.

Correlated decision variables

Another difficulty in identifying action-value neurons is that they are correlated with other decision

variables such as policy, state and chosen-value. Therefore, finding a neuron that is significantly

correlated with action-value could be the byproduct of its being modulated by other decision

variables, in particular policy. The problem is exacerbated by the fact that standard analyses (e.g.,

Fig. 1D-E) are biased towards classifying neurons as representing action-values at the expense of

policy or state. However, because of the correlation between policy and action-value, even

unbiased analyses may erroneously identify a significant fraction of neurons as representing

action-value (Fig. S11B).

It is important to emphasize that policy representation (and more generally, the representation of

other decision variables) does not imply the explicit or implicit representation of action-values

anywhere in the brain. As demonstrated in Figs. S1 and 5, learning of the operant task is possible

using direct policy gradient algorithms, that are devoid of action-value representation [23,30].

Differentiating action-value from other decision variables

As shown in Fig. 6, policy representation can be ruled out by finding representations that are

orthogonal to policy, namely state representation. However, this leads us to a more conceptual

issue. All analyses discussed so far are based on significance tests: we divide the space of

hypothesis into the "scientific claim" (e.g., neurons represent action-values) and the null

hypothesis (e.g., neural activity is independent of the task). An observation that is not consistent

with the null hypothesis is taken to support the alternative hypothesis.

The problem we faced with correlated variables is that the null hypothesis and the "scientific

claim" were not complementary. A neuron that represents policy is expected to be inconsistent

with the null hypothesis that neural activity is independent of the task but it is not an action-value

neuron. The solution proposed was to devise a statistical test that identifies representation that is

orthogonal to policy thus ruling out policy neurons.

However, this does not rule out other representations. A "pure" action-value neuron is modulated

only by Q_1 or by Q_2. A "pure" policy neuron is modulated exactly by $Q_1 - Q_2$. More generally,

we may want to consider the hypotheses that the neuron is modulated by a different combination

of the action values, $a \cdot Q_1 + b \cdot Q_2$, where a and b are parameters. For every such set of parameters

a and b we can devise a statistical test to reject this hypothesis by considering the direction that is

orthogonal to the vector (a, b). In principle, this procedure should be repeated for every pair of

parameters a and b that in not consistent with the action-value hypothesis.

Put differently, in order to find neurons that represent action-values, we first need to *define* a set

of parameters a and b such that a neuron whose activity is modulated by $a \cdot Q_1 + b \cdot Q_2$ will be

considered as representing an action-value. Only after this (arbitrary) definition is given, can we

construct a set of statistical tests that will rule out the competing hypotheses, namely will rule out

all values of a and b that are not in this set.

Finally, considering the distribution of t-values of regression coefficients across the population of

neurons may provide important context to the question of representation of specific variables. For

example, the reader can assess the plausibility of a specific population of neurons representing

action-values, rather than a general, less specific modulation by a combination of decision

variables from observing this distribution in [4,8].

Action value representation in the striatum requires further evidence

Considering the literature, both confounds have been partially acknowledged. Moreover, there

have been some attempts to address them. However, as discussed above, even when these

confounds were acknowledged and solutions were proposed, these solutions do not prevent the

erroneous identification of action-value representation (see Figs. S4, S5, S10, S11). We therefore

conclude that to the best of our knowledge, **all** studies that have claimed to provide direct evidence

that neuronal activity in the striatum is specifically modulated by action-value were either

susceptible to the temporal correlations confound [4,7,8,10,11], or reported results in a manner

indistinguishable from policy [14,21]. Indeed, many studies were susceptible to both confounds

[5,6,9,12,13,17,19]. Furthermore, it should be noted that not all studies investigating the relation

between striatal activity and action-value representation have reported positive results. Several

studies have reported that striatal activity is more consistent with direct policy learning than with

action-value learning [43,44] and one noted that lesions to the dorsal striatum do not impair action-

value learning [45].

The fact that the basal ganglia in general and the striatum in particular play an important role in

operant learning, planning and decision-making is indisputable [3,15,46–50]. However, our results

show that special caution should be applied when relating activity in neurons there with

reinforcement learning algorithms and specific variables therein. Therefore, the prevailing belief

that neurons in the striatum represent action-values must await further tests that address the

potential confounds discussed here.

Materials and Methods

Literature search

In order to thoroughly examine the finding of action-value neurons in the striatum, we conducted a literature search to find all the different approaches used to identify action-value representation in the striatum and see whether they are subject to at least one of the two confounds we described here.

Key words "action-value" and "striatum" were searched for in Web-of-Knowledge, Pubmed and Google Scholar, returning 43, 21 and 980 results, respectively. In the first screening stage, we excluded all publications that did not report new experimental results (e.g., reviews and theoretical papers), focused on other brain regions, or did not address value-representation or learning. In the remaining publications, the abstract of the publication was read and the body of the article was searched for "action-value" and "striatum". After this step, articles in which it was possible to find description of action-value representation in the striatum were read thoroughly. The search included PhD theses, but none were found to report new relevant data, not found in articles. We identified 22 papers that directly related neural activity in the striatum to action-values. These papers included reports of single-unit recordings, functional magnetic resonance imaging (fMRI) experiments and manipulation of striatal activity.

Of these, 3 papers have used the term action-value to refer to the value of the *chosen* action (also known as chosen-value) [51–53] and therefore we will not discuss them any further.

A second group of 12 papers did not distinguish between action-value and policy representations [5,6,9,12–14,19], or reported policy representation [17,18,20–22] in the striatum and therefore

their findings do not necessarily imply action-value representation, rather than policy

representation in the striatum (see confound 2).

In 2 additional papers, it was shown that the activation of striatal neurons changes animals'

behavior, and the results were interpreted in the action-value framework [15,16]. However, a

change in policy does not entail an action-value representation (see, for example, Figs. 5 and S1).

Therefore, these papers were not taken as strong support to the striatal action-value representation

hypothesis.

Finally, 5 papers correlated action-values, separately from other decision variables, with neuronal

activity in the striatum [4,7,8,10,11]. All of them used electrophysiological recordings of single

units in the striatum. From these papers, only one utilized an analysis which is not biased towards

identifying action-value neurons at the expense of policy and state neurons [8]. All papers used

block-design experiments where action-values are temporally correlated.

Taken together, we concluded that previous reports on action-value representation in the striatum

could reflect the representation of other decision variables or temporal correlations in the spike

count that are not related to action-value learning.

The action-value neurons model (Fig. 1)

To model neurons whose firing rate is modulated by an action-value (Fig. 1), we considered

neurons whose firing rate changes according to:

$f(t) = B + K \cdot r \cdot (Q_i(t) - 0.5)$ (4)

Where $f(t)$ is the firing rate in trial t, $B = 2.5$Hz is the baseline firing rate, $Q_i(t)$ is the action-

value associated with one of the targets $i \in \{1,2\}$, $K = 2.35$Hz is the maximal modulation and r

denotes the neuron-specific level of modulation, drawn from a uniform distribution, $r \sim U[-1,1]$.

The spike count in a trial was drawn from a Poisson distribution, assuming a 1 sec-long trial.

The policy neurons model (Fig. 5)

To model neurons whose firing rate is modulated by a policy, we considered neurons whose firing

rate changes according to:

$f(t) = B + K \cdot r \cdot \Pr(a(t) = 1)$ (5)

Where $f(t)$ is the firing rate in trial t, $B = 2.5$Hz is the baseline firing rate, $\Pr(a(t) = 1)$ is the

probability of choosing action 1 in trial t that changes in accordance with REINFORCE [37] (see

also Fig. 5 and corresponding text). $K = 3$Hz is the maximal modulation and r denotes the neuron-

specific level of modulation, drawn from a uniform distribution, $r \sim U[-1,1]$. The spike count in a

trial was drawn from a Poisson distribution, assuming a 1 sec-long trial.

The covariance neurons model (Fig. S1)

In the covariance based plasticity model the decision-making network is composed of two

populations of Poisson neurons: each neuron is characterized by its firing rate and the spike count

of a neuron in a trial (1 sec) is randomly drawn from a Poisson distribution. The chosen action

corresponds to the population that fires more spikes in a trial [26,28]. At the end of the trial, the

firing rate of each of the neurons (in the two population) is updated according to $f(t + 1) =$

$f(t) + \eta \cdot R(t) \cdot (s(t) - f(t))$, where $f(t)$ is the firing rate in trial t, $\eta = 0.07$ is the learning

rate, $R(t)$ is the reward delivered in trial t ($R(t) \in \{0,1\}$ in our simulations) and $s(t)$ is the

measured (realized) firing rate in that trial, that is the spike count in the trial. The initial firing rate

of all simulated neurons is 2.5Hz. The network model was tested in the operant learning task of

Fig. 1. A session was terminated (without further analysis) if the model was not able to choose the

better option more than 14/20 times for at least 200 trials. This occurred on 20% of the sessions.

Simulated neurons were excluded due to low spike rate if the mean spike count was lower than 1

for all blocks. This occurred on 0.03% of the sessions. Note that because on average, the empirical

firing rate is equal to the true firing rate, $f(t) = \langle s(t) \rangle$, changes in the firing rate are driven, on

average, by the covariance of reward and the empirical firing rate: $\langle \Delta f(t) \rangle \equiv \langle f(t+1) - f(t) \rangle =$

$\eta \cdot \text{cov}\big(R(t), s(t)\big)$[26]. The estimated Q-values in Fig. S1 were computed from the actions and

rewards of the covariance model by assuming the Q-learning model (Eqs. 1 and 2).

The motor cortex recordings (Fig. S2)

The data in Fig. S2A-B was recorded from one female monkey (Macaca fascicularis) at 3 years of

age, using a 10x10 microelectrode array (Blackrock Microsystems) with 0.4mm inter-electrode

distance. The array was implanted in the arm area of M1, under anesthesia and aseptic conditions.

Behavioral Task: The Monkey sat in a behavioral setup, awake and performing a Brain Machine

Interface (BMI) and sensorimotor combined task. Spikes and Local Field Potentials were extracted

from the raw signals of 96 electrodes. The BMI was provided through real time communication

between the data acquisition system and a custom-made software, which obtained the neural data,

analyzed it and provided the monkey with the desired visual and auditory feedback, as well as the

food reward. Each trial began with a visual cue, instructing the monkey to make a small hand move

to express alertness. The monkey was conditioned to enhance the power of beta band frequencies

(20-30Hz) extracted from the LFP signal of 2 electrodes, receiving a visual feedback from the BMI

algorithm. When a required threshold was reached, the monkey received one of 2 visual cues and

following a delay period, had to report which of the cues it saw by pressing one of two buttons.

Food reward and auditory feedback were delivered based on correctness of report. The duration of

a trial was on average 14.2s. The inter-trial-interval was 3s following a correct trial and 5s after

error trials. The data used in this paper, consists of spiking activity of 89 neurons recorded during

the last second of inter-trial-intervals, taken from 600 consecutive trials in one recording session.

Pairwise correlations were comparable to previously reported[54], $r_{SC} = 0.047 \pm 0.17$ (SD),

($r_{SC} = 0.037 \pm 0.21$ for pairs of neurons recorded from the same electrode).

Animal care and surgical procedures complied with the National Institutes of Health Guide for the

Care and Use of Laboratory Animals and with guidelines defined by the Institutional Committee

for Animal Care and Use at the Hebrew University.

The auditory cortex recordings (Fig. S2)

The auditory cortex recordings appearing in Fig. S2C-D are described in detail in [55]. In short,

membrane potential was recorded intracellularly from 39 neurons in the auditory cortex of

anesthetized rats. 125 experimental sessions were considered. Each session consisted of 370 50

msec tone bursts, presented every 300-1000 msec. For each session, all trials were either 300 msec

or 500 msec long. Trial length remained identical throughout a session and depended on smallest

interval between two tones in each session. Trials began 50 msec prior to tone burst. For spike

detection, data was high pass filtered with a corner frequency of 30Hz. Maximum points that were

higher than 60 times the median of the absolute deviation from the median were classified as

spikes.

The Basal ganglia recordings (Fig. 3 and S3)

The basal ganglia recordings that are analyzed in Figs. 3 and S3 are described in detail in [7]. In

short, rats performed a combination of a tone discrimination task and a reward-based free-choice

task. Extracellular voltage was recorded in the behaving rats from the NAc and VP using an

electrode bundle. Spike sorting was done using principal component analysis. In total, 148 NAc

and 66 VP neurons across 52 sessions were used for analyses (In 18 of the 70 behavioral sessions

there were no neural recordings).

Estimation of action-values from model choices and rewards

To imitate experimental procedures, we regressed the spike count on estimates of the action-

values, rather than the subjective action-values that underlay model behavior (to which the

experimentalist has no direct access). For that goal, for each session, we assumed that $Q_i(1) = 0.5$

and found the set of parameters $\hat{\alpha}$ and $\hat{\beta}$ that yielded the estimated action-values that best fit the

sequences of actions in each experiment by maximizing the likelihood of the sequence. Action-

values were estimated from Eq. (1), using these estimated parameters and the sequence of actions

and rewards. Overall, the estimated values of the parameters α and β were comparable to the actual

values used: on average, $\hat{\alpha} = 0.12 \pm 0.09$ (standard deviation) and $\hat{\beta} = 2.6 \pm 0.7$ (compare with

α=0.1 and β=2.5).

Exclusion of neurons

Following standard procedures, a sequence of spike-counts, either simulated or experimentally

measured was excluded due to low firing rate if the mean spike count in all blocks was smaller

than 1. This procedure excluded 0.02% (4/20,000) of the random-walk neurons. Considering the

auditory cortex recordings, we assigned each of the 125 spike counts to 40 randomly-selected

sessions. 23% of the neural recordings (29/125) were excluded in all 40 sessions. Of the remaining

96 recordings, 14% of the recordings × sessions were also excluded. Similarly, considering the

basal ganglia neurons, we assigned each of the 642 recordings (214×3 epochs) to 40 randomly-

selected sessions. 11% (74/(214×3)) of the recordings were excluded in all 40 sessions. Of the

remaining 568 recordings, 9% of the recordings × sessions were also excluded. None of the

simulated action-value neurons (0/20,000) or the motor cortex neurons (0/89) were excluded.

Statistical analyses

The computation of the t-values of the regression of the spike counts on the estimated Q-values

(as in Figs. 1, 2, 5, S1, S2, S3) was done using the following regression model:

$s(t) = \beta_0 + \beta_1 Q_1(t) + \beta_2 Q_2(t) + \epsilon(t)$ (6)

Where s(t) is the spike count in trial t, $Q_1(t)$ and $Q_2(t)$ are the estimated action-values in trial t,

$\epsilon(t)$ is the residual error in trial t and β_{0-2} are the regression parameters.

The computation of the t-values of the regression of the spike counts on the reward probabilities

in the trial design experiment (as in Fig. 4) was done using the following regression model:

$s(t) = \beta_0 + \beta_1 RP_1(t) + \beta_2 RP_2(t) + \epsilon(t)$ (7)

Where $s(t)$ is the mean spike count in the last 80 trials of the session, $RP_1(t)$ and $RP_2(t)$ are the

reward probabilities given action 1 or action 2 were chosen, respectively in the last 80 trials of the

session (in this experimental design RP could be 0.1,0.5 or 0.9), $\epsilon(t)$ is the residual error and β_{0-2}

are the regression parameters.

The computation of the t-values of the regression of the spike counts on *state* and *policy* in a trial

design experiment was done using the following regression model:

$s(t) = \beta_0 + \beta_1 [RP_1(t) + RP_2(t)] + \beta_2 [RP_1(t) - RP_2(t)] + \epsilon(t)$ (8)

All variables and parameters are the same as in Eq. (7)

All regression analyses were done using *regstats* in MATLAB (version 2016A).

To find neurons whose spike count in the last 20 trials is modulated by reward probability (Figs.

1B, 2A) we executed the Wilcoxon rank sum test, using *ranksum* in MATLAB. All tests were two-

tailed.

Permutation test (Fig. 3)

For each neuron, we computed the t-values of the regressions of its spike-count on estimated

action-values from sessions in Fig. 1E. Because the number of trials can affect the distribution of

t-values, we only considered in our analysis the first 170 trials of the 504 sessions longer or equal

to 170 trials. This number, which is approximately the median of the distribution of number of

trials per session, was chosen as a compromise between the number of trials per session and

number of sessions.

Two points are noteworthy. First, the distribution of the t-values of the regression of the spike

count of a neuron on all action-values depends on the neuron (see Fig. 3A). Similarly, the

distribution of the t-values of the regression of the spike counts of all neurons on an action-value

depends on the action-value (not shown). Therefore, the analysis could be biased in favor (or

against) finding action-value neurons if the number of neurons per session is different between

sessions. Second, this analysis does not address the correlated decision variables confound.

Comparison with permuted spike counts (Fig. S4)

In Fig. S4 we considered the experiment and analysis described in [6]. That experiment consisted

of four blocks, each associated with a different pair of reward probabilities, (0.72, 0.12), (0.12,

0.72), (0.21, 0.63) and (0.63, 0.21), appearing in a random order, with the better option changing

location with each block change. The number of trials in a block was preset, ranging between 35

and 45 with a mean of 40 (this is unlike the experiment described in Fig. 1, in which termination

of a block depended on performance).

First, we used Eqs. (1) and (2) to model learning behavior in this protocol. Then, we estimated the

action-values according to choice and reward sequences, as in Fig. 1. These estimated action-

values were used for regression of the spike counts of the random-walk, motor cortex, auditory

cortex, and basal ganglia neurons in the following way: each spike count sequence was randomly

assigned to a particular estimation of a pair of action-values from one session. The spike count

sequence was regressed on these estimated action-values. The resultant t-values were compared

with the t-values of 1,000 regressions of the spike-count, permuted within each block, on the same

Q-values. The p-value of this analysis was computed as the percentage of t-values from the

permuted spike-counts that were higher in absolute value than the t-value from the regression of

the original spike count. The significance boundary was set at $p<0.025$ [6]. Neurons with at least

one significant regression coefficient (rather than exactly one significant regression coefficient)

were classified as action-value modulated neurons[6].

Data Availability

The data of the basal ganglia recordings is available online at https://groups.oist.jp/ncu/data and

was analyzed with permission from the authors. Motor cortex and auditory cortex data is available

upon request.

Code Availability

Custom MATLAB scripts used to create simulated neurons and to analyze data are also available

upon request.

Author Contributions

L.E.D. and Y.L. designed the analysis and wrote the paper

Acknowledgements

We are extremely grateful to Oren Peles, Eilon Vaadia and Uri Werner-Reiss for providing us with

their motor cortex recordings, Bshara Awwad, Itai Hershenhoren, Israel Nelken for providing us

with their auditory cortex recordings, Kenji Doya and Makoto Ito for providing us with their basal

ganglia recordings, Mati Joshua, Gianluigi Mongillo and Roey Schurr for careful reading of the

manuscript and helpful comments and Inbal Goshen and Hanan Shteingart for discussions. This

work was supported by the Israel Science Foundation (Grant No. 757/16), DFG and the Gatsby

Charitable Foundation.

References

1.   Sutton RS, Barto AG. Reinforcement Learning: An Introduction. MIT Press; 1998.

2.   Louie K, Glimcher PW. Efficient coding and the neural representation of value. Ann N Y Acad Sci. 2012;1251: 13–32. doi:10.1111/j.1749-6632.2012.06496.x

3.   Schultz W. Neuronal Reward and Decision Signals: from Theories to Data. Physiol Rev. 2015;95: 853–951. doi:10.1152/physrev.00023.2014

4.   Samejima K, Ueda Y, Doya K, Kimura M. Representation of Action-Specific Reward Values in the Striatum. Science. 2005;310: 1337–1340. doi:10.1126/science.1115270

5.   Lau B, Glimcher PW. Value Representations in the Primate Striatum during Matching Behavior. Neuron. 2008;58: 451–463. doi:10.1016/j.neuron.2008.02.021

6.   Kim H, Sul JH, Huh N, Lee D, Jung MW. Role of Striatum in Updating Values of Chosen Actions. J Neurosci. 2009;29: 14701–14712. doi:10.1523/JNEUROSCI.2728-09.2009

7.   Ito M, Doya K. Validation of Decision-Making Models and Analysis of Decision Variables in the rat basal ganglia. J Neurosci. 2009;29: 9861–9874. doi:10.1523/JNEUROSCI.6157-08.2009

8.   Wang AY, Miura K, Uchida N. The dorsomedial striatum encodes net expected return, critical for energizing performance vigor. Nat Neurosci. 2013;16: 639–47. doi:10.1038/nn.3377

9.   Kim H, Lee D, Jung MW. Signals for Previous Goal Choice Persist in the Dorsomedial , but Not Dorsolateral Striatum of Rats. J Neurosci. 2013;33: 52–63. doi:10.1523/JNEUROSCI.2422-12.2013

10.   Ito M, Doya K. Distinct Neural Representation in the Dorsolateral, Dorsomedial, and

Ventral Parts of the Striatum during Fixed- and Free-Choice Tasks. J Neurosci. 2015;35:

3499–3514. doi:10.1523/JNEUROSCI.1962-14.2015

11.   Ito M, Doya K. Parallel Representation of Value-Based and Finite State-Based Strategies

in the Ventral and Dorsal Striatum. PLoS Comput Biol. 2015;11: 1–25.

doi:10.1371/journal.pcbi.1004540

12.   Funamizu A, Ito M, Doya K, Kanzaki R, Takahashi H. Condition interference in rats

performing a choice task with switched variable- and fixed-reward conditions. Front

Neurosci. 2015;9: 1–14. doi:10.3389/fnins.2015.00027

13.   Her ES, Huh N, Kim J, Jung MW. Neuronal activity in dorsomedial and dorsolateral

striatum under the requirement for temporal credit assignment. Sci Rep. Nature Publishing

Group; 2016;6: 1–11. doi:10.1038/srep27056

14.   Cai X, Kim S, Lee D. Heterogeneous Coding of Temporally Discounted Values in the

Dorsal and Ventral Striatum during Intertemporal Choice. Neuron. Elsevier Inc.; 2011;69:

170–182. doi:10.1016/j.neuron.2010.11.041

15.   Tai L-H, Lee AM, Benavidez N, Bonci A, Wilbrecht L. Transient stimulation of distinct

subpopulations of striatal neurons mimics changes in action value. Nat Neurosci. 2012;15:

1281–9. doi:10.1038/nn.3188

16.   Lee E, Seo M, Monte OD, Averbeck BB. Injection of a Dopamine Type 2 Receptor

Antagonist into the Dorsal Striatum Disrupts Choices Driven by Previous Outcomes , But

Not Perceptual Inference. J Neurosci. 2015;35: 6298–6306.

doi:10.1523/JNEUROSCI.4561-14.2015

17.   Fitzgerald THB, Friston KJ, Dolan RJ. Action-Specific Value Signals in Reward-Related

847   Regions   of   the   Human   Brain.   J   Neurosci.   2012;32:   16417–16423.

848   doi:10.1523/JNEUROSCI.3254-12.2012

849   18.   Guitart-Masip M, Huys QJM, Fuentemilla L, Dayan P, Duzel E, Dolan RJ. Go and no-go

850   learning in reward and punishment: Interactions between affect and effect. Neuroimage.

851   Elsevier Inc.; 2012;62: 154–166. doi:10.1016/j.neuroimage.2012.04.024

852   19.   Wunderlich K, Rangel A, O'Doherty JP. Neural computations underlying action-based

853   decision making in the human brain. PNAS. 2009;106: 17199–17204.

854   20.   Kim YB, Huh N, Lee H, Baeg EH, Lee D, Jung MW. Encoding of Action History in the

855   Rat Ventral Striatum. J Neurophysiol. 2007;98: 3548–3556. doi:10.1152/jn.00310.2007.

856   21.   Kim S, Cai X, Hwang J, Lee D. Prefrontal and striatal activity related to values of objects

857   and locations. Front Comput Neurosci. 2012;6: 1–13. doi:10.3389/fnins.2012.00108

858   22.   Stalnaker TA, Calhoon GG, Ogawa M, Roesch MR, Schoenbaum G. Neural correlates of

859   stimulus – response and response – outcome associations in dorsolateral versus dorsomedial

860   striatum. Front Integr Neurosci. 2010;4: 1–18. doi:10.3389/fnint.2010.00012

861   23.   Shteingart H, Loewenstein Y. Reinforcement learning and human behavior. Curr Opin

862   Neurobiol. 2014;25: 93–98. doi:10.1016/j.conb.2013.12.004

863   24.   Granger CWJ, Newbold P. Spurious Regressions in Econometrics. J Econom. 1974;2: 111–

864   120. doi:10.1016/0304-4076(74)90034-7

865   25.   Phillips PCB. Understanding spurious regressions in econometrics. J Econom. 1986;33:

866   311–340. doi:10.1016/0304-4076(86)90001-1

26.   Loewenstein Y, Seung HS. Operant matching is a generic outcome of synaptic plasticity
based on the covariance between reward and neural activity. PNAS. 2006;103: 15224–
15229.

27.   Loewenstein Y. Robustness of Learning That Is Based on Covariance- Driven Synaptic
Plasticity. PLoS Comput Biol. 2008;4: 1–10. doi:10.1371/journal.pcbi.1000007

28.   Loewenstein Y. Synaptic theory of Replicator-like melioration. Front Comput Neurosci.
2010;4: 1–12. doi:10.3389/fncom.2010.00017

29.   Neiman T, Loewenstein Y. Covariance-Based Synaptic Plasticity in an Attractor Network
Model Accounts for Fast Adaptation in Free Operant Learning. J Neurosci. 2013;33: 1521–
1534. doi:10.1523/JNEUROSCI.2068-12.2013

30.   Mongillo G, Shteingart H, Loewenstein Y. The misbehavior of reinforcement learning. Proc
IEEE. 2014;102: 528–541. doi:10.1109/JPROC.2014.2307022

31.   Padoa-Schioppa C. Neurobiology of Economic choice: A Good-Based Model. Annu Rev
Neurosci. 2011;34: 333–359. doi:10.1146/annurev-neuro-061010-113648

32.   Seung HS. Learning in Spiking Neural Networks by Reinforcement of Stochastic Synaptic
Transmission. Neuron. 2003;40: 1063–1073. doi:10.1016/S0896-6273(03)00761-X

33.   Fiete IR, Fee MS, Seung HS. Model of Birdsong Learning Based on Gradient Estimation
by Dynamic Perturbation of Neural Conductances. J Neurophysiol. 2007;98: 2038–2057.
doi:10.1152/jn.01311.2006.

34.   Urbanczik R, Senn W. Reinforcement learning in populations of spiking neurons. Nat
Neurosci. 2009;12: 250–252. doi:10.1038/nn.2264

35.   Darshan R, Leblois A, Hansel D. Interference and Shaping in Sensorimotor Adaptations
with Rewards. PLoS Comput Biol. 2014;10: 1–20. doi:10.1371/journal.pcbi.1003377

36.   Fremaux N, Sprekeler H, Gerstner W. Functional Requirements for Reward-Modulated
Spike-Timing-Dependent Plasticity. J Neurosci. 2010;30: 13326–13337.
doi:10.1523/JNEUROSCI.6249-09.2010

37.   Williams RJ. Simple Statistical Gradient-Following Algorithms for Connectionist
Reinforcement Learning. Mach Learn. 1992;8: 229–256.

38.   Woolrich MW, Ripley BD, Brady M, Smith SM. Temporal autocorrelation in univariate
linear modeling of FMRI data. Neuroimage. 2001;14: 1370–86.

39.   Arbabshirani MR, Damaraju E, Phlypo R, Plis S, Allen E, Ma S, et al. Impact of
autocorrelation on functional connectivity. Neuroimage. 2014;102: 294–308.

40.   Wilson RC, Niv Y. Is Model Fitting Necessary for Model-Based fMRI? PLoS Comput Biol.
2015;11: 1–21. doi:10.1371/journal.pcbi.1004237

41.   Gouvea TS, Monteiro T, Motiwala A, Soares S, Machens CK, Paton JJ. Striatal dynamics
explain duration judgments. Elife. 2015;4: e11386. doi:10.7554/eLife.11386

42.   Mello GBM, Soares S, Paton JJ. A Scalable Population Code for Time in the Striatum. Curr
Biol. 2015;25: 1113–1122. doi:10.1016/j.cub.2015.02.036

43.   Li J, Daw ND. Signals in Human Striatum Are Appropriate for Policy Update Rather than
Value Prediction. J Neurosci. 2011;31: 5504–5511. doi:10.1523/JNEUROSCI.6316-
10.2011

44.   FitzGerald THB, Schwartenbeck P, Dolan RJ. Reward-Related Activity in Ventral Striatum

Is Action Contingent and Modulated by Behavioral Relevance. J Neurosci. 2014;34: 1271–

1279. doi:10.1523/JNEUROSCI.4389-13.2014

45.   Vo K, Rutledge RB, Chatterjee A, Kable JW. Dorsal striatum is necessary for stimulus-

912   value but not action-value learning in humans. Brain. 2014;137: 3129–3135.

913   doi:10.1093/brain/awu277

914   46.   McDonald R. J, White N. M. A Triple Dissociation of Memory Systems: Hippocampus,

915   Amygdala, and Dorsal Striatum. Behav Neurosci. 1993;107: 3–22. doi:10.1037/0735-

916   7044.107.1.3

917   47.   O'Doherty J, Dayan P, Schultz J, Deichmann R, Friston K, Dolan RJ. Dissociable Roles of

918   Ventral and Dorsal Striatum in Instrumental Conditioning. Science. 2004;304: 452–454.

919   doi:10.1126/science.1094285

920   48.   Thorn CA, Atallah H, Howe M, Graybiel AM. Differential Dynamics of Activity Changes

921   in Dorsolateral and Dorsomedial Striatal Loops During Learning. Neuron. 2010;66: 781–

922   795. doi:10.1016/j.neuron.2010.04.036.

923   49.   Yarom O, Cohen D. Putative cholinergic interneurons in the ventral and dorsal regions of

924   the striatum have distinct roles in a two choice alternative association task. Front Syst

925   Neurosci. 2011;5: 1–9. doi:10.3389/fnsys.2011.00036

926   50.   Ding L, Gold JI. Caudate encodes multiple computations for perceptual decisions. J

927   Neurosci. 2010;30: 15747–15759. doi:10.1523/JNEUROSCI.2894-10.2010.Caudate

928   51.   Pasquereau B, Nadjar A, Arkadir D, Bezard E, Goillandeau M, Bioulac B, et al. Shaping of

929   Motor Responses by Incentive Values through the Basal Ganglia. 2007;27: 1176–1183.

doi:10.1523/JNEUROSCI.3745-06.2007

52.   Day JJ, Jones JL, Carelli RM. Nucleus accumbens neurons encode predicted and ongoing

reward costs in rats. Eur J Neurosci. 2011;33: 308–321. doi:10.1111/j.1460-

9568.2010.07531.x

53.   Seo M, Lee E, Averbeck BB. Article Action Selection and Action Value in Frontal-Striatal

Circuits. Neuron. 2012;74: 947–960. doi:10.1016/j.neuron.2012.03.037

54.   Cohen MR, Kohn A. Measuring and interpreting neuronal correlations. Nat Neurosci.

2011;14: 811–819. doi:10.1038/nn.2842

55.   Hershenhoren I, Taaseh N, Antunes FM, Nelken I. Intracellular Correlates of Stimulus-

Specific Adaptation. J Neurosci. 2014;34: 3303–3319. doi:10.1523/JNEUROSCI.2166-

13.2014

Figure 1 Model of action-value neurons **(A)** Behavior of model in example session, composed of four blocks (separated by a dashed vertical line). The probabilities of reward for choosing actions 1 and 2 are denoted by the pair of numbers above the block. Black line denotes the probability of choosing action 1; vertical lines denote choices in individual trials, where red and blue denote actions 1 and 2, respectively, and long and short lines denote rewarded and unrewarded trials, respectively. **(B)** Neural activity. Firing rate (line) and spike-count (dots) of two example simulated action-value neurons in the session depicted in **(A)**. The red and blue-labeled neurons represent Q_1 and Q_2, respectively. Black horizontal lines denote the mean spike count in the last 20 trials of the block. Error bars denote the standard error of the mean. The two asterisks denote p<0.01 (rank sum test). **(C)** Values. Thick red and blue lines denote Q_1 and Q_2, respectively. Note that the firing rates of the two neurons in **(B)** are a linear function of these values. Thin red and blue lines denote the estimations of Q_1 and Q_2, respectively, based on the choices and rewards in **(A)**. The similarity between the thick and thin lines indicates that the parameters of the model can be accurately estimated from the behavior (see also Materials and Methods). **(D)** and **(E)** Population analysis. **(D)** Example of 500 simulated action-value neurons from randomly chosen sessions. Each dot corresponds to a single neuron and the coordinates correspond to the t-values of regression of the spike counts on the estimated values of the two actions. Color of dots denote significance: dark red and blue denote significant regression coefficient only on one estimated action-value, action 1 and action 2, respectively; light blue – significant regression coefficients on both estimated action-values with similar signs (ΣQ), orange - significant regression coefficients on both estimated action-values with opposite signs (ΔQ). Black – no significant regression coefficients. The two simulated neurons in **(B)** are denoted by squares. **(E)** Fraction of neurons in each category, estimated from 20,000 simulated neurons in 1,000 sessions. Error bars denote the standard error of the mean.

Figure 2 Erroneous detection of action-value representation in random-walk neurons **(A)** Two example random-walk neurons that appear is if they represent action-values. The red (top) and blue (bottom) lines denote the estimated action-values 1 and 2, respectively that were depicted in Fig. 1B,C. Gray lines and gray dots denote the firing rates and the spike counts of two example random-walk neurons that were randomly assigned to this simulated session. Black horizontal lines denote the mean spike count in the last 20 trials of the block. Error bars denote the standard error of the mean. The two asterisks denote p<0.01 (rank sum test). **(B)** and **(C)** Population analysis. Each random-walk neuron was regressed on the two estimated action-values, as in Figs.1D and 1E. The two random-walk neurons in **(A)** are denoted by squares in **(B)**. **(D)** Fraction of random-walk neurons classified as action-value neurons (red), and classified as state neurons (ΣQ) or policy neurons (ΔQ) (green) as a function of the magnitude of the diffusion parameter of random-walk (σ). Light red and light green are standard error of the mean. Dashed lines mark the results for σ=0.1, which is the value of the diffusion parameter used in **(A)-(C)**. Initial firing rate for all neurons in the simulations is $f(1) = 2.5$Hz.

Figure 3 Permutation analysis **(A)** Red and blue correspond to red and blue - labeled neurons in fig. 1B, respectively. Up-pointing arrows denote the t-values from regressions on the estimated action values from the session in which the neuron was simulated (depicted in Fig. 1A). The red and blue histograms denote the t-values of the regressions of the spike-count on estimated action-values from *different* sessions in Fig. 1E (Materials and Methods). Dashed black lines denote the 5% significance boundary. We say that the regression coefficient of neural activity on an action-value is significant in this analysis if it exceeds these significance boundaries. Note that because of the temporal correlations, these significance boundaries are larger than ±2 (the significance boundaries in Figs. 1,2). According to this permutation test the red-labeled but not the blue-labeled neuron is classified as an action-value neuron **(B)** Fraction of neurons classified in each category using the permutation analysis for the action-value neurons (green, Fig. 1), random-walk neurons (yellow, Fig. 2). Dashed line denotes chance level for action-value 1 or 2 classification. Error bars denote the standard error of the mean. The test correctly identifies 29% of actual action-value neurons as such, while classification of random walk neurons was at chance level **(C)** Light orange, fraction of basal ganglia neurons from [7] classified in each category when regressing the spike count of 214 basal ganglia neurons in three different experimental epochs on the estimated action-values associated with their experimental session. This analysis classified 32% of neurons as representing action-value. Dark orange, fraction of basal ganglia neurons classified in each category when applying the permutation analysis. This test classified 3.6% of neurons as representing action-value. Dashed line denotes significance level of p<0.01.

Figure 4 A solution for the temporal correlations confound that is based on trial design. **(A)** A Q-learning model was simulated in sessions of 280 trials, where the original reward probabilities from Fig. 1A were associated with different cues and appeared randomly. Learning was done separately for each cue. Top panel: The first 20 trials in an example session. Background colors denote the reward probabilities in each trial. Black circles denote the learned value of action-value 1 in each trial. Top and bottom black lines denote choices of action 1 and 2, respectively. Long and short lines denote rewarded and unrewarded trials, respectively. Bottom panels: Two examples of the grouping of trials with the same reward probabilities to show the continuity in learning. Note that the action-value changes only when action 1 is chosen because it is the action-value associated with action 1. **(B-E)** population analysis. Action-value neurons were simulated from the model in **(A)** similarly to the action-value neurons in Fig. 1. Random-walk neurons were simulated similarly to the random-walk neurons in Fig. 2. For each neuron, the spike-counts in the last 80 trials of the session were regressed on the reward probabilities (see Materials and Methods). **(B)** and **(C)** population analysis for action-value neurons. Legend and number of neurons is the same in Figs. 1D-e. This analysis correctly identifies 54% of action-value neurons as such **(D)** and **(E)** population analysis for random-walk neurons. Legend and number of neurons is the same in Figs. 2B-C. Classification of random-walk neurons as representing action-value is at 10%, as would be expected from chance level with two tests with 5% significance threshold.

Figure 5 Erroneous detection of action-value representation in policy neurons. **(A)** Behavior of model in example session, same as in Fig. 1A for the direct policy model. **(B)** Red and blue lines denote "action-value" 1 and 2 respectively, calculated from the choices and rewards in **(A)**. Note that the model learned without any explicit or implicit calculation of action-values. The calculation of these action-values is based on the fitting of Eq. (1) to the learning behavior. **(C)** Strong correlation between policy from the direct policy algorithm depicted in **(A)** and action-values calculated by fitting Eq. (1) to behavior. The three panels depict probability of choice as a function of the difference between the calculated action-values (left), "Q_1" (center) and "Q_2" (right). This correlation can cause policy neurons to be erroneously classified as representing action-value **(D)** and **(E)** Population analysis, same as in Figs.1D and 1E for the policy neurons.

Figure 6 Possible solution for the policy and state confounds. **(A)** The Q-learning behavioral model (Eqs. 1 and 2) was simulated in 1,000 sessions of 280 trials each, where the reward probabilities were associated with different cues and appeared randomly, as in Fig. 4. Learning occurred separately for each cue. In each session 20 action-value neurons, whose firing rate is proportional to the action-values (as in Fig. 1) were simulated. For each neuron, the spike-counts in the last 80 trials of each session were regressed on the sum of the reward probabilities (ΣQ; state) and the difference of the reward probabilities (ΔQ; policy, see Materials and Methods). Each dot denotes the t-values of the two regression coefficients of a single neuron (only 500, randomly-selected neurons are shown). Dashed lines denote the 5% significance boundaries of the regression coefficients. Neurons that had significant regression coefficients with *both* policy and state were identified as action-value neurons. Colors as in Fig. 1D. **(B)** Population analysis revealed that 22% of the action-value neurons were identified as such. Error bars are the standard error of the mean. **(C)** Same as in **(A)** with random-walk neurons, as in Fig. 2. **(D)** Population analysis revealed that less than 1% of the random-walk neurons were erroneously classified as representing action-values. **(E)** To test the policy neurons, we simulated a direct-policy learning algorithm (as in Fig. 5) in the same sessions as in **(A-D)**. Learning occurred separately for each cue. In each session 20 policy neurons, whose firing rate is proportional to the probability of choice (as in Fig. 5) were simulated. As in **(A-D)**, the spike-counts in the last 80 trials of each session were regressed on the sum and difference of the reward probabilities and each dot denotes the t-values of the two regression coefficients of a single neuron. **(F)** As expected, only 4% of the policy neurons were erroneously classified as representing action-values.