The type IVa pilus machinery is pre-installed during cell division

Tyson Carter[1], Ryan N.C. Buensuceso[1], Stephanie Tammam[2], Ryan P. Lamers[1],

Hanjeong Harvey[1], P. Lynne Howell[2,3#], and Lori L. Burrows[1#].

[1]Department of Biochemistry and Biomedical Sciences and the Michael G. DeGroote

Institute for Infectious Disease Research, McMaster University, ON, CANADA;

[2]Program in Molecular Structure & Function, The Hospital for Sick Children; and

[3]Department of Biochemistry, University of Toronto, ON, CANADA.

Running title: Pre-installation of the type IV pilus machinery

[#]To whom correspondence should be addressed:

Dr. Lori L. Burrows, 4H18 Health Sciences Centre, 1200 Main St. West, Hamilton, ON,

L8N 3Z5 CANADA, Tel: (905)-525-9140 ext: 22029, Fax: (905) 522-9033, E-mail:

burrowl@mcmaster.ca

Dr. P. Lynne Howell, 20-9-715 Peter Giligan Centre for Research and Learning, 686 Bay

St., Toronto, ON M5G 0A4 CANADA, Tel: (416) 813-5378, Fax: (416) 813-5022, E-mail:

howell@sickkids.ca

ABSTRACT

Type IV pili (T4aP) are ubiquitous microbial appendages used for adherence, twitching

motility, DNA uptake, and electron transfer. Many of these functions depend on dynamic

assembly and disassembly of the pilus by a megadalton-sized, cell envelope-spanning

protein complex located at the poles of rod-shaped bacteria. How the T4aP assembly

complex becomes integrated into the cell envelope in the absence of dedicated

peptidoglycan (PG) hydrolases is unknown. After ruling out potential involvement of

housekeeping PG hydrolases in installation of the T4aP machinery in *P. aeruginosa*, we

discovered that key components of inner (PilMNOP) and outer (PilQ) membrane

subcomplexes are recruited to future sites of cell division. Mid-cell recruitment of a

fluorescently tagged alignment subcomplex component, mCherry-PilO, depended on

PilQ secretin monomers - specifically, their N-terminal PG-binding AMIN domains. PilP,

which connects PilO to PilQ, was required for recruitment, while PilM, which is

structurally similar to divisome component FtsA, was not. Recruitment preceded secretin

oligomerization in the outer membrane, as loss of the PilQ pilotin, PilF, had no effect on

localization. These results were confirmed in cells chemically blocked for cell division

prior to outer membrane invagination. The hub protein FimV and a component of the

Polar Organelle Coordinator complex - PocA - were independently required for mid-cell

recruitment of PilO and PilQ. Together, these data reveal an integrated, energy-efficient

strategy for the targeting and pre-installation – rather than retrofit - of the T4aP system

into nascent poles, without the need for dedicated PG-remodelling enzymes.

IMPORTANCE

  The peptidoglycan (PG) layer of bacterial cell envelopes has limited porosity, representing a physical barrier to insertion of large protein complexes involved in secretion and motility. Many systems include dedicated PG hydrolase components that create space for their insertion, but the ubiquitous type IVa pilus (T4aP) system lacks such an enzyme. Instead, we found that components of the T4aP system are recruited to future sites of cell division where they can be incorporated into the cell envelope during the formation of new poles, eliminating the need for PG hydrolases. Targeting depends on the presence of septal PG-binding motifs in specific components, as removal of those motifs causes delocalization. This pre-installation strategy for the T4aP assembly system ensures that both daughter cells are poised to extrude pili from new poles as soon as they separate from one another.

INTRODUCTION

  The cell envelopes of most gram-negative bacteria are comprised of an inner membrane, a peptidoglycan (PG) layer found in the periplasm, and an outer membrane (OM). A variety of large protein complexes, including motility machines, secretion systems, and efflux pumps span all these layers (1), but in many cases the mechanisms used to integrate these systems into the cell envelope remain uncharacterized. In many cases, they are integrated at specific locations in the cell such as the poles, which is important for their function.

  The machinery that assembles and disassembles type IVa pili (T4aP) is among the most common cell envelope-spanning complexes found in gram-negative bacteria.

T4aP are long, thin protein fibres used for adherence, biofilm formation and a flagellum-

independent form of motility known as twitching, defined as repeated cycles of pilus

extension, adherence to a surface, and retraction of the pilus fibre (2, 3). *P. aeruginosa*

T4aP are important for surface mechanosensing and consequent up-regulation of

virulence factor expression (4), and for the delivery of toxins by other secretion systems

(5, 6). In the absence of T4aP, *P. aeruginosa* has reduced pathogenicity (5).

The T4aP machinery is composed of four subcomplexes that together form a

dynamic cylindrical machine (3, 7, 8). The inner membrane motor and alignment

subcomplexes form the platform for assembly and disassembly of the pilus, and interact

directly with the secretin that allows for pilus extrusion through the OM. The OM

lipoprotein PilF promotes formation of the secretin, a highly stable oligomer of 14 PilQ

subunits that form a gated channel in the OM (9, 10). The N-terminus of each PilQ

monomer contains two tandem amidase N-terminal (AMIN) domains, involved in binding

peptidoglycan (11-13). The alignment subcomplex composed of PilMNOP connects the

motor subcomplex with the secretin, and participates in pilus extension and retraction

(14-17). PilM is a cytoplasmic, actin-like protein that clamps tightly onto the short amino-

terminus of the inner membrane protein, PilN, which in turn forms heterodimers with

inner membrane protein, PilO (14, 18, 19). The inner membrane lipoprotein PilP

interacts with PilNO heterodimers via its long unstructured N-terminus, and with the

periplasmic N0 domain of PilQ via its C-terminal homology region (HR) domain,

completing the transenvelope complex (17, 20).

T4aP are usually located at the poles of rod-shaped cells, which promotes

adherence by minimizing surface area and thus electrostatic repulsion (3). How the

multi-protein T4aP system is initially targeted to and integrated into the poles of *P.*

aeruginosa cells are not well understood. Type III (T3SS) and Type IV (T4SS) secretion

systems have dedicated PG-remodelling enzymes that allow for retrofitting of the

complexes into the cell envelope (1). There are no such enzymes associated with the

T4aP system, suggesting that it uses an alternate pathway for cell envelope integration.

Here we used fluorescent fusions to an informative alignment subcomplex component,

PilO, and to the secretin monomer, PilQ, to track and quantify their localization in *P.*

aeruginosa wild type and mutant backgrounds. The results show that PilO and

associated components are recruited to future sites of cell division by the concerted

action of least three pathways. We propose that the T4aP machinery is preinstalled

during formation of nascent cell poles, a strategy that eliminates the need for dedicated

cell-wall-processing enzymes.

RESULTS

mCherry-PilO localizes to cell poles and future cell division sites

When selecting components of the T4aP machinery to track using fluorescent

fusions, we considered the known interactions and stoichiometry of these proteins and

how a fusion might affect function. PilM is cytoplasmic, but has multiple interaction

partners (14, 19, 20), including PilN, whose cytoplasmic N-terminus binds to a groove on

PilM; thus, we excluded both as candidates. PilN interacts along most of its remaining

length with PilO (15, 21), but PilO's cytoplasmic N-terminus is poorly conserved among

T4aP-expressing species (16), supporting the observation that it appears to have no

cytoplasmic interaction partners. Therefore, PilO was selected as a fusion candidate for

localization of the alignment subcomplex. To maintain the physiological expression

levels and stoichiometry that are important for function (14), a *pilO* construct encoding

an in-frame fusion of mCherry to the N-terminus of PilO was used to replace the WT

version at the native *pilO* locus (**Fig. 1A**). Stability and functionality of the fusion protein

were verified by western blot with anti-PilO and anti-mCherry antisera, and assessment

of twitching motility, respectively (**Fig. 1B**).

mCherry-PilO localized to both poles, as well as to the midpoint of cells

undergoing division (**Fig. 1C**). To more clearly visualize the predivisional localization

pattern of mCherry-PilO, cells were treated with sub-inhibitory levels of cefsulodin, which

inhibits PBP3 (FtsI), a late-stage PG transpeptidase, leading to division arrest and

filamentation (22). In filamented cells, mCherry-PilO localized to the poles and to

regularly spaced foci, suggestive of recruitment to future sites of cell division (**Fig. 1D**).

Quantification of mCherry-PilO localization in wild type cells (**Fig. 1E**) confirmed its

predominately bipolar localization.

To provide further evidence that mCherry-PilO was recruited to sites of cell

division, the *minC* gene, which encodes a key component of the oscillating Min system

required for inhibition of FtsZ polymerization at non-mid-cell locations (23, 24) was

deleted in the mCherry-PilO strain. Cells lacking MinC exhibit aberrantly localized septa,

producing mini-cells due to unequal cell division (**Fig. 2A**). As predicted, the mCherry-

PilO fusion continued to localize to polar and septal sites despite their aberrant

placement (**Fig. 2B**). Together, these data suggest that when expressed under

physiological conditions, PilO – and by inference, its interaction partners, PilMN and PilP

(20) – are recruited to future sites of cell division.

PilQ monomers are required for T4aP alignment subcomplex localization

To define the mechanism of recruitment of the T4aP alignment subcomplex to

sites of cell division, we first examined the effects of deleting specific subcomplex

components. The cytoplasmic member of the alignment subcomplex, PilM, has

pronounced structural similarity ($C\alpha$ root mean square deviation = 1.9 Å over 257

residues) to the early divisome protein, FtsA, a peripheral inner membrane protein

whose interaction with FtsZ tethers the latter to the membrane (18, 19). Based on its

structural mimicry of FtsA and its peripheral membrane localization via binding of PilN's

amino terminus, we previously hypothesized (25) that PilM – and thus the alignment

subcomplex – might be recruited to mid-cell through interactions with the FtsZ ring.

However, mCherry-PilO exhibited polar and mid-cell localization in a *pilM* mutant (**Fig.

3AB**), ruling out PilM as a driver of recruitment.

The PilMNOP alignment subcomplex connects to the secretin via interaction of

the C-terminal HR domain of PilP with the periplasmic N0 domain of PilQ (20).

Preceding the N0 domain in *P. aeruginosa* PilQ are two tandem AMIN domains,

predicted to bind septal PG (11-13). To test their PG-binding ability, we cloned,

expressed, and purified 3 fragments of PilQ's N-terminal region (the AMIN domains

alone; the N0-N1 domains alone; and all 4 domains together), and performed pulldown

assays using purified *P. aeruginosa* PG (**Supplementary Fig. S1**). In the absence of

PG, all fragments remained in the soluble fraction; however, when insoluble PG was

present, only those fragments containing the AMIN domains were also found in the

insoluble fraction, indicating that they bind PG.

Since *pilMNOPQ* are expressed from a single operon, we hypothesized that

recruitment of PilMNOP to sites of cell division might occur through a series of PilMNOP-

PilQ-septal PG interactions. We first confirmed that PilQ-mCherry was localized to the

poles and to regularly spaced foci in filamented cells, a pattern similar to mCherry-PilO

(**Supplementary Fig. S2**). When we deleted *pilQ*, mCherry-PilO became delocalized

(**Fig. 3CD**). To confirm that loss of mCherry-PilO localization in the *pilQ* strain was

indirectly due to loss of PilP-PilQ interactions, *pilP* was deleted in the mCherry-PilO

strain. In the *pilP* mutant, mCherry-PilO was delocalized (**Fig. 3EF**), supporting the

hypothesis that recruitment of PilQ to mid-cell is the primary event, and the remaining

alignment subcomplex components are recruited via their interaction with PilP, and its

interaction with PilQ.

To test whether its AMIN domains were responsible for localization of PilQ and its

interaction partners to mid-cell and polar positions, we complemented a *pilQ* mutant

expressing mCherry-PilO with full length *pilQ*, or a construct expressing only the N0-N1

and secretin domains, previously shown to form stable but non-functional secretin

oligomers (**Fig. 4A**) (9). Full length PilQ restored the wild type pattern of mCherry-PilO

localization (**Fig. 4BC**), but the AMIN-deficient PilQ did not (**Fig. 4DE**). These data show

that localization of T4aP assembly components is dependent upon PilQ's AMIN

domains, which interact with PG (**Supplementary Fig. S1**).

Localization of mCherry-PilO precedes PilQ oligomerization

The *Myxococcus xanthus* T4aP system was proposed to assemble in an 'outside-

in' manner, with PilMNOP alignment subcomplexes docking onto the PilQ secretin after

its assembly in the OM (26). Those data are not consistent with our observation that

mCherry-PilO localized to regularly spaced foci in chemically-filamented *P. aeruginosa*

cells (**Fig. 1**), where the OM is not yet invaginated due to arrest of cell division. To test

whether secretin oligomerization was a necessary prerequisite for recruitment, we

examined localization of PilQ-mCherry in mutants lacking the OM lipoprotein PilF. In the

absence of PilF, PilQ remains in a monomeric state in the inner membrane (9), but its

localization pattern was unchanged compared to wild type (**Supplementary Fig. S2AB**).

Similarly, localization of mCherry-PilO in the *pilF* background resembled wild type (**Fig.**

5). These data confirm that localization of T4aP components to mid-cell – while PilQ-

dependent – precedes secretin oligomerization.

Additional factors modulate T4aP assembly machinery localization

FimV is a T4aP-associated protein required for twitching motility and has a highly

conserved PG-binding LysM motif in its periplasmic N-terminal domain (27, 28). We

showed previously (28) that PilQ multimer formation is reduced in a *fimV* mutant, which

made FimV of interest for this study. In the absence of FimV, mCherry-PilO was

delocalized (**Fig. 6AB**), as was PilQ-mCherry (**Supplementary Fig. S2C**). We

previously generated a strain expressing a variant of FimV with an in-frame deletion of

its LysM motif (28). mCherry-PilO (**Fig. 6CD**) was delocalized in the $fimV_{\Delta LysM}$ strain, as

was the majority of PilQ-mCherry (**Supplementary Fig. S2D**). These data suggest that

interactions of both PilQ and FimV with PG are important steps in the localization

process, and that polar localization of PilQ depends in part on FimV.

During a mutant screen looking for factors required for correct localization of *P.*

aeruginosa's single polar flagellum, Cowles and colleagues (29) identified a putative

protein complex that they named PocAB -TonB3 for 'polar organelle coordinator'. Loss

of TonB3 had previously been shown to affect twitching motility (30), but the mechanism

was unknown. All three components were required for expression and/or polar

localization of T4aP. *pocA* mutants made a few, non-polar pili, while *pocB* and *tonB3*

mutants failed to produce pili. The mechanism by which this system controls polar

localization of motility organelles remains unclear, since the Poc proteins themselves

are not localized to the poles (29).

We focused on *pocA* as the mutant's ability to make a few pili suggested that the

T4aP assembly system was functional, and deleted it from cells expressing either

mCherry-PilO (**Fig. 7AB**) or PilQ-mCherry (**Supplementary Fig. S2E**). mCherry-PilO

was delocalized in the absence of *pocA*, and its wild type localization was re-established

upon complementation with *pocA in trans* (**Fig. 7CD**). PilQ-mCherry was delocalized in

the *pocA* background although some polar fluorescence remained (**Supplementary Fig.**

S2E). Interestingly, while both PocA and FimV were required for T4P assembly

component localization to polar and midcell locations, they appear to act independently

of one another. The fluorescent FimV fusion, which exhibits bipolar localization in wild

type cells, remained localized to the poles of a *pocA* mutant (**Fig. 8**).

DISCUSSION

The T4aP machinery spans all layers of the gram-negative cell envelope at the

poles of rod-shaped cells, but how this megadalton protein complex is installed in the

absence of dedicated PG hydrolyzing enzymes was unclear. Here we showed that

degradation of PG is unnecessary, because components of the *P. aeruginosa* T4P

alignment subcomplex and the secretin are recruited to sites of cell division where they

can be pre-installed, rather than retrofitted, into nascent poles. These data are

consistent with our observation that mutants lacking multiple housekeeping PG

hydrolases – which could potentially provide PG lytic activities for retrofitting in the

absence of system-specific enzymes – have correctly inserted, functional T4aP

machines, as demonstrated by their ability to twitch (**Supplementary Fig. S3**). Instead,

at least 3 different mechanisms participate in the process of T4aP polar localization in *P.*

aeruginosa, two of which (PilQ and FimV) depend on recognition of PG, as deletion of

PilQ's AMIN domains or FimV's LysM motif results in delocalization.

Cell poles are characterized by PG that has fewer stem peptides compared to

lateral wall PG, due to the activity of *N*-acetylmuramyl-L-alanine amidases that cleave

the amide bond between *N*-acetylmuramic acid and the stem peptides during daughter

cell separation (31-33). This denuded form of PG is the binding target for multiple

protein motifs, among them AMIN, SPOR, and LysM (28, 34-36). *P. aeruginosa* and *M.*

xanthus PilQ monomers have N-terminal AMIN domains (2 per monomer in *P.*

aeruginosa and 3 in *M. xanthus*), which in addition to targeting (37), may anchor secretin

complexes to the PG layer to counter the substantial forces generated during twitching

motility. *Neisseria gonorrhoeae* and *N. meningitidis* PilQ monomers also contain 2 AMIN

domains at their amino termini (11), yet they express peritrichous T4aP. *Neisseria* spp.

lack the cytoskeletal element MreB which patterns elongation of rod-shaped bacteria

(38), thus have a coccoid morphology. It's possible that the mechanism of T4aP

assembly system recruitment to sites of cell division is similar in *Neisseria* spp., but due

to their shape – the equivalent of two closely apposed poles – they appear to have

peritrichous T4aP distribution.

In the *P. aeruginosa* T4aP system, at least two different PG-binding proteins were

needed for correct positioning. We showed previously (28) that FimV binds PG via its

LysM motif, important for optimal secretin assembly. A number of recent studies showed

that FimV and related HubP proteins – which contain multiple protein-protein interaction

domains in addition to conserved LysM motifs – act as landmarks, responsible for polar

localization of other proteins, including those involved in regulation of T4aP function,

flagellar placement, and chromosome segregation (39-43). For example, *P. aeruginosa*

FimL – a regulatory protein important for T4aP assembly via its regulation of intracellular

cAMP levels – interacts with the C-terminal TPR motif of FimV (39, 44). Loss of FimV

also results in mislocalization of the core T4aP machinery (**Fig. 6**); thus FimV appears to

have both physical and regulatory roles in T4aP assembly and function.

Cowles *et al.* (29) reported that both the flagellum and T4aP were mislocalized in

P. aeruginosa mutants lacking the polar organelle coordinating (Poc) proteins, but the

mechanism remains enigmatic. The T4aP alignment subcomplex was delocalized in a

pocA mutant (**Fig. 7**) while FimV remained bipolar (**Fig. 8**), implying that they act

independently of one another – with the caveat that the FimV fusion used here was non-

functional in twitching motility. PocAB-TonB3 are homologous to the ExbBD-TonB

complex in *E. coli* (29) which energizes siderophore uptake across the OM via direct

interactions with the 'Ton box' on one beta strand of TonB-dependent receptors. The N0

domains of secretin monomers are structurally similar (2.3 Å root mean square deviation

over 65 residues) to the signalling domain of FpvA, a TonB-dependent siderophore

receptor in *P. aeruginosa* (45). This similarity hints at potential interactions between the

TonB3 component of the Poc complex and the N0 domain of PilQ monomers. We noted

that although deletion of *pocA* had no significant effect on cell morphology,

complementation with a plasmid-borne copy of the gene increased average cell length

by ~50% (**Supplementary Fig. S4**), suggesting that the Poc system might influence the

timing of cell division or affect PG remodelling. Subtle structural differences in PG

architecture in the absence of PocA could impact binding of proteins with PG targeting

motifs.

Mid-cell recruitment of mCherry-PilO was dependent on the secretin monomer

PilQ (**Fig. 3CD**) but deletion of the pilotin PilF had no effect (**Fig. 5**), showing that

recruitment and confinement of PilMNOPQ to cell division sites occur prior to secretin

oligomerization. Prior studies of *M. xanthus* T4aP assembly system formation suggested

that PilQ multimerization was essential for the recruitment of additional T4aP assembly

components, because loss of Tgl – the *M. xanthus* PilF homolog – abrogated polar

localization of PilQ and other T4aP proteins. These data supported an 'outside in' model

for *M. xanthus* T4aP system formation, dependent on PilQ oligomerization (26).

Interestingly, *M. xanthus* PilQ was suggested in that study to localize to the septum

either "late during cell division, or immediately after". These data are consistent with the

pattern of PilQ localization in *P. aeruginosa*: however, the pilotin proteins in *M. xanthus*

and *P. aeruginosa* appear to play somewhat different roles.

Together, the data led us to propose a pathway for T4aP assembly machinery

integration into the *P. aeruginosa* cell envelope in the absence of dedicated PG

hydrolases. We suggest that PilMNOPQ are co-translated, forming an IM-bound

subcomplex that diffuses laterally until the AMIN domains of PilQ monomers bind to

septal PG. FimV, which has the capacity to interact with septal PG via its Lys motif, may

also interact with components of the subcomplex to retain them at cell division sites. The

next steps are still unclear, as the oligomerization pathway for PilQ in the OM remains

unknown. We previously proposed (9) that co-translocation of the lipoprotein PilF and

PilQ monomers by the Lol system might promote PilQ insertion into the OM, where it

oligomerizes. In this new model, we suggest that PilF is translocated independently by

the Lol system to the OM, where it then scans for, binds to, and promotes translocation

of PilQ monomers from the inner to the OM during invagination in the final stages of cell

division.

Two lines of evidence support this new hypothesis. First, in *M. xanthus*, co-

incubation of *pilQ* and *tgl* (*pilF*) mutants, neither of which are able to twitch, allowed for

intercellular transfer of Tgl from the *pilQ* mutant to the *tgl* mutant and oligomerization of

the latter's PilQ monomers, restoring piliation and motility (46). Thus, Tgl already

present in the OM of the *pilQ* mutant could promote insertion and oligomerization of PilQ

monomers in the *tgl* mutant. Second, the ability of OM lipoproteins to interact with inner

membrane proteins across the periplasm is now well established, as exemplified by the

E. coli OM lipoproteins, LpoA and LpoB (47, 48). These proteins are essential for

stimulating the activities of the inner membrane PG synthases PBP1a and PBP1b,

respectively, through direct interactions with regulatory domains of those enzymes.

Once PilQ monomers reach the OM with the assistance of PilF, they may first form

conformational intermediates prior to maturation into stable, SDS-resistant oligomers. In

N. gonorrhoeae, certain point mutations in the secretin domain were associated with

formation of immature, leaky oligomers, rendering the cells more susceptible to beta

lactam antibiotics (49).

The above scenario for PilQ translocation suggests that the ~ 2 nm porosity of

PG (50) would not pose a physical barrier to secretin assembly, as individual monomers

(~77 kDa) are small enough to pass through its natural gaps. Retention of PilQ AMIN

interactions with septal PG during translocation and insertion of the C-terminal secretin

domain in the OM would ultimately place the AMIN domains 'under' the PG layer, where

they would be ideally positioned to counteract retraction forces imposed during twitching

motility. Because polar PG is less likely to be remodelled due to its limited side chain

content, the secretin would remain stably fixed to the cell wall. Studies of the half-life of

a single bacterial cell pole show that it can be hundreds of generations old (51); thus, it

is perhaps not surprising that secretin complexes are highly resistant to denaturation,

allowing them to last as long as the poles in which they are embedded.

Among the questions remaining is whether the interaction interface between PilP

and PilQ is altered during translocation of PilQ monomers to the OM. We established

that mid-cell localization of mCherry-PilO depends on PilP (**Fig. 3EF**), which interacts

with the N0 domain of PilQ monomers via its C-terminal HR domain (11, 20). PilP also

interacts with PilNO heterodimers via its long unstructured N-terminus (17). This flexible

tether might allow the C-terminal domain of PilP to remain connected to the N0 domain

of PilQ as secretin monomers transit from the inner to OM. Alternatively, PilP may be

repositioned relative to PilQ when the latter translocates to the OM. In the evolutionarily

related T2SS, multiple interaction interfaces between the HR domain of GspC (PilP) and

N0 of GspD (PilQ) have been identified through a combination of structural, biochemical,

and functional studies (45, 52, 53). These various interfaces – some of which are

mutually exclusive – were attributed to differences in experimental setup, but could also

reflect capture of different interaction states pre- and post-secretin oligomerization.

Studies to address these possibilities are currently underway.

MATERIALS & METHODS

Strains, media and growth conditions

Bacterial strains, plasmids, and primers used in this study are listed in Tables 1

and 2. *Escherichia coli* and *P. aeruginosa* were grown at 37°C in LB Lennox broth or on

LB 1.5% agar plates supplemented with antibiotics. The antibiotics and their respective

concentrations were: ampicillin (Ap), 100 µg/mL; carbenicillin (Cb), 200 µg/mL;

gentamicin (Gm), 15 µg/mL for *E. coli* strain and 30 µg/mL for *P. aeruginosa* strains,

unless otherwise stated. Plasmids were transformed by heat shock into chemically

competent *E. coli* cells or by electroporation into *P. aeruginosa* suspended in sterile

water. All constructs were verified by DNA sequencing (MOBIX Lab, McMaster

University).

Generation of *P. aeruginosa* mutants

Mutants were made using previously described methods (54). Deletion constructs

for generation of *minC* and *pilF* mutants were designed to include 500 nucleotides up

and downstream of the gene to be deleted, as well as the first and last 50 nucleotides of

the gene. Constructs were synthesized and cloned into pUC57 (Genscript, Piscataway,

NJ). Deletion inserts in pUC57 were subcloned into the pEX18Gm suicide vector (55)

using the 5' BamHI and the 3' HindIII restriction sites, and verified by DNA sequencing.

After verification, suicide vectors containing the mutant genes (*pilF* or *minC*) were

transformed into *E. coli* SM10 cells. Plasmids were then transferred by conjugation in a

1:9 ratio of *P. aeruginosa* to *E. coli*. The mixed culture was pelleted for 1 min at 2292 x

g, and the pellet was resuspended in a 100 µL aliquot of LB, spot-plated on LB agar,

and incubated overnight at 37 °C. The mating mixture was removed from the LB agar

using a sterile toothpick, resuspended in 1 mL of LB and the *E. coli* SM10 donor was

counterselected by plating the mixture on *Pseudomonas* isolation agar (PIA; Difco)

containing Gm (100 µg/ml). Gm-resistant *P. aeruginosa* colonies were streaked on LB

no salt plates containing sucrose (1% (w/v) bacto-tryptone, 5% (w/v) sucrose, 1.5%

agar, 0.5% (w/v) bacto-yeast extract) and incubated for 16 h at 30 °C. Select colonies

were then subcultured on LB agar and LB agar plus Gm, and Gm-sensitive colonies

were screened by PCR using appropriate primers (Table 2) to confirm integration.

Amplicons from colonies with the desired PCR profile were sequenced to confirm

incorporation of the mutation, and the *pilF* mutant was analyzed by western blot analysis

using an α-PilF antibody to verify loss of the gene product.

A *pilQ* KO construct containing an FRT-flanked Gm cassette was designed

previously (9). This construct was integrated into the PAK chromosome using a Flp-FRT

recombination system (55) and selected for using the methods described above with the

following modifications: following growth on sucrose, colonies were cultured on LB agar,

LB agar with Gm and LB agar plus Cb. Colonies that had undergone double

recombination, integrating *pilQ*::GmFRT into the chromosome without pEX18Ap (Cb

sensitive colonies), were selected. The Gm cassette was removed by Flp recombinase-

catalyzed excision, by conjugally transferring the Flp-expressing pFLP2 from SM10 cells

into PAK cells carrying the GmFRT-disrupted *pilQ* gene. SM10 cells were

counterselected on PIA containing Cb (200 µg/ml). pFLP2 was removed from *P.*

aeruginosa by streaking Cb-resistant colonies on LB agar, no salt, containing 5% (w/v)

sucrose for 16 h at 30 °C. Colonies were then cultured on LB agar and LB-agar

containing Gm or Cb. Cb and Gm sensitive colonies were screened by PCR using the

PilQ FRT-Chk primers listed in Table 2 to confirm the retention of the FRT scar. The *pilQ*

gene of mutant strains were sequenced to confirm correct incorporation of the mutation,

and analyzed by western blot with α-PilQ sera to confirm loss of the gene product.

The *pocA* deletion construct was created previously in the pEX18Tc suicide

vector (29). The *pocA* deletion insert was subcloned into pEX18Gm using EcoRI and

XbaI, and constructs verified by DNA sequencing. After verification, the suicide vector

containing the *pocA* deletion construct transformed into *E. coli* SM10 cells and the

mutation introduced into *P. aeruginosa* as described above.

Generation of a chromosomally encoded mCherry-PilO fusion

A construct encoding an fusion of mCherry (56) to the N-terminus of PilO, plus

700 nucleotides upstream of *pilO* and 700 nucleotides downstream of *pilO,* was

synthesized and cloned into pUC57 (Genscript, Piscataway, NJ). The mCherry-PilO-

encoding insert was subcloned into the suicide vector pEX18Gm using the 5' EcoRI and

the 3' HindIII restriction sites, constructs verified by DNA sequencing, and introduced

into chemically competent *E. coli* SM10 by heat shock. The plasmid was then

transferred by conjugation in a 1:9 ratio of *P. aeruginosa* (wild type or various T4aP

mutants) to *E. coli* as described above. Following counter selection and curing of

merodiploids by sucrose selection, integration of the mCherry-PilO fusion in Gm-

sensitive colonies was confirmed by PCR and DNA sequencing (Table 2). mCherry-PilO

expressing strains were analyzed using western blot analysis and twitching assays to

verify expression of an intact fluorescent fusion protein and – in the wild type – function

of the T4aP machinery, respectively.

Generation of complementation constructs

The *pilQ, pocA* and *pilF* genes were amplified by PCR using PAK chromosomal

DNA as the template (Table 2) and cloned into pBADGr (Table 1). The digested DNA

was purified and ligated into pBADGr with T4 DNA ligase, according to the

manufacturer's instructions. Ligation mixtures were then transformed into *E. coli* DH5α

cells and grown overnight at 37 °C on LB agar supplemented with the appropriate

antibiotics.

A construct encoding a truncated version of PilQ lacking its AMIN domains was

previously generated in the PAO1 strain (9). The same boundaries were used here to

generate a truncated version of PAK PilQ. The PilQ Clone Fwd and PilQ 5' Rev primers

were used to amplify the 5' end of the gene, excluding the first AMIN domain, and the

PilQ N0 Fwd and PilQ Clone Rev primers (Table 2) were used to amplify the remainder

of the gene beginning at the N0 domain and excluding the second AMIN domain.

Purified amplicons were digested with XbaI and ligated into pBADGr with T4 DNA

Ligase. The ligation product and pBADGr were then digested with EcoRI and HindIII and

ligated following the same protocol. The fidelity of the pBADGr-*pilQ* construct was

confirmed by sequencing using the pBADGr mcs Fwd and Rev primers (Table 2).

Since PilQ has a cleavable N-terminal signal sequence, and a C-terminus that is

exposed to the extracellular environment (9, 57), we designed a construct encoding an

internal fusion of mCherry to PilQ (between the first and second AMIN domains,

residues 132 and 133), which was synthesized and subcloned into pUC57 (Genscript).

The PilQ-mCherry-encoding insert was subcloned into the complementation vector

pBADGr (58) using the 5' EcoRI and the 3' HindIII restriction sites and verified by DNA

sequencing. The construct was then introduced into PAK *pilQ*::FRT cells by

electroporation. Expression of the PilQ-mCherry fusion was verified using western blot

analysis with α-PilQ and α-mCherry antibodies and function was verified using twitching

assays (59).

The *yfp* gene encoding Yellow Fluorescent Protein (YFP) was cloned into the

HindIII restriction site of pBADGr. Correct insertion of the pBADGr-*yfp* construct was

confirmed by DNA sequencing using mcs Fwd/Rev primers (Table 2). A version of *fimV*

lacking its stop codon was amplified using the FimV Clone Fwd/Rev primers (Table 2)

and cloned into pBADGr in frame with the *yfp* gene by digesting both the vector and

insert with KpnI and XbaI restriction enzymes and ligating with T4 DNA ligase.

Successful ligation of the insert was confirmed using the pBADGr multiple cloning site

flanking primers (Table 2) and DNA sequencing. The construct was then electroporated

into *ΔfimV* and *ΔpocA* cells. A stable fusion of YFP to the C-terminus of FimV was

confirmed by western blotting with α-AFP and α-FimV antibodies. This fusion is

considered non-functional as it failed to restore twitching in the *fimV* background.

Twitching motility assays

Twitching motility assays were performed as previously described (59). Single

colonies were stab inoculated to the bottom of a 1% LB agar plate. Plates were

incubated for 24 h at 30 °C. Following incubation, the agar was removed and the

adherent bacteria stained with 1% (w/v) crystal violet for 30 min, followed by washing

with water to remove unbound dye. Twitching zones were photographed and their areas

measured using Fiji (60). All experiments were performed in triplicate with at least three

independent replicates.

Analysis of whole cell lysates by western blot

Cultures were grown overnight at 37 °C in LB supplemented with appropriate

antibiotics and diluted to an OD_{600} of 0.6. A 1 mL aliquot of cells was collected by

centrifugation at 2292 x g for 1 min. and the cell pellet was resuspended in 100 µl of

SDS sample buffer (80 mM Tris (pH 6.8), 5.3% (v/v) 2- mercaptoethanol, 10% (v/v)

glycerol, and 0.02% (w/v) bromophenol blue, and 2% (w/v) SDS) and boiled for 10 min.

Equal volumes of whole cell lysates were separated by 12.5% SDS-PAGE at 80-

150 V and transferred to nitrocellulose membranes for 1 h at 225 mA. Membranes were

blocked using 5% (w/v) low fat skim milk powder in phosphate buffered saline solution

(PBS) at pH 7.4 for 1 h at room temperature, followed by incubation with the appropriate

polyclonal (for PilF, PilM, PilO, PilP, PilQ, FimV) or monoclonal (for mCherry) antisera

for 1 h at room temperature. Membranes were washed twice in 10 mL of PBS for 5 min

then incubated in either goat-anti-rabbit or goat-anti-mouse IgG-alkaline phosphatase

conjugated secondary antibody (Bio-Rad) at a dilution of 1:3000 for 1 h at room

temperature. Membranes were washed twice in PBS for 5 min, and developed in a

solution containing 100 μL nitro-blue tetrazolium (NBT) and 100 μL 5-bromo-4-chloro-3-

indolyl phosphate (BCIP) in 10 mL of alkaline phosphatase buffer (100 mM NaCl, 5 mM

MgCl$_2$, 100 mM Tris, pH 9.5).

Cell preparation for transmission electron microscopy

Cells from a 1 mL overnight culture in LB were pelleted at 2292 x g and

resuspended in a 1 mL solution of 2% (w/v) gluteraldehyde in 0.1M sodium cacodylate

buffer at a pH of 7.4. Cells were fixed in solution for 2 h at 4 °C, and washed twice with

cold sodium cacodylate buffer (pH 7.4). Samples were then treated with 1% osmium

tetroxide in 0.1M sodium cacodylate for 1 h and stained with 2% uranyl acetate

overnight. After staining, cells were dehydrated gradually with ethanol, treated with

propylene oxide, embedded in Spurr's resin, and polymerized at 60 °C overnight. The

polymerized samples were sectioned using a Leica UCT ultramicrotome and post-

stained with 2% uranyl acetate and lead citrate. The prepared specimens were

examined in McMaster University's Electron Microscopy Facility using the JEOL JEM

1200 EX TEMSCAN microscope (JEOL, Peabody, MA, USA) operating at an

accelerating voltage of 80kV. Images were acquired with an AMT 4-megapixel digital

camera (Advanced Microscopy Techniques, Woburn, MA).

Fluorescence Microscopy

Strains were incubated overnight at 37 °C in 5 mL of LB supplemented with the

appropriate antibiotics. One mL of cells were pelleted at 2292 x g, and resuspended in 1

mL of sterile water containing 10 µg/ml of either Fm1-43 Fx (for cells expressing

mCherry) or Fm1-64 Fx (for cells expressing YFP) membrane stain (Invitrogen). Cells

were mixed with dye by gentle pipetting, and pelleted at 2292 x *g*. Pelleted cells were

then stab-inoculated using a pipette tip to the coverslip-media interface at the bottom of

an 8-well microscope glass-coverslip slide containing 200 µl 1% LB agarose per well

(Lab-Tek). Cells were then incubated for 75 min at 37 °C prior to imaging. In

filamentation experiments, 1 mL of cells grown overnight were sub-cultured into 4 mL of

LB containing 40 µg/ml of cefsulodin and incubated for 3 h at 37 °C. Cells were then

stained and mounted for microscopy using the above protocol. Cells were imaged using

an EVOS Fl-auto microscope in the McMaster Biophotonics Facility. Images were

acquired using a 60x oil immersion objective using a Texas Red filter, YFP filter, or

transmitted white light (no filter). Fiji (60) was used to adjust image brightness and

contrast, and to overlay brightfield and fluorescent images.

Quantification of mCherry-PilO fluorescence

Fluorescence was quantified using Fiji software (60). Each field in the Texas Red

(for mCherry) and YFP (for Fm1-43 Fx) filter sets were overlaid, and used with the

micrograph from the YFP filter for image analysis. A 250 µm^2 grid was added to each

field for precise documentation of the quantified cells. Moving systematically between

squares in the grid, cells that fulfilled the following criteria were selected: 1) not

contacting other cells; 2) normal rod morphology; and 3) a stable mCherry-PilO fusion

(diffuse cytoplasmic fluorescence implies cleavage of the tag in that cell). Cell lengths

and pixel intensities were measured from one cell pole to the other and normalized to 1

(i.e., each data point associated with length divided by the total length of the cell to yield

normalized 0 to 1 scale). Pixel intensity measurements in the YFP field (cell outline)

were then subtracted from the overlay field, producing mCherry-PilO pixel intensity

measurements alone in each cell on a scale from 0 to 1. From the data, an XY scatter

plot was generated to show the distribution of mCherry-PilO fluorescence over cell

length.

ACKNOWLEDGEMENTS

We thank Dr. Zemer Gitai for the *pocA* deletion construct, and Dr. Ray Truant and

McMaster Biophotonics for access to the EVOS microscope, and the McMaster Electron

Microscopy facility for assistance with TEM.

REFERENCES

1. **Scheurwater EM, Burrows LL.** 2011. Maintaining network security: how

macromolecular structures cross the peptidoglycan layer. FEMS Microbiology

Letters **318:**1-9.

2. **Mattick JS.** 2002. Type IV Pili and Twitching Motility. Annual Review of

Microbiology **56:**289-314.

3. **Burrows LL.** 2012. *Pseudomonas aeruginosa* Twitching Motility: Type IV Pili in

Action. Annual Review of Microbiology **66:**493-520.

4. **Siryaporn A, Kuchma SL, O'Toole GA, Gitai Z.** 2014. Surface attachment

induces *Pseudomonas aeruginosa* virulence. Proc Natl Acad Sci U S A

111:16860-16865.

5. **Heiniger RW, Winther-Larsen HC, Pickles RJ, Koomey M, Wolfgang MC.**

2010. Infection of human mucosal tissue by *Pseudomonas aeruginosa* requires

sequential and mutually dependent virulence factors and a novel pilus-associated

adhesin. Cellular Microbiology **12:**1158-1173.

6. **Golovkine G, Faudry E, Bouillot S, Elsen S, Attree I, Huber P.** 2016.

Pseudomonas aeruginosa Transmigrates at Epithelial Cell-Cell Junctions,

Exploiting Sites of Cell Division and Senescent Cell Extrusion. PLoS Pathog

12:e1005377.

7. **Chang YW, Rettberg LA, Treuner-Lange A, Iwasa J, Sogaard-Andersen L,**

Jensen GJ. 2016. Architecture of the type IVa pilus machine. Science

351:aad2001.

8. **Gold VA, Salzer R, Averhoff B, Kuhlbrandt W.** 2015. Structure of a type IV

pilus machinery in the open and closed state. Elife **4**.

9. **Koo J, Tang T, Harvey H, Tammam S, Sampaleanu L, Burrows LL, Howell**

PL. 2013. Functional Mapping of PilF and PilQ in the *Pseudomonas aeruginosa*

Type IV Pilus System. Biochemistry **52:**2914-2923.

10. **Koo J, Lamers RP, Rubinstein JL, Burrows LL, Howell PL.** 2016. Structure of

the *Pseudomonas aeruginosa* Type IVa Pilus Secretin at 7.4 A. Structure

24:1778-1787.

11. **Berry JL, Phelan MM, Collins RF, Adomavicius T, Tonjum T, Frye SA, Bird L,**

Owens R, Ford RC, Lian LY, Derrick JP. 2012. Structure and assembly of a

trans-periplasmic channel for type IV pili in *Neisseria meningitidis*. PLoS Pathog

8:e1002923.

12.   **Leighton TL, Buensuceso RNC, Howell PL, Burrows LL.** 2015. Biogenesis of

Pseudomonas aeruginosa type IV pili and regulation of their function.

Environmental Microbiology **17:**4148-4163.

13.   **Rocaboy M, Herman R, Sauvage E, Remaut H, Moonens K, Terrak M,**

Charlier P, Kerff F. 2013. The crystal structure of the cell division amidase AmiC

reveals the fold of the AMIN domain, a new peptidoglycan binding domain.

Molecular Microbiology **90:**267-277.

14.   **Ayers M, Sampaleanu LM, Tammam S, Koo J, Harvey H, Howell PL, Burrows**

LL. 2009. PilM/N/O/P Proteins Form an Inner Membrane Complex That Affects

the Stability of the *Pseudomonas aeruginosa* Type IV Pilus Secretin. J Mol Biol

394:128-142.

15.   **Leighton TL, Dayalani N, Sampaleanu LM, Howell PL, Burrows LL.** 2015.

Novel Role for PilNO in Type IV Pilus Retraction Revealed by Alignment

Subcomplex Mutations. J Bacteriol **197:**2229-2238.

16.   **Sampaleanu LM, Bonanno JB, Ayers M, Koo J, Tammam S, Burley SK, Almo**

SC, Burrows LL, Howell PL. 2009. Periplasmic Domains of *Pseudomonas*

aeruginosa PilN and PilO Form a Stable Heterodimeric Complex. J Mol Biol

394:143-159.

17.   **Tammam S, Sampaleanu LM, Koo J, Sundaram P, Ayers M, Chong AP,**

Forman-Kay JD, Burrows LL, Howell PL. 2011. Characterization of the PilN,

PilO and PilP type IVa pilus subcomplex. Molecular Microbiology **82:**1496-1514.

18.   **Karuppiah V, Derrick JP.** 2011. Structure of the PilM-PilN inner membrane type

IV pilus biogenesis complex from *Thermus thermophilus*. J Biol Chem

286:24434-24442.

19.   **McCallum M, Tammam S, Little DJ, Robinson H, Koo J, Shah M, Calmettes**

C, Moraes TF, Burrows LL, Howell PL. 2016. PilN binding modulates the

structure and binding partners of the *Pseudomonas aeruginosa* Type IVa Pilus

protein PilM. J Biol Chem **291:**11003-11015.

20.   **Tammam S, Sampaleanu LM, Koo J, Manoharan K, Daubaras M, Burrows**

LL, Howell PL. 2013. PilMNOPQ from the *Pseudomonas aeruginosa* type IV

pilus system form a transenvelope protein interaction network that interacts with

PilA. J Bacteriol **195:**2126-2135.

21.   **Leighton TL, Yong DH, Howell PL, Burrows LL.** 2016. Type IV Pilus Alignment

Subcomplex Components PilN and PilO Form Homo- and Heterodimers In Vivo.

J Biol Chem doi:10.1074/jbc.M116.738377.

22.   **Gotoh N, Nunomura K, Nishino T.** 1990. Resistance of *Pseudomonas*

aeruginosa to cefsulodin: modification of penicillin-binding protein 3 and mapping

of its chromosomal gene. J Antimicrob Chemo **25:**513-523.

23.   **Davis BM, Waldor MK.** 2013. Establishing polar identity in gram-negative rods.

Curr Opin Microbiol **16:**752-759.

24.   **Hu Z, Mukherjee A, Pichoff S, Lutkenhaus J.** 1999. The MinC component of

the division site selection system in *Escherichia coli* interacts with FtsZ to prevent

polymerization. PNAS **96:**14819-14824.

25.   **Burrows LL.** 2013. A new route for polar navigation. Molecular Microbiology
90:919-922.

26.   **Friedrich C, Bulyha I, Søgaard Andersen L.** 2014. Outside-in assembly
pathway of the type IV pilus system in *Myxococcus xanthus*. J Bacteriol **196:**378-
390.

27.   **Semmler ABT, Whitchurch CB, Leech AJ, Mattick JS.** 2000. Identification of a
novel gene, *fimV*, involved in twitching motility in *Pseudomonas aeruginosa*.
Microbiology **146:**1321-1332.

28.   **Wehbi H, Portillo E, Harvey H, Shimkoff AE, Scheurwater EM, Howell PL,**
Burrows LL. 2011. The peptidoglycan-binding protein FimV promotes assembly
of the *Pseudomonas aeruginosa* type IV pilus secretin. J Bacteriol **193:**540-550.

29.   **Cowles KN, Moser TS, Siryaporn A, Nyakudarika N, Dixon W, Turner JJ,**
Gitai Z. 2013. The putative Poc complex controls two distinct *Pseudomonas*
aeruginosa polar motility mechanisms. Molecular Microbiology **90:**923-938.

30.   **Huang B, Ru K, Yuan Z, Whitchurch CB, Mattick JS.** 2004. tonB3 is required
for normal twitching motility and extracellular assembly of type IV pili. J Bacteriol
186:4387-4389.

31.   **Bernhardt TG, De Boer PAJ.** 2003. The *Escherichia coli* amidase AmiC is a
periplasmic septal ring component exported via the twin-arginine transport
pathway. Molecular Microbiology **48:**1171-1182.

32.   **Heidrich C, Templin MF, Ursinus A, Merdanovic M, Berger J, Schwarz H, de**
Pedro MA, Holtje J-V. 2001. Involvement of N-acetylmuramyl-L-alanine

amidases in cell separation and antibiotic induced autolysis of *Escherichia coli*.

Molecular Microbiology **41:**167-178.

33. **Peters NT, Dinh T, Bernhardt TG.** 2011. A fail-safe mechanism in the septal ring

assembly pathway generated by the sequential recruitment of cell separation

amidases and their activators. J Bacteriol **193:**4973-4983.

34. **Buist G, Steen A, Kok J, Kuipers OP.** 2008. LysM, a widely distributed protein

motif for binding to (peptido)glycans. Molecular Microbiology **68:**838-847.

35. **Duncan TR, Yahashiri A, Arends SJR, Popham DL, Weiss DS.** 2013.

Identification of SPOR domain amino acids important for septal localization,

peptidoglycan binding, and a disulfide bond in the cell division protein FtsN. J

Bacteriology **195:**5308-5315.

36. **Mesnage Sep, Dellarole M, Baxter NJ, Rouget J-B, Dimitrov JD, Wang N,**

Fujimoto Y, Hounslow AM, Lacroix-Desmazes Seb, Fukase K, Foster SJ,

Williamson MP. 2014. Molecular basis for bacterial peptidoglycan recognition by

LysM domains. Nat Communications **5:**1-11.

37. **de Souza RF, Anantharaman V, de Souza SJ, Aravind L, Gueiros-Filho FJ.**

2008. AMIN domains have a predicted role in localization of diverse periplasmic

protein complexes. Bioinformatics **24:**2423-2426.

38. **Ramirez-Arcos S, Szeto J, Beveridge T, Victor C, Francis F, Dillon J.** 2001.

Deletion of the cell-division inhibitor MinC results in lysis of *Neisseria*

gonorrhoeae. Microbiology **147:**225-237.

39. **Buensuceso RNC, Nguyen Y, Zhang K, Daniel-Ivad M, Sugiman-Marangos**

SN, Fleetwood AD, Zhulin IB, Junop MS, Howell PL, Burrows LL. 2016. The

conserved TPR-containing C-terminal domain of *Pseudomonas aeruginosa* FimV

is required for its cAMP-dependent and independent functions. J Bacteriol

198:2263-2274.

40.   **Kirkpatrick CL, Viollier PH.** 2012. Cell polarity: ParA-logs gather around the

Hub. Curr Biol **22:**R1055-1057.

41.   **Rossmann F, Brenzinger S, Knauer C, Dorrich AK, Bubendorfer S, Ruppert**

U, Bange G, Thormann KM. 2015. The role of FlhF and HubP as polar landmark

proteins in *Shewanella putrefaciens* CN-32. Mol Microbiol **98:**727-742.

42.   **Takekawa N, Kwon S, Nishioka N, Kojima S, Homma M.** 2016. HubP, a polar

landmark protein, regulates flagellar number by assisting in the proper polar

localization of FlhG in *Vibrio alginolyticus*. J Bacteriol doi:10.1128/JB.00462-16.

43.   **Yamaichi Y, Bruckner R, Ringgaard S, Moll A, Cameron DE, Briegel A,**

Jensen GJ, Davis BM, Waldor MK. 2012. A multidomain hub anchors the

chromosome segregation and chemotactic machinery to the bacterial pole.

Genes Dev **26:**2348-2360.

44.   **Inclan YF, Persat A, Greninger A, Von Dollen J, Johnson J, Krogan N, Gitai**

Z, Engel JN. 2016. A scaffold protein connects type IV pili with the Chp

chemosensory system to mediate activation of virulence signaling in

Pseudomonas aeruginosa. Mol Microbiol **101:**590-605.

45.   **Korotkov KV, Pardon E, Steyaert J, Hol WG.** 2009. Crystal structure of the N-

terminal domain of the secretin GspD from ETEC determined with the assistance

of a nanobody. Structure **17:**255-265.

46.   **Nudleman E, Wall D, Kaiser D.** 2005. Cell-to-cell Transfer of Bacterial Outer

Membrane Lipoproteins. Science **309:**125-127.

47.   **Typas A, Banzhaf M, van den Berg van Saparoea B, Verheul J, Biboy J,**

Nichols RJ, Zietek M, Beilharz K, Kannenberg K, von Rechenberg M,

Breukink E, den Blaauwen T, Gross CA, Vollmer W. 2010. Regulation of

peptidoglycan synthesis by outer-membrane proteins. Cell **143:**1097-1109.

48.   **Jean NL, Bougault CM, Lodge A, Derouaux A, Callens G, Egan AJ, Ayala I,**

Lewis RJ, Vollmer W, Simorre JP. 2014. Elongated structure of the outer-

membrane activator of peptidoglycan synthesis LpoA: implications for PBP1A

stimulation. Structure **22:**1047-1054.

49.   **Nandi S, Swanson S, Tomberg J, Nicholas RA.** 2015. Diffusion of antibiotics

through the PilQ secretin in *Neisseria gonorrhoeae* occurs through the immature,

sodium dodecyl sulfate-labile form. J Bacteriol **197:**1308-1321.

50.   **Demchick P, Koch AL.** 1996. The Permeability of the Wall Fabric of *Escherchia*

coli and *Bacillus subtilis*. J Bacteriol **178:**768-773.

51.   **Wang P, Robert L, Pelletier J, Dang WL, Taddei F, Wright A, Jun S.** 2010.

Robust growth of *Escherichia coli*. Curr Biol **20:**1099-1103.

52.   **Korotkov KV, Johnson TL, Jobling MG, Pruneda J, Pardon E, Heroux A,**

Turley S, Steyaert J, Holmes RK, Sandkvist M, Hol WG. 2011. Structural and

functional studies on the interaction of GspC and GspD in the type II secretion

system. PLoS Pathog **7:**e1002228.

53.   **Wang X, Pineau C, Gu S, Guschinskaya N, Pickersgill RW, Shevchik VE.**

2012. Cysteine scanning mutagenesis and disulfide mapping analysis of

arrangement of GspC and GspD protomers within the type 2 secretion system. J

Biol Chem **287:**19082-19093.

54. **Lamers RP, Nguyen UT, Nguyen Y, Buensuceso RN, Burrows LL.** 2015. Loss

of membrane-bound lytic transglycosylases increases outer membrane

permeability and beta-lactam sensitivity in *Pseudomonas aeruginosa*.

Microbiologyopen **4:**879-895.

55. **Hoang TT, Karkhoff-Schweizer RR, Kutchma AJ, Schweizer HP.** 1998. A

broad-host-range Flp-FRT recombination system for site-specific excision of

chromosomally-located DNA sequences: application for isolation of unmarked

Pseudomonas aeruginosa mutants. Gene **212:**77-86.

56. **Shu X, Shaner NC, Yarbrough CA, Tsien RY, Remington SJ.** 2006. Novel

Chromophores and Buried Charges Control Color in mFruits. Biochemistry

45:9639-9647.

57. **Lieberman JA, Petro CD, Thomas S, Yang A, Donnenberg MS.** 2015. Type IV

pilus secretins have extracellular C termini. MBio **6:**e00322-15.

58. **Asikyan ML, Kus JV, Burrows LL.** 2008. Novel proteins that modulate type IV

pilus retraction dynamics in *Pseudomonas aeruginosa*. Journal of Bacteriology

190:7022-7034.

59. **Kus JV, Tullis E, Cvitkovitch DG, Burrows LL.** 2004. Significant differences in

type IV pilin allele distribution among *Pseudomonas aeruginosa* isolates from

cystic fibrosis (CF) versus non-CF patients. Microbiology **150:**1315-1326.

60. **Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T,**

Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ,

Hartenstein V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source

platform for biological-image analysis. Nature Methods **9:**676-682.

61.   **Simon R, Priefer U, Puhler A.** 1983. A Broad Host Range Mobilization System

for in vivo Genetic Engineering: Transposon Mutagenesis in Gram Negative

Bacteria. Nature Biotechnology **1:**784-791.

62.   **Takhar HK, Kemp K, Kim M, Howell PL, Burrows LL.** 2013. The platform

protein is essential for type IV pilus biogenesis. J Biol Chem **288:**9721-9728.

63.   **Kus JV, Kelly J, Tessier L, Harvey H, Cvitkovitch DG, Burrows LL.** 2008.

Modification of *Pseudomonas aeruginosa* Pa5196 Type IV Pilins at Multiple Sites

with D-Araf by a Novel GT-C Family Arabinosyltransferase, TfpW. Journal of

Bacteriology **190:**7464-7478.

64.   **Nguyen Y, Harvey H, Sugiman-Marangos S, Bell SD, Buensuceso RNC,**

Junop MS, Burrows LL. 2015. Structural and Functional Studies of the

Pseudomonas aeruginosa Minor Pilin, PilE. Journal of Biological Chemistry

290:26856-26865.

 TABLES

 Table 1. Bacterial strains and plasmids

Strain	Description	Source
E. coli		
DH5α	F-, φ80lacZ, M15, Δ(*lacZYA-argF*), U169, *recA1, endA1, hsdR17(rk-,mk+), phoAsupE44, thi-1, gyrA96, relA1*, λ-	Invitrogen
SM10	*thi-1, thr, leu, tonA, lacy, supE, recA*, RP4-2-Tcr::Mu, Kmr; mobilizes plasmids into *P. aeruginosa* via conjugation	(61)
P. aeruginosa		
PAK	Wild type	J. Boyd
ΔpilM	Deletion of *pilM*	(62)
pilP::FRT	FRT scar at nucleotide 86 of *pilP*	(14)

pilQ::FRT	FRT scar at nucleotide 571 of *pilQ*	(9)
pilA::FRT	FRT scar at SphI site within *pilA*	(63)
ΔpilF	*pilF* deletion strain	This study
ΔminC	*minC* deletion strain	This study
mCherry-PilO	In frame fusion of *mCherry* to the 5' end of *pilO* on the PAK chromosome	This study
ΔpocA	*pocA* deletion strain	(29)
ΔfimV	*fimV* deletion strain	(39)
fimVΔLysM	*fimV* with in frame deletion of LysM domain	(28)
mCherry-PilO *ΔpilM*	deletion of *pilM* in mCherry-PilO strain	This study
mCherry-PilO *pilP*::FRT	FRT scar in *pilP* at nucleotide 86 in mCherry-PilO strain	This study
mCherry-PilO *pilQ*::FRT	FRT scar at position 571 within *pilQ* in mCherry-PilO strain	This study
mCherry-PilO *ΔpilF*	deletion of *pilF* in	This study

	mCherry-PilO strain	
mCherry-PilO *ΔminC*	deletion of *minC* in mCherry-PilO strain	This study
mCherry-PilO *ΔpocA*	deletion of *pocA* in mCherry-PilO background	This study
mCherry-PilO *ΔfimV*	deletion of *fimV* in mCherry-PilO background	This study
mCherry-PilO *fimVΔLysM*	Insertion of *mCherry* at the 5' end of *pilO* with *fimV* containing deletion of LysM domain	This study
Plasmids	**Description**	**Source**
pEX18Gm	Suicide vector used for gene replacement, Gm	(55)
pEX18Ap	Suicide vector used for gene replacement, Ap	(55)
pBADGr	Arabinose inducible protein expression vector	(58)
pFLP2	Suicide vector containing	(55)

	Flp recombinase	
pEX18Gm::*mCherry-pilO*	Suicide vector containing *mCherry-pilO*, subcloned from pUC57 (Genscript) using EcoRI/HindIII	This study
pEX18Gm::*ΔpilF*	Suicide vector containing *pilF* deletion, cloned using BamHI/HindIII	This study
pEX18Gm::*ΔminC*	Suicide vector containing *minC* deletion, cloned using BamHI/HindIII	This study
pEX18Gm::*ΔpocA*	Suicide vector containing *pocA* deletion, cloned using EcoRI/XbaI	This study
pEX18Ap-*pilQ*::GmFRT	Suicide vector containing PAK *pilQ* disrupted with FRT-flanked Gm cassette at position 571	This study
pBADGr::*pilQ-mCherry*	*pilQ-mCherry* complementation construct, cloned using EcoRI/HindIII	This study
pBADGr::*pilF*	*pilF* complementation	This study

	construct, cloned using SphI/HindIII	
pBADGr::*pilQ*	*pilQ* complementation construct, cloned using EcoRI/HindIII	This study
pBADGr::*pilQΔAMIN*	*pilQΔAMIN* complementation construct, cloned using EcoRI/HindIII	This study
pBADGr::*pocA*	*pocA* complementation construct, cloned using EcoRI/HindIII	This study
pBADGr::*fimV-yfp*	*fimV-yfp* complementation construct, cloned using KpnI/XbaI	This study

759 **Table 2. Primer sequences**

Name	Oligonucleotide sequence
PilF-Fwd	5'ACGCCTTGCAAGATCAACCTGATTCCG3'
PilF-Rev	5'TCGAACGCGCCGTTTTCCAGCTGACGC3'
PilF mid-Rev	5'GGCCGGTTTCTTCATTTGCAGCG3'
PilF Clone-Fwd	5'TCATGCATGCATGACTGTACGCGCCGCGCTGG3'
PilF Clone-Rev	5'TCATAAGCTTTCATTTTTCCGCCTGGAATTCCTG3'
MinC Fwd	5'GGATACGGCAACTGCACCACCAGCCAG3'
MinC Rev	5'GACGACGTTGACGAAGTCGTACACCAC3'
MinC mid-Rev	5'CGTGGCGGCGACAGACCTCGAGGA3'
PilN 3'-Fwd	5'GCCAACGTGTTCCAACTG3'
PilO Rev	5'CCACGCTGATCTGGATCG3'
PocA Fwd	5'GAATTCGGGGATATGCCACGTGTGGGA3'
PocA Rev	5'AAGCTTTGCGGCGGAATTTCACGCTTTGC3'
PocA Mid-Rev	5'GAGGATCTCTCCCAGCGG3'
PocA Clone-Fwd	5'TCATGAATTCGTGTGGGAACTGGTTCAAGCCGG3'

PocA Clone-Rev	5'TCATAAGCTTTCACGCTTTGCCTTCCTCGACGTAG3'
PilQ Clone-Fwd	5'AGAATTCCAACAGCAGTCTGTACAA3'
PilQ Clone-Rev	5'TCATAAGCTTTCAGCGACCGATTGCGATGGCCTG3'
PilQ 910-Fwd	5'CGACCTGAATCTGGTGGC3'
PilQ FRT-Chk Fwd	5'TTGATCATCAACCTGACCGCGCTGTCG3'
PilQ FRT-Chk Rev	5'TCATCGGCTTGATGCTGACGGTCAGG3'
pBAD mcs-Fwd	5'AAGTGTCTATAATCACGGCAGA3'
pBAD mcs-Rev	5'TCACTTCTGAGTTCGGCATGG3'
mCherry Rev	5'TCATGCATGCGCTTCCGCCGCTCTTGTACAGCTCGTCCATGCCGC3'
PilQ 5' Rev	5'TCATTCTAGAGTCCGCGGCGAGCAATGCCGGCGC3'
PilQ N0 Fwd	5'TCATTCTAGAGGCGAGAAACTGTCGCTGAACTTC3'
FimV Clone Fwd	5'GCGGGTACCATGGTTCGGCTTCG3'
FimV Clone Rev	5'GCGTCTAGAGGCCAGGCGCTCCA3'

FIGURE LEGENDS

Figure 1. mCherry-PilO localizes to cell poles and future sites of cell division in *P. aeruginosa*. A. Map of the T4aP alignment subcomplex operon showing *mCherry-pilO* integration. **B**. The mCherry-PilO fusion was stable (~48 kDa product recognized by both anti-PilO and anti-mCherry antibodies; a plasmid-encoded PilE-mCherry fusion – open triangle – was used as a positive control for the latter; (64)) and functional for twitching motility. **C**. mCherry-PilO localized to the poles of wild type cells and to the septum in late-stage dividing cells. **D**. When cells were filamented using 40 µg/ml cefsulodin, mCherry-PilO localized to the poles and at regularly spaced foci (arrowheads). Scale bar = 3 µm. **E**. mCherry pixel intensity in 50 untreated cells after normalization of length to 1 as described in the Methods.

Figure 2. mCherry-PilO localizes to aberrantly localized septa in a *minC* mutant. A. Transmission electron micrograph of a *minC* mutant of *P. aeruginosa* strain PAK. In the absence of *minC*, cells have aberrantly located division sites (red arrows). Scale bar = 1 µm. **B.** mCherry-PilO polar foci are indicated with white arrowheads while foci localized to aberrantly placed septa are indicated with yellow arrows.

Figure 3. Polar localization of T4aP alignment subcomplex requires PilQ. In the absence of PilM, mCherry-PilO is localized to **A.** poles and septa in untreated cells, and **B.** poles and regularly spaced foci in cells treated with cefsulodin. In the absence of PilQ, mCherry-PilO is delocalized in both **C.** untreated and **D.** antibiotic-filamented cells. Similarly, in the absence of PilP, mCherry-PilO is delocalized in **E.** untreated and **F.**

antibiotic-treated cells. The last 3 panels show quantification of mCherry pixel intensity

in the absence of **G.** *pilM*, **H.** *pilQ*, or **I.** *pilP*, in 50 untreated cells after normalization of

length to 1 as described in the Methods. Scale bar = 3 µm.

Figure 4. The AMIN domains of PilQ are required for polar localization of mCherry-

PilO. A. Domain map of PilQ showing its PG-binding AMIN domains in blue.

Complementation of a *pilQ* rmutant expressing mCherry-PilO with full length *pilQ* re-

established its polar and septal localization (white arrowheads) in **B.** untreated cells, and

C. localization to the poles and regularly spaced foci in cefsulodin treated cells. When

the same mutant was complemented with a truncated version of *pilQ* that encodes only

N0-N1 and the secretin domains, mCherry-PilO was delocalized in both **D.** untreated

and **E.** antibiotic-treated cells. The last 3 panels show quantification of mCherry pixel

intensity in the presence of **F.** PilQ or **G.** PilQ lacking its AMIN domains, in 50 untreated

cells after normalization of length to 1 as described in the Methods. Scale bar = 3 µm.

Figure 5. mCherry-PilO is polarly localized in the absence of *pilF*. A. mCherry-PilO

localized to the poles and septa of late-stage dividing cells, marked by white

arrowheads. **B.** When cells were filamented with cefsulodin, mCherry-PilO localized to

the poles and to foci (arrows). Scale bar = 3 µm. Panel **C.** shows quantification of

mCherry pixel intensity in the *pilF* background, in 50 untreated cells after normalization

of length to 1 as described in the Methods.

Figure 6. FimV is required for polar localization of mCherry-PilO. In the absence of

FimV, mCherry-PilO was delocalized in **A.** untreated cells and **B.** cells treated with

cefsulodin. In cells expressing a version of FimV with an in-frame deletion of its PG-

binding domain (LysM), mCherry-PilO was delocalized in **C.** untreated and **D.** antibiotic-

treated cells. The last two panels show quantification of mCherry pixel intensity **F.** in the

absence of FimV or **G.** in the strain expressing FimV$_{\Delta LysM}$, in 50 untreated cells after

normalization of length to 1 as described in the Methods. Scale Bar = 3 µm.

Figure 7. mCherry-PilO is delocalized in a *pocA* mutant. In the absence of the polar

organelle coordinating protein, PocA, mCherry-PilO was delocalized in **A.** untreated

cells and **B.** cells treated with cefsulodin. When *pocA* was reintroduced *in trans*,

mCherry-PilO localization to the poles and septum was recovered (white arrowheads) in

C. untreated cells, as well as to **D.** the poles and regularly spaced foci in antibiotic-

treated cells. The last two panels show quantification of mCherry pixel intensity **E.** in the

absence of PocA and **F.** in the *pocA*-complemented mutant, in 50 untreated cells after

normalization of length to 1 as described in the Methods. Scale bar = 3 µm.

Figure 8. FimV remains polarly localized in the absence of PocA. A. FimV-YFP

remained localized to the cell poles (arrowheads) in a *pocA* mutant. **B.** FimV-YFP

localized to the poles in cefsulodin treated cells. Scale bar = 3 µm.

Carter et al. Figure 1

A.

B.

Brightfield   Membrane   mCherry-PilO   Overlay

C.

D.

E.

Carter et al. Figure 2

Carter et al. Figure 5

Carter et al. Figure 6

	Brightfield	Membrane	mCherry-PilO	Overlay

Carter et al. Figure 4

Carter et al. Figure 7

Carter et al. Figure 8

| Brightfield | Membrane | FimV-YFP | Overlay |