RUNNING TITLE: Deterministic assembly of hyporheic microbiomes

Deterministic influences exceed dispersal effects on hydrologically-connected microbiomes

Emily B. Graham,[1*] Alex R. Crump,[1] Charles T. Resch,[2] Sarah Fansler,[1] Evan Arntzen,[3] David

W. Kennedy,[1] Jim K. Fredrickson,[1] James C. Stegen[1]

[1]Biological Sciences Division, Pacific Northwest National Laboratory, Richland, WA USA

[2]Geochemistry Department, Pacific Northwest National Laboratory, Richland, WA USA

[3]Environmental Compliance and Emergency Preparation, Pacific Northwest National

Laboratory, Richland, WA USA

[*]**Corresponding author:** Emily B. Graham, Pacific Northwest National Laboratory, PO Box

999, Richland, WA 99352, 509-372-6049, emily.graham@pnnl.gov

Data accessibility statement: Upon acceptance for publication, all data will be made publically

available, and the DOI will be provided in-text.

Conflict of Interest: The authors declare no conflict of interest.

Originality-Significance Statement. Subsurface zones of groundwater and surface water

mixing (hyporheic zones) are hotspots of biogeochemical activity and strongly influence carbon,

nutrient and contaminant dynamics within riverine ecosystems. Hyporheic zone microbiomes are

responsible for up to 95% of riverine ecosystem respiration, yet the ecology of these

microbiomes remains poorly understood. While significant progress is being made in the

development of microbially-explicit ecosystem models, poor understanding of hyporheic zone

microbial ecology impedes development of such models in this critical zone. To fill the

knowledge gap, we present a comprehensive analysis of biogeographical patterns in hyporheic

microbiomes as well as the ecological processes that govern their composition and function

through space and time. Despite pronounced hydrologic connectivity throughout the hyporheic

zone—and thus a strong potential for dispersal—we find that ecological selection

deterministically governs microbiome composition within local environments, and we identify

specific groups of organisms that correspond to seasonal changes in hydrology. Based on our

results, we propose a conceptual model for hyporheic zone metabolism in which comparatively

high-organic C conditions during surface water intrusion into the hyporheic zone support

heterotrophic metabolisms that succumb to autotrophy during time periods of groundwater

discharge. These results provide new opportunities to develop microbially-explicit ecosystem

models that incorporate the hyporheic zone and its influence over riverine ecosystem function.

Keywords: hyporheic zone, selection, subsurface, microbial community structure, community

assembly, freshwater biology, microbial ecology, stochastic assembly

Summary.

Subsurface groundwater-surface water mixing zones (hyporheic zones) have enhanced

biogeochemical activity, but assembly processes governing subsurface microbiomes remain a

critical uncertainty in understanding hyporheic biogeochemistry. To address this obstacle, we

investigated (a) biogeographical patterns in attached and waterborne microbiomes across three

hydrologically-connected, physicochemically-distinct zones (inland hyporheic, nearshore

hyporheic, and river); (b) assembly processes that generated these patterns; (c) groups of

organisms that corresponded to deterministic changes in the environment; and (d) correlations

between these groups and hyporheic metabolism. All microbiomes remained dissimilar through

time, but consistent presence of similar taxa suggested dispersal and/or common selective

pressures among zones. Further, we demonstrated a pronounced impact of deterministic

assembly in all microbiomes as well as seasonal shifts from heterotrophic to autotrophic

microorganisms associated with increases in groundwater discharge. The abundance of one

statistical cluster of organisms increased with active biomass and respiration, revealing

organisms that may strongly influence hyporheic biogeochemistry. Based on our results, we

propose a conceptualization of hyporheic zone metabolism in which increased organic carbon

concentrations during surface water intrusion support heterotrophy, which succumbs to

autotrophy under groundwater discharge. These results provide new opportunities to enhance

microbially-explicit ecosystem models describing hyporheic zone biogeochemistry and its

influence over riverine ecosystem function.

Introduction.

Groundwater-surface water mixing zones (i.e., hyporheic zones) are characterized by

temporally and spatially dynamic, redox-active environments with elevated rates of

biogeochemical cycling (Hedin et al., 1998; McClain et al., 2003; Hancock et al., 2005; Boulton

et al., 2010). Yet, hyporheic microbiomes have rarely been examined (Robertson and Wood,

2010; Marmonier et al., 2012; Gonzalez and Bell, 2013), constituting a key uncertainty in

understanding the biogeochemistry of these dynamic zones (Robertson and Wood, 2010;

Marmonier et al., 2012; Gonzalez and Bell, 2013). In particular, microbial community assembly

processes drive microbiome composition and are thought to be imperative in coupling

microbiomes with ecosystem function (Ferrenberg et al., 2013; Nemergut et al., 2013; Graham et

al., 2016a; Graham et al., 2016b). Deterministic assembly (*e.g.*, selection) may either sort species

such that the microbiome is optimized for a given environment or exclude taxa that may

otherwise enhance ecosystem function (Knelman and Nemergut, 2014; Graham et al., 2016a),

while stochasticity (*e.g.*, dispersal, drift) can regulate microbiome composition and function via

mechanisms such as such as priority effects (Fukami, 2004; Fukami et al., 2010) and dispersal

limitation (Lindström and Langenheder, 2012; Adams et al., 2013; Cline and Zak, 2014).

Knowledge on the balance of assembly processes in determining microbiome composition and

function in response to hydrologic exchange in the hyporheic zone is critical to understanding

river corridor function, as hyporheic zones account for up to 95% of riverine ecosystem

respiration (Naegeli and Uehlinger, 1997).

Determinism has been widely shown to impact environmental microbiomes (Fierer and

Jackson, 2006; Lauber et al., 2009), while stochastic processes can strongly influence microbial

community composition when environmental heterogeneity and/or species sorting is low (Stegen

et al., 2012; Wang et al., 2013; Woodcock et al., 2013; Battin et al., 2016). Both of these

processes operate in concert to structure microbiomes, and recent work has focused on

deciphering the balance of deterministic vs. stochastic assembly processes through space and

time (Dini-Andreote et al., 2015; Gianuca et al., 2016; Graham et al., 2016a). Further, assembly

processes function at an organismal level; determinism selects for traits contained within certain

groups of organisms and dispersal disproportionately affects organisms expressing traits for

motility (Martiny et al., 2006; Martiny et al., 2013). Understanding microbiome assembly at both

the community and sub-community levels is therefore essential to fully comprehending

microbiome assembly and subsequent ecosystem functioning. Nonetheless, influences of

assembly processes on specific groups of microorganisms are poorly understood, and linking a

process-based understanding of variation in microbial physiologies to the metabolism of carbon

and nutrients remains a vital unknown in ecosystem science.

The role of community assembly processes in determining microbiome composition and

function may be particularly important within hyporheic zones that experience frequent

groundwater-surface water mixing (hereafter, hydrologic mixing) and harbor a variety of

complementary resources (Boulton et al., 2010; Stegen et al., 2016). For instance, dispersal may

be paramount to structuring hyporheic microbiomes, as hydrologic mixing induces physical

transport of microbial cells across physicochemically-distinct environments. A heightened

importance of stochasticity has been hypothesized to generate a disconnect between microbiome

composition and biogeochemical function (Nemergut et al., 2013; Graham et al., 2016b), and

such a scenario would suggest microbiome composition is critical to accurately predicting

biogeochemical function. Although most biogeochemical activity occurs within hyporheic

sediments (Boulton et al., 1998; Hancock et al., 2005; Boulton et al., 2010), waterborne

hyporheic microbiomes serve as dispersal vectors among attached microbiomes. Both types of

microbiomes therefore convey mechanistic information regarding ecological dynamics that

influence hyporheic function. Finally, strong local selective filters may negate dispersal effects

by limiting successful immigration into physicochemically-distinct environments. In such a case,

gradients in physicochemistry both determines microbiomes and drives hyporheic

biogeochemical cycling (Graham et al., 2014; Graham et al., 2016b).

Our research has four primary objectives: to determine (a) biogeographical patterns in

hyporheic microbiomes across three physicochemically-distinct zones and to infer associated

trends in microbial physiologies; (b) assembly processes that generate these patterns through

space and time; (c) specific groups of organisms whose relative abundances are deterministically

driven by changes in the environment; and (d) how each identified group corresponds to

hyporheic biogeochemical function. We hypothesize that (a) physicochemical differences across

zones and microbiomes types (i.e., attached and waterborne) generate differences in microbiome

composition and physiology; (b) hydrologic connectivity facilitates microbial dispersal across

hyporheic zones despite steep physicochemical gradients; (c) the relative abundance of

heterotrophic vs. autotrophic organisms corresponds to changes in organic carbon concentration

associated with hydrologic mixing conditions; and (d) increases in the relative abundance of

heterotrophic taxa corresponds with enhanced hyporheic zone biogeochemical activity.

Results.

Hydrology and Physicochemical Conditions.

The Hanford Reach of the Columbia River exhibits frequent hydrologic mixing and steep

physicochemical gradients, presenting a prime opportunity for investigating the impacts of

assembly processes on environmental microbiomes. We sampled distinct but hydrologically-

connected zones that lie in close proximity (<250m) – the inland groundwater aquifer, the

nearshore subsurface environment, and the Columbia River surface water – across seasonally

variable hydrologic mixing. We observed variation in temporal dynamics of hydrologic mixing

and physicochemistry among zones (Fig. 1A-C, Table S1, Fig. S1). Increases in chloride (Cl⁻)

concentration in nearshore and inland zones at and beyond our July 22 sampling event indicated

discharge of groundwater through the hyporheic zone (Graham et al. 2016a, Stegen et al. 2016,

Fig. 1A), coincident with decreases in non-purgable organic carbon (NPOC) concentration in all

three zones (Fig. 1B). Temperature followed a seasonal trend in nearshore and river zones but

remained stable in the inland zone (Fig. 1C).

Microbiome composition.

River (PERMANOVA, $R^2 = 0.22$, P = 0.001), nearshore attached (PERMANOVA, $R^2 =$

0.44, P = 0.001), nearshore waterborne (PERMANOVA, $R^2 = 0.44$, P = 0.001), and inland

waterborne (PERMANOVA, $R^2 = 0.41$, P = 0.001) microbiomes changed seasonally. Inland

attached microbiomes (PERMANOVA, P = 0.29) were stable through time. Comparing all 5

data subsets to each other revealed distinct microbiomes (PERMANOVA $R^2 = 0.25$, P = 0.001,

Fig. S3), and within each zone the composition of waterborne and attached communities differed

(PERMANOVA, nearshore $R^2 = 0.19$, P = 0.001, inland $R^2 = 0.14$, P = 0.001).

The ten most abundant phyla and classes in each data subset are presented in Table 1.

Members of *Proteobacteria* were ubiquitous, and *Bacteroidetes* was highly abundant in all river

and nearshore subsets but had lower abundance in inland zone habitats. The *Planctomycetes-*

Verrucomicrobia-Chlamydiae (PVC) superphylum was prevalent, though the composition of

organisms within this phylum varied among data subsets. *Chloracidobacteria* was widespread in

attached microbiomes in both nearshore and inland zones but was not highly abundant in

waterborne microbiomes. In the inland zone, classes *PBS-25* and *koll11* of the *OP3* candidate

phylum were among the ten most abundant classes in the waterborne microbiome. Further, at the

class level, the ammonia-oxidizing *Thaumarcheaota* was highly abundant in both inland

waterborne and attached microbiomes, joined by a nitrite-oxidizer (*Nitrospira*) in attached

microbiomes.

Microbiome assembly.

We used ecological modeling to infer the relative influences of deterministic selection vs.

stochastic processes on microbiome composition—and futher parsed out the influences of high

and low levels of dispersal within the stochastic component—in attached and waterborne

communities. Deterministic assembly processes had more influence on microbiome composition

than stochastic processes (Table 2). Below, we refer to homogenous selection (i.e., consistent

selection pressures between locations or points in time), variable selection (i.e., divergent

selection pressures between locations or points in time), dispersal limitation (i.e., inhibition of

transport between microbiomes coupled with stochastic demographics), and homogenizing

dispersal (i.e., heightened transport between microbiomes, leading to establishment) as per

definitions outlined by Stegen et al. (2015). In attached microbiomes, both nearshore and inland,

homogenous selection accounted for 82-100% of assembly processes, and for river microbiomes

homogenous selection account for >95% of assembly. While waterborne microbiomes in

nearshore and inland zones exhibited higher stochasticity than their attached counterparts,

selection still accounted for 59.8% and 35.7% of assembly, respectively (Table 2).

Because we hypothesized dispersal mechanisms to have the greatest impact on

waterborne microbiomes compared across zones, we investigated assembly processes

influencing across-zone comparisons within waterborne microbiomes only. Selection comprised

57-92% of across-zone assembly processes, with dispersal being of secondary importance.

Variable selection was responsible for >90% of dissimilarity in microbiomes between nearshore

and inland zones, while nearshore-to-river comparisons showed an even balance of variable

(34.84%) and homogenous (29.18%) selection. The influence of spatial processes was most

evident as dispersal limitation (26.76%) in inland-to-river comparisons and as homogenizing

dispersal (15.30%) in nearshore-to-river comparisons.

Spatiotemporal co-occurrence networks, environmental correlations, and keystone taxa.

Finally, we used co-occurrence networks to infer relationships between deterministic

assembly and specific groups (i.e., statistical clusters) of organisms, as well as to identify taxa

vital in determining the composition of each cluster of organisms (i.e., keystone taxa). We also

investigated relationships between each cluster of organisms and changes in hyporheic

biogeochemical functioning by correlating the abundance of each cluster with active biomass

(ATP) and aerobic respiration (resasurin assay, referred to below as 'Raz').

Properties of full networks at the family level are listed in Table S2, and the composition

of clusters are listed in Table S3. In the nearshore zone, attached cluster 1 was negatively

correlated with time, Cl-, and temperature (Fig. 2A-D); waterborne cluster 9 was positively

correlated with time, Cl-, and temperature and negatively correlated with NPOC (Fig. 2E-H). In

the inland zone, clusters of attached organisms did not display consistent trends with time or

physicochemistry. Cluster 3 (Fig. 2I-L) and cluster 5 (Fig. 2M-P) of inland waterborne

microbiomes exhibited contrasting relationships with time (positive/negative), Cl⁻

(positive/negative), NPOC (negative/positive), and temperature (positive/negative). River

waterborne cluster 2 was negatively correlated with time and positively correlated with Cl⁻, with

no evident relationships to NPOC or temperature (Fig. 2Q-T). Only one cluster –nearshore

attached cluster 1– exhibited correlations with ATP and Raz (Fig. 3).

Nearshore attached cluster 1 contained keystone taxa belonging to *Verrucomicrobia* and

Thaumarcheaota, as well as families of *Gammaproteobacteria*, *Alphaproteobacteria*, and

Chloracidobacteria with a secondary importance (Fig. 4A, Table S3). Nearshore waterborne

cluster 9 contained two keystone families—one family of unassigned organisms and one

belonging to the candidate phylum *OP3* (Fig. 4B). No keystone taxa were identified in inland

waterborne cluster 5 (Fig. 4C), but inland waterborne cluster 3 contained two keystone families

belonging to *Chloracidobacteria* and *Chloroflexi* as well as organisms with secondary

importance (Fig. 4D). No keystone taxa were identified in river cluster 2 (Fig. 4E).

Discussion.

Hydrologic mixing conditions changed through time and corresponded to shifts in

aqueous physicochemistry and microbiome composition across all zones (Fig. 1, Fig. S1). In the

following sub-sections, we parse (a) biogeographical patterns in microbiomes, (b) microbiome

assembly processes, (c) network clusters linked with seasonal changes in physicochemistry, and

(d) associations between network clusters and hyporheic metabolism.

Microbiome biogeography.

Despite hydrologic connectivity across zones, distinct microbiomes were maintained

within each zone and within attached vs. waterborne environments (Febria et al., 2012;

Niederdorfer et al., 2016), supporting our hypothesis that physicochemical differences would

lead to distinct microbiomes.

Among differences in microbiomes, we observed higher abundances of

Chloracidobacteria in attached vs. waterborne microbiomes within both nearshore and inland

zones (Table 1). This suggests a potentially important role for *Chloracidobacteria* in microbial

biofilms attached to sediment particles. Biofilm communities, which are common in hyporheic

zones, consist of a mixture of eukaryotic and prokaryotic organisms embedded in a porous

extracellular matrix and contain diverse organisms with complementary metabolisms (Battin et

al., 2016). At our sampling scale, we capture the diversity of organisms in biofilms but were

unable to resolve spatial organization that may be key to elucidating specific relationships among

these organisms. Still, *Chloracidobacteria* were prevalent in attached microbiomes (*i.e.*, within

biofilms), possibly indicating a role in biofilm stability and functioning. These organisms are

thought to be photoheterotrophs (Bryant et al., 2007; Tank and Bryant, 2015) that can utilize

non-visible near-infrared light as an energy source for growth (Behrendt et al., 2012). While

these properties may confer advantages in subsurface sediment conditions with low levels of

light, our environments experience no light, and the prevalence of *Chloracidobacteria* suggests

possible alternative physiologies for these organisms or dormant lifestyles to allow persistence in

non-optimal environments (Gasol et al., 1995; Lennon and Jones, 2011).

Further, inland communities (characterized by a low organic C, high ionic strength

environment) also contained many unique taxa relative to river and nearshore microbiomes,

reflective of a distinct selective envrionment (Table 1). In particular, we observed members of

the candidate phylum *OP3* in high abundance in the inland waterborne microbiome. *OP3* has

been shown to be abundant in other subsurface communities with high inorganic C

concentrations (Emerson et al., 2015) and appears to thrive in low oxygen environments,

potentially utilizing anaerobic metabolisms reliant on iron, manganese, and/or sulfur (Glöckner

et al., 2010). Our inland zone was characterized by high levels of inorganic C but remained

oxygenated year round (Table S1), and although anaerobic microniches can form in interstitial

water, the relative abundance of this taxon (~13% of microbiome, 2^{nd} most abundant phylum)

suggests that *OP3* may be able to persist under a broader range of redox conditions than

currently recognized. Further, nitrifying organisms *Thaumarcheaota* and *Nitrospira* were

prevalent in the inland zone. These organisms are chemosynthetic autotrophs, not dependent on

external organic material, and instead fix CO_2 through autotrophic pathways involving NH_4^+ and

NO_2^- oxidation. A higher level of variable selection in the inland zone relative to other zones

(Table 2) is consistent with niche diversification, and the unique physicochemical environment

of the inland zone may allow for diversification of oligotrophic organisms belonging to *OP3*,

Thaumarcheaota, and *Nitrospira*, while excluding heterotrophic organisms that are dominant in

other zones.

Despite pronounced biogeographical differences in microbiomes, our results indicate

some common selective features and/or dispersal routes across zones and habitats. We observed

high abundances of *Proteobacteria* and members of the PVC superphylum in all data subsets,

and *Bacteroidetes* was among the most abundant phyla in both waterborne and attached

microbiomes in nearshore and river zones (Table 1).

The ability of *Proteobacteria* to assimilate low-energy substrates, such as sulfate, nitrate,

and inorganic C—in addition to various organic C compounds—may contribute to their abilities

to thrive across physicochemically-distinct aquatic environments (Cottrell and Kirchman, 2000;

Brenner et al., 2005; Battin et al., 2016). For example, *Alphaproteobacteria* and

Betaproteobacteria exhibit diverse heterotrophic metabolisms (Sato et al., 2009; Newton et al.,

2011; Battin et al., 2016). *Gammaproteobacteria* can fix inorganic C and digest organic C

(Nikrad et al., 2014; Dyksma et al., 2016), while *Deltaproteobacteria* utilize organic C as carbon

sources and as electron donors (Lovley et al., 1998; Torres et al., 2010). Microbiomes with a

diverse community of *Proteobacteria* may therefore be well-adapted to fluctuations in organic C

availability, facilitating their persistence across all zones. Moreover, *Bacteriodetes* has been

shown to be prevalent in both waterborne and biofilm microbiomes in other systems—

preferentially degrading recalcitrant C compounds and recycling biomass (Cottrell and

Kirchman, 2000; Martin et al., 2015; Bennke et al., 2016). The presence of *Bacteroidetes* in

addition to *Proteobacteria* within nearshore and river microbiomes may therefore indicate niche

complementarity within these higher-C environments, possibly coincident with spatial

organization that is not captured in our measurements.

Additionally, members of the PVC superphylum were abundant in all data subsets and

possess distinctive physiologies. Many of these organisms have a proteinaceous membrane

conveying antibiotic resistance (Fuerst and Sagulenko, 2011; Speth et al., 2012) and/or

metabolize C1 compounds such as methane (Dunfield et al., 2007; Fuerst and Sagulenko, 2011;

Sharp et al., 2013). Although the biogeochemical implications of selection for these organisms

are unexplored, their unique ecology, ability to consume methane, and abundance within our

system merits future investigating into their role in C cycling in hyporheic environments.

Microbiome assembly.

Hydrodynamics can be a driver for both stochastic and deterministic assembly processes,

as advection can physically transport microorganisms across geographic barriers as well as

generate changes in physicochemical selective environments. However, biogeographical patterns

in microbiomes suggest a limited influence of dispersal relative to selection in our system. We

empirically assessed assembly processes structuring microbiomes using ecological null models

and, contrary to our hypothesis, our results indicate that selective pressures overwhelmed

dispersal processes in structuring microbiomes.

In particular, attached communities were nearly entirely assembled by homogeneous

selective pressures and displayed network topologies indicating a highly-structured microbiome

(*i.e.*, high clustering coefficient and low heterogeneity, Table S3). While our measurements do

not provide sufficient resolution to describe fine-scale spatial organization of biofilms, our

measurement scale likely captures multiple levels of organization and niche partitioning in each

sample, allowing for inferences regarding community-level interactions. Our results are

consistent with a recent review that noted the prevalence of niche-based assembly processes over

stochasticity in structuring stream-associated biofilm microbiomes (Battin et al., 2016). Battin *et*

al. (2016) suggest micro-niches in biofilms select for specific organisms and repel poorly-

adapted immigrating species. Indeed, substrate features may impose strong environmental filters

on microbiomes—sediment geochemistry (Carson et al., 2007; Jorgensen et al., 2012) and matrix

structure (Vos et al., 2013; Breulmann et al., 2014) can select for traits that enhance attachment

on a particular substrate.

Likewise, river microbiomes were assembled via homogeneous selection but were also

differentiated from nearshore and inland zone microbiomes via a small influence of dispersal

limitation (Table 2). Network structure was also consistent with a highly-organized river

microbiome (*i.e.*, high clustering coefficient and low heterogeneity, Table S3). A relatively

stable geochemical environment in the river may provide a steady selective environment despite

seasonal changes in temperature (Table S1, Fig. 1C). In particular, NPOC concentration was

approximately 2-5 times higher in river water than the nearshore and inland waterborne

environments. Furthermore, organic C associated with surface water intrusion has been shown to

be correlated with short-term increases in deterministic assembly processes and changes in

microbiome composition in our nearshore environment (Stegen et al., 2016). Thus, consistently

high NPOC concentration in the river may impose environmental filters for waterborne

heterotrophic organisms within river microbiomes, resulting in homogeneous selection through

time. Moreover, distinct river physicochemistry and physical filtering of microorganisms

(Brunke, 1999; Hartwig and Borchardt, 2014), as compared to nearshore and inland zones (Table

S1), may limit successful colonization of river-associated taxa within other zones.

Although deterministic assembly processes were prevalent forces in structuring

microbiomes, nearshore and inland waterborne microbiomes showed a greater influence of

stochasticity than other zones. Within the nearshore, significant fluxes of both groundwater and

surface water may contribute to spatial assembly processes (e.g., dispersal). We observed the

impact of both dispersal limitation and homogenizing dispersal on nearshore waterborne

microbiomes, suggesting that this zone may experience periods of comparatively low and high

rates of hydrologic transport (i.e., hydrologic mixing and interstial flow). Furthermore, sampling

locations for inland waterborne microbiomes were distributed across a broader spatial extent than

any other zone, perhaps facilitating our ability to detect dispersal limitation in these samples.

When we examined assembly processes differentiating waterborne microbiomes across

zones, we anticipated more impact from spatial processes due to an increase in geographic scale.

In contrast to this expectation, selection was responsible for more than 50% of community

dissimilarity in all comparisons. While we have suggested physical inhibition of microbial

dispersal as a possible mechanism generating microbiome dissimilarity across zones, our

analyses indicate that an overarching influence of selection generally outweighed influences of

spatial processes. Our results indicate that although spatial processes play a role in shaping

microbiomes, local physicochemical environments often limit the ability of physically-

transported microorganisms to outcompete local biota.

Indeed, nearshore-to-inland zones comparisons suggested that distinct selective pressures

in each environment were the dominant cause (>90%) of differences among waterborne

microbiomes. Likewise, spatial processes were overwhelmed by selection in nearshore-to-river

comparisons. Differences among these microbiomes were due to relatively even proportions of

homogenous and variable selection, with no evident seasonal trends in the balance between these

processes. Our system is characterized by pronounced geomorphic heterogeneity in the

subsurface environment that creates preferential flow paths for hydrologic mixing (Johnson et

al., 2015). The presence of both homogeneous and variable selection may indicate spatial

variation in surface water intrusion or local biogeochemical conditions across our sampling

locations. Homogenizing dispersal also had a small but detectable influence in nearshore-to-river

comparisons, supporting some hydrologic transport of microorganisms between these zones.

Lastly, assembly estimates between the inland and river zones provided the strongest

evidence for spatial processes in our system (42.5% stochasticity). In particular, inland-to-river

comparisons yielded the largest impact of dispersal limitation (26.8%), supporting a decay of

community similarity across increasing spatial distances (Green and Bohannan, 2006). Dispersal

limitation is, therefore, likely to play a more significant role in subsurface microbiome assembly

at larger spatial scales.

Network clusters, keystone taxa, and seasonal physicochemical change.

Given pervasive deterministic assembly, we used network analysis to identify specific

clusters of organisms whose relative abundances changed deterministically with changes in the

environment within each zone. Co-occurrence networks fragment into clusters that are sensitive

to variation in hydrological regimes (Widder et al., 2014; Febria et al., 2015) and have been used

to identify keystone taxa (González et al., 2010; Vick-Majors et al., 2014; Banerjee et al., 2015;

Banerjee et al., 2016). Our sampling scale aggregates multiple niches within each sample,

allowing us to infer niches with similar vs. contrasting responses to environmental stimuli. In

general, our results were consistent with our hypothesis that higher NPOC concentrations in

surface water would shift microbiomes towards heterotophy. Specifically, we found that the

prevelance of heterotrophic organisms increased with increased surface water contribution in

each zone, and below we discuss microbial physiologies in each data subset that corresponded to

seasonal changes in hydrology.

In attached microbiomes no relationships were present in the inland zone, but we found

that nearshore attached cluster 1 was favored during early season conditions with pronounced

surface water intrusion and low temperature (Fig. 2A-D). Clusters of organisms can denote both

positive and negative co-occurrence patterns, signifying similar or opposite (respectively)

ecological dynamics influencing taxa. For instance, organisms sharing a cluster may constitute

an ecological niche, whereby organisms either co-occur in similar environments due to shared

environmental requirements or be involved in beneficial interactions with each other (Shi et al.,

2016). Alternatively, clusters may contain organisms that are favored under opposing

environmental conditions or have negative interactions (e.g., competition, predation). Nearshore

attached cluster 1 contained two anticorrelated groups of organisms—group 1 consisted of

heterotrophic organisms (*Oxalobacteraceae*, *Comamonadaceae*, *Verrucomicrobiaceae*, and

Flavobacteriaceae) that decreased through time, while organisms in group 2 were primarily

autotrophic oligotrophs (*Crenarchaeaceae*, *Ectothiorhodospiraceae*, *Pelagibacteraceae*, and

families of *Alphaproteobacteria*) that increased through time. Because of the abundance of

organisms in group 1 compared to group 2 (4498 vs. 1987 total OTU counts), group 1 dictated

correlations between this cluster and the environment, supporting an association between

heterotrophic organisms and periods experiencing relatively high-organic C surface water

intrusion. However, as the abundance of cluster 1 declined in concert with groundwater intrusion

into the nearshore environment (Fig. 3A-D), organisms in group 2 increased, putatively

outcompeting heterotrophic organisms in group 1 under more oligotrophic conditions. The co-

association of heterotrophic and autotrophic organisms within a single cluster may indicate

comparatively strong tradeoffs between these lifestyles and merits further investigation into

niche dynamics and competitive interactions between these taxa. Further, we identified primary

keystone taxa as *Verrucomicrobiaceae* and *Crenarchaeaceae*, possibly denoting the principal

heterotrophic and autotrophic organisms in groups 1 and 2, respectively.

In waterborne microbiomes, we observed relationships between clusters of organisms in

all zones with hydrologic mixing and physicochemistry. Nearshore waterborne cluster 9

displayed a seasonal association with groundwater discharging conditions and high temperature

(Fig. 2E-H). Organisms were assigned to families highly abundant in the inland zone (*e.g.*,

Thaumarchaeota, candidate phylum *OP3*, Table 1, Table S3), and keystone taxa, in particular,

belonged to *OP3*. Trends in the relative abundance of this cluster may therefore denote enhanced

dispersal from the inland zone during groundwater discharge and/or a change in selection

favoring these organisms corresponding to physicochemical shifts in nearshore environment. The

inland waterborne microbiome also revealed two clusters related to environmental change.

Cluster 3 was present in low abundance, and increased with higher groundwater contribution

and temperature (Fig. 2I-L), and contained facultative anaerobic microorganisms (*Lentisphaeria*,

Gemmatimonadetes, *Anaerolineae, Alphaproteobacteria*, *Gammaproteobacteria*, *Chloroflexi*,

Clostridia, *Spirochaetes*) and methylotrophs (*Betaproteobacteria*, Table S3). Such organisms are

largely distinct from organisms found in the nearshore and river environments, supporting the

inference that physical and geochemical isolation drives differences between microbiomes. In

contrast, cluster 5 was associated with surface water intrusion and low temperature (cluster 5,

Fig. 2M-P). An influx of surface water-associated NPOC (Fig. 1B) should favor heterotrophic

microorganisms, and cluster 5 contained methanotrophs (*Methylacidiphilae, Betaproteobacteria)*

and diverse *Actinobacteria* and *Acidimicrobiia* involved in C cycling, possibly denoting a shift

towards heterotrophy under comparatively high-organic C conditions. While no keystone taxa

were found in cluster 5, *Chloracidobacteria* and a family of *Chloroflexi* were identified in cluster

3, meriting further investigation into their role in the hyporheic environment.

Finally, we identified one cluster of diverse river organisms that was positively

associated with Cl⁻ concentration, which may indicate significant groundwater discharge into the

river (Fig. 2Q-T, Table S3). In particular, *Cytophagia* was the most abundant organism in this

cluster and was common in the nearshore and river zones, but all other families in this cluster

were primarily found in the inland zone. These inland taxa were, however, in low abundance in

the river and may therefore indicate a small contribution of the inland microbiome to river

communities under discharging condtions. No keystone taxa were found in this cluster.

Linkages between network clusters and hyporheic metabolism.

Finally, we further investigated the involvement of each network cluster in microbial

metabolism and found only one cluster—cluster 1 of nearshore attached microorganisms—that

correlated with active biomass (ATP) and aerobic respiration (Raz). Positive correlations

between cluster 1 and both ATP and Raz (Fig. 3) indicate that organisms within this cluster are

either responsible for biomass growth and aerobic nutrient cycling or are facilitated by

conditions under which these processes are promoted. While skewed towards heterotrophy,

nearshore attached cluster 1 contained organisms with a broad range of metabolic capabilities,

only partially supporting our hypothesis that increases in heterotrophic organisms would

generate higher rates of biogeochemical activity. Complementary microbial resource use and

niche diversification resulting from physical heterogeneity in sediments has been proposed to

increase the rate and diversity of compounds metabolized by attached organisms in other systems

(Singer et al., 2010; Battin et al., 2016). Moreover, spatial organization and excretion of

extracellular enzymes that can be utilized across organisms in biofilms can allow for

complementarity that enhances rates of metabolism (Loreau et al., 2001; Naeem et al., 2012).

Nearshore attached cluster 1 may therefore constitute the foundation (or portion thereof) of this

complementary resource structure.

Conclusions.

Our research generates a framework describing the roles of selection and dispersal in

influencing microbal physiologies within hyporheic environments (Fig. 5A-B). We advance that

spatiotemporal patterns in ecological selection impose strong environmental filters in hyporheic

zones, limiting successful immigration of organisms across physicochemical gradients. For

example, in Fig. 5A, hydrologic processes transport microorganisms from two communities to a

third location, and the establishment of dispersing organisms in the third location is strongly

impacted by local selective filters. Each community consists of largely distinct organisms, but

contains a small proportion of similar organisms common to both dispersing communities, as

evidenced by high levels of determistic assembly processes in our system but overlapping taxa

between each zone. These selective filters also change with hydrologic mixing, inducing a shift

from heterotrophic to autotrophic metabolisms with a change from surface water-dominated to

groundwater-dominated hydrology (Fig 5B). This inference is supported by associations of

nearshore attached and waterborne clusters as well as inland waterborne clusters with hydrologic

mixing. All correlations between these clusters show a descrease in heterotrophs and an increase

autotrophic organisms associated with increased groundwater contributions to physicochemistry.

Thus, despite largely deterministically-assembled microbiomes, a small proportion of

common taxa across a continuum of groundwater- to surface water-dominated habitats indicate

some influence of microbial dispersal on microbiome composition. We also identify organisms

that change through seasonally dynamic hydrologic mixing conditions, providing insight into

hyporheic microbial physiology. In particular, our research suggests a tradeoff between

heterotrophic and autotrophic organisms under surface water- vs. groundwater-dominated

hydrology. We therefore propose that comparatively high-organic C conditions during surface

water intrusion into the hyporheic zone support heterotrophy, succumbing to autotrophy during

time periods of groundwater discharge. Enhanced rates of activity associated with a nearshore

attached cluster may also indicate a central biogeochemical role for taxa, and in particular

keystone taxa (*Verrucomicrobiaceae* and *Crenarchaeaceae*), within this cluster. As a whole, our

work shows that deterministic assembly may overwhelms dispersal processes in hyporheic

environments despite pronounced hydrologic connectivity, and provides a process-based

understanding of spatiotemporal patterns in microbiome composition and function in a zone of

elevated biogeochemical cycling.

Experimental Procedures.

Study Design.

This study was conducted in three zones (inland, nearshore, river) within the Hanford

300A (approximately 46^o 22' 15.80"N, 119^o 16' 31.52"W) in eastern Washington State (Slater et

al., 2010; Zachara et al., 2013; described in Graham et al., 2016a). Previous research has shown

redox conditions as well as microbiome composition and activity to be variable across space and

time in this system (Lin et al., 2012a; Lin et al., 2012b; Stegen et al., 2012; Graham et al.,

2016a). sThe inland zone lies within 250m of the Columbia River and is characterized by an

unconfined groundwater aquifer in the Hanford formation. It displays relatively stable

temperatures ($\sim15^oC$) and elevated concentrations of anions and inorganic carbon relative to

other zones (Table S1). In contrast, the Columbia River contains higher organic carbon and

lower ion concentrations with seasonally variable temperatures. Surface water from the river and

groundwater from the inland zone mix in the nearshore hyporheic zone. At high river stages,

surface water intrudes into the inland zone; and at low river stages, groundwater discharges into

the Columbia River. To monitor hydrologic mixing, we employed Cl⁻ as a conservative

groundwater tracer per Stegen *et al.* (2016).

Detailed sampling and analytical methods are in the Supplemental Material. Briefly,

attached (nearshore and inland only) and waterborne (all zones) microbiomes were obtained

from deployed colonization substrate and waterborne samples, respectively. Colonization

substrate was incubated *in situ* six weeks prior to removal. Samples were collected at three-week

intervals from March through November 2014, with the first waterborne samples collected in

March and the first attached samples collected in April after a six-week incubation period.

Numbers of samples obtained at each timepoint are list in Table S4. Attached samples were

obtained from fully screened stainless steel piezometers installed to 1.2m depth below the

riverbed (nearshore, 5.25cm inside diameter (MAAS Midwest, Huntley, IL)) or from established

groundwater wells (inland). Piezometers were installed across a zone of known groundwater-

surface water exchange—and given the very significant cost of installing groundwater wells into

the broader aquifer—we made use of existing groundwater wells that were as close as possible to

the river shoreline. The need to use multiple pre-existing near-river wells resulted in a larger

spatial extent covered by the wells than by the piezometers. While a larger inland spatial domain

may have contributed to higher incidence of spatial assembly processes, this effect would only

impact comparisons of within zone assembly processes (vs. across zone comparisons), and we

note a dominance of selection within the inland zone similar to other zones (Table 2).

Waterborne samples were obtained from galvanized piezometers located <1m from the

piezometers used to sample attached microbiomes. For each inland location, a single well was

used for both waterborne and attached samples. River water was sampled adjacent to the

piezometers. DNA was extracted from each sample using the MoBio PowerSoil kit (MoBio

Laboratories, Inc., Carlsbad, CA), and the 16S rRNA gene was sequenced on the Illumina MiSeq

platform as described in the Supplemental Material. Physicochemical properties, active biomass

(ATP), and aerobic respiration (Raz) were determined as per the Supplemental Material.

Statistical analysis.

All analyses were conducted in *R* software (http://cran.r-project.org/) unless otherwise

noted. Relative abundances of major microbial taxa at phylum- and class- levels were calculated

in each of five data subsets (nearshore attached, nearshore waterborne, inland attached, inland

waterborne, river). An apparent breakpoint in hydrology at July 22 was assessed by comparing

Cl⁻ and NPOC concentration pre- and post- July 22 with one-sided Mann Whitney U tests.

Temporal changes in temperature in each subset were assessed with quadratic regressions.

Microbiome dissimilarities were estimated as Bray-Curtis distance and analyzed through time

and across data subsets with PERMANOVA in QIIME (Caporaso et al., 2010).

Modeling assembly processes.

We implemented null modeling methods developed by Stegen *et al.* (2013; 2015) to

disentangle community assembly processes. The approach uses pairwise phylogenetic turnover

between communities, calculated using the mean-nearest-taxon-distance (βMNTD) metric

(Webb et al., 2008; Fine and Kembel, 2011), to infer the strength of selection. Communities were

evaluated for significantly less turnover than expected (βNTI < -2, homogeneous selection) or

more turnover than expected (βNTI > 2, variable selection) by comparing observed βMNTD

values to the mean of a null distribution of βMNTD values—and normalizing by its standard

deviation—to yield the beta-nearest taxon index (βNTI) (Stegen et al., 2012). Pairwise

community comparisons that did not deviate from the null βMNTD distribution were evaluated

for the influences of dispersal limitation and homogenizing dispersal by calculating the Raup-

Crick metric extended to account for species relative abundances (RC_{bray}), as per Stegen *et al.*

(2013; 2015). Observed Bray-Curtis dissimilarities were compared to the null distribution to

derive RC_{bray}. RC_{bray} values that were > 0.95, > -0.95 and < 0.95, or < -0.95 were interpreted as

indicating dispersal limitation, no dominant assembly process, or homogenizing dispersal,

respectively. Significance levels for βNTI and RC_{bray} are respectively based on standard

deviations—|βNTI| = 2 denotes two standard deviations from the mean of the null distribution—

and alpha values—|RC_{bray}| = 0.95 reflects significance at the 0.05 level. Inferences from both

βNTI and RC_{bray} have previously been shown to be robust (Dini-Andreote et al., 2015; Stegen et

al., 2015).

Co-occurrence network analysis.

Co-occurrence networks were constructed using Spearman's correlation at four

taxonomic levels – class, order, family, and OTU (singletons removed). To examine taxa with

moderate to strong co-occurrence patterns, correlations with *rho* $> +/- 0.6$ and FRD-corrected *P*

< 0.01 were imported into Cytoscape v3.3 for visualization and calculation of network

parameters. Networks were examined for deviations from randomness by comparing network

parameters to random networks generated with Network Randomiser 1.1 in Cytoscape (Barabasi-

Albert model with same number of nodes). Keystone taxa were identified using the Betweenness

Centrality metric (BC > 0), whereby increasing BC values indicate greater contribution of nodes

to network structure (González et al., 2010; Vick-Majors et al., 2014; Banerjee et al., 2016).

We identified clusters in Cytoscape using MCODE, using default parameters as per

Banerjee *et al.* (2015; 2016). We then calculated Pearson's momentum correlation between the

relative abundance of each cluster and selected parameters (Cl⁻, NPOC, temperature, ATP, and

Raz) to screen for clusters of interest ($P < 0.05$). Regression models (linear and quadratic, as

appropriate) were then fit between cluster abundance (dependent variable) and the selected

parameters (independent variables). Analyses were conducted at all taxonomic levels, but we

present family-level correlations of microbiomes with time and physicochemistry due to the

consistency of patterns. For correlations with microbiome activity and aerobic respiration, results

with were similar at family- and class-levels (Fig. S2 and Table S5), and we present class-level

results that exhibited better model fit.

Acknowledgements.

This research was supported by the US Department of Energy (DOE), Office of

Biological and Environmental Research (BER), as part of Subsurface Biogeochemical

Research Program's Scientific Focus Area (SFA) at the Pacific Northwest National

Laboratory (PNNL). PNNL is operated for DOE by Battelle under contract

DE-AC06-76RLO 1830. A portion of the research was performed using Institutional

Computing at PNNL.

585
References.

Adams, R.I., Miletto, M., Taylor, J.W., and Bruns, T.D. (2013) Dispersal in microbes: fungi in
indoor air are dominated by outdoor air and show dispersal limitation at short distances. *The*
ISME Journal **7**: 1262-1273.
Banerjee, S., Kirkby, C.A., Schmutter, D., Bissett, A., Kirkegaard, J.A., and Richardson, A.E.
(2016) Network analysis reveals functional redundancy and keystone taxa amongst bacterial
and fungal communities during organic matter decomposition in an arable soil. *Soil Biology*
and Biochemistry **97**: 188-198.
Banerjee, S., Baah‑Acheamfour, M., Carlyle, C.N., Bissett, A., Richardson, A.E., Siddique, T.
et al. (2015) Determinants of bacterial communities in Canadian agroforestry systems.
Environmental Microbiology.
Battin, T.J., Besemer, K., Bengtsson, M.M., Romani, A.M., and Packmann, A.I. (2016) The
ecology and biogeochemistry of stream biofilms. *Nature Reviews Microbiology* **14**: 251-263.
Behrendt, L., Larkum, A.W., Trampe, E., Norman, A., Sørensen, S.J., and Kühl, M. (2012)
Microbial diversity of biofilm communities in microniches associated with the didemnid
ascidian Lissoclinum patella. *The ISME Journal* **6**: 1222-1237.
Bennke, C.M., Krüger, K., Kappelmann, L., Huang, S., Gobet, A., Schüler, M. et al. (2016)
Polysaccharide utilisation loci of Bacteroidetes from two contrasting open ocean sites in the
North Atlantic. *Environmental Microbiology*.
Boulton, A.J., Findlay, S., Marmonier, P., Stanley, E.H., and Valett, H.M. (1998) The functional
significance of the hyporheic zone in streams and rivers. *Annual Review of Ecology and*
Systematics: 59-81.
Boulton, A.J., Datry, T., Kasahara, T., Mutz, M., and Stanford, J.A. (2010) Ecology and
management of the hyporheic zone: stream-groundwater interactions of running waters and
their floodplains. *Journal of the North American Benthological Society* **29**: 26-40.
Brenner, D.J., Krieg, N., Staley, J., and Garrity, G. (2005) Bergey's Manual of Systematic
Bacteriology, Vol 2: The Proteobacteria. *NY: Springer*.
Breulmann, M., Masyutenko, N.P., Kogut, B.M., Schroll, R., Dörfler, U., Buscot, F., and Schulz,
E. (2014) Short-term bioavailability of carbon in soil organic matter fractions of different
particle sizes and densities in grassland ecosystems. *Science of The Total Environment* **497**:
29-37.
Brunke, M. (1999) Colmation and depth filtration within streambeds: retention of particles in
hyporheic interstices. *International Review of Hydrobiology* **84**: 99-117.
Bryant, D.A., Costas, A.M.G., Maresca, J.A., Chew, A.G.M., Klatt, C.G., Bateson, M.M. et al.
(2007) Candidatus Chloracidobacterium thermophilum: an aerobic phototrophic
acidobacterium. *Science* **317**: 523-526.
Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K. et al.
(2010) QIIME allows analysis of high-throughput community sequencing data. *Nature*
Methods **7**: 335-336.
Carson, J.K., Rooney, D., Gleeson, D.B., and Clipson, N. (2007) Altering the mineral
composition of soil causes a shift in microbial community structure. *FEMS Microbiology*
Ecology **61**: 414-423.
Cline, L.C., and Zak, D.R. (2014) Dispersal limitation structures fungal community assembly in
a long‑term glacial chronosequence. *Environmental Microbiology* **16**: 1538-1548.

Cottrell, M.T., and Kirchman, D.L. (2000) Natural assemblages of marine proteobacteria and
members of the Cytophaga-Flavobacter cluster consuming low-and high-molecular-weight
dissolved organic matter. *Applied and Environmental Microbiology* **66**: 1692-1697.
Dini-Andreote, F., Stegen, J.C., van Elsas, J.D., and Salles, J.F. (2015) Disentangling
mechanisms that mediate the balance between stochastic and deterministic processes in
microbial succession. *Proceedings of the National Academy of Sciences* **112**: E1326-E1332.
Dunfield, P.F., Yuryev, A., Senin, P., Smirnova, A.V., Stott, M.B., Hou, S. et al. (2007) Methane
oxidation by an extremely acidophilic bacterium of the phylum Verrucomicrobia. *Nature*
450: 879-882.
Dyksma, S., Bischof, K., Fuchs, B.M., Hoffmann, K., Meier, D., Meyerdierks, A. et al. (2016)
Ubiquitous Gammaproteobacteria dominate dark carbon fixation in coastal sediments. *The*
ISME Journal.
Emerson, J.B., Thomas, B.C., Alvarez, W., and Banfield, J.F. (2015) Metagenomic analysis of a
high carbon dioxide subsurface microbial community populated by chemolithoautotrophs
and bacteria and archaea from candidate phyla. *Environmental Microbiology*.
Febria, C.M., Beddoes, P., Fulthorpe, R.R., and Williams, D.D. (2012) Bacterial community
dynamics in the hyporheic zone of an intermittent stream. *The ISME Journal* **6**: 1078-1088.
Febria, C.M., Hosen, J.D., Crump, B.C., Palmer, M.A., and Williams, D.D. (2015) Microbial
responses to changes in flow status in temporary headwater streams: a cross-system
comparison. *Frontiers in Microbiology* **6**: 522.
Ferrenberg, S., O'Neill, S.P., Knelman, J.E., Todd, B., Duggan, S., Bradley, D. et al. (2013)
Changes in assembly processes in soil bacterial communities following a wildfire
disturbance. *The ISME Journal* **7**: 1102-1111.
Fierer, N., and Jackson, R.B. (2006) The diversity and biogeography of soil bacterial
communities. *Proceedings of the National Academy of Sciences* **103**: 626-631.
Fine, P.V.A., and Kembel, S.W. (2011) Phylogenetic community structure and phylogenetic
turnover across space and edaphic gradients in western Amazonian tree communities.
Ecography **34**: 552-565.
Fuerst, J.A., and Sagulenko, E. (2011) Beyond the bacterium: planctomycetes challenge our
concepts of microbial structure and function. *Nature Reviews Microbiology* **9**: 403-413.
Fukami, T. (2004) Assembly history interacts with ecosystem size to influence species diversity.
Ecology **85**: 3234-3242.
Fukami, T., Dickie, I.A., Paula Wilkie, J., Paulus, B.C., Park, D., Roberts, A. et al. (2010)
Assembly history dictates ecosystem functioning: evidence from wood decomposer
communities. *Ecology Letters* **13**: 675-684.
Gasol, J., Del Giorgio, P., Massana, R., and Duarte, C. (1995) Active versus inactive bacteria:
size-dependence in a coastal marine plankton community. *Marine Ecology Progress Series*
128: 91-97.
Gianuca, A.T., Declerck, S.A., Lemmens, P., and De Meester, L. (2016) Effects of dispersal and
environmental heterogeneity on the replacement and nestedness components of β‐
diversity. *Ecology*.
Glöckner, J., Kube, M., Shrestha, P.M., Weber, M., Glöckner, F.O., Reinhardt, R., and Liesack,
W. (2010) Phylogenetic diversity and metagenomics of candidate division OP3.
Environmental Microbiology **12**: 1218-1229.

Gonzalez, A., and Bell, G. (2013) Evolutionary rescue and adaptation to abrupt environmental
change depends upon the history of stress. *Philosophical Transactions of the Royal Society of*
London B: Biological Sciences **368**: 20120079.
González, A.M.M., Dalsgaard, B., and Olesen, J.M. (2010) Centrality measures and the
importance of generalist species in pollination networks. *Ecological Complexity* **7**: 36-43.
Graham, E., Crump, A.R., Resch, C.T., Fansler, S., Arntzen, E., Kennedy, D. et al. (2016a)
Coupling spatiotemporal community assembly processes to changes in microbial
metabolism. *Frontiers in Microbiology* **7**: 1949.
Graham, E.B., Wieder, W.R., Leff, J.W., Weintraub, S.R., Townsend, A.R., Cleveland, C.C. et
al. (2014) Do we need to understand microbial communities to predict ecosystem function?
A comparison of statistical models of nitrogen cycling processes. *Soil Biology and*
Biochemistry **68**: 279-282.
Graham, E.B., Knelman, J.E., Schindlbacher, A., Siciliano, S., Breulmann, M., Yannarell, A. et
al. (2016b) Microbes as engines of ecosystem function: when does community structure
enhance predictions of ecosystem processes? *Frontiers in Microbiology* **7**.
Green, J., and Bohannan, B.J. (2006) Spatial scaling of microbial biodiversity. *Trends in ecology*
& evolution **21**: 501-507.
Hancock, P.J., Boulton, A.J., and Humphreys, W.F. (2005) Aquifers and hyporheic zones:
towards an ecological understanding of groundwater. *Hydrogeology Journal* **13**: 98-111.
Hartwig, M., and Borchardt, D. (2014) Alteration of key hyporheic functions through biological
and physical clogging along a nutrient and fine‐sediment gradient. *Ecohydrology*.
Hedin, L.O., von Fischer, J.C., Ostrom, N.E., Kennedy, B.P., Brown, M.G., and Robertson, G.P.
(1998) Thermodynamic constraints on nitrogen transformations and other biogeochemical
processes at soil-stream interfaces. *Ecology* **79**: 684-703.
Johnson, T., Versteeg, R., Thomle, J., Hammond, G., Chen, X., and Zachara, J. (2015) Four‐
dimensional electrical conductivity monitoring of stage‐driven river water intrusion:
Accounting for water table effects using a transient mesh boundary and conditional inversion
constraints. *Water Resources Research* **51**: 6177-6196.
Jorgensen, S.L., Hannisdal, B., Lanzén, A., Baumberger, T., Flesland, K., Fonseca, R. et al.
(2012) Correlating microbial community profiles with geochemical data in highly stratified
sediments from the Arctic Mid-Ocean Ridge. *Proceedings of the National Academy of*
Sciences **109**: E2846-E2855.
Knelman, J.E., and Nemergut, D.R. (2014) Changes in community assembly may shift the
relationship between biodiversity and ecosystem function. *Frontiers in Microbiology* **5**.
Lauber, C.L., Hamady, M., Knight, R., and Fierer, N. (2009) Pyrosequencing-based assessment
of soil pH as a predictor of soil bacterial community structure at the continental scale.
Applied and Environmental Microbiology **75**: 5111-5120.
Lennon, J.T., and Jones, S.E. (2011) Microbial seed banks: the ecological and evolutionary
implications of dormancy. *Nature Reviews Microbiology* **9**: 119-130.
Lin, X., Kennedy, D., Fredrickson, J., Bjornstad, B., and Konopka, A. (2012a) Vertical
stratification of subsurface microbial community composition across geological formations at
the Hanford Site. *Environmental Microbiology* **14**: 414-425.
Lin, X., McKinley, J., Resch, C.T., Kaluzny, R., Lauber, C.L., Fredrickson, J. et al. (2012b)
Spatial and temporal dynamics of the microbial community in the Hanford unconfined
aquifer. *The ISME Journal* **6**: 1665-1676.

Lindström, E.S., and Langenheder, S. (2012) Local and regional factors influencing bacterial
community assembly. *Environmental Microbiology Reports* **4**: 1-9.
Loreau, M., Naeem, S., Inchausti, P., Bengtsson, J., Grime, J., Hector, A. et al. (2001)
Biodiversity and ecosystem functioning: current knowledge and future challenges. *science*
294: 804-808.
Lovley, D., Fraga, J.L., Blunt-Harris, E.L., Hayes, L., Phillips, E., and Coates, J.D. (1998)
Humic substances as a mediator for microbially catalyzed metal reduction. *Acta*
hydrochimica et hydrobiologica **26**: 152-157.
Marmonier, P., Archambaud, G., Belaidi, N., Bougon, N., Breil, P., Chauvet, E. et al. (2012) The
role of organisms in hyporheic processes: gaps in current knowledge, needs for future
research and applications. In *Annales de Limnologie-International Journal of Limnology*:
Cambridge Univ Press, pp. 253-266.
Martin, M., Barbeyron, T., Martin, R., Portetelle, D., Michel, G., and Vandenbol, M. (2015) The
Cultivable Surface Microbiota of the Brown Alga Ascophyllum nodosum is Enriched in
Macroalgal-Polysaccharide-Degrading Bacteria. *Frontiers in Microbiology* **6**.
Martiny, A.C., Treseder, K., and Pusch, G. (2013) Phylogenetic conservatism of functional traits
in microorganisms. *The ISME Journal* **7**: 830-838.
Martiny, J.B.H., Bohannan, B.J., Brown, J.H., Colwell, R.K., Fuhrman, J.A., Green, J.L. et al.
(2006) Microbial biogeography: putting microorganisms on the map. *Nature Reviews*
Microbiology **4**: 102-112.
McClain, M.E., Boyer, E.W., Dent, C.L., Gergel, S.E., Grimm, N.B., Groffman, P.M. et al.
(2003) Biogeochemical hot spots and hot moments at the interface of terrestrial and aquatic
ecosystems. *Ecosystems* **6**: 301-312.
Naeem, S., Duffy, J.E., and Zavaleta, E. (2012) The functions of biological diversity in an age of
extinction. *Science* **336**: 1401-1406.
Naegeli, M.W., and Uehlinger, U. (1997) Contribution of the hyporheic zone to ecosystem
metabolism in a prealpine gravel-bed-river. *Journal of the North American Benthological*
Society: 794-804.
Nemergut, D.R., Schmidt, S.K., Fukami, T., O'Neill, S.P., Bilinski, T.M., Stanish, L.F. et al.
(2013) Patterns and processes of microbial community assembly. *Microbiology and*
Molecular Biology Reviews **77**: 342-356.
Newton, R.J., Jones, S.E., Eiler, A., McMahon, K.D., and Bertilsson, S. (2011) A guide to the
natural history of freshwater lake bacteria. *Microbiology and Molecular Biology Reviews* **75**:
14-49.
Niederdorfer, R., Peter, H., and Battin, T.J. (2016) Attached biofilms and suspended aggregates
are distinct microbial lifestyles emanating from differing hydraulics. *Nature Microbiology* **1**:
16178.
Nikrad, M.P., Cottrell, M.T., and Kirchman, D.L. (2014) Uptake of dissolved organic carbon by
gammaproteobacterial subgroups in coastal waters of the West Antarctic Peninsula. *Applied*
and Environmental Microbiology **80**: 3362-3368.
Robertson, A.L., and Wood, P.J. (2010) Ecology of the hyporheic zone: origins, current
knowledge and future directions. *Fundamental and Applied Limnology/Archiv für*
Hydrobiologie **176**: 279-289.
Sato, K., Kato, Y., Taguchi, G., Nogawa, M., Yokota, A., and Shimosaka, M. (2009)
Chitiniphilus shinanonensis gen. nov., sp. nov., a novel chitin-degrading bacterium belonging
to Betaproteobacteria. *The Journal of General and Applied Microbiology* **55**: 147-153.

Sharp, C.E., den Camp, H.J.M.O., Tamas, I., and Dunfield, P.F. (2013) Unusual Members of the
PVC Superphylum: The Methanotrophic Verrucomicrobia Genus "Methylacidiphilum". In
Planctomycetes: Cell Structure, Origins and Biology: Springer, pp. 211-227.
Shi, S., Nuccio, E.E., Shi, Z.J., He, Z., Zhou, J., and Firestone, M.K. (2016) The interconnected
rhizosphere: High network complexity dominates rhizosphere assemblages. *Ecology Letters*.
Singer, G., Besemer, K., Schmitt-Kopplin, P., Hödl, I., and Battin, T.J. (2010) Physical
heterogeneity increases biofilm resource use and its molecular diversity in stream
mesocosms. *PLoS One* **5**: e9988.
Slater, L.D., Ntarlagiannis, D., Day‐Lewis, F.D., Mwakanyamale, K., Versteeg, R.J., Ward, A.
et al. (2010) Use of electrical imaging and distributed temperature sensing methods to
characterize surface water–groundwater exchange regulating uranium transport at the
Hanford 300 Area, Washington. *Water Resources Research* **46**.
Speth, D.R., van Teeseling, M.C.F., and Jetten, M.S.M. (2012) Genomic analysis indicates the
presence of an asymmetric bilayer outer membrane in Planctomycetes and Verrucomicrobia.
Frontiers in Microbiology **3**.
Stegen, J.C., Lin, X., Konopka, A.E., and Fredrickson, J.K. (2012) Stochastic and deterministic
assembly processes in subsurface microbial communities. *The ISME Journal* **6**: 1653-1664.
Stegen, J.C., Lin, X., Fredrickson, J.K., and Konopka, A.E. (2015) Estimating and mapping
ecological processes influencing microbial community assembly. *Frontiers in Microbiology*
6.
Stegen, J.C., Lin, X., Fredrickson, J.K., Chen, X., Kennedy, D.W., Murray, C.J. et al. (2013)
Quantifying community assembly processes and identifying features that impose them. *The
ISME Journal* **7**: 2069-2079.
Stegen, J.C., Fredrickson, J.K., Wilkins, M.J., Konopka, A.E., Nelson, W.C., Arntzen, E.V. et al.
(2016) Groundwater-surface water mixing shifts ecological assembly processes and
stimulates organic carbon turnover. *Nature Communications* **7**.
Tank, M., and Bryant, D.A. (2015) Nutrient requirements and growth physiology of the
photoheterotrophic Acidobacterium, Chloracidobacterium thermophilum. *Frontiers in
Microbiology* **6**: 226.
Torres, C.I., Marcus, A.K., Lee, H.-S., Parameswaran, P., Krajmalnik-Brown, R., and Rittmann,
B.E. (2010) A kinetic perspective on extracellular electron transfer by anode-respiring
bacteria. *FEMS Microbiology Reviews* **34**: 3-17.
Vick-Majors, T.J., Priscu, J.C., and Amaral-Zettler, L.A. (2014) Modular community structure
suggests metabolic plasticity during the transition to polar night in ice-covered Antarctic
lakes. *The ISME Journal* **8**: 778-789.
Vos, M., Wolf, A.B., Jennings, S.J., and Kowalchuk, G.A. (2013) Micro-scale determinants of
bacterial diversity in soil. *FEMS Microbiology Reviews* **37**: 936-954.
Wang, J., Shen, J., Wu, Y., Tu, C., Soininen, J., Stegen, J.C. et al. (2013) Phylogenetic beta
diversity in bacterial assemblages across ecosystems: deterministic versus stochastic
processes. *The ISME Journal* **7**: 1310-1321.
Webb, C.O., Ackerly, D.D., and Kembel, S.W. (2008) Phylocom: software for the analysis of
phylogenetic community structure and trait evolution. *Bioinformatics* **24**: 2098-2100.
Widder, S., Besemer, K., Singer, G.A., Ceola, S., Bertuzzo, E., Quince, C. et al. (2014) Fluvial
network organization imprints on microbial co-occurrence networks. *Proceedings of the
National Academy of Sciences* **111**: 12799-12804.

Woodcock, S., Besemer, K., Battin, T.J., Curtis, T.P., and Sloan, W.T. (2013) Modelling the
effects of dispersal mechanisms and hydrodynamic regimes upon the structure of microbial
communities within fluvial biofilms. *Environmental Microbiology* **15**: 1216-1225.
Zachara, J.M., Long, P.E., Bargar, J., Davis, J.A., Fox, P., Fredrickson, J.K. et al. (2013)
Persistence of uranium groundwater plumes: Contrasting mechanisms at two DOE sites in
the groundwater–river interaction zone. *Journal of Contaminant Hydrology* **147**: 45-72.
816
817

818
Figures and Tables.

Figure 1. Changes in waterborne (A) Cl⁻ concentration, (B) NPOC concentration, and (C)

temperature through time in each zone are presented in Figure 1. Lowess smoothers are plotted

in panels (A) and (B) and quadratic regressions are plotted in (C) to aid in visualization. Data

from nearshore, inland, and river zones are plotted with open, gray, and black circles,

respectively. River stage dynamics are presented in Figure S1.

Figure 2. Regressions between the abundance of selected clusters and day of year (column 1),

Cl⁻ concentration (column 2), non-purgable organic carbon (NPOC) (column 3), and temperature

(column 4). Nearshore attached cluster 1 is presented in row 1 (panels A-D), nearshore

waterborne cluster 9 is in row 2 (E-H), inland waterborne cluster 3 is in row 3 (I-L), inland

waterborne cluster 5 is in row 4 (M-P), and river cluster 2 is in row 5 (Q-T).

836   **Figure 3.** The abundance of nearshore attached cluster 1 was positively associated with (A)

837   active microbial biomass and (B) aerobic respiration. Class-level analyses are presented here;

838   family level-analyses are presented in Figure S2.

Figure 4. Network topology of clusters of organisms that correlated with time, Cl⁻, NPOC,

temperature, ATP, and/or Raz (see Figure 3) are presented in (A) nearshore attached cluster 1,

(B) nearshore waterborne cluster 9, (D) inland waterborne cluster 3, (E) inland waterborne

cluster 5, and (E) river cluster 2. Edge thickness denotes the strength of Spearman's correlation

between two nodes (ranging from 0.6 to 1.0), and keystone taxa are denoted as black nodes with

labels. All other nodes are numbered according to Table S3, and BC values for all nodes are also

listed in table S3. Solid edges represent positive correlations and dashed edges represent negative

correlations.

Figure 5. Conceptual diagrams depicting (A) influences of assembly processes on micriobiomes in our system and (B) microbiome shifts associated with hydrologic change in the nearshore hyporheic zone. In (A), hydrologic processes transport microorganisms from two communities to a third location. Each community consists of largely distinct organisms denoted by color (red vs. yellow), but contains a small proportion of similar organisms represented by the blue organism common to both dispersing communities. The establishment of dispersing organisms in the third location (purple zone) is strongly impacted by selective filters that limit successful immigration of organisms across environments. The local environment therefore selects for a unique microbiome (purple organisms) with the exception of a blue organism common to all three communities. Panel (B) depicts changes in microbiomes from heterotrophic to autotrophic metabolisms as hydrologic mixing conditions shift from surface water-dominated to groundwater-dominated. Mixing conditions are displayed as the background color, with green indicating surface water and blue indicating groundwater. Heterotrophic organisms are displayed in red, and autotrophic organisms are displayed in yellow.

864 **Table 1.** Phyla and classes with the highest relative abundance during our study period for each data subset. Mean relative abundance

865 is presented with standard deviations listed in parentheses.

Phyla

Nearshore attached		Nearshore waterborne		Inland attached		Inland waterborne		River	
Proteobacteria	34.82% (4.49%)	Proteobacteria	29.35% (0.01%)	Proteobacteria	60.66% (1.85%)	Proteobacteria	27.66% (7.40%)	Bacteroidetes	28.82% (7.58%)
Bacteroidetes	19.12% (6.78%)	Bacteroidetes	17.38% (0.01%)	Bacteroidetes	8.13% (4.19%)	Unassigned	17.32% (7.65%)	Actinobacteria	23.35% (7.76%)
Planctomycetes	13.74% (4.09%)	Actinobacteria	14.67% (7.21%)	Planctomycetes	5.27% (2.80%)	OP3	12.91% (4.13%)	Proteobacteria	23.31% (4.82%)
Verrucomicrobia	11.91% (3.35%)	Verrucomicrobia	8.16% (13.22%)	Acidobacteria	4.41% (1.64%)	Actinobacteria	7.45% (2.30%)	Verrucomicrobia	10.07% (5.71%)
Acidobacteria	6.03% (1.84%)	Planctomycetes	6.68% (8.08%)	Crenarchaeota	4.07% (1.28%)	Bacteroidetes	5.98% (1.99%)	Planctomycetes	5.18% (3.87%)
Crenarchaeota	2.68% (2.68%)	Unassigned	5.13% (3.81%)	Verrucomicrobia	2.93% (1.70%)	Planctomycetes	4.43% (1.44%)	Cyanobacteria	4.66% (2.79%)
Unassigned	2.50% (0.80%)	Acidobacteria	4.11% (3.76%)	Actinobacteria	2.76% (1.23%)	Crenarchaeota	4.00% (1.80%)	Acidobacteria	1.37% (3.15%)
Actinobacteria	1.59% (0.64%)	OP3	3.15% (3.70%)	Unassigned	2.44% (1.68%)	Verrucomicrobia	2.87% (1.45%)	Armatimonadetes	7.51% (1.09%)
Nitrospirae	1.56% (1.18%)	Cyanobacteria	2.44% (3.87%)	Nitrospirae	2.29% (1.00%)	Acidobacteria	2.58% (1.21%)	Unassigned	6.08% (0.41%)
Chloroflexi	1.28% (0.87%)	Crenarchaeota	2.1% (3.46%)	Chloroflexi	1.72% (0.50%)	Nitrospirae	2.17% (1.36%)	Chloroflexi	5.4% (0.38%)
Remainder	4.76% (1.10%)	Remainder	6.83% (2.86%)	Remainder	5.32% (1.31%)	Remainder	12.65% (0.22%)	Remainder	1.35% (0.05%)

Class

Nearshore attached		Nearshore waterborne		Inland attached		Inland waterborne		River	
Gammaproteobacteria	12.49% (2.95%)	Actinobacteria	12.06% (6.67%)	Betaproteobacteria	30.50% (12.56%)	Unassigned	17.32% (5.71%)	Actinobacteria	21.19% (7.52%)
Planctomycetia	12.07% (3.74%)	Betaproteobacteria	10.22% (3.38%)	Gammaproteobacteria	17.78% (9.71%)	Betaproteobacteria	8.68% (5.98%)	Betaproteobacteria	10.95% (2.27%)
Betaproteobacteria	9.20% (3.06%)	Alphaproteobacteria	9.98% (5.22%)	Alphaproteobacteria	9.40% (3.33%)	Alphaproteobacteria	7.71% (8.69%)	Alphaproteobacteria	9.80% (6.27%)
Alphaproteobacteria	9.11% (2.40%)	Saprospirae	5.81% (3.15%)	Planctomycetia	4.37% (3.23%)	PBS-25	7.34% (4.37%)	Flavobacteriia	9.46% (3.02%)
Pedosphaerae	6.93% (1.49%)	Planctomycetia	5.18% (3.28%)	Thaumarchaeota	4.07% (4.19%)	Actinobacteria	6.41% (6.85%)	Saprospirae	9.45% (3.01%)
Saprospirae	5.79% (1.74%)	Unassigned	5.13% (3.70%)	Chloracidobacteria	3.09% (1.82%)	Gammaproteobacteria	5.53% (3.77%)	Sphingobacteriia	5.94% (4.30%)
Flavobacteriia	5.05% (3.88%)	Flavobacteriia	4.73% (5.44%)	Unassigned	2.44% (1.28%)	koll11	5.52% (3.35%)	Planctomycetia	4.55% (3.92%)
Cytophagia	4.15% (1.78%)	Gammaproteobacteria	4.44% (2.29%)	Deltaproteobacteria	2.35% (1.54%)	Deltaproteobacteria	5.45% (1.87%)	Synechococcophycideae	4.10% (2.82%)
Chloracidobacteria	3.84% (1.52%)	Deltaproteobacteria	4.42% (2.57%)	Saprospirae	2.29% (1.90%)	Thaumarchaeota	3.85% (1.95%)	Cytophagia	3.77% (1.82%)
Deltaproteobacteria	3.81% (0.85%)	Sphingobacteriia	4.01% (3.10%)	Nitrospira	2.29% (1.70%)	Planctomycetia	2.59% (1.67%)	Opitutae	3.65% (4.75%)
Remainder	27.54% (1.49%)	Remainder	34.02% (2.74%)	Remainder	21.43% (1.84%)	Remainder	29.58% (2.47%)	Remainder	17.15% (1.25%)

866
867

869 **Table 2.** Assembly processes influencing microbiome composition as percentage of assembly mechanisms. Assembly processes are

870 defined according to Stegen et al. (2015, see main text), with the "undominated" fraction being the residual fraction of comparions

871 after accounting for all other assembly processes.

	Variable Selection	Homogeneous Selection	Total Selection	Dispersal Limitation	Homogenizing Dispersal	Undominated	Total Stochasticity
Within Geographic Zone							
Nearshore attached	0	100	100	0.0	0.0	0.0	0
Nearshore waterborne	6.8	52.9	59.8	6.6	14.9	18.7	40.2
Inland attached	0.4	82.1	82.5	1.7	1.7	14.0	17.5
Inland waterborne	14.3	21.5	35.7	22.1	7.2	35.6	64.3
River	0	95.5	95.5	0.0	3.0	1.5	4.5
Across Geographic Zones							
Inland-to-nearshore waterborne	91.96	0	91.96	1.79	1.79	4.46	8.04
Nearshore-to-river waterborne	34.84	29.18	64.02	2.83	15.3	17.85	35.98
Inland-to-river waterborne	51.17	6.33	57.5	26.76	1.64	14.1	42.5

872