TITLE

Germ granules prevent accumulation of somatic transcripts in the adult *C. elegans* germline

AUTHORS

Andrew Kekūpa'a Knutson*[†], Thea Egelhofer*[†], Andreas Rechtsteiner*, and Susan Strome*

AFFILIATIONS

*Department of Molecular, Cell & Developmental Biology, University of California Santa Cruz, Santa

Cruz, CA 95064

[†]These authors contributed equally

GEO ACCESSION NUMBER

RUNNING TITLE

P granules antagonize somatic RNAs

KEYWORDS

Germline, *C. elegans*, P granules, PGL-1, GLH-1, germ-soma antagonism

CORRESPONDING AUTHOR INFORMATION

Susan Strome

Molecular, Cell, and Developmental Biology

1156 High St., Santa Cruz, CA 95064

(831) 459-4615

E-mail: sstrome@ucsc.edu

ABSTRACT

The germ cells of multicellular organisms protect their developmental potential through

specialized mechanisms. A shared feature of germ cells from worms to humans is the presence of

non-membrane-bound ribonucleoprotein organelles called germ granules. Depletion of germ

granules in *Caenorhabditis elegans* (i.e., P granules) leads to sterility and in some germlines

expression of the neuronal transgene *unc-119::gfp* and the muscle myosin MYO-3. Thus, P

granules are hypothesized to maintain germ cell totipotency by preventing somatic development,

although the mechanism by which P granules carry out this function is unknown. In this study, we

performed transcriptome and single molecule RNA-FISH analyses of dissected P-granule-depleted

gonads at different developmental stages. Our results demonstrate that P granules are necessary

for adult germ cells to down-regulate spermatogenesis RNAs and to prevent the accumulation of

numerous soma-specific RNAs. P-granule-depleted gonads that express the *unc-119::gfp*

transgene also express many other genes involved in neuronal development and concomitantly

lose expression of germ cell fate markers. Finally, we show that removal of either of two critical P-

granule components, PGL-1 or GLH-1, is sufficient to cause germ cells to express UNC-119::GFP

and MYO-3 and to display RNA accumulation defects similar to those observed after depletion of P

granules. Our data identify P granules as critical modulators of the germline transcriptome and

guardians of germ cell fate.

INTRODUCTION

A cell's fate and function are determined by its transcriptome and proteome. The

transcriptomes of germ cells are unique and are comprised of RNAs that encode proteins essential

for specialized, germ cell-specific functions, such as meiosis, spermatogenesis, and oogenesis

(Lesch and Page 2012). Throughout development, germ cells must maintain expression of

germline-specific genes as well as repress genes that would lead to somatic differentiation (Strome

and Updike 2015). Failure to maintain appropriate gene expression programs can lead to sterility,

preventing an organism from passing its genetic information to future generations.

A conserved feature of germ cells is the presence of electron-dense, non-membrane-bound

organelles called germ granules, which include polar granules in *Drosophila*, chromatoid bodies in

mouse spermatids, Balbiani bodies in mouse oocytes, and P granules in *C. elegans* (reviewed in

Voronina *et al*. 2011). Germ granules are typically associated with the cytoplasmic face of germ

cell nuclei and have been observed to overlie clusters of nuclear pores in worms, zebrafish, and

mice (Pitt *et al*. 2000; Knaut *et al.* 2000; Chuma *et al.* 2009). Many essential germ-granule factors

are known or predicted to bind RNA. Thus, germ granules are suspected to serve as

posttranscriptional processing centers to modify, degrade, and/or store mRNAs as they exit germ

cell nuclei (Voronina *et al*. 2011; Updike and Strome 2010).

In *C. elegans*, P granules are maternally loaded into the 1-cell embryo and then

asymmetrically segregated to the germ lineage during embryogenesis (Updike and Strome 2010).

Although segregation of the bulk of maternal P granules to the germline blastomeres is not required

for germ cell specification (Gallo *et al.* 2010), P granules are required for proper germ cell

development and may protect germ cells from stressful conditions such as high temperature

(Kawasaki *et al*. 1998; Kawasaki *et al*. 2004; Spike *et al*. 2008; Gallo *et al*. 2010). The PGL and

GLH proteins make up the core constitutive components of P granules. The three PGL proteins are

worm-specific. The two most important PGL proteins, PGL-1 and PGL-3, contain a predicted RNA-

binding motif at their C-terminus called an RGG (Arg-Gly-Gly) box (Kawasaki *et al*. 1998; Kawasaki

et al. 2004). The four GLH proteins are homologs of the highly conserved Vasa protein and contain

DEAD-box helicase domains, which may also bind and modulate RNAs (Gruidl *et al.* 1996; Kuznicki

et al. 2000; Spike *et al.* 2008). The most important GLH protein, GLH-1, is necessary for P-granule

retention at the nuclear periphery (Updike *et al*. 2011). In addition to the PGL and GLH core

components, P granules contain proteins involved in small RNA biogenesis and the RNAi pathway,

suggesting that P granules are involved in small-RNA regulation in germ cells (Updike and Strome 2010; Kasper *et al.* 2014).

Recently P granules have been implicated in maintaining germ cell fate (Updike *et al*. 2014). Germ cells that are depleted of the most important P-granule components (PGL-1, PGL-3, GLH-1, and GLH-4) sometimes express a neuron-specific transgene and a muscle-specific myosin, supporting the hypothesis that P granules prevent somatic development in the *C. elegans* germline. These findings raise many questions, including: 1) What is the extent of somatic development in P-granule-depleted germlines? 2) Is any particular somatic fate favored over others? 3) How early do P-granule-depleted germ cells start expressing somatic markers? 4) What is the mechanism by which P granules prevent expression of somatic genes in the germline?

In this study, we investigated the role of P granules in maintaining appropriate transcript accumulation patterns in the *C. elegans* germline. Using transcript profiling and single molecule RNA-FISH of dissected gonads, we show that upon depletion of P granules, major changes in mRNA levels occur in the germline. Those changes include persistence of sperm transcripts past the normal sperm-production period in hermaphrodites and progressive up-regulation of transcripts from numerous genes normally expressed only in somatic cells. P-granule-depleted germ cells express numerous genes involved in neuronal fate and differentiation, and those cells lose their germ cell identity, highlighting the antagonistic relationship between germ and somatic fates. Finally, we show that germlines that lack either PGL-1 or GLH-1 also misexpress somatic markers, suggesting that these factors are critical for P-granule function. Our studies show that a key role of P granules is to maintain germ cell identity, perhaps by regulating the stability and/or transcription of target mRNAs.

MATERIALS AND METHODS

Strains and worm husbandry: *C. elegans* strains were maintained at 15° or 20° on OP50 bacteria as described (Brenner 1974). Strains used in this study include SS1174 wild type, DP132

edIs6[unc119p::unc119::gfp + rol-6] IV, SS0002 *pgl-1(ct131) him-3(e1147) IV,* SS0804 *glh-*

1(gk100) I/hT2g(I;III), SS1193 *glh-1(ok439) I/hT2g(I;III); edIs6[unc-119p::unc-119::gfp + rol-6] IV,*

and SS1162 *glh-1(ok439) glh-4(gk225) I/hT2g(I; III); edIs6[unc-119p::unc-119::gfp + rol-6] IV*.

Gonad isolation and RNA extraction: For experiments using RNAi, synchronized *unc-*

119::gfp L1 worms (P0) were placed on HT115 bacteria containing empty RNAi vector (control) or

expressing double strand RNA that targets P granules (i.e., *pgl-1*, *pgl-3*, *glh-1*, and *glh-4*) or targets

pgl-1 or *pgl-1*; *pgl-3* (Updike *et al*. 2014; Campbell and Updike 2015) at 24° (Day 0). Three days

later, F_1 larvae were transferred to new control or P-granule RNAi plates and returned to 24°. The

majority of these F_1 larvae were at the L1 or L2 stage at the time of transfer. The next day (Day 4),

L4 worms were visually identified by their size and the presence of a white crescent at their vulva.

Gonads were dissected from L4 control worms and L4 P-granule RNAi worms and cut at the gonad

bend with 30½ gauge needles in egg buffer (27.5 mM HEPES, 130 mM NaCl, 2.2 mM $MgCl_2$, 2.2

mM $CaCl_2$, and 0.528 mM KCl) containing 0.5% Tween-20 and 1 mM levamisole. Sterility could not

be assessed in L4 worms. However, greater than 80% sterility was observed in P-granule RNAi

adults that were aged from the population of worms from which L4 gonads were collected. For Day

1 adults, worms were allowed to age one day past the L4 stage and then dissected. For Day 2

adults, worms were allowed to age two days past the L4 stage and then dissected. In Day 2 adults,

gonads that visually expressed the *unc-119::gfp* transgene were collected as a separate sample

from those that did not detectably express *unc-119::gfp*. For adult control gonads, gonads were

dissected from fully fertile worms and cut at the gonad bend. For adult P-granule RNAi worms,

gonads were collected only from sterile worms, as judged by the lack of embryos in the uterus.

Large P-granule RNAi gonads were cut at the gonad bend; stunted P-granule RNAi gonads were

collected in their entirety. Once dissected, gonad arms were collected in Trizol, and total RNA was

extracted. For control samples, 75 to 100 gonads were collected per replicate. For P-granule RNAi

samples, 125 to 150 gonads were collected per replicate. Three biological replicates were

performed for each treatment and time point.

For experiments using P-granule mutants, wild-type (control), *pgl-1* single mutants, or *glh-1* single mutant adults (isolated from *glh-1(gk100)/hT2g* balanced parents) were allowed to lay eggs overnight on OP50 bacteria at 24°. The day after the egg lay, the adults were removed and the F_1 progeny were returned to 24° for two days. Day 1 adult gonads were dissected, collected into Trizol, and total RNA was extracted. Only distal gonad arms were collected from fertile wild-type worms or sterile P-granule mutant worms, as judged by the lack of embryos in the uterus. Three biological replicates were performed for the wild-type and *pgl-1* samples, and two biological replicates were performed for the *glh-1* sample.

Transcript profiling and analysis: Ribosomal RNA was depleted from each RNA sample using an NEBNext rRNA Depletion Kit (Human/Mouse/Rat) (Cat. #E6310), and libraries were constructed using an NEBNext Ultra RNA Library Prep Kit for Illumina sequencing (Cat. #E7530). Libraries were sequenced at the Vincent J. Coates Genomics Sequencing Laboratory at UC Berkeley using the Illumina HiSeq 2000, 2500, and 4000 platforms. Raw sequences were mapped to transcriptome version WS220 using Tophat2 (Kim *et al.* 2013). Only reads with one unique mapping were allowed; otherwise default options were used. Reads mapping to ribosomal RNAs were removed from further analysis. DESeq2 (Love *et al.* 2014) was used for differential expression analysis. A Benjamini-Hochberg multiple hypothesis corrected p-value cutoff of 0.05 was used as a significance cutoff. Read counts for each gene were normalized using the size factor normalization function in DESeq2 to ~3.5 million reads genome-wide per condition, after which replicate values were averaged. Gene Ontology analysis was done using the DAVID Bioinformatics Resource version 6.7.

Gene categories: Gene categories were defined using published microarray and SAGE datasets that profiled specific tissues or whole worms that contained or lacked a germline (Baugh *et al.* 2003; Reinke *et al.* 2004; Meissner *et al.* 2009; Wang *et al.* 2009), as described in (Rechtsteiner *et al*. 2010; Knutson *et al.* 2016). Ubiquitous genes (1,895 genes) are genes that are expressed (tag > 0) in all SAGE datasets that profiled germline, muscle, neuronal, and gut tissue, but that are

not in the germline-enriched category. Germline-enriched genes (2,230 genes) are genes with

transcripts enriched in adults with a germline compared with adults lacking a germline as assessed

by microarray analysis. Germline-specific genes (169 genes) are genes whose transcripts are

expressed exclusively in the adult germline and accumulate in embryos strictly by maternal

contribution. Sperm-specific genes (858 genes) are genes in the spermatogenesis-enriched

category as described in (Reinke et al. 2004) minus genes in the germline-enriched and germline-

specific categories. Soma-specific genes (1,181 genes) are genes expressed (tag > 4) in muscle,

neuronal, and/or gut tissue, but not expressed (tag = 0) in germline SAGE datasets and also not in

the germline-enriched category.

Unsupervised clustering analysis: The k-means clustering algorithm in R was used to

cluster the $\log_2$ fold changes shown in Figure S3. 14,571 genes significant in at least one of ten

comparisons (shown as fold change comparisons in Figure S3) at an adjusted p-value of 0.05 and

a $\log_2$ fold change of at least one were clustered. The number of clusters was increased until the

number of clusters and their patterns stabilized but were distinct, resulting in eight clusters.

Expression values in the various conditions are shown as DESeq2 normalized $\log_{10}$ read counts;

they were scaled by a factor of two, so that they would have a similar range as the fold change

values.   The DESeq2 size factor normalization scaled the samples to a total of ~3.5 million

genome-wide reads per condition. Within each cluster the genes were sorted from highest to

lowest average expression values across conditions, but expression values were not used to

cluster the genes. Our gene set categories are also shown, red indicating the gene being part of

the gene set. Boxplots of DESeq normalized $\log_{10}$ read counts for each gene cluster are shown

across conditions in Figure S4.

Single molecule RNA fluorescence in situ hybridization (smFISH): smFISH analysis

was performed on L4, Day 1 Adult, and Day 2 Adult control and P-granule RNAi gonads dissected

from *unc-119::gfp* worms as described (Lee *et al*. 2016). Stellaris probes were designed,

synthesized, and labeled by Biosearch Technologies. The *let-2* and *aqp-2* probes were coupled to

Cal Fluor Red 610, and the *cebp-1*, *puf-9*, and *lag-1* probes were coupled to TAMRA. Each probe

was resuspended in 250 µL of TE buffer pH 8.0 and then further diluted for hybridization (*let-2* and

lag-1 probes were diluted 1:10, *aqp-2* 1:30, and *cebp-1* and *puf-9* 1:50). Gonads were imaged on a

Solamere spinning disk confocal system controlled by Micro-Manager software (Edelstein *et al.*

2014). The set-up was as follows: Yokogawa CSUX-1 scan head, Nikon TE2000-E inverted stand,

Hamamatsu ImageEM x2 camera, 561 nm laser, and Plan Apo 60x/1.4 n.a. oil objective. Figures 4,

6E, S5, and S6B contain montages generated by splicing together contiguous images acquired with

identical settings. Control and P-granule RNAi pairs used identical confocal settings. All images

were processed with Image J and Adobe Illustrator.

Immunocytochemistry: For antibody staining, gonads were dissected using 30½ gauge

needles in egg buffer containing 0.5% Tween-20 and 1 mM levamisole. Dissected gonads were

permeabilized using the freeze-crack method, fixed for 10 min in methanol and 10 min in acetone,

and stained as described (Strome and Wood 1983). Antibody dilutions were 1:4000 rabbit anti-

UNC-64 (Saifee *et al*. 1998), 1:1000 rabbit anti-REC-8 (Liu *et al.* 2011), 1:500 guinea pig anti-HTP-

3 (MacQueen *et al*. 2005), 1:2000 mouse anti-GFP (Roche, Cat. #11814460001), 1:2000 rabbit

anti-GFP (Novus, Cat. #NB600-308), 1:2000 mouse anti-MYO-3 (Priess and Thomson 1987), and

1:500 Alexa Fluor secondary antibodies (Life Technologies). Images were acquired on the

Solamere spinning disk confocal system described above. Figures 5C, 6A, 6C, and S7B contain

montages generated by splicing together contiguous images acquired using identical settings. All

images were processed with ImageJ and Adobe Illustrator.

For quantification of REC-8 or HTP-3 signal relative to UNC-119::GFP signal, three optical

slices were projected for both control and P-granule RNAi gonads. Using the color histogram tool

in ImageJ, the intensity of the red channel (REC-8 or HTP-3 staining) was recorded and plotted as

a function of the intensity of the green channel (UNC-119::GFP) from random individual cells within

the gonad. For each experiment, one control gonad and three P-granule RNAi gonads were

sampled.

Data availability: Strains and reagents are available upon request. File S1 contains processed gene expression data including fold changes and average read counts. Gene expression data are available at GEO with the accession number XXX.

RESULTS

P-granule RNAi effectively depletes P-granule mRNAs from the germline: Depletion of P granules from the *C. elegans* germline causes sterility and, in some sterile worms, expression of the pan-neuronal *unc-119::gfp* transgene (Updike *et al.* 2014). Among UNC-119::GFP(+) gonads, approximately 60% also co-expressed the muscle myosin MYO-3 (Updike *et al.* 2014). UNC-119::GFP and MYO-3 were often observed in the same regions of the germline, and those regions showed morphological changes not typical of germ cells (i.e., loss of the rachis, loss of proper germ cell organization, and abnormal nuclear size). These features raise the possibility that depletion of P granules causes germ cells to undergo large-scale developmental changes in gene expression.

To identify genes that become misregulated upon depletion of P granules and to investigate the extent of reprogramming toward somatic cell types, we performed transcriptome analysis of dissected gonads from P-granule-depleted worms compared to control worms. We treated L1 larvae that harbored the *unc-119::gfp* transgene with P-granule RNAi and manually dissected and collected gonads from their F_1 progeny at different developmental stages: L4 larvae, Day 1 sterile adults, and Day 2 sterile adults (Figure 1A, see Materials and Methods). Throughout this report, we use the shorthand "L4" to mean the L4 larval stage and "PG(-)" to mean P-granule depleted. Of the Day 2 sterile adults, approximately 10% expressed the *unc-119::gfp* transgene in one or both gonad arms. These "transformed" gonads (i.e., Day 2 Adult PG(-) UNC-119::GFP(+) gonads) were collected as a separate sample from Day 2 gonads that did not detectably express the *unc-119::gfp* transgene (i.e., Day 2 Adult PG(-) UNC-119::GFP(-) gonads; Figure 1B). For comparison, worms were fed empty vector RNAi instead of P-granule RNAi, and control gonads were collected from fertile F_1 worms at the same developmental stages (L4, Day 1 adult, and Day 2 adult). From

control worms, only the distal halves of gonads were profiled by cutting at the gonad bend so as to

avoid oocytes, which are not present in P-granule-depleted gonads. From P-granule-depleted

worms, large gonads were cut at the gonad bend and stunted gonads were collected in their

entirety. Three biological replicates were collected for each treatment at each time point. Total

RNA was isolated from these replicates, depleted of ribosomal RNA, and used to construct libraries

for next-generation sequencing.

Our P-granule depletion method was highly effective. Compared to control gonads, P-

granule-depleted gonads at all three developmental stages showed between a 9- and 133-fold

reduction in *pgl-1*, *pgl-3*, *glh-1*, and *glh-4* transcripts (Figure S1A and B). In addition to the four P-

granule targets included in our RNAi construct, our P-granule RNAi also depleted *glh-2* and the

DEAD-box helicase domain-containing pseudogene *T08D2.3*, likely due to their extensive

sequence identity with *glh-1* and *glh-4*. UCSC genome browser tracks of these P-granule loci

revealed read pile-ups spanning approximately 350 base pairs (Figure S1A). Based on their

mapped locations, we infer that these reads originate from the RNAi construct. We also observed a

4-fold increase of a portion of *unc-119* transcripts in Day 2 Adult PG(-) UNC-119::GFP(+) gonads

compared to Day 2 Adult control gonads (Figure S1C). The reads that mapped to *unc-119* are

most likely from the *unc-119::gfp* transgene and not the endogenous *unc-119* locus, since only

exons 2, 3, and half of 4 were up-regulated and only those exons are present in the transgene.

Taken together, these data show that our P-granule RNAi treatment is highly effective at depleting

P-granule mRNAs and that Day 2 Adult PG(-) UNC-119::GFP(+) gonads up-regulate transcription

of the *unc-119::gfp* transgene.

P-granule-depleted adult gonads display progressive changes in transcript

accumulation: We observed genome-wide changes in transcript accumulation in P-granule-

depleted adult gonads compared to control gonads. Using cut-offs of an adjusted p-value of less

than 0.05 and a fold change of more than 2-fold, we identified 2,383 genes as significantly up-

regulated and 127 genes as significantly down-regulated in Day 1 Adult PG(-) UNC-119::GFP(-)

gonads compared to Day 1 Adult control gonads (Figure 2A). Allowing worms to age to Day 2 of

adulthood resulted in 6,833 genes significantly up-regulated and 965 genes significantly down-

regulated in Day 2 Adult PG(-) UNC-119::GFP(-) gonads compared to Day 2 Adult control gonads

(Figure 2A). Day 2 Adult PG(-) UNC-119::GFP(+) gonads showed even greater transcript

accumulation changes when compared to Day 2 Adult control gonads: 8,777 genes significantly up-

regulated and 3,049 genes significantly down-regulated (Figure 2A). Comparison of the different

up- and down-regulated genes from each adult sample revealed high overlap but also unique

genes within later gonad samples (Figure 2B). The nested Venn diagrams suggest that gene

misexpression in P-granule-depleted adult gonads becomes progressively more severe over time

and especially in gonads that express a somatic transgene (Figure 2B).

P-granule-depleted L4 larval gonads display little change in transcript accumulation:

In contrast to Day 1 and Day 2 Adult PG(-) gonads, L4 PG(-) gonads showed essentially no

transcript accumulation changes compared to L4 control gonads; only nine genes were significantly

down-regulated and 12 genes significantly up-regulated (Figure 2A). Of the nine down-regulated

genes, five genes were direct P-granule RNAi targets (i.e., *pgl-1*, *pgl-3*, *glh-1*, *glh-4*, *glh-2*) and two

genes were likely the result of 3' to 5' transitive RNAi spreading (*him-3* and *hil-4*, which are at the 5'

ends of *pgl-1* and *pgl-3*, respectively; Alder *et al.* 2003). Another gene, *T12F5.2*, is down-regulated

as a result of being in the same operon as *glh-4.* The 12 genes significantly up-regulated in P-

granule-depleted L4 gonads did not fall into any specific Gene Ontology category.

Principle component analysis (PCA) of all seven samples revealed clustering of the Day 1

and Day 2 Adult control gonad samples close to each other and far from the L4 control sample and

all of the PG(-) samples along PC1 (Figure 2C). Surprisingly, the control L4 gonad sample is

similar to all of the PG(-) samples along PC1 (Figure 2C). This along with results discussed below

suggests that gonads isolated from P-granule-depleted adult worms are developmentally more

similar to L4 gonads than adult gonads and possibly maintain mRNAs that are normally restricted to

the L4 stage. These results suggest that P granules are necessary for the normal progression of

germline development from the L4 stage to adult. The PG(-) samples spread out along PC2, with

the Day 2 Adult PG(-) UNC-119::GFP(+) gonad sample being farthest from the others. This likely

reflects differences in transcript accumulation that may underlie somatic development in

transformed Day 2 Adult PG(-) UNC-119::GFP(+) gonads.

P-granule-depleted adult gonads down-regulate germline transcripts and up-regulate

somatic and sperm transcripts: To determine the types of genes that are misregulated in

response to P-granule depletion, we analyzed the levels of mapped reads across different gene

categories: ubiquitously expressed, enriched expression in the germline, expressed specifically in

the germline, expressed specifically in sperm, or expressed specifically in somatic cells. These

categories were defined previously using microarray and serial analysis of gene expression (SAGE)

data from different tissues and from animals that possessed or lacked a germline (see Materials

and Methods). After normalizing for sequencing depth across all seven samples, we compared

average read counts across different gene categories.

Ubiquitous genes displayed constant levels of read counts across all seven samples (Figure

3A). Closer examination of individual genes revealed a small subset of ubiquitous genes that

showed up-regulation in P-granule-depleted gonads, especially in Day 2 Adult PG(-) UNC-

119::GFP(+) gonads (Figure S2A). However, for the majority of ubiquitous genes, including *ama-1*

which encodes the large subunit of RNA Polymerase II, read levels were not changed by P-granule

depletion (Figure 3B).

Germline-enriched and germline-specific gene read counts in control gonads increased

progressively from L4 larvae to Day 1 adults to Day 2 adults (Figure 3A). Upon P-granule

depletion, L4 PG(-), Day 1 Adult PG(-), and Day 2 Adult PG(-) UNC-119::GFP(-) gonads showed

levels of reads across germline genes that were similar to levels observed in L4 control gonads

(Figure 3A). Those levels dropped in Day 2 Adult PG(-) UNC-119::GFP(+) gonads compared to

Day 2 Adult control gonads (Figures 3A and S2B). Examples of germline transcripts that were

down-regulated include transcripts encoding the P-granule-associated argonaute CSR-1 and the P-

granule-associated translational repressor GLD-1 (Figure 3C). These data suggest that P granules

are required for the normal accumulation of germline transcripts and prevent the decrease in

germline mRNAs in gonads that express the *unc-119::gfp* transgene.

Sperm genes displayed elevated read counts in P-granule-depleted adult gonads compared

to control adult gonads (Figure 3A and D). In hermaphrodite worms, spermatogenesis occurs

during the L4 larval stage and ends at the L4-to-adult molt, at which time oogenesis begins (Hirsh

et al. 1976). Consistent with this, transcripts from sperm-specific genes, such as the *msp* genes

that encode MSPs or Major Sperm Proteins, accumulated in L4 gonads and then abruptly declined

from control adult gonads. However, these transcripts remained elevated in P-granule-depleted

adult gonads (Figure 3D and Figure S2C), supporting the suggestion from PCA analysis that P

granules are necessary for germlines to progress from L4 to adulthood. These data are consistent

with a recent study that also detected perdurance of sperm transcripts in P-granule RNAi Day 1

adult gonads (Campbell and Updike 2015).

Soma-specific genes showed several patterns of increased transcript accumulation in P-

granule-depleted adult gonads (Figures 3E, S3, and S4). Some soma-specific transcripts

accumulated in a pattern resembling sperm transcripts (transcripts detected in L4 gonads, down-

regulated in control adult gonads, and present at elevated levels in P-granule-depleted adult

gonads), including the alpha 2(IV) collagen mRNA, *let-2* (Figure 3E)*.* Other transcripts, like *cebp-1*,

which encodes a transcription factor important in axon regeneration in neurons (Yan *et al.* 2009),

and *puf-9*, which encodes a Pumilio-related RNA-binding protein that is expressed in the

hypodermis and neurons (Nolde *et al*. 2007), had very low read levels in control gonads but were

significantly up-regulated in P-granule-depleted adult gonads at all stages (Figures 3E and S6A). In

contrast to genes like *let-2, cebp-1*, and *puf-9*, some soma-specific transcripts were only

significantly up-regulated in Day 2 Adult PG(-) UNC-119::GFP(+) gonads, including *aqp-2,* which

encodes a membrane-bound aquaporin that is normally expressed in the excretory cell, muscle

cells, motor neurons, and the hypodermis (Huang *et al.* 2007; Figure 3E). Although different

somatic genes displayed different patterns of up-regulation (Figure 3E), as a whole, the levels of somatic gene transcripts progressively increased over time and with expression of the *unc-119::gfp* transgene (Figures 3A and S2D).

As an unbiased, genome-wide approach to determine different gene expression patterns among our samples, we also performed k-means clustering using the fold changes of RNAs in P-granule-depleted gonads versus their paired controls (see Materials and Methods). The 14,571 genes that showed a significant fold change in at least one of our comparisons grouped into eight clusters. The majority of ubiquitous genes, germline-enriched genes, and germline-specific genes grouped into Cluster 1, and the majority of sperm-specific genes grouped into Cluster 4 (Figures S3 and S4). This shows that these genes act similarly within their assigned gene category; that is, the majority of germline genes show similar responses to P-granule depletion, as do the majority of sperm genes. Somatic genes were spread out among the rest of the clusters (Clusters 2, 3, 5, 6, 7, and 8; Figure S3), supporting our observation that P-granule depletion causes several patterns of misregulation of soma-specific transcripts. Some somatic transcripts became more prevalent at progressively later stages (e.g., *cebp-1* in Cluster 2), while others were detected in L4 gonads, were reduced in control adult gonads, and persisted in PG(-) adult gonads (e.g., *let-2* in Cluster 6). We suspect that the somatic genes that become highly expressed in UNC-119::GFP(+) gonads (e.g., *aqp-2* and certain genes in Clusters 2, 3, and 6) are due to indirect, secondary effects of P-granule loss (see Discussion). Taken together, our transcript profiling shows that P granules are necessary to promote appropriate transcript accumulation patterns in gonads after the L4 larval stage.

Single molecule RNA-FISH analysis confirms that somatic transcripts accumulate in P-granule-depleted gonads: Although our RNA-seq method is highly sensitive, it is unable to provide spatial information about transcript accumulation patterns within gonads. As an independent method to test if, when, and where soma-specific transcripts are up-regulated in P-granule-depleted germ cells, we performed single molecule RNA-FISH (smFISH) using probes to

detect several soma-specific mRNAs (*cebp-1*, *aqp-2, let-2,* and *puf-9*) in control and P-granule-

depleted gonads.

By RNA-seq, *cebp-1* transcripts were not detected in control L4 or adult gonads but were

detected in P-granule-depleted gonads at all stages (Figure 3E). By smFISH, very few *cebp-1*

transcripts were detected in control gonads at any stage (Figure 4A). We did detect some *cebp-1*

transcripts around the nuclei of somatic gonad cells, including the very distal tip of the gonad, likely

the somatic distal tip cell, in Day 1 and Day 2 Adult control gonads, but no signal was detected in

the germline. Upon P-granule depletion, individual *cebp-1* mRNA molecules were present

throughout the germ cell region of Day 1 and Day 2 Adult PG(-) UNC-119::GFP(-) gonads as well

as Day 2 Adult PG(-) UNC-119::GFP(+) gonads (Figure 4A). Indeed, 58% of Day 1 Adult PG(-)

UNC-119::GFP(-) gonads (n=24) and 100% of Day 2 Adult PG(-) UNC-119::GFP(+) gonads (n=13)

had *cebp-1* transcripts throughout the germ cell region. Unlike *cebp-1* transcripts, which were

detected in younger P-granule-depleted gonads by RNA-seq, *aqp-2* transcripts were only up-

regulated in Day 2 Adult PG(-) UNC-119::GFP(+) gonads (Figure 3E). Similarly, by smFISH, *aqp-2*

transcripts were only detected in Day 2 Adult PG(-) UNC-119::GFP(+) gonads (Figure 4B), and

100% of Day 2 Adult PG(-) UNC-119::GFP(+) gonads had *aqp-2* transcripts (n=23). Like *cebp-1*

transcripts, *aqp-2* transcripts were distributed throughout the germline and sometimes concentrated

in regions around nuclei (Figure 4B).

We also used smFISH to analyze *let-2* and *puf-9* transcripts. By RNA-seq, *let-2* transcripts

in control gonads behaved in a similar manner to sperm transcripts; transcripts were detected in L4

gonads and decreased in adult gonads (Figure 3E). After P-granule depletion, *let-2* transcripts

persisted in adults and were especially elevated in Day 2 Adult PG(-) UNC-119::GFP(+) gonads

(Figure 3E). Interestingly, by smFISH, the *let-2* signal in both L4 control and L4 PG(-) gonads

mostly originated from *let-2* transcripts concentrated in the very distal tip of the gonad, likely in the

somatic distal tip cell and not in germ cells (Figure S5). In Day 2 Adult control gonads, *let-2*

transcripts were mostly gone, with only a few transcripts remaining at the distal tip (Figure S5). In

contrast, in 54% (n=36) of Day 2 Adult PG(-) UNC-119::GFP(+) gonads, *let-2* transcripts were

present throughout the gonad, frequently concentrated in regions around the nuclei of germ cells

(Figure S5). By RNA-seq, levels of *puf-9* transcripts were low in control gonads and elevated in

both adult stages lacking P granules (Figure S6A). By smFISH, *puf-9* transcripts were also

detected throughout the distal half of the germline in P-granule-depleted adult gonads: 71% of Day

1 Adult PG(-) (n=45) and 90% of Day 2 Adult PG(-) UNC-119::GFP(+) (n=19) (Figure S6B). Very

few *puf-9* transcripts were detected in the distal half of control gonads. We did detect *puf-9*

transcripts in the proximal half of both control and P-granule RNAi gonads. Taken together, our

smFISH analysis corroborates our RNA-seq results and confirms that soma-specific transcripts are

up-regulated in P-granule-depleted germ cells.

Our smFISH results shed light on the subset of soma-specific transcripts that are detected in

L4 gonads, are reduced in control adult gonads, but persist in P-granule-depleted adult gonads

(e.g., *let-2* transcripts in Figure 3E; Cluster 6 in Figures S3 and S4). Based on our RNA-seq data

alone, two explanations are possible. One possibility is that these transcripts are expressed in

germ cells at the L4 stage and then disappear in a P-granule-dependent manner in adult gonads,

similar to sperm transcripts. An alternative possibility is that these transcripts reside in the somatic

gonad and appear to be down-regulated in control adult gonads and up-regulated in P-granule-

depleted adult gonads because of the altered ratio of somatic gonad cells to germ cells, i.e. a lower

ratio in control adults and a higher ratio in P-granule-depleted adults with stunted germlines.

Illustrating the latter possibility, our smFISH analysis detected *let-2* transcripts in the somatic gonad

of L4s. However, *let-2* transcripts were later expressed in the germ cells of UNC-119::GFP(+)

adults, along with many other somatic transcripts (e.g., *cebp-1*, *aqp-2*, and *puf-9*). Thus, even

somatic gonad markers can become expressed in the germlines of reprogrammed adults. Further

research is needed to identify the direct versus indirect effects that P granules have on individual

genes.

UNC-119::GFP(+) gonads turn on numerous genes involved in neuronal development:

Most of the genes misregulated in Day 2 Adult PG(-) UNC-119::GFP(-) gonads were also misregulated in Day 2 Adult PG(-) UNC-119::GFP(+) gonads (Figure 2B). Day 2 Adult PG(-) UNC-119::GFP(+) gonads turned on many additional genes as well. Direct comparison of Day 2 Adult PG(-) UNC-119::GFP(+) and UNC-119::GFP(-) gonads identified 2,268 genes significantly up-regulated and 692 genes significantly down-regulated in UNC-119::GFP(+) gonads compared to UNC-119::GFP(-) gonads (Figure 5A). Gene Ontology analysis of the 2,268 up-regulated genes identified processes involved in neuron differentiation, synaptic transmission, and axon guidance, as well as a large number of genes involved in regulation of RNA metabolism (Figure 5B). Interestingly, we did not detect processes involved in development of other somatic cell types like hypodermis, intestine, or muscle, even though 60% of UNC-119::GFP(+) gonads also express the muscle myosin MYO-3 (Updike *et al*. 2014).

As an independent test of whether neuronal genes other than the *unc-119::gfp* transgene are up-regulated in Day 2 Adult PG(-) UNC-119::GFP(+) gonads, we immunostained dissected gonads for the *C. elegans* syntaxin homolog UNC-64. UNC-64 is a component of the core synaptic vesicle fusion machinery and is expressed throughout the nervous system (Ogawa *et al.* 1998; Saifee *et al.* 1998). In our transcriptome data, *unc-64* transcripts were significantly up-regulated in Day 2 Adult PG(-) UNC-119::GFP(+) gonads compared to Day 2 Adult control and Day 2 Adult PG(-) UNC-119::GFP(-) gonads. In control gonads, some UNC-64 staining was detected in the somatic sheath and distal tip cell but not in the germ cell region (Figure 5C). In contrast, in Day 2 Adult PG(-) UNC-119::GFP(+) gonads, UNC-64 staining was observed throughout the germ cell region and stained in a filamentous pattern (Figure 5C), which may indicate cell membranes since syntaxin is a membrane-anchored receptor (Saifee *et al.* 1998). These data show that Day 2 Adult PG(-) UNC-119::GFP(+) gonads up-regulate many genes involved in neuronal development, in addition to the *unc-119::gfp* transgene.

UNC-119::GFP(+) germ cells lose germline identity: Gene Ontology analysis of the 692 genes down-regulated in Day 2 Adult PG(-) UNC-119::GFP(+) versus UNC-119::GFP(-) gonads revealed processes involved in the cell cycle, meiosis, and chromosome segregation (Figure 5B). Interestingly, a large number of down-regulated genes were categorized as important for embryonic development, including genes whose gene products are maternally loaded into developing oocytes to support and guide early embryogenesis (e.g., *mex-5*, *mex-6*, *pie-1*, *nos-1*). To test if UNC-119::GFP(+) cells lose germ cell identity upon acquisition of a neuronal fate, we stained Day 2 Adult PG(-) UNC-119::GFP(+) gonads for the germline proteins REC-8 and HTP-3 and performed smFISH to detect *lag-1* transcripts. REC-8 is a cohesin found in the mitotic stem cell region of germlines (Hansen *et al*. 2004), and HTP-3 is an integral component of the synaptonemal complex present in the meiotic region (Severson *et al*. 2009; Figure 6A and 6C). LAG-1 is a transcription factor that works downstream of Notch signaling to control germ cell entry into meiosis (Crittenden *et al*. 2003). In Day 2 Adult PG(-) UNC-119::GFP(+) gonads, we observed a mutually exclusive pattern of UNC-119::GFP and REC-8 staining; cells that expressed UNC-119::GFP lacked REC-8 and vice versa (Figure 6A and B). This was also the case for HTP-3 (Figure 6C and D) and *lag-1* transcripts (Figure 6E). This demonstrates the antagonistic relationship between somatic and germline fates and highlights the importance of P granules in maintaining germ cell identity.

Taken together, our transcript profiling, immunostaining, and smFISH experiments demonstrate that changes to the transcriptome and proteome are occurring in at least a fraction of the germ cells in P-granule-depleted gonads. These changes reflect both the loss of germline identity (e.g., loss of REC-8, HTP-3, and *lag-1* mRNA) and acquisition of somatic identity (e.g., expression of the *unc-119::gfp* transgene and UNC-64).

The germlines from *pgl-1* and *glh-1* single mutants also express somatic markers: Our P-granule RNAi construct simultaneously targets four important P-granule genes: *pgl-1*, *pgl-3*, *glh-1*, and *glh-4*. We tested if loss of one or two P-granule components is sufficient to cause germline expression of somatic genes by crossing the *unc-119::gfp* transgene into *glh-1* single and

glh-1 glh-4 double mutant backgrounds and by performing single *pgl-1* RNAi or double *pgl-1; pgl-3*

RNAi on *unc-119::gfp* transgenic worms. We detected UNC-119::GFP expression in all

backgrounds tested (Figures 7A and S7). Additionally, of the germlines that expressed UNC-

119::GFP, a fraction also expressed the muscle myosin MYO-3 (Figures 7A and S7). To determine

if *pgl-1* and *glh-1* mutant gonads misregulate a similar set of genes, we performed transcript

profiling of gonads collected from sterile Day 1 Adult *pgl-1* mutants and sterile Day 1 Adult *glh-1*

mutants; neither mutant contained the *unc-119::gfp* transgene. Comparison to wild-type gonads

revealed more transcript accumulation changes in *pgl-1* mutants than in *glh-1* mutants; *pgl-1* single

mutants up-regulated 2,894 genes and down-regulated 273 genes, while *glh-1* single mutants up-

regulated 626 genes and down-regulated 110 genes (Figure 7B). The majority of the genes up-

regulated in *glh-1* gonads were also up-regulated in *pgl-1* gonads (592 out of 626 genes, 94%),

suggesting that PGL-1 and GLH-1 may work together to control certain transcripts (Figure 7C). The

majority of genes up-regulated in both *pgl-1* and *glh-1* mutant gonads were also up-regulated in

Adult P-granule RNAi gonads (Figures S3 and S4; data not shown). These results suggest that

PGL-1 and GLH-1 are both critical P-granule factors and that both are necessary to prevent

somatic development within the germline.

18   **DISCUSSION**

Although germ granules were first observed over 100 years ago, their functions are only

starting to be understood at the molecular level. *C. elegans* germ granules or "P granules" are

necessary for germline development and fertility (Kawasaki *et al*. 1998; Kawasaki *et al*. 2004; Spike

et al. 2008) and protect the germline from expressing somatic fate markers (Updike *et al*. 2014). In

this study, we analyzed the progressive transcript changes that occur in P-granule-depleted

germlines. We treated worms with P-granule RNAi for two generations and performed

transcriptome analysis of dissected gonads from the second-generation sterile hermaphrodites. In

worms that lack P granules, germlines appear relatively healthy at the L4 stage but deteriorate in

adults. Germline deterioration in adults is accompanied by and perhaps caused by aberrant

persistence of sperm-specific transcripts after the sperm-production period in L4, as previously

observed (Campbell and Updike 2015), and elevated levels of soma-specific transcripts. The

persistence of sperm-specific transcripts into adulthood suggests that P granules promote

progression of germ cells from the L4 larval stage to adulthood, and the healthy appearance and

normal RNA profile of L4 germlines suggest that P granules are not necessary for germline

development from embryogenesis to L4. In adults, the majority of P-granule-depleted germlines

become reduced in size and disorganized, while a minority of germlines appear to proliferate; both

categories of germlines can contain cells that have lost their previous germ cell fate and appear to

be on a neuronal developmental path. Taken together, these findings reveal that *C. elegans* germ

granules are major regulators of the germline transcriptome and prevent germ cells from

undergoing somatic development, possibly by preventing the expression of somatic factors in the

germline (Figure 8). The mechanism by which P granules perform these functions remains to be

determined.

P granules reside on the outer periphery of germ cell nuclei and overlie nuclear pores (Pitt *et*

al 2000; Sheth *et al*. 2010; Updike *et al*. 2011). It has been proposed that as mRNAs exit the

nucleus, P granules intercept and perhaps process these transcripts, thus influencing the

transcriptome and proteome of the germline (Kasper *et al*. 2014; Figure 8). Indeed, some germline

transcripts have been shown to at least transiently reside in P granules (Schisa *et al*. 2001; Sheth

et al. 2010). Recently, PGL-1 was shown to dimerize and to have guanosine-specific RNase

activity (Aoki *et al*. 2016). This raises the possibility that PGL-1 directly influences the

transcriptome by cleaving its associated mRNAs, although PGL-1's catalytic activity is not essential

for its role in fertility (Aoki *et al*. 2016). If PGL-1 indeed has RNase activity *in vivo*, then an

attractive scenario is that it recognizes mRNAs inappropriate for germline development and cleaves

them, while allowing germline-appropriate mRNAs to pass through P granules and be translated in

the cytoplasm (Figure 8). Alternatively, PGL-1 and the paralogous protein PGL-3 may not have a

preference for transcript type. Indeed, PGL-3 was recently shown to bind mRNA in extracts and *in vitro* in a sequence-independent manner (Saha *et al*. 2016). One possibility is that P granules intercept all types of transcripts, including germline-appropriate transcripts, but effectively prevent low levels of stochastically expressed soma-specific transcripts from producing levels of proteins that would threaten germ cell identity.

Our study extends a recent analysis of transcripts from Day 1 adult P-granule RNAi gonads (Campbell and Updike 2015) and documents the progressive appearance of somatic transcripts that underlies the germ-toward-soma reprogramming we previously observed in older P-granule RNAi adults (Updike *et al*. 2014). Campbell and Updike documented perdurance of sperm-specific transcripts in Day 1 adult P-granule RNAi gonads, as we describe in both Day 1 and Day 2 P-granule RNAi adults, but they did not detect increased accumulation of soma-specific transcripts in their samples. The Day 1 adult worms sampled by Campbell and Updike experienced weaker RNAi (based on RNA read counts from RNAi target genes) and contained healthier germlines than in our experiments. Thus, the defects observed by Campbell and Updike are probably the earliest defects and likely include the primary defects caused by P-granule depletion. Our stronger RNAi and additional analysis of Day 2 adults with progressively reprogrammed germlines identified additional consequences, probably many secondary consequences, of P-granule depletion. Based on their observations, Campbell and Updike concluded that P granules likely repress translation of soma-specific mRNAs that are already present at low levels in the germline (Figure 8, Model A). Our data suggest that P granules additionally affect the levels of soma-specific mRNAs in the germline, possibly by degrading soma-specific mRNAs and/or repressing transcription of soma-specific genes (Kasper *et al*. 2014; Figure 8, Model B).

Consistent with P granules functioning as regulators of mRNA accumulation in the germline, many RNAi factors localize to P granules, including Dicer/DCR-1, the RNA-dependent RNA polymerase EGO-1, and the argonaute proteins PRG-1 and CSR-1 (Batista *et al*. 2008; Claycomb *et al*. 2009). PRG-1 is a Piwi-related argonaute that prevents the expression of foreign genetic

elements, including transposon and transgene mRNAs. PRG-1 recognizes these mRNAs through

imperfect base-pairing with its associated 21U-RNAs, resulting in heritable silencing of transposon

and transgene loci (Batista *et al*. 2008; Ashe *et al*. 2012; Bagijn *et al*. 2012; Shirayama *et al*. 2012;

Lee *et al*. 2012). To maintain germline identity, PRG-1 may work with P-granule proteins to

recognize and degrade soma-specific transcripts identified as "foreign". This information could then

be relayed back into the nucleus to repress inappropriately activated soma-specific genes through

epigenetic mechanisms (Kasper *et al*. 2014; Figure 8). CSR-1, another P-granule associated

argonaute, recognizes its targets through associated 22G-RNAs that base-pair with germline-

expressed mRNAs including those important for embryogenesis (Claycomb *et al*. 2009).

Interestingly, CSR-1 was recently found to tune the levels of certain germline mRNAs in a manner

that is dependent on its slicer activity (Gerson-Gurwitz *et al*. 2016). Importantly, depletion of CSR-1

and depletion of P granules cause similar transcript accumulation changes (Campbell and Updike

2015), strengthening the link between argonautes and P granules in regulating the germline

transcriptome.

The development of different neuronal subtypes can be induced in the *C. elegans* germline

by simultaneous removal of certain chromatin factors and forced expression of master neuronal

transcription factors (Tursun *et al*. 2011; Patel *et al*. 2012), highlighting the plasticity of the germline.

Notably, other genetic perturbations have been shown to cause *C. elegans* germ cells to express

neuronal and muscle markers, without the need for driving transcription factors. These include loss

of the H3K4 methyltransferase SET-2, loss of the nuclear argonaute HRDE-1, and concomitant loss

of the NuRD complex component LET-418 and the H3K4 demethylase SPR-5 (Robert *et al*. 2014;

Kaser-Pebernard *et al*. 2014). Because these factors are chromatin-based, they are likely to have

direct effects on gene transcription. One possibility is that these factors work downstream of P

granules and interpret the RNA-processing events that occur within P granules. Comparing the

transcriptome of gonads lacking P granules to the transcriptomes of *set-2* or *let-418; spr-5* mutant

gonads may provide insight into how these regulatory mechanisms operate and influence each

other to protect germ cell identity.

Our Gene Ontology analysis of genes uniquely up-regulated in P-granule-depleted gonads

from Day 2 adults expressing the *unc-119::gfp* neuronal transgene revealed processes in neuronal

development. Interestingly, processes involved in muscle development and function were not

observed, despite the fact that 60% of UNC-119::GFP(+) gonads also express the muscle myosin

MYO-3 (Updike *et al*. 2014). Perhaps many fewer P-granule-depleted germ cells up-regulate

genes involved in muscle development than up-regulate genes involved in neuronal development,

which would bias our transcriptome analysis in favor of genes involved in neuron-related processes.

The expression of many neuronal genes in P-granule-depleted Day 2 adult germlines is

likely to be a secondary effect of P-granule loss. The burning challenges are to identify the primary

effect(s) of P-granule loss, and to determine why P-granule-depleted germ cells appear to favor a

neuronal path. One possibility is that these germ cells are reverting to a "default" neuronal

differentiation program, which has been observed in the contexts of dissociated *Xenopus* embryos

and mammalian embryonic stem cells (ESCs) grown in culture (Sato and Sargent 1989; Tropepe *et*

al. 2001). To acquire neuronal fate through this "default" program, ESCs must be cultured at low-

density and in growth factor-free conditions, suggesting an inherent propensity of ESCs to

differentiate into neurons (Smukler *et al*. 2006). Recently, the transcription factor BLIMP1 was

found to be essential to reprogram human induced pluripotent stem cells (hiPSCs) into primordial

germ cell (PGC)-like cells (Sasaki *et al*. 2015). Interestingly, when Sasaki *et al*. attempted to derive

PGC-like cells from *BLIMP1-/-* hiPSCs, essential PGC genes were not expressed and genes

involved in neuronal differentiation were up-regulated instead. These and our findings suggest that

neuronal fate is the preferred somatic fate of cells that lose their pluripotent stem cell or germline

identity. Future work should investigate if *C. elegans* germlines that lack P granules revert to a

"default" neuronal program and if so, how the neuronal program becomes activated and why it

becomes activated in adults and not larval germlines.

The breadth of ribonucleoprotein complexes and the regulatory circuits they form within the *C. elegans* germline highlight the complex nature of this tissue. In order to fully understand P-granule function, a thorough analysis of individual factors like PGL-1 and GLH-1 will need to be performed, especially at the biochemical level. Additionally, identification of P-granule-associated mRNAs may provide insight into how P granules prevent somatic development in the germline. Finally, because many P-granule proteins are conserved in other species, it is important to test if germ granules in other organisms including vertebrates prevent somatic development of germ cells.

AUTHOR CONTRIBUTIONS

A.K.K., T.E., and S.S. designed the research. A.K.K. performed the RNA-seq and immunostaining experiments. T.E. performed the mutant analyses and smFISH experiments. A.K.K. and A.R. performed the RNA-seq bioinformatic analysis. A.K.K., T.E. and S.S. wrote the paper.

ACKNOWLEDGEMENTS

We thank Geraldine Seydoux, David Greenstein, and Judith Kimble for advice on smFISH, Judith Kimble for the *lag-1* smFISH probe, Mike Nonet for the UNC-64 antibody, Dustin Updike for the *pgl-1; pgl-3* double RNAi construct, Ben Abrams at the UCSC Life Sciences Microscopy Facility for training, and past and current members of the Strome lab for helpful discussions. This work used the Vincent J. Coates Genomics Sequencing Laboratory at UC Berkeley, supported by NIH S10 Instrumentation Grants (S10RR029668, S10RR027303, and S10OD018174). Some strains used in this study were provided by the CGC, which is funded by the NIH Office of Research Infrastructure Programs (P40 OD010440). This work was supported by a CIRM predoctoral fellowship (TG2-01157) to A.K.K. and an NIH R01 grant (GM34059) and an Ellison Senior Scholar in Aging Award (AG-SS-3098-12) to S.S.

REFERENCES

Alder, M. N., S. Dames, J. Gaudet, and S. E. Mango, 2003 Gene silencing in *Caenorhabditis*

elegans by transitive RNA interference. RNA **9**: 25–32.

Aoki S. T., A. M. Kershner, C. A. Bingman, M. Wickens, and J. Kimble, 2016 PGL germ granule

assembly protein is a base-specific, single-stranded RNase. Proc. Natl. Acad. Sci. USA **113**:

1279–1284.

Ashe A., A. Sapetschnig, E. M. Weick, J. Mitchell, M. P. Bagijn *et al.*, 2012 piRNAs can trigger a

multigenerational epigenetic memory in the germline of *C. elegans*. Cell **150**: 88–99.

Bagijn M. P., L. D. Goldstein, A. Sapetschnig, E. Weick, S. Bouasker *et al.*, 2013 Function, targets,

and evolution of *Caenorhabditis elegans* piRNAs. Science **337**: 574–578.

Batista P. J., J. G. Ruby, J. M. Claycomb, R. Chiang, N. Fahlgren *et al.*, 2008 PRG-1 and 21U-

RNAs interact to form the piRNA complex required for fertility in *C. elegans*. Mol. Cell **31**: 67–

78.

Baugh L. R., A. A. Hill, D. K. Slonim, E. L. Brown, and Hunter C. P., 2003 Composition and

dynamics of the *Caenorhabditis elegans* early embryonic transcriptome. Development **130**:

889–900.

Brenner S., 1974 The genetics of *Caenorhabditis elegans*. Genetics **77**: 71–94.

Campbell A. C. and D. L. Updike, 2015 CSR-1 and P granules suppress sperm-specific

transcription in the *C. elegans* germline. Development **142**: 1745–1755.

Chuma S., M. Hosokawa, T. Tanaka, and N. Nakatsuji, 2009 Ultrastructural characterization of

spermatogenesis and its evolutionary conservation in the germline: Germinal granules in

mammals. Mol. Cell. Endocrinol. **306**: 17–23.

Claycomb J. M., P. J. Batista, K. M. Pang, W. Gu, J. J. Vasale *et al.*, 2009 The argonaute CSR-1

and its 22G-RNA cofactors are required for holocentric chromosome segregation. Cell **139**:

123–134.

Crittenden S. L., C. R. Eckmann, L. Wang, D. S. Bernstein, M. Wickens *et al.*, 2003 Regulation of

the mitosis/meiosis decision in the *Caenorhabditis elegans* germline. Philos. Trans. R. Soc. L.

B. Biol. Sci. **358**: 1359–1362.

Edelstein A. D., M. A. Tsuchida, N. Amodaj, H. Pinkard, R. D. Vale *et al.*, 2014 Advanced methods

of microscope control using µManager software. J. Biol. Methods **1**: 10.

Gallo C. M., J. T. Wang, F. Motegi, and G. Seydoux, 2010 Cytoplasmic partitioning of P granule

components is not required to specify the germline in *C. elegans*. Science **330**: 1685–1689.

Gerson-Gurwitz A., S. Wang, S. Sathe, R. Green, G. W. Yeo *et al.*, 2016 A Small RNA-Catalytic

Argonaute Pathway Tunes Germline Transcript Levels to Ensure Embryonic Divisions. Cell

165: 396–409.

Gruidl M. E., P. A. Smith, K. A. Kuznicki, J. S. McCrone, J. Kirchner *et al.*, 1996 Multiple potential

germ-line helicases are components of the germ-line-specific P granules of *Caenorhabditis*

elegans. Proc. Natl. Acad. Sci. USA **93**: 13837–13842.

Hansen D., E. J. A. Hubbard, and T. Schedl, 2004 Multi-pathway control of the proliferation versus

meiotic development decision in the *Caenorhabditis elegans* germline. Dev. Biol. **268**: 342–

357.

Hirsh D., D. Oppenheim, and M. Klass, 1976 Development of the reproductive system of

Caenorhabditis elegans. Dev. Biol. **49**: 200–219.

Huang C. G., T. Lamitina, P. Agre, and K. Strange, 2007 Functional analysis of the aquaporin gene

family in *Caenorhabditis elegans*. Am. J. Physiol. Cell Physiol. **292**: C1867–C1873.

Kaser-Pebernard S., F. Muller, and C. Wicky, 2014 LET-418/Mi2 and SPR-5/LSD1 cooperatively

prevent somatic reprogramming of *C. elegans* germline stem cells. Stem Cell Reports **2**: 547–

559.

Kasper D. M., K. E. Gardner, and V. Reinke, 2014 Homeland security in the *C. elegans* germ line:

Insights into the biogenesis and function of piRNAs. Epigenetics **9**: 62–74.

1. Kawasaki I., Y. H. Shim, J. Kirchner, J. Kaminker, W. B. Wood *et al.*, 1998 PGL-1, a predicted RNA-binding component of germ granules, is essential for fertility in *C. elegans*. Cell **94**: 635–645.

2. Kawasaki I., A. Amiri, Y. Fan, N. Meyer, S. Dunkelbarger *et al.*, 2004 The PGL family proteins associate with germ granules and function redundantly in *Caenorhabditis elegans* germline development. Genetics **167**: 645–661.

3. Kim D., G. Pertea, C. Trapnell, H. Pimentel, R. Kelley *et al.*, 2013 TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. Genome Biol. **14**: R36.

4. Knaut H., F. Pelegri, K. Bohmann, H. Schwarz, and C. Nüsslein-Volhard, 2000 Zebrafish vasa RNA but not its protein is a component of the germ plasm and segregates asymmetrically before germline specification. J. Cell Biol. **149**: 875–888.

5. Knutson A. K., A. Rechtsteiner, and S. Strome, 2016 Reevaluation of whether a soma-to-germ-line transformation extends lifespan in *Caenorhabditis elegans*. Proc. Natl. Acad. Sci. USA **113**: 3591–3596.

6. Kuznicki K. A., P. A. Smith, W. M. Leung-Chiu, A. O. Estevez, H. C. Scott *et al.*, 2000 Combinatorial RNA interference indicates GLH-4 can compensate for GLH-1; these two P granule components are critical for fertility in *C. elegans*. Development **127**: 2907–2916.

7. Lee H., W. Gu, M. Shirayama, E. Youngman, D. Conte *et al.*, 2012 *C. elegans* piRNAs mediate the genome-wide surveillance of germline transcripts. Cell **150**: 78–87.

8. Lee C., E. B. Sorensen, T. R. Lynch, and J. Kimble, 2016 *C. elegans* GLP-1/Notch activates transcription in a probability gradient across the germline stem cell pool. eLife **5**: e18370.

9. Lesch B. J. and D. C. Page, 2012 Genetics of germ cell development. Nat. Rev. Genet. **13**: 781–794.

10. Liu T., A. Rechtsteiner, T. A. Egelhofer, A. Vielle, I. Latorre *et al.*, 2011 Broad chromosomal domains of histone modification patterns in *C. elegans*. Genome Res. **21**: 227–236.

Love M. I., W. Huber, and S. Anders, 2014 Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol. **15**: 550.

MacQueen A. J., C. M. Phillips, N. Bhalla, P. Weiser, A. M. Villeneuve *et al.*, 2005 Chromosome

sites play dual roles to establish homologous synapsis during meiosis in *C. elegans*. Cell **123**:

1037–1050.

Meissner B., A. Warner, K. Wong, N. Dube, A. Lorch *et al.*, 2009 An integrated strategy to study

muscle development and myofilament structure in *Caenorhabditis elegans*. PLoS Genet. **5**:

e1000537

Nolde M., N. Saka, K. L. Reinert, and F. J. Slack, 2007 The *C. elegans pumilio* homolog, *puf-9*, is

required for the 3'UTR mediated repression of the *let-7* microRNA target gene, *hbl-1*. Dev. Biol.

305: 551–563.

Ogawa H., S. I. Harada, T. Sassa, H. Yamamoto, and R. Hosono, 1998 Functional properties of the

unc-64 gene encoding a *Caenorhabditis elegans* syntaxin. J. Biol. Chem. **273**: 2192–2198.

Patel T., B. Tursun, D. P. Rahe, and O. Hobert, 2012 Removal of Polycomb Repressive Complex 2

makes *C. elegans* germ cells susceptible to direct conversion into specific somatic cell types.

Cell Rep. **2**: 1178–1186.

Pitt J. N., J. A. Schisa, and J. R. Priess, 2000 P granules in the germ cells of *Caenorhabditis*

elegans adults are associated with clusters of nuclear pores and contain RNA. Dev. Biol. **219**:

315–333.

Priess J. R., and J. N. Thomson, 1987 Cellular interactions in early *C. elegans* embryos. Cell **48**:

241–250.

Rechtsteiner A., S. Ercan, T. Takasaki, T. M. Phippen, T. A. Egelhofer *et al.*, 2010 The histone

H3K36 methyltransferase MES-4 acts epigenetically to transmit the memory of germline gene

expression to progeny. PLoS Genet. **6**: e1001091.

Reinke V., I. S. Gil, S. Ward, and K. Kazmer, 2004 Genome-wide germline-enriched and sex-

biased expression profiles in *Caenorhabditis elegans*. Development **131**: 311–323.

Robert V. J., M. G. Mercier, C. Bedet, S Janczarski, J. Merlet *et al*., 2014 The SET-2/SET1 histone

H3K4 methyltransferase maintains pluripotency in the *Caenorhabditis elegans* germline. Cell

Rep. **9**: 443–450.

Saha S., C. A. Weber, M. Nousch, O. Adame-Arana, C. Hoege *et al*., 2016 Polar positioning of

phase-separated liquid compartments in cells regulated by an mRNA competition mechanism.

Cell **166**: 1572-1584.

Saifee O., L. Wei, and M. L. Nonet, 1998 The *Caenorhabditis elegans unc-64* locus encodes a

syntaxin that interacts genetically with synaptobrevin. Mol Biol Cell **9**: 1235–1252.

Sasaki K., S. Yokobayashi, T. Nakamura, I. Okamoto, Y. Yabuta *et al*., 2015 Robust in vitro

induction of human germ cell fate from pluripotent stem cells. Cell Stem Cell **17**: 178–194.

Sato S. M. and T. D. Sargent, 1989 Development of neural inducing capacity in dissociated

Xenopus embryos. Dev. Biol. **134**: 263–266.

Schisa J. A., J. N. Pitt, and J. R. Priess, 2001 Analysis of RNA associated with P granules in germ

cells of *C. elegans* adults. Development **128**: 1287–1298.

Severson A. F., L. Ling, V. van Zuylen, and B. J. Meyer, 2009 The axial element protein HTP-3

promotes cohesin loading and meiotic axis assembly in *C. elegans* to implement the meiotic

program of chromosome segregation. Genes Dev. **23**: 1763–1778.

Sheth U., J. Pitt, S. Dennis, and J. R. Priess, 2010 Perinuclear P granules are the principal sites of

mRNA export in adult *C. elegans* germ cells. Development **137**: 1305–1314.

Shirayama M., M. Seth, H. C. Lee, W. Gu, T. Ishidate *et al*., 2012 piRNAs initiate an epigenetic

memory of nonself RNA in the *C. elegans* germline. Cell **150**: 65–77.

Smukler S. R., S. B. Runciman, S. Xu, and D. van der Kooy, 2006 Embryonic stem cells assume a

primitive neural stem cell fate in the absence of extrinsic influences. J. Cell Biol. **172**: 79–90.

Spike C., N. Meyer, E. Racen, A. Orsborn, J. Kirchner *et al*., 2008 Genetic analysis of the

Caenorhabditis elegans GLH family of P-granule proteins. Genetics **178**: 1973–1987.

Strome S. and W. B. Wood, 1983 Generation of asymmetry and segregation of germ-line granules

in early *C. elegans* embryos. Cell **35**: 15–25.

Strome S. and D. Updike, 2015 Specifying and protecting germ cell fate. Nat. Rev. Mol. Cell Biol.

16: 406–416.

Tropepe V., S. Hitoshi, C. Sirard, T. W. Mak, J. Rossant *et al.*, 2001 Direct neural fate specification

from embryonic stem cells: a primitive mammalian neural stem cell stage acquired through a

default mechanism. Neuron **30**: 65–78.

Tursun B., T. Patel, P. Kratsios, and O. Hobert, 2011 Direct conversion of *C. elegans* germ cells

into specific neuron types. Science **331**: 304–308.

Updike D. and S. Strome, 2010 P granule assembly and function in *Caenorhabditis elegans* germ

cells. J. Androl. **31**: 53–60.

Updike D. L., S. J. Hachey, J. Kreher, and S. Strome, 2011 P granules extend the nuclear pore

complex environment in the *C. elegans* germ line. J. Cell Biol. **192**: 939–948.

Updike D. L., A. K. Knutson, T. A. Egelhofer, A. C. Campbell, and S. Strome, 2014 Germ-granule

components prevent somatic development in the *C. elegans* germline. Curr. Biol. **24**: 970–975.

Voronina E., G. Seydoux, P. Sassone-Corsi, and I. Nagamori, 2011 RNA granules in germ cells.

Cold Spring Harb. Perspect. Biol. **3**: a002774.

Wang X., Y. Zhao, K. Wong, P. Ehlers, Y. Kohara *et al.*, 2009 Identification of genes expressed in

the hermaphrodite germ line of *C. elegans* using SAGE. BMC Genomics **10**: 213.

Yan D., Z. Wu, A. D. Chisholm, and Y. Jin, 2009 The DLK-1 kinase promotes mRNA stability and

local translation in *C. elegans* synapses and axon regeneration. Cell **138**: 1005–1018.

                                    FIGURE LEGENDS

Figure 1. Method to profile transcripts from P-granule-depleted gonads. **A)** Experimental scheme

and timeline to analyze the transcriptomes of control and P-granule-depleted gonads from L4 larval,

Day 1 adult, and Day 2 adult stages of worms grown at 24^o. **B)** Seven different samples were

collected (three control, and four P-granule RNAi) all in triplicate. PG(-) is shorthand for P-granule

depletion. Throughout this report, the samples are color-coded: control samples are in light blue;

L4 and Day 1 Adult PG(-) UNC-119::GFP(-) samples are in pink; Day 2 Adult PG(-) UNC-

119::GFP(-) samples are in red; and Day 2 Adult PG(-) UNC-119::GFP(+) samples are in green.

Figure 2. Large, progressive changes in transcript accumulation are observed in adult P-granule-

depleted gonads but not in L4 larval gonads. **A)** Volcano plots comparing P-granule-depleted

gonads to control gonads at different time points. Dashed lines indicate the significance cut-off of

q=0.05 (horizontal line) and a 2-fold change in transcript level (vertical lines). The numbers of

significantly misregulated genes in each quadrant are shown at the top of each plot. **B)** Venn

diagrams showing the overlap of genes down- or up-regulated in Day 1 Adult PG(-) UNC-

119::GFP(-) gonads (pink), Day 2 Adult PG(-) UNC-119::GFP(-) gonads (red), and Day 2 Adult

PG(-) UNC-119::GFP(+) gonads (green). The overlap of all three samples is shaded in orange. **C)**

Principle component analysis (PCA) of all seven samples (P-granule-depleted and control samples)

across two principle components (PC1 and PC2).

Figure 3. P-granule-depleted gonads down-regulate germline genes and up-regulate sperm and

somatic genes. **A)** Boxplots of normalized RNA-seq reads from ubiquitous genes, genes with

germline-enriched expression, genes with germline-specific expression, sperm genes, and soma-

specific genes in control and P-granule-depleted gonads at different stages. Each box extends

from the 25th to the 75th percentile of read values. The whiskers extending from each box indicate

the 2.5th and the 97.5th percentiles. Wedges around the median indicate 95% confidence intervals

for the medians. **B-E)** UCSC genome browser tracks showing the abundance of RNA-seq reads

from control and P-granule-depleted gonad samples across (B) the ubiquitous gene *ama-1*, (C)

germline-specific genes *gld-1* and *csr-1*, (D) sperm-specific genes *msp-81*, *-10*, and *-56*, *nspd-2*,

and *ssp-10*, and (E) soma-specific genes *let-2*, *cebp-1*, and *aqp-2*. The sample names in the

boxplots and the genome browser tracks are abbreviated with L4 denoting L4 larvae, Ad1 denoting

Day 1 Adults, and Ad2 denoting Day 2 Adults. Additionally, UNC(-) denotes no expression and

UNC(+) denotes expression of the *unc-119::gfp* transgene. The boxplots and browser tracks are

color-coded as described for Figure 1.

Figure 4. Soma-specific transcripts accumulate in P-granule-depleted germlines. **A)** Single

molecule RNA-FISH (smFISH) analysis of *cebp-1* transcripts (red) in dissected Day 1 and Day 2

Adult control gonads, a Day 1 Adult PG(-) gonad, and a Day 2 Adult PG(-) UNC-119::GFP(+)

gonad. **B)** smFISH analysis of *aqp-2* transcripts (red) in a Day 2 Adult control gonad and a Day 2

Adult PG(-) UNC-119::GFP(+) gonad. For each gonad in (A) and (B), UNC-119::GFP signal is in

green and DAPI-stained DNA is in blue. The distal half of each gonad is shown as a z-stack

projection, with the distal tip indicated by an asterisk. Boxed regions are expanded in the right-most

panels. The zoomed-in regions are projections of 3 slices taken from the middle section of the

gonad.

Figure 5. Day 2 Adult PG(-) UNC-119::GFP(+) gonads up-regulate genes involved in neuronal

processes and down-regulate genes involved in germline processes. **A)** Volcano plot comparing

Day 2 Adult PG(-) UNC-119::GFP(+) gonads to Day 2 Adult PG(-) UNC-119::GFP(-) gonads.

Dashed lines indicate the significance cut-off of q=0.05 (horizontal line) and a 2-fold change in

transcript level (vertical lines). The numbers of significantly misregulated genes in each quadrant

are shown at the top of the plot. **B)** Gene Ontology analysis of the 2,268 up-regulated genes and

692 down-regulated genes in Day 2 Adult PG(-) UNC-119::GFP(+) gonads compared to Day 2

Adult PG(-) UNC-119::GFP(-) gonads. The Gene Ontology processes are organized based on their

p-values, and the numbers of genes in each ontology category are shown in parentheses. **C)**

Gonads from Day 2 Adult control and Day 2 Adult PG(-) UNC-119:GFP(+) worms were dissected

and immunostained for UNC-64 (red), GFP (green), and DNA (blue). The distal half of each gonad

is shown as a z-stack projection, with the distal tip indicated by an asterisk. Boxed regions are

expanded in the right-most panels. The zoomed-in regions are projections of 3 slices taken from

the middle section of the gonad.

Figure 6. Acquisition of neuronal fate in P-granule-depleted germ cells is accompanied by loss of

germ cell markers. **A)** Gonads from Day 2 Adult control and Day 2 Adult PG(-) UNC-119::GFP(+)

worms were dissected and immunostained for REC-8 (red), GFP (green), and DNA (blue). **B)**

Quantification of UNC-119::GFP signal relative to REC-8 signal in germ cells from control and P-

granule-depleted gonads. **C)** Same as in (A) except gonads were stained for HTP-3 (red). **D)**

Same as in (B) except gonads were quantified for UNC-119::GFP signal relative to HTP-3 signal.

E) Single molecule RNA-FISH (smFISH) analysis of *lag-1* transcripts (red) in a dissected Day 2

Adult control gonad and a Day 2 Adult P granule(-), UNC-119::GFP(+) gonad. For (A), (C), and (E),

the distal half of each gonad is shown as a z-stack projection, with the distal tip indicated by an

asterisk. Boxed regions are expanded in the right-most panels. The zoomed-in regions are

projections of 3 slices taken from the middle section of the gonad, except for the control HTP-3

gonad (C), which shows the top section of the gonad. In the expanded panels in (C), the red

channel of the PG(-) gonad was increased to better show relative HTP-3 levels.

Figure 7. Germlines lacking PGL-1 or GLH-1 also misexpress somatic markers. **A)** Percent of

germlines expressing the neuronal marker UNC-119::GFP (green bars) in *glh-1* single mutants, *glh-*

1 glh-4 double mutants, *pgl-1* single RNAi, and *pgl-1; pgl-3* double RNAi animals containing the

unc-119::gfp transgene. Of the germlines that expressed UNC-119::GFP, a fraction also expressed

the muscle marker MYO-3 (green and red striped bars). **B)** Volcano plots comparing *glh-1* (gold) or

pgl-1 (blue) single mutant gonads to wild-type gonads. All animals lacked the *unc-119::gfp*

transgene. Dashed lines indicate the significance cut-off of q=0.05 (horizontal line) and a 2-fold

change in transcript levels (vertical lines). The numbers of significantly misregulated genes in each

quadrant are shown at the top of the plot. **C)** Venn diagrams showing the overlap of genes down-

or up-regulated in *glh-1* and *pgl-1* single mutant gonads.

Figure 8. Models of P-granule function. P granules reside over nuclear pores, where they intercept

stochastically misexpressed soma-specific mRNAs (red mRNAs), while allowing germline-

appropriate mRNAs (green mRNAs) to transit to the cytoplasm for translation. In Model A, P

granules prevent the translation of somatic mRNAs. In Model B, P granules slice somatic mRNAs

into small RNAs, which in turn induce heritable silencing of somatic genes. In the absence of P

granules (right panel), somatic mRNAs accumulate and are translated into proteins (red) that

specify somatic identity and antagonize germline identity.

Figure 1

A

Day 0 — Place *unc-119::gfp* L1 larvae (**P₀ parents**) on control RNAi or P-granule RNAi

Day 3 — Transfer L1/L2 larvae (**F₁ progeny**) to new RNAi plates

Day 4 — Dissect gonads from L4 larvae

Day 5 — Dissect gonads from Day 1 adults

Day 6 — Dissect gonads from Day 2 adults

B

L4 larval gonads	Day 1 Adult gonads	Day 2 Adult gonads
control	control	control
	Fertile	Fertile
PG(-) UNC-119::GFP(-)	PG(-) UNC-119::GFP(-)	PG(-) UNC-119::GFP(-)
	Sterile	Sterile
		PG(-) UNC-119::GFP(+)
		Sterile

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8