Soft sweeps are the dominant mode of adaptation in the human genome

Daniel R. Schrider[*,†,1] and Andrew D. Kern[*,†]

[*]Department of Genetics, Rutgers University, Piscataway, NJ, 08854, USA

[†]Human Genetics Institute of New Jersey, Rutgers University, Piscataway, NJ, 08554, USA

[1]Corresponding author: Department of Genetics, Rutgers University, 604 Allison Rd.,

Piscataway, NJ 08854. E-mail: dan.schrider@rutgers.edu

Keywords: adaptation, selective sweeps, soft sweeps, machine learning, population genomics

Running title: Adaptation via soft sweeps is prevalent in humans

ABSTRACT

The degree to which adaptation in recent human evolution shapes genetic variation remains controversial. This is in part due to the limited evidence in humans for classic "hard selective sweeps," wherein a novel beneficial mutation rapidly sweeps through a population to fixation. However, positive selection may often proceed via "soft sweeps" acting on mutations already present within a population. Here we examine recent positive selection across six human populations using a powerful machine learning approach that is sensitive to both hard and soft sweeps. We found evidence that soft sweeps are widespread and account for the vast majority of recent human adaptation. Surprisingly, our results also suggest that linked positive selection affects patterns of variation across much of the genome, and may increase the frequencies of deleterious mutations. Our results also reveal insights into the role of sexual selection, cancer risk, and central nervous system development in recent human evolution.

INTRODUCTION

Spurred by the ongoing revolution in DNA sequencing capacity, human population genetic datasets have grown exponentially in size over the past five years (Auton et al. 2015; Consortium 2015). Such growth enables insight into the evolutionary histories of human populations with hitherto unrivaled precision. A central question in the study of human evolution is the extent to which adaptation has driven recent evolution and affected patterns of genetic diversity (Akey 2009). This can be addressed by scanning genomic data for evidence of selective sweeps, wherein a beneficial mutation is favored by natural selection and therefore rapidly increases in frequency within a population. Such selective sweeps leave a characteristic footprint in variation; they create a valley of diversity around the selected site (Maynard Smith and Haigh 1974;

Kaplan et al. 1989; Stephan et al. 1992), a deficit of both low- and high-frequency derived alleles

at linked sites (Fay and Wu 2000), and an increase in linkage disequilibrium in flanking regions

(Kim and Nielsen 2004). Thus there are multiple population genetic signals to exploit.

Accordingly numerous theoretical and methodological advances (Kaplan et al. 1989; Stephan et

al. 1992; Fu 1997; Kim and Stephan 2002; Nielsen et al. 2005b; Voight et al. 2006) in the study

of selective sweeps have given researchers the ability to uncover the genetic basis of adaptation

on a genome-wide scale.

43   There are two complimentary approaches to studying the impact of adaptive evolution on

genetic variation. The first approach aims to infer genome-wide rates of adaptive evolution by

estimating the mean effects of selective sweeps across the genome (Wiehe and Stephan 1993;

Kern et al. 2002; Andolfatto 2007; Jensen et al. 2008; Hernandez et al. 2011; Sattath et al. 2011).

Such approaches may estimate the rates of sweeps or their effects with respect to the genomic

background, but do not focus on the targets of sweeps themselves. An alternative approach is to

focus on finding individual selective sweeps throughout the genome, and in so doing characterize

specific cases of the adaptation with hopes of gaining general insight into the adaptive process

(Sabeti et al. 2002; Voight et al. 2006; Williamson et al. 2007). Such approaches have yielded

valuable insights into the recent evolutionary histories of natural populations in various species,

such as revealing the pervasive impact of adaptive evolution on polymorphism in *Drosophila*

melanogaster (Begun et al. 2007; Macpherson et al. 2007; Langley et al. 2012; Lee et al. 2013;

Garud et al. 2015). In humans, the picture remains less clear: while scans for selective sweeps

have discovered numerous compelling candidates for strong positive selection (e.g. Ruwende et

al. 1995; Stephens et al. 1998; Tishkoff et al. 2007; Bryk et al. 2008; Huerta-Sánchez et al.

2014), some recent studies have suggested that the impact of adaptation on patterns of variation

genome-wide is quite limited (Hernandez et al. 2011; Lohmueller et al. 2011). Conversely, Enard

et al. (2014) argue that the genome-wide reduction in diversity around substitutions is driven in

part by positive selection.

One possible explanation for the difficulty in characterizing the contributions of adaptive

and non-adaptive forces in human populations is that genetic hitchhiking effects may be muted

by human demographic history. Many human populations appear to have experienced

bottlenecks and/or recent growth (Marth et al. 2004; Fagundes et al. 2007; Gravel et al. 2011;

Auton et al. 2015), which cause much of the genome to resemble selective sweeps (Nielsen et al.

2005b). Moreover, positive selection has historically been modeled as the process of a *de novo*

beneficial mutation rapidly sweeping to fixation, a process now referred to as a hard sweep.

However selection may act on previously segregating neutral or weakly deleterious variants (Orr

and Betancourt 2001; Innan and Kim 2004). Selection on standing variation will produce

qualitatively different skews in linkage disequilibrium and allele frequencies, along with a

shallower valley in diversity (Hermisson and Pennings 2005; Przeworski et al. 2005; Berg and

Coop 2015; Schrider et al. 2015)—such an event is thus referred to as a soft sweep. If selection

typically proceeds through soft sweeps, as may be the case in Drosophila (Garud et al. 2015),

then many sweeps may have been missed by previous scans that were designed to detect

signatures produced under a hard sweep model.

We sought to address the controversy over the impact of adaptation on human genomic

variation by conducting a genome-wide scan for both hard and soft selective sweeps across

human populations. We previously developed S/HIC (Soft/Hard Inference through

Classification), a machine learning method capable of detecting completed sweeps and inferring

their mode of selection with unparalleled accuracy and robustness to non-equilibrium

demography (Schrider and Kern 2016). Here we apply S/HIC to uncover hard and soft sweeps in

six population samples from the 1000 Genomes Project (Auton et al. 2015), thereby performing

the most comprehensive investigation of completed selective sweeps in humans to date.

Surprisingly, our results suggest that patterns of polymorphism across much of the human

genome may be affected by linked positive selection—primarily soft sweeps. Moreover, we find

evidence that the mode of selection differs substantially across populations, with non-African

populations adapting via hard sweeps to a much greater extent than African populations. Finally,

we investigate the biological targets of selection in recent human evolution, with particular

processes such as immunity, cancer, and sexual reproduction playing outsized roles.

RESULTS

We set out to detect completed hard and soft selective sweeps in six populations from Phase 3 of

the 1000 Genomes Project: two West-African populations (YRI and GWD from Yoruba and The

Gambia, respectively), one East-African population (LWK from Kenya), one European

population (CEU, from Utah, USA), one East Asian population (JPT from Japan), and one from

the Americas (PEL from Peru). For each population we trained and applied a S/HIC classifier to

identify hard and soft selective sweeps across the genome (Methods), distinguishing them from

neutrally evolving regions as well as those linked to sweeps (Schrider and Kern 2016). Briefly,

S/HIC is a machine learning method that leverages spatial patterns of a variety of statistics across

a large genomic window in order to infer the mode of evolution at the center of the window. We

previously showed that S/HIC is exceptionally robust to the confounding effect of linked

selection (i.e. the "soft shoulder" effect; Schrider et al. 2015), as well as non-equilibrium

demographic histories, making it well suited for a survey of positive selection in humans. We

also assessed the accuracy of our classifiers on simulated test data with the same demographic

history used to generate training data, finding that S/HIC achieved good power for each

demographic history, with somewhat higher accuracy for histories inferred from the African than

non-African populations (Figure S1). We also examined values of Garud et al.'s (2015) H_{12} and

H_2/H_1 within windows classified by S/HIC as hard, soft, or neutral, noting that as expected, H_{12}

is higher in sweeps than neutral regions, while H_2/H_1 is higher for soft sweeps than hard sweeps

(Figure S2). Below, we begin with a brief overview of the broad patterns of adaptation we

observe across populations, before discussing genomic features and biological pathways with a

strong enrichment of selective sweeps, as well as compelling novel candidates for recently

completed selective sweeps.

The majority of sweeps in humans resemble selection on standing variation

We found a total of 1,927 distinct selective sweeps merged across all six populations (Methods).

190 (9.9%) of these are present in all populations, 59 (3.1%) are shared among the African

populations, 71 (3.7%) are shared among the non-African populations, and 701 (36.4%) are

population-specific. The remaining 906 (47.0%) sweeps were present in more than one

population but do not fit into any of the categories above. We observe that across populations,

the vast majority (1,776, or 92.2%) of sweeps were classified as soft. While hard sweeps appear

to be quite rare globally, the fraction of hard sweeps is significantly higher in non-African than

African populations (Table 1). For example, when comparing PEL to GWD, we observe a

significantly higher fraction of hard sweeps in PEL (4.7% versus 1.6%; p=0.05). For each other

African vs. non-African comparison we see an even greater (and more significant) disparity.

Further, we observe a suggestive correlation between the fraction sweeps in a population that

were classified as soft and the harmonic mean of its population size within the last $4N$

generations (Pearson's ρ=-0.96; Methods). Though taken at face value this correlation appears to

be highly significant, we note that due to the six populations' shared evolutionary history a

statistical test of this correlation would be invalid.

Comparing our results to those of previous scans we find that 519 of S/HIC's sweep calls

(26.9%) have previously been identified according to dbPSHP, a database of candidate regions

for recent positive selection across human populations (Li et al. 2013). This accounts for 10.9%

of the loci in the dbPSHP set (ignoring regions not classified by S/HIC). The remaining 1,408

sweeps called by S/HIC (73.1% of calls) represent potentially novel selective sweeps. There are

several possible explanations for the modest overlap between our set of sweep candidates and

those in dbPSHP. First, the sweep candidates in dbPSHP have been identified by a variety of

methods, some of which are designed to detect selective scenarios other than completed sweeps

(e.g. partial sweeps, spatially varying selection). Second, when comparing results from methods

designed to detect the same type of sweeps, the intersection between studies is often fairly small

(Akey 2009). The presence of false positives may in part be responsible for the imperfect

concordance between scans.

145   Selective sweeps preferentially target genes involved in cancer and viral infection

Examining the locations of selective sweeps across the genome, we find that regions classified as

selective sweeps are significantly overrepresented for both coding sequence and untranslated

regions ($q<0.05$ in several populations for hard sweeps, and each population for soft sweeps;

Figure 1A, B; Table S1), relative to data sets with permuted classifications (see Methods).

Enrichment for transcription factor binding sites was less pronounced, and only significant in

soft sweeps for the three African populations along with PEL. The most striking result we

observed was a dramatic enrichment of sweep windows for mutations in the COSMIC data set of

somatic mutations that have been observed in cancer cells (Forbes et al. 2015) and may therefore

play a role in tumor suppression/progression. Averaged across populations, the number of

COSMIC mutations found in soft sweeps represents a 3.7-fold increase relative to that observed

in permuted data sets; this enrichment was significant in each population, and peaked at 4.5-fold

in PEL. For hard sweeps, this enrichment was 12-fold on average, reaching as high as 21-fold in

CEU, though this was the only population for which the enrichment was statistically significant.

We also observed a sizeable overrepresentation of genes encoding virus-interacting proteins

(VIPs) curated by (Enard et al. 2016) in soft sweeps, with a 1.9-fold increase relative to

permuted sets (averaged across populations). VIPs show a similar magnitude of enrichment in

hard sweeps for some populations, but does not achieve significance at $q<0.05$.

Selective sweeps increase linked deleterious variation

Because S/HIC not only detects selective sweeps, but also attempts to identify regions of the

genome that appear to be linked to recent sweeps, our classifications allow us to examine the

effect of linked selection in a principled way. We found that while a minority of genomic

windows were classified as selective sweeps (7.6% on average across all populations), a large

fraction of windows were classified as linked to a completed selected sweep, either hard or soft

(56.4% on average). These estimates range from 41.5% in JPT to 74.0% in GWD (Figure 2).

We also asked whether selective sweeps have a detectable impact on linked deleterious

variation. As beneficial alleles increase in frequency in a population, they may carry along with

them linked deleterious polymorphisms as hitchhikers, potentially increasing the frequency of

deleterious variants over what would be expected given mutation-selection-drift equilibrium

(Birky and Walsh 1988; Hartfield and Otto 2011). To this end we asked whether relatively

common candidate deleterious mutations were enriched in regions classified as either hard-

linked or soft-linked. Indeed, we observed a fairly subtle but significant overrepresentation of

SNPs with derived allele frequencies of at least 0.01 but predicted to by damaging by SIFT

(Kumar et al. 2009) in both the hard-linked (mean enrichment across populations: 1.3-fold) and

soft-linked (mean enrichment: 1.1-fold) classes for most populations (Fig 1C, D; Table S1). We

find a similar enrichment in these sweep-linked classes of common SNPs in regions inferred to

be conserved across primates according to phastCons (Siepel et al. 2005). Phenotype-associated

variants from the GWAS catalogue (Welter et al. 2014) were also significantly overrepresented

sweep-linked regions in several populations (Fig 1C, D).

Sexual reproduction, the central nervous system, and immunity are targets of recent

sweeps

In order to determine if positive selection preferentially acts on particular organismal functions,

we asked which Gene Ontology (GO) terms were enriched in our sweep calls relative to the

permuted data (Methods). In soft sweeps, we found a sizeable and significant enrichment

($q<0.05$) of terms related to sperm development, structure, and function. Terms enriched in at

least one population include spermatogenesis (4.4-fold enrichment averaged across populations),

"sperm flagellum" (3.9-fold enrichment), and "sperm-egg recognition" (3.3-fold enrichment).

These terms were not significantly enriched for hard sweeps in any population, though this may

be due in part to reduced power because of the smaller number of sweeps.

We also observed an overrepresentation of genes involved in the "glutamate receptor

signaling pathway" in our soft sweep sets ($q<0.05$ for each population; 4-fold mean enrichment).

Glutamate receptors are the primary excitatory neurotransmitter in the central nervous system,

and important for both proper brain development and function (Luján et al. 2005). Indeed, soft

sweeps are enriched for "central nervous system development" in multiple populations (1.6-fold

mean enrichment). Numerous terms related to immune response, especially adaptive immunity,

were also significantly enriched among soft sweeps. These, and the many other GO terms

overrepresented in our sweep sets are listed in Table S2. We also found numerous KEGG

pathways enriched for sweeps, including many pathways related to immunity, cancer

progression/tumor suppression (see Table S2 for full list).

Positive selection on interacting gene pairs

We examined three types of gene interaction networks: protein-protein interactions (PPIs),

transcription factor-target gene interactions, and genetic interactions where one gene modifies

the effect of another (Methods). Interestingly, we observed a dramatic enrichment of sweeps in

genes that encode proteins that physically interact with one another (Figure 3A–B): if a gene

experienced a soft sweep, genes that interact with this gene were on average 3.3 times more

likely to have experienced a soft sweep than expected by chance ($p<0.0001$ for each population;

Figure 3B). Despite the smaller number of candidate regions, we found a significant enrichment

for PPIs in hard sweeps, though this was only significant in for non-African populations (4.0-fold

enrichment averaged across populations; $p<0.05$ in CEU, JPT, YRI; Figure 3A). For

transcription factor-target interactions, we observe no overrepresentation of soft sweeps, but a

significant enrichment of hard sweeps in non-African populations ($p<0.05$ for each; 8.5-fold

enrichment on average; Figure 3C–D). There were no populations exhibiting an

overrepresentation of pairs of genes with genetic interactions and experiencing sweeps of either

type (Figure 3E–F).

Examples of novel selective sweep candidates

In this section we describe several sweep candidates that exemplify the set of sweeps, and

functions of putative targets of selection, that we were able to detect. As discussed above, our

sets of sweeps were highly enriched for glutamate receptor-encoding genes. In Figure S3, we

show a sweep candidate region on chromosome 4 that encompasses the glutamate receptor gene

GRIA2. This sweep was previously detected in non-African populations by Pickrell et al. (2009),

who did not find any evidence of selection in Africa. However, S/HIC infers that this region has

experienced a soft sweep that is found in GWD and YRI, as well as the non-African populations.

Consistent with this, Europeans, Asians, and African populations show a reduction in π, a trough

in Tajima's *D* (Tajima 1989), and a peak in Nielsen et al.'s SweepFinder composite likelihood

ratio (CLR) test statistic, which captures regions that appear to be at the epicenter of the spatial

skew in the SFS expected around sweeps (Nielsen et al. 2005b). Intriguingly, *GRIA2* interacts

with the *GRID2* glutamate receptor gene (Kohda et al. 2003), which itself is classified as a soft

sweep in CEU, LWK, PEL, and GWD. The remaining glutamate receptors overlapping identified

sweeps are *GRIA4*, *GRID1*, *GRIK1*, *GRIK3*, *GRM2*, and *GRM7*. Of these genes, *GRIA4* and

GRID2 were shown by Liu et al. (2012) to have evolved a human-specific developmental

expression profile.

Figure 4 shows a region on chromosome 9 that exhibits strong evidence of a previously

undetected hard sweep in each of our six populations. This region contains several members of

the spermatogenesis associated 31 gene family: *SPATA31B1*, *SPATA31D1*, *SPATA31D3*, and

SPATA31D4. Across populations this region shows dramatic valleys in π and Tajima's *D*, as well

as an elevated CLR near the center of the sweep window. These genes are highly testis-specific

according to data from the GTEx project (Lonsdale et al. 2013), and male mice are infertile when

lacking Spata31, another member of these gene family (Wu et al. 2015). Figure 4 also shows that

each of these genes overlaps a cluster of non-repetitive piRNAs (data from piRBase; Zhang et al.

2014). Also near this region is *DDX10P2*, which GENCODE annotates as a processed

pseudogene (Pei et al. 2012). *DDX10P2*, which is located at the center of the CLR peak for CEU,

is expressed with a high degree of testis-specificity according to GTEx data, similar to the

neighboring *SPATA31* genes. A BLAT search (Kent 2002) revealed that this putative

pseudogene exhibits 99.5% sequence identity to the orthologous sequence in chimpanzees. The

parent gene of *DDX10P2*, *DDX10*, is expressed in many tissues, but shows highest expression in

the testis.

On chromosome 11 we detected what appear to be several novel soft sweeps present in

and upstream of *CADM1* (cell adhesion molecule 1; Figure 5), one of which is present in each

population. This gene is essential for spermatogenesis in mice (Van Der Weyden et al. 2006),

and is also a tumor suppressor that is hypermethelated in various cancers (Kuramochi et al. 2001;

Allinen et al. 2002; Fukuhara et al. 2002), as it works with the adaptive immune system to

suppress metastasis (Faraji et al. 2012). *CADM1* is also active in the brain where it is involved in

synaptic adhesion and has been linked to autism (Zhiling et al. 2008; Fujita et al. 2010). *CADM1*

forms a complex with those other genes: the GABA receptor *GABBR2*, which has a soft sweep in

YRI, and *MUPP1*, which has a soft sweep found in each population; this complex appears to

localize to Purkinje cell dendrites (Fujita et al. 2012). Thus, this example encompasses many of

the functions that we find are highly enriched across our sweep sets: adaptation in multiple

interacting genes (one of which is a neurotransmitter), spermatogenesis, and tumor suppression

(via adaptive immunity).

DISCUSSION

Understanding the history of human adaptation at the genetic level is a central goal of population

genomics and human evolutionary biology. Accordingly, since the completion of the human

genome assembly (Lander et al. 2001) and subsequent proliferation of population genomic data,

numerous genome-wide scans for selection have been conducted using differing methodologies

(Sabeti et al. 2002; Voight et al. 2006; Sabeti et al. 2007; Pickrell et al. 2009; Field et al. 2016).

The majority of these studies searched primarily for partial selective sweeps—the signature of a

beneficial mutation currently sweeping through a population (see Williamson et al. 2007 for a

notable exception)—and rightly so, as these sweeps can reveal the targets of ongoing adaptation

in human populations. However, because the sojourn of an adaptive mutation to fixation should

be rapid, the success of efforts to detect ongoing selection implies the presence of a larger

number of recently completed sweeps. We have therefore focused on completed sweeps in order

to complement previous studies and to construct a more comprehensive catalogue of the loci

underpinning recent human adaptation. Using a powerful and robust machine learning method

that we have recently introduced (S/HIC; Schrider and Kern 2016) for finding completed

selective sweeps, we performed a genome-wide search for the targets of recent positive selection

in six human populations. Furthermore, we sought to determine the mode of positive selection,

distinguishing between selection on *de novo* mutations and on previously standing variation.

Soft sweeps dominate human adaptation

Perhaps our most consequential result is the finding that the majority of our candidate sweeps

resemble soft sweeps on standing variation. This result implies that adaptation in humans may

not be mutation-limited (Gillespie 1991; Karasov et al. 2010): rather than waiting for a novel

mutation to arise, human populations may often be able to respond via selection on previously

segregating polymorphisms, thereby more rapidly responding to novel environmental challenges.

This may be surprising given the apparently small effective population size and low nucleotide

diversity levels in humans. However, if the mutational target for the trait to be selected on is

fairly large, then the probability of a population harboring a mutation affecting that trait may be

appreciable.

While soft sweeps appear to be the dominant mode of selection globally, there is a

significant increase in the proportion of putative hard sweeps in non-African populations relative

to African populations. This is consistent with theoretical expectations, as larger populations

have more standing variation for selection to act on (Hermisson and Pennings 2005). Moreover,

the human migration out of Africa was associated with a severe population bottleneck (Marth et

al. 2004; Fagundes et al. 2007). Soft selective sweeps may be "hardened" by a reduction in

population size, which can result in the stochastic loss of some genetic backgrounds harboring

the adaptive allele so that only a single haplotype reaches fixation (Wilson et al. 2014). Thus,

though one might expect selection on segregating neutral or nearly neutral variation when a

population enters a new environment with novel selective pressures, if the migration event is

accompanied by a bottleneck then the population may experience a somewhat counterintuitive

increase in the proportion of hard sweeps. Moreover, the causal relationship between population

size and mode of adaptation may not be unidirectional. As Orr and Unckless (2014) have shown

in the context of evolutionary rescue, when faced with a changing environment, a population

which does not harbor standing variation that is beneficial may experience a more protracted

decline in size while it waits for an adaptive *de novo* mutation.

Our genome-wide results amplify results of earlier studies that by design have tried to

infer the mode of adaptation in a smaller number of targeted loci. For instance Peter *et al.* (2012)

attempted to infer the mode of adaptation among 7 loci previously identified to be under

selection in human populations. They report that half of the loci that they could confidently

classify supported selection on standing variation. In *Drosophila melanogaster,* when looking

among strong outliers of haplotype homozgosity, Garud *et al.* (2015) found that patterns of

variation in those regions were consistent with recent soft selective sweeps. Our finding, that the

vast majority of sweeps in human populations are soft sweeps, thus underscores the ubiquity of

selection from standing variation in natural populations. Indeed it seems plausible that adaptation

from standing variation might be the rule, rather than the exception.

There are two caveats affecting our ability to discriminate between selection on standing

variation and on *de novo* mutations. First, while we have trained our classifier to detect soft

sweeps on previously segregating mutations, soft sweeps may also occur via recurrent mutation

to the adaptive allele (Pennings and Hermisson 2006b; Pennings and Hermisson 2006a). Though

there are some qualitative differences between these two models of soft sweeps (Berg and Coop

2015; Schrider et al. 2015), these are fairly subtle in comparison to the differences between the

other models we consider. Thus, our classifiers may have sensitivity to both types of sweeps. If

this is so, then some of the soft sweeps that we detect may result from recurrent mutation.

Additionally, gene conversion during a sweep can transfer the adaptive mutation on to new

genetic backgrounds (Jones and Wakeley 2008), thereby "softening" the sweep (Schrider et al.

2015). This implies that selection on a single de novo mutation could sometimes appear to be a

soft sweep in our classification. In any case, our finding that most sweeps in humans do not

appear to be hard sweeps underscores the importance of using methods that are sensitive to soft

sweeps.

Extensive impact of linked positive selection

Our analysis demonstrates that the impact of linked positive selection on genetic variation is

considerable, with roughly half of the genome classified by S/HIC as being influenced by a

nearby sweep. This result has important implications for efforts to infer demographic histories

from patterns of genetic polymorphism, as most inference methods hinge on the assumption of

neutrality. Indeed, we have recently shown that linked positive selection has the potential to

severely confound demographic inferences (Schrider et al. 2016). Similarly, Ewing and Jensen

(2016) have found that background selection, the effect of purifying selection on polymorphism

at linked unselected sites (Charlesworth et al. 1993), can also bias demographic estimates. One

strategy is to use only those polymorphisms that are distant from genes and conserved noncoding

elements to mitigate these effects (Gazave et al. 2014). One could further supplement such an

approach by using S/HIC to directly ask which intergenic regions are unaffected by hitchhiking

in order to further diminish the bias introduced by linked selection.

If linked positive selection affects much of the genome, then that implies that the

frequencies of many neutral or weakly deleterious mutations may be altered by genetic draft

(Gillespie 2000). That is to say, deleterious mutations that happen to reside on chromosomes that

begin to sweep may be able to reach higher frequencies than expected from mutation-selection-

drift equilibrium. Consistent with this, we observe a slight but significant excess of potentially

deleterious polymorphisms in windows classified as linked to selective sweeps. Previously, Chun

and Fay (2011) found evidence that the ratio of deleterious to neutral polymorphisms is elevated

in sweep regions, concluding that hitchhiking carries linked deleterious variants to higher

frequencies. Our finding that SNPs from the GWAS catalogue are also enriched regions linked to

selective sweeps lends further support to this hypothesis. Indeed, several compelling examples of

hitchhiking mutations known or suspected of causing disease have been described in the

literature (Helgason et al. 2007; Chun and Fay 2011; Huff et al. 2012). It seems that the

phenomenon of deleterious alleles hitchhiking along with strongly beneficial alleles is not

restricted to humans: a recent study also uncovered evidence that selection during domestication

increased the frequency of deleterious polymorphisms in dogs (Marsden et al. 2016).

Targets of recent human selective sweeps

Our catalogue of sweep candidates allowed us to characterize the biological functions that are

overrepresented in sweeps. Notably, we found a strong excess of spermatogenesis genes within

sweep regions, a phenomenon previously observed by Voight et al. (2006). This signature may

be a result of sexual selection, sexual conflict, and/or sperm competition (Swanson and Vacquier

2002). We also observed a significant enrichment of cancer-related genes among our sweep

candidates. Nielsen et al. (2005a) found a similar enrichment of candidate genes under selection

related to cancer when examining protein divergence between humans and chimpanzees. These

authors found that some of these genes are also involved in spermatogenesis (much like our

CADM1 example), and concluded that genomic conflict between tumor suppression and the

advantage of avoiding apoptosis during spermatogenesis may explain the selection on cancer

genes. An alternative (and non-mutually exclusive) explanation is that the increase in longevity

along the human lineage has created an immense selective pressure to reduce the rate of cancer

progression by orders of magnitude (Nunney and Muir 2015).

We also observed a significant excess of glutamate receptor genes targeted by sweeps,

suggesting that these loci may underlie some of the dramatic neurological changes that have

occurred along the human lineage. Consistent with this, we previously found evidence

suggesting some of these glutamate receptor genes may have recently gained novel regulatory

elements in humans (Schrider and Kern 2015). The most striking examples of glutamate

receptors experiencing sweeps are *GRIA2* and *GRID2*, which show strong signatures of selection

in multiple populations and physically interact with one another. The action of positive selection

on multiple members of the protein complex appears to be a general phenomenon (Figure 3). For

a more in-depth examination of positive selection in the PPI network, see Qian et al. (2015), who

found that genes in candidate regions for positive selection were more likely to lie close together

in the PPI network.

Conclusions

Our investigation has revealed several valuable insights into the adaptive process in human

populations. The success of our approach exemplifies the potential of machine learning methods

to elucidate the adaptive process in humans and other species (Fan et al. 2016). To date several

machine learning methods have been devised to detect selective sweeps (Pavlidis et al. 2010; Lin

et al. 2011; Ronen et al. 2013; Pybus et al. 2015; Sheehan and Song 2016), and they tend to

substantially outperform more traditional approaches (see Schrider and Kern 2016). We suspect

that machine learning could be used to make important inroads in answering a variety of

evolutionary questions.

Finally, Hernandez et al. (2011) argued that hard selective sweeps might be rare in human

populations, and instead suggested that the majority of adaptation might be a consequence of

selection on standing variation or selection on polygenic traits. We here find direct evidence that

indeed this is the case—the vast bulk of human adaptation is occurring as a consequence of soft

sweeps. Our observation thus reconciles Hernandez et al.'s findings with those of Enard et al.,

who conclude that the reduction in diversity around amino acid substitutions is caused by

widespread selective sweeps (Enard et al. 2014). Moreover, while our scan leveraged a method

that performs very well in detecting both hard and soft sweeps, it was not trained to detect cases

of polygenic selection (e.g. Berg and Coop 2014). It is fair to assume that a large majority of

phenotypes are determined by multiple loci, thus polygenic selection should be expected to be

common. If that were the case, then it could very well be that an even larger portion of genetic

variation is influenced by natural selection and its linked effects throughout the genome.

METHODS

Sequence and annotation data

We downloaded phased genotype data from Phase 3 of the 1000 Genomes Project (Auton et al.

2015). This data set consists of 26 population samples from Africa, East Asia, South Asia,

Europe, and the Americas. We wished to include only populations where the influence of

admixture/migration on genetic variation appeared to be minimal, while still allowing us to

characterize selection across multiple continents. We therefore chose to scan the following

populations for selective sweeps: the GWD (Gambians in Western Divisions in The Gambia) and

YRI (Yoruba in Ibadan, Nigeria) populations from West Africa, LWK (Luhya in Webuye,

Kenya) from East Africa, JPT (Japanese in Tokyo, Japan) from Asia, CEU (Utah residents with

Northern and Western European Ancestry) from Europe, and PEL (Peruvians from Lima, Peru)

from the Americas. Examining Auton et al.'s results from running ADMIXTURE (Alexander et

al. 2009), we see that for most values of K, each of these populations appears to correspond

primarily to a single ancestral population rather than displaying multiple clusters of ancestry (see

Extended Data Figure 5 from Auton et al. 2015). One exception may be the PEL population, but

among the highly admixed American samples it appears to exhibit the smallest amount of

possible mixed ancestry (for most values of K), so we retained this population in order to have

some representation from the Americas. We opted not to examine any South Asian population,

as for each of these samples ADMIXTURE inferred evidence of ancestry from three or more

ancestral populations.

We downloaded numerous annotation data sets containing genomic features to test for

enrichment/depletion of selective sweeps and perform other downstream analyses. These

included GENCODE gene model release 19 (Harrow et al. 2012) including pseudogenes (Pei et

al. 2012), virus-interacting proteins from Enard et al. (2016), enhancers gained or along the

human lineage since diverging from Old World monkeys (Cotney et al. 2013), and SIFT's

(Kumar et al. 2009) predictions of damaging amino acid polymorphisms from dbNSFP version

3.2a (Liu et al. 2016). We obtained Gene Ontology (GO) annotations from ENSEMBL release

75 (Yates et al. 2016). We also downloaded coordinates of previously identified selective sweeps

from dbPSHP (Li et al. 2013).

We used the UCSC Table Browser (Karolchik et al. 2004) to obtain the following data

sets: phenotype-associated SNPs from the GWAS Catalog (accessed Apr 12, 2016; Welter et al.

2014), ClinVar pathogenic SNPs and indels ≤ 20 bp in length (Apr 26, 2016; Landrum et al.

2016), COSMIC somatic mutations in cancer (accessed Feb 25, 2014; Forbes et al. 2015),

phastCons elements conserved across primates (accessed Jun 2, 2013; Siepel et al. 2005),

ENCODE transcription factor binding sites version 3 (accessed Aug 25, 2013; Dunham et al.

2012), tables of genes and SNPs implicated in Mendelian phenotypes from OMIM (accessed

May 2, 2016; Amberger et al. 2015), and KEGG pathway annotations (accessed Apr 27, 2016;

Kanehisa et al. 2015). For each of these data sets we used GRCh37/hg19 coordinates.

In order to examine the prevalence of selective sweeps within interacting gene networks,

we downloaded physical and genetic interactions from BioGRID version 3.4.136 (Chatr-

Aryamontri et al. 2015). Our set of genetic interactions consisted of those annotated as "synthetic

genetic interaction defined by inequality," "suppressive genetic interaction defined by

inequality," or "additive genetic interaction defined by inequality." Physical interactions

included those annotated as "direct interaction," "association," or "physical association." We

extracted transcription factor-target interactions from ORegAnno (accessed Dec 22, 2015;

Griffith et al. 2008), retaining only interacting pairs where the ENSEMBL gene identifier were

provided for both genes in order to avoid ambiguity.

Building classifiers to detect selective sweeps

To detect sweeps we used S/HIC (https://github.com/kern-lab/shIC), a machine learning

approach we previously described and showed to be remarkably powerful and robust to non-

equilibrium demography (Schrider and Kern 2016). Briefly, the S/HIC machine learning

approach leverages spatial patterns (along a genome) of a variety of population genetic summary

statistics to classify genomic windows as being the target of a completed hard sweep (hard),

being closely linked to a hard sweep (hard-linked), a completed soft sweep (soft), linked to a soft

sweep (soft-linked), or evolving neutrally (neutral). While this classification approach allows

inference when considering a large number of features jointly, it necessitates training from a

large number of data instances known to belong to each class. Because the number of genomic

windows known to belong to each our five classes is limited, we must rely on simulation to

generate our training data. To this end we used the program discoal (Kern and Schrider 2016) to

simulate large chromosomal regions, subdivided into 11 sub-windows. Training examples for the

hard class experienced a hard sweep in the center of the central sub-window (i.e. the 6[th]

window), while examples for the hard-linked class experienced a hard sweep in the center of one

of the remaining sub-windows (selected randomly). Analogous simulations with soft sweeps

were generated for the soft and soft-linked classes, respectively. Finally, neutrally evolving

examples did not experience any selective sweep.

We sought to train a classifier for each population under a demographic model that offers

a better approximation to the population size history than the standard neutral model. For this we

used Auton et al.'s (2015) population histories inferred by PSMC (Li and Durbin 2011). The

1000 Genomes Project's PSMC output did not contain estimates of θ, the population mutation

rate parameter. Thus for each population we conducted a grid search by simulating genomic

windows with the appropriate sample size under each demographic model with varying values of

$\theta=4NuL$ (where L is the length of the locus, which we set to 100 kb); the grid of θ values raged

from 10 to 250, examining multiples of 10. For each value of θ, we compared the values of π

(Nei and Li 1979), $\hat{\theta}_w$ (Watterson 1975), $\hat{\theta}_H$ (Fay and Wu 2000), H_2/H_1 (Garud et al. 2015), and

Z_{nS} (Kelly 1997) from 1000 simulations to those from 1000 randomly selected genomic loci

(calculated as described below), calculating the mean of each statistic in the real and simulated

datasets. We chose as the final values of θ that for which the sum of the percent deviations of the

simulated from the observed means of each statistic was minimized. This estimate of θ allowed

us to calculate estimated population sizes and times scaled by the number generations for each

time point in the history inferred by PSMC. The harmonic mean of each population's size was

calculated by taking the estimated population size for each of the last $4N$ generations. We note

that these models may not accurately capture the demographic histories of the populations we

examined due to the confounding effects of positive (Schrider et al. 2016) and negative (Ewing

and Jensen 2016) selection. However, because of S/HIC's robustness to demographic

misspecification, we do not expect this to severely impact our analysis (Schrider and Kern 2016).

For each population we simulated a total of 2000 regions for each of our five classes. For

simulations involving sweeps, we drew the selection coefficient from $U(0.005, 0.1)$, the sweep

completion time from $U(0, 2000)$, the initial selected frequency for soft sweeps from $U(1/N, 0.2)$.

We drew values of θ uniformly from a range spanning exactly one order of magnitude, specified

so that the mean value of θ was equal to that estimated for the population as described above. We

drew recombination rates from an exponential distribution with mean 1×10^{-8}, truncated at triple

the mean due to memory constraints. The simulation program discoal requires some of these

parameters to be scaled by the present-day effective population size; we did this by taking the

mean value of θ and dividing by $4uL$, where u was set to 1.2×10^{-8} (Kong et al. 2012). The full

command lines we used to generate 1.1 Mb regions (to be subdivided into 11 windows each 100

kb in length) for each population are shown in Table S3. We also simulated 1000 test examples

for each population in the same manner as for the training data.

Our feature vector for each simulated region examined the spatial patters (following

Schrider and Kern 2016) of each of the following statistics: π (Nei and Li 1979), $\hat{\theta}_w$ (Watterson

1975), $\hat{\theta}_H$ (Fay and Wu 2000), the number of distinct haplotypes, average haplotype

homozygosity, Garud et al.'s (2015) H_{12} and H_2/H_1 statistics, Z_{nS}, ω (Kim and Nielsen 2004), and

the maximum frequency of derived mutations (Li 2011). Before calculating these summary

statistics we masked a number of sites within each simulation by randomly selecting a 1.1 Mb

region from our empirical windows sampled throughout the genome and masking the same

regions in the simulated window as were masked in the genomic window (see below). Thus our

simulated windows exhibit the same distribution of regions of missing data as the windows to

which we applied our classifiers. We then used S/HIC to train extra-trees classifiers (Geurts et al.

2006), one for each population.

Classifying genomic windows in each population

Having trained our classifiers, we then applied them to genomic data from the corresponding

population. We inferred ancestral states of polymorphisms and masked inaccessible sites

(whether polymorphic or not) in the same manner as described previously (Schrider and Kern

2016). We then used S/HIC to classify the central 100 kb sub-window of 1.1 Mb windows across

the autosomes, while taking the stringent approach of omitting those for which any sub-window

was less than 25% accessible, before sliding 100 kb downstream to examine the next window.

We also removed windows where any of the three central sub-windows had a mean

recombination rate of zero (using data from Kong et al. 2010). Importantly, for each retained 1.1

Mb window, we recorded the locations of all sites deemed inaccessible for use in masking our

training data (see above). In total we classified 13,968 windows, accounting for 48.5% of the

assembled autosomes.

In order to count the number of distinct sweep candidates found within our set of

populations , we simply merged all 100 kb windows classified as a sweep of either type that were

located either at the exact same coordinates or adjacent to one another, repeating this until no

more sweep regions could be merged. If all constituent windows were classified as soft, we

counted the sweep as soft; otherwise we counted it as a hard sweep. We used a similar approach

but examining classifications from only one population at a time in order to count the number of

sweeps of each type in that population. If a gene found within a sweep window identified by

S/HIC was not found in an entry of dbPSHP (Li et al. 2013), we referred to it as a novel sweep.

Visualization of sweep candidates was performed using the UCSC Genome Browser (Kent et al.

2002) , along with custom tracks showing values of various population genetic summary

statistics and selection scan scores for the CEU, YRI, and JPT populations from the Human

Positive Selection Browser (Pybus et al. 2013). Our classification results are available at

http://kerndev.rutgers.edu/~dan/shicHumanScan/.

Permutation tests for enrichment of annotation features in sweeps

To determine whether certain annotation features were enriched within any of our five classes,

we carefully designed a permutation test to account for the subset of the genome that we

examined with S/HIC, as well as the spatial correlation of S/HIC's classifications (i.e. adjacent

windows are especially likely to receive the same classification). Briefly, the permutation

algorithm begins by examining our classification results for a given population and keeping track

of the length of runs of consecutive windows assigned to each class. The permutation algorithm

then selects a chromosome, and begins at its first classified window (i.e. not removed by data

filtering). A run length and associated class assignment is then randomly drawn without

replacement. This process continues until the end of the chromosome, and then another

chromosome is selected until the end of the final chromosome is reached, at which point the

permutation has been completed. We then repeated this permutation procedure 10,000 times for

each population. Note that this process preserves the run length distribution of our classifications

while permuting them across the set of genomic windows that had enough unmasked data to be

included in our scan.

After constructing our permuted data sets, we conducted one-sided enrichment tests by

counting the number of base pairs in the intersection between the S/HIC class of interest and the

annotation feature of interest, and comparing this number to its distribution among the permuted

data sets. The fraction of permuted data sets where this intersect was greater than or equal to that

observed for the real data is the p-value. Because we tested each of S/HIC's five classes for

enrichment of a fairly large number of genomic features (Table S1), we corrected for multiple

testing using false discovery rate q-values following Storey (2002). When testing GO terms and

KEGG pathways for enrichment, we considered only the hard and soft sweep classes, corrected

for calculating q-values separately for each class.

We also asked whether the number of pairs of interacting genes both overlapping

windows classified as sweeps was greater than in our permuted data sets. To ensure that our

results were not inflated by the spatial clustering of interacting genes, we only counted

interacting pairs overlapping sweep windows if they were separated by at least 10 Mb or on

separate chromosomes. In addition, if we observed an interaction between two genes, A and B,

that each overlapped sweeps, and a third sweep candidate gene, C, was found, to avoid

redundancy we counted at most one interaction between A and C and B and C, even if C was

found interact with both other genes. As with GO and KEGG terms, we only searched the hard

and soft classes for enrichments before calculating one-sided q-values as described above.

ACKNOWLEDGMENTS

We thank Matthew Hahn for feedback on the manuscript. A.D.K. was supported in part by NIH

award no. R01GM078204.

DISCLOSURE DECLARATION

591   The authors have no conflicts of interest to declare.

REFERENCES

Akey JM. 2009. Constructing genomic maps of positive selection in humans: Where do we go from here? *Genome Res* **19**: 711-722.

Alexander DH, Novembre J, Lange K. 2009. Fast model-based estimation of ancestry in unrelated individuals. *Genome Res* **19**: 1655-1664.

Allinen M, Peri L, Kujala S, Lahti-Domenici J, Outila K, Karppinen SM, Launonen V, Winqvist R. 2002. Analysis of 11q21–24 loss of heterozygosity candidate target genes in breast cancer: indications of TSLC1 promoter hypermethylation. *Genes Chromosomes Cancer* **34**: 384-389.

Amberger JS, Bocchini CA, Schiettecatte F, Scott AF, Hamosh A. 2015. OMIM. org: Online Mendelian Inheritance in Man (OMIM®), an online catalog of human genes and genetic disorders. *Nucleic Acids Res* **43**: D789-D798.

Andolfatto P. 2007. Hitchhiking effects of recurrent beneficial amino acid substitutions in the Drosophila melanogaster genome. *Genome Res* **17**: 1755-1762.

Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, Korbel JO, Marchini JL, McCarthy S, McVean GA, Abecasis GR et al. 2015. A global reference for human genetic variation. *Nature* **526**: 68-74.

Begun DJ, Holloway AK, Stevens K, Hillier LW, Poh Y-P, Hahn MW, Nista PM, Jones CD, Kern AD, Dewey CN. 2007. Population genomics: whole-genome analysis of polymorphism and divergence in Drosophila simulans. *PLoS Biol* **5**: e310.

Berg JJ, Coop G. 2014. A population genetic signal of polygenic adaptation. *PLoS Genet* **10**: e1004412.

Berg JJ, Coop G. 2015. A Coalescent Model for a Sweep of a Unique Standing Variant. *Genetics* **201**: 707-725.

Birky CW, Walsh JB. 1988. Effects of linkage on rates of molecular evolution. *Proceedings of the National Academy of Sciences* **85**: 6414-6418.

Bryk J, Hardouin E, Pugach I, Hughes D, Strotmann R, Stoneking M, Myles S. 2008. Positive selection in East Asians for an EDAR allele that enhances NF-κB activation. *PLoS ONE* **3**: e2209.

Charlesworth B, Morgan M, Charlesworth D. 1993. The effect of deleterious mutations on neutral molecular variation. *Genetics* **134**: 1289-1303.

Chatr-Aryamontri A, Breitkreutz B-J, Oughtred R, Boucher L, Heinicke S, Chen D, Stark C, Breitkreutz A, Kolas N, O'Donnell L. 2015. The BioGRID interaction database: 2015 update. *Nucleic Acids Res* **43**: D470-D478.

Chun S, Fay JC. 2011. Evidence for hitchhiking of deleterious mutations within the human genome. *PLoS Genet* **7**: e1002240.

Consortium UK. 2015. The UK10K project identifies rare variants in health and disease. *Nature* **526**: 82-90.

Cotney J, Leng J, Yin J, Reilly SK, DeMare LE, Emera D, Ayoub AE, Rakic P, Noonan JP. 2013. The Evolution of Lineage-Specific Regulatory Activities in the Human Embryonic Limb. *Cell* **154**: 185-196.

Dunham I Kundaje A Aldred SF Collins PJ Davis C Doyle F Epstein CB Frietze S Harrow J Kaul R et al. 2012. An integrated encyclopedia of DNA elements in the human genome. *Nature* **489**: 57-74.

Enard D, Cai L, Gwennap C, Petrov DA. 2016. Viruses are a dominant driver of protein adaptation in mammals. *eLife* **5**: e12469.

Enard D, Messer PW, Petrov DA. 2014. Genome-wide signals of positive selection in human evolution. *Genome Res*.

Ewing GB, Jensen JD. 2016. The consequences of not accounting for background selection in demographic inference. *Mol Ecol* **25**: 135-141.

Fagundes NJ, Ray N, Beaumont M, Neuenschwander S, Salzano FM, Bonatto SL, Excoffier L. 2007. Statistical evaluation of alternative models of human evolution. *Proceedings of the National Academy of Sciences* **104**: 17614-17619.

Fan S, Hansen ME, Lo Y, Tishkoff SA. 2016. Going global by adapting local: A review of recent human adaptation. *Science* **354**: 54-59.

Faraji F, Pang Y, Walker RC, Borges RN, Yang L, Hunter KW. 2012. Cadm1 is a metastasis susceptibility gene that suppresses metastasis by modifying tumor interaction with the cell-mediated immunity. *PLoS Genet* **8**: e1002926.

Fay JC, Wu C-I. 2000. Hitchhiking under positive Darwinian selection. *Genetics* **155**: 1405-1413.

Field Y, Boyle EA, Telis N, Gao Z, Gaulton KJ, Golan D, Yengo L, Rocheleau G, Froguel P, McCarthy MI et al. 2016. Detection of human adaptation during the past 2000 years. *Science* doi:10.1126/science.aag0776.

Forbes SA, Beare D, Gunasekaran P, Leung K, Bindal N, Boutselakis H, Ding M, Bamford S, Cole C, Ward S. 2015. COSMIC: exploring the world's knowledge of somatic mutations in human cancer. *Nucleic Acids Res* **43**: D805-D811.

Fu Y-X. 1997. Statistical tests of neutrality of mutations against population growth, hitchhiking and background selection. *Genetics* **147**: 915-925.

Fujita E, Dai H, Tanabe Y, Zhiling Y, Yamagata T, Miyakawa T, Tanokura M, Momoi M, Momoi T. 2010. Autism spectrum disorder is related to endoplasmic reticulum stress induced by mutations in the synaptic cell adhesion molecule, CADM1. *Cell death & disease* **1**: e47.

Fujita E, Tanabe Y, Imhof BA, Momoi MY, Momoi T. 2012. A complex of synaptic adhesion molecule CADM1, a molecule related to autism spectrum disorder, with MUPP1 in the cerebellum. *J Neurochem* **123**: 886-894.

Fukuhara H, Kuramochi M, Fukami T, Kasahara K, Furuhata M, Nobukuni T, Maruyama T,
Isogai K, Sekiya T, Shuin T. 2002. Promoter methylation of TSLC1 and tumor
suppression by its gene product in human prostate cancer. *Jpn J Cancer Res* **93**: 605-609.
Garud NR, Messer PW, Buzbas EO, Petrov DA. 2015. Recent selective sweeps in North
American Drosophila melanogaster show signatures of soft sweeps. *PLoS Genet* **11**:
e1005004.
Gazave E, Ma L, Chang D, Coventry A, Gao F, Muzny D, Boerwinkle E, Gibbs RA, Sing CF,
Clark AG et al. 2014. Neutral genomic regions refine models of recent rapid human
population growth. *Proceedings of the National Academy of Sciences* **111**: 757-762.
Geurts P, Ernst D, Wehenkel L. 2006. Extremely randomized trees. *Machine Learning* **63**: 3-42.
Gillespie JH. 1991. *The causes of molecular evolution*. Oxford University Press, Oxford.
Gillespie JH. 2000. Genetic drift in an infinite population: the pseudohitchhiking model.
Genetics **155**: 909-919.
Gravel S, Henn BM, Gutenkunst RN, Indap AR, Marth GT, Clark AG, Yu F, Gibbs RA,
Bustamante CD, Altshuler DL. 2011. Demographic history and rare allele sharing among
human populations. *Proceedings of the National Academy of Sciences* **108**: 11983-11988.
Griffith OL, Montgomery SB, Bernier B, Chu B, Kasaian K, Aerts S, Mahony S, Sleumer MC,
Bilenky M, Haeussler M. 2008. ORegAnno: an open-access community-driven resource
for regulatory annotation. *Nucleic Acids Res* **36**: D107-D113.
Harrow J, Frankish A, Gonzalez JM, Tapanari E, Diekhans M, Kokocinski F, Aken BL, Barrell
D, Zadissa A, Searle S. 2012. GENCODE: The reference human genome annotation for
The ENCODE Project. *Genome Res* **22**: 1760-1774.
Hartfield M, Otto SP. 2011. Recombination and hitchhiking of deleterious alleles. *Evolution* **65**:
2421-2434.
Helgason A, Pálsson S, Thorleifsson G, Grant SF, Emilsson V, Gunnarsdottir S, Adeyemo A,
Chen Y, Chen G, Reynisdottir I. 2007. Refining the impact of TCF7L2 gene variants on
type 2 diabetes and adaptive evolution. *Nat Genet* **39**: 218-225.
Hermisson J, Pennings PS. 2005. Soft sweeps molecular population genetics of adaptation from
standing genetic variation. *Genetics* **169**: 2335-2352.
Hernandez RD, Kelley JL, Elyashiv E, Melton SC, Auton A, McVean G, Sella G, Przeworski M.
2011. Classic selective sweeps were rare in recent human evolution. *Science* **331**: 920-
924.
Huerta-Sánchez E, Jin X, Bianba Z, Peter BM, Vinckenbosch N, Liang Y, Yi X, He M, Somel
M, Ni P. 2014. Altitude adaptation in Tibetans caused by introgression of Denisovan-like
DNA. *Nature* **512**: 194-197.
Huff CD, Witherspoon DJ, Zhang Y, Gatenbee C, Denson LA, Kugathasan S, Hakonarson H,
Whiting A, Davis CT, Wu W. 2012. Crohn's disease and genetic hitchhiking at IBD5.
Mol Biol Evol **29**: 101-111.
Innan H, Kim Y. 2004. Pattern of polymorphism after strong artificial selection in a
domestication event. *Proc Natl Acad Sci U S A* **101**: 10667-10672.
Jensen JD, Thornton KR, Andolfatto P. 2008. An approximate Bayesian estimator suggests
strong, recurrent selective sweeps in Drosophila. *PLoS Genet* **4**: e1000198.
Jones DA, Wakeley J. 2008. The influence of gene conversion on linkage disequilibrium around
a selective sweep. *Genetics* **180**: 1251-1259.
Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. 2015. KEGG as a reference
resource for gene and protein annotation. *Nucleic Acids Res*: gkv1070.

Kaplan NL, Hudson R, Langley C. 1989. The" hitchhiking effect" revisited. *Genetics* **123**: 887-899.

Karasov T, Messer PW, Petrov DA. 2010. Evidence that adaptation in Drosophila is not limited by mutation at single sites. *PLoS Genet* **6**: e1000924.

Karolchik D, Hinrichs AS, Furey TS, Roskin KM, Sugnet CW, Haussler D, Kent WJ. 2004. The UCSC Table Browser data retrieval tool. *Nucleic Acids Res* **32**: D493-D496.

Kelly JK. 1997. A test of neutrality based on interlocus associations. *Genetics* **146**: 1197-1206.

Kent WJ. 2002. BLAT—the BLAST-like alignment tool. *Genome Res* **12**: 656-664.

Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, Haussler D. 2002. The human genome browser at UCSC. *Genome Res* **12**: 996-1006.

Kern AD, Jones CD, Begun DJ. 2002. Genomic effects of nucleotide substitutions in Drosophila simulans. *Genetics* **162**: 1753-1761.

Kern AD, Schrider DR. 2016. discoal: flexible coalescent simulations with selection. *Bioinformatics*: btw556.

Kim Y, Nielsen R. 2004. Linkage disequilibrium as a signature of selective sweeps. *Genetics* **167**: 1513-1524.

Kim Y, Stephan W. 2002. Detecting a local signature of genetic hitchhiking along a recombining chromosome. *Genetics* **160**: 765-777.

Kohda K, Kamiya Y, Matsuda S, Kato K, Umemori H, Yuzaki M. 2003. Heteromer formation of δ2 glutamate receptors with AMPA or kainate receptors. *Molecular Brain Research* **110**: 27-37.

Kong A, Frigge ML, Masson G, Besenbacher S, Sulem P, Magnusson G, Gudjonsson SA, Sigurdsson A, Jonasdottir A, Jonasdottir A. 2012. Rate of *de novo* mutations and the importance of father's age to disease risk. *Nature* **488**: 471-475.

Kong A, Thorleifsson G, Gudbjartsson DF, Masson G, Sigurdsson A, Jonasdottir A, Walters GB, Jonasdottir A, Gylfason A, Kristinsson KT et al. 2010. Fine-scale recombination rate differences between sexes, populations and individuals. *Nature* **467**: 1099-1103.

Kumar P, Henikoff S, Ng PC. 2009. Predicting the effects of coding non-synonymous variants on protein function using the SIFT algorithm. *Nat Protoc* **4**: 1073-1081.

Kuramochi M, Fukuhara H, Nobukuni T, Kanbe T, Maruyama T, Ghosh HP, Pletcher M, Isomura M, Onizuka M, Kitamura T. 2001. TSLC1 is a tumor-suppressor gene in human non-small-cell lung cancer. *Nat Genet* **27**: 427-430.

Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, Devon K, Dewar K, Doyle M, FitzHugh W. 2001. Initial sequencing and analysis of the human genome. *Nature* **409**: 860-921.

Landrum MJ, Lee JM, Benson M, Brown G, Chao C, Chitipiralla S, Gu B, Hart J, Hoffman D, Hoover J. 2016. ClinVar: public archive of interpretations of clinically relevant variants. *Nucleic Acids Res* **44**: D862-D868.

Langley CH, Stevens K, Cardeno C, Lee YCG, Schrider DR, Pool JE, Langley SA, Suarez C, Corbett-Detig RB, Kolaczkowski B. 2012. Genomic variation in natural populations of *Drosophila melanogaster*. *Genetics* **192**: 533-598.

Lee YCG, Langley CH, Begun DJ. 2013. Differential strengths of positive selection revealed by hitchhiking effects at small physical scales in Drosophila melanogaster. *Mol Biol Evol*: mst270.

Li H. 2011. A new test for detecting recent positive selection that is free from the confounding impacts of demography. *Mol Biol Evol* **28**: 365-375.

Li H, Durbin R. 2011. Inference of human population history from individual whole-genome sequences. *Nature* **475**: 493-496.

Li MJ, Wang LY, Xia Z, Wong MP, Sham PC, Wang J. 2013. dbPSHP: a database of recent positive selection across human populations. *Nucleic Acids Res*: gkt1052.

Lin K, Li H, Schlötterer C, Futschik A. 2011. Distinguishing positive selection from neutral evolution: boosting the performance of summary statistics. *Genetics* **187**: 229-244.

Liu X, Somel M, Tang L, Yan Z, Jiang X, Guo S, Yuan Y, He L, Oleksiak A, Zhang Y. 2012. Extension of cortical synaptic development distinguishes humans from chimpanzees and macaques. *Genome Res* **22**: 611-622.

Liu X, Wu C, Li C, Boerwinkle E. 2016. dbNSFP v3. 0: A One‑Stop Database of Functional Predictions and Annotations for Human Nonsynonymous and Splice‑Site SNVs. *Hum Mutat* **37**: 235-241.

Lohmueller KE, Albrechtsen A, Li Y, Kim SY, Korneliussen T, Vinckenbosch N, Tian G, Huerta-Sanchez E, Feder AF, Grarup N. 2011. Natural selection affects multiple aspects of genetic variation at putatively neutral sites across the human genome. *PLoS Genet* **7**: e1002326.

Lonsdale J, Thomas J, Salvatore M, Phillips R, Lo E, Shad S, Hasz R, Walters G, Garcia F, Young N. 2013. The genotype-tissue expression (GTEx) project. *Nat Genet* **45**: 580-585.

Luján R, Shigemoto R, Lopez-Bendito G. 2005. Glutamate and GABA receptor signalling in the developing brain. *Neuroscience* **130**: 567-580.

Macpherson JM, Sella G, Davis JC, Petrov DA. 2007. Genomewide spatial correspondence between nonsynonymous divergence and neutral polymorphism reveals extensive adaptation in Drosophila. *Genetics* **177**: 2083-2099.

Marsden CD, Ortega-Del Vecchyo D, O'Brien DP, Taylor JF, Ramirez O, Vilà C, Marques-Bonet T, Schnabel RD, Wayne RK, Lohmueller KE. 2016. Bottlenecks and selective sweeps during domestication have increased deleterious genetic variation in dogs. *Proceedings of the National Academy of Sciences* **113**: 152-157.

Marth GT, Czabarka E, Murvai J, Sherry ST. 2004. The allele frequency spectrum in genome-wide human variation data reveals signals of differential demographic history in three large world populations. *Genetics* **166**: 351-372.

Maynard Smith J, Haigh J. 1974. The hitch-hiking effect of a favourable gene. *Genet Res* **23**: 23-35.

Nei M, Li W-H. 1979. Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proceedings of the National Academy of Sciences* **76**: 5269-5273.

Nielsen R, Bustamante C, Clark AG, Glanowski S, Sackton TB, Hubisz MJ, Fledel-Alon A, Tanenbaum DM, Civello D, White TJ. 2005a. A scan for positively selected genes in the genomes of humans and chimpanzees. *PLoS Biol* **3**: e170.

Nielsen R, Williamson S, Kim Y, Hubisz MJ, Clark AG, Bustamante C. 2005b. Genomic scans for selective sweeps using SNP data. *Genome Res* **15**: 1566-1575.

Nunney L, Muir B. 2015. Peto's paradox and the hallmarks of cancer: constructing an evolutionary framework for understanding the incidence of cancer. *Phil Trans R Soc B* **370**: 20150161.

Orr HA, Betancourt AJ. 2001. Haldane's sieve and adaptation from the standing genetic variation. *Genetics* **157**: 875-884.

Orr HA, Unckless RL. 2014. The population genetics of evolutionary rescue. *PLoS Genet* **10**: e1004551.

Pavlidis P, Jensen JD, Stephan W. 2010. Searching for footprints of positive selection in whole-genome SNP data from nonequilibrium populations. *Genetics* **185**: 907-922.

Pei B, Sisu C, Frankish A, Howald C, Habegger L, Mu XJ, Harte R, Balasubramanian S, Tanzer A, Diekhans M. 2012. The GENCODE pseudogene resource. *Genome Biol* **13**: 1.

Pennings PS, Hermisson J. 2006a. Soft sweeps II—molecular population genetics of adaptation from recurrent mutation or migration. *Mol Biol Evol* **23**: 1076-1084.

Pennings PS, Hermisson J. 2006b. Soft sweeps III: the signature of positive selection from recurrent mutation. *PLoS Genet* **2**: e186.

Pickrell JK, Coop G, Novembre J, Kudaravalli S, Li JZ, Absher D, Srinivasan BS, Barsh GS, Myers RM, Feldman MW. 2009. Signals of recent positive selection in a worldwide sample of human populations. *Genome Res* **19**: 826-837.

Przeworski M, Coop G, Wall JD. 2005. The signature of positive selection on standing genetic variation. *Evolution* **59**: 2312-2323.

Pybus M, Dall'Olio GM, Luisi P, Uzkudun M, Carreño-Torres A, Pavlidis P, Laayouni H, Bertranpetit J, Engelken J. 2013. 1000 Genomes Selection Browser 1.0: a genome browser dedicated to signatures of natural selection in modern humans. *Nucleic Acids Res*: gkt1188.

Pybus M, Luisi P, Dall'Olio GM, Uzkudun M, Laayouni H, Bertranpetit J, Engelken J. 2015. Hierarchical boosting: a machine-learning framework to detect and classify hard selective sweeps in human populations. *Bioinformatics*: btv493.

Qian W, Zhou H, Tang K. 2015. Recent Coselection in Human Populations Revealed by Protein–Protein Interaction Network. *Genome Biol Evol* **7**: 136-153.

Ronen R, Udpa N, Halperin E, Bafna V. 2013. Learning natural selection from the site frequency spectrum. *Genetics* **195**: 181-193.

Ruwende C, Khoo S, Snow R, Yates S, Kwiatkowski D, Gupta S, Warn P, Allsopp C, Gilbert S, Peschu N. 1995. Natural selection of hemi-and heterozygotes for G6PD deficiency in Africa by resistance to severe malaria. *Nature* **376**: 246-249.

Sabeti PC, Reich DE, Higgins JM, Levine HZ, Richter DJ, Schaffner SF, Gabriel SB, Platko JV, Patterson NJ, McDonald GJ. 2002. Detecting recent positive selection in the human genome from haplotype structure. *Nature* **419**: 832-837.

Sabeti PC, Varilly P, Fry B, Lohmueller J, Hostetter E, Cotsapas C, Xie X, Byrne EH, McCarroll SA, Gaudet R. 2007. Genome-wide detection and characterization of positive selection in human populations. *Nature* **449**: 913-918.

Sattath S, Elyashiv E, Kolodny O, Rinott Y, Sella G. 2011. Pervasive adaptive protein evolution apparent in diversity patterns around amino acid substitutions in Drosophila simulans. *PLoS Genet* **7**: e1001302.

Schrider D, Shanku A, Kern A. 2016. Effects of Linked Selective Sweeps on Demographic Inference and Model Selection. *Genetics* doi:10.1534/genetics.116.190223.

Schrider DR, Kern AD. 2015. Inferring selective constraint from population genomic data suggests recent regulatory turnover in the human brain. *Genome Biol Evol* **7**: 3511-3528.

Schrider DR, Kern AD. 2016. S/HIC: Robust Identification of Soft and Hard Sweeps Using Machine Learning. *PLoS Genet* **12**: e1005928.

Schrider DR, Mendes FK, Hahn MW, Kern AD. 2015. Soft shoulders ahead: spurious signatures of soft and partial selective sweeps result from linked hard sweeps. *Genetics* **200**: 267-284.

Sheehan S, Song YS. 2016. Deep learning for population genetic inference. *PLoS Comput Biol* **12**: e1004845.

Siepel A, Bejerano G, Pedersen JS, Hinrichs AS, Hou M, Rosenbloom K, Clawson H, Spieth J, Hillier LW, Richards S. 2005. Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes. *Genome Res* **15**: 1034-1050.

Stephan W, Wiehe TH, Lenz MW. 1992. The effect of strongly selected substitutions on neutral polymorphism: analytical results based on diffusion theory. *Theor Popul Biol* **41**: 237-254.

Stephens JC, Reich DE, Goldstein DB, Shin HD, Smith MW, Carrington M, Winkler C, Huttley GA, Allikmets R, Schriml L. 1998. Dating the origin of the CCR5-Δ32 AIDS-resistance allele by the coalescence of haplotypes. *The American Journal of Human Genetics* **62**: 1507-1515.

Storey JD. 2002. A direct approach to false discovery rates. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* **64**: 479-498.

Swanson WJ, Vacquier VD. 2002. The rapid evolution of reproductive proteins. *Nature Reviews Genetics* **3**: 137-144.

Tajima F. 1989. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* **123**: 585-595.

Tishkoff SA, Reed FA, Ranciaro A, Voight BF, Babbitt CC, Silverman JS, Powell K, Mortensen HM, Hirbo JB, Osman M. 2007. Convergent adaptation of human lactase persistence in Africa and Europe. *Nat Genet* **39**: 31-40.

Van Der Weyden L, Arends MJ, Chausiaux OE, Ellis PJ, Lange UC, Surani MA, Affara N, Murakami Y, Adams DJ, Bradley A. 2006. Loss of TSLC1 causes male infertility due to a defect at the spermatid stage of spermatogenesis. *Mol Cell Biol* **26**: 3595-3609.

Voight BF, Kudaravalli S, Wen X, Pritchard JK. 2006. A map of recent positive selection in the human genome. *PLoS Biol* **4**: e72.

Watterson G. 1975. On the number of segregating sites in genetical models without recombination. *Theor Popul Biol* **7**: 256-276.

Welter D, MacArthur J, Morales J, Burdett T, Hall P, Junkins H, Klemm A, Flicek P, Manolio T, Hindorff L. 2014. The NHGRI GWAS Catalog, a curated resource of SNP-trait associations. *Nucleic Acids Res* **42**: D1001-D1006.

Wiehe T, Stephan W. 1993. Analysis of a genetic hitchhiking model, and its application to DNA polymorphism data from Drosophila melanogaster. *Mol Biol Evol* **10**: 842-854.

Williamson SH, Hubisz MJ, Clark AG, Payseur BA, Bustamante CD, Nielsen R. 2007. Localizing recent adaptive evolution in the human genome. *PLoS Genet* **3**: e90.

Wilson BA, Petrov DA, Messer PW. 2014. Soft selective sweeps in complex demographic scenarios. *Genetics* **198**: 669-684.

Wu YY, Yang Y, Xu YD, Yu HL. 2015. Targeted disruption of the spermatid‑specific gene Spata31 causes male infertility. *Mol Reprod Dev* **82**: 432-440.

Yates A, Akanni W, Amode MR, Barrell D, Billis K, Carvalho-Silva D, Cummins C, Clapham P, Fitzgerald S, Gil L. 2016. Ensembl 2016. *Nucleic Acids Res* **44**: D710-D716.

Zhang P, Si X, Skogerbø G, Wang J, Cui D, Li Y, Sun X, Liu L, Sun B, Chen R. 2014. piRBase: a web resource assisting piRNA functional study. *Database* **2014**: bau110.

Zhiling Y, Fujita E, Tanabe Y, Yamagata T, Momoi T, Momoi MY. 2008. Mutations in the gene encoding CADM1 are associated with autism spectrum disorder. *Biochem Biophys Res Commun* **377**: 926-929.

898
899

900   **Table 1: Number of sweeps of each type detected in each population sample.**

Population	# of Hard Sweeps	# of Soft Sweeps
JPT	61 (5.8%)	998 (94.2%)
CEU	66 (6.5%)	947 (93.5%)
PEL	32 (4.7%)	655 (95.3%)
GWD	5 (0.6%)	795 (99.4%)
YRI	13 (1.6%)	797 (98.4%)
LWK	3 (0.4%)	805 (99.6%)

Figure 1: Enrichment of various annotation features in regions classified as sweeps or linked to sweeps relative. The fold enrichment is the ratio of the number of base pairs in the intersection between windows assigned to a given class and an annotation feature divided by the mean of this intersection across the permuted data sets (Methods). This was calculated separately for each population. (A) Enrichment of elements in windows classified as hard sweeps. (B) Same as A, but for soft sweeps. (C) Enrichment of elements in windows classified as affected by linked hard sweeps. (D) Linked soft sweeps.

922
923 **Figure 2: The number of windows assigned to each class by S/HIC in each population.**
924
925

Figure 3: Enrichment of pairs of interacting genes each falling within a window classified
as a sweep. The fold enrichment is the ratio of the number of pairs of interacting genes
overlapping a window classified as a sweep of a given type divided by the mean of this number
across the permuted data sets (Methods). This was calculated separately for each population.
When no pairs of interacting sweep genes were observed in our true data set or a population, no
bar was drawn. (A) Enrichment of pairs of genes encoding protein products that physically
interact with each other (data from BioGRID) and both overlap hard sweep windows. (B) Same
as A, but for soft sweeps. (C) Enrichment of pairs of genes, one of which is encodes a
transcription factor that affects expression of the other (data from ORegAnno), where both
overlap hard sweep windows. (D) Same as D, but for soft sweeps. (E) Enrichment of pairs of
genes for which a genetic interaction has been observed (data from BioGRID) and both overlap
hard sweep windows. (F) Same as E, but for soft sweeps.

940
Figure 4: Hard selective sweep near several *SPATA31* spermatogenesis-associated genes.
The S/HIC classification tracks show the raw classifier output for each population (red=hard
sweep, blue=soft sweep, light red=hard-linked, light blue=soft-linked, black=neutral). We also
show the values of various population genetic summary and test statistics (π, Tajima's D, Kelly's
Z_{nS}, and the SweepFinder composite likelihood ratio, or CLR). To avoid clutter, we only show
statistics from CEU.
947
948

949
950 **Figure 5: Soft selective sweeps near *CADM1*.** The same tracks are shown as in Figure 4.

SUPPLEMENTARY FIGURE AND TABLE LEGENDS

Figure S1: Heatmaps showing the accuracy of our six classifiers on test data, one for each population. On the *y*-axis, we show the location of the sweep relative to the classified window (i.e. the central sub-window), with the exception of the "Neutral" case where there is no sweep. The test data were simulated under the same demographic models used for training. On the *x*-axis we show the class inferred by S/HIC. A perfect classifier would infer "Hard" for 100% test instance where a hard sweep is in the focal sub-window (and analogously for soft sweeps), "Hard-linked" for 100% of cases where a hard sweep occurs elsewhere (and analogously for soft sweeps not located in the central sub-window), and "Neutral" for 100% of cases with no sweep. (A) Test results for CEU. (B) GWD. (C) JPT. (D) LWK. (E) PEL. (F) YRI.

Figure S2: Histograms of H_{12} and H_2/H_1 within windows classified has hard sweeps, soft sweeps, or neutral for each population.

Figure S3: Soft selective sweep in *GRIA2*. The same tracks are shown as in Figures 4 and 5.

Table S1: Enrichment of various sequence annotations in each S/HIC class.

Table S2: Enrichment of annotation terms in hard and soft sweeps (only terms with $q<0.05$ for at least one sweep type in at least one population are shown).

Table S3: Example command lines used to generate training data for each population, with a soft sweep occurring in the central sub-window.