MtrA is an essential regulator that coordinates antibiotic production

and sporulation in *Streptomyces* species

Nicolle F. Som[1#], Daniel Heine[2#], John T. Munnoch[1], Neil A. Holmes[1], Felicity Knowles[1],

Govind Chandra[2], Ryan F. Seipke[1,3], Paul A. Hoskisson[4], Barrie Wilkinson[2*] and Matthew I

Hutchings[1*]

# Authors contributed equally; * Correspondence

[1]School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich,

United Kingdom. NR4 7TJ.

[2]Department of Molecular Microbiology, John Innes Centre, Norwich Research Park, Norwich,

United Kingdom. NR4 7TJ.

[3] School of Molecular & Cellular Biology, Astbury Centre for Structural Molecular Biology,

University of Leeds, Leeds, LS2 9JT, UK

[4]Strathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, 161,

Cathedral Street, Glasgow, G4 0RE, UK

Abstract. *Streptomyces* bacteria make numerous secondary metabolites, including half of all

known antibiotics. Understanding the global regulation of secondary metabolism is important

because most *Streptomyces* natural products are not made under laboratory conditions and

unlocking 'cryptic' biosynthetic gene clusters (BGCs) is a major focus for natural product

discovery. Production is coordinated with sporulation but the regulators that coordinate

development with antibiotic biosynthesis are largely unknown. Here we characterise a highly

conserved actinobacterial response regulator called MtrA in antibiotic-producing

Streptomyces species. We show that MtrA is an essential global regulator of secondary

metabolism that directly activates antibiotic production in in *S. coelicolor* and *S. venezuelae*.

MtrA also controls key developmental genes required for DNA replication and cell division and

we propose that MtrA is the missing link that coordinates secondary metabolism with

development in *Streptomyces* species.

Introduction.

Streptomyces secondary metabolites account for two thirds of all known antibiotics and

numerous other compounds that are used in human medicine as anticancer, anti-parasitic,

antiviral and immunosuppressant drugs. Discovery of these natural products peaked in the

1950s but there has been a resurgence of interest in the 21[st] century, driven by genome

sequencing and the increasing threat of drug resistant infections [1]. Despite their importance

however, we still have a poor understanding of how *Streptomyces* bacteria control the

production of their secondary metabolites. This is important because ≥75% of their secondary

metabolite biosynthetic gene clusters (BGCs) are not expressed in laboratory culture and

activating cryptic BGCs could facilitate the discovery of new antibiotics and other useful

compounds [2,3].

The major way in which bacteria sense and respond to their environment is through

two-component systems and several have been implicated in the regulation of antibiotic

production in *Streptomyces* species [4]. Two component systems typically consist of a

bifunctional sensor kinase and a cognate response regulator [5]. The sensor kinase perceives

an extracellular signal and activates its cognate response regulator through a two-step

phosphorylation. The phosphorylated regulator (RR~P) brings about a response to the original

signal, usually by modulating target gene expression. In the absence of signal, the bifunctional

sensor kinase dephosphorylates its response regulator to keep the response switched off [6].

The delicate balance of kinase and phosphatase activities is crucial in modulating the activity

of the response regulator and its target genes during the bacterial cell cycle [5]. Cross-talk

between systems is rare in wild-type cells but removal of a sensor kinase can result in

constitutive activation of its response regulator by a non-cognate sensor kinase or by the

cellular pool of acetyl phosphate [7]. Similarly, altering a sensor kinase to block its phosphatase

activity might result in a response regulator that cannot be dephosphorylated and is

constitutively active [5].

Here we report characterisation of the highly conserved actinobacterial response

regulator MtrA in the model organism *Streptomyces venezuelae* [8]. MtrA was first identified as

an essential regulator in *Mycobacterium tuberculosis* (<u>M</u>ycobacterium <u>t</u>uberculosis <u>r</u>egulator

<u>A</u>) [9]. *M. tuberculosis* MtrA (TB-MtrA) regulates the expression of *dnaA* and *dnaN*, which are

essential for DNA replication, and sequesters the origin of DNA replication, *oriC*, in dividing

cells [10]. TB-MtrA also regulates the expression of cell division genes and interacts directly with

the DnaA protein [10]. MtrA is activated when the MtrB sensor kinase localises to the site of cell

division through interaction with FtsI and DivIVA. These data have led to a model in which

oscillations in TB-MtrA~P levels play a key role in cell cycle progression by repressing DNA

replication and activating cell division [10-12]. In the closely related *Corynebacterium glutamicum*,

Cg-MtrA controls genes involved in cell wall remodelling and the osmotic stress response [13,14]

but deletion of the *mtrAB* genes gives rise to elongated cells which are indicative of a defect

in cell division [15].

Streptomyces bacteria are filamentous saprophytes which grow through the soil as

branching substrate mycelium that extends at the hyphal tips. Nutrient starvation triggers the

production of reproductive aerial hyphae that accelerate DNA replication, generating up to 200

copies of the chromosome in each aerial hypha, before undergoing cell division to form chains

of unigenomic spores. Aerial hyphae production and sporulation are coordinated with the

production of bioactive secondary metabolites, including antibiotics. *S. coelicolor* has

traditionally been used to study development and antibiotic production because it makes

pigmented antibiotics and spores but it only differentiates into aerial hyphae and spores on

solid agar. *S. venezuelae* has emerged as an alternative model because it completes a full

developmental life cycle in liquid growth medium in ~20 hours [16]. Here we report that MtrA is

essential for the growth of *S. venezuelae* and directly regulates the expression of genes

involved in DNA replication, cell division and secondary metabolism. ChIP-seq throughout the

S. venezuelae life-cycle shows that Sv-MtrA activity displays a biphasic plasticity such that it

is active during vegetative growth and sporulation, but inactive during formation of aerial

hyphae. We propose that in *Streptomyces* species, MtrA represses DNA replication and cell

division during active growth and following sporulation. We also show that Sv-MtrA binds to

sites spanning ~85% of the BGCs in *S. venezuelae* and directly modulates expression of

target genes in at least 75% of these. ChIP-seq in *S. coelicolor* yielded similar results and key

developmental genes bound by MtrA in both streptomycetes include *bldM, ftsZ, ssgA, smc,*

smeA, whiI, whiB, whiD and *wblE.* We propose that MtrA is an essential master regulator that

coordinates differentiation and secondary metabolism in *Streptomyces* species.

Results

MtrA is essential in *Streptomyces venezuelae*.

To investigate the function of the MtrAB two component system in *S. venezuelae* we deleted

either *mtrA* or *mtrB* [17]. *mtrB* was deleted easily but all attempts to delete *mtrA* were

unsuccessful until we introduced a second copy of *mtrA in trans* suggesting MtrA is essential

in *S. venezuelae.* It follows that MtrA must be active in the absence of MtrB, otherwise deleting

mtrB would be lethal. We propose that deletion of *mtrB* leads to constitutive phosphorylation

of MtrA, either by the cellular pool of acetyl phosphate or another sensor kinase. We observed

a similar result with *Streptomyces coelicolor* VanRS, where VanR and its target genes are

constitutively active in a ΔvanS mutant [7]. Surprisingly, deletion of *mtrB* has no effect on the

growth rate of *S. venezuelae* in liquid medium (Figure S1).

MtrA activity changes during the life cycle of *S. venezuelae*.

Microarray data [18] shows that *mtrA* expression levels remain fairly constant during the *S.*

venezuelae life cycle and we do not detect any significant change in MtrA protein levels during

the life cycle using immunoblotting (Figure S2). Attempts to compare levels of phosphorylated

and non-phosphorylated MtrA using Phostag were unsuccessful and so, to examine MtrA

activity during the *S. venezuelae* life cycle, we performed ChIP-seq at two-hour time points (8-

20 hours) from active growth through to sporulation (accession number GSE84311, Table S1).

We expressed MtrA-3xFlag *in trans* under the control of its own promoter and deleted the

native *mtrA* gene*.* Survival of this strain shows that MtrA-3xFlag is functional and there is no

significant difference in growth rate between this strain and the wild-type (Figure S1). Analysis

of the ChIP-seq data using a P value > 0.05 revealed that only one target is enriched at all

time points, the promoter region of the *ectABCD* operon, which is the BGC for the secondary

metabolites ectoine and 5' hydroxyectoine (5HE). We do not detect ectoine or 5HE in the wild-

type or Δ*mtrB* mutant using LCMS with authentic standards so we predict that MtrA represses

ectABCD expression throughout the life cycle. Analysis of all the MtrA ChIP-seq datasets

using $P > 0.05$ shows that the pattern of binding of MtrA is dynamic throughout the lifecycle

with peak DNA binding activity seen at 10 and 20 hours, coinciding with mid-exponential

growth, and at 20 hours, in spores (Figure S3)*.* Mapping ChIP-seq reads at targets such as

the *mtrA* promoter provides a good illustration of this biphasic activity (Figure 1). Differential

RNA sequencing shows that the major *mtrAB-lpqB* transcript is leaderless, with an MtrA

binding site immediately upstream of the -35 site, suggesting positive autoregulation. A minor

transcript starts at -79bp suggesting two promoters might drive expression of the *mtrAB-lpqB*

operon (Figure 1).

Figure 1. MtrA autoregulates its own expression. *ChIP-seq during the S. venezuelae life cycle shows that MtrA has highest activity in actively growing mycelium (10 and 12 hours) and spores (20 hours). The peaks shown are at the mtrA promoter, which has two transcript start sites at +1 (P1) and -79 (P2). Deletion of mtrB increases mtrA transcript levels ~3-fold indicating positive autoregulation.*

Identifying the MtrA binding site. To identify an MtrA consensus binding sequence we used

MEME [19] to analyse three MtrA target promoters identified by ChIP-seq and confirmed *in vitro*

using electrophoretic mobility shift assays (EMSAs). This analysis identified an AT-rich 7 bp

motif present at all three promoters (Figure 2). We then used MEME to analyse 50 bp of

sequence from beneath each peak at the 14h and the 16/18h time points and this analysis

identified a conserved sequence at each target which is effectively a direct repeat of the AT

rich motif (Figure 2). However, many of the sequences enriched in the MtrA ChIP-seq dataset

do not obviously contain this motif and we hypothesise that MtrA may interact with other

transcription factors. Immunoprecipitation experiments using anti-Flag beads pulled down four

different regulators in the MtrA-3xFlag cultures that are not immunoprecipitated in the wild-

type control. The upstream regions of the genes encoding these regulators (*sven15_0243*,

sven15_2691, *sven15_3571* and *sven15_4644*) are all bound by MtrA (Table S1).

Sven15_3571 is DnaA, which initiates DNA replication and acts as a transcription factor in

bacteria, and is also bound *in vivo* by TB-MtrA {Purushotham:2015iq}. Thus, our data suggests

(but does not prove) that MtrA forms complexes with other DNA binding proteins and this could

explain why there are so many enrichment peaks in the 20-hour ChIP-seq dataset. Interaction

between regulatory proteins to form heterodimers is not unprecedented in *S. venezuelae.* The

response regulator BldM modulates one set of target genes as a homodimer and another set

by forming heterodimers with the WhiI response regulator [20]. The developmental regulators

WhiA and WhiB are also proposed to interact and have identical ChIP-seq regulons [21].

MtrA regulates the expression of DNA replication and cell division genes.

ChIP-seq shows that MtrA activity changes during the *S. venezuelae* life cycle (Figure 1) and

RNA sequencing of wild-type and Δ*mtrB* strains shows that expression of *mtrA* is 3-fold

upregulated in the Δ*mtrB* mutant (Table S2) which is consistent with activation by MtrA~P.

Given the essentiality of MtrA and its altered activities at different stages of the life cycle we

predicted that it could play a key role in regulating cell cycle progression in *S. venezuelae*.

Indeed, ChIP-seq shows that many key developmental genes are bound by MtrA in *S.*

venezuelae and *S. coelicolor* (see Tables S1 and S2 for complete lists). Remarkably, *M.*

tuberculosis and *S. venezuelae* MtrA proteins share some common targets, including the

promoter regions of *wblE*, *dnaA* and *dnaN*, and the *oriC* region between *dnaA* and *dnaN*

(Figure 3). RNA-seq shows that deletion of *mtrB* does not affect the expression levels of *dnaA*

or *dnaN* under the conditions we used but deletion of *mtrB* causes a two-fold increase in *wblE*

transcript levels, suggesting MtrA-dependent activation. MtrA also directly represses *adpA*

expression, which is down-regulated nearly 4-fold in the Δ*mtrB* mutant (Table S2).

Target	P value	Binding site	
Sven15_0205 (ectA)	4.23e-5	GCGTTACCCATCCGTTATAAACGGTCAT	
Sven15_0880 (cmlF)	4.23e-5	ATTCGCCATCTACGTTCTCGGGCTGGCC	
Sven15_3571 (dnaA)	1.51e-5	CACAATCTTTTCCGTTCTGTCCTTACCT	

14h   16 and 18h

Figure 2. MtrA binds to 6-5-6bp direct repeat motif. The table shows three MtrA ChIP targets that are bound in vitro by purified MtrA protein. They are the promoter regions of ectABCD operon (ectoine BGC), cmlF (chloramphenicol transporter) and dnaA (transcription factor and initiator of DNA replication). MEME analysis identified a conserved 7bp motif (coloured) that is present as a 6-5-6bp repeat motif at the ectA promoter (second half underlined). MEME analysis of different subsets of ChIP targets identified the same direct repeat motif (shown for the targets identified at 14 hours and 16 and 18 hours.

AdpA controls key developmental genes and represses DNA replication in *Streptomyces*

species [22,23]. *wblE* encodes a WhiB-like transcription factor of unknown function that appears

to be essential [24]. Sv-MtrA also binds upstream of the cell division genes *ssgA* and *ssgB*,

which are upregulated in the Δ*mtrB* mutant, and the *ftsZ* gene, which is unaffected by loss of

MtrB, probably because this gene is subject to complex regulation (Figure 3B and Table S2).

SsgAB localise the divisome marker protein FtsZ to the correct positions in aerial hyphae to

mark the sites of cell division, prior to sporulation [25,26]. Sv-MtrA also regulates expression of

the *smeA-ssfA* operon, which is elevated between 2- and 4-fold in the Δ*mtrB* mutant. SmeA

targets the DNA pump SffA to the cell division septa [16]. MtrA also binds upstream of (and

activates some) key developmental regulators, including *whiB*, *whiD*, *whiI* and *bldM* (Table

S2) [20,21,27].

Figure 3. MtrA binds to targets required for DNA replication and cell division in S. venezuelae. *Top. MtrA ChIP peaks upstream of dnaA, which encodes the initiator of DNA replication, and dnaN which encodes the DNA polymerase beta clamp subunit and at the origin of DNA replication, oriC. Bottom. MtrA ChIP peaks upstream of ftsZ, which encodes the Z ring forming FtsZ protein and ssgA, whose product helps localise FtsZ to the sites of cell division during sporulation.*

Many of the Sv-MtrA target genes are also repressed by BldD (in complex with c-di-GMP).

Sv-MtrA does not regulate *bldD* expression but it does regulate the expression of genes

encoding proteins that metabolise cyclic di-GMP [28] (Table S2).

MtrA regulates global BGC expression in *Streptomyces venezuelae*.

ChIP-seq shows that MtrA binds to sites spanning 28 of the 31 BGCs predicted by antiSMASH

3.0 in the *S. venezuelae* genome [29] (Table S3). The only BGCs with no MtrA enrichment are

those encoding biosynthesis of the desferrioxamine siderophores, the WhiE polyketide spore

pigment and an unknown natural product. Of these three clusters, the WhiE BGC is

upregulated in the Δ*mtrB* mutant suggesting indirect regulation by MtrA, probably via BldM [20].

The other two BGCs are unaffected by loss of MtrB (Table S3). Of the 28 BGCs that are bound

by MtrA, nine have genes that are positively regulated by MtrA, 10 have genes that are

negatively regulated by MtrA and three are subject to both positive and negative regulation by

MtrA at individual genes within the gene cluster (Table S3). The other six BGCs have sites

that are enriched in MtrA ChIP-seq but expression of the genes nearest the ChIP-seq peaks

are not affected by deletion of *mtrB* under the conditions we used (Table S3). We predict that

one of these, the *ectABCD* operon, is repressed by MtrA because the *ectA* promoter is bound

by MtrA at all time points. Consistent with this, we cannot detect ectoine or 5HE in the wild-

type or Δ*mtrB* strains. Since most of the *S. venezuelae* BGCs are uncharacterised we know

little about their cluster specific regulation, or their products. The antiSMASH predictions we

have used here are also likely to include additional flanking genes that may not be part of the

BGC so the actual number of BGCs bound by Sv-MtrA may be slightly lower.

MtrA activates chloramphenicol production in *S. venezuelae*.

Phenotypic screening of the Δ*mtrB* mutant revealed that a cryptic antibacterial is activated by

removing MtrB but not by simply over-expressing *mtrA* in wild-type *S. venezuelae* (Figure S4).

This is probably because MtrA~P levels only increase in the absence of MtrB. *S. venezuelae*

encodes the biosynthetic pathway for the antibiotic chloramphenicol, which according to

previous studies is silent in the wild-type strain [30]. ChIP-seq shows that Sv-MtrA binds in the

intergenic region between the divergent *cmlF* (*sven15_0879*) and *cmlN* (*sven15_0880*) genes

(Figure 4) and expression of *cmlN* is 6-fold upregulated in the Δ*mtrB* mutant, consistent with

direct activation by MtrA. CmlN is an efflux permease, predicted to export chloramphenicol [30].

HPLC confirmed that chloramphenicol is produced in the Δ*mtrB* mutant but we also detected

very low levels in the wild-type strain suggesting the cluster is not completely silent (Figure 4).

Cultivation for 24 hours in biological and technical triplicates confirmed an increased

production of chloramphenicol in the Δ*mtrB* mutant with a mean concentration of 0.41 mg/L

which is >30 times higher than the wild-type strain (0.013 mg/L) or wild-type over-expressing

mtrA (0.010 mg/L).

Deleting *mtrB* has a global effect on the *S. venezuelae* metabolome.

While loss of MtrB clearly leads to increased production of chloramphenicol, we were

interested to know if there are any other effects on the metabolome. We cultivated the wild

type strain and three independently isolated Δ*mtrB* mutants in biological and technical

triplicates and analysed the extracts by UPLC/HRMS using an untargeted metabolomics

approach. Runs were aligned to compensate for between-run variation and a peak-picking

algorithm was applied to allow for the immaculate matching of each feature (a discrete *m/z*

value and its retention time) among all runs. Following normalisation, features could be

compared quantitatively and their putative identity proposed based on their high-resolution

MS-signal. Comparing the level of metabolite signals, it appeared that all Δ*mtrB* mutants

showed increased production of a considerable portion of their putative secondary

metabolites. To display multidimensional data, we used Principle Component Analysis (PCA)

(Figure 5).

Figure 4. MtrA controls chloramphenicol production. *Top. MtrA ChIP peaks at the chloramphenicol (Cm) BGC show two significant peaks between cmlN and cmlF and a smaller peak upstream of the regulator cmlR which is P<0.05. Middle. Expression data for wild-type and ∆mtrB strains at 14 and 20 hours shows cmlF is 6-fold upregulated in ∆mtrB suggesting it is activated by MtrA. All the other Cm genes are downregulated in the mutant which may be due to negative feedback. Bottom left. HPLC shows the Cm standard and extracts of wild-type, ∆mtrB, wild-type plus empty pMS82 vector, and wild-type plus pMS82 expressing Sv-Y99C MtrA, TB-MtrA and TB-Y102C MtrA. Right. Zooming in shows that Sv-Y99C MtrA induces production of low levels of Cm production.*

243 Each sphere in the 3D Plot represents one dataset obtained from a particular UPLC-HRMS

244 run. Data from the *ΔmtrB* mutant strains clearly group together, and are distinct from data

245 obtained from the wild type, while variations within each group are comparably small. The 3D

246 Plot therefore shows consistent and global changes in the metabolome upon loss of MtrB

247 (Figure 5).

Figure 5. Loss of MtrB results in a global shift in the S. venezuelae metabolome.
Principle Component Analysis on the wild-type (blue dots) and triplicate samples of the
ΔmtrB strain (red, green and orange dots). Data from ΔmtrB mutant strains clearly group
together, and are distinct from data obtained from the wildtype while variations within each
group are comparably small.

MtrA directly activates antibiotic production in *Streptomyces coelicolor*.

MtrA is conserved in all sequenced *Streptomyces* genomes so we reasoned that it might

activate antibiotic production in other streptomycetes. To test this, we deleted *mtrB* in the

model organism *Streptomyces coelicolor* [31]. The 16S rDNA phylogenetic tree of the family

Streptomycetaceae shows that *S. venezuelae* (clade 40) is highly divergent from *S. coelicolor*

(clade 112) so we reasoned that if deleting *mtrB* switches on antibiotic production in these

distantly related species it is probably universal to all streptomycetes [32]. The *S. coelicolor*

Δ*mtrB* mutant over-produces the red antibiotic undecylprodigiosin and the blue antibiotic

actinorhodin and we also detected significant amounts of metacycloprodigiosin, a potent

anticancer compound encoded by the undecylprodigiosin BGC but not previously reported

from *S. coelicolor* (Figure 6). Other secondary metabolites are clearly down-regulated in the

Δ*mtrB* mutant, including the siderophores desferrioxamine B and E and germicidin A. We

performed ChIP-seq in *S. coelicolor* because its BGCs and their natural products are the most

well defined of any *Streptomyces* species [31,33,34]. The results show that MtrA binds to sites

spanning 21 of its 29 BGCs (Tables S4 and S5). MtrA binds upstream of the genes encoding

the actinorhodin activator ActII-4 and the undecylprodigiosin activator RedZ (Figure S11).

MtrA does not bind to the desferrioxamine BGC (*desABCD*) in *S. coelicolor* (Table S4) but

does bind upstream of *sco4394*, which encodes DesR, an iron dependent repressor of

desferrioxamine biosynthesis (Figure S5). A single type III polyketide gene, *SCO7221*, is

responsible for germicidin biosynthesis [35] and it is not regulated by MtrA, so the effect on

germicidin biosynthesis must be indirect.

TB-MtrA activates chloramphenicol production in *S. venezuelae*.

Modulating MtrA activity by deleting *mtrB* is time consuming and impractical when applied to

multiple *Streptomyces* strains. Instead we tested whether alternative, gain-of-function MtrA

proteins might be used to activate antibiotic production. We constructed expression vectors

for wild-type TB-MtrA, gain-of-function Y102C TB-MtrA [36] and a Y99C Sv-MtrA variant which

has the equivalent change to Y102C TB-MtrA (Figure S6). We were unable to delete the native

mtrA gene in strains carrying these constructs but all three MtrA constructs induced

chloramphenicol production in wild-type *S. venezuelae* (Figure 4). Y99C Sv-MtrA switches on

the production of relatively small amounts of chloramphenicol in wild-type *S. venezuelae* but

the TB-MtrA proteins switch on much higher levels of production, with the strain expressing

Y102C-TB-MtrA making more chloramphenicol than the Δ*mtrB* mutant (Figure 4). These

results suggest that expressing different MtrA proteins might be a useful avenue to explore in

terms of activating BGCs in *Streptomyces* species and our constructs are available from

AddGene (IDs 85988-94).

Figure 6. Deleting mtrB switches on antibiotic production in S. coelicolor. *(a) HPLC on extracts of wild-type S. coelicolor M145 cultures shows production of desferrioxamines and germicidin. (b) HPLC on extracts of an S. coelicolor ΔmtrB mutant shows production of actinorhodin and undecylprodigiosin is switched on while the production of the desferrioxamine siderophores and germicidin are switched off.*

Discussion.

Our data, and previously published work on MtrA in mycobacteria and corynebacteria, support

a model in which removing MtrB results in constitutive phosphorylation and activation of MtrA

which then constitutively activates or represses its target genes in the cell. In *S. coelicolor* and

S. venezuelae these include key developmental genes and there is a significant overlap with

the regulon controlled by the master repressor of development, BldD, which is activated for

DNA binding by the secondary messenger cyclic-di-GMP [28]. Exactly where MtrA fits in to the

hierarchy of developmental regulation is not yet clear but MtrA has an essential role in *S.*

venezuelae and shares some functions with *M. tuberculosis* MtrA [10]. Here we have focused

on the role of MtrA in regulating secondary metabolism and how it might be exploited to

activate cryptic BGCs. The effects on antibiotic production appear to be direct in both *S.*

coelicolor and *S. venezuelae* because MtrA activates expression of the chloramphenicol

transporter CmlN and switches on chloramphenicol production. It also binds to the promoters

of the cluster specific activators ActII-4 and RedZ in *S. coelicolor* and activates the products

under their control, actinorhodin and undecylprodigiosin. In a developmental time course it is

remarkable that MtrA binds to sites spanning 27 of the 31 BGCs in *S. venezuelae* and directly

affects the expression of target genes in at least 22 of these BGCs. In vegetatively growing *S.*

coelicolor MtrA binds to sites spanning 21 out of 29 BGCs. We propose therefore that MtrA is

a master regulator of secondary metabolism in *Streptomyces* species. Consistent with this,

unbiased metabolomics analysis of *S. venezuelae* shows that deleting *mtrB* results in a global

change in the metabolome. Given that secondary metabolite BGCs are often subject to

complex and multilevel regulation, and given the energy costs associated with making these

natural products, it is perhaps not surprising that we do not see obvious over-production of

other compounds in the Δ*mtrB* strains. We predict that deletion of known BGCs in the Δ*mtrB*

mutants and / or depletion of *mtrA* mRNA will have positive effects on the production of other

secondary metabolites. In summary, our study has revealed an important role for MtrA in the

life cycles of streptomycetes and our data suggest it might play a similar role to CtrA in

Caulobacter crescentus, acting as a master regulator of the cell cycle alongside other master

regulators such as BldD. Rapid progress has been made recently in elucidating the roles of

developmental regulators using *S. venezuelae* as a model system [16]. Our study shows that *S.*

venezuelae can also be used as a model to study global effects on secondary metabolism

and to elucidate the regulatory cascades that link secondary metabolism and differentiation.

Understanding these genetic circuits is essential if we are to unlock the full potential of these

bacteria.

Materials and Methods

Strains, plasmids and primers. The bacterial strains, plasmids and cosmids and primers used in this study are listed in Tables S6-8. The Sv- and TB-MtrA expression vectors have been deposited with AddGene (ID 85988-94). *S. venezuelae* NRRL B-65442 is deposited in the USDA Agricultural Research Services (ARS) Culture Collection (http://nrrl.ncaur.usda.gov/cgi-bin/usda/prokaryote/report.html?nrrlcodes=B%2d65442).

Plasmid stocks were prepared using Qiagen miniprep kits and cosmids were prepared as described previously [37]. Genes were deleted using the ReDirect PCR targeting method [17] and an *S. venezuelae* NRRL B-65442 cosmid library provided by Professor Mark Buttner at the John Innes Centre, Norwich. All of the expression constructs used in this work were made by Genscript by synthesising the *mtrA* alleles with or without 3' tags and then cloning into the required vectors. Liquid cultures of *E. coli* were routinely grown shaking at 220 rpm in Lennox Broth at 37°C. Liquid cultures of *S. coelicolor* or *S. venezuelae* were grown in Mannitol Yeast Extract Malt Extract (MYM) at 30°C, shaking at 220 rpm. Cultures grown on solid MYM agar were grown at 30°C. Spore stocks of *S. coelicolor* were prepared from cultures grown on Soya Flour + Mannitol (SFM) agar. All media recipes have been published previously and *Streptomyces* spores were prepared as described [37]. To determine the developmental growth in liquid culture *S. venezuelae* and mutant strains were grown, shaking in 35ml MYM in 250ml conical flasks containing springs at 30°C at 220rpm. A spore inoculum sufficient to reach an OD_{600} of 0.35 after 8 hours of growth was added to 35ml of MYM media in 250 ml flasks containing springs. The culture densities were measured at OD_{600}. Development in liquid cultures was monitored using an GXML3000B microscope from GX optical. Pictures of agar plate grown colonies were taken with a Zeiss SVII stereo microscope. SEM images were taken at the bioimaging facility at the John Innes Centre.

Purification of MtrA-His. The *mtrA* gene was cloned into pETDuet1 (Novagene) to express the protein with a C-terminal hexa-His tag and purified using a batch method. Cell pellets were resuspended in 25ml lysis buffer (75mM Tris-HCl pH8, 20mM NaCl, 0.1% Triton X100, 50 µl 10mg/ml lysozyme, 3 x Pierce EDTA-free Protease Inhibitor Mini Tablets (Thermo Scientific)

in 1L) and incubated for 30 minutes at room temperature. The cell lysate was sonicated 2 x

40 seconds at 50Hz with 1 minute in between sonication steps. Cell debris was removed by

centrifugation at 18,000rpm for 20 minutes at 4°C in Beckman Coulter Avanti® J-20 high

performance centrifuge using a JLA-25-50 rotor (Beckman Coulter). The supernatant was

transferred in a fresh 50ml Falcon tube and 350µl of Ni-NTA agarose beads (Qiagen) were

added and incubated under gentle agitation for 1 hour at 4°C. The Ni-NTA agarose beads

were spun down gently (maximum of 1200rpm, 4°C) and the supernatant was discarded. The

Ni-NTA beads were resuspended in 2ml wash buffer (75mM Tris-HCl pH8, 200mM NaCl, 10%

glycerol, 10mM $MgCl_2$, 0.1mM DTT, 20mM 2-mercaptoethanol in 1L) and transferred in

Polypropylene columns (1ml, Qiagen). The beads in the column were washed with 20ml wash

buffer. The protein was eluted from the beads with 2.5ml elution buffer (wash buffer plus

350mL imidazole in 1L). Polyclonal antiserum was raised by Cambridge Research

Biochemicals. Immunoblotting was performed as described previously [38].

DNA binding studies. Chromatin Immunoprecipitation followed by sequencing (ChIP-seq)

was performed on *S. coelicolor* M145 ∆*mtrA* + MtrA-3xFlag and *S. venezuelae* NRRL B-65442

∆*mtrA* + MtrA-3xFlag grown in liquid MYM medium [37]. Note that *S. coelicolor* M145 grows

vegetatively and does not differentiate under these growth conditions whereas *S. venezuelae*

NRRL B-65442 undergoes a full life cycle in 20 hours. Samples were taken at 16 and 20 hours

for *S. coelicolor* and at 8, 10, 12, 14, 16, 18 and 20 hours for *S. venezuelae*. ChIP-seq and

electrophoretic mobility shift assays (EMSAs) were performed as described previously [39] using

probes generated by PCR with 6-Fam labelled primers from Integrated DNA Technologies

(Table S8). ChIP-seq Gels were visualised using a Typhoon FLA 9500 laser scanner (GE

Healthcare) with LBP/BPB1 emission filter, Exmax 495nm Emmax 576nm, at 50µM resolution.

ChIP-seq data was analysed as described previously [40] and peaks were visually inspected

using integrated genome browser [41]. Binding sites were identified using MEME [19].

RNA-sequencing. Duplicate wild-type and Δ*mtrB* cultures were grown for 14 and 20 hours in

MYM medium and RNA was prepared as described previously [42]. Libraries were prepared and

sequenced by Vertis Biotechnologie and analysed using CLC Genomics Workbench.

LCMS and PCA.

Analytical HPLC was performed on an HPLC 1100 system (Agilent Technologies) connected

to a Gemini® 3 μm NX-C18 110 Å, 150×4.6 mm column (Phenomenex) and on a Synergi™

4 μm Fusion-RP 80 Å LC column 250×10 mm. UPLC-MS for metabolic profiling was performed

on a Nexera X2 liquid chromatograph (LC-30AD) LCMS system (Shimadzu) connected to an

autosampler (SIL-30AC), a Prominence column oven (CTO-20AC) and a Prominence photo

diode array detector (SPD-M20A). A Kinetex® 2.6 μm C18 100 Å, 100×2.1 mm column

(Phenomenex) was used. The UPLC-System was connected with a LCMS-IT-TOF Liquid

Chromatograph mass spectrometer (Shimadzu). UPLC-HRMS Data was acquired on an

Acquity UPLC system (Waters Corporation) equipped with an ACQUITY UPLC® BEH 1.7 μm

C18, 1.0 × 100 mm column (Waters Corporation) and connected to a Synapt G2-Si high

resolution mass spectrometer (Waters Corporation). A gradient between mobile phase A (H_2O

with 0.1 % formic acid) and mobile phase B (acetonitrile with 0.1 % formic acid) at a flow rate

of 80 μL/min was used. Initial conditions were 1 % B for 1 min, ramped to 90 % B within

6 minutes, ramped to 100 % B within 0.5 min, held for 0.5 min, returned to 1 % B within 0.1 min

and held for 1.9 min. MS spectra were acquired with a scan time of one second in the range

of m/z = 150 - 1200 in positive MSe-Resolution mode. The following instrument parameters

were used: capillary voltage of 3 kV, sampling Cone 40, source offset: 80, source temperature

of 120 °C, desolvation temperature 350 °C, desolvation gas flow 800 L/h. A solution of sodium

formate was used for calibration. A solution of leucine encephalin (H_2O/MeOH/formic acid:

49.95/49.95/0.1) was used as lock mass and was injected every 15 sec. The lock mass has

been acquired with a scan time of 0.3 sec and 3 scans were averaged each time. The lock

mass (m/z = 556.2766) has been applied during data acquisition.For processing

metabolomics data we used the Software Progenesis QI (Waters). All solvents for analytical

HPLC and UPLC-MS were obtained commercially at least in HPLC grade from Fisher

Scientific and were filtered prior to use. Formic acid (EZInfo 3.0 (MKS Umetrics AB) was used

for plotting PCA data.

References

1.   Katz, L. & Baltz, R. H. Natural product discovery: past, present, and future. *J Ind*

Microbiol Biotechnol **43,** 1–22 (2016). doi:10.1007/s10295-015-1723-5

2.   Hosaka, T. *et al.* Antibacterial discovery in actinomycetes strains with mutations in

RNA polymerase or ribosomal protein S12. *Nat Biotechnol* **27,** 462–464 (2009).

3.   Rutledge, P. J. & Challis, G. L. Discovery of microbial natural products by activation of

silent biosynthetic gene clusters. *Nat Rev Micro* **13,** 509–523 (2015).

4.   Hsiao, N.-H. & Kirby, R. Two-component signal transduction systems in Streptomyces

and related organisms studied using DNA comparative microarray analysis. *Antonie*

Van Leeuwenhoek **95,** 189–206 (2009).

5.   Salazar, M. E. & Laub, M. T. Temporal and evolutionary dynamics of two-component

signaling pathways. *Curr Opin Microbiol* **24,** 7–14 (2015).

6.   Capra, E. J. & Laub, M. T. Evolution of Two-Component Signal Transduction

Systems. *Annu Rev Microbiol* **66,** 325–347 (2012).

7.   Hutchings, M. I., Hong, H.-J. & Buttner, M. J. The vancomycin resistance VanRS two-

component signal transduction system of Streptomyces coelicolor. *Mol Microbiol* **59,**

923–935 (2006).

8.   Hoskisson, P. A. & Hutchings, M. I. MtrAB-LpqB: a conserved three-component

system in actinobacteria? *Trends Microbiol* **14,** 444–449 (2006).

9.   Zahrt, T. C. & Deretic, V. An essential two-component signal transduction system in

Mycobacterium tuberculosis. *J Bacteriol* **182,** 3832–3838 (2000).

10. Purushotham, G., Sarva, K. B., Blaszczyk, E., Rajagopalan, M. & Madiraju, M. V.

Mycobacterium tuberculosis oriC sequestration by MtrA response regulator. *Mol*

Microbiol **98,** 586–604 (2015).

11.   Plocinska, R. *et al.* Septal Localization of the Mycobacterium tuberculosis MtrB

Sensor Kinase Promotes MtrA Regulon Expression. *Journal of Biological Chemistry*

287, 23887–23899 (2012).

12.   Plocinska, R. *et al.* Mycobacterium tuberculosis MtrB Sensor Kinase Interactions with

FtsI and Wag31 Proteins Reveal a Role for MtrB Distinct from That Regulating MtrA

Activities. *J Bacteriol* **196,** 4120–4129 (2014).

13.   Brocker, M., Mack, C. & Bott, M. Target genes, consensus binding site, and role of

phosphorylation for the response regulator MtrA of Corynebacterium glutamicum. *J*

Bacteriol **193,** 1237–1249 (2011).

14.   Krämer, R. Osmosensing and osmosignaling in Corynebacterium glutamicum. *Amino*

Acids **37,** 487–497 (2009).

15.   Möker, N. *et al.* Deletion of the genes encoding the MtrA-MtrB two-component system

of Corynebacterium glutamicum has a strong influence on cell morphology, antibiotics

susceptibility and expression of genes involved in osmoprotection. *Mol Microbiol* **54,**

420–438 (2004).

16.   Bush, M. J., Tschowri, N., Schlimpert, S., Flärdh, K. & Buttner, M. J. c-di-GMP

signalling and the regulation of developmental transitions in streptomycetes. *Nature*

Reviews Microbiology, **13,** 1–12 (2015).

17.   Gust, B., Challis, G. L., Fowler, K., Kieser, T. & Chater, K. F. PCR-targeted

Streptomyces gene replacement identifies a protein domain needed for biosynthesis

of the sesquiterpene soil odor geosmin. *Proc Natl Acad Sci USA* **100,** 1541–1546

(2003).

18.   Bibb, M. J., Domonkos, Á., Chandra, G. & Buttner, M. J. Expression of the chaplin

and rodlin hydrophobic sheath proteins in Streptomyces venezuelae is controlled by

σ(BldN) and a cognate anti-sigma factor, RsbN. *Mol Microbiol* **84,** 1033–1049 (2012).

19.   Bailey, T. L., Johnson, J., Grant, C. E. & Noble, W. S. The MEME Suite. *Nucleic Acids*

Res **43,** W39–49 (2015).

20. Bassam, Al, M. M., Bibb, M. J., Bush, M. J., Chandra, G. & Buttner, M. J. Response

regulator heterodimer formation controls a key stage in Streptomyces development.

PLoS Genet **10,** e1004554 (2014).

21. Bush, M. J., Chandra, G., Bibb, M. J., Findlay, K. C. & Buttner, M. J. Genome-Wide

Chromatin Immunoprecipitation Sequencing Analysis Shows that WhiB Is a

Transcription Factor That Cocontrols Its Regulon with WhiA To Initiate Developmental

Cell Division in *Streptomyces*. *mBio* **7,** e00523–16 (2016).

22. Wolański, M., Jakimowicz, D. & Zakrzewska-Czerwińska, J. AdpA, key regulator for

morphological differentiation regulates bacterial chromosome replication. *Open*

Biology **2,** 120097 (2012).

23. Wolański, M. *et al.* The level of AdpA directly affects expression of developmental

genes in Streptomyces coelicolor. *J Bacteriol* **193,** 6358–6365 (2011).

24. Stapleton, M. R., Smith, L. J., Hunt, D. M., Buxton, R. S. & Green, J. Mycobacterium

tuberculosis WhiB1 represses transcription of the essential chaperonin GroEL2.

Tuberculosis (Edinb) **92,** 328–332 (2012).

25. Schwedock, J., McCormick, J. R., Angert, E. R., Nodwell, J. R. & Losick, R. Assembly

of the cell division protein FtsZ into ladder-like structures in the aerial hyphae of

Streptomyces coelicolor. *Mol Microbiol* **25,** 847–858 (1997).

26. Willemse, J., Mommaas, A. M. & Wezel, G. P. Constitutive expression of ftsZ

overrides the whi developmental genes to initiate sporulation of Streptomyces

coelicolor. *Antonie Van Leeuwenhoek* **101,** 619–632 (2011).

27. Molle, V., Palframan, W. J., Findlay, K. C. & Buttner, M. J. WhiD and WhiB,

Homologous Proteins Required for Different Stages of Sporulation in Streptomyces

coelicolor A3(2). *J Bacteriol* **182,** 1286–1295 (2000).

28. Tschowri, N. *et al.* Tetrameric c-di-GMP Mediates Effective Transcription Factor

Dimerizationto Control Streptomyces Development. *Cell* **158,** 1136–1147 (2014).

29. Weber, T. *et al.* antiSMASH 3.0-a comprehensive resource for the genome mining of

biosynthetic gene clusters. *Nucleic Acids Res* **43,** W237–43 (2015).

30.   Fernández-Martínez, L. T. *et al.* New Insights into Chloramphenicol Biosynthesis in

493   Streptomyces venezuelae ATCC 10712. *Antimicrob Agents Chemother* **58,** 7441–

494   7450 (2014).

495   31.   Challis, G. L. Exploitation of the Streptomyces coelicolor A3(2) genome sequence for

496   discovery of new natural products and biosynthetic pathways. *J Ind Microbiol*

497   *Biotechnol*. **41,** 219-32 (2013).

498   32.   Labeda, D. P. *et al.* Phylogenetic study of the species within the family

499   Streptomycetaceae. *Antonie van Leeuwenhoek* **101,** 73-104 (2011).

500   33.   Bentley, S. D. *et al.* Complete genome sequence of the model actinomycete

501   Streptomyces coelicolor A3(2). *Nature* **417,** 141–147 (2002).

502   34.   van Keulen, G. & Dyson, P. J. In Advances in Applied Microbiology. Elsevier, pp.

503   217–266 (2014).

504   35.   Song, L. *et al.* Type III polyketide synthase beta-ketoacyl-ACP starter unit and

505   ethylmalonyl-CoA extender unit selectivity discovered by Streptomyces coelicolor

506   genome mining. *J Am Chem Soc* **128,** 14754–14755 (2006).

507   36.   Satsangi, A. T., Pandeeti, E. P., Sarva, K., Rajagopalan, M. & Madiraju, M. V.

508   Tuberculosis. *Tuberculosis (Edinb)* **93,** S28–S32 (2013).

509   37.   Kieser, T., Bibb, M. J., Buttner, M. J., Chater, K. F. & Hopwood, D. A. Practical

510   Streptomyces Genetics. Published by the John Innes Foundation. ISBN 0-7084-0623-

511   8 (2000).

512   38.   Hutchings, M. I., Hong, H.-J., Leibovitz, E., Sutcliffe, I. C. & Buttner, M. J. The

513   sigma(E) cell envelope stress response of Streptomyces coelicolor is influenced by a

514   novel lipoprotein, CseA. *J Bacteriol* **188,** 7222–7229 (2006).

515   39.   Crack, J. C. *et al.* NsrR from Streptomyces coelicoloris a Nitric Oxide-Sensing [4Fe-

516   4S] Cluster Protein with a Specialized Regulatory Function. *Journal of Biological*

517   *Chemistry* **290,** 12689-704 (2015).

518   40.   Bush, M. J., Bibb, M. J., Chandra, G., Findlay, K. C. & Buttner, M. J. Genes Required

519   for Aerial Growth, Cell Division, and Chromosome Segregation Are Targets of WhiA

before Sporulation in Streptomyces venezuelae. *mBio* **4,** e00684–13–e00684–13

(2013).

41.   Nicol, J. W., Helt, G. A., Blanchard, S. G., Raja, A. & Loraine, A. E. The Integrated

Genome Browser: free software for distribution and exploration of genome-scale

datasets. *Bioinformatics* **25,** 2730–2731 (2009).

42.   Munnoch, J. T. *et al.* Characterization of a putative NsrR homologue in. *Sci. Rep.* **6,**

1–14 (2016).

Acknowledgements

We thank Kim Findlay for the Scanning Electron Microscopy, Mark Buttner for the gift of *S.*

venezuelae NRRL B-65442, cosmid Sv-6-A04, access to microarray data prior to publication

and helpful discussions, Matt Bush for critical reading of the manuscript and helpful

discussions and Elaine Patrick for technical support. This research was supported by a

BBSRC PhD studentship to NS, a UEA-funded PhD studentship to FK, MRC Milstein award

G0801721 to MIH, Norwich Research Park translational funding, NPRONET Proof of Concept

funding and Natural Environment Research Council responsive mode grants NE/M015033/1

and NE/M014657/1 to MIH and BW and NE/M001415/1 to PAH.

Author contributions

FK and RFS constructed the *S. coelicolor* mutants and FK undertook phenotype analysis, NS

constructed all the *S. venezuelae* strains and constructs, analysed phenotypes and performed

ChIP-seq experiments, JTM performed RNA-seq and dRNA-seq experiments, JTM and GC

analysed the ChIP- and RNA-seq datasets, NH made constructs and prepared strain extracts

for LCMS, DH purified and analysed all natural products including LCMS and PCA analyses,

PAH, MIH and BW conceived the study and all the authors analysed data and wrote the

manuscript.