1

The NCA-1 ion channel functions downstream of G_q and Rho

to regulate locomotion in *C. elegans*

Irini Topalidou[*], Pin-An Chen[†], Kirsten Cooper[*,1], Shigeki Watanabe[†,2],

Erik M. Jorgensen[†], Michael Ailion[*]

[*]Department of Biochemistry, University of Washington, Seattle, WA 98195

[†]Howard Hughes Medical Institute, Department of Biology, University of Utah, Salt Lake City, UT 84112

[1]Current Address: Fred Hutchinson Cancer Research Center, Seattle, WA 98109

[2]Current Address: Department of Cell Biology, Johns Hopkins University, Baltimore, MD 21205

Running Title: Gq signaling activates NCA-1 through Rho

key words: G_q signaling, Rho small GTPase, NCA/NALCN ion channel, *C. elegans*, G protein

Corresponding author:

Michael Ailion

Department of Biochemistry

University of Washington

Box 357350

1705 NE Pacific St

Seattle, WA 98195

Phone: 206-685-0111

email: mailion@uw.edu

Abstract

The heterotrimeric G protein G_q positively regulates neuronal activity and synaptic transmission. Previously, the Rho guanine nucleotide exchange factor Trio was identified as a direct effector of G_q that acts in parallel to the canonical G_q effector phospholipase C. Here we examine how Trio and Rho act to stimulate neuronal activity downstream of G_q in the nematode *Caenorhabditis elegans*. Through two forward genetic screens, we identify the cation channel NCA-1/NALCN as a downstream target of the G_q/Rho pathway. By performing genetic epistasis analysis using dominant activating mutations and recessive loss-of-function mutations in the members of this pathway, we show that NCA-1 acts downstream of G_q in a linear pathway. Through cell-specific rescue experiments, we show that function of this channel in head acetylcholine neurons is sufficient for normal locomotion in *C. elegans*. Our results suggest that NCA-1 is a physiologically relevant target of neuronal G_q/Rho signaling in *C. elegans*.

Introduction

Heterotrimeric G protein pathways play central roles in altering neuronal activity and

synaptic transmission in response to experience or changes in the environment. G_q is one of the

four types of heterotrimeric G proteins alpha subunits in animals (Wilkie *et al.* 1992) and is widely

expressed in the mammalian brain (Wilkie *et al.* 1991) where it typically acts to stimulate neuronal

activity and synaptic transmission (Krause *et al.* 2002; Gamper *et al.* 2004; Coulon *et al.* 2010).

These roles are conserved in the nematode *C. elegans*. Unlike mammals which have four

members of the G_q family, *C. elegans* has only a single $G_q\alpha$ (Brundage *et al.* 1996). In *C. elegans*,

loss-of-function and gain-of-function mutants in the single $G_q\alpha$ gene *egl-30* are viable but have

strong neuronal phenotypes, affecting locomotion, egg-laying, and sensory behaviors (Brundage

et al. 1996; Lackner *et al.* 1999; Bastiani *et al.* 2003; Matsuki *et al.* 2006; Esposito *et al.* 2010;

Adachi *et al.* 2010). We aim to identify the signal transduction pathways and downstream targets

by which G_q signaling alters neuronal activity.

In the canonical G_q pathway, G_q activates phospholipase Cβ (PLC) to cleave the lipid

phosphatidylinositol 4,5,-bisphosphate (PIP2) into diacylglycerol (DAG) and inositol trisphosphate

(IP3), each of which can act as a second messenger. This pathway operates in both worms and

mammals, but in both systems, a number of PLC-independent effects of G_q have been described,

indicating the existence of additional G_q signal transduction pathways (Lackner *et al.* 1999; Miller

et al. 1999; Vogt *et al.* 2003; Bastiani *et al.* 2003; Sánchez-Fernández *et al.* 2014). Using a

genetic screen for suppressors of activated G_q, we identified the Rho guanine nucleotide

exchange factor (GEF) Trio as a direct effector of G_q in a second major G_q signal transduction

pathway independent of the canonical PLC pathway (Williams *et al.* 2007). Biochemical and

structural studies demonstrated that G_q directly binds and activates RhoGEF proteins in both

worms and mammals, indicating that this new G_q pathway is conserved (Lutz *et al.* 2005; Williams

et al. 2007; Lutz *et al.* 2007).

G_q activation of the Trio RhoGEF leads to activation of the small GTP-binding protein Rho,

a major cellular switch that affects a number of cellular processes, ranging from regulation of the

cytoskeleton to transcription (Etienne-Manneville and Hall 2002; Jaffe and Hall 2005). In *C.*

elegans neurons, Rho has been shown to regulate synaptic transmission downstream of the G_{12}-

class G protein GPA-12 via at least two pathways, one dependent on the diacylglycerol kinase

DGK-1 and one independent of DGK-1 (McMullan *et al.* 2006; Hiley *et al.* 2006). Here we

investigate what targets operate downstream of Rho in the G_q signaling pathway to regulate

neuronal activity. Through two forward genetic screens, we identify the cation channel NCA-1

(NALCN in mammals) as a downstream target of the G_q/Rho pathway. The NCA-1/NALCN

channel is a relative of voltage-gated cation channels that has been suggested to be a sodium

leak channel required for the propagation of neuronal excitation and the fidelity of synaptic

transmission (Lu *et al.* 2007; Jospin *et al.* 2007; Yeh *et al.* 2008). However, there is controversy

over the current carried by NALCN and whether it is indeed a sodium leak channel (Senatore *et*

al. 2013; Senatore and Spafford 2013; Boone *et al.* 2014). It is also unclear how NALCN is gated

and what pathways activate the channel. Two studies have shown that NALCN-dependent

currents can be activated by G protein-coupled receptors, albeit independently of G proteins (Lu *et*

al. 2009; Swayne *et al.* 2009), and another study showed that the NALCN leak current can be

activated by low extracellular calcium via a G protein-dependent pathway (Lu *et al.* 2010). Our

data presented here suggest that NCA-1 can be activated by Rho acting downstream of G_q in a

linear pathway.

Materials and Methods

Strains

Worm strains were cultured and maintained using standard methods (Brenner, 1974). A complete list of strains and mutations used is provided in the strain list (Table S1).

Isolation of suppressors of activated G_q

We performed an ENU mutagenesis to screen for suppressors of the hyperactive locomotion of an activated G_q mutant, *egl-30(tg26)* (Ailion *et al.* 2014). From approximately 47,000 mutagenized haploid genomes, we isolated 10 mutants that had a fainter phenotype when outcrossed away from the *egl-30(tg26)* mutation. By mapping and complementation testing, we assigned these ten mutants to three genes: three *unc-79* mutants (*yak37*, *yak61*, and *yak73*), six *unc-80* mutants (*ox329*, *ox330*, *yak8*, *yak35*, *yak36*, and *yak56*) and one *nlf-1* mutant (*ox327*).

Isolation of suppressors of activated G_o

We first isolated suppressors of the activated G_o mutant *unc-109(n499)* by building double mutants of *unc-109(n499)* with the activated G_q allele *egl-30(tg26)*. Unlike *unc-109(n499)* homozygotes which are lethal, *egl-30(tg26) unc-109(n499)* homozygotes are viable, but paralyzed and sterile, indicating that activated G_q partially suppresses activated G_o, consistent with the model that these two G proteins act antagonistically. We built a balanced heterozygote strain *egl-30(tg26) unc-109(n499)/egl-30(tg26) unc-13(e51) gld-1(q126)* which has an "unmotivated" phenotype in which the worms move infrequently and slowly. We mutagenized these animals with ENU and screened for F1 progeny that moved better. From a screen of approximately 16,000 mutagenized haploid genomes, we isolated two apparent *unc-109* intragenic mutants, *ox303* and *ox304*. *ox303* is a strong *unc-109* loss-of-function allele, as evidenced by the fact that *egl-30(tg26) unc-109(n499 ox303)/egl-30(tg26) unc-13(e51) gld-1(q126)* mutants resembled *egl-30(tg26)*

mutants (i.e. hyperactive). Additionally, *unc-109(n499 ox303)* mutants themselves are hyperactive

in an otherwise wild-type background. *ox304*, however, appears to be a partial loss-of-function

mutant, because the *egl-30(tg26) unc-109(n499 ox304)/egl-30(tg26) unc-13(e51) gld-1(q126)*

mutant moves better than the *egl-30(tg26) unc-109(n499)/egl-30(tg26) unc-13(e51) gld-1(q126)*

parent strain, but is not hyperactive like the *egl-30(tg26)* strain. Also, *unc-109(n499 ox304)*

homozygote animals on their own are viable but are almost paralyzed and show very little

spontaneous movement, with a typical straight posture. However, when stimulated by transfer to a

new plate, the *unc-109(n499 ox304)* mutant is surprisingly capable of coordinated movements.

This strain was used for mapping and sequencing experiments that demonstrated that *unc-109* is

allelic to *goa-1*, encoding the worm G_o ortholog (see below). The *unc-109(n499 ox304)* strain was

also used as the starting point for a second screen to isolate extragenic suppressors of activated

G_o.

        Previously, a screen for suppressors of activated *goa-1* was performed using heat-shock

induced expression of an activated *goa-1* transgene (Hajdu-Cronin *et al.* 1999). This screen

isolated many alleles of *dgk-1*, encoding diacyclglycerol kinase, but only a single allele of *eat-16*,

encoding a regulator of G protein signaling (RGS) protein that negatively regulates G_q (Hajdu-

Cronin *et al.* 1999), along with mutants in three genes needed for expression of the heat-shock

goa-1 transgene (Hajdu-Cronin *et al.* 2004). Because of the strong bias of this screen for isolating

alleles of *dgk-1*, we used the *goa-1(n499 ox304)* strain to perform a screen for suppressors of

activated G_o that did not involve overexpression of *goa-1* or rely on expression of a heat-shocked

induced *goa-1* transgene. We performed ENU mutagenesis of *goa-1(n499 ox304)* and isolated F2

animals that were not paralyzed. From a screen of approximately 24,000 mutagenized haploid

genomes, we isolated 17 suppressors, 9 with a relatively stronger suppression phenotype and 8

that were weaker. Of the 9 stronger suppressors, we isolated two alleles of *eat-16* (*ox359*, *ox360*),

three alleles of the BK type potassium channel *slo-1* (*ox357*, *ox358*, *ox368*), one allele of the

innexin *unc-9* (*ox353*), one gain-of-function allele in the ion channel gene *nca-1* (*ox352*), and two

141 mutants that were mapped to chromosomal regions distinct from the other mutations listed above,

142 but not further characterized (*ox356*, *ox364*).

Mapping and cloning *unc-109(n499)*

unc-109 was shown to be allelic to *goa-1*. First, we performed SNP mapping of both *unc-*

109(n499) and its intragenic revertant *unc-109(n499 ox303)*, using the Hawaiian strain CB4856 as

described (Davis *et al.* 2005). These experiments mapped *unc-109* to an approximately 1 Mb

region in the middle of chromosome I between SNPs on cosmids D2092 and T24B1 (SNPs CE1-

15 and T24B1[1]). A good candidate in this region was *goa-1*. We confirmed that *unc-109* was

indeed *goa-1* by sequencing the three *unc-109* mutants: the gain-of-function allele *n499* carries a

point mutation that leads to an R179C missense mutation, affecting a conserved arginine residue

shown to be important for the GTPase activity of G proteins (Coleman *et al.* 1994); the partial loss-

of-function allele *ox304* carries a point mutation leading to a W259R missense mutation; and the

strong loss-of-function allele *ox303* carries a one basepair deletion that leads to a premature stop

32 amino acids from the C-terminal.

Mapping and cloning *nlf-1(ox327)*

We mapped the *ox327* mutation using single nucleotide polymorphisms (SNPs) in the

Hawaiian strain CB4856 as described (Davis et al., 2005). The *ox327* mutation was mapped to an

approximately 459 kb region on the left arm of the X chromosome between SNPs on cosmids

F39H12 and C52B11 (SNPs F39H12[4] and pkP6101). This region included 74 predicted protein-

coding genes. We injected cosmids spanning this region and found that injection of cosmid F55A4

rescued the *ox327* mutant phenotype. We performed RNAi to the genes on this cosmid in the *eri-*

1(mg366) lin-15(n744) strain that has enhanced RNAi and found that RNAi of the gene F55A4.2

caused a weak fainter phenotype. We sequenced F55A4.2 in the *ox327* mutant and found a T to

A transversion mutation in exon 1, leading to a premature stop codon at amino acid C59. We also

rescued the *ox327* mutant with a transgene carrying only F55A4.2, confirming the gene

identification. We subsequently obtained a deletion allele *tm3631* that has fainter and G_q

suppression phenotypes indistinguishable from *ox327*. F55A4.2 was given the gene name *nlf-1*

(Xie *et al.* 2013).

We obtained six independent *nlf-1* cDNAs that were predicted to be full-length: yk1105g4,

yk1159a2, yk1188d11, yk1279a1, yk1521f8, and yk1709b10. Restriction digests suggested that

all six were of the same size. We sequenced yk1159a2 and yk1279a1 and both gave the same

nlf-1 exon-intron structure, which differed from the gene-structure on Wormbase WS253 in several

ways: *nlf-1* is 4 bp shorter at the 3' end of exon 5, and has a new 154 bp exon (now exon 6) not

predicted on Wormbase. *nlf-1* consists of 8 exons and is predicted to encode a protein of 438

amino acids. This is identical to the gene structure reported independently (Xie *et al.* 2013). Both

sequenced cDNAs had 5'UTRs of 64 bp, with yk1159a2 (but not yk1279a1) being trans-spliced to

the SL1 splice leader, and 3'UTRs of 424 bp (yk1279a1) or 429 bp (yk1159a2). The yk1279a1

cDNA was mutation-free and was cloned into a Gateway entry vector for use in rescue

experiments. The full-length sequence of the yk1279a1 *nlf-1* cDNA was deposited in GenBank

under accession # KX808524.

Molecular biology and transgenes

A complete list of constructs is provided in the plasmid list (Table S2). Most of the

constructs were made using the three slot multisite Gateway system (Invitrogen). For C3

transferase constructs, a promoter, an FRT-mCherry-FRT-GFP cassette (pWD178), and the C3

transferase-*unc-54* 3'UTR (cloned into a Gateway entry vector from plasmid QT#99) were

combined into the pDEST R4-R3 destination vector. For *nlf-1* tissue-specific rescue constructs, a

promoter, the *nlf-1* coding sequence (genomic DNA or cDNA), and a C-terminal GFP tag were

cloned along with the *unc-54* 3'UTR into the pDEST R4-R3 destination vector. Promoters used

were *nlf-1p* (5.7 kb upstream of the ATG), *rab-3p* (all neurons), *unc-17p* (acetylcholine neurons),

unc-17Hp (head acetylcholine neurons) (Hammarlund *et al.* 2007), *acr-2p* (acetylcholine motor

neurons), *unc-17βp* (acetylcholine motor neurons) (Charlie *et al.* 2006), and *glr-1p* (glutamate-

receptor interneurons). Extrachromosomal arrays were made by standard transformation methods

(Mello *et al.* 1991). Constructs of interest were injected at 10 ng/μl with marker and carrier DNAs

added to make a final total concentration of at least 100 ng/μl. For most constructs, we isolated

multiple independent insertions that behaved similarly. C3 transferase extrachromosomal arrays

were integrated into the genome using X-ray irradiation (4000 rads). Integrated transgenes were

mapped to chromosomes and outcrossed twice before further analysis.

Locomotion assays

We performed two different assays to measure locomotion. Body bend assays measured

the rate of locomotion. Radial locomotion assays measured the radial distance animals moved

from a point in a given unit of time, which provides a combined measurement of different aspects

of the locomotion phenotype including the rate of locomotion, waveform, and frequency of

reversals. Both types of assays were performed on 10 cm plates seeded with thin lawns of OP50

bacteria. These plates were prepared by seeding 1.5 ml of stationary phase OP50 bacteria to

evenly cover the entire plate surface, and then growing the bacteria for two days at room

temperature. Plates were stored at 4° for up to one month before being used. For body bend

assays, first-day adult worms were picked to an assay plate, allowed to rest for 30 seconds, and

then body bends were counted for one minute. A body bend was defined as the movement of the

worm from maximum to minimum amplitude of the sine wave (Miller *et al.* 1999). To minimize

variation, all animals in a body bend experiment were assayed on the same plate. For radial

locomotion assays, five to eight first-day adults were picked together to the center of a plate to

begin the assay (time 0). Positions of the worms were marked on the lid of the plate every ten

minutes for up to forty minutes. Following the assay, the distance of each point to the center was

measured. For most strains, radial distances did not increase after the first ten minutes of the assay and all data presented here are for the ten-minute time point. Analysis of the data at later time points leads to the same conclusions. For all locomotion assays, the experimenter was blind to the genotypes of the strains being assayed.

For Rho inhibition experiments (Figure 1), expression of C3 transferase (C3T) was induced by FLP-mediated recombination. Expression of FLP was induced by heat shock for 1 hr at 34°C, plates were returned to room temperature, and animals were scored for locomotion 4 hrs after the end of the heat shock period. For heat-shock induction of activated Rho (Figure S5), worms were heat shocked for 1 hr at 34°C, returned to room temperature, and scored for locomotion 2 hrs after the end of the heat-shock period.

Fainting assays

Backward fainting times were measured by touching a worm on the head with a worm pick to stimulate movement, and measuring the time to faint with a stopwatch. Forward fainting time was measured following a touch on the tail. Fainting was defined by an abrupt stop of movement along with a characteristic straightening of the head (see Figure 5). Alternatively, we touched worms on the head or tail with a pick and counted the number of body bends until the worm fainted. If a worm moved 10 body bends without fainting, we stopped the assay.

Imaging and image analysis

Worms were mounted on 2% agarose pads and anesthetized with sodium azide. Images were obtained using a Zeiss Pascal confocal microscope. For quantitative imaging of NCA-1::GFP and NCA-2::GFP (Figure S2), Z-stack projections of the nerve ring axons on one side of the animal were collected and quantified in ImageJ as described (Jospin *et al.* 2007). Dissecting microscope photographs of first-day adult worms were taken at 50X using a Nikon SMZ18 microscope equipped with a DS-L3 camera control system.

Statistics

P values were determined using GraphPad Prism 5.0d (GraphPad Software). Normally

distributed data sets with multiple comparisons were analyzed by a one-way ANOVA followed by

a Bonferroni or Tukey posthoc test to examine selected comparisons or by Dunnett's test if all

comparisons were to the wild type control. Non-normally distributed data sets with multiple

comparisons were analyzed by a Kruskal-Wallis nonparametric ANOVA followed by Dunn's test to

examine selected comparisons. Pairwise data comparisons were analyzed by a two-tailed

unpaired t test for normally distributed data or by a two-tailed Mann-Whitney test for non-normally

distributed data.

Reagent and data availability

Strains and plasmids are shown in Table S1 and Table S2 and are available from the

Caenorhabditis Genetics Center (CGC) or upon request. The full-length sequence of the

yk1279a1 *nlf-1* cDNA was deposited in GenBank under accession # KX808524. The authors state

that all data necessary for confirming the conclusions presented in the article are represented fully

within the article and Supplemental Material.

Results

Inhibition of Rho suppresses activated G_q

Two pieces of data suggested that G_q may regulate locomotion in *C. elegans* through activation of the small G protein Rho. First, both the hyperactive locomotion and tightly wound body posture of the activated G_q mutant *egl-30(tg26)* are suppressed by loss-of-function mutations in the *unc-73* RhoGEF Trio, an activator of Rho (Williams *et al.* 2007). Second, expression of activated Rho as a transgene causes worms to adopt a posture characterized by a tightly coiled, high-amplitude waveform (McMullan *et al.* 2006), reminiscent of the waveform of activated G_q worms (Bastiani *et al.* 2003; Ailion *et al.* 2014). To determine directly whether G_q signals through Rho, we tested whether Rho inhibition suppresses an activated G_q mutant.

In *C. elegans,* there is a single gene encoding Rho (*rho-1*). Loss of *rho-1* causes numerous pleiotropic developmental phenotypes and embryonic lethality (Jantsch-Plunger *et al.* 2000). Thus, to bypass the developmental roles and study Rho function in the adult nervous system, we expressed the Rho inhibitor C3 transferase (C3T) only in adult neurons by using the FLP recombinase/FRT system (Davis *et al.* 2008). In this system, temporal control of C3T is achieved through induction of FLP via heat-shock. FLP in turn promotes recombination between FRT sites to lead to expression of C3 transferase in specific neurons (Figure 1A). We expressed C3T in the following classes of neurons: all neurons (*rab-3p*), acetylcholine neurons *(unc-17p)*, head acetylcholine neurons (*unc-17Hp*), and acetylcholine motor neurons (*unc-17βp*). mCherry fluorescence confirmed expression in the expected neurons and GFP expression was used to monitor induction of C3T following FLP-mediated recombination.

Inhibition of *rho-1* in adult neurons caused a decreased locomotion rate (Figure 1B). This effect was greatest when Rho was inhibited in all neurons, but Rho inhibition in acetylcholine subclasses of neurons also led to slower locomotion. Thus, Rho acts in multiple classes of neurons to promote locomotion in adult worms. In the absence of heat-shock, all strains showed

normal wild-type rates of locomotion and did not express any detectable GFP, indicating that

these transgenes do not provide leaky expression of C3T in the absence of heat-shock. This

confirms that Rho acts post-developmentally in mature neurons to regulate locomotion behavior

(McMullan *et al.* 2006).

To determine whether Rho acts downstream of G_q signaling, we crossed the C3 transgenes

into the background of the activated G_q mutant *egl-30(tg26)*. Inhibition of *rho-1* in all adult neurons

suppressed both the posture and hyperactive locomotion of the activated G_q mutant (Figure

1C,D). Inhibition of *rho-1* in acetylcholine neurons suppressed the hyperactivity of activated G_q

(Figure 1C), but the worms still had a coiled, high-amplitude waveform (Figure 1D). Thus, *rho-1*

exhibits genetic interactions consistent with a role in the G_q signaling pathway.

Mutations in NCA channel subunits suppress activated G_q

What acts downstream of Rho in the G_q signal transduction pathway? We screened for

suppressors of activated Gq and found mutants in three categories: (1) the canonical G_q pathway

(such as the PLC *egl-8*); (2) the RhoGEF *unc-73* (Williams *et al.* 2007), and (3) genes that affect

dense-core vesicle function (*e.g. unc-31*, *rab-2*, *rund-1*) (Ailion *et al.* 2014; Topalidou *et al.* 2016).

Mutations in *unc-73* strongly suppress the high-amplitude waveform of the activated G_q mutant,

but mutations in the canonical G_q pathway or in genes that affect dense-core vesicle function only

weakly suppress the high-amplitude waveform of the activated G_q mutant (Figure S1). Thus,

strong suppression of the high-amplitude waveform of activated G_q may be a specific

characteristic of mutations in the Rho pathway, and we hypothesized that downstream targets of

Rho in this pathway would also suppress the high-amplitude waveform.

To identify possible downstream targets of Rho in the G_q pathway, we examined other

mutants isolated from our screen for suppression of the high-amplitude waveform of activated G_q

(Figure S1). When crossed away from the activating G_q mutation, several of these mutants had a

"fainter" phenotype. Fainter mutants respond to a touch stimulus by moving away, but abruptly

stop, that is "faint", after only a few body bends. The fainter phenotype has been observed only in

mutants that reduce the function of the NCA-1 and NCA-2 ion channels (Humphrey *et al.* 2007;

Jospin *et al.* 2007; Yeh *et al.* 2008). We found that all our G_q suppressors that strongly

suppressed the high-amplitude waveform and had a fainter phenotype were mutants in either *unc-*

79 or *unc-80*, two genes required for function of the NCA channels (Humphrey *et al.* 2007; Jospin

et al. 2007; Yeh *et al.* 2008). We also isolated a single mutant in the gene *nlf-1* that only weakly

suppressed the high-amplitude waveform of activated G_q (Figure S1), but also gave fainters after

outcrossing away from the activated G_q mutation. *unc-79* and *unc-80* mutants have a strong

fainter phenotype, equivalent to that of a double mutant in *nca-1* and *nca-2*, two genes that

encode pore-forming subunits of the NCA channels in *C. elegans* (Humphrey *et al.* 2007; Jospin

et al. 2007; Yeh *et al.* 2008). Like *unc-79* or *unc-80*, an *nca-1 nca-2* double mutant suppressed the

activated G_q mutant. Additionally, *nca-1* on its own suppressed activated G_q, but *nca-2* did not.

This suggests that although *nca-1* and *nca-2* are only redundantly required for normal worm

locomotion, G_q activation may be specific to channels containing the NCA-1 pore-forming subunit.

Cloning and characterization of *nlf-1(ox327)*

In addition to the previously known NCA channel subunits *unc-79* and *unc-80*, we also

isolated the *ox327* mutant in a gene that had not been previously characterized at the time of our

study. We cloned *ox327* by single nucleotide polymorphism (SNP) mapping, RNAi and transgenic

rescue experiments (see Materials and Methods), showing that it carries an early stop mutation in

the gene *nlf-1*. The *nlf-1(ox327)* and *nlf-1(tm3631)* deletion mutants failed to complement each

other and had indistinguishable fainting and G_q suppression phenotypes. *nlf-1* was independently

identified by others and encodes an endoplasmic reticulum-localized protein probably involved in

proper assembly of the NCA channel, since *nlf-1* mutants had reduced expression levels of GFP-

tagged NCA-1 and NCA-2 (Xie *et al.* 2013). We also found that *nlf-1* is required for normal axonal

levels of both GFP-tagged NCA-1 and NCA-2 in the nerve ring (Figure S2). Additionally, an *nlf-1*

mutation strongly suppressed both the slow locomotion and coiled posture of an activated *nca-1*

mutant (Figure 2), demonstrating that *nlf-1* is important for NCA-1 function.

nlf-1 mutants have a weaker fainter phenotype than mutants of *unc-79* or *unc-80*, or the

nca-1 nca-2 double mutant. The *nlf-1* fainting phenotype differs in two ways from those of the

stronger fainting mutants. First, *nlf-1* mutants take a longer time to faint following stimulation

(Figure 3). Second, while the strong fainting mutants show a similarly strong fainting phenotype in

either the forward or backward direction, *nlf-1* mutants faint reliably in the backward direction but

take much longer and have more variable fainting in the forward direction (Figure 3), suggesting

that *nlf-1* mutations cause a partial loss of function of the NCA channels. Additionally, *nlf-1*

mutants have greater spontaneous locomotion activity than *unc-79*, *unc-80* or *nca-1 nca-2* double

mutants. To determine how *nlf-1* interacts with NCA mutants, we built *nlf-1(ox327)* double mutants

with *unc-79, unc-80*, *nca-1* and *nca-2*. The *nlf-1* mutation did not enhance the *unc-79* or *unc-80*

fainter phenotype, suggesting that these mutants act in the same pathway to control fainting.

However, *nca-1* strongly enhanced the *nlf-1* fainter phenotype, while *nca-2* had a weak effect on

nlf-1 fainting that was not statistically significant (Figure 3). Neither *nca-1* nor *nca-2* single mutants

have a fainter phenotype on their own, but the fact that an *nca-1 nlf-1* double mutant has a strong

fainter phenotype suggests that either *nca-1* contributes more than *nca-2* for normal locomotion or

that *nlf-1* specifically perturbs function of *nca-2*. Our data presented below are more consistent

with the possibility that *nca-1* contributes more than *nca-2* to wildtype locomotion behavior.

We determined the cellular expression pattern of *nlf-1* by fusing its promoter to GFP. *nlf-1p::GFP* was expressed in most or all neurons, but was not detected in other tissues (Figure S3).

This agrees with the expression pattern reported elsewhere (Xie *et al.* 2013). To determine the

neuronal focus of the fainter phenotype, we performed rescue experiments in which we

determined whether an *nlf-1* mutant could be rescued by expression of a wild-type *nlf-1* cDNA

under the control of neuron-specific promoters. Expression of *nlf-1(+)* in all neurons (using the *rab-3* promoter) or acetylcholine neurons (using the *unc-17* promoter) fully rescued the *nlf-1* mutant

fainter phenotype (Figure 4A). Expression in acetylcholine motor neurons (using the *acr-2* or *unc-17β* promoters) did not rescue the fainter phenotype, but expression driven by a head-specific derivative of the *unc-17* promoter (*unc-17Hp*) fully rescued the fainter phenotype, indicating that the action of *nlf-1* in head acetylcholine neurons is sufficient to prevent fainting (Figure 4A).

Previously, it was reported that expression of *nlf-1* in premotor interneurons is sufficient to rescue the *nlf-1* mutant fainter phenotype (Xie *et al.* 2013). However, we found that expression of *nlf-1* in premotor interneurons using the *glr-1* promoter (*glr-1p*) only partially rescued the *nlf-1* mutant fainter phenotype and did not fully restore wild-type locomotion (Figure 4B). Fainting behavior was difficult to score in the *glr-1* promoter-rescued *nlf-1* mutant animals because they had sluggish movement and stopped frequently, though generally not with the suddenness and characteristic posture typical of fainters. Though we could blindly score the rescue of fainting in these animals by eye, we saw only weak rescue of the *nlf-1* mutant by *glr-1* promoter expression in our quantitative fainting assays and the effect was not statistically significant (Figure 4B). When we instead measured fainting as the percentage of animals that fainted within ten body bends, we did see a marginally significant rescue by *glr-1* promoter expression (backward fainting: *nlf-1* = 92%, *nlf-1; glr-1p::nlf-1(+)* = 68%, P=0.0738, Fisher's exact test; forward fainting: *nlf-1* = 80%, *nlf-1; glr-1p::nlf-1(+)* = 48%, P=0.0378, Fisher's exact test). We may be underestimating the rescue of the fainting phenotype by *glr-1* promoter expression due to the difficulty distinguishing the frequent pausing from true fainting. Nevertheless, rescue of the *nlf-1* mutant is clearly stronger by expression in head acetylcholine neurons using the *unc-17H* promoter (Figure 4A). Though our data seem to contradict the previous study reporting that *nlf-1* acts in premotor interneurons (Xie *et al.* 2013), there are several possible explanations. First, like our data, the data in the previous study in fact showed only partial rescue of fainting behavior by expression in premotor interneurons (Xie *et al.* 2013). Second, the premotor interneuron promoter combination used in the previous study (*nmr-1p* + *sra-11p*) leads to expression in several other head interneurons that may contribute to the phenotype. Third, it is possible that rescue is sensitive to expression level

391 and that different levels of expression were achieved in the two studies, leading to different levels

392 of rescue. We conclude that NLF-1 acts in head acetylcholine neurons, including the premotor

393 interneurons, to promote sustained locomotion in the worm. Consistent with this, the premotor

394 command interneurons have recently been shown to have acetylcholine (Pereira *et al.* 2015).

Mutations in NCA channel subunits suppress activated Rho

To determine whether NCA mutants act downstream of Rho, we took advantage of an

activated Rho mutant (G14V) expressed specifically in the acetylcholine neurons (McMullan *et al.*

2006). Like an activated G_q mutant, this activated Rho mutant has an exaggerated waveform. We

built double mutants of the activated Rho mutant with mutations in *unc-79*, *unc-80*, *nlf-1* and

mutations in the NCA channel subunit genes *nca-1* and *nca-2*. The *nca-1* and *nca-2* genes are

redundant for the fainter phenotype because neither mutant has a fainter phenotype individually,

but the double mutant has a fainter phenotype indistinguishable from *unc-79* and *unc-80* mutants

(Humphrey *et al.* 2007; Jospin *et al.* 2007; Yeh *et al.* 2008). We found that mutations in *unc-79* or

unc-80 strongly suppressed the high-amplitude waveform of activated Rho, as did an *nca-1 nca-2*

double mutant (Figure 5). In all cases, the resulting double (or triple) mutants had a fainter

phenotype like the single mutants. By contrast, a mutation in *nlf-1* strongly but incompletely

suppressed the high-amplitude waveform of activated Rho, consistent with the *nlf-1* mutation

causing a partial but incomplete loss of NCA channel function. Unlike an *nlf-1* single mutant, an

nlf-1 double mutant with activated Rho did not have a fainter phenotype. Additionally, a mutation

in *nca-1* also strongly but incompletely suppressed the high-amplitude waveform of activated Rho,

but a mutation in *nca-2* did not suppress the high-amplitude waveform (Figure 5). Thus, activated

Rho may signal primarily through NCA channels formed from the NCA-1 pore-forming subunit.

Because an activated Rho mutant has slow locomotion (Figure 6) and *unc-79*, *unc-80*, and

nca-1 nca-2 double mutants also have slow locomotion, it is difficult to determine whether these

NCA channel mutants suppress the locomotion phenotype of activated Rho in addition to its

waveform. We performed radial locomotion assays that provide a combined measurement of

several aspects of the locomotion phenotype, including the rate of movement, frequency of

reversals, and amplitude of the waveform (see Materials and Methods). By these assays,

mutations in *unc-80* or *nca-1 nca-2* lead to only small increases in the radial distance traveled by

an activated Rho mutant and an *nca-2* mutant had no effect (Figure S4). However, a mutation in

nlf-1 much more strongly increased the radial distance traveled by an activated Rho mutant.

Because the *nlf-1* mutant is not as slow on its own, we could also directly assay its effect on the

rate of locomotion of an activated Rho mutant by counting the number of body bends per minute.

An *nlf-1* mutation strongly increased the rate of locomotion of an activated Rho mutant (Figure 6).

In fact, the *nlf-1* double mutant with activated Rho had a faster rate of locomotion than either

activated Rho or *nlf-1* on its own, similar to the effect of *nlf-1* on the locomotion of an activated

nca-1 mutant (Figure 2B). Additionally, a mutation in *nca-1* strongly increased the locomotion rate

of the activated Rho mutant, but a mutation in *nca-2* had no effect (Figure 6), further supporting

the idea that NCA channels consisting of the NCA-1 subunit act downstream of G_q and Rho in this

pathway.

      Rho regulates worm locomotion independently of effects on development, as demonstrated

by the fact that heat-shock induction of an activated Rho transgene in adults leads to a high-

amplitude waveform similar to that seen in worms that express activated Rho in acetylcholine

neurons (McMullan *et al.* 2006). Consistent with the idea that NCA-1 acts downstream of Rho,

mutations in the fainter genes *unc-79*, *unc-80*, and *nlf-1* suppress the high-amplitude waveform

phenotype of heat-shock induced activated Rho (Figure S5A). Additionally, *nlf-1* also suppresses

the locomotion defect of heat-shock induced activated Rho (Figure S5B). Thus, Rho regulates

worm locomotion via NCA-1 by acting in a non-developmental pathway that operates in adult

neurons.

A dominant NCA-1 mutation suppresses activated G_o

In *C. elegans*, the G_q pathway is opposed by signaling through the inhibitory G_o protein

GOA-1 (Hajdu-Cronin *et al.* 1999; Miller *et al.* 1999). Thus, loss-of-function mutants in *goa-1* are

hyperactive and have a high-amplitude waveform, similar to the gain-of-function G_q mutant *egl-*

30(tg26). We found that the uncloned dominant mutant *unc-109(n499)* which is paralyzed and

resembles loss-of-function mutants in *egl-30* (Park and Horvitz 1986) carries an activating

mutation in *goa-1* (see Materials and Methods). The *goa-1(n499)* mutant is paralyzed as a

heterozygote, and is lethal as a homozygote (Park and Horvitz 1986). We performed a screen for

suppressors of *goa-1(n499)* and isolated a partial intragenic suppressor, *goa-1(n499 ox304)*

(Materials and Methods). The *goa-1(n499 ox304)* mutant has a strong movement defect similar to

goa-1(n499), but is viable as a homozygote and thus easier to work with than *goa-1(n499)*.

We used the *goa-1(n499 ox304)* strain to perform a screen for suppressors of activated G_o

(Materials and Methods). Among the suppressors, we isolated a mutant (*ox352*) that was found to

be a dominant gain-of-function mutation in the *nca-1* gene (Fig. 2A, Chen et al., personal

communication). Thus, activation of NCA-1 suppresses activated G_o whereas loss of NCA-1

suppresses activated G_q, both consistent with the model that G_o inhibits G_q and that NCA-1 is a

downstream effector of the G_q pathway.

Discussion

In this study, we identify the NCA-1 ion channel as a downstream effector of the heterotrimeric G protein G_q. Our genetic epistasis data indicate that G_q activates the RhoGEF Trio and the small GTPase Rho. Rho then acts via an unknown mechanism to activate the NCA-1 ion channel, which is required for normal neuronal activity and synaptic transmission in *C. elegans* (Jospin *et al.* 2007; Yeh *et al.* 2008; Xie *et al.* 2013; Gao *et al.* 2015). Thus, this work identifies a new genetic pathway from G_q to an ion channel that regulates neuronal excitability and synaptic release: $G_q \rightarrow$ RhoGEF $\rightarrow$ Rho $\rightarrow$ NCA-1 (Figure 7). NCA-1 has not been previously identified as an effector of the G_q pathway, so this pathway may give insight into how NCA-1 is activated or regulated.

Mutants that eliminate the NCA channels such as *unc-80* or the *nca-1 nca-2* double mutant completely suppress the locomotion and body posture phenotypes of an activated Rho mutant (Figure 5, Figure S4), suggesting that NCA channels act downstream of Rho in a linear pathway. Additionally, channels composed of the pore-forming subunit NCA-1 may be the main target of G_q/Rho signaling. First, loss-of-function mutations in *nca-1* but not *nca-2* partially suppress activated G_q and activated Rho mutants. Second, activating gain-of-function mutants of *nca-1* have phenotypes reminiscent of activated G_q and activated Rho mutants, but no activated mutants in *nca-2* have been isolated. Third, loss-of-function mutations in *nca-1* but not *nca-2* enhance the weaker fainting phenotype of an *nlf-1* mutant which does not fully eliminate function of the NCA channels. Together, these data suggest that although either NCA-1 or NCA-2 activity is sufficient for wild-type locomotion, NCA-1 is likely to be the target of G protein regulation.

It is not clear, however, whether Rho activation of NCA-1 is direct or indirect. G_q directly interacts with and activates the RhoGEF and Rho, as shown by the crystal structure of a complex between $G_q\alpha$, RhoGEF, and Rho (Lutz *et al.* 2007). Rho is known to have many possible effectors and actions in cells (Etienne-Manneville and Hall 2002; Jaffe and Hall 2005). In *C. elegans* neurons, Rho has been previously shown to regulate synaptic transmission via at least two

pathways, one involving a direct interaction of Rho with the DAG kinase DGK-1 and one that is DGK-1 independent (McMullan *et al.* 2006). A candidate for the link between Rho and NCA-1 is the type I phosphatidylinositol 4-phosphate 5-kinase (PIP5K) that synthesizes the lipid phosphatidylinositol 4,5,-bisphosphate (PIP2). PIP5K is an intriguing candidate for several reasons. First, activation of G_q in mammalian cells has been shown to stimulate the membrane localization and activity of PIP5K via a mechanism that depends on Rho (Chatah and Abrams 2001; Weernink *et al.* 2004). Second, in *C. elegans*, mutations eliminating either the NCA channels (*nca-1 nca-2* double mutant) or their accessory subunits (*unc-79* or *unc-80*) suppress mutants in the PIP2 phosphatase synaptojanin (*unc-26*) and also suppress phenotypes caused by overexpression of the PIP5K gene *ppk-1* (Jospin *et al.* 2007). Loss of a PIP2 phosphatase or overexpression of a PIP5K are both predicted to increase levels of PIP2. Since loss of NCA channels suppresses the effects of too much PIP2, it is possible that excessive PIP2 leads to overactivation of NCA channels and that PIP2 might be part of the normal activation mechanism. There are numerous examples of ion channels that are regulated or gated by phosphoinositide lipids such as PIP2 (Balla 2013).

The NCA/NALCN ion channel was discovered originally by bioinformatic sequence analyses (Lee *et al.* 1999; Littleton and Ganetzky 2000). It is conserved among all metazoan animals and is evolutionarily-related to the family of voltage-gated sodium and calcium channels (Liebeskind *et al.* 2012), forming a new branch in this super-family. Although the cellular role of the NCA/NALCN channel and how it is gated are not well understood, NALCN and its orthologs are expressed broadly in the nervous system in a number of organisms (Lee *et al.* 1999; Lear *et al.* 2005; Humphrey *et al.* 2007; Lu *et al.* 2007; Jospin *et al.* 2007; Yeh *et al.* 2008; Lu and Feng 2011; Lutas *et al.* 2016). Moreover, mutations in this channel or its auxiliary subunits lead to defects in rhythmic behaviors in multiple organisms (Lear *et al.* 2005; Lu *et al.* 2007; Jospin *et al.* 2007; Yeh *et al.* 2008; Pierce-Shimomura *et al.* 2008; Lear *et al.* 2013; Xie *et al.* 2013; Funato *et al.* 2016). Thus, NCA/NALCN is likely to play an important role in controlling membrane

excitability. Additionally, NALCN currents have been reported to be activated by two different G protein-coupled receptors (GPCRs), the muscarinic acetylcholine receptor and the substance P receptor, albeit in a G protein-independent fashion (Lu *et al.* 2009; Swayne *et al.* 2009), and by low extracellular calcium via a G protein-dependent pathway (Lu *et al.* 2010). The latter study further showed that expression of an activated G_q mutant inhibited the NALCN sodium leak current, suggesting that high extracellular calcium tonically inhibits NALCN via a G_q-dependent pathway and that low extracellular calcium activates NALCN by relieving this inhibition (Lu *et al.* 2010). By contrast, we find that in *C. elegans*, the physiological role of G_q is to activate NCA-1 and show that NCA-1 is a physiologically relevant target of a G_q signaling pathway that acts through Rho.

In the last few years, both recessive and dominant human diseases characterized by a range of neurological symptoms including hypotonia, intellectual disability, and seizures have been shown to be caused by mutations in either *NALCN* or *UNC80* (Köroğlu *et al.* 2013; Al-Sayed *et al.* 2013; Chong *et al.* 2015; Aoyagi *et al.* 2015; Shamseldin *et al.* 2016; Stray-Pedersen *et al.* 2016; Gal *et al.* 2016; Fukai *et al.* 2016; Karakaya *et al.* 2016; Perez *et al.* 2016; Bend *et al.* 2016; Wang *et al.* 2016). Notably, dominant disease-causing mutations in the NALCN channel were modeled in worms and resemble either dominant activated or loss-of-function NCA mutants such as the ones we use in this study (Aoyagi *et al.* 2015; Bend *et al.* 2016). Human mutations in other components of the pathway we have described may cause similar clinical phenotypes.

Acknowledgments

We thank Shohei Mitani for the *nlf-1(tm3631)* mutant; Steve Nurrish for worm strains and plasmids carrying activated Rho or C3 transferase; Ken Miller for a plasmid with the *unc-17β* promoter; Wayne Davis for FLP/FRT plasmids; Yuji Kohara for cDNA clones; the Sanger Center for cosmids; Brooke Jarvie, Jill Hoyt, and Michelle Giarmarco for the isolation of *unc-79* and *unc-80* mutations in the G_q suppressor screen; and Dana Miller for the use of her microscope and

camera to take worm photographs. Some strains were provided by the CGC, which is funded by

NIH Office of Research Infrastructure Programs (P40 OD010440). M.A. is an Ellison Medical

Foundation New Scholar. E.M.J. is an Investigator of the Howard Hughes Medical Institute. This

work was supported by NIH grants R00 MH082109 to M.A and R01 NS034307 to E.M.J.

Figure Legends

Figure 1 Inhibition of Rho in neurons suppresses an activated G_q mutant. (A) Schematic of the

FLP/FRT system we used for temporal and spatial expression of the Rho inhibitor C3 transferase

(C3T). The transgene in the "off" configuration expresses the mCherry reporter under the control

of the promoter sequence but terminates transcription in the *let-858* 3'UTR upstream of GFP-C3T.

Expression of the FLP recombinase is induced by heat shock and leads to recombination between

the FRT sites and deletion of the intervening *mCherry::let-858* 3'UTR fragment. This leads to

transcription of GFP-C3T in the cells driven by the adjacent promoter. We used the pan-neuronal

promoter *rab-3p,* acetylcholine neuron promoter *unc-17p,* and head acetylcholine neuron

promoter *unc-17Hp.* (B) Inhibition of Rho in adult neurons causes a locomotion defect. Inhibition of

Rho by C3 transferase (C3T) in all adult neurons (*rab-3p, oxIs412* transgene), acetylcholine

neurons (*unc-17p, oxIs414* transgene) or head acetylcholine neurons (*unc-17Hp, oxIs434*

transgene) reduces the locomotion rate of wild type animals. ***, P<0.001, Dunnett's test. Error

bars = SEM; n = 19-20. (C) Inhibition of Rho in all adult neurons (*rab-3p, oxIs412* transgene) or

acetylcholine neurons (*unc-17p, oxIs414* transgene) led to a significant reduction in the rate of

locomotion of the activated G_q mutant *egl-30(tg26)* (here written Gq*). Inhibition of Rho in head

acetylcholine neurons (*unc-17Hp, oxIs434* transgene) led to a small decrease in Gq* locomotion

that was not statistically significant. ***, P<0.001; ns, not significant, P>0.05, Dunnett's test. Error

bars = SEM; n = 10-20. (D) Inhibition of Rho in all adult neurons (*rab-3p*) suppresses the deep

body bends of the activated G_q mutant *egl-30(tg26)* (Gq*). Rho inhibition in acetylcholine (*unc-*

17p) or head acetylcholine (*unc-17Hp*) neurons did not suppress the waveform of Gq*.

Figure 2 The *nlf-1(ox327)* mutation suppresses the coiled posture and locomotion defect of the

activated NCA-1 mutant *nca-1(ox352)* (Nca*). (A) Photos of first-day adult worms. (B) Body bend

assay. (C) Radial locomotion assay. Statistics in B and C: ***, P<0.001, two-tailed Mann-Whitney

test. Error bars = SEM, n = 4-8 (panel B), n= 8-26 (panel C).

Figure 3 *nlf-1* mutants are weak fainters. *nlf-1(ox327)* mutant animals have a much weaker fainter

phenotype than *unc-80(ox330)* or *nca-2(gk5)*; *nca-1(gk9)* double mutants in the forward direction

(A) and a slightly weaker fainter phenotype in the backward direction (B). Additionally, *nlf-1*

mutants are enhanced by a mutation in *nca-1*, but not *nca-2*. ***, P<0.001; ns, not significant,

P>0.05, Dunn's test. Error bars = SEM; n = 20.

Figure 4 *nlf-1* acts in head acetylcholine neurons to control locomotion. The *nlf-1* cDNA was

expressed in an *nlf-1(ox327)* mutant background using the following promoters: *rab-3* (all neurons,

oxEx1146 transgene), *unc-17* (acetylcholine neurons, *oxEx1149* transgene), *unc-17H* (a

derivative of the *unc-17* promoter that lacks the enhancer for ventral cord expression and thus

expresses only in head acetylcholine neurons, *oxEx1155* transgene), *unc-17β* (a derivative of

the *unc-17* promoter that expresses only in ventral cord acetylcholine neurons, *oxEx1323*

transgene), *acr-2* (ventral cord acetylcholine motor neurons, *oxEx1151* transgene), and *glr-1*

(interneurons including premotor command interneurons, *oxEx1152* transgene). (A) Expression

driven by the *rab-3*, *unc-17*, and *unc-17H* promoters rescued the fainting phenotype of an *nlf-1*

mutant in the backward direction. ***, P<0.001, Dunn's test. Error bars = SEM; n = 15. (B)

Expression driven by the *glr-1* promoter partially rescued the fainting phenotype of an *nlf-1* mutant

in either the backward or forward direction, though the effect was not statistically significant (ns,

P>0.05, two-tailed Mann-Whitney tests. Error bars = SEM; n = 25). However, assays of this strain

were complicated by slow movement and frequent pauses that were hard to distinguish from true

fainting behavior (see text for details).

Figure 5 Mutations in NCA channel subunits suppress the high-amplitude waveform of animals

expressing activated Rho. Animals expressing an activated Rho mutant (G14V) in acetylcholine

neurons (*nzIs29* transgene, written as Rho*) have a high-amplitude waveform. Mutations that

eliminate NCA-1 and NCA-2 channel function (*unc-80(ox330)* or *nca-2(gk5)*; *nca-1(gk9)*) strongly

suppress the high-amplitude waveform of activated Rho and convert the activated Rho mutants to

fainters. Mutations in *nlf-1(ox327)* and *nca-1(gk9)* strongly but incompletely suppress the high-

amplitude waveform of activated Rho and these double mutants do not faint. Mutations in *nca-*

2(gk5) do not suppress the high-amplitude waveform of activated Rho. Mutants in *unc-80*, *nlf-1*,

and the *nca-2 nca-1* double mutant have the typical fainter posture characterized by a

straightened anterior part of the body.

Figure 6 Mutations in *nlf-1* and *nca-1* suppress the locomotion defect of animals expressing

activated Rho. Animals expressing an activated Rho mutant (G14V) in acetylcholine neurons

(*nzIs29* transgene, written as Rho*) have a slow locomotion rate. The *nlf-1(ox327)* and *nca-1(gk9)*

mutations, but not *nca-2(gk5)*, strongly suppress the slow locomotion rate of activated Rho. ***,

P<0.001, Bonferroni test. Error bars = SEM; n = 16.

Figure 7 Model. G_q activates NCA-1 in a linear pathway via the RhoGEF Trio and small G protein

Rho. G_q activation of Trio and Rho is direct (solid arrow). Rho activation of NCA-1 is likely to be

indirect (dashed arrow). G_o inhibits G_q via the RGS protein EAT-16 (not shown). Thus, loss-of-

function mutations in NCA-1, its associated subunits UNC-79 and UNC-80, or the ER-localized

assembly factor NLF-1 suppress an activated G_q mutant, whereas a gain-of-function activating

mutation in NCA-1 suppresses an activated G_o mutant.

Figure S1 Mutations affecting the NCA-1 channel suppress the high-amplitude waveform of the

activated G_q mutant *egl-30(tg26)* (Gq*). The *unc-79(yak37)* and *unc-80(ox330)* mutations strongly

suppress the high-amplitude waveform of Gq*. The *nca-1(gk9)* and *nlf-1(ox327)* mutations

partially suppress the high-amplitude waveform of Gq*. Mutations in genes required for dense-

core vesicle biogenesis such as *rund-1(tm3622)* or the canonical G_q pathway such as *egl-*

8(ox333) only weakly suppress the high-amplitude waveform of Gq*. The *nca-2(gk5)* mutation

does not suppress Gq*.

Figure S2 *nlf-1* mutants have reduced *nca-1::gfp* and *nca-2::gfp* fluorescence in the nerve ring.

Quantification of the nerve ring fluorescence of C-terminally tagged *nca-1::gfp* (transgene *vaIs46*)

and *nca-2::gfp* (transgene *vaIs41*) in wild type and *nlf-1(ox327)* mutants. ***, P<0.001, two-tailed

unpaired t tests where each mutant strain is compared to its associated wild-type control. Error

bars = SEM; n = 11-12.

Figure S3 *nlf-1* is expressed widely in the nervous system. A fusion of the *nlf-1* promoter to GFP

(transgene *oxEx1144*) is expressed throughout the nervous system, including head and tail

neurons, and motor neurons in the ventral nerve cord. No expression is seen in non-neuronal

tissues. An L1 stage larval animal is shown with its head to the left.

Figure S4 Mutations in NCA channel subunits suppress the locomotion defect of animals

expressing activated Rho. Animals expressing an activated Rho mutant (G14V) in acetylcholine

neurons (*nzIs29* transgene, written as Rho*) have reduced locomotion as measured in radial

locomotion assays. Mutations that eliminate function of the NCA channels (*unc-80(ox330)* or *nca-*

2(gk5);nca-1(gk9)) increase the radial distance traveled by activated Rho mutants, but the effect is

subtle since *unc-80* and *nca-2; nca-1* mutants have reduced locomotion on their own. The *nlf-*

1(ox327) mutation that partially but incompletely reduces function of the NCA channels strongly

suppresses the locomotion defect of activated Rho. ***, P<0.0001, Bonferroni test. Error bars =

SEM; n = 5-18.

Figure S5 Mutations in NCA channel subunits suppress the high-amplitude waveform and

locomotion of adult animals expressing activated Rho. (A) Animals expressing an activated Rho

mutant (G14V) only in adults by heat-shock induced expression (*nzIs1* transgene, *hs::Rho**) have

a high-amplitude waveform. The *unc-80(ox330)* and *nlf-1(ox327)* mutations strongly suppress the

high-amplitude waveform of activated Rho. *unc-80*, but not *nlf-1*, makes the activated Rho animals

faint. (B) Animals expressing an activated Rho mutant (G14V) only in adults by heat-shock

induced expression (*nzIs1* transgene, *hs::Rho**) have a reduced locomotion rate. The *nlf-1(ox327)*

mutation suppresses the reduced locomotion rate of activated Rho. *, P=0.0157, two-tailed

unpaired t test. Error bars = SEM; n = 18-30)

Literature Cited

Adachi T., Kunitomo H., Tomioka M., Ohno H., Okochi Y., Mori I., Iino Y., 2010 Reversal of salt preference

is directed by the insulin/PI3K and Gq/PKC signaling in Caenorhabditis elegans. Genetics **186**:

1309–1319.

Ailion M., Hannemann M., Dalton S., Pappas A., Watanabe S., Hegermann J., Liu Q., Han H.-F., Gu M.,

Goulding M. Q., Sasidharan N., Schuske K., Hullett P., Eimer S., Jorgensen E. M., 2014 Two Rab2

interactors regulate dense-core vesicle maturation. Neuron **82**: 167–180.

Al-Sayed M. D., Al-Zaidan H., Albakheet A., Hakami H., Kenana R., Al-Yafee Y., Al-Dosary M., Qari A., Al-

Sheddi T., Al-Muheiza M., Al-Qubbaj W., Lakmache Y., Al-Hindi H., Ghaziuddin M., Colak D., Kaya

N., 2013 Mutations in NALCN cause an autosomal-recessive syndrome with severe hypotonia,

speech impairment, and cognitive delay. Am. J. Hum. Genet. **93**: 721–726.

Aoyagi K., Rossignol E., Hamdan F. F., Mulcahy B., Xie L., Nagamatsu S., Rouleau G. A., Zhen M.,

Michaud J. L., 2015 A Gain-of-Function Mutation in NALCN in a Child with Intellectual Disability,

Ataxia, and Arthrogryposis. Hum. Mutat. **36**: 753–757.

Balla T., 2013 Phosphoinositides: tiny lipids with giant impact on cell regulation. Physiol. Rev. **93**: 1019–

1137.

Bastiani C. A., Gharib S., Simon M. I., Sternberg P. W., 2003 Caenorhabditis elegans Galphaq regulates

egg-laying behavior via a PLCbeta-independent and serotonin-dependent signaling pathway and

likely functions both in the nervous system and in muscle. Genetics **165**: 1805–1822.

Bend E. G., Si Y., Stevenson D. A., Bayrak-Toydemir P., Newcomb T. M., Jorgensen E. M., Swoboda K. J.,

2016 NALCN channelopathies: Distinguishing gain-of-function and loss-of-function mutations.

Neurology **87**: 1131–1139.

Boone A. N., Senatore A., Chemin J., Monteil A., Spafford J. D., 2014 Gd3+ and calcium sensitive, sodium

leak currents are features of weak membrane-glass seals in patch clamp recordings. PLoS ONE **9**:

e98808.

Brundage L., Avery L., Katz A., Kim U. J., Mendel J. E., Sternberg P. W., Simon M. I., 1996 Mutations in a

C. elegans Gqalpha gene disrupt movement, egg laying, and viability. Neuron **16**: 999–1009.

Charlie N. K., Schade M. A., Thomure A. M., Miller K. G., 2006 Presynaptic UNC-31 (CAPS) is required to

activate the G alpha(s) pathway of the Caenorhabditis elegans synaptic signaling network. Genetics

172: 943–961.

Chatah N. E., Abrams C. S., 2001 G-protein-coupled receptor activation induces the membrane

translocation and activation of phosphatidylinositol-4-phosphate 5-kinase I alpha by a Rac- and

Rho-dependent pathway. J. Biol. Chem. **276**: 34059–34065.

Chong J. X., McMillin M. J., Shively K. M., Beck A. E., Marvin C. T., Armenteros J. R., Buckingham K. J.,

Nkinsi N. T., Boyle E. A., Berry M. N., Bocian M., Foulds N., Uzielli M. L. G., Haldeman-Englert C.,

Hennekam R. C. M., Kaplan P., Kline A. D., Mercer C. L., Nowaczyk M. J. M., Klein Wassink-Ruiter

J. S., McPherson E. W., Moreno R. A., Scheuerle A. E., Shashi V., Stevens C. A., Carey J. C.,

Monteil A., Lory P., Tabor H. K., Smith J. D., Shendure J., Nickerson D. A., University of

Washington Center for Mendelian Genomics, Bamshad M. J., 2015 De novo mutations in NALCN

cause a syndrome characterized by congenital contractures of the limbs and face, hypotonia, and

developmental delay. Am. J. Hum. Genet. **96**: 462–473.

Coleman D. E., Berghuis A. M., Lee E., Linder M. E., Gilman A. G., Sprang S. R., 1994 Structures of active

conformations of Gi alpha 1 and the mechanism of GTP hydrolysis. Science **265**: 1405–1412.

Coulon P., Kanyshkova T., Broicher T., Munsch T., Wettschureck N., Seidenbecher T., Meuth S. G.,

Offermanns S., Pape H.-C., Budde T., 2010 Activity Modes in Thalamocortical Relay Neurons are

Modulated by G(q)/G(11) Family G-proteins - Serotonergic and Glutamatergic Signaling. Front Cell

Neurosci **4**: 132.

Davis M. W., Hammarlund M., Harrach T., Hullett P., Olsen S., Jorgensen E. M., 2005 Rapid single

nucleotide polymorphism mapping in C. elegans. BMC Genomics **6**: 118.

Davis M. W., Morton J. J., Carroll D., Jorgensen E. M., 2008 Gene activation using FLP recombinase in C.

elegans. PLoS Genet. **4**: e1000028.

Esposito G., Amoroso M. R., Bergamasco C., Di Schiavi E., Bazzicalupo P., 2010 The G protein regulators

EGL-10 and EAT-16, the Giα GOA-1 and the G(q)α EGL-30 modulate the response of the C.

elegans ASH polymodal nociceptive sensory neurons to repellents. BMC Biol. **8**: 138.

Etienne-Manneville S., Hall A., 2002 Rho GTPases in cell biology. Nature **420**: 629–635.

Fukai R., Saitsu H., Okamoto N., Sakai Y., Fattal-Valevski A., Masaaki S., Kitai Y., Torio M., Kojima-Ishii

K., Ihara K., Chernuha V., Nakashima M., Miyatake S., Tanaka F., Miyake N., Matsumoto N., 2016

De novo missense mutations in NALCN cause developmental and intellectual impairment with

hypotonia. J. Hum. Genet. **61**: 451–455.

Funato H., Miyoshi C., Fujiyama T., Kanda T., Sato M., Wang Z., Ma J., Nakane S., Tomita J., Ikkyu A.,

Kakizaki M., Hotta-Hirashima N., Kanno S., Komiya H., Asano F., Honda T., Kim S. J., Harano K.,

Muramoto H., Yonezawa T., Mizuno S., Miyazaki S., Connor L., Kumar V., Miura I., Suzuki T.,

Watanabe A., Abe M., Sugiyama F., Takahashi S., Sakimura K., Hayashi Y., Liu Q., Kume K.,

Wakana S., Takahashi J. S., Yanagisawa M., 2016 Forward-genetics analysis of sleep in randomly

mutagenized mice. Nature **539**: 378–383.

Gal M., Magen D., Zahran Y., Ravid S., Eran A., Khayat M., Gafni C., Levanon E. Y., Mandel H., 2016 A

novel homozygous splice site mutation in NALCN identified in siblings with cachexia, strabismus,

severe intellectual disability, epilepsy and abnormal respiratory rhythm. Eur J Med Genet **59**: 204–

209.

Gamper N., Reznikov V., Yamada Y., Yang J., Shapiro M. S., 2004 Phosphatidylinositol [correction] 4,5-

bisphosphate signals underlie receptor-specific Gq/11-mediated modulation of N-type Ca2+

channels. J. Neurosci. **24**: 10980–10992.

Gao S., Xie L., Kawano T., Po M. D., Pirri J. K., Guan S., Alkema M. J., Zhen M., 2015 The NCA sodium
leak channel is required for persistent motor circuit activity that sustains locomotion. Nat Commun
6: 6323.

Hajdu-Cronin Y. M., Chen W. J., Patikoglou G., Koelle M. R., Sternberg P. W., 1999 Antagonism between
G(o)alpha and G(q)alpha in Caenorhabditis elegans: the RGS protein EAT-16 is necessary for
G(o)alpha signaling and regulates G(q)alpha activity. Genes Dev **13**: 1780–1793.

Hajdu-Cronin Y. M., Chen W. J., Sternberg P. W., 2004 The L-type cyclin CYL-1 and the heat-shock-factor
HSF-1 are required for heat-shock-induced protein expression in Caenorhabditis elegans. Genetics
168: 1937–1949.

Hammarlund M., Palfreyman M. T., Watanabe S., Olsen S., Jorgensen E. M., 2007 Open syntaxin docks
synaptic vesicles. PLoS Biol. **5**: e198.

Hiley E., McMullan R., Nurrish S. J., 2006 The Galpha12-RGS RhoGEF-RhoA signalling pathway regulates
neurotransmitter release in C. elegans. EMBO J. **25**: 5884–5895.

Humphrey J. A., Hamming K. S., Thacker C. M., Scott R. L., Sedensky M. M., Snutch T. P., Morgan P. G.,
Nash H. A., 2007 A putative cation channel and its novel regulator: cross-species conservation of
effects on general anesthesia. Curr. Biol. **17**: 624–629.

Jaffe A. B., Hall A., 2005 Rho GTPases: biochemistry and biology. Annu. Rev. Cell Dev. Biol. **21**: 247–269.

Jantsch-Plunger V., Gönczy P., Romano A., Schnabel H., Hamill D., Schnabel R., Hyman A. A., Glotzer M.,
2000 CYK-4: A Rho family gtpase activating protein (GAP) required for central spindle formation
and cytokinesis. J. Cell Biol. **149**: 1391–1404.

Jospin M., Watanabe S., Joshi D., Young S., Hamming K., Thacker C., Snutch T. P., Jorgensen E. M.,
Schuske K., 2007 UNC-80 and the NCA ion channels contribute to endocytosis defects in
synaptojanin mutants. Curr. Biol. **17**: 1595–1600.

Karakaya M., Heller R., Kunde V., Zimmer K.-P., Chao C.-M., Nürnberg P., Cirak S., 2016 Novel Mutations

in the Nonselective Sodium Leak Channel (NALCN) Lead to Distal Arthrogryposis with Increased

Muscle Tone. Neuropediatrics **47**:273-277.

Köroğlu Ç., Seven M., Tolun A., 2013 Recessive truncating NALCN mutation in infantile neuroaxonal

dystrophy with facial dysmorphism. J. Med. Genet. **50**: 515–520.

Krause M., Offermanns S., Stocker M., Pedarzani P., 2002 Functional specificity of G alpha q and G alpha

11 in the cholinergic and glutamatergic modulation of potassium currents and excitability in

hippocampal neurons. J. Neurosci. **22**: 666–673.

Lackner M. R., Nurrish S. J., Kaplan J. M., 1999 Facilitation of synaptic transmission by EGL-30 Gqalpha

and EGL-8 PLCbeta: DAG binding to UNC-13 is required to stimulate acetylcholine release. Neuron

24: 335–346.

Lear B. C., Lin J.-M., Keath J. R., McGill J. J., Raman I. M., Allada R., 2005 The ion channel narrow

abdomen is critical for neural output of the Drosophila circadian pacemaker. Neuron **48**: 965–976.

Lear B. C., Darrah E. J., Aldrich B. T., Gebre S., Scott R. L., Nash H. A., Allada R., 2013 UNC79 and

UNC80, Putative Auxiliary Subunits of the NARROW ABDOMEN Ion Channel, Are Indispensable

for Robust Circadian Locomotor Rhythms in Drosophila. PLoS ONE **8**: e78147.

Lee J. H., Cribbs L. L., Perez-Reyes E., 1999 Cloning of a novel four repeat protein related to voltage-

gated sodium and calcium channels. FEBS Lett. **445**: 231–236.

Liebeskind B. J., Hillis D. M., Zakon H. H., 2012 Phylogeny unites animal sodium leak channels with fungal

calcium channels in an ancient, voltage-insensitive clade. Mol. Biol. Evol. **29**: 3613–3616.

Littleton J. T., Ganetzky B., 2000 Ion channels and synaptic organization: analysis of the Drosophila

genome. Neuron **26**: 35–43.

Lu B., Su Y., Das S., Liu J., Xia J., Ren D., 2007 The neuronal channel NALCN contributes resting sodium

permeability and is required for normal respiratory rhythm. Cell **129**: 371–383.

Lu B., Su Y., Das S., Wang H., Wang Y., Liu J., Ren D., 2009 Peptide neurotransmitters activate a cation

channel complex of NALCN and UNC-80. Nature **457**: 741–744.

Lu B., Zhang Q., Wang H., Wang Y., Nakayama M., Ren D., 2010 Extracellular calcium controls

background current and neuronal excitability via an UNC79-UNC80-NALCN cation channel

complex. Neuron **68**: 488–499.

Lu T. Z., Feng Z.-P., 2011 A sodium leak current regulates pacemaker activity of adult central pattern

generator neurons in Lymnaea stagnalis. PLoS ONE **6**: e18745.

Lutas A., Lahmann C., Soumillon M., Yellen G., 2016 The leak channel NALCN controls tonic firing and

glycolytic sensitivity of substantia nigra pars reticulata neurons. eLife **5**: e15271.

Lutz S., Freichel-Blomquist A., Yang Y., Rümenapp U., Jakobs K. H., Schmidt M., Wieland T., 2005 The

guanine nucleotide exchange factor p63RhoGEF, a specific link between Gq/11-coupled receptor

signaling and RhoA. J. Biol. Chem. **280**: 11134–11139.

Lutz S., Shankaranarayanan A., Coco C., Ridilla M., Nance M. R., Vettel C., Baltus D., Evelyn C. R.,

Neubig R. R., Wieland T., Tesmer J. J. G., 2007 Structure of Galphaq-p63RhoGEF-RhoA complex

reveals a pathway for the activation of RhoA by GPCRs. Science **318**: 1923–1927.

Matsuki M., Kunitomo H., Iino Y., 2006 Goalpha regulates olfactory adaptation by antagonizing Gqalpha-

DAG signaling in Caenorhabditis elegans. Proc. Natl. Acad. Sci. U.S.A. **103**: 1112–1117.

McMullan R., Hiley E., Morrison P., Nurrish S. J., 2006 Rho is a presynaptic activator of neurotransmitter

release at pre-existing synapses in C. elegans. Genes Dev. **20**: 65–76.

Mello C. C., Kramer J. M., Stinchcomb D., Ambros V., 1991 Efficient gene transfer in C.elegans:

extrachromosomal maintenance and integration of transforming sequences. EMBO J **10**: 3959–

3970.

Miller K. G., Emerson M. D., Rand J. B., 1999 Goalpha and diacylglycerol kinase negatively regulate the

Gqalpha pathway in C. elegans. Neuron **24**: 323–333.

Park E. C., Horvitz H. R., 1986 Mutations with dominant effects on the behavior and morphology of the

nematode Caenorhabditis elegans. Genetics **113**: 821–852.

Pereira L., Kratsios P., Serrano-Saiz E., Sheftel H., Mayo A. E., Hall D. H., White J. G., LeBoeuf B., Garcia

L. R., Alon U., Hobert O., 2015 A cellular and regulatory map of the cholinergic nervous system of

C. elegans. eLife **4**: e12432.

Perez Y., Kadir R., Volodarsky M., Noyman I., Flusser H., Shorer Z., Gradstein L., Birnbaum R. Y., Birk O.

S., 2016 UNC80 mutation causes a syndrome of hypotonia, severe intellectual disability, dyskinesia

and dysmorphism, similar to that caused by mutations in its interacting cation channel NALCN. J.

Med. Genet. **53**: 397–402.

Pierce-Shimomura J. T., Chen B. L., Mun J. J., Ho R., Sarkis R., McIntire S. L., 2008 Genetic analysis of

crawling and swimming locomotory patterns in C. elegans. Proc. Natl. Acad. Sci. U.S.A. **105**:

20982–20987.

Sánchez-Fernández G., Cabezudo S., García-Hoz C., Benincá C., Aragay A. M., Mayor F., Ribas C., 2014

Gαq signalling: the new and the old. Cell. Signal. **26**: 833–848.

Senatore A., Monteil A., Minnen J. van, Smit A. B., Spafford J. D., 2013 NALCN ion channels have

alternative selectivity filters resembling calcium channels or sodium channels. PLoS ONE **8**:

e55088.

Senatore A., Spafford J. D., 2013 A uniquely adaptable pore is consistent with NALCN being an ion sensor.

Channels (Austin) **7**: 60–68.

Shamseldin H. E., Faqeih E., Alasmari A., Zaki M. S., Gleeson J. G., Alkuraya F. S., 2016 Mutations in

UNC80, Encoding Part of the UNC79-UNC80-NALCN Channel Complex, Cause Autosomal-

Recessive Severe Infantile Encephalopathy. Am. J. Hum. Genet. **98**: 210–215.

Stray-Pedersen A., Cobben J.-M., Prescott T. E., Lee S., Cang C., Aranda K., Ahmed S., Alders M.,

Gerstner T., Aslaksen K., Tétreault M., Qin W., Hartley T., Jhangiani S. N., Muzny D. M., Tarailo-

Graovac M., Karnebeek C. D. M. van, Care4Rare Canada Consortium, Baylor-Hopkins Center for

Mendelian Genomics, Lupski J. R., Ren D., Yoon G., 2016 Biallelic Mutations in UNC80 Cause

Persistent Hypotonia, Encephalopathy, Growth Retardation, and Severe Intellectual Disability. Am.

J. Hum. Genet. **98**: 202–209.

Swayne L. A., Mezghrani A., Varrault A., Chemin J., Bertrand G., Dalle S., Bourinet E., Lory P., Miller R. J.,

Nargeot J., Monteil A., 2009 The NALCN ion channel is activated by M3 muscarinic receptors in a

pancreatic beta-cell line. EMBO Rep. **10**: 873–880.

Topalidou I., Cattin-Ortolá J., Pappas A. L., Cooper K., Merrihew G. E., MacCoss M. J., Ailion M., 2016 The

EARP Complex and Its Interactor EIPR-1 Are Required for Cargo Sorting to Dense-Core Vesicles.

PLoS Genet. **12**: e1006074.

Vogt S., Grosse R., Schultz G., Offermanns S., 2003 Receptor-dependent RhoA activation in G12/G13-

deficient cells: genetic evidence for an involvement of Gq/G11. J. Biol. Chem. **278**: 28743–28749.

Wang Y., Koh K., Ichinose Y., Yasumura M., Ohtsuka T., Takiyama Y., 2016 A de novo mutation in the

NALCN gene in an adult patient with cerebellar ataxia associated with intellectual disability and

arthrogryposis. Clin. Genet. **90**: 556–557.

Weernink P. A. O., Meletiadis K., Hommeltenberg S., Hinz M., Ishihara H., Schmidt M., Jakobs K. H., 2004

Activation of type I phosphatidylinositol 4-phosphate 5-kinase isoforms by the Rho GTPases, RhoA,

Rac1, and Cdc42. J. Biol. Chem. **279**: 7840–7849.

Wilkie T. M., Scherle P. A., Strathmann M. P., Slepak V. Z., Simon M. I., 1991 Characterization of G-

protein alpha subunits in the Gq class: expression in murine tissues and in stromal and

hematopoietic cell lines. Proc. Natl. Acad. Sci. U.S.A. **88**: 10049–10053.

Wilkie T. M., Gilbert D. J., Olsen A. S., Chen X. N., Amatruda T. T., Korenberg J. R., Trask B. J., Jong P.

de, Reed R. R., Simon M. I., 1992 Evolution of the mammalian G protein alpha subunit multigene

family. Nat. Genet. **1**: 85–91.

Williams S. L., Lutz S., Charlie N. K., Vettel C., Ailion M., Coco C., Tesmer J. J. G., Jorgensen E. M.,

Wieland T., Miller K. G., 2007 Trio's Rho-specific GEF domain is the missing Galpha q effector in C.

elegans. Genes Dev **21**: 2731–2746.

Xie L., Gao S., Alcaire S. M., Aoyagi K., Wang Y., Griffin J. K., Stagljar I., Nagamatsu S., Zhen M., 2013

NLF-1 delivers a sodium leak channel to regulate neuronal excitability and modulate rhythmic

locomotion. Neuron **77**: 1069–1082.

Yeh E., Ng S., Zhang M., Bouhours M., Wang Y., Wang M., Hung W., Aoyagi K., Melnik-Martinez K., Li M.,

Liu F., Schafer W. R., Zhen M., 2008 A putative cation channel, NCA-1, and a novel protein, UNC-

80, transmit neuronal activity in C. elegans. PLoS Biol. **6**: e55.

Figure 1

Figure 2

A

WT
Nca*
Nca*; *nlf-1*

B

C

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure S1

Gq*

Gq*; *rund-1*

Gq*; *egl-8*

Gq*; *nlf-1*

Gq*; *nca-1*

Gq*; *nca-2*

Gq*; *unc-79*

Gq*; *unc-80*

Figure S2

Figure S3

Figure S4

Figure S5

A

WT   unc-80   nlf-1

hs::Rho*   hs::Rho*; unc-80   hs::Rho*; nlf-1

B

Table S1. List of strains

CB4856 Hawaiian wild isolate
EG303 *goa-1(n499 ox303) I*
EG304 *goa-1(n499 ox304) I*
EG327 *nlf-1(ox327) X*
EG330 *unc-80(ox330) V*
EG352 *nca-1(ox352) IV*
EG3447 *egl-30(tg26) I goa-1(n499) I / egl-30(tg26) I goa-1(+) I*
EG3498 *egl-30(tg26) I goa-1(n499) I / egl-30(tg26) I unc-13(e51) I gld-1(q126) I*
EG3647 *egl-30(tg26) I ; nlf-1(ox327) X*
EG3649 *egl-30(tg26) I ; unc-80(ox329) V*
EG3650 *egl-30(tg26) I ; unc-80(ox330) V*
EG3761 *unc-79(e1068) III*
EG3771 *egl-30(tg26) I ; unc-80(e1272) V*
EG3798 *egl-30(tg26) I ; egl-8(ox333) V*
EG3804 *nca-2(gk5) III ; nca-1(gk9) IV*
EG4036 *egl-30(tg26) I ; nca-2(gk5) III ; nca-1(gk9) IV*
EG4067 *goa-1(n499 ox304) I; nca-1(ox352) IV*
EG4494 *nca-1(ox352) IV ; nlf-1(ox327) X*
EG4530 *nzIs29[unc-17p::rho-1(G14V), unc-122p::gfp] II ; unc-79(e1068) III*
EG4531 *nzIs29[unc-17p::rho-1(G14V), unc-122p::gfp] II ; unc-80(ox330) V*
EG4532 *egl-30(tg26) I*
EG4681 *nzIs29[unc-17p::rho-1(G14V), unc-122p::gfp] II ; nlf-1(ox327) X*
EG4684 *nzIs1[hs::rho-1(G14V), ttx-3p::gfp] I ; unc-79(e1068) III*
EG4685 *nzIs1[hs::rho-1(G14V), ttx-3p::gfp] I ; unc-80(ox330) V*
EG4686 *nzIs1[hs::rho-1(G14V), ttx-3p::gfp] I ; nlf-1(ox327) X*
EG4690 *nzIs29[unc-17p::rho-1(G14V), unc-122p::gfp] II ; nca-2(gk5) III ; nca-1(gk9) IV*
EG4694 *unc-80(ox330) V ; nlf-1(ox327) X*
EG4697 *unc-79(e1068) III ; nlf-1(ox327) X*
EG4700 *nzIs29[unc-17p::rho-1(G14V), unc-122p::gfp] II ; nca-1(gk9) IV*
EG4718 *nlf-1(ox327) X ; oxEx1041[cosmid F55A4, unc-17p::mCherry::H2B, lin-15(+)]*
EG4733 *nca-1(gk9) IV ; vaIs46[NCA-1::GFP, lin-15+] X*
EG4761 *nzIs29[unc-17p::rho-1(G14V), unc-122p::gfp] II ; nca-2(gk5) III*
EG4782 *nzIs29[unc-17p::rho-1(G14V), unc-122p::gfp] II*
EG4783 *nca-2(gk5) III ; vaIs41[NCA-2::GFP, lin-15+]*
EG4785 *egl-30(tg26) I ; nca-1(gk9) IV*
EG4786 *egl-30(tg26) I ; nca-2(gk5) III*
EG4834 *nca-1(gk9) IV; nlf-1(ox327) X*
EG4835 *nca-2(gk5) IV; nlf-1(ox327) X*
EG4836 *nlf-1(ox327) X ; vaIs46[NCA-1::GFP, lin-15+]*
EG4837 *nlf-1(ox327) X ; vaIs41[NCA-2::GFP, lin-15+]*
EG4943 *nlf-1(ox327) X ; oxEx1146[rab-3p::nlf-1(+) cDNA::GFP, myo-2p::mCherry]*
EG5217 *oxIs412[rab-3p-FRT-mCherry-FRT-GFP-C3T, lin-15(+), hsp-FLP] II*
EG5218 *oxIs414[unc-17p-FRT-mCherry-FRT-GFP-C3T, lin-15(+), hsp-FLP] II*
EG5252 *egl-30(tg26) I ; oxIs412[rab-3p-FRT-mCherry-FRT-GFP-C3T, lin-15(+), hsp-FLP] II*
EG5263 *egl-30(tg26) I ; oxIs414[unc-17p-FRT-mCherry-FRT-GFP-C3T, lin-15(+), hsp-FLP] II*
EG5279 *oxIs434[unc-17Hp-FRT-mCherry-FRT-GFP-C3T, lin-15(+), hsp-FLP] X*
EG5287 *egl-30(tg26) I ; oxIs434[unc-17Hp-FRT-mCherry-FRT-GFP-C3T, lin-15(+), hsp-FLP] X*
EG5320 *oxIs435[unc-17βp-FRT-mCherry-FRT-GFP-C3T, lin-15(+), hsp-FLP] II*
EG5504 *nlf-1(tm3631) X*
EG5609 *egl-30(tg26) I ; rund-1(tm3622) X*

EG5611 *egl-30(tg26) I ; nlf-1(tm3631) X*
EG7249 *eri-1(mg366) IV ; lin-15(n744) X*
MT1102 *goa-1(n499) I / +*
N2 Bristol isolate, standard lab wild type
QT47 *nzIs1[hs::rho-1(G14V), ttx-3p::gfp] I*
VC9 *nca-2(gk5) III*
VC12 *nca-1(gk9) IV*
XZ1080 *egl-30(tg26) I ; unc-80(yak8) V*
XZ1113 *egl-30(tg26) I ; unc-80(yak35) V*
XZ1114 *egl-30(tg26) I ; unc-80(yak36) V*
XZ1115 *egl-30(tg26) I ; unc-79(yak37) III*
XZ1151 *egl-30(tg26) I*
XZ1152 *egl-30(tg26) I ; nuIs183[unc-129p::NLP-21-Venus, myo-2p::NLS-GFP] III*
XZ1186 *egl-30(tg26) I ; nuIs183[unc-129p::NLP-21-Venus, myo-2p::NLS-GFP] III; unc-80(yak56) V*
XZ1191 *egl-30(tg26) I ; unc-79(yak61) III nuIs183[unc-129p::NLP-21-Venus, myo-2p::NLS-GFP] III*
XZ1229 *egl-30(tg26) I ; unc-79(yak73) III nuIs183[unc-129p::NLP-21-Venus, myo-2p::NLS-GFP] III*
XZ2008 *lin-15(n765ts) X ; oxEx1144[nlf-1p::GFP, lin-15(+)]*
XZ2009 *nlf-1(ox327) X ; oxEx1147[nlf-1p::nlf-1(+)::GFP, myo-2p::mCherry]*
XZ2010 *nlf-1(ox327) X ; oxEx1149[unc-17p::nlf-1(+) cDNA::GFP, myo-2p::mCherry]*
XZ2011 *nlf-1(ox327) X ; oxEx1155[unc-17Hp::nlf-1(+) cDNA::GFP, myo-2p::mCherry]*
XZ2012 *nlf-1(ox327) X ; oxEx1151[acr-2p::nlf-1(+) cDNA::GFP, myo-2p::mCherry]*
XZ2013 *nlf-1(ox327) X ; oxEx1323[unc-17βp::nlf-1(+) cDNA::GFP, myo-2p::mCherry]*
XZ2014 *nlf-1(ox327) X ; oxEx1152[glr-1p::nlf-1(+) cDNA::GFP, myo-2p::mCherry]*

Table S2. List of plasmids

Miscellaneous plasmids
pCFJ90	*myo-2::mCherry*
pGH47	*unc-17p::mCherry::H2B*
pL15EK	*lin-15(+)*
pWD79-2RV	*hsp-16.48p::FLP*
QT#99	*Phs::GFP-C3T*
F55A4	cosmid carrying the *nlf-1* gene F55A4.2, used to make *oxEx1041* (10 ng/μl)
yk1279a1	*nlf-1* full length cDNA
Litmus 28i	
Litmus 38i	

Gateway entry clones
p_C06E1.4-93	*glr-1p* [4-1] (727 bp of the *glr-1* promoter upstream of and including the ATG)
pADA180	*unc-17Hp* [4-1] (derived from the *unc-17* promoter of pGH1, carrying a 127 bp deletion that removes the nerve cord *β* enhancer, expressed in head cholinergic neurons)
pCFJ31	*acr-2p* [4-1] (3362 bp of the *acr-2* promoter upstream of the ATG)
pCR110	*GFP* [1-2]
pCR185	*GFP::unc-54* 3'UTR [2-3]
pEntry[2-3]-UTR[unc-54]	*unc-54 3'UTR* [2-3]
pEntry[4-1]-P[rab-3]	*rab-3p* [4-1] (1224 bp of the *rab-3* promoter upstream of and including the ATG)
pGH1	*unc-17p* [4-1] (3229 bp of the *unc-17* promoter upstream of and including the ATG)
pMA1	*nlf-1p* [4-1] (5665 bp of the *nlf-1* promoter upstream of and including the ATG)
pMA2	*nlf-1(+)* gene with introns [1-2]
pMA3	*nlf-1* cDNA [1-2] (derived from yk1279a1)
pMA19	*C3T::unc-54 3'UTR* [2-3]
pMA23	*unc-17βp* [4-1] (a fusion of the *unc-17* nerve cord *β* enhancer to the minimal *unc-17* promoter consisting of the 382 bp upstream of and including the ATG, expressed in ventral nerve cord cholinergic neurons)
pWD178	*FRT::mCherry::let-858 3'UTR::FRT::GFP* [1-2]

Gateway expression constructs
pMA4	*nlf-1p::GFP*, used to make *oxEx1144* (10 ng/μl)
pMA5	*rab-3p::nlf-1 cDNA::GFP*, used to make *oxEx1146* (10 ng/μl)
pMA6	*nlf-1p::nlf-1(+)::GFP*, used to make *oxEx1147* (10 ng/μl)
pMA8	*unc-17p::nlf-1 cDNA::GFP*, used to make *oxEx1149* (10 ng/μl)
pMA9	*acr-2p::nlf-1 cDNA::GFP*, used to make *oxEx1151* (10 ng/μl)
pMA10	*glr-1p::nlf-1 cDNA::GFP*, used to make *oxEx1152* (10 ng/μl)
pMA11	*unc-17Hp::nlf-1 cDNA::GFP*, used to make *oxEx1155* (10 ng/μl)
pMA27	*rab-3p::FRT::mCherry::let-858 3'UTR::FRT::GFP::C3T::unc-54 3'UTR*, used to make *oxIs412* (10 ng/μl)
pMA28	*unc-17p::FRT::mCherry::let-858 3'UTR::FRT::GFP::C3T::unc-54 3'UTR*, used to make *oxIs414* (10 ng/μl)
pMA39	*unc-17Hp::FRT::mCherry::let-858 3'UTR::FRT::GFP::C3T::unc-54 3'UTR*, used to make *oxIs434* (10 ng/μl)
pMA40	*unc-17βp::FRT::mCherry::let-858 3'UTR::FRT::GFP::C3T::unc-54 3'UTR*, used to make *oxIs435* (10 ng/μl)
pMA52	*unc-17β::nlf-1 cDNA::GFP*, used to make *oxEx1323* (10 ng/μl)