1

9 **High levels of antibiotic tolerance and persistence are induced by the**

10 **commercial anti-microbial triclosan**

11 Corey S. Westfall and Petra Anne Levin*

12 Department of Biology, Washington University in St. Louis, St. Louis, MO 63130, USA

*Correspondence: plevin@wustl.edu

Antimicrobial tolerance and bacterial persistence are confounding factors in the treatment

of recurrent and chronic infections. Here we report that the anti-microbial triclosan

increased antibiotic tolerance up to 10,000 fold and amplified persister populations in both

***Escherichia coli* and methicillin-resistant *Staphylococcus aureus*. Triclosan-mediated**

protection was dependent on the alarmone ppGpp but independent of growth rate, arguing

against a simplistic model in which slow growth is sufficient to confer antibiotic tolerance.

The prophylactic use of antibiotics in consumer goods ranging from animal feed to personal care

products is widely believed to be a major contributor to the epidemic increase in antibiotic

resistant pathogens[1-3]. While the inverse relationship between antibiotic use and antibiotic

efficacy is largely attributable to the selection of heritable traits, non-heritable traits such as

antibiotic tolerance and persistence are also likely to be involved[4].

In contrast to genetically resistant bacteria, which grow in the presence of an antibiotic, tolerant

bacteria are able to survive antibiotic challenge for longer periods of time than their more

sensitive counterparts[5]. Persister cells are the small sub-set of an otherwise-sensitive population

($\sim$1 in 10^6) that exhibit levels of tolerance sufficient to protect them from otherwise lethal

concentrations of antimicrobial compounds[6]. Increases in antibiotic tolerance and persistence are

confounding factors in the treatment of chronic *P. aeruginosa*[7] and *S. aureus*[8] infections and are

thought to contribute to the refractile nature of medically relevant biofilms. Reduced growth rate

is associated with increased antibiotic tolerance[9] and is a defining trait of persister cells.

Banned from consumer soaps effective September 2017 by the US Food and Drug

Administration, the antimicrobial triclosan remains approved for use in products ranging from

toothpaste to cleansers employed in healthcare settings[10]. In contrast to bactericidal antibiotics,

which kill pathogens outright, triclosan is a bacteriostatic drug that inhibits growth by targeting

enoyl-acyl carrier protein reductase to interfere with early steps in fatty acid synthesis[11].

Based on its ability to inhibit growth, we wondered if physiologically relevant levels of triclosan

might provide protection against bactericidal antibiotics. To address this possibility we examined

the relative sensitivity of *E. coli* (MG1655) and *S. aureus* (FPR3757 an USA-300 MRSA strain)

cultured in minimal inhibitory concentrations (MIC) of triclosan to a panel of bactericidal

antibiotics. Triclosan MICs for *E. coli* and MRSA were 200 ng/mL and 100 ng/mL, respectively

under our growth conditions, equivalent to the urine concentration of individuals using triclosan-

containing consumer products[12,13]. In all cases, triclosan was added 30 minutes prior to the

addition of the specified bactericidal antibiotic and both antibiotics maintained in the culture for

the remainder of the experiment.

We utilized two approaches to evaluate the impact of triclosan on bacterial tolerance and

persistence: a qualitative, end-point plating efficiency assay and a quantitative kinetic kill-curve.

Plating efficiency assesses relative antimicrobial tolerance and the kinetic kill-curve

distinguishes between changes in total antibiotic tolerance and changes in persister frequency[4].

Triclosan had a dramatic protective effect on *E. coli* in the end point assay, increasing survival

by several orders of magnitude in the presence of three bactericidal antibiotics and providing

nearly complete protection against a fourth (Figure 1). *E. coli* treated with triclosan exhibited a

1000-fold increase in survival in the presence of 50 μg/mL (~5x MIC) kanamycin, an inhibitor of

peptide bond formation, and a 10,000-fold increase in survival in the presence of streptomycin

(50 μg /mL: ~2x MIC), an inhibitor of tRNA-ribosome interaction, and ciprofloxacin (100

ng/mL: ~3x MIC) a gyrase inhibitor. (Figure 1). Strikingly, triclosan rendered *E. coli* almost

completely refractile to treatment with the cell wall active antibiotic ampicillin (100 μg /mL;

~10x MIC). Viable cell numbers were essentially identical in triclosan and triclosan-ampicillin

treated cultures at 2 hours, and 10% of cells in triclosan-ampicillin cultures were viable at 20

hours, suggesting triclosan qualitatively increased persister frequency.

To further assess the protective effect of Triclosan, we next performed a kinetic kill curve, in

which we measured colony forming units (CFU) over a 20-hour time, focusing on ciprofloxacin,

the broad spectrum antibiotic used to treat *E. coli* related UTIs. Consistent with the results of the

end point assay, triclosan substantially protected *E. coli* from ciprofloxacin-induced cell death

throughout the duration of the time course. Protection was particularly pronounced at the 2-hour

time point, where the slope of the kill curve for pre-treated cells diverges substantially from that

of untreated cells (Figure 2). A divergent slope is diagnostic for the presence of persister cells in

the pre-treated population[13].

Persister population size was proportional to the concentration of the bacteriocidal antibiotic:

10% of triclosan treated cells cultured at 100 ng/mL ciprofloxacin remained viable at 2 hours,

while only 0.1% cultured at the more clinically relevant 1,000 ng/mL ciprofloxacin were viable

at the same time point. For perspective, 0.1% of the population is equivalent to 1 in 10^3 cells,

1000-fold higher than the expected frequency of persisters in an untreated population[4]. At the 20

hour time point, 90,000 cells per mL were viable in 100 ng/mL ciprofloxacin and 30 cells per

mL in 1000ng/mL ciprofloxacin. In contrast, we observed only 20 cells/ml after 20 hours of

growth in 100 ng/mL ciprofloxacin alone. Cells cultured in 1,000 ng/mL ciprofloxacin alone had

no observable colonies (<10 cells per mL).

Remarkably, triclosan also protected MRSA cells from high concentration of the cell wall

inhibitor vancomycin over the course of a 20-hour experiment (Figure 2C). MRSA treated with

100 ng/ml of triclosan were essentially refractile to 50 ng/ml vancomycin (10x the MIC) at 4

hours and exhibited a viable cell count 200-times that of untreated cells at 8 hours. Although not

statistically significant, at 20 hours the viable cell count was several times higher in the presence

of both triclosan and vancomycin than vancomycin alone (360,000 +/- 200,000 cells/ml versus

130,000 +/- 80,000 cells/ml), consistent with induction of a persistent state.

The ability of triclosan to protect cells from bactericidal antibiotics appears to be independent of

its impact on bacterial growth rate. Instead, genetic data support a model in which triclosan-

mediated tolerance stems from accumulation of the alarmone ppGpp in response to defects in

fatty acid synthesis[15]. ppGpp has been repeatedly implicated in antibiotic tolerance and

persistence in several organisms including *E. coli* and *S. aureus* [14-16]. To determine the

contribution of ppGpp accumulation to triclosan mediated tolerance, we compared the relative

viability of wild type and $ppGpp^0$ (*spoT*::*cat* Δ*relA*) *E. coli* cells after 2 hours antibiotic

challenge in the presence or absence of triclosan. Triclosan was unable to substantially protect

$ppGpp^0$ cells from either ampicillin or ciprofloxacin (Figure 2E). For kanamycin and

streptomycin, the ppGpp^0 cells showed no viable cells after 2 hours for both the triclosan-treated

and untreated cells. In contrast to previous studies which relied on either carbon starvation or the

addition of serine hydroxamate to induce accumulation of high concentrations of ppGpp [100x

above baseline][16], defects in fatty acid synthesis have at best a modest impact on ppGpp

levels[~5x over baseline][17]. Thus even relatively low levels of the alarmone appear to be

sufficient to protect cells from a panel of antimicrobials.

Triclosan inhibited growth in the presence and absence of the alarmone, arguing against a

simplistic model in which reductions in growth rate are inherently protective (Figure 2F).

Specifically how modest increases in [ppGpp] might confer tolerance to different antibiotics thus

remains an open question. We favor the idea that rather than exerting a global effect, ppGpp

mediates changes in individual biosynthetic pathways that render them tolerant of their cognate

antimicrobial. For example, ppGpp-dependent down regulation of ribosomal RNA synthesis

significantly curtails translation, conferring resistance to the translational inhibitors kanamycin

and streptomycin. Similarly, increases in [ppGpp] are reported to curtail DNA replication—both

elongation and initiation—providing a straightforward explanation for ppGpp-mediated

ciprofloxacin resistance.

Regardless of mechanism, these data highlight an unexpected and potentially important role for

triclosan as a contributor to antibiotic tolerance and persistence in both community and health

care settings. While the impact of triclosan on persistence and tolerance *in vivo* remains an open

question, the high concentration of the compound in the urine of individuals using triclosan-

containing products raises the possibility that it might interfere with treatment of bladder and

kidney associated infections. Along with previous work identifying a connection between

triclosan exposure and antibiotic resistance[18,19], our findings reinforce the need for substantial

caution—as well as consideration of unintended consequences—in evaluating the costs and

benefit of antimicrobial additives in consumer products.

Methods

Materials and Strains

Triclosan, ampicillin, kanamycin, streptomycin, ciprofloxacin, and vancomycin were purchased

from Sigma-Aldrich. Stock solutions were made in water for ampicillin (100 mg/mL),

kanamycin (50 mg/mL), streptomycin (100 mg/mL) and ciprofloxacin (10 mg/mL). Triclosan

was dissolved in ethanol (10 mg/mL), and vancomycin was dissolved in DMSO (100 mg/mL). *E.*

coli MG1655 and *S. aureus* FPR3575 were both lab strains. The ppGpp0 strain was created by

transducing spoT::cat (the kind gift of Dr. Jue D. Wang) and relA::ka$n$[20] into MG1655 using the

bacteriophage P1*vir*. To test kanamycin tolerance of the ppGpp0, we first cured the kanamycin

cassette using pCP20, and the pCP20 plasmid was then removed through growth of multiple

generations at 42°C[21]. *E. coli* was grown in Luria-Bertani broth (LB) and *S. aureus* was grown

in tryptic soy broth (TSB). Growth temperature was 37°C for all experiments.

Determination of Minimum Inhibitory Concentration (MIC)

To determine the MIC for the panel of antibiotics utilized in this study, *E. coli* and *S. aureus*

were grown to OD-600 = 0.1 in LB or TSB respectively. Cells were then back-diluted 1000-fold

and transferred to a 96-well plate containing 2-fold dilutions of respective antibiotics and

cultured at 37° for 16 additional hours with vigorous shaking in a BioTek Eon plate reader. MIC

was calculated as the lowest antibiotic concentration preventing development of detectable

turbidity at OD-600.

Assays for antibiotic Tolerance and Persistence

To assay tolerance and persistence, *E. coli* and *S. aureus* were grown to an OD 600 = 0.2 in LB

or TSB respectively. Cells were then back-diluted into media containing triclosan at indicated

concentrations to an OD-600 = 0.1, cultured for an additional 30 minutes, before being

challenged with bactericidal antibiotics. For dot plating, 10 μL of a 10-fold dilution series was

plated on antibiotic free LB-agar or TSB-agar as appropriate. For determination of colony

forming units (CFU), 100 μL of a 10-fold dilution series was spread on antibiotic free LB-agar or

TSB-agar plates. Cells were incubated for ~12 hours at 37° prior to quantification. CFUs were

normalized to CFUs at t0 to correct for the ~2-fold increase in cell number in untreated cultures

during the 30 minute pre-treatment period. Relative persistence is defined as the CFU's of the

triclosan treated sample divided by the CFU's of the non-treated sample.

Growth Curve

176   MG1655 and MG1655 ppGpp^0 cells were grown to an OD600=0.1-0.2 in LB before back-

177   diluting to an OD600=0.02 into LB with or without 200 ng/mL triclosan. Growth was monitored

178   every 30 minutes at OD600.

Statistical Analysis

181   All values are expressed as the mean ± standard deviation of n=3 replicates. Data was analyzed

182   using a one-tailed Student's t-test. Statistical significance was determined when $p < .05$.

Acknowledgements

This work was supported by National Institutes of Health grants GM64671 to P.A.L. C.S.W was supported by an Arnold O. Beckman Postdoctoral Fellowship. The authors thank Dr. Robyn Klein and members of the Levin and Zaher groups for insightful suggestions and critical comments on the manuscript. We are grateful to Dr. Jue Wang for the kind gift of the *spoT*::*cat* strain.

Author Contributions

C.S.W. and P.A.L designed the experiments. C.S.W. performed the experiments. C.S.W. and P.A.L. analyzed the data and prepared the manuscript.

References

1. Bell, B.G. *et al*. *BMC Infections Diseases* **14**, 13 (2014).
2. McEwen, S.A. & Fedorka-Cray, P.J. *Clin. Infect. Dis.* **34**, S93-106 (2002).
3. Yazdankhaha, S.P. *et al*. *Microb. Drug Resist.* **12,** 83-90 (2006).
4. Lewis, K. *Annu. Rev. Microbiol.* **64**, 357-372 (2010).
5. Brauner, A., Fridan, O., Gefen, O., & Balaban, N.Q. *Nat. Rev. Microbiol.* **14,** 320-330 (2016).
6. Bigger, J.W. *Lancet* **244,** 497-500 (1944).
7. Mulcahy, L.R., Burns, J.L., Lory, S., & Lewis, K. *J. Bacteriol.* **192,** 6191-6199 (2010).
8. Proctor, R.A. *et al*. *Clin. Infect. Dis.* **20**, 95-102 (1995).
9. Balaban, N.Q. *et al*. *Science* **305**, 1622-1625 (2004).
10. FDA Safety and Effectiveness of Consumer Antiseptics; Topical Antimicrobial Drug Products for Over-the-Counter Human Use, 21 C.F.R. § 310 (2016).
11. McMurry, L.M., Oethinger, M., & Levy, S.B. *Nature* **394**, 531-532 (1998).
12. Calafat, A.M. *et al*. *Environ. Health Perspect.* **116**, 303-307 (2008).
13. MacIsaac, J.K. *et al*. *J. Occup. Environ. Med.* **56**, 834-839 (2014).
14. Nguyen, D. *et al*. *Science* **334**, 982-986 (2011).
15. Battesti, A. & Bouveret E. *Mol Microbiol.* **62**, 1048-1063 (2006).
16. Maisonneuve, E., Castro-Camargo M., Gerdes, K. *Cell* **154**, 1140-1150 (2013).
17. Seyfzadeh, M., Keener, J., & Nomura, M. *PNAS* 90, 11004-8 (1993).
18. Karatzas K.A. *et al*. *J. Antimicrob. Chemother*. **60**, 947-955 (2007).
19. Webber, M.A., Randall, L.P., Cooles, S., Woodward, M.J., Piddock L.J. *J. Antimicrob. Chemother*. **62**, 83-91 (2008).
20. Baba T. *et al. Mol. Syst. Biol.* **2**, 2006.0008 (2006).
21. Cherepanox, P.P. & Wackernagel W. *Gene* **156**, 9-14 (1995).

233 **Figures**

249 **Figure 1: Triclosan induces tolerance to multiple antibiotics.** *E. coli* (MG1655) were cultured

250 to OD600 = 0.2, split and cultured for an additional 30 minutes with (+T) or without 200ng/ml

251 triclosan (-T). Indicated bactericidal antibiotics were then added and cells cultured for an

252 additional 2 to 20 hours prior to dilution plating. Each experiment was replicated three

253 independent times with only representative data shown.

Figure 2: Kinetic analysis of triclosan-induced persistence. *E. coli* (MG1655) and MRSA

(FPR3757) cells were cultured to OD600 = 0.2, split and cultured for an additional 30 minutes

with (black line, closed squares) or without triclosan (grey line, open squares). At t=0, 100

ng/mL (A) or 1 µg/mL ciprofloxacin (B) was added to *E. coli* cultures and 50ng/ml vancomycin

was added to MRSA cultures (C). Relative persistence in the presence of triclosan

(CFU+T/CFU-T) was calculated for the 4 and 20 hour time points (D). Cell viability of ppGpp^0

E. coli with (+T) or without (-T) pretreatment with triclosan after 2 hour challenge with

ampicillin or vancomycin (E). Growth curves of MG1655 (black curve) or ppGpp^0 (gray curve)

in LB with (dashed lines) or without (solid lines) triclosan (F). Values are the mean of three

independent biological replicates with error bars indicating one standard deviation. Asterisks

represent significant difference between the triclosan treated and non-treated using a Student's t-

test with $p < 0.05$.