1

The commercial antimicrobial triclosan induces high levels of antibiotic

tolerance *in vitro* and reduces antibiotic efficacy up to 100-fold *in vivo*

Corey Westfall[1,2], Ana Lidia Flores-Mireles[1,3,5], John Isaac Robinson[4], Aaron J.L. Lynch[3], Scott

Hultgren[3], Jeffrey P. Henderson[4], & Petra Anne Levin[2,6],

[2]Department of Biology, Washington University in St. Louis, St. Louis, MO 63130, USA

[3]Department of Molecular Microbiology and Microbial Pathogenesis, Washington University
School of Medicine, St. Louis, MO 63110, USA

[4]Division of Infectious Diseases, John T. Milliken Department of Internal Medicine, Washington
University School of Medicine, St. Louis, MO 63110, USA

[5]Department of Biological Sciences, University of Notre Dame, Notre Dame, Indiana, USA.

16
17

[1]These two authors contributed equally to this study

[6]To whom correspondence should be addressed: plevin@wustl.edu

26
27
28

Abstract

The antimicrobial triclosan is used in a wide range of consumer products ranging from

toothpaste, cleansers, socks, and baby toys. A bacteriostatic inhibitor of fatty acid synthesis,

triclosan is extremely stable and accumulates in the environment. Approximately 75% of adults

in the US have detectable levels of the compound in their urine, with a sizeable fraction of

individuals (>10%) having urine concentrations equal to or greater than the minimal inhibitory

concentration for *Escherichia coli* and methicillin-resistant *Staphylococcus aureus* (MRSA).

Previous work has identified connections between defects in fatty acid synthesis and

accumulation of the alarmone guanosine tetraphosphate (ppGpp), which has been repeatedly

associated with antibiotic tolerance and persistence. Based on these data, we hypothesized that

triclosan exposure may inadvertently drive bacteria into a state in which they are able to tolerate

normally lethal concentrations of antibiotics. Here we report that clinically relevant

concentrations of triclosan increased *E. coli* and MRSA tolerance to bactericidal antibiotics as

much as 10,000 fold *in vitro* and reduced antibiotic efficacy up to 100-fold in a mouse urinary

tract infection model. Genetic analysis indicated that triclosan-mediated antibiotic tolerance

requires ppGpp synthesis, but is independent of growth. These data highlight an unexpected and

certainly unintended consequence of adding high concentrations of antimicrobials in consumer

products, supporting an urgent need to reevaluate the costs and benefits of the prophylactic use

of triclosan and other bacteriostatic compounds.

Importance

Added as a prophylactic to a wide range of consumer products, the fatty acid synthesis inhibitor

triclosan accumulates to high levels in humans and the environment. Based on links between

defects in fatty acid synthesis and accumulation of the alarmone ppGpp, we hypothesized that

triclosan would render cells tolerant to bactericidal compounds due to ppGpp-mediated

inhibition of biosynthetic capacity. Our data indicate that clinically relevant concentrations of

triclosan induces higher tolerance of *E. coli* and methicillin resistant *S. aureus* (MRSA) to a

panel of bactericidal antibiotics up to 10,000-fold. In a urinary tract infection model, mice

exposed to triclosan exhibited bacterial loads ~100-fold higher in the bladder than control

animals following ciprofloxacin challenge. These findings highlight an unexpected consequence

of antimicrobials in consumer products and support an urgent need to reevaluate the costs and

benefits of the prophylactic use of triclosan and other bacteriostatic compounds.

Introduction

The prophylactic use of antibiotics in consumer goods, ranging from animal feed to personal care

products, is widely believed to be a major contributor to the epidemic increase in antibiotic-

resistant pathogens (1-3). Prominent among these prophylactics are triclosan and triclocarban,

polychlorinated aromatic antimicrobials targeting fatty acid synthesis. Triclosan in particular is

found in a wide variety of consumer products, including: toothpaste, cleansers, socks, and baby

toys (1). Although the US Food and Drug Administration effectively banned the use of triclosan

in household soap in 2017, as of this writing Canada and Australia, among other countries, have

not elected to take similar action.

An inhibitor of enoyl-acyl carrier protein reductase, (4), at low concentrations (200 ng/ml)

triclosan is bacteriostatic, preventing cell growth but having little effect on viability over the

short term. At high concentrations (>10 μg/ml), triclosan is bactericidal, most likely killing cells

through disruption of plasma membrane integrity (4). Recent work from the Waters lab suggests

that at these higher concentrations triclosan can serve as an adjuvant, acting synergistically with

tobramycin and other drugs, increasing killing by ~100-fold in a *Pseudomonas aeruginosa*

biofilm model for CF lung infections (5). Triclosan is typically used as an antimicrobial additive

at these higher, bactericidal concentrations.

Because of its widespread use as a prophylactic, the high concentrations at which it is employed,

and its inherent stability, triclosan accumulates to high levels in the environment (6, 7).

Approximately 75% of adults in the US have detectable levels of the compound in their urine,

and >10% have urine concentrations greater than or equal to the minimal inhibitory

concentration for *Escherichia coli* (200 ng/mL) and methicillin-resistant *Staphylococcus aureus*

(MRSA) (100 ng/mL) (8, 9).

While the inverse relationship between antibiotic use and antibiotic efficacy is largely

attributable to the selection of heritable traits, non-heritable traits such as antibiotic tolerance and

persistence are also likely to be involved (10). In contrast to genetically-resistant bacteria, which

grow in the presence of an antibiotic, tolerant bacteria are able to survive antibiotic challenge for

longer periods of time than their more sensitive counterparts (10). Persister cells are the small

sub-set of an otherwise-sensitive population (~1 in 10^6) that exhibit levels of tolerance sufficient

to protect them from otherwise lethal concentrations of antimicrobial compounds (11). Increases

in antibiotic tolerance and persistence are confounding factors in the treatment of chronic *P*.

aeruginosa (12) and *S. aureus* (13) infections and are thought to contribute to the refractory

nature of medically relevant biofilms (14). Reduced growth rate and metabolic activity is

associated with increased antibiotic tolerance (10) and is a defining trait of persister cells.

Based on previous work identifying connections between defects in fatty acid synthesis and

accumulation of the alarmone guanosine tetraphosphate (ppGpp) (15), and the possible link

between ppGpp and antibiotic tolerance (16), we hypothesized that triclosan exposure may

inadvertently drive bacteria into a metabolically depressed state in which they are able to tolerate

normally lethal concentrations of antibiotics (17, 18). In particular, inhibiting fatty acid

synthesis stimulates interaction between acyl carrier protein and the hydrolase domain of the

bifunctional ppGpp synthase SpoT, resulting in accumulation of the alarmone and the

concomitant inhibition of biosynthetic capacity (19).

Here we report that clinically-relevant bacteriostatic concentrations of triclosan increased *E. coli*

and methicillin resistant *Staphylococcus aureus* (MRSA) tolerance to bactericidal antibiotics as

much as 10,000-fold *in vitro* and reduced antibiotic efficacy ~100-fold in a mouse urinary tract

infection model. Triclosan-mediated antibiotic tolerance is dependent on ppGpp synthesis:

although triclosan inhibited the growth of both wild-type and ppGpp mutant cells, only the latter

were highly susceptible to challenge with bactericidal compounds. In contrast, pretreatment with

another bacteriostatic drug, spectinomycin, a translation inhibitor that does not impact ppGpp

accumulation (20), induced high levels of antibiotic tolerance in both wild-type and ppGpp

mutant cells. Together, these data highlight an unexpected and certainly unintended

consequence of employing triclosan as a commercial antimicrobial, and support an urgent need

to reevaluate the costs and benefits of the addition of triclosan, and potentially other

bacteriostatic compounds, to consumer products.

Results

Triclosan pretreatment results in high levels of tolerance to bactericidal antibiotics *in vitro*

To assess if physiologically relevant levels of triclosan are sufficient to promote tolerance to

bactericidal antibiotics, we examined the relative sensitivity of *E. coli* (MG1655) and *S. aureus*

(FPR3757 an USA-300 MRSA strain) cultured in minimal inhibitory concentrations (MIC) of

triclosan to a panel of bactericidal antibiotics. Triclosan MICs for *E. coli* and MRSA were 200

ng/mL and 100 ng/mL, respectively under our growth conditions; similar to the triclosan

concentration found in the urine from individuals using triclosan-containing products (8, 9). In

all cases, triclosan was added 30 minutes prior to the addition of the specified bactericidal

antibiotic and both antibiotics were maintained in the culture for the remainder of the

experiment.

Triclosan had a dramatic protective effect on *E. coli* in an end point assay, increasing survival by

several orders of magnitude in the presence of three bactericidal antibiotics and providing nearly

complete protection against a fourth (Fig. 1). *E. coli* treated with triclosan exhibited a 1,000-fold

increase in survival in the presence of 50 μg/mL (~5x MIC) kanamycin, an inhibitor of peptide

bond formation. It also showed a 10,000-fold increase in survival in the presence of streptomycin

(50 μg /mL: ~2x MIC), an inhibitor of tRNA-ribosome interaction, and ciprofloxacin (100

ng/mL: ~3x MIC) a gyrase inhibitor (Fig. 1). Strikingly, triclosan rendered *E. coli* almost

completely refractory to treatment with the cell wall active antibiotic ampicillin (100 μg /mL;

~10x MIC). Viable cell numbers were essentially identical in triclosan and triclosan-ampicillin

treated cultures at 2 hours, and 10% of cells in triclosan-ampicillin cultures were viable at 20

hours, suggesting triclosan increased persister frequency to all tested antibiotics.

Triclosan also protected MRSA cells from high concentrations of the glycopeptide antibiotic,

vancomycin, over the course of a 20-hour experiment (Fig. 2c). MRSA treated with 100 ng/ml of

triclosan were essentially refractory to 50 ng/ml vancomycin (10x the MIC) at 4 hours and

exhibited a viable cell count 200x that of untreated cells at 8 hours. Even at 20 hours the viable

cell count was several times higher in the presence of both triclosan and vancomycin than

vancomycin alone. This delayed reduction in viable cell count is consistent with induction of a

persistent state (10).

Triclosan increases persister cell frequency

To further assess the protective effect of triclosan, we next performed kinetic kill curves, in

which we measured colony-forming units (CFU) over a 20-hour time frame. If persister cells are

present, we expect to observe two slopes, one corresponding to the kill rate of the general

population and the second slope corresponding to the slower kill rate of persister cells (10). In

this manner, the kill curve is able to separate antibiotic tolerance at the population level from the

impact of triclosan treatment on persister levels.

For these experiments, we focused on ciprofloxacin, the broad-spectrum antibiotic used to treat

urinary tract infections (UTIs). Consistent with the results of the end point assay (Fig. 1),

triclosan substantially protected *E. coli* from ciprofloxacin-induced cell death throughout the

duration of the time course (Fig. 2a, b). Protection was particularly pronounced at the 2-hour

time point, where the slope of the kill curve for pre-treated cells diverged substantially from that

of untreated cells (Fig. 2a). A reduced kill rate suggests that the pre-treated population contains a

larger proportion of persister cells (21).

In agreement with previous work (16), persister population size was proportional to the

concentration of ciprofloxacin. 10% of triclosan treated MG1655 cells cultured in 100 ng/mL

ciprofloxacin remained viable after 2 hours (Fig. 2a), while only 0.1% cultured in the more

clinically-relevant 1,000 ng/mL ciprofloxacin were viable at the same time point (Fig. 2b). For

perspective, 0.1% is 1,000-fold higher than the expected frequency of persisters in an untreated

population (10). After 20 hours, 90,000 CFU/mL were viable in cultures treated with both

triclosan and 100 ng/mL ciprofloxacin (Fig. 2a), compared to 20 cells/mL in cultures treated with

ciprofloxacin alone. At 1,000 ng/mL ciprofloxacin, cultures treated with both triclosan and

ciprofloxacin contained 30 viable cells per mL (Fig. 2b). In contrast, no viable cells were

detected (<10 cells/mL) in cultures treated with 1,000 ng/mL ciprofloxacin alone. We observed

an increase in the abundance of persisters at drug concentrations above 1,000 ng/mL (Fig S1)

(16). Although this finding initially appeared counterintuitive, it is consistent with previous

reports suggesting that prophage induction in response to DNA damage is responsible for cell

death at lower concentrations of ciprofloxacin, while higher concentrations kill cells before

prophage are induced (16).

Triclosan-mediated tolerance requires ppGpp

Based on the well-established connection between defects in fatty acid synthesis and

accumulation of ppGpp, we speculated that triclosan-mediated tolerance was dependent on the

synthesis of the alarmone (15). To test this idea, we compared the relative viability of wild-type

E. coli and mutants unable to synthesize the alarmone (ppGpp0; *spoT*::*cat ΔrelA*) 2 hours after

antibiotic challenge in the presence or absence of triclosan.

Although triclosan inhibited the growth of both wild-type and ppGpp0 cells, it was unable to

substantially protect ppGpp0 cells from any of the four bactericidal antibiotics we tested. These

include ampicillin and ciprofloxacin, as well as the translation inhibitors, kanamycin and

streptomycin (Fig. 3a, b). The ppGpp0 cells were more sensitive to kanamycin and streptomycin,

showing no viable cells after 60-minute treatment, thus measurements were performed at 30

minutes. Importantly, triclosan alone was not bactericidal to either wild-type or ppGpp0 cells

(Fig. S2).

In contrast to triclosan, pre-treatment with another bacteriostatic compound, spectinomycin,

increased tolerance to kanamycin, streptomycin, ampicillin, and ciprofloxacin in both wild-type

and ppGpp0 mutant cells (Fig. 3C). A translation inhibitor, spectinomycin does not impact

ppGpp levels in *E. coli* (20). While spectinomycin was still protective in the ppGpp0 cells, levels

of protection were slightly decreased compared to WT cells.

Triclosan drives tolerance to ciprofloxacin in a murine model

Due to its widespread use and inherent stability, triclosan is present in both human populations

and the environment (8). Thus, a key question is whether the tolerance we observed *in vitro* is

relevant *in vivo*. To determine the physiological relevance of triclosan-mediated tolerance, we

employed a mouse model of *E. coli* UTI. UTIs are one of the most prevalent bacterial infections,

impacting approximately 150 million people annually (22). Uropathogenic *E. coli* (UPEC) is the

main causative agent of both uncomplicated and complicated UTI (23). Pretreatment with

triclosan rendered the well-characterized *E. coli* cystitis isolate UTI89 ~10-fold more tolerant to

1,000 ng/mL ciprofloxacin than untreated cells at 2 hours, a level equivalent to the tolerance we

observed for *E. coli* MG1655 at the same time point (Fig. 1 and Fig. S3).

For *in vivo* experiments, we provided six-week old female wild-type C3H/HeN mice with

drinking water containing 1,000 ng/mL triclosan for 21 days. Control mice were given plain

water for the same duration. At 21 days, experimental and control mice were trans-urethrally

infected with ~5 x10^7 CFU of *E. coli* UTI89. At 24 hours post-infection, a subset of the mice was

treated with intraperitoneal ciprofloxacin (25 mg/kg). At 48 hours post-infection, all mice were

sacrificed and bacterial colonization assessed in urine and bladder.

After ciprofloxacin treatment, bacterial titers were >100-fold higher in the urine (p<0.0001) and

>10-fold higher (p<0.0001) in the bladders of triclosan-treated mice versus control animals (Fig.

4a, b), consistent with triclosan-induced tolerance occurring *in vivo*. Bacterial load at 24 hours

post-infection was nearly equivalent in triclosan-treated and control mice, indicating that

triclosan did not significantly impair UTI89 viability (Fig. 4a, b). Treated mice had triclosan

levels between 70-750 ng/mL in their urine, comparable to the MIC for *E. coli* (200 ng/mL) and

similar to reported triclosan levels in human urine (2.4 to 3,790 ng/mL) (8) (Fig. 4c). We also

detected two putative metabolized forms of triclosan, one with a mass consistent with the

previously reported sulfonated triclosan (24), and the other 96 daltons larger (Fig. S4). Whether

or not the modified forms of the drug are active against bacteria is unclear. Control mice had no

observable triclosan (below 1.6 ng/mL limit of detection).

Discussion

Our data indicate that environmentally relevant concentrations of triclosan reduce antibiotic

efficacy as much a 100-fold *in vivo* (Fig. 4) and highlight an unexpected and potentially

important role for triclosan as a contributor to antibiotic tolerance and bacterial persistence in

both community and healthcare settings. Triclosan-mediated tolerance is dependent on ppGpp

synthesis, most likely in response to inhibition of fatty acid synthesis (Fig. 3a) (15). This finding

is consistent with prior work implicating ppGpp in antibiotic tolerance and persister development

(16).

In contrast to previous studies of ppGpp-induced persistence that relied on either carbon

starvation or the addition of serine hydroxamate to induce accumulation of high concentrations

of ppGpp (100x above baseline) (25, 26), defects in fatty acid synthesis have at best a modest

impact on ppGpp levels (~5x over baseline) (15). This suggests that even relatively low levels of

ppGpp are sufficient to protect cells from a panel of antimicrobials. Specifically how modest

increases in ppGpp might confer tolerance to different antibiotics thus remains an open question.

We favor the hypothesis that ppGpp mediates changes in individual biosynthetic pathways that

render them tolerant of their cognate antimicrobial. For example, ppGpp-dependent down-

regulation of ribosomal RNA synthesis significantly curtails translation (27), potentially

conferring tolerance to the translational inhibitors kanamycin and streptomycin. Similarly,

increases in ppGpp are reported to curtail DNA replication (28)—both elongation and

initiation—providing a straightforward explanation for ppGpp-mediated ciprofloxacin resistance.

It is generally recognized as poor practice to prescribe a bacteriostatic compound prior to or

along with delivery of a bactericidal one (29), because of the potential that the former will

interfere with the activity of the latter. At the same time, their mechanisms of action, and

therefore the mechanisms by which these drugs drive tolerance, are likely to differ widely. While

the translation inhibitor spectinomycin provides protection against bactericidal compounds, it

does not induce accumulation of ppGpp (20), a fact supported by our finding that spectinomycin

induces tolerance to bactericidal compounds in both wild-type and ppGpp0 cells (Fig. 3c). At the

same time, ppGpp-dependent induction of antibiotic tolerance is likely to be a feature triclosan

shares with a related compound, triclocarban, which also inhibits an early step in fatty acid

synthesis, and is also a common additive in consumer products. Triclosan also stands out from

other bacteriostatic compounds by virtue of its widespread use and sheer abundance in the

environment. ~1 kg of Triclosan is produced for every 3 kg of other antimicrobials and estimates

indicate that ~100 metric tonnes are being deposited annually in the environment through waste

water treatment in the US alone (30).

Although triclosan has low toxicity (LD-50 4,350 mg/kg orally) (31), accumulating data link

long-term exposure with antibiotic resistance (32) and there are reports that triclosan may also

function as an endocrine disrupter (33, 34). Our analysis of the impact of triclosan on antibiotic

efficacy in a mouse UTI model (Fig. 4) highlights yet another deleterious "side effect" of this

ubiquitous antimicrobial. UTIs alone impact 150 million people worldwide (22) at a cost of $3.5

billion per year in the US alone (35). Complications associated with UTIs include pyelonephritis

with sepsis, renal damage, pre-term birth, *Clostridium difficile* colitis, sepsis, and death

particularly in the very old and the very young (23). Coupled with the well established

283 connection between antibiotic tolerance and recurrent/chronic infections (12, 13), our findings

284 reinforce the need for substantial caution—as well as consideration of unintended

285 consequences—in evaluating the costs and benefits of antimicrobial additives in consumer

286 products.

288 **Methods**

289 **Materials and Strains**

290 Triclosan, ampicillin, kanamycin, streptomycin, ciprofloxacin, and vancomycin were purchased

291 from Sigma-Aldrich. Stock solutions were made in water for ampicillin (100 mg/mL),

292 kanamycin (50 mg/mL), streptomycin (100 mg/mL) and ciprofloxacin (10 mg/mL). Triclosan

293 was dissolved in ethanol (10 mg/mL), and vancomycin was dissolved in DMSO (100 mg/mL). *E.*

294 *coli* MG1655 and *S. aureus* FPR3575 were both lab strains and *E. coli* UTI89 was isolated from

295 a patient with a urinary tract infection (36). *E. coli* were grown in Luria-Bertani broth (LB) and

296 *S. aureus* was grown in tryptic soy broth (TSB). Growth temperature was 37 °C for all

297 experiments.

299 **Determination of Minimum Inhibitory Concentration (MIC)**

300 To determine the MIC for the panel of antibiotics utilized in this study, *E. coli* and *S. aureus*

301 were grown to OD-600 = 0.1 in LB or TSB, respectively. Cells were then back-diluted 1,000-

302 fold and transferred to a 96-well plate containing 2-fold dilutions of respective antibiotics and

303 cultured at 37 °C for 16 additional hours with vigorous shaking in a BioTek Eon plate reader.

304 MIC was calculated as the lowest antibiotic concentration, preventing development of detectable

305 turbidity at OD-600.

Assays for antibiotic tolerance and persistence

To assay tolerance and persistence, *E. coli* and *S. aureus* were grown to an OD-600 = 0.2 in LB or TSB, respectively. Cells were then back-diluted to an OD-600 = 0.1 in media containing triclosan at indicated concentrations and cultured for an additional 30 minutes, before being challenged with bactericidal antibiotics. For dot plating, 10 µL of a 10-fold dilution series was plated on antibiotic-free LB agar or TSB agar as appropriate. For determination of colony forming units (CFU), 100 µL of a 10-fold dilution series was spread on antibiotic-free LB agar or TSB agar plates. Cells were incubated for ~12 hours at 37 °C prior to quantification. CFUs were normalized to CFUs at the initial time point to correct for the ~2-fold increase in cell number in untreated cultures during the 30-minute pre-treatment period. Relative persistence is defined as the CFU's of the triclosan treated sample divided by the CFU's of the non-treated sample.

UTI mouse work

Six-week old female wild-type C3H/HeN mice were obtained from Envigo. Mice were treated with or without 1,000 ng/mL (100 ppm) triclosan in the drinking water for 21 days. At 21 days, mice were anesthetized by inhalation of 4% isoflurane and mouse bladders were trans-urethrally infected with approximately 5 x10^7 CFU of *E. coli* UTI89 in 50µl PBS(37). Briefly, a single UTI89 colony was inoculated in 20 ml of Luria Broth (LB) and incubated at 37 °C under static conditions for 24 hours. Bacteria were then diluted (1:1,000) into fresh LB and incubated at 37 °C under static conditions for 18 to 24 hours. Bacteria were subsequently washed three times with PBS and then concentrated to approximately 5 x10^7 CFU per 50 µL. At 24 hours post-infection, mice received 25mg/kg of ciprofloxacin intraperitoneal. 48 hours post-infection, mice were euthanized, and bladders were harvested and urine was collected. Bladders were

homogenized in PBS and bacterial load present in bladders and urines was determined by plating

serial dilutions on LB agar supplemented with antibiotics when appropriate. Statistical analyses

were performed using the Mann–Whitney U test with GraphPad Prism software (version 6.0 for

Mac). All animal studies were performed in accordance with the guidelines of the Committee for

Animal Studies at Washington University School of Medicine.

Measurement of triclosan and metabolites in mouse urine

Since triclosan has been observed to adsorb to plastic surfaces, sample handling was performed

in glass vessels whenever possible (24). A stock solution of 1 mg/mL triclosan (Sigma) was

prepared in methanol and a 100 μg/mL $^{13}C_{12}$-triclosan (99%) internal standard in MTBE was

purchased from Cambridge Isotope Laboratories (Andover, MA). A dilution series of 1,000, 200,

40, 8, 1.6, and 0.32 ng/mL triclosan was prepared in pooled, untreated mouse urine and spiked

with 100 ng/mL $^{13}C_{12}$-triclosan internal standard. Samples were diluted 1:1 in methanol, spun

down at 20,000×g for 10 minutes, and filtered through 0.45 μm 13 mm diameter PVDF syringe

filters (Millipore). Finally, cleaned samples were diluted 1:1 in HPLC-grade water (Sigma).

Using a Shimadzu UFLC (Kyoto, Japan), 10 μL of each sample was injected onto a fused core

phenyl-hexyl column (100 mm × 2 mm × 2.7 μm) with a 0.4 mL/min flow rate (Ascentis

Express, Supelco). Triclosan was eluted from the column as follows: Solvent A (0.1% formic

acid) and Solvent B (90% acetonitrile with 0.1% formic acid) were held constant at 80% and

20%, respectively, for 0.1 minutes. Solvent B was increased to 98% by 5 minutes, held at 98%

for 1 minutes, and then reduced again to 20% in 0.1 minutes. The column was equilibrated in

20% Solvent B for 3 minutes between runs.

Triclosan was detected using an AB Sciex API 4000 QTrap mass spectrometer (AB Sciex, Foster

City, CA) running in negative ion electrospray ionization mode (ESI) using a Turbo V ESI ion

source. Triclosan was detected using the instrument settings listed in Supplementary Data Table

1. A precursor ion scan was performed for the 35 m/z product ion to determine the mass

spectrum of triclosan, $^{13}C_{12}$-triclosan, and any potential metabolites (Fig. S4a). Because triclosan

contains three chlorine atoms, its mass spectrum includes prominent isotope peaks (M+2, M+4)

corresponding to the natural abundance of ^{37}Cl (Fig. S4b). To improve sensitivity, product ions

from the two most abundant isotopologues were detected and added together prior to peak

integration. Peaks for triclosan and internal standard were integrated with Analyst software (AB

Sciex) and normalized. Normalized peak areas varied linearly with triclosan concentration above

1.6 ng/mL.

Pooled urines from 3 to 4 mice were spiked with 100 ng/mL internal standard and cleaned as

described above. Samples were analyzed by LC-MS/MS and triclosan was quantified using the

standard curve (Fig. S4e).

Statistical analysis

Values for the *in vitro* data are expressed as the mean ± standard error of the mean from n=3

replicates. *In vitro* data was analyzed using a two-tailed Student's t-test with statistical

significance determined when p<.05. For the mouse data, the Mann-Whitney U test was used to

test for statistical significance. Values represent means ± SEM derived from at least 3

independent experiments. *, $P<0.05$; **, $P<0.005$; ***, $P<0.0005$; ****, $P<0.00005$; ns,

difference not significant.

Acknowledgements

This work was supported by National Institutes of Health grants R01-GM64671 and R35-

GM127331 to P.A.L. and R01-DK051406, R01-AI108749-01, and P50-DK0645400 grants to

A.L.F-M and S.J.H. C.S.W was supported in part by an Arnold O. Beckman Postdoctoral

Fellowship. J.I.R. and J.P.H. were supported by National Institutes of Health grant

RO1DK111930. The authors thank Dr. Robyn Klein and members of the Levin, Hultgren, and

Zaher groups for insightful suggestions and critical comments on the manuscript.

Author Contributions

CSW, PAL, ALFM, SJH, JIR and JPH designed the research studies. CSW performed the *in*

vitro assays. ALFM and AJLL performed the animal experiments and acquired data. JIR and

JPH performed triclosan detection and quantification experiments and acquired data. CSW

ALFM, JIR, and PAL analyzed data. CSW, ALFM, and JIR prepared the figures. CSW, PAL,

and ALFM wrote the manuscript. CSW, PAL, ALFM, SJH, JIR and JPH reviewed and edited the

manuscript.

390
391

References

1. **Food and Drug Administration, HHS**. 2016. Safety and Effectiveness of Consumer
Antiseptics; Topical Antimicrobial Drug Products for Over-the-Counter Human Use. Final
rule. Fed Regist **81**:61106–61130.

2. **McEwen SA**, **Fedorka-Cray PJ**. 2002. Antimicrobial use and resistance in animals. Clin
Infec Dis **34 Suppl 3**:S93–S106.

3. **Yazdankhah SP**, **Scheie AA**, **Høiby EA**, **Lunestad B-T**, **Heir E**, **Fotland TØ**,
Naterstad K, **Kruse H**. 2006. Triclosan and antimicrobial resistance in bacteria: an
overview. Microb Drug Resist **12**:83–90.

4. **McMurry LM**, **Oethinger M**, **Levy SB**. 1998. Triclosan targets lipid synthesis. Nature
394:531–532.

5. **Maiden MM**, **Hunt AMA**, **Zachos MP**, **Gibson JA**, **Hurwitz ME**, **Mulks MH**, **Waters**
CM. 2018. Triclosan Is an Aminoglycoside Adjuvant for Eradication of Pseudomonas
aeruginosa Biofilms. Antimicrobial Agents and Chemotherapy **62**:e00146–18.

6. **Drury B**, **Scott J**, **Rosi-Marshall EJ**, **Kelly JJ**. 2013. Triclosan exposure increases
triclosan resistance and influences taxonomic composition of benthic bacterial
communities. Environ Sci Technol **47**:8923–8930.

7. **Singer H**, **Müller S**, **Tixier C**, **Pillonel L**. 2002. Triclosan: occurrence and fate of a
widely used biocide in the aquatic environment: field measurements in wastewater
treatment plants, surface waters, and lake sediments. Environ Sci Technol **36**:4998–5004.

8. **Calafat AM**, **Ye X**, **Wong L-Y**, **Reidy JA**, **Needham LL**. 2008. Urinary concentrations
of triclosan in the U.S. population: 2003-2004. Environ Health Perspect **116**:303–307.

9. **MacIsaac JK**, **Gerona RR**, **Blanc PD**, **Apatira L**, **Friesen MW**, **Coppolino M**, **Janssen**
S. 2014. Health care worker exposures to the antibacterial agent triclosan. J Occup
Environ Med **56**:834–839.

10. **Balaban NQ**, **Merrin J**, **Chait R**, **Kowalik L**, **Leibler S**. 2004. Bacterial persistence as a
phenotypic switch. Science **305**:1622–1625.

11. **Lewis K**. 2010. Persister cells. Annu Rev Microbiol **64**:357–372.

12. **Mulcahy LR**, **Burns JL**, **Lory S**, **Lewis K**. 2010. Emergence of Pseudomonas aeruginosa
strains producing high levels of persister cells in patients with cystic fibrosis. J Bacteriol
192:6191–6199.

13. **Proctor RA**, **van Langevelde P**, **Kristjansson M**, **Maslow JN**, **Arbeit RD**. 1995.
Persistent and Relapsing Infections Associated with Small-Colony Variants of
Staphylococcus aureus. CLIN INFECT DIS **20**:95–102.

14.   **Høiby N**, **Bjarnsholt T**, **Givskov M**, **Molin S**, **Ciofu O**. 2010. Antibiotic resistance of
bacterial biofilms. Int J Antimicrob Agents **35**:322–332.

15.   **Seyfzadeh M**, **Keener J**, **Nomura M**. 1993. spoT-dependent accumulation of guanosine
tetraphosphate in response to fatty acid starvation in Escherichia coli. P Natl Acad Sci
USA **90**:11004–11008.

16.   **Harms A**, **Fino C**, **Sørensen MA**, **Semsey S**, **Gerdes K**. 2017. Prophages and Growth
Dynamics Confound Experimental Results with Antibiotic-Tolerant Persister Cells. MBio
8:e01964–17.

17.   **Korch SB**, **Henderson TA**, **Hill TM**. 2003. Characterization of the hipA7 allele of
Escherichia coli and evidence that high persistence is governed by (p)ppGpp synthesis.
Mol Microbiol **50**:1199–1213.

18.   **Viducic D**, **Ono T**, **Murakami K**, **Susilowati H**, **Kayama S**, **Hirota K**, **Miyake Y**. 2006.
Functional analysis of spoT, relA and dksA genes on quinolone tolerance in Pseudomonas
aeruginosa under nongrowing condition. Microbiol Immunol **50**:349–357.

19.   **Battesti A**, **Bouveret E**. 2006. Acyl carrier protein/SpoT interaction, the switch linking
SpoT-dependent stress response to fatty acid metabolism. Mol Microbiol **62**:1048–1063.

20.   **Muto A**, **Kimura A**, **Osawa S**. 1975. Effects of some antibiotics on the stringent control
of RNA synthesis in Escherichia coli. Mol Gen Genet **139**:321–327.

21.   **Brauner A**, **Fridman O**, **Gefen O**, **Balaban NQ**. 2016. Distinguishing between
resistance, tolerance and persistence to antibiotic treatment. Nat Rev Microbiol **14**:320–
330.

22.   **Harding GK**, **Ronald AR**. 1994. The management of urinary infections: what have we
learned in the past decade? Int J Antimicrob Agents **4**:83–88.

23.   **Flores-Mireles AL**, **Walker JN**, **Caparon M**, **Hultgren SJ**. 2015. Urinary tract
infections: epidemiology, mechanisms of infection and treatment options. Nat Rev
Microbiol **13**:269–284.

24.   **Provencher G**, **Bérubé R**, **Dumas P**, **Bienvenu J-F**, **Gaudreau E**, **Bélanger P**, **Ayotte
P**. 2014. Determination of bisphenol A, triclosan and their metabolites in human urine
using isotope-dilution liquid chromatography-tandem mass spectrometry. J Chromatogr A
1348:97–104.

25.   **Amato SM**, **Orman MA**, **Brynildsen MP**. 2013. Metabolic control of persister formation
in Escherichia coli. Mol Cell **50**:475–487.

26.   **Amato SM**, **Brynildsen MP**. 2015. Persister Heterogeneity Arising from a Single
Metabolic Stress. Curr Biol **25**:2090–2098.

27.   **Durfee T**, **Hansen A-M**, **Zhi H**, **Blattner FR**, **Jin DJ**. 2008. Transcription profiling of

461   the stringent response in Escherichia coli. J Bacteriol **190**:1084–1096.

462 28. **Schreiber G**, **Ron EZ**, **Glaser G**. 1995. ppGpp-mediated regulation of DNA replication
463   and cell division in Escherichia coli. Curr Microbiol **30**:27–32.

464 29. **Ocampo PS**, **Lázár V**, **Papp B**, **Arnoldini M**, **Abel zur Wiesch P**, **Busa-Fekete R**,
465   **Fekete G**, **Pál C**, **Ackermann M**, **Bonhoeffer S**. 2014. Antagonism between
466   bacteriostatic and bactericidal antibiotics is prevalent. Antimicrobial Agents and
467   Chemotherapy, 2nd ed. **58**:4573–4582.

468 30. **McNamara PJ**, **LaPara TM**, **Novak PJ**. 2014. The impacts of triclosan on anaerobic
469   community structures, function, and antimicrobial resistance. Environ Sci Technol
470   **48**:7393–7400.

471 31. **Lyman FL**, **Furia T**. 1969. Toxicology of 2, 4, 4"-trichloro-2-"hydroxy-diphenyl ether.
472   IMS Ind Med Surg **38**:64–71.

473 32. **Karatzas KAG**, **Webber MA**, **Jorgensen F**, **Woodward MJ**, **Piddock LJV**, **Humphrey**
474   **TJ**. 2007. Prolonged treatment of Salmonella enterica serovar Typhimurium with
475   commercial disinfectants selects for multiple antibiotic resistance, increased efflux and
476   reduced invasiveness. J Antimicrob Chemother **60**:947–955.

477 33. **Cavanagh J-AE**, **Trought K**, **Mitchell C**, **Northcott G**, **Tremblay LA**. 2018.
478   Assessment of endocrine disruption and oxidative potential of bisphenol-A, triclosan,
479   nonylphenol, diethylhexyl phthalate, galaxolide, and carbamazepine, common
480   contaminants of municipal biosolids. Toxicol In Vitro **48**:342–349.

481 34. **Louis GW**, **Hallinger DR**, **Braxton MJ**, **Kamel A**, **Stoker TE**. 2017. Effects of chronic
482   exposure to triclosan on reproductive and thyroid endpoints in the adult Wistar female rat.
483   J Toxicol Environ Health Part A **80**:236–249.

484 35. **Foxman B**, **Brown P**. 2003. Epidemiology of urinary tract infections: transmission and
485   risk factors, incidence, and costs. Infect Dis Clin North Am **17**:227–241.

486 36. **Mulvey MA**, **Schilling JD**, **Hultgren SJ**. 2001. Establishment of a persistent Escherichia
487   coli reservoir during the acute phase of a bladder infection. Infect Immun **69**:4572–4579.

488 37. **Hung C-S**, **Dodson KW**, **Hultgren SJ**. 2009. A murine model of urinary tract infection.
489   Nat Protoc **4**:1230–1243.

Figures

Figure 1: Triclosan induces tolerance to multiple antibiotics. *E. coli* (MG1655) were cultured

to OD600 = 0.2, split and cultured for an additional 30 minutes with (+T) or without 200 ng/ml

triclosan (-T). Indicated bactericidal antibiotics were then added and cells cultured for an

additional 2 to 20 hours prior to dilution plating. Each experiment was replicated three

independent times with only representative data shown.

Figure 2: Kinetic analysis of triclosan-induced persistence. *E. coli* (MG1655) and MRSA

(FPR3757) cells were cultured to OD-600 = 0.2, split and cultured for an additional 30 minutes

with (black line, closed squares) or without triclosan (grey line, open squares). At t=0, 100

ng/mL (A) or 1,000 ng/mL ciprofloxacin (B) was added to *E. coli* cultures and 50 ng/ml

vancomycin was added to MRSA cultures (C). Relative persistence in the presence of triclosan

(CFU+T/CFU-T) was calculated from the 4- and 20-hour time points (D). Values are the mean of

three independent biological replicates with error bars representing the standard error of the

mean. Asterisks represent significant difference between the triclosan treated and non-treated

using a Student's two-tailed t-test with * = $p < 0.05$.

Figure 3. ppGpp is needed for triclosan induced tolerance. Cell viability of MG1655 and

ppGpp0 *E. coli* with (+T) or without (-T) pretreatment with triclosan after challenge with

antibiotic (A). Growth curves of MG1655 (black curve) or ppGpp^0 (gray curve) in LB with

(dashed lines) or without (solid lines) triclosan (B). Cell viability of MG1655 and ppGpp0 *E. coli*

with (+T) or without (-T) pretreatment with spectinomycin after challenge with antibiotic (C).

Values are the mean of three independent biological replicates with error bars representing the

standard error of the mean. Asterisks represent significant difference between the triclosan

treated and non-treated using a Student's two-tailed t-test with * = $p < 0.05$ and ** = $p < .001$.

Figure 4: Triclosan reduces ciprofloxacin efficacy up to 100-fold in a mouse UTI model.

For each round of the experiment, 15 mice were given water containing triclosan (1,000

ng/mL) for 21 days, and 15 control mice received plain water. At 21 days, all mice were

infected with *E. coli* UTI89 ($\sim$5 x10^7 CFU). 24 hours post-infection, 10 mice from each

group were given 25 mg/kg of ciprofloxacin intraperitoneally. 48 hours post infection, urine

(A) was taken and mice were sacrificed and bladders (B) were harvested. To compare the

mice groups, Mann-Whitney U test was used, $p < 0.05$ was considered statistically significant:

*, $p < 0.05$; **, $p < 0.005$; ***, $p < 0.0005$; ns, values were not statistically different. The

horizontal bar represents the median value. The horizontal broken line represents the limit of

detection (LOD) of viable bacteria. Data are from 3 independent experiments. (C) Free

triclosan levels measured by LC-MS/MS of triclosan in untreated and treated mouse urine.

(45 mice were untreated and 45 mice were treated, these mice were divided in three

independent experiments. Urine from 3-4 mice was pooled for triclosan analysis). The

horizontal broken line represents the limit of detection (LOD) of triclosan.

Supplementary Data

Cipro (μg/mL):

Supplementary Data Figure 1. Triclosan protects at high concentrations of ciprofloxacin.

MG1655 cells were grown to OD-600 = 0.1 before triclosan was added for a final concentration

of 200 ng/mL for 30 minutes. Ciprofloxacin was added at the labeled concentration.

Ciprofloxacin was washed off, and cell viability was determined. Values are shown as averages

of three replicates with error bars showing the standard error of the mean.

Supplementary Data Figure 2. Triclosan is not bactericidal to the ppGpp0 cells. ppGpp0

cells were grown to OD-600 = 0.2 before triclosan was added for a final concentration of 200

ng/mL. Cells were plated at each time pointed and colony-forming units were quantified. Each

point represents the average of three biological replicates with the error bars representing the

standard error of the mean.

Ciprofloxacin 100 ng/mL

Supplementary Data Figure 3. Triclosan induces ciprofloxacin tolerance in the

uropathogentic *E. coli* UTI89. UTI89 cells were grown up to OD-600 = 0.2, split and cultured

for an additional 30 minutes with (+T) or without 200ng/ml triclosan (-T). Ciprofloxacin was

added to obtain a final concentration of 100 ng/mL. Cells were dot-plated at the 0- and 2-hour

time points. The plating efficiency was repeated three independent times with a representative

image shown.

Supplementary Data Figure 4. Measurement of free and metabolized triclosan in triclosan-

treated mouse urine. Chromatogram of a representative LC-MS/MS experiment showing the

elution profile of metabolite 1 (blue), triclosan (red) and metabolite 2 (green) (A). Mass spectra

of triclosan, metabolite 1, and metabolite 2, respectively, measured using precursor ion scans for

616   the chloride product ion (35 m/z) (B, C, and D). Triclosan peak area detected by LC-MS/MS

617   varies linearly with concentration down to 1.6 ng/mL (E).

Instrument Settings		
Ion Spray Voltage	-4.5 kV	
Heater Temperature	500 °C	
Nebulizer Gas	40	
Auxiliary Gas	40	
Declustering Potential	-10 V	
Collision Energy	-40 V	
Ions Detected (m/z)		
Triclosan	$286.8 \rightarrow 35$	$288.7 \rightarrow 35$
$^{13}C_{12}$-Triclosan	$298.84 \rightarrow 35$	$300.74 \rightarrow 35$
Metabolite 1	$382.8 \rightarrow 35$	$384.8 \rightarrow 35$
Metabolite 2	$366.7 \rightarrow 35$	$368.8 \rightarrow 35$

640 **Supplementary Data Table 1. MS/MS settings for triclosan detection**
641