1 Nutrient dynamics and stream order influence microbial community patterns along a 2914 km
2 transect of the Mississippi River
3
4

5 Michael W. Henson[1], Jordan Hanssen[2], Greg Spooner[2], Patrick Fleming[2], Markus Pukonen[2],
6 Frederick Stahr[3], and J. Cameron Thrash[1*]
7
8 [1] Department of Biological Sciences, Louisiana State University, Baton Rouge, LA 70803,
9 U.S.A.
10 [2] O.A.R. Northwest, Seattle, WA 98103, U.S.A.
11 [3] School of Oceanography, University of Washington, Seattle, WA 98195, U.S.A.
12
13
14
15 *Correspondence:
16
J. Cameron Thrash
Louisiana State University
Department of Biological Sciences
202 Life Sciences Bldg.
Baton Rouge, LA 70803
Phone: 225-578-8210
Fax: 225-578-2597
thrashc@lsu.edu

37 Running title: Microbial regime changes on the Mississippi River
38
39
40 Keywords: Mississippi River, WGCNA, microbial ecology, eutrophication

Abstract

Draining 31 states and roughly 3 million km^2, the Mississippi River (MSR) and its tributaries

constitute an essential resource to millions of people for clean drinking water, transportation,

agriculture, and industry. Since the turn of the 20[th] century, MSR water quality has continually

rated poorly due to human activity. Acting as first responders, microorganisms can mitigate,

exacerbate, and/or serve as predictors for water quality, yet we know little about their community

structure or ecology at the whole river scale for large rivers. We collected both biological (16S

and 18S rRNA gene amplicons) and physicochemical data from 38 MSR sites over nearly 3000

km from Minnesota to the Gulf of Mexico. Our results revealed a microbial community

composed of similar taxa to other rivers but with unique trends in the relative abundance patterns

among phyla, OTUs, and the core microbiome. Furthermore, we observed a separation in

microbial communities that mirrored the transition from an 8[th] to 10[th] Strahler order river at the

Missouri River confluence, marking a different start to the lower MSR than the historical

distinction at the Ohio River confluence in Cairo, IL. Within MSR microbial assemblages we

identified subgroups of OTUs from the phyla Acidobacteria, Bacteroidetes, Oomycetes, and

Heterokonts that were associated with, and predictive of, the important eutrophication nutrients

nitrate and phosphate. This study offers the most comprehensive view of MSR microbiota to

date, provides important groundwork for higher resolution microbial studies of river

perturbation, and identifies potential microbial indicators of river health related to

eutrophication.

Introduction

By connecting terrestrial, lotic, and marine systems, rivers perform vital roles in both the transport and processing of compounds in all major global biogeochemical cycles (Richey et al. 2002; Ensign and Doyle 2006; Withers and Jarvie 2008; Battin et al. 2009; Savio et al. 2015). Within the carbon cycle alone, rivers collectively discharge organic carbon to the oceans at over 0.4 Pg C yr^{-1} (Cauwet et al. 2002). Perhaps more importantly, rivers are generally net heterotrophic (Cole et al. 2007), indicating that they not only transport organic matter but host active metabolic processing of it as well. Conservative estimates place heterotrophic output of the world's fluvial networks (streams, rivers, and estuaries) at 0.32 Pg C yr^{-1} (Cole and Caraco 2001; Battin et al. 2009). Although rivers contain a small minority of global fresh water at any given moment, the considerable volumes that pass through these systems make them relevant to models attempting to quantify global elemental transformations. However, despite the fact that microbial functions likely play a vital role in ecosystem health for both rivers themselves and their places of discharge, microbial functions in rivers remain understudied.

At 3734 km, the Mississippi River (MSR) is the fourth longest on earth, draining 31 U.S. states and two Canadian provinces- a watershed consisting of 41% of the continental U.S (Turner and Rabalais 2003; Dagg et al. 2004). The MSR is a major source of drinking water for many U.S. cities; a critical thoroughfare for transportation, commerce, industry, agriculture, and recreation; and conveys the vestiges of human activity to the Gulf of Mexico (GOM). In New Orleans, the average flow rate is over 600,000 cubic feet s^{-1} (cfs) (Rabalais et al. 1996), but can exceed 3 million cfs during flood stages (Singh 2012), and carries over 150 x 10^9 kg of suspended sediment into the northern GOM annually (Dagg et al. 2004, 2005). This massive discharge includes excess nutrients (nitrogen and phosphorus), primarily from agricultural runoff

(McIsaac et al. 2001; Turner and Rabalais 2004; Schilling et al. 2010; Duan et al. 2014; Staley et

al. 2014a), that fuels one of the largest marine zones of seasonal coastal hypoxia in the world

(Rabalais et al. 2002, 2007; Bianchi et al. 2010; Bristow et al. 2015). Studying microbial

relationships to river eutrophication will improve our understanding of their contributions to

either mitigating or exacerbating nutrient input.

Far from a homogenous jumble of organisms ferried downriver, microbial community

composition changes with distance from the river mouth and/or from the influence of tributaries

(Lemke *et al.*, 2009; Winter *et al.*, 2007; Kolmakova *et al.*, 2014) (Read et al. 2015; Savio et al.

2015), resulting from altered nutrient concentrations (Staley et al. 2014b; Van Rossum et al.

2015; Meziti et al. 2016), differing dissolved organic matter (DOM) sources (Ruiz-González et

al. 2013; Blanchet et al. 2016), and land use changes (Staley et al. 2014a; Van Rossum et al.

2015; Zeglin 2015; Meziti et al. 2016). Past studies of the Thames, Danube, Yenisei, and

Columbia Rivers have found that planktonic river microbial assemblages were dominated by the

phyla Actinobacteria, Proteobacteria, and Bacteriodetes, and taxa such as acI Actinobacteria,

Polynucleobacter spp., GKS9 and LD28 *Betaproteobacteria*, CL500-29 Bacteriodetes, LD12

SAR11 *Alphaproteobacteria*, and *Novosphingobium* spp.(Zwart et al. 2002; Savio et al. 2015).

More recent 16S rRNA gene amplicon and metagenomic studies of the Minnesota portion of the

MSR corroborated previous research in other rivers that identified an increased proportion (Savio

et al. 2015) or richness (Read et al. 2015) of freshwater taxa with river distance, and an increased

abundance of "core" river microbes (Staley et al. 2013) with cumulative residence time (Savio *et*

al., 2015, Winter *et al.*, 2007; Read *et al.*, 2015; Staley *et al.*, 2013; Crump *et al.*, 1996).

Researchers have suggested that these patterns supported the application of the River

Continuum Concept (RCC) (Vannote *et al.*, 1980) to river microbiota. The RCC postulates that

as a river increases in size, the influences of riparian and other inputs will decrease as the river

establishes a dominant core community. Richness will increase with stream order complexity

before decreasing in higher order rivers (Vannote *et al.*, 1980). Therefore, as continuous systems

with increasing volumes and residence times, river microbiota should transition from

experiencing strong influences of mass effects from terrestrial and tributary sources to systems

where species sorting plays a more important role (Fortunato et al. 2012; Besemer et al. 2013;

Savio et al. 2015). Analysis of MSR microbial communities suggests the importance of point

source inputs on local communities while regional scale differences are more influenced by mass

immigration from sediment and tributaries (Staley et al. 2013, 2016; Payne et al. 2017).

Complicating matters, particle-associated communities in rivers (frequently defined as

those found on filters of > ~3 μm) remain distinct from their free-living counterparts (Jackson *et*

al., 2014; Crump *et al.*, 1999; Riemann and Winding, 2001), potentially due to increased

production rates from access to readily obtainable carbon (Crump *et al.*, 1999; Crump and

Baross, 1996). Typical taxa associated with particles include OTUs related to the Bacteroidetes

clades *Flavobacteria* and *Cytophaga*, Planctomycetes, *Rhizobium* spp., and *Methylophilaceae*

spp. (Crump and Baross 1996; Allgaier and Grossart 2006; D'Ambrosio et al. 2014; Jackson et

al. 2014). However, consistent trends in particle community composition are murky, with recent

evidence suggesting organisms may switch between free-living and particle-associated lifestyles

depending on substrate availability and chemical queues (Grossart 2010; D'Ambrosio et al.

2014). Thus, rivers constitute complex and highly dynamic ecosystems from a metacommunity

perspective.

Although important insights have been gained from recent research on portions of the

MSR (Staley et al. 2013, 2015, 2016; Payne et al. 2017), microbiological transects at the whole-

river or catchment scale have yet to be completed. This study aimed to *i)* compare size-

fractionated MSR microbial community populations to those in other rivers, *ii)* examine within-

river heterogeneity of microbial communities, and *iii)* identify MSR microorganisms most

strongly associated with eutrophication- all at a near whole-river scale. During the fall of 2014,

we completed the most extensive microbiological survey of the Mississippi River to date with a

continual rowed transect over two months (70 days). Rowers from the adventure education non-

profit OAR Northwest collected samples from Minneapolis, MN to the Gulf of Mexico (2918

km) (Fig. 1A). Our findings expand the current information available on microbial assemblages

in major lotic ecosystems; further delineate the relationships between microbial structure and

stream order, nutrients, and volume; and identify MSR taxa predictive of the eutrophication

nutrients nitrate and phosphate.

Materials and Methods

Sampling and Cell Counts

We used rowboats and a simple filtration protocol (Supplementary Information) to collect water

from 39 sites along a continually rowed transect of the MSR, starting in Lake Itasca and ending

in the GOM, over 70 days from September 18th to November 26th, 2014. Sites were chosen to be

near major cities and above and below large tributaries. After some samples were removed due

to insufficient sequence data, contamination, or incomplete metadata (see below), the final

usable set of samples included 38 sites starting at Minneapolis (Fig. 1A, Table S1). Most

sampling occurred within the body of the river, although due to safety issues, three samples were

collected from shore (Table S1). We collected duplicate samples at each site, but because

separate rowboat teams frequently collected these sometimes several dozen meters apart, they

cannot be considered true biological replicates and we have treated them as independent samples.

At each site, we filtered 120 mL of water sequentially through a 2.7 µm GF/D filter (Whatman

GE, New Jersey, USA) housed in a 25 mm polycarbonate holder (TISCH, Ohio, USA) followed

by a 0.2 µm Sterivex filter (EMD Millipore, Darmstadt, Germany) with a sterile 60 mL syringe

(BD, New Jersey, USA). We refer to fractions collected on the 2.7 µm and 0.22 µm filters as >

2.7 µm and 0.2-2.7 µm, respectively. Flow-through water from the first 60 mL was collected in

autoclaved acid-washed 60 mL polycarbonate bottles. Both filters were wrapped in parafilm, and

together with the filtrate, placed on ice in Yeti Roadie 20 coolers (Yeti, Austin, TX) until

shipment to LSU. Further, 9 mL of whole water for cell counts was added to sterile 15 mL

Falcon tubes containing 1 mL of formaldehyde and placed into the cooler. We monitored cooler

temperature with HOBO loggers (Onset, Bourne, MA) to ensure samples stayed at $\leq$ 4ºC. The

final cooler containing samples from sites P-Al had substantial ice-melt. Though our filters were

wrapped in parafilm, we processed melted cooler water alongside our other samples to control

for potential contamination in these filters. Given that some of our samples were expected to

contain low biomass, we also included duplicate process controls for kit contamination (Salter et

al. 2014; Weiss et al. 2014) with unused sterile filters. Flow-through 0.2 µm filtered water from

each collection was analyzed for SiO_4, PO_4^{3-}, NH_4^+, NO_3^-, and NO_2^- (µg/L) at the University of

Washington Marine Chemistry Laboratory

(http://www.ocean.washington.edu/story/Marine+Chemistry+Laboratory). Aboard-rowboat

measurements were taken for temperature and turbidity. We determined turbidity by deploying a

secchi disk (Wildco, Yulee, FL), while drifting with the current so the line hung vertically. It was

lowered until no longer visible, then raised until just visible, and measured for its distance below

the waterline. We then calculated secchi depth from the average of two measurements.

Temperature was measured with probes from US Water Systems (Indianapolis, IN), rinsed with

distilled water between samples. Samples for cell counts were filtered through a 2.7 μm GF/D

filter, stained with 1x Sybr Green (Lonza), and enumerated using the Guava EasyCyte

(Millipore) flow cytometer as previously described (Thrash et al. 2015).

DNA extraction and Sequencing

DNA was extracted from both filter fractions and controls using a MoBio PowerWater DNA kit

(MoBio Laboratories, Carlsbad, CA) following the manufacturer's protocol with one minor

modification: in a biosafety cabinet (The Baker Company, Stanford, ME), Sterivex filter

housings were cracked open using sterilized pliers and filters were then removed by cutting

along the edge of the plastic holder with a sterile razor blade before being placed into bead-

beating tubes. DNA was eluted with sterile MilliQ water, quantified using the Qubit2.0

Fluorometer (Life Technologies, Carlsbad, CA), and stored at -20° C. Bacterial and archaeal

sequences were amplified at the V4 region of the 16S rRNA gene using the 515f and 806r primer

set (Caporaso et al. 2012), and eukaryotic sequences from the V9 region of the 18S rRNA gene

using the 1391r and EukBR primer set (Amaral-Zettler et al. 2009). Amplicons were sequenced

on an Illumina MiSeq as paired-end 250 bp reads at Argonne National Laboratory. Sequencing

of the 16S and 18S rRNA gene amplicons resulted in 13,253,140 and 13,240,531 sequences,

respectively.

Sequence Analysis

We analyzed amplicon data with Mothur v.1.33.3 (Schloss et al. 2009) using the Silva v.119

database (Pruesse et al. 2007; Quast et al. 2013). Briefly, 16S and 18S rRNA gene sequences

were assembled into contigs and discarded if the contig had any ambiguous base pairs, possessed

repeats greater than 8 bp, or were greater than 253 or 184 bp in length, respectively. Contigs

were aligned using the Silva rRNA v.119 database, checked for chimeras using UCHIME (Edgar

et al. 2011), and classified also using the Silva rRNA v.119 database. Contigs classified as

chloroplast, eukaryotes, mitochondria, or "unknown;" or as chloroplast, bacteria, archaea,

mitochondria, or "unknown;" were removed from 16S or 18S rRNA gene data, respectively. The

remaining contigs were clustered using the cluster.split() command into operational taxonomic

units (OTUs) using a 0.03 dissimilarity threshold ($OTU_{0.03}$). After these steps, 146,725 and

131,352 OTUs remained for the 16S and 18S rRNA gene communities, respectively.

Sample quality control

To evaluate the potential for contamination from extraction kits, cooler water in the last set of

samples, or leaking/bursting of pre-filters, all samples were evaluated with hierarchical

clustering and NMDS analysis. Hierarchical clustering was preformed in R using the *hclust*

function with methods set to "average", from the vegan package (Oksanen et al. 2015). Samples

were removed from our analysis if they were observed to be outliers in both the NMDS and

hierarchical clustering such that they grouped with our process controls. The process and cooler

water controls were extreme outliers in both, as was sample L2 (Fig. S1, S2). Sterivex and

prefilter samples generally showed strong separation with the exception of three 16S rRNA gene

samples- STER X2, W2, S2 (Fig. S1, S2). The only other samples that were removed were due

to missing chemical data (Lake Itasca1-2, A1-2) or failed sequencing (16S STER Af1; 16S PRE

S2, X2; 18S PRE O1). Not including process or cooler water controls, 152 samples were

sequenced each for prokaryotic and eukaryotic communities. After these QC measures, 144 and

149 samples remained in the analyses from the 16S and 18S rRNA gene amplicons, respectively.

Further, to control for potential contaminants, any OTU with greater 20 reads in the process or

cooler controls was removed from the data set. 146,725 and 131,327 OTUs remained after these

steps for 16S and 18S rRNA gene communities, respectively.

Alpha and Beta Diversity

$OTU_{0.03}$ analyses were completed with the R statistical environment v.3.2.1(R Developement

Core Team 2015). Using the package PhyloSeq (McMurdie and Holmes 2013), alpha-diversity

was first calculated on the unfiltered OTUs using the "estimate richness" command within

PhyloSeq, which calculates Chao1 (McMurdie and Holmes 2013). After estimating chao1,

potentially erroneous rare OTUs, defined here as those without at least two sequences in 20% of

the data, were discarded. After this filter, the dataset contained 950 and 724 16S and 18S rRNA

gene OTUs, respectively. For site-specific community comparisons, OTU counts were

normalized using the package DESeq2 (Love et al. 2014) with a variance stabilizing

transformation (Learman et al. 2016). Beta-diversity between samples was examined using Bray-

Curtis distances via ordination with non-metric multidimensional scaling (NMDS). Analysis of

similarity (ANOSIM) was used to test for significant differences between groups of samples (e.g

lower versus upper MSR) using the *anosim* function in the vegan package (Oksanen et al. 2015).

The influence of environmental parameters on beta-diversity was calculated in R with the *envfit*

function.

Network analyses and modeling

To identify specific OTUs with strong relationships to environmental parameters (e.g. turbidity,

NO_3^-), we employed weighted gene co-expression network analysis (WGCNA) (Langfelder and

Horvath 2008) as previously described for OTU relative abundances (Guidi et al. 2016). First, a

similarity matrix of nodes (OTUs) was created based on pairwise Pearson correlations across

samples. This was transformed into an adjacency matrix by raising the similarity matrix to a soft

threshold power (p; p = 6 for 16S and 18S > 2.7 µm, p = 4 for 16S 0.2-2.7 µm, p = 9 for 18S 0.2-

2.7 µm) that ensured scale-free topology. Submodules of highly co-correlating OTUs were

defined with a topological overlap matrix and hierarchical clustering. Each submodule,

represented by an eigenvalue, was pairwise Pearson correlated to individual environmental

parameters (Figs. S7-10A). To explore the relationship of submodule structure to these

parameters, OTUs within the submodule were plotted using their individual correlation to the

parameter of interest (here nitrate or phosphate) and their submodule membership, defined as the

number of connections within the module (Figs. S7-10B, D). Strong correlations between

submodule structure and an environmental parameter facilitate identification of OTUs that are

highly correlated to that parameter. To evaluate the predictive relationship between a submodule

and a parameter, we employed partial least square regression (PLS) analysis. PLS maximizes the

covariance between two parameters (e.g., OTU abundance and nitrate concentration) to define

the degree to which the measured value (OTU abundance) can predict the response variable

(nutrient concentration). The PLS model was permutated 1000 times and Pearson correlations

were calculated between the response variable and leave-one-out cross-validation (LOOCV)

predicted values. Modeled values were then compared with measured values to determine the

explanatory power of the relationships (Figs. S7-10C, E). Relative contributions of individual

OTUs to the PLS regression were calculated using value of importance in the projection (VIP)

(Chong and Jun 2005) determination. PLS was run using the R package *pls* (Mevik and Wehrens

2007), while VIP was run using additional code found here:

http://mevik.net/work/software/VIP.R.

Environmental Ontology

Environmental ontology of individual 16S rRNA gene OTUs was determined using the SEQenv

(v1.2.4) pipeline (https://github.com/xapple/seqenv; Sinclair et al. 2016) as previously described

(Savio et al. 2015). Briefly, representative sequences of our OTUs were searched against the

NCBI nt database (updated on 07/01/2016) using BLAST and filtered for hits with a minimum of

99% identity. From each hit, a text query of the metadata was performed to identify terms

representing the sequence's environmental source. The text was mined for EnvO terms

(http://environmentontology.org/) and the frequency in which the terms appeared for each OTU

was recorded. Using the seq_to_names output provided by SEQenv, EnvO terms were formed

into six groups: Freshwater, Salt Water, Anthropogenic, Terrestrial, Sediment, and Unclassified

(Table S1). To be assigned a group, an OTU had to have the majority (> 50%) of its hits

classified to that term, while OTUs with equal distribution between two or more groups were

termed as Unclassified. OTUs that returned no significant hits to an EnvO term were assigned to

a seventh category, NA. OTUs and their corresponding relative abundances were merged based

on the assigned group and plotted with R.

Accession numbers

Community 16S and 18S rRNA gene sequence fastq files are available at the NCBI Sequence

Read Archive under the accession numbers: SRR3485674 - SRR3485971 and SRR3488881 -

SRR3489315, respectively.

Code Availability

All code used for Mothur, SeqENV, PhyloSeq, WGCNA, and PLS regression analyses can be

found on the Thrash lab website (http://thethrashlab.com/publications) with the reference to this

manuscript linked to "Supplemental Information".

<u>Results</u>

Using rowboats and a simple syringe-based filtration protocol, we measured 12 biological,

chemical, and physical parameters (e.g. 16S and 18S rRNA gene communities, NH_4^+, river

speed, etc.) from 38 sites along a 2918 km transect of the MSR (Fig. 1A). River order increases

dramatically at the Missouri confluence (eighth to tenth Strahler order (Pierson et al. 2008),

which corresponded to overall discharge (Fig. 1A) and beta diversity changes discussed below.

Therefore, we refer to this juncture as the separator between the upper (0 km – 1042 km, Sites A-

S) and lower MSR (1075-2914 km, Sites T-Al). Within the upper MSR, NO_3^-, PO_4^{3-}, and NO_2^-

were variable but generally increased downriver until peak concentrations near the confluences

of the Illinois and Missouri Rivers. This gave way to lower and more consistent concentrations

along the lower MSR (Fig. 1B). Turbidity (inversely related to secchi disk visibility) increased

steadily downriver to a maximum near the Illinois and Missouri River confluences (1042 km,

Site S) (Fig. 1B), then trended downwards for the rest of the transect. Planktonic (< 2.7 µm) cell

counts varied between 1 and 3×10^6 cells $\cdot$ mL^{-1} in the upper MSR, and decreased to high 10^5 cells

·mL^{-1} in the lower MSR (Fig. 1B). Water temperature ranged from 19°C (133km, Site E) to

11.7°C (2552 km, Site Ag), and river speed, excluding three sites sampled from shore, was

between 5.5 mph at Site Y and 0.4 mph (597 km, Site L) (Table S1). Spearman rank correlations

of the measured environmental parameters showed strong positive correlations between nitrate,

phosphate, distance, and increased turbidity; while nitrate and phosphate both strongly correlated

negatively to water temperature and river speed (Table S1).

Bacterial and archaeal communities

We observed a clear distinction between the 0.2-2.7 μm and > 2.7 μm 16S rRNA gene

communities (ANOSIM R = 0.65, P = 0.001) (Fig. S1A). Both size fractions had comparable

species richness and evenness that trended upwards downriver (Figs. S3A, B), although a peak

occurred for both fractions at sites O-Q (761-999 km) below the Des Moines River and above the

Illinois River (Fig. S3A, B). Both size fractions (NMDS stress = 0.14 for each) also showed a

significant separation between sites above and below the Missouri River confluence (ANOSIM,

> 2.7 μm: R = 0.44, P=0.001; 0.2-2.7 μm: R = 0.48, P = 0.001) (Fig. 2A, B), that the Eukaryotic

fractions mirrored (below). Phosphate and turbidity had the highest correlation with the

separation between the upper and lower > 2.7 μm communities (r = 0.57, r= 0.54, respectively),

with water temperature and distance (r > 0.40) also contributing (Fig. 2A). At an OTU level, taxa

related to the acI clade (Actinobacteria) and unclassified *Bacillaceae* (r > 0.77, P = 0.001)

contributed most to the separation between the upper and lower > 2.7 communities, with OTUs

related to the *Bacillales, Gemmatimonadaceae, Peptococcaceae*, and *Micromonosporaceae*

clades also a factor (r > 0.70, P = 0.001) (Figure 2A, Table S1). For the 0.2-2.7 μm fraction,

distance and nitrate were the strongest correlating environmental factors with the distinction

between upper and lower communities (r = 0.59 and r = 0.47, respectively), although phosphate,

turbidity, and water temperature (r > 0.40 for each) also contributed (Fig. 2B). OTUs related to

Flavobacterium and an unclassified Bacterium (closest NCBI BLAST hit *Acidovorax* sp.,

KM047473), most strongly contributed to the separation between the 0.2-2.7 μm communities (r

> 0.52, P = 0.001). Other important OTUs driving the separation between upper and lower

communities belonged to the clades Bacteroidetes, *Microbacteriaceae*, *Clostridiales*, and

Holophagaceae (r > 0.49, P = 0.001) (Figure 2B, Table S1).

At the phylum level, Proteobacteria, Actinobacteria, and Bacteroidetes dominated

bacterial communities in both fractions (Figs. 3A and B) (Table S1). Proteobacteria in the > 2.7

μm fraction fluctuated widely in abundance (Fig. 3A), whereas their 0.2-2.7 μm counterparts

generally increased in relative abundance downriver (Fig. 3B). 0.2-2.7 μm Bacteroidetes and

Actinobacteria generally decreased in the upper river and stabilized in the lower river. These

phyla showed considerable abundance variation in the > 2.7 μm fraction. Cyanobacteria in the >

2.7 μm fraction negatively correlated with increased turbidity (Spearman rank = 0.67), consistent

with lower irradiance. Both > 2.7 μm and 0.2-2.7 μm Acidobacteria increased in abundance

downriver and positively correlated with river distance (Wilcoxon single ranked test, P = < 0.01)

(Fig. 3A, B). Within the 0.2-2.7 μm fraction, the five most abundant OTUs were classified as a

LD12 (OTU11), two acI clade OTUs (OTU4, OTU7), a *Limnohabitans* sp. (OTU2), and a LD28

(OTU8) (Table S1). Comparatively, an unclassified *Methylophilaceae* (OTU1), a *Planktothrix*

sp. (OTU12), a NS11-12 marine group (OTU21), an *Aquirestis* sp. (OTU17), and an unclassified

Sphingobacteriales (OTU25) were the most abundant OTUs in the > 2.7 μm fraction (Table S1).

Archaea occurred at much lower relative abundances: we found only eight OTUs belonging to

the phyla Euryarchaeota and Thaumarchaeota, collectively. Thaumarchaeota increased in

abundance along the transect more in the > 2.7 µm fraction compared to the 0.2-2.7 µm (Fig. 3A,

B). In both fractions, we only detected Euryarcheota at very low abundances. Importantly, the

primers used in this study may miss some archaeal taxa (Parada et al. 2015).

Microbial eukaryotic communities

Eukaryotic communities, observed via the 18S rRNA gene, also showed a significant separation

between > 2.7 µm and 0.2-2.7 µm fractions (ANOSIM R = 0.689, P = 0.001) (Fig. S1B). As

expected due to generally larger cell sizes in microbial eukaryotes compared to prokaryotes,

species richness remained higher in the > 2.7 µm vs. 0.2-2.7 µm fractions (Fig. S3C, D).

Richness in the > 2.7 µm fraction gradually increased downriver, similarly to prokaryotic

communities, but remained relatively stable within the 0.2-2.7 µm fraction. Both the > 2.7 µm

(stress = 0.113) and 0.2-2.7 µm (stress = 0.146) fractions also showed a significant separation

between the upper and lower MSR (ANOSIM, > 2.7 µm: R = 0.696, P = 0.001; 0.2-2.7 µm: R =

0.576, P = 0.001) (Fig. 2C, D). Distance and phosphate were the top two environmental factors

influencing this distinction (r = 0.75, r = 0.48; r = 0.70, r = 0.54, respectively) (Fig. 2C, D; Table

S1). No other factors had correlations > 0.40 (Table S1). At the OTU level, taxa related to an

unclassified Ochrophyta (OTU63) and an unclassified Eukaryote (OTU1) separated the 0.2-2.7

µm communities (r > 0.63, P= 0.001), while the same unclassified Eukaryote OTU (OTU1) and

a second unclassified Eukaryote (OTU222) contributed most to separating the > 2.7 µm

communities (r > 0.80, P = 0.001) (Figure 2C and D, Table S1).

Stramenopiles (or Heterokonts), encompassing diatoms and many other forms of algae,

and OTUs that could not be classified at the phylum level dominated both the > 2.7 µm and 0.2-

2.7 µm communities (Fig. 4). Stramenopiles accounted for over 25% of both communities, with

higher abundances in the upper vs. lower river. We observed a similar trend of disparate

abundances between the upper and lower river for > 2.7 µm Cryptomonadales and 0.2-2.7 µm

Nucletmycea, the latter of which include fungi (Fig. 4A; Table S1). Within the 0.2-2.7 µm

fraction, we identified the five most abundant OTUs as three unclassified *Bacillariophytina*

(OTU7, OTU14, OTU9), a *Pythium* sp. (OTU170), and an unclassified *Cryptomonas* (OTU11)

(Table S1). Comparatively, two unclassified *Eukaryotes* (OTU2 and OTU1), an unclassified

Stramenopiles (OTU3), an unclassified *Perkinsidae* (OTU13), and an unclassified

Chrysophyceae (OTU6) had the highest abundance in the > 2.7 µm fraction (Table S1).

The Mississippi River Core Microbiome

We defined the core microbiome as those OTUs detectable after normalization in greater than

90% of the samples. The 16S rRNA gene > 2.7 µm and 0.2-2.7 µm core microbiomes consisted

of 82 and 98 OTUs, respectively, classified into eight different phyla- Proteobacteria,

Actinobacteria, Bacteroidetes, Cyanobacteria, Verrucomicrobia, Chloroflexi, Chlorobi,

Gemmatimonadetes- and composed of taxa such as freshwater SAR11 (LD12), *Limnohabitans*

sp., *Polynucleobacter* sp., acI clade, LD28 clade, and *Planktothrix* sp. (Table S1). Core

microbome relative abundance in both fractions decreased along the upper river but stabilized in

the lower river (Fig. 5A). We confirmed this effect by analyzing the upper and lower core

microbiomes separately. Although the total OTU numbers changed (81 and 116 OTUs in the

upper MSR and 160 and 144 OTUs in the lower MSR for the > 2.7 µm and 0.2-2.7 µm fractions,

respectively), the trends remained the same (Fig. S4).

Eighty OTUs comprised the > 2.7 µm 18S rRNA gene core microbiome (Fig. 5B). We

classified these as *Alveolata*, *Cryptophceae*, *Nucletmycea*, *Stramenopiles*, or unclassified

Eurkaryota (Table S1). Again, consistent with larger organism sizes, and thus fewer OTUs

overall, the 0.2-2.7 µm Eukaryotic core microbiome comprised only 21 OTUs (Fig. 5B). These

OTUs consisted of *Alveolata*, *Nucletmycea*, *Stramenopiles*, or unclassified Eurkaryota (Table

S1). While the 0.2-2.7 µm core microbiome remained relatively stable along the river, the > 2.7

µm core decreased along the upper MSR before stabilizing in the lower river, similarly to that of

the prokaryotes (Fig. 5B).

16S rRNA gene environmental ontology

We successfully classified 136 of the 950 16S rRNA gene OTUs with EnvO terminology (Table

S1). Freshwater organisms dominated the OTU designations that we could assign, although their

relative abundance in both the > 2.7 µm and 0.2-2.7 µm fractions decreased along the upper river

before stabilizing in the lower MSR (Fig. S5A, B). However, LD12 (OTU11), the most abundant

OTU in our dataset and a well-established freshwater organism (Zwart et al. 2002; Newton et al.

2011; Salcher et al. 2011), did not receive an EnvO classification, demonstrating the limitations

of this technique with current database annotations. Terrestrial organisms from the > 2.7 µm

fraction decreased along the transect (Fig. S5A), while those in the 0.2-2.7 µm fraction remained

stable (Fig. S5B). Although representing a minor fraction of total OTUs, sediment-associated >

2.7 µm microorganisms fluctuated widely in abundance along the upper river before stabilizing

in the lower, whereas those in the 0.2-2.7 µm fraction occurred in similar abundances throughout

the river. Taxa associated with anthropogenic sources in both fractions increased along the river

(Fig. S5A-B).

Network analyses identify taxa associated with and predictive of eutrophication

We applied Weighted Gene Correlation Network Analysis (WGCNA) to identify co-occurring

groups of OTUs (submodules) that also had significant associations with the eutrophication

nutrients phosphate and nitrate. Of the submodules identified through WGCNA as being most

strongly correlated to phosphate and nitrate, we restrict our discussion to those modeled via PLS

analysis to predict > 50% of the measured nutrient concentration. Our additional PLS analyses

for those submodules predicting < 50% of nitrate and/or phosphate concentrations are included

in figure S6.

Three submodules in the prokaryotic and eukaryotic fractions were strongly associated

with phosphate. The 0.2-2.7 μm prokaryotic submodule most associated with phosphate was

composed of 51 OTUs (Table 1), and had moderate correlation between the submodule structure

and phosphate (Fig. S7D). Strong correlations between submodule structure and the measured

nutrient suggests that individual submodule OTUs that also have strong correlations to the

nutrient are the most important organisms associated with that nutrient (Langfelder and Horvath

2008). PLS modeling determined that this prokaryotic submodule predicted 80% of measured

phosphate concentrations (Table 1). Variable importance in the projection (VIP) analysis found

OTUs corresponding to an unclassified *Holophagaceae,* an unclassified *Gemmatimonadaceae*,

and an unclassified *Burkholderiaceae* were the three most important in the PLS model for

phosphate (Fig. 6A, Table S1). OTU322 (Acidobacteria subgroup 6), had moderate correlation to

phosphate but high node centrality (n = 51; Fig. 6A, Table S1), corroborating evidence that

freshwater sediment Acidobacteria subgroup 6 often occur in co-culture with

Alphaproteobacteria and may be metabolically connected (Kielak et al. 2016 and refs. within). A

submodule from the 0.2-2.7 μm eukaryotic fraction was also highly predictive of phosphate

(Table 1, Fig. S8D). We identified the top four VIP taxa as an unclassified *Peronosporomycetes*,

an unclassified *Ochrophyta*, an unclassified Eukaryote, and an unclassified *Stramenopiles* (Fig.

6F, Table S1). All four OTUs had Pearson correlations with phosphate greater than 0.60, among

which two were negative (Fig. 7A, Table S1). Unfortunately, the most highly interconnected

OTU within the submodule remained unclassified even at the Phylum level (Fig. 7A Table S1).

A > 2.7 μm eukaryotic submodule could predict 62% of measured phosphate variance (Table 1).

An unclassified Eukaryote and an unclassified *Peronosporomycetes* occupied top two VIP

positions (Fig. 7B, Table S1).

Only two submodules had strong associations with nitrate. The > 2.7 μm prokaryotic

submodule most strongly correlated with nitrate contained 133 OTUs (Table 1, Fig. S9A).

However, despite a low structural correlation to nitrate, the submodule strongly predicted

measured nitrate in the PLS model (Figure S9B, Table 1). The four highest VIP scoring OTUs

were an *Anabaena* sp., a *Flavobacterium* sp., an unclassified bacterium, and a member of the

Sphingobacteriales NS11-12 marine group (Fig. 6B, Table S1). OTUs with the highest node

centrality (> 20) belonged to the *Sphingomonadales* (*Alphaproteobacteria*) and

Sphingobacteriales (Bacteroidetes), and each positively correlated with nitrate (r > 0.48, Fig. 6B,

Table S1). The > 2.7 μm eukaryotic submodule most associated with nitrate predicted 57% of

observed variation in nitrate concentrations (Table 1). OTUs with the top VIP scores were two

unclassified *Chrysophyceae*, an unclassified *Ochrophyta*, and an unclassified Diatom (Fig. 7C,

S10; Table S1).

Discussion

Understanding microbial communities on a basin scale is essential for unveiling the complex

linkage between organisms, metabolism, and water quality. Our near whole river 2914 km

transect revealed that while the MSR comprised similar aquatic microbial taxa found within

other rivers (Ghai et al. 2011; Fortunato et al. 2013; Jackson et al. 2014; Read et al. 2015; Savio

et al. 2015; Meziti et al. 2016), the river hosted unique relative abundance and core microbiome

trends. Physio-chemical measurements demonstrated continual increases in nutrients and

sediment along the upper river, whereas in the lower river, nitrate, phosphate, nitrite, and

turbidity stayed more consistent (Fig. 1B). These distinct geographic patterns mirrored the

observed separation of two microbial regimes at the Missouri River confluence (Fig. 2). This

differed from the historical distinction of the upper and lower MSR at the Ohio River confluence

in Cairo, IL, but matched the changes in Strahler order from eight to ten (Pierson et al. 2008). In

general, both prokaryotic fractions and the > 2.7 μm fraction of eukaryotic communities

increased in richness downriver (Fig. S3). The relative abundance of core community taxa

decreased in the upper MSR before settling in the lower river (Fig. 5). Co-occurrence network

analyses identified taxa associated with, and predictive of, the eutrophication nutrients nitrate

and phosphate that provide important targets for future study and may assist in detecting and

quantifying imminent changes in river water quality.

Some qualities of the MSR matched other rivers. MSR microbial communities separated

into two distinct fractions of microbial assemblages, 0.2-2.7 μm and > 2.7 μm (ANOSIM, R =

0.65, P = 0.001), throughout the river transect (DeLong et al. 1993; Crump and Baross 1996;

Allgaier and Grossart 2006; D'Ambrosio et al. 2014; Jackson et al. 2014). These fractions were

composed of OTUs classified as LD12 (freshwater SAR11), *Limnohabitans* sp.

(Betaproteobacteria), LD28 (Betaproteobacteria), acI clade (Actinobacteria), and *Algoriphagu*s

sp. (Bacteriodetes). These taxa also dominated in other rivers (Ghai et al. 2011; Fortunato et al.

2013; Kolmakova et al. 2014; Read et al. 2015; Savio et al. 2015; Meziti et al. 2016).

However, despite the MSR generally sharing similar taxa with other rivers, abundance

patterns indicated a unique system. Phylum level abundances contrasted a previous study limited

to the Minnesota portion of the MSR (Staley et al. 2013), as well as studies of the Thames (Read

et al. 2015), Danube (Savio et al. 2015), and Columbia Rivers (Crump et al. 1999). Specifically,

within the 0.2-2.7µm MSR prokaryotic fraction, Proteobacteria remained the most abundant

phylum throughout the river, while Bacteroidetes and Actinobacteria decreased downriver (Fig.

3). In contrast, the other river studies identified Bacteroidetes-dominated headwaters that gave

way to more abundant Actinobacteria and Proteobacteria further downriver (Crump et al. 1999;

Staley et al. 2014a; Read et al. 2015; Savio et al. 2015). Although comparison beyond phylum

level is important for understanding community changes, most studies report only OTU

taxonomy making comparisons of OTU relative abundances difficult. OTUs belonging to the acI

and LD12 freshwater clades continually increased in abundance towards the river mouth in the

Thames and Danube Rivers (Read et al. 2015; Savio et al. 2015), whereas MSR OTUs classified

as acI (OTUs 3, 4 and 7) and LD12 (OTU 11) did not change in abundance throughout the

transect. Further, MSR OTU richness generally increased with distance, and core community

abundance decreased along the upper river in both fractions, contrasting predictions by the RCC

(Vannote *et al.*, 1980) and observations in previous studies from other rivers (Read et al. 2015;

Savio et al. 2015) where these trends were reversed. Importantly, we did not sample the true

headwaters of the MSR (Lake Itasca to above St. Cloud), and at the point of first sampling, the

MSR already constituted an eighth order river. Therefore, some of the trends predicted by the

RCC may have been missed. Ultimately, the variant observations between this MSR survey and

other river studies may result from different sampling methodologies or the particular timing of

sampling. However, these differences may also stem from biological signal related to unique

environmental conditions, human impacts, and changes in the level of river engineering with

distance.

Within the MSR, we observed distinct prokaryotic and eukaryotic microbial community

regimes separated at the Missouri River confluence. This separation matched physio-chemical

changes observed for the MSR such as increased Strahler river order (8 to 10) and more

consistent nutrient concentrations (Fig. 1). We hypothesize that the separation between the upper

and lower MSR communities was driven by changes in the importance of immigration,

emigration, and resource gradient dynamics. Specifically, our data suggests that mass effects

play a role in structuring microbial communities in the upper MSR, although instead of only in

the headwaters, this process continues for almost a third of the length of the river. Increased

turbidity correlated with decreases in freshwater associated bacteria (Spearman rank correlation,

> 2.7 µm r $= 0.55$; 0.2-2.7 µm r $= 0.51$) and the core microbiome relative abundance (Spearman

rank correlation, > 2.7 µm r $= 0.53$; 0.2-2.7 µm r $= 0.63$) in the upper MSR. Similarly, nutrients

like phosphate and nitrate continually increased along the upper river (Fig. 1). In contrast, these

nutrients, turbidity, and core microbiome relative abundance all statbilized in the lower MSR.

These patterns are consistent with communities under the influence of mass effects from

tributaries in the upper MSR. However, once the MSR grew to a tenth order river, the large

volume and size potentially buffered it from allochthonous influences, allowing species sorting

effects to have greater influence over mass immigration. The lower river represents a more stable

environment (e.g. nutrient concentrations) with its increased size and volume, contrasting the

more variable upper MSR. Though river speed increased, the effective residence time also

increased since taxa no longer experience rapidly changing environmental variables.

That we still observed increasing richness and variation between microbial communities

along the lower MSR might indicate the increased influence of environmental filtering

attributable to unmeasured bottom-up factors, such as the quality and quantity of DOM, or top-

down influences such as predation or viral lysis. An overall community shift from a mixture of

allochthonous members to a "native" population requires growth rates greater than residence

time over a given distance (Crump et al. 2004). The extent to which a local or regional event

impacts downriver populations would be dependent on the success of allochthonous taxa

associated with those events to establish within the autochthonous population. The lower river

provides ample opportunities for microbial community differentiation based on average

prokaryotic growth rates (Savio et al. 2015), especially among particle-associated (> 2.7 µm)

taxa (Crump et al. 1999). Thus, although the aggregate patterns in particular phyla and

taxonomic richness within the MSR differ from other systems, similar ecological processes may

still occur, but the relative proportion of the river whereby mass effects vs. species sorting

dominate fosters unique community dynamics.

Microorganisms most associated with eutrophication nutrients are important target taxa

for future study of nitrate and phosphate uptake and/or metabolizism, and also represent

plausible biological indicators of river trophic state. By focusing on OTUs with the highest VIP

scores in the PLS models, we identified several bacterial and eukaryotic taxa that fit these criteria

(Figs. 6, 7). Organisms in the *Sphingomonadaceae* (e.g., *Novosphingobium* spp.) contributed

strongly to the PLS models predicting nitrate with both > 2.7 µm (Fig. 6B) and 0.2-2.7 µm (Fig.

S6A) size fractions (Table S1). *Novosphingobium* spp. isolates have previously been associated

with eutrophic environments (Trusova and Gladyshev 2002; Zwart et al. 2002; Addison et al.

2007; Li et al. 2012) and some can reduce nitrate (Addison et al. 2007; Li et al. 2012), making

these specific OTUs candidates for nitrate metabolism in the river water column. An *Anabaena*

sp. (OTU40) from the core microbiome had the top VIP score within the > 2.7 μm submodule

that could predict 69% of nitrate concentrations, and correlated negatively with nitrate (Table 1,

Fig. 6B). The nitrogen-fixing *Anabaena* (Allen and Arnon 1955) typically bloom in low

dissolved inorganic nitrogen (DIN) conditions, making the absence of these consistent with high

DIN. An unclassified *Holophagaceae* OTU (OTU33) had the highest VIP score within the

prokaryotic 0.2-2.7 μm submodule that predicted 80% of the variance in phosphate

concentrations (Table 1, Fig. 6A). This same OTU was also a key driver of the beta diversity

separation between upper and lower river communities (Fig. 2B), and a member of the core

microbiome (Table S1). The *Holophagaceae* belong to the Acidobacteria phylum, and the Ohio

River has been reported to contain a much higher abundance of Acidobacteria relative to other

tributaries (Jackson et al. 2014). Notably, Acidobacteria increased with river distance in our

study as well (Fig. 3) with a peak in the 0.2-2.7 μm fraction around the Arkansas river. This

increase in "free-living" Acidobacteria downriver is unique among other whole river studies,

making these organisms, and the *Holophagaceae* OTUs in particular, potentially important

organisms for the MSR river basin specifically.

Within Eukaryotes, multiple different algae, diatom, and Oomycetes OTUs occupied

submodules highly predictive of nitrate and phosphate (Fig. 6A-C, Table S1), and specifically

Chrysophyceae taxa from both size fractions correlated strongly with nitrate (Fig. 7C, Fig. S6E).

Chrysophyceae (golden algae) commonly occupy river systems (Necchi Jr 2016) including the

MSR (Korajkic et al. 2015), can be autotrophic and mixotrophic (Jansson et al. 1996), and may

serve as predators of prokaryotes (Caron et al. 1990). Further, multiple OTUs classified as

Peronosporomycetes were important in predicting phosphate in both size fractions (Fig 7A and

B, Table S1). *Peronosporomycetes* are fungus-like eukaryotic organisms known to be pathogenic

in fish, plants, and mammals (Dick 2003; Islam and von Tiedemann 2011). While we also

identified many other eukaryotic OTUs as important predictors of nutrients, poor taxonomic

resolution hindered our ability to discuss them further. Improved cultivation and systematics of

key microbial eukaryotes will be vital to understanding river nutrient dynamics.

While the most geographically comprehensive analysis to date for the MSR, this study

only encompasses a snapshot in time. Seasonal changes that have been observed in the

Minnesota portion of the upper MSR and the Columbia River (Smith et al. 2010; Fortunato et al.

2013; Staley et al. 2015) undoubtedly influence this dynamic system. Future studies should

incorporate microbial responses, at a whole-river scale, to seasonal pulse events (e.g. rain, snow

melt) and how river size and volume may buffer local microbial communities from

allochthonous inputs (Zeglin 2015). Our current research highlights the distinctiveness of MSR

microbial communities and the complexities influencing their structure within the MSR

ecosystem. While similar to other riverine studies in some respects, the 2014 MSR microbes

differed in alpha diversity and core community membership compared to the other rivers. The

observed association between changes in Strahler's river order, nutrient dynamics, and

community composition indicates the importance of hydrology (Freimann et al. 2015; Zeglin

2015; Niño-García et al. 2016) on the spatial dynamics structuring microbial communities and

provides baseline information for future MSR studies that incorporate greater temporal and

spatial resolution. With water quality and river health of growing local and global importance

(Vorosmarty et al. 2010; Russell and Weller 2013), the determination of candidate taxa

predictive of eutrophic nutrients provides impetus for targeted research on their larger potential

in predicting, and potentially ameliorating, river quality.

Acknowledgements

This work was supported by the Department of Biological Sciences, College of Science, and the

Office of Research and Economic Development at Louisiana State University; and the College of

the Environment at University of Washington. Portions of this research were conducted with

high performance computing resources provided by Louisiana State University

(http://www.hpc.lsu.edu). The authors thank Dr. Gary King and Dr. Caroline Fortunato for

thoughtful comments on the manuscript. The authors also thank the countless volunteers,

schools, and organizations that facilitated the research. We specifically thank Pete Weess, Jessica

Zimmerman, Katy Welch, Brian Moffitt, and David Cheney for helping organize the shipment of

coolers between sites, and the OAR Northwest sponsors- Seattle Yacht Club Foundation, Yeti

Coolers, and the National Mississippi River Museum and Aquarium. A full list of sponsors and

volunteers can be found on the OAR Northwest website (oarnorthwest.org). We would also like

to thank Dr. Matthew Sullivan and Dr. Simon Roux for their help with coding the WGCNA and

sPLS analyses. Lastly, we would like to thank Mrs. Ginger Thrash, who connected the Thrash

lab to OAR Northwest.

Conflict of Interest

The authors declare no competing financial interests.

References

Addison, S. L., S. M. Foote, N. M. Reid, and G. Lloyd-Jones. 2007. Novosphingobium

nitrogenifigens sp. nov., a polyhydroxyalkanoate-accumulating diazotroph isolated from a

New Zealand pulp and paper wastewater. Int. J. Syst. Evol. Microbiol. **57**: 2467–2471.

Allen, M. B., and D. I. Arnon. 1955. Studies on nitrogen-fixing blue-green algae. I. Growth and

nitrogen fixation by Anabaena cylindrica Lemm. Plant Physiol. **30**: 366.

Allgaier, M., and H.-P. Grossart. 2006. Seasonal dynamics and phylogenetic diversity of free-

living and particle-associated bacterial communities in four lakes in northeastern Germany.

Aquat. Microb. Ecol. **45**: 115–128.

Amaral-Zettler, L. A., E. A. McCliment, H. W. Ducklow, and S. M. Huse. 2009. A Method for

Studying Protistan Diversity Using Massively Parallel Sequencing of V9 Hypervariable

Regions of Small-Subunit Ribosomal RNA Genes. PLoS One **4**: e6372.

Battin, T. J., L. a. Kaplan, S. Findlay, C. S. Hopkinson, E. Marti, A. I. Packman, J. D. Newbold,

and F. Sabater. 2009. Biophysical controls on organic carbon fluxes in fluvial networks.

Nat. Geosci. **2**: 595–595.

Besemer, K., G. Singer, C. Quince, E. Bertuzzo, W. Sloan, and T. J. Battin. 2013. Headwaters

are critical reservoirs of microbial diversity for fluvial networks. Proc. Biol. Sci. **280**:

20131760.

Bianchi, T. S., S. F. DiMarco, J. H. Cowan, R. D. Hetland, P. Chapman, J. W. Day, and M. A.

Allison. 2010. The science of hypoxia in the northern Gulf of Mexico: A review. Sci. Total

Environ. **408**: 1471–1484.

Blanchet, M., O. Pringault, C. Panagiotopoulos, D. Lefèvre, B. Charrière, J.-F. Ghiglione, C.

Fernandez, F. L. Aparicio, C. Marrasé, P. Catala, L. Oriol, J. Caparros, and F. Joux. 2016.

When riverine dissolved organic matter (DOM) meets labile DOM in coastal waters:

changes in bacterial community activity and composition. Aquat. Sci. , doi:10.1007/s00027-

016-0477-0

Bristow, L. A., N. Sarode, J. Cartee, A. Caro-Quintero, B. Thamdrup, and F. J. Stewart. 2015.

Biogeochemical and metagenomic analysis of nitrite accumulation in the Gulf of Mexico

hypoxic zone. Limnol. Oceanogr. **60**: 1733–1750.

Caporaso, J. G., C. L. Lauber, W. a Walters, D. Berg-Lyons, J. Huntley, N. Fierer, S. M. Owens,

J. Betley, L. Fraser, M. Bauer, N. Gormley, J. a Gilbert, G. Smith, and R. Knight. 2012.

Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq

platforms. ISME J. **6**: 1621–1624.

Caron, D. A., K. G. Porter, and R. W. Sanders. 1990. Carbon, nitrogen, and phosphorus budgets

for the mixotrophic phytoflagellate Poterioochromonas malhamensis (Chrysophyceae)

during bacterial ingestion. Limnol. Oceanogr. **35**: 433–443.

Cauwet, G., D. A. Hansell, and C. A. Carlson. 2002. DOM in the coastal zone, p. 579–609. *In*

Biogeochemistry of marine dissolved organic matter. Academic Press, San Diego, CA.

Chong, I. G., and C. H. Jun. 2005. Performance of some variable selection methods when

multicollinearity is present. Chemom. Intell. Lab. Syst. **78**: 103–112.

Christian Winter Gerhard Kavka, Robert L. Mach, and, T. H., and A. H. Farnleitner. 2007.

Longitudinal Changes in the Bacterial Community Composition of the Danube River: a

Whole-River Approach. AEM

Cole, J. J., and N. F. Caraco. 2001. Carbon in catchments: Connecting terrestrial carbon losses

with aquatic metabolism. *Marine and Freshwater Research*. 101–110.

Cole, J. J., Y. T. Prairie, N. F. Caraco, W. H. McDowell, L. J. Tranvik, R. G. Striegl, C. . Duarte,

P. Kortelainen, J. a. Downing, J. J. Middelburg, and J. M. Melack. 2007. Plumbing the

global carbon cycle: Integrating inland waters into the terrestrial carbon budget. Ecosystems

10: 171–184.

Crump, B. C., E. V. Armbrust, and J. A. Baross. 1999. Phylogenetic Analysis of Particle-

Attached and Free-Living Bacterial Communities in the Columbia River, Its Estuary, and

the Adjacent Coastal Ocean. Appl. Environ. Microbiol. **65**: 3192–3204.

Crump, B. C., and J. A. Baross. 1996. Particle-attached bacteria and heterotrophic plankton

associated with the Columbia River estuarine turbidity maxima. Part. Bact. heterotrophic

Plankt. Assoc. with Columbia River Estuar. Turbid. maxima

Crump, B. C., C. S. Hopkinson, M. L. Sogin, and J. E. Hobbie. 2004. Microbial Biogeography

along an Estuarine Salinity Gradient: Combined Influences of Bacterial Growth and

Residence Time. Appl. Environ. Microbiol. **70**: 1494–1505.

D'Ambrosio, L., K. Ziervogel, B. Macgregor, A. Teske, and C. Arnosti. 2014. Composition and

enzymatic function of particle-associated and free-living bacteria: a coastal/offshore

comparison. ISME J. 1–13.

Dagg, M., R. Benner, S. Lohrenz, and D. Lawrence. 2004. Transformation of dissolved and

particulate materials on continental shelves influenced by large rivers: Plume processes.

Cont. Shelf Res. **24**: 833–858.

Dagg, M. J., T. S. Bianchi, G. A. Breed, W.-J. Cai, S. Duan, H. Liu, B. A. McKee, R. T. Powell,

and C. M. Stewart. 2005. Biogeochemical characteristics of the lower Mississippi River,

USA, during June 2003. Estuaries **28**: 664–674.

DeLong, E. F., D. G. Franks, and A. Alldredge. 1993. Phylogenetic diversity of aggregate-

attached vs. free-living marine bacteria assemblages. Limnol. Ocean. **38**: 924–934.

Dick, M. . 2003. Straminipilous Fungi: Systematics of the Peronosporomycetes Including

Accounts of the Marine Straminipilous Protists, the Plasmodiophorids and Similar

704  Organisms. Mycopathologia **156**: 385–386.

705 Duan, S., R. T. Powell, and T. S. Bianchi. 2014. High frequency measurement of nitrate

706  concentration in the Lower Mississippi River, USA. J. Hydrol. **519**: 376–386.

707 Edgar, R. C., B. J. Haas, J. C. Clemente, C. Quince, and R. Knight. 2011. UCHIME improves

708  sensitivity and speed of chimera detection. Bioinformatics **27**: 2194–2200.

709 Ensign, S. H., and M. W. Doyle. 2006. Nutrient spiraling in streams and river networks. J.

710  Geophys. Res. Biogeosciences **111**, doi:10.1029/2005JG000114

711 Fortunato, C. S., A. Eiler, L. Herfort, J. A. Needoba, T. D. Peterson, B. C. Crump, F. CS, A.

712  Eiler, L. Herfort, N. JA, P. TD, and C. BC. 2013. Determining indicator taxa across spatial

713  and seasonal gradients in the Columbia River coastal margin. ISME J **7**: 1899–1911.

714 Fortunato, C. S., L. Herfort, P. Zuber, A. M. Baptista, and B. C. Crump. 2012. Spatial variability

715  overwhelms seasonal patterns in bacterioplankton communities across a river to ocean

716  gradient. ISME J **6**: 554–563.

717 Freimann, R., H. Bürgmann, S. E. G. Findlay, and C. T. Robinson. 2015. Hydrologic linkages

718  drive spatial structuring of bacterial assemblages and functioning in alpine floodplains TL -

719  6. Front. Microbiol. **6 VN-re**, doi:10.3389/fmicb.2015.01221

720 Van Geel, B., L. R. Mur, M. Ralska-Jasiewiczowa, and T. Goslar. 1994. Fossil akinetes of

721  Aphanizomenon and Anabaena as indicators for medieval phosphate-eutrophication of Lake

722  Gosciaz (Central Poland). Rev. Palaeobot. Palynol. **83**: 97–105.

723 Ghai, R., F. Rodŕíguez-Valera, K. D. McMahon, D. Toyama, R. Rinke, T. C. S. de Oliveira, J.

724  W. Garcia, F. P. de Miranda, and F. Henrique-Silva. 2011. Metagenomics of the water

725  column in the pristine upper course of the Amazon river. PLoS One **6**,

726  doi:10.1371/journal.pone.0023785

Grossart, H. P. 2010. Ecological consequences of bacterioplankton lifestyles: Changes in

concepts are needed. Environ. Microbiol. Rep. **2**: 706–714.

Guidi, L., S. Chaffron, L. Bittner, D. Eveillard, A. Larhlimi, S. Roux, Y. Darzi, S. Audic, L.

Berline, and J. R. Brum. 2016. Plankton networks driving carbon export in the oligotrophic

ocean. Nature

Islam, M. T., and A. von Tiedemann. 2011. 2,4-Diacetylphloroglucinol suppresses

zoosporogenesis and impairs motility of Peronosporomycete zoospores. World J. Microbiol.

Biotechnol. **27**: 2071–2079.

Jackson, C. R., J. J. Millar, J. T. Payne, and C. a Ochs. 2014. Free-living and particle-associated

bacterioplankton in large rivers of the Mississippi River Basin demonstrate biogeographic

patterns. Appl. Environ. Microbiol. **80**: 7186–7195.

Jansson, M., P. Blomqvist, A. Jonsson, and A. BergstrÖm. 1996. Nutrient limitation of

bacterioplankton, autotrophic and mixotrophic phytoplankton, and heterotrophic

nanoflagellates in Lake Örträsket. Limnol. Oceanogr. **41**: 1552–1559.

Kielak, A. M., C. C. Barreto, G. A. Kowalchuk, J. A. van Veen, and E. E. Kuramae. 2016. The

Ecology of Acidobacteria: Moving beyond Genes and Genomes. Front. Microbiol. **7**: 744.

Kolmakova, O. V., M. I. Gladyshev, A. S. Rozanov, S. E. Peltek, and M. Y. Trusova. 2014.

Spatial biodiversity of bacteria along the largest Arctic river determined by next-generation

sequencing. FEMS Microbiol. Ecol. **89**: 442–450.

Korajkic, A., L. W. Parfrey, B. R. McMinn, Y. V. Baeza, W. VanTeuren, R. Knight, and O. C.

Shanks. 2015. Changes in bacterial and eukaryotic communities during sewage

decomposition in Mississippi river water TL - 69. Water Res. **69 VN-r**: 30–39.

Langfelder, P., and S. Horvath. 2008. WGCNA: an R package for weighted gene co-expression

network analysis. BMC Bioinformatics **9**: 559.

Learman, D. R., M. W. Henson, J. C. Thrash, B. Temperton, P. M. Brannock, S. R. Santos, A. R.

Mahon, and K. M. Halanych. 2016. Biogeochemical and microbial variation across 5500

km of Antarctic surface sediment implicates organic matter as a driver of benthic

community structure. Front. Microbiol. **7**, doi:10.3389/fmicb.2016.00284

Lemke, M. J., E. K. Lienau, J. Rothe, T. A. Pagioro, J. Rosenfeld, and R. Desalle. 2009.

Description of freshwater bacterial assemblages from the upper Paran?? river floodpulse

system, Brazil. Microb. Ecol. **57**: 94–103.

Li, H.-F., Z. T. Zou, B. Z. Li, X. Wang, J. S. Yang, and H. L. Yuan. 2012. Novosphingobium

sediminis sp. nov., isolated from the sediment of a eutrophic lake. J. Gen. Appl. Microbiol.

58: 357–362.

Love, M. I., S. Anders, and W. Huber. 2014. Differential analysis of count data - the DESeq2

package. Genome Biol. **15**: 550.

McIsaac, G. F., M. B. David, G. Z. Gertner, and D. a Goolsby. 2001. Nitrate flux in the

Mississippi River. Nature **414**: 166–167.

McMurdie, P. J., and S. Holmes. 2013. phyloseq: An R Package for Reproducible Interactive

Analysis and Graphics of Microbiome Census Data. PLoS One **8**: e61217.

Mevik, B. H., and R. Wehrens. 2007. The pls Package: Principle Component and Partial Least

Squares Regression in R. J. Stat. Softw. **18**: 1–24.

Meziti, A., D. Tsementzi, K. Ar. Kormas, H. Karayanni, and K. T. Konstantinidis. 2016.

Anthropogenic effects on bacterial diversity and function along a river-to-estuary gradient

in Northwest Greece revealed by metagenomics. Environ. Microbiol.

Necchi Jr, O. 2016. Heterokonts (Xanthophyceae and Chrysophyceae) in Rivers, p. 153–158. *In*

River Algae. Springer.

Newton, R. J., and S. L. McLellan. 2015. A unique assemblage of cosmopolitan freshwater

bacteria and higher community diversity differentiate an urbanized estuary from

oligotrophic Lake Michigan. Front. Microbiol. **6**: 1–13.

Newton, R., S. Jones, A. Eiler, K. McMahon, and S. Bertilsson. 2011. A guide to the natural

history of freshwater lake bacteria. Microbiol Mol Biol R **75**: 14–49.

Niño-García, J. P., C. Ruiz-González, and P. A. del Giorgio. 2016. Interactions between

hydrology and water chemistry shape bacterioplankton biogeography across boreal

freshwater networks. ISME J. , doi:10.1038/ismej.2015.226

Oksanen, J., F. G. Blanchet, R. Kindt, P. Legendre, P. R. Minchin, R. B. O'Hara, G. L. Simpson,

P. Solymos, M. H. H. Stevens, and H. Wagner. 2015. vegan: Community Ecology Package.

R package version 2.2-1. R Packag. version **1**: R package version 2.2-1.

Parada, A. E., D. M. Needham, and J. A. Fuhrman. 2015. Every base matters: assessing small

subunit rRNA primers for marine microbiomes with mock communities, time series and

global field samples. Environ. Microbiol.

Payne, J. T., J. J. Millar, C. R. Jackson, and C. A. Ochs. 2017. Patterns of variation in diversity

of the Mississippi river microbiome over 1,300 kilometers. PLoS One **12**: e0174890.

Pierson, S. M., B. J. Rosenbaum, L. D. McKay, and T. G. Dewald. 2008. Strahler Stream Order

and Strahler Calculator Values in NHDPlus.

Pruesse, E., C. Quast, K. Knittel, B. M. Fuchs, W. Ludwig, J. Peplies, F. O. Glockner, and F. O.

Glöckner. 2007. SILVA: a comprehensive online resource for quality checked and aligned

ribosomal RNA sequence data compatible with ARB. Nucleic Acids Res. **35**: 7188–7196.

Quast, C., E. Pruesse, P. Yilmaz, J. Gerken, T. Schweer, P. Yarza, J. Peplies, and F. O. Glöckner.

2013. The SILVA ribosomal RNA gene database project: Improved data processing and

web-based tools. Nucleic Acids Res. **41**: 590–596.

Rabalais, N. N., R. E. Turner, B. K. Sen Gupta, D. F. Boesch, P. Chapman, and M. C. Murrell.

2007. Hypoxia in the northern Gulf of Mexico: Does the science support the Plan to

Reduce, Mitigate, and Control Hypoxia? Estuaries and Coasts **30**: 753–772.

Rabalais, N. N., R. E. Turner, D. Justić, Q. Dortch, W. J. Wiseman, B. K. Sen Gupta, and D.

Justic. 1996. Nutrient Changes in the Mississippi River and System Responses on the

Adjacent Continental Shelf. Estuaries **19**: 386.

Rabalais, N. N., R. E. Turner, and W. J. Wiseman. 2002. Gulf of Mexico Hypoxia, a.K.a. "the

Dead Zone." Annu. Rev. Ecol. Syst. **33**: 235–263.

R Developement Core Team. 2015. R: A Language and Environment for Statistical Computing.

R Found. Stat. Comput. **1**: 409.

Read, D. S., H. S. Gweon, M. J. Bowes, L. K. Newbold, D. Field, M. J. Bailey, and R. I.

Griffiths. 2015. Catchment-scale biogeography of riverine bacterioplankton. ISME J **9**:

516–526.

Richey, J. E., J. M. Melack, A. K. Aufdenkampe, V. M. Ballester, and L. L. Hess. 2002.

Outgassing from Amazonian rivers and wetlands as a large tropical source of atmospheric

CO2. Nature **416**: 617–620.

Van Rossum, T., M. A. Peabody, M. I. Uyaguari-Diaz, K. I. Cronin, M. Chan, J. R. Slobodan,

M. J. Nesbitt, C. A. Suttle, W. W. L. Hsiao, P. K. C. Tang, N. A. Prystajecky, and F. S. L.

Brinkman. 2015. Year-long metagenomic study of river microbiomes across land use and

water quality   . Front. Microbiol.   **6**: 1–15.

Ruiz-González, C., L. Proia, I. Ferrera, J. M. Gasol, and S. Sabater. 2013. Effects of large river

dam regulation on bacterioplankton community structure. FEMS Microbiol. Ecol. **84**: 316–

331.

Russell, T. A., and L. Weller. 2013. State of the River Report: Water Quality and River Health in

the Metro Mississippi River. Friends of the Mississippi River.

Salcher, M. M., J. Pernthaler, and T. Posch. 2011. Seasonal bloom dynamics and ecophysiology

of the freshwater sister clade of SAR11 bacteria "that rule the waves"(LD12). ISME J. **5**:

1242–1252.

Salter, S. J., M. J. Cox, E. M. Turek, S. T. Calus, W. O. Cookson, M. F. Moffatt, P. Turner, J.

Parkhill, N. J. Loman, and A. W. Walker. 2014. Reagent and laboratory contamination can

critically impact sequence-based microbiome analyses. BMC Biol. **12**: 87.

Savio, D., L. Sinclair, U. Z. Ijaz, J. Parajka, G. H. Reischer, P. Stadler, A. P. Blaschke, G.

Blöschl, R. L. Mach, A. K. T. Kirschner, A. H. Farnleitner, and A. Eiler. 2015. Bacterial

diversity along a 2600 km river continuum. Environ. Microbiol. **17**: n/a-n/a.

Schilling, K. E., K.-S. Chan, H. Liu, and Y.-K. Zhang. 2010. Quantifying the effect of land use

land cover change on increasing discharge in the Upper Mississippi River. J. Hydrol. **387**:

343–345.

Schloss, P. D., S. L. Westcott, T. Ryabin, J. R. Hall, M. Hartmann, E. B. Hollister, R. a.

Lesniewski, B. B. Oakley, D. H. Parks, C. J. Robinson, J. W. Sahl, B. Stres, G. G.

Thallinger, D. J. Van Horn, and C. F. Weber. 2009. Introducing mothur: Open-source,

platform-independent, community-supported software for describing and comparing

microbial communities. Appl. Environ. Microbiol. **75**: 7537–7541.

Sinclair, L., U. Z. Ijaz, L. J. Jensen, M. J. L. Coolen, C. Gubry-Rangin, A. Chroňáková, A.

Oulas, C. Pavloudi, J. Schnetzer, A. Weimann, A. Ijaz, A. Eiler, C. Quince, and E. Pafilis.

2016. <tt>Seqenv</tt> : linking sequences to environments through text mining. PeerJ **4**:

e2690.

Singh, V. 2012. Application of Frequency and Risk in Water Resources: Proceedings of the

International Symposium on Flood Frequency and Risk Analyses, 14–17 May 1986,

Louisiana State University, Baton Rouge, USA, Springer Science & Business Media.

Smith, M. W., L. Herfort, K. Tyrol, D. Suciu, V. Campbell, B. C. Crump, T. D. Peterson, P.

Zuber, A. M. Baptista, and H. M. Simon. 2010. Seasonal Changes in Bacterial and Archaeal

Gene Expression Patterns across Salinity Gradients in the Columbia River Coastal Margin.

PLoS One **5**: e13312.

Staley, C., T. J. Gould, P. Wang, J. Phillips, J. B. Cotner, and M. J. Sadowsky. 2014a. Bacterial

community structure is indicative of chemical inputs in the Upper Mississippi River. Front

Microbiol **5**: 524.

Staley, C., T. J. Gould, P. Wang, J. Phillips, J. B. Cotner, and M. J. Sadowsky. 2014b. Core

functional traits of bacterial communities in the Upper Mississippi River show limited

variation in response to land cover. Front. Microbiol. **5**: 414.

Staley, C., T. J. Gould, P. Wang, J. Phillips, J. B. Cotner, and M. J. Sadowsky. 2015. Species

sorting and seasonal dynamics primarily shape bacterial communities in the Upper

Mississippi River. Sci. Total Environ. **505**: 435–45.

Staley, C., T. J. Gould, P. Wang, J. Phillips, J. B. Cotner, and M. J. Sadowsky. 2016. Sediments

and Soils Act as Reservoirs for Taxonomic and Functional Bacterial Diversity in the Upper

Mississippi River. Microb. Ecol. **71**: 814–824.

Staley, C., T. Unno, G. TJ, B. Jarvis, J. Phillips, C. JB, T. J. Gould, B. Jarvis, J. Phillips, J. B.

Cotner, and M. J. Sadowsky. 2013. Application of Illumina next-generation sequencing to

characterize the bacterial community of the Upper Mississippi River. J. Appl. Microbiol.

115: 1147–1158.

Thrash, J. C., J. L. Weckhorst, and D. M. Pitre. 2015. Cultivating Fastidious Microbes, p. 1–22.

In Protocols for Metagenomic Library Generation and Analysis in Petroleum Hydrocarbon

Microbe Systems. Humana Press.

Trusova, M. Y., and M. I. Gladyshev. 2002. Phylogenetic diversity of winter bacterioplankton of

eutrophic siberian reservoirs as revealed by 16S rRNA gene sequence. Microb. Ecol. **44**:

252–259.

Turner, R. E., and N. N. Rabalais. 2003. Linking landscape and water quality in the Mississippi

river basin for 200 years. Bioscience **53**: 563–572.

Turner, R. E., and N. N. Rabalais. 2004. Suspended sediment, C, N, P, and Si yields from the

Mississippi River Basin. Hydrobiologia **511**: 79–89.

Vannote, R. L., G. W. Minshall, K. W. Cummins, J. R. Sedell, and C. E. Cushing. 1980. River

continuum concept. Can J Fish Aquat Sci **37**: 130–137.

Vorosmarty, C. J., P. B. McIntyre, M. O. Gessner, D. Dudgeon, A. Prusevich, P. Green, S.

Glidden, S. E. Bunn, C. A. Sullivan, C. R. Liermann, and P. M. Davies. 2010. Global

threats to human water security and river biodiversity. Nature **467**: 555–561.

Weiss, S., A. Amir, E. R. Hyde, J. L. Metcalf, S. J. Song, and R. Knight. 2014. Tracking down

the sources of experimental contamination in microbiome studies. Genome Biol. **15**: 564.

Withers, P. J. A., and H. P. Jarvie. 2008. Delivery and cycling of phosphorus in rivers: A review.

Sci. Total Environ. **400**: 379–395.

Wood, S. A., M. J. Prentice, K. Smith, and D. P. Hamilton. 2010. Low dissolved inorganic

nitrogen and increased heterocyte frequency: precursors to Anabaena planktonica blooms in

a temperate, eutrophic reservoir. J. Plankton Res. **32**: 1315–1325.

Zeglin, L. H. 2015. Stream microbial diversity in response to environmental changes: review and

synthesis of existing research. Front. Microbiol. **6**: 454.

Zwart, G., B. C. Crump, M. P. Kamst-van Agterveld, F. Hagen, and S. K. Han. 2002. Typical

freshwater bacteria: An analysis of available 16S rRNA gene sequences from plankton of

lakes and rivers. Aquat. Microb. Ecol. **28**: 141–155.

927   **Tables**

Table 1. Correlations between prokaryotic and eukaryotic submodules to nitrate and phosphate, as well as PLS model results

Fraction	Gene	Nutrient	Submodule	Eigen Correlations	Total OTUs	PLS model
0.2-2.7 μm	16S	NO_3^{2-}	Blue	r = 0.6, P = 7e-08	77	R^2=0.35; corr = 0.65, P =1.e-09
> 2.7 μm	16S	NO_3^{2-}	Brown	r = 0.56, P = 3e-07	133	R^2 = 0.69; corr = 0.83, P= < 2.2e-16
0.2-2.7 μm	16S	PO_4^{3-}	Turquoise	r = 0.53, P = 2e-06	51	R^2=0.80; corr = 0.89, P = < 2.3e-16
> 2.7 μm	16S	PO_4^{3-}	Yellow	r = 0.53, P = 1e-06	80	R^2= 0.48; corr= 0.77, P = 4.88e-15
0.2-2.7 μm	18S	NO_3^{2-}	Green	r = 0.39, P = 6e-04	39	R^2=0.38; corr = 0.62, P =2.97e-9
> 2.7 μm	18S	NO_3^{2-}	Brown	r = 0.52, P = 2e-06	59	R^2=0.572; corr = 0.759, P= 6.7e-15
0.2-2.7 μm	18S	PO_4^{3-}	Turquoise	r = 0.56, P = 2e-07	56	R^2=0.80; corr = 0.89, P = < 2e-16
> 2.7 μm	18S	PO_4^{3-}	Red	r = 0.60, P = 2e-08	39	R^2 = 0.618; corr = 0.799, P = < 2e-16

931   **Figure Legends**
932
Figure 1. Sampling map (A) with graph inserts that represent the measured discharge rate (cubic

feet second^{-1} [CFS]) as recorded on the USGS gauges, and six environmental parameters

measured along the transect, according to concentration or visible depth of secchi disk by

distance (B). Throughout the figure and text, blue and red dots represent sampling

locations above and below the Missouri River confluence, respectively, and are designated

throughout as "Upper" and "Lower." Cell Counts only represent the < 2.7 μm fraction.

Figure 2. Non-metric multidimensional scaling (NMDS) results for whole community with

correlations to environmental parameters (phosphate, nitrate, nitrite, ammonia, distance

(km), water temperature, turbidity (cm), river speed (mph)) and the top ten OTUs based on

significance (P) and strength of correlation (r). The four plots represent > 2.7 μm (A, C)

and 0.2-2.7 μm (B, D) fractions for the 16S (A, B) and 18S (C, D) rRNA gene

communities. Vector length is proportional to the strength of the correlation.

Figure 3. Relative abundance, by phylum, according to transect distance, for phyla accounting

for > 0.1% of the total reads for the 16S rRNA gene > 2.7 μm (A) and 0.2-2.7 μm (B)

communities. Non-linear regressions with 95% CI (gray shading) are provided for

reference.

Figure 4. Relative abundance, by phylum, according to transect distance, for phyla accounting

for > 0.1% of the total reads for the 18S rRNA gene > 2.7 μm (A) and 0.2-2.7 μm (B)

communities. Non-linear regressions with 95% CI (gray shading) are provided for

reference.

Figure 5. Core microbiome aggregate abundance for the 16S (A) and 18S (B) rRNA gene. In

each, triangles and circles points represent 0.2-2.7 μm and > 2.7 μm fractions,

respectively. Non-linear regressions with 95% confidence intervals (CI) (gray shading) are

provided for reference.

Figure 6. PLS results for the 16S rRNA gene community submodules most associated with

phosphate (A) and nitrate (B). OTU correlation with a given nutrient is indicated on the y-

axis according to the number of co-correlations (node centrality) on the x-axis.

Community fractions: 0.2-2.7 μm (A) and > 2.7 μm (B). Circle size is proportional to VIP

scores, with top 10 VIP scoring and top node centrality OTUs labeled with their highest-

resolution taxonomic classification and OTU number. Colors represent the taxonomic

classification the phylum level.

Figure 7. PLS results for the 18S rRNA gene community submodules most associated with

phosphate and nitrate. OTU correlation with a given nutrient is indicated on the y-axis

according to the number of co-correlations (node centrality) on the x-axis. Community

fractions: 0.2-2.7 μm (A) and > 2.7 μm (B,C). Circle size is proportional to VIP scores,

with top 10 VIP scoring and top node centrality OTUs labeled with their highest-

resolution taxonomic classification and OTU number. Colors represent the taxonomic

classification the phylum level.

 Supplemental Figure Legends

Figure S1. NMDS results for the 16S (A) and 18S (B) rRNA gene communities. In each,

circles and triangles represent the > 2.7 µm and 0.2-2.7 µm fractions, respectively.

Figure S2. Hclust results for the 16S (A) and 18S (B) rRNA gene communities.

Figure S3. Richness and Evenness indexes for the 16S > 2.7 µm (A) and 0.2-2.7 µm (B) and

18S > 2.7 (C) and 0.2-2.7 µm (D) communities. Non-linear regressions with 95% CI (gray

shading) are provided for reference.

Figure S4. Separated upper and lower MSR core microbiome aggregate abundance for the 16S

rRNA gene communities. For the upper and lower river, the core microbiome was defined

separately requiring OTUs to have greater than one read in 90% of the samples. > 2.7 µm

(A) and 0.2-2.7 µm (B) 16S rRNA gene communities in the upper and lower MSR.

Figure S5. Relative abundance, by environmental ontology group, according to transect

distance for the 16S rRNA gene > 2.7 µm (A) and 0.2-2.7 µm (B) communities. Non-

linear regressions with 95% CI (gray shading) are provided for reference.

Figure S6. PLS results for the 0.2-2.7 µm and > 2.7 µm 16S (A-D) and 18S (E-H) rRNA gene

community for selected submodules with nitrate and phosphate and a VIP score > 1.

Correlation of submodule OTUs to nitrate and phosphate according to the number of co-

correlations (node centrality) for 0.2-2.7 µm (A,B, E,F) and > 2.7 µm (C,D,G,H) 16S and

18S rRNA gene communities. Circle size is proportional to VIP scores, with top 10 VIP

scoring and top node centrality OTUs labeled with their highest-resolution taxonomic

classification and OTU number. Colors represent the taxonomic classification the phylum

level.

Figure S7. WCGNA results for 0.2-2.7 µm 16S rRNA gene community submodules of interest

based on Pearson correlations to nitrate and phosphate (A, B, D) and the resultant PLS

regression of the predicted versus measured nutrient concentrations (C, E). Linear

regressions with 95% CI (gray shading) are provided for reference.

Figure S8. WCGNA results for 0.2-2.7 µm 18S rRNA gene community submodules of interest

based on Pearson correlations to nitrate and phosphate (A, B, D) and the resultant PLS

regression of the predicted versus measured nutrient concentrations (C, E). Linear

regressions with 95% CI (gray shading) are provided for reference.

Figure S9. WCGNA results for > 2.7 µm 16S rRNA gene community submodules of interest

based on Pearson correlations to nitrate and phosphate (A, B, D) and the resultant PLS

regression of the predicted versus measured nutrient concentrations (C, E). Linear

regressions with 95% CI (gray shading) are provided for reference.

Figure S10. WCGNA results for > 2.7 µm 18S rRNA gene community submodules of interest

based on Pearson correlations to nitrate and phosphate (A, B, D) and the resultant PLS

regression of the predicted versus measured nutrient concentrations (C, E). Linear

regressions with 95% CI (gray shading) are provided for reference.

1024

1025

1026

1027

1028

1035 **Supplemental Tables**
1036
1037 Supplemental Table S1 is a spreadsheet, TableS1.xlsx. This includes site data, NMDS

1038 correlations for environmental data, Relative abundances of the Core taxa, SeqENV data, 16S

1039 and 18S rRNA OTU tables, and VIP scores for all samples and datum.

1040

1041 Additional Supplemental Information, including R scripts and our Mothur workflow, are hosted

1042 on the Thrash Lab website at: thethrashlab.com/publications.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5