Differential Adaptation to a Harsh Granite Outcrop Habitat between Sympatric *Mimulus*

Species

by

Kathleen G. Ferris[1,*,δ] and John H. Willis[1]

[1] Department of Biology, Duke University, 125 Science Drive Durham, NC 27705

kgferris@ucdavis.edu, jwillis@duke.edu

Running Title: Differential Habitat Adaptation *Mimulus*

Abstract: 149 words, Main text: 4,891 words

Figures: 3, Tables: 2

*For correspondence kgferris@ucdavis.edu, (314) 550-1882

δCurrent Address: *Center for Population Biology, 2320 Storer Hall, University of California*

Davis, One Shields Avenue Davis, CA 95616

Word counts: Total 741; Abstract 178

Figures: 5, Tables: 4, Supporting Information: 2 Figures & 5 Tables

ABSTRACT

A primary goal in evolutionary biology is to understand which environmental variables and traits drive adaptation to harsh environments. This is difficult since many traits evolve simultaneously as populations or species diverge. Here we investigate the ecological variables and traits that underlie *Mimulus laciniatus'* adaptation to granite outcrops compared to its sympatric, mesic-adapted progenitor *M. guttatus*. We use fine scale measurements of soil moisture and herbivory to examine differences in selective forces between the species' habitats, and measure selection on flowering time, flower size, plant height, and leaf shape in a reciprocal transplant using *M. laciniatus x M. guttatus* F_4 hybrids. We find that differences in drought & herbivory drive survival differences between habitats, that *M. laciniatus* and *M. guttatus* are each better adapted to their native habitat, and differential habitat selection on flowering time, plant stature, and leaf shape. We conclude that while early flowering time, small stature, and lobed leaf shape underlie plant fitness in *M. laciniatus'* seasonally dry environment, increased plant size is advantageous in a competitive mesic environment replete with herbivores like *M. guttatus'*.

Key Words: differential adaptation, flowering time, *Mimulus*, plasticity, drought, phenotypic selection

INTRODUCTION

Closely related species often occupy ecologically disparate habitats. How do different

taxa adapt to these new and often stressful environments? What traits evolve in response

to novel selection pressures as a species colonizes a new habitat? And which

environmental variables act as selective forces? These are exciting questions, especially

in sympatric species since adaptation to different habitats can lead to ecological

speciation (Mayr 1947; Mayr 1949; Schluter 2001; Coyne & Orr 2004; Rundle & Nosil

2005) or reinforce reproductive isolation between species that have come into secondary

contact through selection against migrants or ecologically unfit hybrids (Coyne & Orr

2004). Reciprocal transplant studies have demonstrated differential habitat adaptation

between many populations and species, however little is still known about the specific

environmental variables and traits behind this adaptation (reviewed in Kawecki & Ebert

2004; Hereford 2009; Salvolainen et al. 2013). This is especially true for closely related

species that occupy different habitats. Identifying particular selective agents causing

adaptive divergence is a challenge because habitats vary in many ecological variables,

and tracking variation in particular ecological variables and their effect on fitness in the

field is labor intensive (Wade & Kalisz 1990, MacColl 2011). Understanding the adaptive

significance of individual phenotypes is difficult since as species diverge many traits

evolve simultaneously due to both adaptive and neutral demographic processes. While

many studies have examined natural selection on phenotypes in the wild using the

multivariate approach developed by Lande and Arnold (1983; reviewed in Kingsolver et

al. 2001), the majority of them measured selection in natural populations where many

traits are genetically correlated with each other due to shared demographic history. This

extensive genetic correlation among traits makes inferring direct selection on any one

phenotype problematic even with a multivariate statistical approach (Hall & Willis 2006).

This problem becomes even larger when selection is measured on phenotypic differences

between species rather than between populations, since species have a longer history of

divergence.

Combining fine scale environmental measurements, controlled crosses between

species, and reciprocal transplant experiments can overcome these obstacles. Collecting

fine scale ecological data in a reciprocal transplant experiment makes it possible to test

whether individual ecological variables are associated with fitness and therefore whether

they act as selective agents in a particular habitat (Huber et al. 2004; Garant et al. 2007;

Lind & Johansson 2007; Quinn et al. 2009; Weese et al. 2010). In a segregating hybrid

population trait associations present in the parental lines are broken up through

recombination. This disentangles phenotypes from their species-specific genetic

background. Using a segregating hybrid population in a reciprocal transplant experiment

therefore allows the measurement of selection on individual traits in the wild (Hall and

Willis 2006; Anderson et al. 2011b, Agren et al. 2016).

Plant species are excellent systems for studying differential habitat adaptation.

Because of their sessile lifestyle, plants often experience strong divergent selection across

heterogeneous environments on a small geographic scale (Kalisz 1986; Schmitt &

Antonovics 1986; Stewart & Schoen 1987; Robichaux 1990). Variation in soil moisture

is a major driver of plant species' distribution and diversification (Stebbins 1957;

Axelrod 1972). Certain traits have evolved repeatedly in plant species that occur in dry,

marginal habitats. Early flowering allows plants to reproduce before the onset of summer

drought and has been shown to be adaptive across plant species (Kiang & Hamrick 1987;

Fox 1990, Macnair & Gardner 1998; Stanton et al. 2000; Hall & Willis 2006; Willis et al.

2008; Anderson et al. 2012). Plant taxa occupying harsh, marginal habitats also often

have self-fertilizing mating systems (Stebbins 1957; Stebbins 1970; Kiang & Hamrick

1987; MacNair & Gardner 1998; Mazer et al. 2010; Wu et al. 2010). Small flower size

and rapid floral development are correlated with self-fertilization and may reduce floral

tissue transpiration in dry environments (Galen 1999). Additionally, self-fertilization

provides reproductive assurance in marginal environments often sparsely populated by

pollinators (Stebbins 1970; Piper et al. 1986; Cunningham 2000; Fausto et al. 2001).

Diminutive plant stature is adaptive in dry environments because small plants require less

water for proper development and therefore are less stressed under drought conditions

(Blum and Sullivan 1997; Blum et al. 1997; Pantuwan et al. 2002). Small plants also tend

to have more rapid development and flowering time (Thornsberry et al. 2001; Bolmgren

and Cowan 2008; Wei et al. 2010). Leaf shape is also hypothesized to be adaptive in dry

environments because it affects the gas and heat exchange traits of the plant. Lobed

leaves have thinner boundary layers and are cooled more efficiently by convection than

round leaves. More efficient convective cooling should decrease the amount of

transpiration required when leaves are heated above ambient temperature by direct

sunlight (Vogel 1968; Givnish 1978; Schuepp 1993; Nobel 2005). Lobed leaves also

have lower hydraulic resistance (R_{leaf}) and therefore less drought stress prone tissue than

round leaves (Nicotra et al. 2011). A thinner boundary layer & lower R_{leaf} should be

advantageous in dry, exposed habitats.

The *Mimulus guttatus* species complex is an excellent system for studying

adaptation to harsh marginal habitats because it is a closely related group of wildflowers

that are largely inter-fertile and occupy a variety of edaphic environments across Western

North America. *Mimulus guttatus*, the putative progenitor of the species complex, occurs

in moist seeps and streambeds, but several smaller ranging sympatric species have

colonized rapidly drying habitats such as serpentine soils, copper mine tailings, and

granite outcrops (MacNair & Gardner 1998; Wu et al. 2007; Ferris et al. 2015; Wright et

al. bioRxiv). Granite outcrops are harsh habitats characterized by shallow rocky soils,

high light intensity, extreme temperatures, and low soil water retention (Ferris et al.

2014). The Sierra Nevada natives *M. laciniatus* (Figure 1b) and the recently described *M.*

filicifolius (Sexton et al. 2013, Ferris et al. 2014) specialize on granitic habitat (Figure

1a). These species grow in thin strips of moss and gravel where few other plant taxa can

thrive. Both granite outcrop specialists are sympatric with *Mimulus guttatus* (Figure 1d)*,*

which occurs in deeper, densely populated soils in moist seeps and meadows (Figure 1e,

Wu et al. 2007). *Mimulus guttatus* can often be found growing in mesic habitat within

meters of *M. laciniatus'* granite outcrops (Ferris et al. 2016). *Mimulus laciniatus* and *M.*

filicifolius share several traits that are likely adaptive in rocky outcrops that dry rapidly

once seasonal snowmelt is gone: early and rapid flowering (Friedman & Willis 2013;

Ferris et al. 2016), a self-fertilizing mating system (Fenster & Ritland 1994; Ferris et al.

2014), small plant size, and lobed leaf shape (Sexton et al. 2013, Ferris et al. 2015, Ferris

et al. 2016). Lobed leaf shape is unique to rocky outcrop *Mimulus* taxa (Ferris et al.

2015), and is seemingly an example of parallel evolution (Ferris et al. 2014), therefore it

is also likely adaptive in this marginal habitat. In contrast *M. guttatus* is round-leaved,

larger, later-flowering, and predominantly outcrossing (Wu et al. 2007, Ferris et al.

2016).

In this study we investigate how *M. laciniatus* has adapted to its marginal granite

outcrop environment using a reciprocal transplant experiment with *M. laciniatus*, *M.*

guttatus, and a large segregating outbred F_4 hybrid population planted in replicated native

granite and meadow habitats. Trait associations present in the parental species are broken

up by recombination and independent assortment in F_4's allowing the measurement of

selection on individual phenotypes. To understand adaption to *M. laciniatus'* harsh

granite outcrop habitats in we asked three questions: (1) What ecological variables act as

selective agents in each species' habitat? (2) Are *M. laciniatus* & *M. guttatus*

differentially adapted and if so are there fitness trade-offs? (3) What traits are involved in

this adaptation? In a previous survey of microclimatic variation we found that *M.*

laciniatus' granite outcrop habitat is significantly drier than adjacent meadow and

streambed habitat occupied by *M. guttatus* (Ferris et al. 2014). We also observed lower

herbivore density in *M. laciniatus'* granite outcrop habitat than in *M. guttatus'* meadows

(K. Ferris personal observation). Soil moisture and herbivory are important factors in

plant distribution and diversification (Stebbins 1952; Ehrlich and Raven 1964; Axelrod

1972; Janz 2011). To characterize differences in selective forces between *M. laciniatus*

and *M. guttatus'* native habitats we measured variation in these two variables on a fine

spatial scale throughout our reciprocal transplant, and tested whether variation in soil

moisture or herbivory was associated with survival in each habitat. To test for differential

adaption and trade-offs between species we recorded fitness differences between *M.*

laciniatus & *M. guttatus* in granite and meadow habitats. To examine individual traits

that contribute to *M. laciniatus'* adaptation to a marginal granite outcrop habitat we

measured selection in the field on phenotypes that the granite specialists, *M. laciniatus*

and *M. filicifolius*, share and which are also likely adaptive in seasonally dry habitats:

flowering time, flower size, leaf size, plant size, and a lobed leaf shape in the *M.*

laciniatus x M. guttatus F_4 hybrids. If *M. laciniatus* and *M. guttatus* are differentially

adapted to their respective microhabitats and flowering time, flower size, leaf size, plant

size, and leaf shape contribute to this adaptation, then we expect there to be divergent

selection on these traits between habitats in the direction of species differences.

MATERIALS AND METHODS

Construction of the hybrid population

In the summer of 2008 we collected individuals from each parental population, White

Wolf (WLF, *M. laciniatus*) and Yosemite Overlook (YVO, *M. guttatus*), along Tioga

Road in Yosemite National Park, CA. The parental inbred lines, WLF47 and YVO6,

were developed through hand pollinated self-fertilization and single seed descent for 7

and 5 generations respectively. The experimental population of outbred F_4 hybrids was

created by first crossing the parental lines WLF47 and YVO6 to generate F_1 hybrids, then

self-fertilizing F_1's to generate a large F_2 population. Two hundred F_2's were randomly

paired and reciprocally crossed to generate 200 maternal families. Then 200 F_3's were

randomly paired and reciprocally crossed (Figure 1c). We pooled ~30 seeds from each of

the 200 maternal F_3's to create one large outbred F_4 family. F_4 seeds were then randomly

distributed throughout experimental blocks.

Reciprocal transplant experiment

In April of 2013, 400 *M. laciniatus* (WLF47), 400 *M. guttatus* (YVO6), and 3000 F_4's

were planted in open flats of soil in the UC Davis greenhouses. Seeds were cold stratified

at 4°C for 10 days, then left in the greenhouse to germinate for one week. At the

cotyledon stage seedlings were transplanted one inch apart into 50 randomized blocks of

19 plants each (2 WLF47, 2 YVO6, 15 F_4) at each of four field sites along Tioga Road in

Yosemite NP, CA. The two *M. laciniatus* habitat sites, Olmstead Point (Granite 1, Figure

1a) and Yosemite Creek (Granite 2), are undisturbed granite outcrops with native *M.*

laciniatus growing on moss at elevations of 8,500 and 7,500 feet respectively. The *M.*

guttatus sites, Little Meadow (Meadow 1) and Crane Flat (Meadow 2, Figure 1e), are

undisturbed meadows with native *M. guttatus* growing near a standing seep. They occur

at 6,200 and 6,000 feet respectively. Sites were chosen to maximize the similarity

between the developmental stages of transplanted and native plants of each species.

Seedlings that died within a week of transplant were replaced so that transplant

shock would not affect survivorship. Experimental plants were censused every other day

May - August 2013 for survival and flowering time. On the day of first flower, flowering

time, corolla width, and plant height were recorded. Morphological measurements were

taken with a small metal ruler in millimeters. Leaves were not collected for leaf shape

measurement until plants began to senesce to avoid damaging plants and imposing

selection before they set seed. Leaf area and degree of lobing were determined by

digitally scanning leaves and performing convex hull analysis in ImageJ as described

previously (Ferris et al. 2015). Briefly, the convex hull analysis consists of comparing the

area of each leaf's convex hull (the shape created by connecting the outermost points of a

leaf) to the leaf's true area and dividing this difference in area by the convex hull area to

control for size. Fruit number was counted after senescence as a measure of lifetime

fitness.

Measuring environmental agents of selection

We collected fine scale soil moisture and herbivore damage data in each block across

each transplant site. This allowed us to test whether these environmental variables are

associated with plant fitness differences on a fine spatial and temporal scale within and

between habitat types. We used survival as our measure of fitness in this analysis because

we had data on survival in every block in each site giving us greater spatial variation in

survival than in fruit number since only a subset of experimental blocks produced fruit.

Percent soil moisture was measured weekly in each experimental block May through

August using a Decagon soil moisture probe. We measured herbivory in each block at the

time of plant senescence by recording the presence or absence of damage on the first true

leaf.

ANALYSES

Selective environment

To determine whether drought and herbivory contributed to the difference in

species' selective environments we measured percent soil moisture and leaf damage in

each experimental block of our reciprocal transplant experiment. We tested if soil

moisture and leaf damage affected plant fitness across habitats with a linear mixed effects

model where soil moisture was treated as a repeated measure using the nlme package in

R (R Core Development Team 2008). In our mixed effects model survival was the

dependent variable with habitat type, soil moisture, leaf damage, and time as fixed

independent variables, and block as a random effect nested within site, nested within

habitat. To determine whether different environmental variables were associated with

plant survival in granite vs. meadow sites we tested for interactions between our

environmental variables and habitat type. We also ran separate mixed effects models on

the survival data from each habitat. We determined the best-fit model using AIC model

selection criteria (Zuur et al. 2009) with the model selection R-package MuMin (Bartoń

2013). To determine whether there was an absolute difference in herbivore damage

between habitat types regardless of its effect on survival, we performed a t-test. If soil

moisture and herbivory are selective agents in either granite or meadow habitats then we

expect a strong association between the environmental variable and survival within a

transplant site. If either variable contributes to differential selection between habitat

types, then we expect a stronger association in one habitat over the other.

Habitat adaptation and trait differences

To test for differential habitat adaptation between *M. laciniatus* and *M. guttatus*

we performed two-tailed t-tests to determine whether mean survival to flowering and fruit

number significantly differed between *M. laciniatus* and *M. guttatus* by habitat. We also

conducted a logistic regression with survival to flowering as the dependent variable and

species, habitat, and their interaction as independent variables. If *M. laciniatus* and *M.*

guttatus are differentially adapted then we expect each species to have a fitness

advantage in its native habitat. If there is a species trade-off between habitats we expect

to see an interaction between habitat type and species in our regression. Based on our

phenotypic results we also decided to look for species differences in the degree and

direction of phenotypic plasticity in each trait. To detect phenotypic plasticity in

flowering time, flower size, plant height, leaf area, and leaf shape we conducted factorial

ANOVAs with habitat (Granite vs. Meadow) & genotypic class (*M. laciniatus, M.*

guttatus, & F$_4$) as independent variables. To detect genotype by environment (GxE)

interactions we tested for interactions between habitat and genotypic class. To correct for

multiple testing we used a Bonferroni correction ($\alpha = 0.01$). If there is a difference in the

degree or direction of phenotypic plasticity between species, then we should see a

significant interaction between genotypic class and habitat (GxE). Phenotypic

correlations were measured among traits for each habitat type using a restricted

maximum likelihood model (REML) in JMP (JMP®, Version *10*. SAS Institute Inc.,

Cary, NC, 1989-2007).

Phenotypic selection analysis

To measure the strength of selection on individual phenotypes we conducted

linear and quadratic selection analysis (Lande & Arnold 1983; Mitchell-Olds & Shaw

1987). We used lifetime fitness (fruit number) of plants that survived to flower as our

fitness measure in our selection analysis because we only have phenotypic values of

plants that survived to flower. Fecundity, fruit number in our case, is also the best

measure of fitness for an annual plant. We used a generalized linear mixed model to

regress lifetime fitness simultaneously on flowering time, corolla width, leaf area, plant

height, and leaf shape in our F_4 population. To detect differential habitat selection we

tested for interactions between trait and habitat type. We also measured the strength of

selection on each phenotype in each habitat by performing a regression on data from each

habit type separately. Block and site were included in the model as random effects where

block was nested within site, which was nested within habitat. The strength of selection

on each trait in each habitat was measured with linear (β') and quadratic (γ) regression

coefficients. All phenotypes were standardized to a mean of 0 and standard deviation of 1

to enable comparison of traits measured in different units and to values from other

phenotypic selection studies (Lande & Arnold 1983).

To control for excess zeros in our lifetime fitness measure we broke our data up

into two distinct analyses: (1) a binomial logistic regression on all plants the flowered

whether or not they produced fruits, and (2) a truncated Poisson regression on only plants

that did produce fruits (as in Anderson et al. 2012, Anderson et al. 2015) using the R

package glmmADMB (Skaug et al. 2006). The binomial distributed logistic regression

determines whether a trait was associated with the probability of a plant setting seed or

not. The truncated Poisson regression on the count data for plants that did fruit tested

whether phenotypes were associated with the number of fruits a plant produced (Ridout

et al. 1998). Model selection was performed using AIC selection criteria (Zuur et al.

2009) with the model selection R-package MuMin (Bartoń 2013). Phenotypes and

interactions included in the top model were considered to be under selection. If the

phenotypes we have measured contribute to differential habitat adaptation between these

species we expect divergent selection in the direction of the native species trait value. For

example, if flowering time is involved in differential adaptation we would expect there to

be selection for earlier flowering in *M. laciniatus'* granite outcrops, but for later

flowering in *M. guttatus'* meadow habitat.

RESULTS

Drought and herbivory are differentially associated with survival differences between

habitats

When the environmental data from all sites was analyzed together our best-fit mixed

effects model included both soil moisture and leaf damage, indicating that both

significantly affected plant survival across habitats (Table 1). Our best model also

included an interaction between soil moisture & habitat and a three-way interaction

between soil moisture, habitat, and time indicating that soil moisture affects survival

differently in each habitat over time (Table 1). Specifically, while soil moisture is in fact

higher in granite outcrops at the beginning of the growing season, it then decays rapidly

to effectively zero over the course of approximately one week mid-summer (Figure 2a).

In contrast, soil moisture in the meadow habitat decays in a gradual linear fashion over

the course of the entire season (Figure 2b). Survival tracks soil moisture with a slight

delay in both granite and meadow habitats (Figure 2c&d). When data from each habitat

type was analyzed separately soil moisture was associated with survival in both, but leaf

damage only affected survival in meadows (Table 1). There was also significantly more

herbivore damage, and therefore more herbivore pressure, in the meadows than in granite

(p-value < 0.001, Figure 3).

***M. laciniatus* & *M. guttatus* are differentially adapted and phenotypically plastic**

There was strong viability selection in each site with the total proportion surviving to

flower ranging from 0.18-0.57 (G1 = 0.45, G2 = 0.22, M1 = 0.18, M2 = 0.57). *M.*

laciniatus had significantly higher survival than *M. guttatus* in its native granite (0.23 >

0.12, p-value < 0.0001), but the species had similar survival in *M. guttatus'* meadows (*M.*

guttatus = 0.09, *M. laciniatus* = 0.10, p-value = 0.72, Figure 4a). However, a fitness

trade-off in fecundity was indicated by crossing reaction norms (Figure 4b). *M. laciniatus*

has significantly higher mean fruit number in granite habitat (mean = 0.48, SE = 0.05)

than *M. guttatus* (mean = 0.10, SE = 0.022, p-value <0.0001), while *M. guttatus* has

marginally significantly higher mean fruit number than *M. laciniatus* in its meadows (*M.*

guttatus = 0.05, SE = 0.022, *M. laciniatus* = 0.007, SE = 0.007, p-value = 0.055).

In addition, our logistic regression of survival to flowering detected a significant

interaction between habitat and species indicating that parental lines are differentially

adapted (Table S1).

To understand how ecologically important traits varied between habitats and

species we calculated the mean trait value and standard error for F_4's, *M. laciniatus*, and

M. guttatus in each habitat and transplant site. In granite habitat phenotypic differences

between genotypes are in the expected direction with *M. laciniatus* flowering earlier (by

3.29 days), being smaller flowered (by 2.88 mm) and leaved (by 51.84 mp), shorter (by

2.64), and more lobed (by 0.07) than *M. guttatus* (Table 2). In the meadow habitat the

direction of trait differences was the same, except that *M. laciniatus* flowered 3.5 days

later than *M. guttatus* (Table 2).

We saw plasticity in our focal traits between habitats and decided to test whether

there was significant phenotypic plasticity and genetic variation for plasticity between *M.*

laciniatus and *M. guttatus*. There was significant phenotypic plasticity in all traits except

leaf area, significant GxE in flowering time (p-value = 0.007), and marginally significant

GxE in plant height (p-value = 0.022). Within each genotypic class (F_4, *M. laciniatus*, *M.*

guttatus) plants flowered earlier (8.57, 8.8, 2.05 days, p-value < 0.001) and were shorter

(4.63, 17.94, 43.23 mm, p-value < 0.001) in *M. laciniatus'* granite habitats than in *M.*

guttatus' meadows (Table 2, Figure 5). Interestingly *M. laciniatus* was significantly more

plastic in flowering time, but less plastic in height than *M. guttatus* (p-value = 0.007,

0.022, Figure 5&c). F_4's and *M. laciniatus* had larger flowers (2.2 mm) in granite, while

M. guttatus had larger flowers (1 mm) in meadows (p-value < 0.001). *M. laciniatus* &

F_4's showed a slight, but significant (p-value = 0.002) increase in leaf lobing in the

granite habitat (Tables 1, Figure 5e).

We examined phenotypic correlations among traits in the F_4's (Table 3).

Phenotypic correlations in a segregating hybrid population can provide insight

concerning the genetic correlations among traits. In both habitats leaf area, flower size,

and height at first flower were highly positively correlated (r > 0.5), while leaf area and

shape are slightly positively correlated (granite r = 0.079, meadow r = 0.136). This

suggests that there is a genetic correlation among these traits in both habitats. Flowering

time was uncorrelated with other traits in granite (Table 3), but positively correlated with

height ($r = 0.110$) and leaf area ($r = 0.128$) in meadows. This difference suggests that the

genetic architecture of flowering time could differ between habitats.

Early flowering time is adaptive in granite outcrops, while being large is adaptive in

meadows

To assess selection on individual phenotypic traits in each habitat we performed linear

and quadratic selection analysis in our outbred F_4 population using a combination of

binomial logistic and truncated Poisson generalized linear mixed models (GLMM). In the

binomial logistic model we tested whether phenotypic values were correlated with the

probability that an individual F_4 produced fruit or not. Our best-fit binomial model

included linear selection gradients on flower size, plant height, leaf shape, flowering

time, a negative quadratic selection gradient on flowering time, an interaction between

flowering time and plant height, and an interaction between habitat type and flowering

time (Table 4). In the granite habitat we found directional selection for larger flower size,

strong selection for smaller plant size and earlier flowering time, and weak selection for

more highly lobed leaves (Table 5). We also found a signature of stabilizing selection on

flowering time in granite indicated by a negative quadratic selection gradient (Table 5;

Lande & Arnold 1983). In the meadow habitat we saw selection for earlier flowering

time, smaller flowers, less lobed leaves, and strong selection for larger plant size (Table

5). We also detected a signature of stabilizing selection on plant height in meadows. The

interaction between flowering time and plant height in our top binomial model suggests

correlated selection on these two traits and a potential functional correlation (Lande &

Arnold 1983). Selection for earlier flowering time in granite was significantly stronger

than selection for early flowering in the meadow habitat indicated by the interaction

between habitat and flowering time in our top binomial model (Tables 4&5). We list all

binomial distributed generalized linear mixed models testing for phenotypic selection in

the supplementary materials (Table S2).

In the truncated Poisson regression, which tested for associations between

phenotypic value and fruit number among plants that did set seed, our best-fit model

included linear selection gradients for flowering time, flower size, plant height, leaf

shape, an interaction term for leaf shape and plant height, and interactions between

flowering time and habitat and leaf shape and habitat (Table 4). There was directional

selection for earlier flowering, larger flowers, taller plants, and more highly lobed leaves

in the granite habitat (Table 6). In the meadow habitat we found weaker selection for

earlier flowering time, selection for larger flowers, and stronger selection for taller plants

(Table 6). We found interactions between habitat type and flowering time and habitat and

leaf shape in our top Poisson model (Table 6) indicating that there is differential habitat

selection on these two phenotypes. In our second to top model we also saw an interaction

between habitat and plant height (Table S3) suggesting that height may also be under

differential selection. We list all Poisson distributed generalized linear mixed models

testing for phenotypic selection in the supplementary materials (Table S3). The

interaction between leaf shape and plant height in our top truncated Poisson model

suggests correlated selection on these two traits and a potential functional correlation

between them (Lande & Arnold 1983). Leaf area was not included in either the top

binomial or Poisson models and therefore we do not include any selection gradient values

for leaf area. We recognize that selection gradient values could be due to selection on

unmeasured correlated characters, although because we used a segregating hybrid

population there should be fewer genetically correlated phenotypes.

DISCUSSION

Since the pioneering reciprocal transplant experiments of Clausen, Keck, and Hiesey

(1941) differential habitat adaptation has been repeatedly demonstrated in plants

(Hereford 2009). However the selective forces and traits driving differential adaptation

are still poorly understood (Salvolainen et al. 2013). This is especially true of

differentially adapted species. When closely related species are sympatric adaptation to

different environments can increase reproductive isolation between them (Coyne & Orr

2004) and promote species coexistence (Wright 2002). To test the adaptive significance

of early flowering time, small flower size, small plant size, and lobed leaf shape in *M.*

laciniatus' marginal rocky outcrop habitat we performed a reciprocal transplant with *M.*

laciniatus, M. guttatus, and outbred F_4's. We collected fine scale soil moisture and

herbivory data, measured fitness of the parental species, and selection on phenotypes in

F_4's in each habitat. Drought and herbivory are differentially associated with survival in

each habitat making them likely agents of divergent selection. *Mimulus laciniatus* is

better adapted to granite than its close relative and progenitor *M. guttatus,* and this

adaptation is underlain by differential selection on flowering time, plant size, and leaf

shape between habitats.

Soil moisture & herbivory are likely selective agents

Determining the selective forces that drive adaptation to different in environments is an

important, but challenging component of the study of adaptive evolution (Wade & Kalisz

1990). It is a challenge because many environmental variables differ between any two

habitats and any number of them may affect fitness (MacColl 2011). To test the

significance of a particular ecological variable it is necessary to isolate the impact of that

variable on fitness either through (1) environmental manipulation (Bright 1998; Stanton

et al. 2000; Weijshede et al. 2008; Campitelli & Stinchcombe 2013a), or (2) through fine

scale environmental measurements within and between habitat types (Huber et al. 2004;

Garant et al. 2007; Lind & Johansson 2007; Quinn et al. 2009; Weese et al. 2010). We

have assessed the selective environments of granite and meadow habitats through the

latter approach. We measured soil moisture and herbivory, which we had prior reason to

believe were selective factors in these habitats (Peterson et al. 2013, Ferris et al. 2014, K.

Ferris personal observation), on a fine spatial scale throughout each reciprocal transplant

site so that we could determine whether variation in these factors was associated with

variation in survival within each habitat type. We also measured soil moisture on a fine

temporal scale. Then we examined whether there was a difference in soil moisture or

herbivory's association with fitness between habitat type.

Soil moisture availability is one of the main drivers of plant distribution and

diversification (Stebbins 1952; Axelrod 1972). From our environmental analysis we

learned that decreasing soil moisture was associated with decreasing survival on a fine

spatial and temporal scale across all sites in our top linear mixed effects model (Table 1;

Figure 2). However, a significant interaction between soil moisture, time, and habitat type

demonstrated that soil moisture decayed and interacted with survival differently in

granite outcrops vs. meadows (Table 1; Figure 2). In *M. guttatus'* meadows soil moisture

decayed in a slow linear fashion (Figure 2c&d), while in *M. laciniatus'* granite outcrops

soil moisture remained high and constant early in the season and then dropped

precipitously mid-summer (Figure 2a&b). Different soil moisture regimes should select

for different plant life histories. A fast cycling, drought escape strategy should be

advantageous in the fast-drying granite outcrops while a longer lived, more water-use-

efficient strategy should be advantageous in meadow habitat (Dudley 1996; Stanton et al.

2000; McKay et al. 2003; Anderson et al. 2011). The granite habitat's water regime

seems to have selected for *M. laciniatus'* rapid, early flowering life history.

We also found a difference in herbivory pressure between habitats with herbivory

impacting survival on a fine spatial scale in *M. guttatus'* meadows, but not in granite

outcrops (Table 1). There was also significantly less total herbivore damage in granite

(Figure 3). Therefore it seems likely that herbivory acts as an important selective force in

M. guttatus' meadows, but not in *M. laciniatus'* granite outcrops. This is supported by the

author's observation of greater insect density in mesic meadows than in granite outcrops

(K. Ferris personal observation). Perhaps *M. laciniatus* is not well adapted to herbivory

which could contribute to the fecundity trade-off between *M. laciniatus* & *M. guttatus* in

the meadow habitat. Herbivory is believed to be a major selective force driving plant

diversification through coevolutionary forces (Ehrlich and Raven 1964; Janz 2011). In

summary our data show that differences in soil moisture regime and herbivory between

granite outcrops and nearby meadows likely contribute to divergent selection between *M.*

laciniatus & *M. guttatus'* habitats.

Differential phenotypic plasticity between species

There is much debate in the literature about the role of phenotypic plasticity in adaptive

evolution (Via et al. 1995; Schmitt et al. 1995; Dudley & Schmitt 1996; Ghalambor et al.

2007; Ghalambor et al. 2015). We discovered both phenotypic plasticity and genetic

variation for plasticity (GxE) in flowering time, plant height, and flower size (Figure 5) in

our reciprocal transplant. The presence of GxE in height, flower size, and flowering time

indicates that phenotypic plasticity has the potential to respond to selection. Plasticity

was in the expected direction, meaning in the same direction as species differences, for

flowering time, plant height, and leaf shape. Plants in all genotypic classes (F_4, *M.*

laciniatus, *M. guttatus*) flowered earlier and were shorter in granite, and both *M.*

laciniatus and F_4's were more highly lobed on average in granite than in meadow habitat

(Table 2). *Mimulus laciniatus* had significantly greater flowering time plasticity than *M.*

guttatus (Figure 5a). This ability to flower more readily in response to environmental

cues may be advantageous given the rapid decreases in soil moisture and survival that

characterize *M. laciniatus'* native granite habitat (Figure 2a&b). On the other hand, the

slow decline in soil moisture throughout the season in *M. guttatus'* meadows would not

require plants to flower as rapidly in response to drought (Figure 2 c&d). Therefore M.

laciniatus's granite environment might select for increased flowering time plasticity

while *M. guttatus'* meadows may not. Flowering time plasticity has been shown to be

adaptive in other plant systems such as *Impatiens campensis* (Donohue et al. 2000;

Donohue et al. 2001).

Mimulus guttatus, on the other hand, has greater environmentally induced

plasticity in plant height than *M. laciniatus*. In a competitive environment like *M.*

guttatus' meadows a plant's ability to plastically change its height could be advantageous

as a shade avoidance mechanism (Dudley & Schmitt 1996, Aphalo et al. 1999, Donohue

et al. 2000, Weinig 2000). In a densely populated environment taller plants are able to

outcompete other individuals for access to light and will therefore have higher

photosynthetic rates and fitness (Schmitt et al. 1987, Aphalo et al. 1999). The

significantly smaller size of *M. laciniatus* plants and leaves (Figure 5c&d) in meadows

compared to the native *M. guttatus* suggests that *M. laciniatus* struggled to compete there.

In the granite outcrop habitat *M. laciniatus* was only slightly smaller in stature and leaf

size than *M. guttatus*. There was also selection for increased plant height in the meadow

habitat, but decreased height in granite (Table 5). *Mimulus guttatus'* meadows have a

much higher plant density than *M. laciniatus'* relatively depauperate granite outcrops (K.

Ferris, personal observation), and small plants would be shaded out by other more

vigorous species. To empirically test whether there is selection for increased flowering

time plasticity in *M. laciniatus'* granite outcrops or height plasticity in *M. guttatus'*

meadows further experiments with replicate hybrid genotypes planted in each habitat

would be necessary. Here we merely hypothesize that these differences in plasticity are

advantageous given the known characteristics of each species' habitat.

Another piece of evidence suggesting that *M. laciniatus* was at a competitive

disadvantage in meadows was the direction of flower size plasticity between habitats.

While *M. guttatus* displayed phenotypic plasticity in the expected direction, that is it had

larger flowers in its native meadows than in the granite outcrops, *M. laciniatus* had larger

flowers in granite outcrops than in meadows (Figure 5b). In our truncated Poisson

distributed model larger flowers were selected for in both habitats, and consequently the

decrease in *M. laciniatus'* flower size in meadows relative to granite suggests that *M.*

laciniatus was at a competitive disadvantage in *M. guttatus'* habitat. Therefore

competition with other species for light and resources could explain *M. laciniatus'* poor

fecundity in *M. guttatus'* habitat and the species trade-off in lifetime fitness.

Flowering time, plant size, and leaf shape contribute to differential habitat adaptation

We found that *M. laciniatus* and *M. guttatus* are differentially adapted to their respective

habitats (Figure 1). *M. laciniatus* parents were significantly more likely to survive to

flowering and produce more fruits in granite outcrops than *M. guttatus* parents (Table 1,

Figure 4a). Interestingly in meadows we saw no difference in survival to flowering

between species, but *M. guttatus* did produce more fruits than *M. laciniatus* in its

meadow habitat (Figure 4b). A previous study also found that *M. laciniatus* had higher

survival to flowering in granite than *M. guttatus*, but that the two survived equally well in

meadows (Peterson et al. 2013). Many reciprocal transplant studies have found that one

species or population does best in both habitats (Hereford et al. 2009) and this

observation elicits the question, why doesn't the population with highest overall fitness

occur in both habitats? One potential reason for not observing a fitness trade-off in a

reciprocal transplant study is that lifetime fitness was not measured. In this study we

found an asymmetric result without a clear trade-off between species when we examined

survival, but then found evidence of a species trade-off when we measured fecundity.

The fact that we find selection against the non-native parent in each habitat indicates that

there is ecological reproductive isolation between these species (Coyne & Orr 2004).

Mimulus laciniatus occurs within the geographic range of *M. guttatus* (Ferris et al. 2014)

and the two often co-occur within meters of each other creating the opportunity for gene

flow (Ferris et al. 2016). Due to the lack of obvious intrinsic barriers between these

species (K. Ferris personal observation) illustrated by the ease of crossing, this ecological

component of reproductive isolation may be important for maintaining species

boundaries in this system.

Our selection analysis found that earlier flowering time, small plant size, and

lobed leaves underlie *M. laciniatus'* adaptation to its harsh granite outcrop habitat (Tables

4, 5, 6). Earlier flowering time is critical for drought escape in many annual plants that

occupy seasonally dry environments (Kiang & Hamrick 1978; Fox 1990; Stanton et al.

2000; Mckay et al. 2003; Eckhart et al. 2004; Hall & Willis 2006; Franks et al. 2007;

Anderson et al. 2012). This is because early flowering allows plants to reproduce before

the rapid onset of seasonal drought in harsh, ephemeral habitats. Work with the

herbaceous annual *Impatiens campensis* has found that early season drought selects for a

rapid flowering, drought escape strategy, while late season drought favors slower

developing genotypes with high WUE (Heschel & Reginos 2005). In continually moist

environments we expect selection for later flowering or stabilizing selection on

midseason flowering since later flowering plants are larger and more fecund (Mitchell-

Olds 1996; Hall & Willis 2006; Anderson et al. 2012). *Mimulus laciniatus'* granite

outcrop habitat dries out a month earlier in the growing season than nearby meadows

where *M. guttatus* occurs (Figure 2; Peterson et al. 2013). In our reciprocal transplant

earlier flowering significantly increased both the probability of a plant setting fruit and

the number of fruits a plant was able to produce, in the granite outcrop habitat. Although

we found evidence that selection acts more strongly on flowering time in granite outcrops

than meadows (Tables 2&3), we did not detect selection for later flowering in the *M.*

guttatus habitat. This may be because meadows dried out earlier than normal due to the

2013 California drought (Swain et al., 2014). Selection for earlier flowering has

previously been detected in inland annual populations of *M. guttatus* (Hall & Willis

2006) indicating that this may be common in populations of this species with an annual

life history.

We also detected differential selection on plant height between granite and

meadow habitats. Plant size is intimately tied to plant fitness (Stearns 1992; Thornsberry

et al. 2001; Falster & Westoby 2003). Larger plants are more fecund, but usually

reproduce later in the growing season (Mitchell-Olds 1992; Mitchell-Olds 1996). In a

seasonally dry environment later flowering places larger plants at risk of dying before

reproduction. However, in a continually moist and densely populated environment larger

plants have a competitive advantage both in terms of access to resources such as light and

water and in increased fecundity due to their increased biomass (Mitchell-Olds 1992;

Dudley & Schmitt 1996). In our reciprocal transplant we saw strong selection for taller

plants in *M. guttatus's* mesic meadow habitat both in terms of whether a plant produced

fruit and the number of fruits it produced. In the seasonally dry granite habitat we also

found that taller plants produced more fruit, but this selection was not as strong as in the

meadow habitat. Importantly, we also found that smaller plants were significantly more

likely to produce fruit in *M. laciniatus'* granite habitat. Therefore we saw differential

selection on plant height at first flower between species' habitats in the direction of

species' differences. This indicates that plant height is an important component of

differential habitat adaptation between *M. laciniatus* and *M. guttatus*.

Lobed leaves have been proposed to be adaptive in dry, marginal habitats because

of their ability to be efficiently cooled by convection (Vogel 1968; Givnish 1978;

Schuepp 1993; Nobel 2005) and because they have less drought stress prone tissue than

entire leaves (Nicotra et al. 2011). However, it has been difficult to definitively

demonstrate the adaptive significance of lobed leaf shape (Nicotra et al. 2011). Nicotra et

al. (2008) demonstrated that lobed species of the South African genus *Pelargonium* have

higher photosynthetic rates under high temperatures than their close entire-leaved

relatives which they hypothesize is a strategy to maximize growth when water is

available in seasonally dry habitats. However many traits are correlated when species

diverge so it is difficult to say whether photosynthetic rates were a direct result of leaf

shape or of evolutionary history. In the ivy leaf morning glory, *Ipomea hederaceae,* leaf

shape has been shown to be under selection along a North-South cline (Campitelli &

Stinchcombe 2013b) and in a segregating experimental field population (Bright &

Rausher 2008). Northern populations of *I. hederaceae* are lobed, while Southern

populations are predominantly entire-leaved (Campitelli & Stinchcombe 2013b). Our

experiment is the first to directly test whether lobed leaf shape is under selection in a

segregating inter-specific hybrid population. In the *M. guttatus* species complex lobed

leaf shape is associated with the occupation of rocky, dry habitat (Ferris et al. 2015).

Within the entire genus *Mimulus,* which consists of over 150 species, only two species,

M. laciniatus and *M. filicifolius*, are fixed for lobed leaves and both specialize on granite

outcrops (Ferris et al. 2014). Therefore we hypothesized that lobed leaf shape is an

adaptation to this specific habitat in *Mimulus*. We are able to confirm this prediction in

our reciprocal transplant where we found that lobed leaf shape was under positive

selection in our F_4 hybrid's in granite, but not in meadow habitat (Tables 5&6).

We detected positive directional selection on flower size in both habitats (Table 6)

in our Poisson regression, indicating this trait does not contribute to differential habitat

adaptation between *M. guttatus* and *M. laciniatus* at this life history stage. . In a study of

M. guttatus inland annuals vs. coastal perennials, Hall and Willis (2006) also detected a

pattern of selection for larger flowers across habitats. Increased flower size likely

facilitates increased outcrossing via pollinator visitation, which would alleviate effects of

inbreeding depression in both habitats. However, we did see evidence of differential

selection on flower size between habitats in our binomial regression on whether a trait

increased the probability of a plant producing fruit. The result was in the opposite

direction of species differences since we observed selection for smaller flowers in the

meadow habitat even though *M. guttatus* has larger flowers, while in granite larger

flowers were more likely to fruit. Instead of being adaptive, the difference in floral size

between the parental species may be due to selection on a correlated trait such as whole

plant size or flowering time.

Conclusions

Understanding the ecological variables and traits driving differential habitat adaptation is

a key component of ecology and evolutionary biology, and is particularly relevant in light

of the future increases in temperature and precipitation stress predicted to accompany

global climate change. Adaptation is also critical for the generation and maintenance of

biodiversity since differential adaptation between closely related sympatric species is an

important component of reproductive isolation. We found that *M. laciniatus* and its

supposed progenitor *M. guttatus* are differentially adapted to microhabitats that differ

dramatically in drought regime and herbivory. Our results indicate that while early

flowering time, rapid development, and lobed leaves are critical for plant fitness in *M.*

laciniatus' harsh seasonally dry environment, traits such as herbivore resistance and large

plant size may be more important in a competitive mesic environment like *M. guttatus'*.

Our study is one of the few to find selection on lobed leaves, and the first to demonstrate

that lobed leaves are adaptive in a seasonally dry environment. Additionally we find

genetic variation for phenotypic plasticity between species in all traits indicating that

plasticity can respond to selection in this system and may be a part of adaptation to each

species habitat. By combining fine scale environmental surveys, large-scale reciprocal

transplants, and phenotypic selection analysis we have reached a greater understanding of

adaptation to a harsh marginal environment.

ACKNOWLEDGEMENTS

We would like to thank Alex Gunderson for giving a month of his life to help the author

drive cross-country and plant 3800 seedlings. The experiment could not have succeeded

without his heroic efforts. We also thank Jenn Coughlan, Kathy Toll, and Annie Jeong

for assistance collecting environmental data and censusing plants. Yosemite National

Park provided permits and the Sierra Nevada Research Institute provided housing at the

Yosemite Field Station. Funding was provided by an NSF DDIG (DEB-1210755), a

California Native Plant Society Educational Grant, and a Sigma Xi Grants-in-Aid of

Research awarded to Kathleen Ferris and John Willis, and by NSF En-Gen (EF-0723814)

and LiT (IOS-1024966) grants awarded to John Willis.

AUTHOR CONTRIBUTIONS

KF designed & performed the controlled crossing and field experiments, collected and

analyzed all data, and wrote the first draft of the manuscript. JW helped with

experimental design and provided feedback on the manuscript.

REFERENCES

Ågren, J., C. G. Oakley, S. Lundemo, & D. W. Schemske. 2016. Adaptive divergence in
flowering time among natural populations of Arabidopsis thaliana: Estimates of selection
and QTL mapping. *Evolution* 71:550-564.
652
653 Anderson, J. T., D. W. Inouye, A.M. McKinney, R.I. Coulatti, T. Mitchell-Olds. 2012.
654 Phenotypic plasticity and adaptive evolution contribute to advancing flowering
655 phenology in response to climate change. *Proc. R. Soc. B* 279:3843-3852.
656
657 Anderson, J. T., C. R. Lee, T. Mitchell-Olds. 2011. Life history QTLs and natural
658 selection on flowering time in *Boechera stricta*, a wild relative of
659 *Arabidopsis*. *Evolution* 65:771-787.
660
661 Anderson, J.T., J. H. Willis, T. Mitchell-Olds. 2011. Evolutionary genetics of plant
662 adaptation. *Trends in Genetics* 27:258-266.
663
664 Anderson, J. T., V. M. Eckhart, & M. A. Geber. 2015. Experimental studies of adaptation
665 in Clarkia xantiana. III. Phenotypic selection across a subspecies border. *Evolution*,
666 69:2249-2261.
667
668 Aphalo, P. J., C. L. Ballare, & , A. L. Scopel. 1999. Plant-plant signalling, the shade-
669 avoidance response and competition. *Journal of Experimental Botany*, 50:1629-1634.
670
671 Axelrod, D. I. 1972. Edaphic aridity as a factor in angiosperm evolution. *The American*
672 *Naturalist*, 106:311-320.
673
674 Bartoń, K. 2013. MuMIn: multi-model inference, R package version 1.9.13
675
676 Blum, A. & C. Y. Sullivan. 1997. The effect of plant size on wheat response to agents of
677 drought stress. I. Root drying. *Functional Plant Biology*, 24:35-41.
678
679 Blum, A., C. Y. Sullivan, & H. Nguyen. 1997. The effect of plant size on wheat response
680 to agents of drought stress. II. Water deficit, heat and ABA. *Functional Plant Biology*,
681 24:43-48.
682
683 Bolmgren, K., & P. D. Cowan. 2008. Time–size tradeoffs: a phylogenetic comparative
684 study of flowering time, plant height and seed mass in a north☐temperate flora. *Oikos*,
685 117:424-429.
686
687 Bright, K. L. 1998. Geographic variation and natural selection on a leaf shape
688 polymorphism in the ivyleaf morning glory (Ipomoea hederacea). Duke University.
689
690 Bright, K. L., M. D. Rausher. 2008. Natural Selection on a leaf-shape polymorphism in
691 the Ivyleaf Morning Glory (*Ipomoea heredacea*). *Evolution* 62:1978-1990.
692

693 Campitelli, B. E., & J. R. Stinchcombe. 2013a. Testing potential selective agents acting
694 on leaf shape in Ipomoea hederacea: predictions based on an adaptive leaf shape cline.
695 *Ecology and Evolution*, 3:2409-2423.
696
697 Campitelli, B. E., & J. R. Stinchcombe. 2013b. Natural selection maintains a
698 single☐locus leaf shape cline in Ivyleaf morning glory, Ipomoea hederacea. *Molecular*
699 *ecology*, 22:552-564.
700
701 Clausen, J., D. D. Keck, W. M. Hiesey. 1941. Regional differentiation in plant species.
702 *The American Naturalist*, 75:231-250.
703
704 Coyne, J. A., and A. H. Orr. 2004. *Speciation*. Sinauer Associates, Inc. Sunderland, MA.
705
706 Cunningham, S. A. 2000. Depressed pollination in habitat fragments causes low fruit set.
707 *Proceedings of the Royal Society of London B: Biological Sciences* 267:1149-1152.
708
709 Des Marais, D. L., K. M. Hernandez, T.E. Juenger. 2013. Genotype by environment
710 interaction and plasticity: exploring genomic responses of plants to the abiotic
711 environment. *Annu. Rev. Ecol. Evol. Syst.,* 44:5-29.
712
713 Donohue, K., D. Messiqua, E. Hammond Pyle, M. S. Heschel, & J. Schmitt. 2000.
714 Evidence of adaptive divergence in plasticity: density-and site-dependent selection on
715 shade-avoidance responses in Impatiens capensis. *Evolution*, 54:1956-1968.
716
717 Donohue, K., E. Hammond Pyle, D. Messiqua, M. S. Heschel, & J. Schmitt. 2001.
718 Adaptive divergence in plasticity in natural populations of Impatiens capensis and its
719 consequences for performance in novel habitats. *Evolution*, 55:692-702.
720
721 Dudley SA. 1996. Differing selection on plant physiological traits in response to
722 environmental water availability: a test of adaptive hypotheses. *Evolution*, 50:92–102.
723
724 Dudley, S. A., J. Schmitt. 1996. Testing the adaptive plasticity hypothesis: density-
725 dependent selection on manipulated stem length in Impatiens capensis. *American*
726 *Naturalist*, 445-465.
727
728 Ehrlich, P. R., & P. H. Raven. 1964. Butterflies and plants: a study in coevolution.
729 *Evolution*, 18:586-608.
730
731 Falster, D. S., & M. Westoby. 2003. Plant height and evolutionary games. *Trends in*
732 *Ecology & Evolution*, 18:337-343.
733
734 Fausto, J. A., V. M. Eckhart, M. A. Geber. 2001. Reproductive assurance and the
735 evolutionary ecology of self-pollination in Clarkia xantiana (Onagraceae). *American*
736 *Journal of Botany* 88:1794-1800.
737

738 Fenster, C. B., K. Ritland. 1994. Evidence for natural selection on mating system in
739 Mimulus (Scrophulariaceae). *International journal of plant sciences*, 588-596.
740
741 Ferris, K.G., L. B. Baldridge, B. K. Blackman, and J. H. Willis. 2016. The Genetic
742 Architecture of Local Adaptation and Reproductive Isolation under Gene Flow in the
743 *Mimulus guttatus* Species Complex. *Molecular Ecology* DOI: *10.1111/mec.13763*
744
745 Ferris, K. G., T. Rushton, A. B. Greenlee, K. Toll, B. K. Blackman, J. H. Willis. 2015.
746 Leaf shape evolution has a similar genetic architecture in three edaphic specialists within
747 the Mimulus guttatus species complex. *Annals of botany* 116:213-223.
748
749 Ferris, K. G., J. P. Sexton, J. H. Willis. 2014. Speciation on a local geographic scale: the
750 evolution of a rare rock outcrop specialist in Mimulus. *Phil. Trans. of the Royal Soc. B*,
751 369:20140001.
752
753 Fishman L, Aagaard J, Tuthill JC. 2008. Toward the evolutionary genomics of
754 gametophytic divergence: patterns of transmission ratio distortion in monkeyflower
755 (Mimulus) hybrids reveal a complex genetic basis for conspecific pollen precedence.
756 *Evolution,* 62: 2958–2970.
757
758 Fox, G. A. 1990. Drought and the evolution of flowering time in desert annuals.
759 *American Journal of Botany,* 77:1508-1518.
760
761 Franks, S. J., S. Sim, A. E. Weis. 2007. Rapid evolution of flowering time by an annual
762 plant in response to a climate fluctuation. *Proceedings of the National Academy of*
763 *Sciences*, 104:1278-1282.
764
765 Friedman, J., J. H. Willis. 2013. Major QTL's for critical photoperiod and vernalization
766 underlie extensive variation in flowering in the *Mimulus guttatus* species complex. *New*
767 *Phytologist* 199:571-583.
768
769 Galen, C. 1999. Are flowers physiological sinks or faucets? Costs and correlates of water
770 use by flowers of Polemonium viscosum. *Oecologia* 118:461-470.
771
772 Garant, D., L. E. Kruuk, R. H. McCleery, & B. C. Sheldon. 2007. The effects of
773 environmental heterogeneity on multivariate selection on reproductive traits in female
774 great tits. *Evolution*, 61:1546-1559.
775
776 Ghalambor, C. K., J. K. McKay, S. P. Carroll, D. N. Reznick. 2007. Adaptive versus
777 non☐adaptive phenotypic plasticity and the potential for contemporary adaptation in new
778 environments. *Functional Ecology* 21:394-407.
779
780 Ghalambor, C. K., K. L. Hoke, E. W. Ruell, E. K. Fischer, D. N. Reznick, K. A. Hughes.
781 2015. Non-adaptive plasticity potentiates rapid adaptive evolution of gene expression in
782 nature. *Nature* 525:372-375.
783

784 Givnish, T. J. 1978. Ecological aspects of plant morphology: Leaf form in relation to
785 environment. pp. 83-142 in R. Sattler (ed.), Theoretical Plant Morphology. Leiden
786 University Press, Leiden.
787
788 Hall, M. C., J. H. Willis. 2006. Divergent selection on flowering time contributes to local
789 adaptation in Mimulus guttatus populations. Evolution 60:2466-2477.
790
791 Hereford, J. 2009. A quantitative survey of local adaptation and fitness tradeoffs. *The*
792 *American Naturalist* 173:579-588.
793
794 Heschel, M. S., & C. Riginos. 2005. Mechanisms of selection for drought stress tolerance
795 and avoidance in Impatiens capensis (Balsaminaceae). *American Journal of Botany*, 92:
796 37-44.
797
798 Huber, H., N. C. Kane, M. S. Heschel, E. J. von Wettberg, J. Banta, A. M. Leuck, & J.
799 Schmitt. 2004. Frequency and microenvironmental pattern of selection on plastic shade-
800 avoidance traits in a natural population of Impatiens capensis. *The American Naturalist*,
801 163:548-563.
802
803 Janz, N. 2011. Ehrlich and Raven revisited: mechanisms underlying codiversification of
804 plants and enemies. *Annual Review of Ecology, Evolution, and Systematics*, 42:71-89.
805
806 Kalisz, S. 1986. Variable Selection on the timing of germination in Collinsia verna
807 (Scrophulariaceae). *Evolution,* 40:479-491.
808
809 Kawecki, T. J., D. Ebert. 2004. Conceptual Issues in Local Adaptation. *Ecology Letters,*
810 7:1225-1241.
811
812 Kiang, Y. T., J. L. Hamrick. 1978. Reproductive isolation in the M. guttatus–M. nasutus
813 complex. *American Midland Naturalist* 100:269–276.
814
815 Kingsolver, J. G., H. E. Hoekstra, J. M. Hoekstra, D. Berrigan, S. N. Vignieri, C. E.
816 Hill,... & P. Beerli. 2001. The strength of phenotypic selection in natural populations. *The*
817 *American Naturalist*, 157:245-261.
818
819 Lande, R., S. J. Arnold. 1983. The measurement of selection on correlated characters.
820 *Evolution* 37:1210-1226.
821
822 Lind, M. I. & F. Johansson. 2007. The degree of adaptive phenotypic plasticity is
823 correlated with the spatial environmental heterogeneity experienced by island populations
824 of Rana temporaria. *Journal of evolutionary biology*, 20:1288-1297.
825
826 MacColl, A. D. C. 2011. The ecological causes of evolution. *Trends Ecol. Evol.* **26**:514–
827 522.
828

829   Macnair, M. R., M. Gardner. 1998. The Evolution of Edaphic Endemics. *Endless Forms:*
830   *Species and Speciation.* edited by D.J. Howard and S.H. Berlocher. Oxford University
831   Press Inc., Oxford.
832
833   Mazer, S. J., L. S. Dudley, A. A. Hove, S. K. Emms, A. S. Verhoeven. 2010.
834   Physiological performance in Clarkia sister taxa with contrasting mating systems: do
835   early-flowering autogamous taxa avoid water stress relative to their pollinator-dependent
836   counterparts? *International Journal of Plant Sciences* 171:1029–1047.
837
838   Mayr, E. 1947. Ecological Factors in Speciation. *Evolution* 1:263-288.
839
840   Mayr, E. 1949. Speciation and Selection. *Proceedings of the American Philosophical*
841   *Society* 93:514-519.
842
843   Mckay, J. K., J. H. Richards, & T. Mitchell☐Olds. 2003. Genetics of drought adaptation
844   in Arabidopsis thaliana: I. Pleiotropy contributes to genetic correlations among ecological
845   traits. *Molecular Ecology*, 12:1137-1151.
846
847   Messmer, R., Y. Fracheboud, M. Bänziger, M. Vargas, P. Stamp, J. M. Ribaut. 2009.
848   Drought stress and tropical maize: QTL-by-environment interactions and stability of
849   QTLs across environments for yield components and secondary traits. *Theoretical and*
850   *Applied Genetics* 119:913-930.
851
852   Mitchell-Olds, T. 1992. Does environmental variation maintain genetic variation? A
853   question of scale. *Trends in ecology & evolution*, 7:397-398.
854
855   Mitchell-Olds, T. 1996. Genetic Constraints on Life-History Evolution: Quantitative-
856   Trait Loci Influencing Growth and Flowering in Arabidopsis thaliana. *Evolution,* 50:140-
857   145.
858
859   Mitchell-Olds, T., R.G. Shaw. 1987. Regression analysis of natural selection: statistical
860   inference and biological interpretation. *Evolution* 41:1149-1161.
861
862   Mizukami, Y. 2001. A matter of size: developmental control of organ size in plants.
863   *Current opinion in plant biology* 4:533-539.
864
865   Mojica, J. P., Y. W. Lee, J. H. Willis, J. K. Kelly. 2012. Spatially and temporally varying
866   selection on intrapopulation quantitative trait loci for a life history trade-off in Mimulus
867   guttatus. *Molecular ecology* 21:3718-3728.
868
869   Nicotra, A. B., M. J.Cosgrove, A. Cowling, C. D. Schlichting, & C. S. Jones. 2008. Leaf
870   shape linked to photosynthetic rates and temperature optima in South African
871   Pelargonium species. *Oecologia* 154:625-635.
872

873   Nicotra, A. B., A. Leigh, C. K. Boyce, C. S. Jones, K. J. Niklas, D. L. Royer, H. Tsukaya.
874   2011. The evolution and functional significance of leaf shape in the angiosperms.
875   *Functional Plant Biology* 38:535-552.
876
877   Niklas, K. J. 1994. Plant Allometry: The Scaling of Form and Process, The University of
878   Chicago Press, Chicago
879
880   Nobel, P. S. 2005. Physicochemical and Environmental Plant Physiology. Elsevier
881   Academic Press. Burlington, MA.
882
883   Pantuwan, G., S. Fukai, M. Cooper, S. Rajatasereekul, & J. C. O'toole. 2002. Yield
884   response of rice (Oryza sativa L.) genotypes to drought under rainfed lowland: 3. Plant
885   factors contributing to drought resistance. *Field Crops Research*, 73:181-200.
886
887   Parkhurst, D. F., P. R. Duncan, D. M. Gates, F. Kreith. 1968. Wind-tunnel modelling of
888   convection of heat between air and broad leaves of plants. *Agricultural Meteorology*
889   5:33-47.
890
891   Paterson, A. H., S. Damon, J. D. Hewitt, D. Zamir, H. D. Rabinowitch, S. E. Lincoln, E.
892   S. Lander, S. D. Tanksley. 1991. Mendelian factors underlying quantitative traits in
893   tomato: comparison across species, generations, and environments. *Genetics* 127:181-
894   197.
895
896   Peterson, M. L., K. J. Rice, J. P. Sexton. 2013. Niche partitioning between close relatives
897   suggests trade☐offs between adaptation to local environments and competition. *Ecology*
898   *and evolution* 3:512-522.
899
900   Piper, J. G., B. Charlesworth, D. Charlesworth. 1986. Breeding system evolution in
901   Primula vulgaris and the role of reproductive assurance. *Heredity* 56:207-217.
902
903   Quinn, J. L., S. C. Patrick, S. Bouwhuis, T. A. Wilkin, & B. C. Sheldon. 2009.
904   Heterogeneous selection on a heritable temperament trait in a variable environment.
905   *Journal of Animal Ecology*, 78:1203-1215.
906
907   R Core Development Team. 2008. R: A language and environment for statistical
908   computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-
909   0, http://www.R-project.org.
910
911   Rausher, M. D. 1983. The evolution of habitat preference in subdivided populations.
912   *Evolution* 38:596-608.
913
914   Ridout, M., C. G. B. Demetrio, J. Hinde. 1998. Models for count data with many zeros.
915   *International Biometric Conference*, Cape Town, SA.
916

917 Robichaux, R. H., G. D. Carr, M. Liebman, R. W. Pearcy. 1990. Adaptive radiation of the
918 Hawaiian Silversword Alliance (Compositae- Madiiane): Ecological, Morphological, and
919 Physiological Diversity. *Annals of the Missouri Botanical Garden* 77:64-72.
920
921 Rundle, H., and P. Nosil. 2005. Ecological specaition. *Ecology Letters* 8:336-352.
922
923 Savolainen, O., M. Lascoux, & J. Merilä. 2013. Ecological genomics of local adaptation.
924 *Nature Reviews Genetics* 14:807-820.
925
926 Schluter, D. 2001. Ecology and the origin of the species. *Trends in Ecology and*
927 *Evolution* 16:372-380.
928
929 Schmitt, J., J. Eccleston, & D. W. Ehrhardt. 1987. Dominance and suppression, size-
930 dependent growth and self-thinning in a natural Impatiens capensis population. *The*
931 *Journal of Ecology*, 75:651-665.
932
933 Schmitt, J., A. C. McCormac, H. Smith. 1995. A test of the adaptive plasticity hypothesis
934 using transgenic and mutant plants disabled in phytochrome-mediated elongation
935 responses to neighbors. *American Naturalist*, 937-953.
936
937 Schmitt, J., and J. Antonovics. 1986. Experimental Studies of the Evolutionary
938 Significance of Sexual Reproduction III. Maternal and Paternal Effects During Seedling
939 Establishment. *Evolution* 40:817-829.
940
941 Schuepp, P. H. 1993. Tansley review no.59: leaf boundary layers. *New Phytologist*
942 125:477-507.
943
944 Skaug, H., D. Fournier, and A. Nielsen. 2006. glmmADMB: generalized linear mixed
945 models using AD Model Builder. R package version 0.3.
946
947 Stanton, M. L., B. A. Roy, & D. A. Thiede. 2000. Evolution in stressful environments. I.
948 Phenotypic variability, phenotypic selection, and response to selection in five distinct
949 environmental stresses. *Evolution*, 54:93-111.
950
951 Stearns, S. C. 1992. *The evolution of life histories.* Vol. 249. Oxford: Oxford University
952 Press.
953
954 Stebbins, G. L. 1952. Aridity as a stimulus to plant evolution. *The American Naturalist*,
955 86:33-44.
956
957 Stebbins, G. L. 1957. Self fertilization and population variability in the higher plants. *The*
958 *American Naturalist*, 91:337-354.
959
960 Stebbins, G. L. 1970. Adaptive radiation of reproductive characteristics in angiosperms,
961 I: pollination mechanisms. *Annual Review of Ecology and Systematics* 307-326.
962

963 Stewart, S. C., and D. J. Schoen. 1987. Pattern of phenotypic viability and fecundity
964 selection in a natural population of *Impatiens pallida*. *Evolution* 41:1290-1301.
965

966 Swain, D. L., M. Tsiang, M. Haugen, D. Singh, A. Charland, B. Rajaratnam, N. S.
967 Diffenbaugh. 2014. The extraordinary California drought of 2013/2014: Character,
968 context, and the role of climate change. *Bulletin of the American Meteorological Society*
969 95:S3.
970

971 Thornsberry, J. M., M. M. Goodman, J. Doebley, S. Kresovich, D. Nielsen, & E. S.
972 Buckler. 2001. Dwarf8 polymorphisms associate with variation in flowering time. *Nature*
973 *genetics*, 28:286-289.
974

975 Via, S., R. Gomulkiewicz, G. De Jong, S. M. Scheiner, C. D. Schlichting, P. H. Van
976 Tienderen. 1995. Adaptive phenotypic plasticity: consensus and controversy. *Trends in*
977 *Ecology & Evolution* 10:212-217.
978

979 Vogel, S. 1968. 'Sun leaves' and 'shade leaves': differences in convective heat
980 dissipation. *Ecology* 49:1203-1204.
981

982 Wade, M. J., and S. Kalisz. 1990. The causes of natural selection. *Evolution* **44**:1947–
983 1955.
984

985 Wan, X. Y., J. M. Wan, J. F. Weng, L. Jiang, J. C. Bi, C. M. Wang, H. Q. Zhai. 2005.
986 Stability of QTLs for rice grain dimension and endosperm chalkiness characteristics
987 across eight environments. *Theoretical and Applied Genetics* 110:1334-1346.
988

989 Weese, D. J., S. P. Gordon, A. P. Hendry, & M. T. Kinnison. 2010. Spatiotemporal
990 variation in linear natural selection on body color in wild guppies (Poecilia reticulata).
991 *Evolution*, 64:1802-1815.
992

993 Wei, X., J. Xu, H. Guo, L. Jiang, S. Chen, C. Yu, Z. Zhou, P. Hu, H. Zhai, and J. Wan.
994 2010. DTH8 Suppresses Flowering in Rice, Influencing Plant Height and Yield Potential
995 Simultaneously 1[W][OA]. Plant Physiol. 153:1747–1758.
996

997 Weijschedé, J., R. Berentsen, H. de Kroon, & H. Huber. 2008. Variation in petiole and
998 internode length affects plant performance in Trifolium repens under opposing selection
999 regimes. *Evolutionary Ecology*, 22:383-397.
1000

1001 Weinig, C. 2000. Plasticity versus canalization: population differences in the timing of
1002 shade-avoidance responses. *Evolution*, 54:441-451.
1003

1004 Willis, J. H. 1993a. Effects of different levels of inbreeding on fitness components in
1005 *Mimulus guttatus. Evolution* 47:864-876.
1006

1007 Willis, C. G., B. Ruhfel, R. B. Primack, A. J. Miller-Rushing, C. C. Davis. 2008.
1008 Phylogenetic patterns of species loss in Thoreau's woods are driven by climate change.
1009 *Proceedings of the National Academy of Sciences* 105:17029-17033.
1010
1011 Wright, J. S. 2002. Plant diversity in tropical forests: a review of mechanisms of species
1012 coexistence. *Oecologia*, 130:1-14.
1013
1014 Wright, K. M., U. Hellsten, C. Xu, A. L. Jeong, A. Sreedasyam, J. A. Chapman, J.
1015 Schmutz, G. Coop, D. S. Rokhsar, and J. H. Willis. Adaptation to heavy-metal
1016 contaminated environments proceeds via selection on pre-existing genetic variation.
1017 bioRxiv doi: http://dx.doi.org/10.1101/029900
1018
1019 Wu, C. A., D. B. Lowry, A. M. Cooley, K. M. Wright, Y. W. Lee, J. H. Willis. 2007.
1020 *Mimulus* is an emerging model system for the integration of ecological and genomic
1021 studies. Heredity, 100:220-230.
1022
1023 Wu, C. A., D. B. Lowry, L. I. Nutter, J. H. Willis. 2010. Natural variation for drought-
1024 response traits in the Mimulus guttatus species complex. *Oecologia,* 162:23–33.
1025
1026 Zuur, A., E. N. Ieno, N. Walker, A. A. Saveliev, G. M. Smith. 2009. Mixed effects
1027 models and extensions in ecology with R. Berlin, Germany: Springer.

1053
1054
Table 1. Results of the linear mixed effects model testing how environmental variables
affected survival across sites and time given as the equations with the highest AIC
criteria. In the model description SM = Soil Moisture, LD = Leaf Damage, T=Time, and
H=Habitat. "k" is the number of parameters in the model, "Log(L)" is the log likelihood,
"AIC" score, "Δi" is the difference between the AIC score of the current & top models,
"wi" is the model weight, Marginal R^2 is the contribution of the fixed effects, Conditional
R^2 is the total contribution of both random and fixed effects.
1062

Model	k	Log(L)	AIC	Δi	wi	Marginal R^2	Conditional R^2
Survival Total ~ SM + LD + T + H + SM*T + SM*H + LD*SM + SM*T*H	10	977.1	-1912	0	1	0.802	0.833
Survival Granite ~ SM + T + SM*T	5	399.6	-774.8	0	1	0.853	0.856
Survival Meadow ~ SM + T + LD + SM*T	6	962.4	-1896	0	0.607	0.627	0.692

1063
1064

1065 **Table 2.** Phenotypic trait means and standard errors calculated for day of first flower
1066 (days), plant height (mm), flower width (mm), leaf are (megapixels), and leaf shape *M.*
1067 *laciniatus's* granite outcrop versus *M. guttatus's* meadow habitat type in F$_4$'s, *M.*
1068 *laciniatus*, and *M. guttatus*. Significant phenotypic plasticity in a trait is indicated by "*"
1069 and significance codes are as follows: p-value < 0.001 = ***, 0.01= **,0.05 =*.

Plant	Trait	Mean Granite	SE Granite	Mean Meadow	SE Meadow
F4	Days to 1st Flower	77.85***	0.263	86.42***	0.774
	Height	40.55***	0.789	45.18***	1.764
	Flower Width	8.45***	0.131	6.292***	0.246
	Leaf Area	2009	157.6	3203	281.2
	Leaf Shape	0.13**	0.002	0.12**	0.004
M. laciniatus	Days to 1st Flower	76.96***	0.505	85.78***	1.327
	Height	28.59***	1.089	46.53***	4.766
	Flower Width	6.075***	0.253	3.875***	0.467
	Leaf Area	1068	144.0	1307	248.2
	Leaf Shape	0.18**	0.011	0.17**	0.018
M. guttatus	Days to 1st Flower	80.25***	0.743	82.3***	1.739
	Height	31.23***	1.667	74.46***	7.032
	Flower Width	8.8***	0.42	9.835***	0.473
	Leaf Area	1120	103.1	3524	647.1
	Leaf Shape	0.11	0.006	0.11	0.01

1070

1071
1072
1073

1074
1075 **Table 3.** Phenotypic correlation matrix from REML analysis among all traits at granite
1076 vs. meadow sites separately. The granite habitat correlations are in the bottom matrix in
1077 black and the meadow habitat correlations are on the top in gray.

	Days to Flowering	Stem Length	Flower Width	Leaf Area	Shape
Days to Flowering	**1**	0.11	0.023	0.128	-0.058
Stem Length	-0.073	**1**	0.557	0.534	0.010
Flower Width	0.047	0.677	**1**	0.305	0.0002
Leaf Area	-0.088	0.525	0.34	**1**	0.135
Shape	-0.052	0.046	0.08	0.08	**1**

Table 4. A list of the top phenotypic selection generalized linear mixed models based on
AIC score. For each analysis we report the complete model, k which is the number of
parameters, Log(L) which is the likelihood score, AIC score, Δi which is the difference
between that model's AIC and the top model's AIC, and wi which is the model weight.
FT stands for flowering time, FW stands for flower width, HT stands for height, LA
stands for leaf area, LS stands for leaf shape, and HB stands for habitat.
1089

Model	k	Log(L)	AIC	Δi	wi
Binomial					
Total Fitness ~ FT + FW + HT + LA + LS + HB + FT^2 + FT*HB + FT*HT	11	-329.388	687.517	0	0.420
Lifetime Fitness Granite ~ FT + FW + HT + LS + FT^2 + FT*HT	10	-238.845	498.270	0	0.886
Lifetime Fitness Meadow ~ FT + FW + HT + LS + HT^2 + LS*HT	10	-82.893	187.002	0	0.372
Truncated Poisson					
Lifetime Fitness ~ FT + FW + HT + LS + HB + FT*HB + LS*HB + LS*HT	12	-123.617	272.659	0	0.293
Lifetime Fitness Granite ~ FT + FW + HT + LS + LS*HT	8	-101.696	220.206	0	0.206
Lifetime Fitness Meadow ~ FT + FW + HT	6	-21.426	57.006	0	0.246

1090
1091
1092
1093
1094 **Table 5.** Results of phenotypic selection analysis on flowering time, flower width, plant
1095 height, leaf area, and leaf shape. This table contains the regression coefficients from the
1096 binomially distributed logistic generalized linear mixed model (glmm) with the

probability that the plant set fruit as the response variable. β represents the linear
selection gradient and γ represents the quadratic selection gradient for each phenotypic
trait in each habitat type. Significant codes are as follows: p-value < 0.001 = ***, 0.01=
**,0.05 =* .

Trait	β Granite	β Meadow	γ Granite	γ Meadow
Flowering Time	-0.957	-0.216	-0.801	--
Flower Width	0.331	-0.279	--	--
Height	-1.256	1.329	--	-0.55
Leaf Shape	0.066	-0.273	--	--
LS*Height	--	--	-0.096	0.062

Table 6. Results of phenotypic selection analysis on flowering time, flower width, plant
height, leaf area, and leaf shape. This table contains the regression coefficients from the
truncated Poisson distributed generalized linear mixed model on fruit count of plants that
did produce fruit. β represents the linear selection gradient and γ represents the quadratic
selection gradient for each phenotypic trait in each habitat type and population.
Significance codes are as follows: p-value < 0.001 = ***, 0.01= **,0.05 =* .

Trait	β Granite	β Meadow	γ Granite	γ Meadow
Flowering Time	-1.535	-0.228	--	--
Flower Width	0.47	0.484	--	--
Height	0.555	0.999	--	--
Leaf Shape	0.135	--	--	--
LS*Height	--	--	-0.479	--

1125   **Figure 1.** Images of (a) granite outcrop habitat (Granite 1), (b) *M. laciniatus* growing in
1126   its native granite habitat, (c) crossing design used to generate the F_4 hybrid population,
1127   (d) meadow habitat (Meadow 2), and (e) *M. guttatus* growing in its native meadow
1128   habitat.
1129
1130   **Figure 2.** Percent soil moisture (gray) and proportion plants surviving (black) measured
1131   weekly over the two month duration of the experiment on a block by block basis across
1132   the (a) *M. laciniatus* habitat (red) Granite 1 and (b) Granite 2 sites (c) *M. guttatus* habitat
1133   (blue) Meadow 1 and (d) Meadow 2 sites.
1134
1135   **Figure 3.** A boxplot of the number of herbivore damaged leaves per experimental block
1136   at the time of plant senescence in granite vs. meadow habitat.
1137
1138   **Figure 4.** Reaction norm plots of (a) mean survival & (b) fecundity of *M. laciniatus* (red)
1139   and *M. guttatus* (blue) parental inbred lines in granite (*M. laciniatus*) and meadow (*M.*
1140   *guttatus*) habitats.
1141
1142   **Figure 5.** Reaction norm plots demonstrating phenotypic plasticity between granite and
1143   meadow habitats for (a) mean flowering time (b) mean corolla width (c) mean plant
1144   height (d) mean leaf area, and (e) leaf shape for each genotypic class: *M. laciniatus* (red),
1145   *M. guttatus* (blue), and F_4's (purple).
1154   **Table S1**. Results of a logistic regression of survival to flowering on species and habitat
1155   to test for local adaptation. The model was Survival to Flowering ~ Habitat (Granite or
1156   Meadow) + Species (*M. guttatus* or *M. laciniatus*) + Habitat*Species.

Independent Variables	β'	z-value	p-value
Plant(L)	0.092	0.356	0.722
Habitat(L)	0.265	1.071	0.284
Plant(L)*Habitat(L)	0.964	2.876	0.004**

1157
1158

Table S2. Complete list of all binomial distributed generalized linear mixed models
testing which phenotypes influenced the probability that a plant produced fruit. For each
we report the complete model, k which is the number of parameters, Log(L) which is the
likelihood score, AIC score, Δi which is the difference between that model's AIC and the
top model's AIC, and wi which is the model weight. FT stands for flowering time, FW
stands for flower width, HT stands for height, LA stands for leaf area, LS stands for leaf
shape, and HB stands for habitat.
1166

Model Number	Model	k	Log(L)	AIC	Δi	wi
Total						
model29b	Fitness ~ FT + FW + HT + LA + LS + HB + FT^2 + FT*HT + FT*HB	14	-329.39	687.52	0	0.2519
model28b	Fitness ~ FT + FW + HT + LA + LS + FT^2 + FT*HT	12	-331.7	687.95	0.44	0.2026
model33b	Fitness ~ FT + FW + HT + LA + LS + HB + FT^2 + FT*HT + FT*HB + LS*HB	15	-328.66	688.17	0.66	0.1816
model32b	Fitness ~ FT + FW + HT + LA + LS + HB + FT^2 + FT*HT + FT*HB + HT*HB	15	-329.01	688.86	1.34	0.1288
model34b	Fitness ~ FT + FW + HT + LA + LS + HB + FT^2 + FT*HT + FT*HB + HT*HB + LS*HB	16	-328.3	689.57	2.05	0.0902
model27b	Fitness ~ FT + FW + HT + LA + LS + HB + FT^2 + FT*HT	13	-331.62	689.88	2.36	0.0773
model30b	Fitness ~ FT + FW + HT + LA + LS + HB + FT^2 + FT*HT + HT*HB	14	-331.54	691.81	4.29	0.0294
model22b	Fitness ~ FT + FW + HT + LA + LS + HB + FT*HT	12	-334.8	694.15	6.64	0.0091
model17b	Fitness ~ FT + FW + HT + LA + LS + FT^2	11	-337.01	696.48	8.96	0.0029
model16b	Fitness ~ FT + FW + HT + LA + LS	10	-338.06	696.51	8.99	0.0028
model13b	Fitness ~ FT + HT + HB + FT*HB + HT*HB	10	-338.08	696.55	9.03	0.0028
model4b	Fitness ~ FT + FW + HT + LA + HB	10	-338.26	696.91	9.39	0.0023
model10b	Fitness ~ FT + FW + HT + LA + HB + FT*HB	11	-337.48	697.43	9.91	0.0018
model18b	Fitness ~ FT + FW + HT + LA + LS + FW^2	11	-337.56	697.58	10.06	0.0016
model21b	Fitness ~ FT + FW + HT + LA + LS + HB + FT^2 + FT^2*HB	13	-335.61	697.85	10.34	0.0014
model31b	Fitness ~ FT + FW + HT + LA + LS + LS^2	11	-337.76	697.97	10.46	0.0014
model14b	Fitness ~ FT + HT + HB + FT*HB	9	-339.84	697.99	10.47	0.0013
model23b	Fitness ~ FT + FW + HT + LA + LS + HB + FW*HT	12	-336.76	698.07	10.55	0.0013
model20b	Fitness ~ FT + FW + HT + LA + LS + LA^2	11	-	698.19	10.67	0.0012

			337.86			
model19b	Fitness ~ FT + FW + HT + LA + LS + HT^2	11	-337.9	698.26	10.74	0.0012
model3b	Fitness ~ FT + FW + HT + HB	9	-339.99	698.31	10.8	0.0011
model5b	Fitness ~ FT + FW + HT + LA + LS + HB	11	-338.05	698.56	11.04	0.001
model11b	Fitness ~ FT + FW + HT + HB + FT*HB	10	-339.36	699.1	11.58	0.0008
model6b	Fitness ~ FT + FW + HT + LA + LS + HB + FT*HB	12	-337.31	699.17	11.65	0.0007
model25b	Fitness ~ FT + FW + HT + LA + LS + HB + LS*HT	12	-337.72	699.99	12.48	0.0005
model26b	Fitness ~ FT + FW + HT + LA + LS + HB + LS*LA	12	-337.72	699.99	12.48	0.0005
model0b	Fitness ~ FT	6	-343.95	700.05	12.54	0.0005
model24b	Fitness ~ FT + FW + HT + LA + LS + HB + LA*HT	12	-337.99	700.54	13.02	0.0004
model1b	Fitness ~ FT + HB	7	-343.69	701.57	14.05	0.0002
model2b	Fitness ~ FT + HB + FW	8	-343.45	703.15	15.64	0.0001
model12b	Fitness ~ FT + HB + FW + FT*HB	9	342.49	703.29	15.77	1.00E-04

Granite

model36b.L	Fitness ~ FT + FW + HT + LS + FT^2 + FT*HT	10	-237.69	495.95	0	0.5744
model37b.L	Fitness ~ FT + FW + HT + LS + FT^2 + LS*HT + FT*HT	11	-237.58	497.85	1.9	0.2221
model30b.L	Fitness ~ FT + FW + HT + LS + FT^2 + LS*HT	10	-238.85	498.27	2.32	0.1802
model27b.L	Fitness ~ FT + FW + HT + LS + FT*HT	9	-243.63	505.72	9.77	0.0043
model22b.L	Fitness ~ FT + FW + HT	7	-245.72	505.73	9.77	0.0043
model29b.L	Fitness ~ FT + FW + HT + LA + LS	9	-243.79	506.06	10.1	0.0037
model23b.L	Fitness ~ FT + FW + HT + LS	8	-245.25	506.88	10.93	0.0024
model35b.L	Fitness ~ FT + FW + HT + LA + LS + LA^2	10	-243.34	507.25	11.3	0.002
model26b.L	Fitness ~ FT + FW + HT + LS + FT^2 + LS*HT + FT*HT	10	-243.47	507.53	11.58	0.0018
model34b.L	Fitness ~ FT + FW + HT + LA + LS + LS*HT	10	-243.76	508.1	12.15	0.0013
model25b.L	Fitness ~ FT + FW + HT + LS + LS*HT	9	-245.1	508.66	12.71	0.001
model20b.L	Fitness ~ FT + FW + LS + FT*FW	8	-246.67	509.73	13.77	0.0006
model33b.L	Fitness ~ FT + FW + HT + LS + LS^2 + LS*HT	10	-244.84	510.27	14.32	0.0004

model32b.L	Fitness ~ FT + FW + HT + LS + HT^2 + LS*HT	10	-245.03	510.64	14.69	0.0004
model31b.L	Fitness ~ FT + FW + HT + LS + FW^2 + LS*HT	10	-245.08	510.73	14.78	0.0004
model28b.L	Fitness ~ FT + FW + HT + LS + LS*HT + FW*HT	10	-245.1	510.77	14.82	0.0003
model18b.L	Fitness ~ FT + FW + LS + FT*FW + FT*LS	9	-246.47	511.42	15.47	0.0003
model21b.L	Fitness ~ FT + FW + LS	7	-251.73	517.76	21.81	0.00E+00
model19b.L	Fitness ~ FT + FW + LS + FT*LS	8	-251.65	519.67	23.72	0.00E+00

Meadow

model32b.G	Fitness ~ FT + FW + HT + LS + HT^2 + LS*HT	10	-82.89	187	0	0.3449
model20b.G	Fitness ~ FT + FW + LS + FT*FW	8	-86.21	189.21	2.21	0.1145
model21b.G	Fitness ~ FT + FW + LS	7	-87.35	189.3	2.3	0.1094
model36b.G	Fitness ~ FT + FW + HT + LS + FT^2 + FT*HT	10	-84.74	190.7	3.7	0.0542
model28b.G	Fitness ~ FT + FW + HT + LS + LS*HT + FW*HT	10	-84.93	191.08	4.08	0.0449
model22b.G	Fitness ~ FT + FW + HT	7	-88.36	191.34	4.33	0.0395
model27b.G	Fitness ~ FT + FW + HT + LS + FT*HT	9	-86.18	191.35	4.35	0.0393
model18b.G	Fitness ~ FT + FW + LS + FT*FW + FT*LS	9	-86.21	191.41	4.4	0.0382
model19b.G	Fitness ~ FT + FW + LS + FT*LS	8	-87.32	191.42	4.42	0.0379
model23b.G	Fitness ~ FT + FW + HT + LS	8	-87.35	191.48	4.48	0.0368
model24b.G	Fitness ~ FT + HT + LS + LS*HT	8	-87.35	191.49	4.48	0.0366
model37b.G	Fitness ~ FT + FW + HT + LS + FT^2 + LS*HT + FT*HT	11	-84.74	192.95	5.95	0.0176
model30b.G	Fitness ~ FT + FW + HT + LS + FT^2 + LS*HT	10	-85.95	193.12	6.12	0.0162
model29b.G	Fitness ~ FT + FW + HT + LA + LS	9	-87.13	193.25	6.25	0.0152
model26b.G	Fitness ~ FT + FW + HT + LS + FT^2 + LS*HT + FT*HT	10	-86.18	193.57	6.57	0.0129
model25b.G	Fitness ~ FT + FW + HT + LS + LS*HT	9	-87.34	193.68	6.68	0.0122
model31b.G	Fitness ~ FT + FW + HT + LS + FW^2 + LS*HT	10	-86.45	194.12	7.12	0.0098
model33b.G	Fitness ~ FT + FW + HT + LS + LS^2 + LS*HT	10	-86.56	194.33	7.33	0.0088
model35b.G	Fitness ~ FT + FW + HT + LA + LS + LA^2	10	-86.93	195.07	8.07	0.0061
model34b.G	Fitness ~ FT + FW + HT + LA + LS + LS*HT	10	-87.12	195.46	8.46	0.005

1175 **Table S3.** Complete list of all truncated Poisson distributed generalized linear mixed
1176 models testing which phenotypes influenced the number of fruits a plant produced. For

each we report the complete model, k which is the number of parameters, Log(L) which is the likelihood score, AIC score, Δi which is the difference between that model's AIC and the top model's AIC, and wi which is the model weight. FT stands for flowering time, FW stands for flower width, HT stands for height, LA stands for leaf area, LS stands for leaf shape, and HB stands for habitat.

Model Number	Model	k	Log(L)	AIC	Δi	wi
Total						
model31t	Fitness ~ FT + FW + HT + LS + FT*HB +LS*HB + LS*HT	12	-123.62	272.68	0	0.2933
model33t	Fitness ~ FT + FW + HT + LS + FT*HB +LS*HB + HT*HB + LS*HT	13	-122.54	272.75	0.091	0.2802
model25t	Fitness ~ FT + FW + HT + LS + FT*HB + LS*HT	11	-126.15	275.5	2.843	0.0708
model11t	Fitness ~ FT + FW + HT + FT*HB	9	-128.88	276.56	3.904	0.0416
model17t	Fitness ~ FT + FW + HT + FT*HB + FT^2	10	-127.88	276.76	4.103	0.0377
model32t	Fitness ~ FT + FW + HT + LS + FT*HB + HT*HB + LS*HT	12	-125.68	276.79	4.13	0.0372
model12t	Fitness ~ FT + HT + FT*HB	8	-130.41	277.46	4.805	0.0265
model26t	Fitness ~ FT + FW + HT + LA + LS + FT*HB + LA*LS	12	-126.11	277.65	4.992	0.0242
model22t	Fitness ~ FT + FW + HT + LS + FT*HB + FT*HT	11	-127.25	277.69	5.035	0.0236
model18t	Fitness ~ FT + FW + HT + LS + FW^2	10	-128.66	278.32	5.663	0.0173
model23t	Fitness ~ FT + FW + HT + LS + FT*HB + FW*HT	11	-127.61	278.42	5.757	0.0165
model10t	Fitness ~ FT + FW + HT + LA + FT*HB	10	-128.8	278.6	5.939	0.0151
model6t	Fitness ~ FT + FW + HT + LA + LS + FT*HB	11	-127.73	278.66	6.001	0.0146
model19t	Fitness ~ FT + FW + HT + FT*HB +HT^2	10	-128.87	278.74	6.079	0.014
model29t	Fitness ~ FT + FW + HT + LA + LS + HB + FT^2 + FT*HB + FT*HT	13	-125.55	278.77	6.111	0.0138
model21t	Fitness ~ FT + FW + HT + LS + LS^2 + FT*HB	11	-127.87	278.93	6.271	0.0127
model15t	Fitness ~ FT + HT + FT*HB +HT*HB	9	-130.1	279.01	6.348	0.0122
model28t	Fitness ~ FT + FW + HT + LA + LS + FT^2 + FT*HT	11	-128	279.2	6.545	0.0111
model24t	Fitness ~ FT + FW + HT + LA + LS + HB + FT*HB + LA*HT	12	-127.06	279.54	6.878	0.0094
model20t	Fitness ~ FT + FW + HT + LA + HB + LA^2	11	-128.21	279.62	6.961	0.009
model7t	Fitness ~ FT + FW + HT + LA + LS + HB + FT*HB + HT*HB	12	-127.3	280.03	7.374	0.0073
model27t	Fitness ~ FT + FW + HT + LA + LS + HB +	12	-	280.91	8.246	0.0047

	FT^2 + FT*HT		127.74			
model8t	Fitness ~ FT + FW + HT + LA + LS + HB + FT*HB + HT*HB + FW*HB	13	-127.21	282.09	9.429	0.0026
model30t	Fitness ~ FT + FW + HT + LA + LS + HB + FT^2 + HT*HB + FT*HT	13	-127.58	282.82	10.161	0.0018
model9t	Fitness ~ FT + FW + HT + LA + LS + HB + FT*HB + HT*HB + FW*HB + LA*HB	14	-127.14	284.22	11.565	0.0009
model13t	Fitness ~ FT + FW + HB + FT*HB	8	-134.18	285.01	12.353	0.0006
model16t	Fitness ~ FT + FW + HT + LA + LS	9	-133.34	285.49	12.832	0.0005
model3t	Fitness ~ FT + FW + HT + HB	8	-135.03	286.7	14.045	0.0003
model5t	Fitness ~ FT + FW + HT + LA + LS + HB	10	-133.31	287.61	14.953	0.0002
model4t	Fitness ~ FT + FW + HT + LA + HB	9	-134.8	288.42	15.758	0.0001
model2t	Fitness ~ FT + FW + HT + HB	7	-141.32	297.14	24.481	0.00E+00
model14t	Fitness ~ FT + HB + FT*HB	7	-143.94	302.37	29.715	0.00E+00
model1t	Fitness ~ FT + HB	6	-149.29	310.95	38.289	0.00E+00
model0t	Fitness ~ FT	5	-151.3	312.86	40.199	0.00E+00
Granite						
model25t.L	Fitness ~ FT + FW + HT + LS + LS*HT	8	-101.7	220.21	0	0.2064
model26t.L	Fitness ~ FT + FW + HT + LS + LS*HT + FT*HT	9	-100.91	220.85	0.639	0.1499
model34t.L	Fitness ~ FT + FW + HT +LA + LS + LS*HT	9	-101.05	221.12	0.909	0.131
model28t.L	Fitness ~ FT + FW + HT + LS + LS*HT + FW*HT	9	-101.19	221.4	1.197	0.1134
model30t.L	Fitness ~ FT + FW + HT + LS + LS*HT + FT^2	9	-101.43	221.87	1.667	0.0897
model31t.L	Fitness ~ FT + FW + HT + LS + LS*HT + FW^2	9	-101.65	222.32	2.111	0.0718
model33t.L	Fitness ~ FT + FW + HT + LS + LS*HT + LS^2	9	-101.66	222.33	2.127	0.0713
model24t.L	Fitness ~ FT + HT + LS + LS*HT	7	-103.88	222.39	2.184	0.0693
model32t.L	Fitness ~ FT + FW + HT + LS + LS*HT + HT^2	9	-101.69	222.4	2.197	0.0688
model22t.L	Fitness ~ FT + FW + HT	6	-106.7	225.87	5.666	0.01215
model23t.L	Fitness ~ FT + FW + HT + LS	7	-106.23	227.08	6.878	0.0066
model29t.L	Fitness ~ FT + FW + HT + LA + LS	8	-106.16	229.14	8.93	0.0024
model27t.L	Fitness ~ FT + FW + HT + LS + FT*HT	8	-106.21	229.23	9.026	0.0023
model35t.L	Fitness ~ FT + FW + HT + LA + LS + LA^2	9	-105.46	229.94	9.737	0.0016

model21t.L	Fitness ~ FT + FW + LS	6	-108.8	230.07	9.862	0.0015
model19t.L	Fitness ~ FT + FW + LS + FT*LS	7	-108.26	231.18	10.974	0.0009
model20t.L	Fitness ~ FT + FW + LS + FT*FW	7	-108.5	231.63	11.426	0.0007
model18t.L	Fitness ~ FT + FW + LS + FT*FW + FT*LS	8	-108.09	232.99	12.782	0.0003
Meadow						
model22t.G	Fitness ~ FT + FW + HT	6	-21.43	57.01	0	0.2463
model23t.G	Fitness ~ FT + FW + HT + LS	7	-20.28	57.5	0.494	0.1924
model32t.G	Fitness ~ FT + FW + HT + LS + LS*HT + HT^2	9	-18.07	59.15	2.142	0.0844
model21t.G	Fitness ~ FT + FW + LS	6	-22.53	59.21	2.199	0.082
model33t.G	Fitness ~ FT + FW + HT + LS + LS*HT + LS^2	9	-18.19	59.37	2.364	0.0755
model24t.G	Fitness ~ FT + HT + LS + LS*HT	7	-21.59	60.13	3.128	0.0515
model25t.G	Fitness ~ FT + FW + HT + LS + LS*HT	8	-20.18	60.25	3.241	0.0487
model27t.G	Fitness ~ FT + FW + HT + LS + FT*HT	8	-20.19	60.27	3.266	0.0481
model29t.G	Fitness ~ FT + FW + HT + LA + LS	8	-20.27	60.43	3.428	0.0444
model20t.G	Fitness ~ FT + FW + LS + FT*FW	7	-22.14	61.23	4.227	0.0298
model19t.G	Fitness ~ FT + FW + LS + FT*LS	7	-22.5	61.94	4.935	0.0209
model30t.G	Fitness ~ FT + FW + HT + LS + LS*HT + FT^2	9	-19.88	62.75	5.745	0.0139
model31t.G	Fitness ~ FT + FW + HT + LS + LS*HT + FW^2	9	-19.97	62.94	5.929	0.0127
model26t.G	Fitness ~ FT + FW + HT + LS + LS*HT + FT*HT	9	-20.05	63.1	6.093	0.0117
model28t.G	Fitness ~ FT + FW + HT + LS + LS*HT + FW*HT	9	-20.14	63.27	6.265	0.0107
model34t.G	Fitness ~ FT + FW + HT +LA + LS + LS*HT	9	-20.17	63.34	6.329	0.0104
model35t.G	Fitness ~ FT + FW + HT + LA + LS + LA^2	9	-20.24	63.47	6.464	0.0097
model18t.G	Fitness ~ FT + FW + LS + FT*FW + FT*LS	8	-22.14	64.17	7.166	0.0068