No evidence for genome editing in mouse zygotes and HEK293T human cell line

using the DNA-guided *Natronobacterium gregoryi* Argonaute (NgAgo).

Short title: No evidence for genome editing using NgAgo

Authors: Nay Chi Khin, Jenna L. Lowe, Lora M. Jensen and Gaetan Burgio

Department of Immunology and Infectious diseases, the John Curtin School of Medical

Research, The Australian National University, Canberra, Australia.

Corresponding author:

Gaetan Burgio:

The John Curtin School of Medical Research

The Australian National University

131 Garran Road, Acton, ACT 2601, Australia

Ph: +61 2 6125 9428

email: Gaetan.burgio@anu.edu.au

Twitter handle @GaetanBurgio

Abstract: A recently published research article reported that the extreme halophile archaebacterium *Natronobacterium gregoryi* Argonaute enzyme (NgAgo) could cleave the cellular DNA under physiological temperature conditions in cell line and be implemented as an alternative to CRISPR/Cas9 genome editing technology. We assessed this claim in mouse zygotes for four loci (*Sptb*, *Tet-1*, *Tet-2* and *Tet-3*) and in the human HEK293T cell line for the EMX1 locus. Over 100 zygotes were microinjected with nls-NgAgo-GK plasmid provided from Addgene and various concentrations of 5'-phosphorylated guide DNA (gDNA) from 2.5 ng/µl to 50 ng/µl and cultured to blastocyst stage of development. The presence of indels was verified using T7 endonuclease 1 assay (T7E1) and Sanger sequencing. We reported no evidence of successful editing of the mouse genome. We then assessed the lack of editing efficiency in HEK293T cell line for the EMX1 endogenous locus by monitoring the NgAgo protein expression level and the editing efficiency by T7E1 assay and Sanger sequencing. We reported that the NgAgo protein was expressed from 8 hours to a maximum expression at 48 hours post-transfection, confirming the efficient delivery of the plasmid and the gDNA but no evidence of successful editing of EMX1 target in all transfected samples. Together our findings indicate that we failed to edit using NgAgo.

Introduction

Type II CRISPR/Cas9 genome editing system offers the ability to efficiently and precisely edit DNA using a combination of the Cas9 endonuclease enzyme and a single guide RNA gRNA [1]. However the requirement of a specific protospacer adjacent motif (PAM) sequence limits the ability of the Cas9 enzyme to edit any nucleotide of a genome. Recently a report described a novel genome editing technology based on the archaebacterium *Natrnobacterium gregoryi* Argonaute (NgAgo) enzyme. Gao et al. described an endonuclease activity for NgAgo, which has the ability to create a site-specific double strand break in the DNA under the guidance of the 24 nucleotide- 5'-phosphorylated single stranded DNA (gDNA) which binds to the endogenous DNA [2]. Gao et al. demonstrated the high efficiency of NgAgo to edit the genome under physiological temperature condition without the requirement of a PAM [2]. Interestingly, Gao et al. demonstrated the intolerance of NgAgo to guide-target mismatches leading to negligible off-target effects. With such ability to target any nucleotide in the genome and with an equivalent efficiency to CRISPR/Cas9 genome editing technology, NgAgo has an undeniably important therapeutic potential [3]. We sought to assess the efficiency of NgAgo in two different systems: mouse zygotes and the HEK293T human cell line to determine the suitability of NgAgo as an alternative to the CRISPR/Cas9 genome editing tool. Here we report our attempts to edit the mouse and human genomes using NgAgo. We synthetized the gDNA and co-microinjected in mouse zygotes with the nls-NgAgo-GK plasmid vector provided by Gao et al. We also co-transfected the gDNA with a modified Flag-nls-NgAgo-GK plasmid into HEK293T cells and assessed the DNA editing using T7 endonuclease assay and Sanger sequencing. We monitored the

expression of the protein for the first 48 hours post-transfection in HEK293T cells. We

found no evidence for a double strand break and editing of the DNA under various

conditions and optimizations. We concluded that we failed to edit the genome using

NgAgo.

Results and discussion

To assess NgAgo efficiency to create a double strand break under the guidance of a

single gDNA as described in Gao et al. [2] we used mouse zygotes as a system model.

We firstly co-injected mouse zygotes with the nls-NgAgo-GK plasmid and the gDNA

targeting four different genes (*Sptb*, *Tet1*, *Tet2* and *Tet3*). The gDNA selected were

previously shown to edit efficiently (over 60% efficiency) as an sgRNA with

CRISPR/Cas9 mediated genome editing [4]. Initially, we targeted exon 26 of *Sptb* in

mouse zygotes (Figure 1A). We titrated the gDNA at various concentrations (2.5, 25 or

50 ng/µl) and co-injected with 5, 10 or 15 ng/µl of nls-NgAgo-GK plasmid, provided by

Gao et al. and available at Addgene, into the pronucleus of the fertilized zygotes. The

zygotes were cultured for 4 days to the blastocyst stage of development. No abnormality

in the development was found in these embryos. From the 49 bastocysts that were

genotyped (Table 1), we found the amplification of a band corresponding to the expected

amplicon length for *Sptb* (Figure 1B presented on NgAgo 5 ng/µl and 50 ng/µl of

gDNA). We performed a T7 endonuclease assay (T7E1) and Sanger sequencing to

identify indels. We could not identify any indels from the T7E1 assay (Figure 1C) and the

Sanger sequencing (Figure 1D) on all blastocysts. We then hypothesized the lack of

editing could be gene specific, hence we decided to assess three other genes; exon 5 of

Tet-1 (Suppl Figure 1A), exon 3 of *Tet-2* (Suppl Figure 1B) and exon 5 of *Tet-3* (Suppl

Figure 1C) as described previously [4]. The zygotes were co-injected with 5ng/µl of nls-

NgAgo-GK plasmid and 2.5 ng/µl of gDNA. We constantly found one band using gel

electrophoresis corresponding to the expected amplicon size for *Tet-1* (10 blastocysts),

Tet-2 (14 blastocysts) and *Tet-3* (13 blastocysts). We performed a T7E1 assay and

genotyped the blastocysts by Sanger sequencing. We found again, no evidence for

developmental phenotype and presence of indels (data not shown). We further assessed

the NgAgo editing efficiency for *Sptb* and *Tet-2* on pups born from the microinjection

sessions. All pups displayed a normal phenotype and developed normally to adulthood.

We performed a T7E1 assay and Sanger sequencing to assess the editing efficiency and

we did not find any indels, suggesting no successful editing for *Sptb* and for *Tet-2* (Table

2). Together, this suggests that we failed to generate indels for the four analyzed mouse

loci (*Sptb*, *Tet-1*, *Tet-2* and *Tet-3*) in over 86 mouse blastocysts and 25 mouse pups using

NgAgo, suggesting NgAgo does not create a double strand break nor that it is capable of

editing the mouse genome. We therefore speculated this lack of editing activity could be

due to the degradation of the protein within the cell. To address this hypothesis, we

tagged the protein with a flag tag upstream of the nls signal (Flag-nls-NgAgo-GK, Figure

1E) and monitored the expression of the protein and whether NgAgo edited the DNA in

the HEK293T cell line at various time points from 8 to 48 hours post lipofection with or

without the gDNA for exon 3 of the EMX1 gene (Figure 1F). We used an anti-flag

antibody to probe for NgAgo expression, and utilized GAPDH as a loading control. We

first conducted a PCR and a T7E1 assay on the NgAgo-EMX1-lipofectamine treated cells

to determine whether the DNA was edited. We found no evidence for editing in any

NgAgo-lipofection samples at 8 and 12 hours post-transfection (Figure 1G). Similarly,

there was no editing for the additional time points of 24 and 48 hours post transfection

(Supplementary Figure 2 A and B). We therefore speculated that the NgAgo protein

could be not expressed post transfection or rapidly degraded after transfection and

required a specific timing to edit the DNA. To verify this hypothesis, we followed the

kinetics of the NgAgo protein production and degradation from 8 to 48 hours post

transfection. We noticed the NgAgo protein expression started at 8 hours and persisted

over 48 hours post transfection (Figure 1H), with maximum expression being observed at

48 hours post transfection compared to 8 hours (p = 0.02) (Figure 1H and Supplementary

Figure 3) suggesting an efficient delivery of NgAgo and the gDNA. Interestingly, we

noticed the presence of small-Flag-tagged fragments by 8 hours post transfection,

suggesting the protein started degrading, with the fragment intensity reaching its peak at

48 hours post-transfection associated with the peak protein expression mentioned above

with and without the co-transfection of the gDNA (Figure 1H). We noted no difference in

protein expression and degradation with or without the gDNA (p = 0.0631 and p = 0.25

respectively) (Figure 1H and Supplementary Figure 3). Therefore, since degradation of

the protein is the same between NgAgo treated samples with and without gDNA, the

difference in protein expression is unlikely to be due to this rapid degradation.

Gao et al. reported that NgAgo creates a double strand break in the DNA using a single

DNA guide [2], with a reported efficiency equivalent to Cas9. Importantly, and in

agreement with recently published reports [5-7], we found no evidence of the mouse and

human DNA editing with NgAgo despite an efficient delivery of NgAgo and the gDNA.

We did not observe an editing event in over 100 mouse embryos injected with NgAgo,

giving an editing efficiency of less than 1%, contradicting the results reported in Gao et

al. Interestingly, we found in the mouse embryos, mouse pups and in the HEK293T

human cell line no evidence of a single indel using a T7E1 assay, and confirmed by

Sanger sequencing. Qi et al reported recently a silencing role of NgAgo that may affects

the phenotype of the Zebrafish embryos [6]. We have not noted such a change in

phenotype in our mouse embryos. The plausible explanation for this lack of phenotype is

the genes targeted were not expressed during early embryonic development [8].

Although, we did not isolate mRNA from these mouse blastocysts and we did not assess

the expression of the genes.

In summary, in contradiction with Gao et al's study, and in agreement with recently

published reports, we found that NgAgo does not edit endogenous genomic DNA under

physiological temperature conditions.

Material and Methods:

Design and preparation of NgAgo, 5'-phosphorylated guide DNA (gDNA):

Four genes were targeted to design the primers and the 5'-phosphorylated

oligonucleotides. These genes were the exon 26 of *Beta-Spectrin1* (*Sptb*), exon 5 of *Tet-*

1, exon 3 of *Tet-2* and exon 5 of *Tet-3*. We choose gDNA previously published to be

highly efficient using the CRISPR/Cas9 genome editing system. The NgAgo plasmid

containing a Nuclear Localisation Signal was obtained from the Addgene repository [2].

The plasmid was cultured as per protocol and the DNA extracted using a PureLink Quick

Plasmid Miniprep Kit (Invitrogen, K210010) according to the manufacturer's

instructions. 22 and 24 bp 5' phosphorylated oligonucleotides and the amplification

primers were synthetized from Integrated DNA Technologies. The sequences of these

oligonucleotides are listed below (Table 3).

Ethics statement :

All animal experiments were approved by The Australian National University Animal

Experimentation Ethics Committee under the permit A2014/058 and the institutional

Biosafety Committee NLRD 15.10 in accordance with the National Health and

Medical Research Council (NHMRC) code of practice.

Mouse husbandry and microinjection:

C57BL/6 and recipient ICR females were purchased from Charles River laboratory

and maintained in a specific pathogen-free environment at the Australian Phenomics

Facility, the Australian National University. Mice were maintained on a 12h light/12h

dark cycle and had ad libitum access to food and water *ad libitum*. Female C57BL/6

(3-4 weeks old, >10g) were superovulated by intraperitoneal injection of 5IU of

pregnant mare serum gonadotropin (PMSG), followed by 5IU of human chorionic

gonadotropin hormone (hCG) 46-48 hours later. Following injection with hCG,

superovulated females were mated with stud C57BL/6 males (10-20 weeks old). The

embryos were collected from oviducts approximately 45 hours after the last injection

and held in M16 medium (Sigma M7292) overlaid with mineral oil at 37ºC and 5%

CO_2 until injection.

Microinjection was performed in M2 medium (Sigma, M7167) under mineral oil using

an inverted microscope (Leica DMi8) and micromanipulators. Pronuclear injection of

fertilized zygotes was performed with the following mixes: circular plasmid DNA at 5

ng/µl and 5'-P oligonucleotide at 2.5, 25 and 50 ng/µl. For Sptb CRISPR injection, the

gRNA was cloned into px330 vector [9] obtained from Addgene (ID 42330) using the

following oligonucleotides 5'- CACCGTGACATGTGGGCGGACCTGC - 3' and 5' –

AAACGCAGGTCCCCACATGTCAC – 3'. 5 ng/µl of px330 circular plasmid was

injected into the fertilized zygotes by pronuclear injection to obtained live mice. For

blastocysts culture, fertilized zygotes were microinjected using the following mix: 50

ng/µl of Cas9 purified protein from PNA BIO (Thoussand Oaks, CA), 0.6 pMol of

CrRNA 5' – UGACAUGUGGGCGGACCUGGUUUAGAGCUAUGCUGUUUUG -

3' and 0.6 pMol of TracRNA. The CrRNA and TracRNA were complexed with the

Cas9 protein by incubating at 37ºC for 10 minutes. Microinjected zygotes were

cultured overnight in M16. Resulting two-cell embryos were surgically transferred into

the ampulla of pseudo-plugged ICR female recipients (8-12 weeks old) or cultured in

M16 media for 4 days at 37ºC.

Genotyping:

A subset of the microinjected zygotes was cultured for 4 days to the blastocyst stage in

M16 medium overlaid with mineral oil at 37ºC and 5% CO_2. The other zygotes were

cultured for 24 hours and surgically transferred into the surrogate mouse ampulla. The

mice were maintained and the resulting pups were maintained and genotyped 15 days

after birth. DNA was extracted from the blastocysts at day 5 or live mouse pups over

15 days old using a crude DNA extraction protocol. In short, the blastocysts were

lysed in Tris-EDTA-Tween lysis buffer (50mMTris HCl, pH8.0, 0.125mM EDTA, 2%

Tween 20) with 1µl of proteinase K (20 mg/ml in 10mM Tris chlorate, 0.1 mM

ethylenediaminetretaacetic acid (EDTA) pH 8.0) and incubated at 56ºC for an hour

before being denatured at 95ºC for 10 minutes. We amplified regions encompassing

the gDNA with 2x MyTaq HS mix (Bioline, cat no. BIO-25045) under the following

PCR conditions: 95ºC for 3 minutes followed by 35 cycles (95ºC for 15", 58ºC for 15"

and 72ºC for 20") and 72ºC for 3 minute. The PCR products were checked on a 1.5%

electrophoresis gel. The PCR products were purified with ExoSAP-IT® (affymetrix,

Cat no. 78202), or cut from the gel and purified using the Wizard® SV Gel and PCR

Clean-Up System (Promega, Cat no. A9282) kit according to the manufacturer's

instructions. The Sanger sequencing was conducted at the Biomedical Resource

Facility at the John Curtin School for Medical Research, The Australian National

University.

T7 endonuclease assay.

After PCR amplification, the PCR fragments were hybridized and digested with a T7

endonuclease (NEB, Cat no. M0302S) for 15-30 minutes at 37ºC. After digestion, the

enzymatic reaction was stopped using 1 µl of 0.25 M EDTA and the digested product

run on a 1.5% agarose gel alongside the undigested PCR product as a control.

Construction of a Flag-nls-NgAgo-GK plasmid:

The nls-NgAgo-GK plasmid was a gift from Chunyu Han (Addgene plasmid 78253).

nls-NgAgo-GK plasmid was digested overnight with AleI (NEB, cat no. R0634S).

Following   the   digestion   a   forward   (5'-

TGGACTATAAGGACCACGACGGAGACTACAAGGATCATGATATTGATTAC

AAAGACGATGACGATAAGA-3')   and   a   reverse   (5'   -

TCTTATCGTCATCGTCTTTGTAATCAATATCATGATCCTTGTAGTCTCCGTC

GTGGTCCTTATAGTCCA-3') oligonucleotide sequence encoding for a flag tag were

annealed and ligated into the nls-NgAgo-GK plasmid upstream to the SV40 Nuclear

Localization Signal (nls). Sanger sequencing was used to assess the correct integration

of the flag-tag in frame with the start codon. The plasmid was transformed into heat-

chock competent BL21 *E. coli*. The plasmid is being deposited at Addgene (Plasmid

#73681) and will be available to the community.

Cell culture and transfection:

HEK293T cells were obtained from ATCC (CRL-11268). The cells were maintained

in Dulbecco's Modified Eagle Medium (DMEM) (Sigma, Cat. No. D6546)

supplemented with 10% heat-inactivated fetal bovine serum (Sigma, Cat. No.

12003C), 2mM L-Glutamine/1%Penicilin/Streptomycin solution (Thermofisher, Cat.

No. 10378016) and incubated at 37ºC with 5% CO_2. The cells were seeded at $2x10^5$

cells per well in 2.5 mL of medium in a 6-well plate, to reach 60-70% confluency

immediately prior to transfection. Per sample well, 3.0ug of plasmid DNA (Flag-nls-

NgAgo-GK) and/or 0.5µg of 5' phosphorylated EMX1 guide oligo was added to

150uL of basal DMEM (without additives) with 3.5uL Plus Reagent (Invitrogen,

Thermofisher, Cat. No. 15338100). 150uL of this diluted DNA mixture was added to

150uL of DMEM (without additives) and 12uL of Lipofectamine LTX® Reagent

(Invitrogen, Thermofisher, Cat. No. 15338100). The mixture was incubated for 15

minutes at room temperature to form DNA-Lipofectamine LTX® Reagent complexes.

After incubation, 250uL of the DNA-Lipofectamine complex was added drop-wise to

each well, and the plate was gently rocked. Negative controls constituted a 'cell

growth' control, with no DNA or lipofectamine reagents, and a 'lipofectamine only'

control with no plasmid or gDNA. Transfected cells were incubated at 37ºC in a 5%

CO_2 incubator for 8, 12, 24 and 48 hours post-transfection before collecting proteins to

assay for transgene expression, and extracting genomic DNA for genotyping assays.

Genomic DNA extraction and protein extraction from HEK293T cells:

The genomic DNA was isolated and purified using ISOLATE II Genomic DNA Kit

(Bioline, BIO-52066) according to the manufacturer's instructions. To extract proteins

from HEK293T cells, the cells were incubated with RIPA buffer (5M NaCl, 0.5M

EDTA, pH 8.0, 1M Tris, pH8.0, 1% TritonX-100, 10% sodium deoxycholate, 10%

SDS, 1x Protease Phosphatase Inhibitor) with gentle rocking on ice for 30 minutes.

The cells were scraped off the plates to dislodge lysate. Lysate was centrifuged at

13,000g for 5 minutes at 4ºC. The supernatant of lysate was stored at -20ºC.

Western blotting:

The nuclear and cytoplasm lysate samples were denatured at 95°C for 5minutes in 1x

Laemmli buffer prior to loading onto a 4-15% SDS-PAGE (4–15% Mini-PROTEAN®

TGX^{TM} Precast Protein Gels, Bio-Rad #4561085). 10uL Precision Plus Protein^{TM}

Kaleidoscope^{TM} Prestained Protein Standards (Bio-Rad, #1610375) or 10-15uL of

samples were loaded on to the gel. The gel was eletrophorised for 35-40mins at

200mV, in 1x running buffer (25mM Tris-Base, 190mM glycine, 0.1% SDS, pH 8.3).

Proteins were transferred onto a nitrocellulose membrane (Bio-Rad, Cat no. 162-

0115) at 400mA, 300V for 1 hour and 30 mins at 4°C, in 1x transfer buffer (25mM

Tris-Base, 190mM glycine, 20% methanol, pH 8.3). Primary antibodies (anti-FLAG

1:1000, Sigma Cat no. F1804-200UG, or anti-GAPDH1:1000, Millipore Cat no

MAB374) were diluted in 1% skim milk in PBS and incubated for 1 hour at RT with

shaking. After incubation, membranes were washed with 1xPBS+0.1%Tween for three

times 5 mins and once for 10 mins. After washing, membranes were incubated with

secondary antibody (1:5000 goat anti- mouse Ig-HRP, Sigma Cat no. A44161ML)

diluted in 1% skim milk in PBS, for 1 hour at RT with shaking, and then washed again

as above. Membrane was visualised under chemiluminescence with HRP substrate

(Millipore, Cat no. WBLUF0500) at various exposure times.

Analysis and Statistics:

Using ImageJ 1.05i software, Western blot membranes were analyzed for the mean

band intensity for anti-FLAG (corresponding to the expression of NgAgo), as well as

for the loading control anti-GAPDH. The relative abundance of anti-FLAG was

determined as a ratio of the loading control to produce a FLAG:GAPDH ratio. These

ratios were then normalized to the negative control, containing no NgAgo. These

values were then analysed as listed below.

A Two-Way ANOVA using Tukey's multiple comparisons test was performed to

quantify changes in protein expression over time for each treatment. Paired T-tests

were performed to compare protein expression over time between NgAgo samples

with and without gDNA. A two-tailed Wilcoxon test was performed to examine

changes in degradation between samples with or without gDNA. All analyses were

conducted using the GraphPad software, Prism 7, with significance at $P \leq 0.05$.

Acknowledgments: This work was supported from the National Collaborative

Research Infrastructure (NCRIS) through the Australian Phenomics Network in

Australia. The funders have no role in study design, data collection and analysis,

decision to publish or preparation of the manuscript. We would like to thank all the

comments, anonymous or not on the initial blogpost, on Twitter and/or on a Google

discussion group dedicated to genome editing technology that have led to stimulating,

interesting and fruitful discussions on this technique. Finally we would like to thanks 3

anonymous reviewers for their helpful comments that has improved the quality of the

manuscript.

Author contribution: Designed the experiments: NCK, JL, LJ and GB. Performed the

experiments NCK, JL, LJ. Analyzed the data: NCK, JL, LJ and GB. Wrote the

manuscript GB with the input from NCK, JL and LJ. All the authors read and

approved the manuscript.

References:

1. Komor AC, Badran AH, Liu DR (2016) CRISPR-Based Technologies for the
Manipulation of Eukaryotic Genomes. Cell.
2. Gao F, Shen XZ, Jiang F, Wu Y, Han C (2016) DNA-guided genome editing using
the Natronobacterium gregoryi Argonaute. Nat Biotechnol 34: 768-773.
3. Martinez-Galvez G, Ata H, Campbell JM, Ekker SC (2016) ssDNA and the
Argonautes: The Quest for the Next Golden Editor. Hum Gene Ther 27: 419-422.
4. Wang H, Yang H, Shivalila CS, Dawlaty MM, Cheng AW, et al. (2013) One-step
generation of mice carrying mutations in multiple genes by CRISPR/Cas-mediated
genome engineering. Cell 153: 910-918.
5. Burgess S, Cheng L, Gu F, Huang J, Huang Z, et al. (2016) Questions about
NgAgo. Protein Cell.
6. Qi J, Dong Z, Shi Y, Wang X, Qin Y, et al. (2016) NgAgo-based fabp11a gene
knockdown causes eye developmental defects in zebrafish. Cell Res.
7. Lee SH, Turchiano G, Ata A, Nowsheen S, Romito M, et al. (2016) Failure to
detect DNA-guided genome editing using Natronobacterium gregoryi Argonaute.
Nat Biotechnol.
8. Richardson L, Stevenson P, Venkataraman S, Yang Y, Burton N, et al. (2014)
EMAGE: Electronic Mouse Atlas of Gene Expression. Methods Mol Biol 1092: 61-
79.
9. Cong L, Ran FA, Cox D, Lin S, Barretto R, et al. (2013) Multiplex genome
engineering using CRISPR/Cas systems. Science 339: 819-823.

362   **Figure Legends:**

Figure 1: No evidence for double strand break cleavage and editing from NgAgo. A) DNA sequence indicating the locus targeted for the exon 26 of *Sptb*. The gDNA sequence is indicated in red. B) Gel electrophoresis (1.5%) of *Sptb* blastocysts (n=6) co-injected with nls-NgAgo-GK plasmid (2.5 ng/µl) and 2.5 ng/µl of gDNA. C57BL/6 DNA (B6) was also amplified as a control. The PCR product is 326 bp. C) T7 endonuclease 1 (T7E1) assay on the *Sptb* blastocysts indicating the absence of heteroduplexes suggesting indels in 6 blastocysts co-injected with nls-NgAgo-GK and gDNA. C57BL/6 (B6) non-edited control DNA was utilized as a negative control. A positive control in mouse zygotes edited with CRISPR/Cas9 was used as a positive control. The arrows indicate the presence of heteroduplexes suggesting a successful editing of the DNA. D) Representative chromatogram of one *Sptb* blastocyst (#5) suggesting no editing of the DNA under the DNA-guided NgAgo. E) Schematic diagram representing the flag-nls-NgAgo-GK plasmid. The expression of NgAgo is driven from a CMV promoter. A Flag tag was inserted in the 5' end of NgAgo sequence. Two Sv40 nuclear localization signals were inserted in the 3' end of the NgAgo sequence and in the 3' end of the flag tag. A Ploy A tail was appended to the sequence in the 3' end a Neomycin cassette was added in the 3' end of the plasmid sequence. F) DNA sequence indicating the targeting of the exon 3 from EMX1 human sequence. The gDNA sequence is indicated in red. G) Gel electrophoresis (2%) of the PCR for EMX1 in HEK293T cells at 8 and 12 hours post lipofection. The control samples were: The DNA without transfection, the lipofection reagent (LTX) and EMX1 DNA. The HEK293T cells were transfected with NgAgo alone, EMX1 gDNA

alone or co-transfected with NgAgo and EMX1 gDNA. A control DNA was

successfully edited with CRISPR/Cas9 (+ Cas9 control) and was utilized as a negative

control (- Cas9 control). The top electrophoresis gel respresents the PCR only

whereas the bottom gel represents the T7E1 assay. The arrows indicate the formation

of heteroduplexes for CRISPR/Cas9 genome editing. H) Western blot of NgAgo

protein production with and without the co-transfection of the gDNA from 8 to 48

hours post lipofection. The staining was performed with a monoclonal Flag anti-

antibody and anti-GAPDH anti-antibody. The top band represents NgAgo at 103KDa.

GAPDH was utilized as a Housekeeper gene. The exposure time was 30 seconds. The

controls were the lipefection agent alone (LTX), lane 1 and the gDNA EMX1 alone

(lane 2). The smears under the NgAgo band show the degradation of the protein

stained with the Flag tag.

409 **Table**

Gene	Microinjection mix	Zygotes injected	Blastocysts cultured	Targeted blastocysts (%)
Sptb	NgAgo 5 ng/μl + 2.5 ng/μl gDNA	8	8	0 (0%)
Sptb	NgAgo 5 ng/μl + 25 ng/μl gDNA	8	8	0 (0%)
Sptb	NgAgo 5 ng/μl + 50 ng/μl gDNA	6	6	0 (0%)
Sptb	NgAgo 10 ng/μl + 50 ng/μl gDNA	12	12	0 (0%)
Sptb	NgAgo 15 ng/μl + 50 ng/μl gDNA	15	15	0 (0%)
Tet-1	NgAgo 5 ng/μl + 25 ng/μl gDNA	10	10	0 (0%)
Tet-2	NgAgo 5 ng/μl + 2.5 ng/μl gDNA	7	7	0 (0%)
Tet-2	NgAgo 5 ng/μl + 25 ng/μl gDNA	7	7	0 (0%)
Tet-3	NgAgo 5 ng/μl + 2.5 ng/μl gDNA	13	13	0 (0%)
Sptb	50 ng/μl Cas9 + 0.6 pMol CrRNA & TracRNA	6	6	3(50%)

411 **Table 1**: Generation of edited mouse blastocyst using NgAgo and CRISPR/Cas9

412 genome editing technologies. NgAgo circular DNA was co-injected into the mouse

413 zygotes with various concentrations of gDNA. Cas9 Ribonucleoprotein (RNP) was

414 injected as controls for *Sptb* targeting the same genomic sequence as NgAgo.

Gene	Microinjection mix	Zygotes injected	Zygotes Trasferred	Newborn pups	Targeted pups (%)
Sptb	NgAgo 5 ng/μl + 50 ng/μl gDNA	68	43	9	0 (0%)
Sptb	NgAgo 5 ng/μl + 25 ng/μl gDNA	27	22	12	0 (0%)
Sptb	5 ng/μl Plasmid DNA px330	49	41	10	6(60%)
Tet2	NgAgo 5 ng/μl + 25 ng/μl gDNA	66	50	4	0 (0%)

418 **Table 2**: Generation of knockout mice using NgAgo and CRISPR/Cas9 genome

419 editing technologies. Various concentrations of gDNA were co-injected with NgAgo

420 plasmid or px330-U6-Chimeric_BB-CBh-hSpCas9 circular DNA as control.

Primer Name	Sequence
Sptb-F	5'-GCTACGTGACAAGTTCCGAGA-3'
Sptb-R	5'- GGTGGAAGGACTCAGCAGTG -3'
5'P-Sptb	5'-P-TGACACCCGCATGCAGCTGCT-3'
Tet1-F	5'-GTGTCAGGTTCAAGGCCATC-3'
Tet1-R	5'-ACGGGCGAGTTAGGGTTAAA-3'
5'P-Tet1	5'-P-GGCTGCTGTCAGGGAGCTCAT-3'
Tet2-F	5'-TGTCCAGCAGGATAAAGCAA-3'
Tet2-R	5'-ACCTGGATTGCATCCTTCAC-3'
5'P-Tet2	5'-P-GAAAGTGCCAACAGATATCCA-3'
Tet3-F	5'-GTGGAACAGGAGCAGAGGAG-3'
Tet3-R	5'-CCCGTGATGGTGAATGTCTA-3'
5'P-Tet3	5'-P-CAAGGAGGGGAAGAGTTCTCG-3'
EMX1-F	5'-GGGGCCCTAACCCTATGTA-3'
EMX1-R	5'-AGGGAGATTGGAGACACGGA-3'
5'P-EMX1	5'P-GAGTCCGAGCAGAAGAAGAAGGGC-3'

425   **Table 3**: List of oligonucleotides used in this study.

A

5' P-GGCTGCTGTCAGGGAGCTCAT
                |||||||||||||||||||||
5'-GGCAGGACCAAGTGTGGCTGCTGTCAGGGAGCTCATGGAGACTAGGTGAGGA- 3'
       |||||||||||||||| |||||||||||||||||
3'-CCGTCCTGGTTCACACCGACGACAGTCCCTCGAGTACCTCTGATCCACTCCT- 5'

B

5' P-GAAAGTGCCAACAGATATCCA
               |||||||||||||||||||||
5'-GTCACAGAAACACGTGAAAGTGCCAACAGATATCCAGGCTGCAGAATCGGAG- 3'
      |||||||||||||||| ||||||||||||||
3'-CAGTGTCTTTGTGCACTTTCACGGTTGTCTATAGGTCCGACGTCTTAGCCTC- 5'

C

5' P-CAAGGAGGGGAAGATTCTGGA
               |||||||||||||||||||||
5'-GGTCATCTACACGGGCAAGGAGGGGAAGATTCTGGAGGCTGTCCCATCGCCA- 3'
      |||||||||||||| ||||||||||||||||
3'-CCAGTAGATGTGCCCGTTCCTCCCCTTCTAAGACCTCCGACAGGGTAGCGGT- 5'

Khin et al. Supplementary Figure 1

A

-ve
LTX
NgAgo
EMX1
NgAgo + EMX1
-ve
LTX
NgAgo
EMX1
NgAgo + EMX1
EMX1
Control + Cas9
Control

24Hrs

48Hrs

B

Negative control
LTX
NgAgo
EMX1
NgAgo + EMX1
Negative control
LTX
NgAgo
EMX1
NgAgo + EMX1
EMX1
Control + Cas9
Control

24Hrs

48Hrs

Khin et al. Supplementary Figure 2