1

Colonization and transmission of the gut microbiota of the burying beetle, *Nicrophorus vespilloides*, through development

Yin Wang and Daniel E. Rozen*

Institute of Biology, Leiden University, Leiden, The Netherlands

* corresponding author: d.e.rozen@biology.leidenuniv.nl

Abstract:

Carrion beetles in the genus *Nicrophorus* rear their offspring on decomposing

carcasses where larvae are exposed to a diverse microbiome of decomposer bacteria.

Parents coat the carcass with antimicrobial secretions prior to egg hatch (defined as

Pre-Hatch care) and also feed regurgitated food, and potentially bacteria, to larvae

throughout development (defined as Full care). Here we partition the roles of pre- and

post-hatch parental care in the transmission and persistence of culturable symbiotic

bacteria to larvae. Using three treatment groups (Full-Care, Pre-Hatch care only, and

No Care), we found that larvae receiving Full-Care are predominantly colonized by

bacteria resident in the maternal gut, while larvae receiving No Care are colonized

exclusively with bacteria from the carcass. More importantly, larvae receiving only

Pre-Hatch care were also predominantly colonized by maternal bacteria; this result

indicates that parental treatment of the carcass, including application of bacteria to the

carcass surface, is sufficient to ensure symbiont transfer even in the absence of direct

larval feeding. Later in development, we found striking evidence that pupae undergo a

aposymbiotic stage, after which they are recolonized at eclosion with bacteria shed in

the moulted larval cuticle and on the wall of the pupal chamber. Our results clarify the

importance of pre-hatch parental care for symbiont transmission in *Nicrophorus*

vespilloides, and suggest that these bacteria successfully outcompete decomposer

bacteria during larval and pupal gut colonization.

Key words: Nicrophorus, parental care, symbiosis, microbiota, transmission

Importance

Here we examine the origin and persistence of the culturable gut microbiota of larvae in the burying beetle *Nicrophorus vespilloides*. This insect is particularly interesting for this study because larvae are reared on decomposing vertebrate carcasses where they are exposed to high-densities of carrion-decomposing microbes. Larvae also receive extensive parental care in the form of carcass preservation and direct larval feeding. We find that parents transmit their gut bacteria to larvae both directly, through regurgitation, and indirectly via their effects on the carcass. In addition, we find that larvae become aposymbiotic during pupation, but are recolonized from bacteria shed onto the insect cuticle before adult eclosion. Our results highlight the diverse interactions between insect behavior and development on microbiota composition. They further indicate that strong competitive interactions mediate the bacterial composition of *Nicrophorus* larvae, suggesting that the bacterial communities of these insects may be highly coevolved with their host species.

Introduction

Animals are colonized by a diverse community of bacterial symbionts that play
crucial roles in their ecology and evolution [1, 2]. This has been especially well
studied in insects, whose bacterial symbionts can influence traits ranging from mate
and diet choice [3] to susceptibility to natural enemies [4]. Bacterial symbionts can
also differ in the fidelity of their associations with their insect hosts. Endosymbionts
like Buchnera in aphids, that serve obligate functions for their insect hosts by
overcoming host nutritional deficiencies, are highly specific and have been aphid-
associated for millions of years [5]. At the opposite extreme, insects can retain
transient associations with bacteria whose effects are more variable [6, 7, 8, 9].
Although different factors may underlie the divergent influences of bacterial
symbionts on insect hosts, one key component is the way that bacteria are transmitted
between insect generations [10, 11]. Whereas obligate symbionts are almost
exclusively transmitted vertically, often via direct passage through eggs, more
transient associations, typical of the gut microbiota, involve an external stage where
bacteria are reacquired horizontally each generation via ingestion [12, 13].

Distinguishing symbionts on the basis of transmission mode (vertical versus
horizontal) has been extremely useful by focusing attention on how this can align the
fitness interests of symbionts and hosts [14, 15]. However, many associations
between insects and their microbial symbionts fall somewhere in the middle of these
strict extremes. Among diverse possibilities, trophallaxis and coprophagy occurs
when bacteria are passed horizontally between individuals via oral-oral/anal contact
or fecal consumption [16, 17, 18]. Similarly, horizontal symbiont transmission can
take place via ingestion of the bacteria-smeared egg-coat or via consumption of
bacteria-rich capsules [19, 20]. While these methods of transfer can effectively
vertically transmit symbionts from parent to offspring [12], the presence of an
environmental component implies that young and developing insects can be
simultaneously colonized by beneficial symbionts as well as environmental bacteria
that can harm the host [20, 21]. In these cases, establishment of the inherited
microbiota will be dependent on the ability for inherited symbionts to competitively
exclude environmental bacteria, as well as the timing and manner of their acquisition
[22, 23]. Additionally, especially for holometabolous insects that undergo a complete

metamorphosis, the manner of acquisition can change markedly throughout

development, at one stage occurring from the mother while at later stages through

alternative transmission routes [24].

Here, we examine the mechanisms of transmission and stability of the culturable gut

microbiota of the carrion beetle, *Nicrophorus vespilloides*, throughout its

development. This system is particularly interesting for addressing these questions

given the peculiar life-history of these organisms. Nicrophorus beetles are reared on

decomposing carrion where they encounter and ingest high densities of microbes [25].

Eggs are laid in the soil in close proximity to the carcass [26]. Upon hatching larvae

migrate to the carcass where they both self-feed and are fed regurgitated material

from the caring parents [27, 28]. Next, following an ~ 6-7 day feeding period upon

the carcass, larvae cease feeding and disperse into the surrounding environment where

they eventually pupate individually in underground chambers. Finally, pupae eclose

into adults, and emerge from the pupal chambers to commence feeding [29, 30].

N. vespilloides larvae may be colonized by a varied microbiota throughout

development [25, 31, 32], and this will likely be influenced by both the presence of

parents and the stage of development [25, 33]. First, parents may modify the carcass

microbiota by coating it in antimicrobial secretions throughout the period of parental

care [21, 34, 35]. Notably, these secretions are not sterile and contain significant

numbers of bacteria that can proliferate on the carcass. Secondly, parents feed larvae

with regurgitated food which may facilitate the transfer of the parental gut microbiota

to offspring (Post-hatch care) [36]. Finally, following dispersal, larvae cease feeding,

thereby preventing continued colonization from diet-borne bacteria; and then during

metamorphosis they shed the larval gut [37, 38]. At present, there is no understanding

of the dynamics of these gut bacterial communities through time.

There is little knowledge of the colonization dynamics of Nicrophorus gut bacteria or

the extent to which colonization is influenced by parental care, a hallmark of this

system. To examine these questions we manipulated *N. vespilloides* parental care and

used a culture-based approach to monitor the dynamics of symbiont colonization and

stability through development. Although culturing can underestimate bacterial

densities when compared to total cell counts or sequence-based approaches (see

Supplemental Figure 2), this approach allowed us to examine the largest set of

experimental conditions, while also identifying the bacterial groups that can be

experimentally manipulated to understand mechanisms of colonization and

community assembly of the microbiota using the *Nicrophorus* model system. Briefly,

our results provide strong evidence that beetle parents play a defining role of the

establishment of the bacteria residing in *Nicrophorus* larval guts; however, continuous

parental care and feeding is not essential for the stable maintenance of this microbiota.

Most strikingly, we also find that pupae undergo a sterile aposymbiotic stage, after

which they are recolonized from the bacteria shed into the pupal chamber. We discuss

these results in the context of the role of the *Nicrophorus* microbiota for beetle fitness.

Methods

General procedures

Experimental beetles were taken from an outbred laboratory population derived

from wild-caught *N. vespilloides* individuals trapped in Warmond near Leiden in The

Netherlands, between May and June 2014. Beetles were maintained in the laboratory at

20°C with a 15:9 hour light:dark cycle. All adults were fed fresh chicken liver twice

weekly. To generate outcrossed broods, non-sibling pairs of beetles were allowed to

mate for 24 hours in small plastic containers with soil. Next the mated pair were

provided with a freshly thawed mouse carcass weighing 24-26 g in a 15cm × 10cm

plastic box filled with approximately 1-2 cm of moist soil. Although fresh carcasses

may differ in bacterial composition from aged carcasses [39], our use fresh carcasses in

this study ensured higher brood success and is consistent with recent data showing that

most mouse carcasses are discovered shortly after they are placed in experimental

forests [40]   Broods were reared in sterile soil until the point of larval dispersal from

the carcass, after which larvae were transferred to new boxes for pupation with

unsterilized peat soil to complete development. Soil was sterilized using two autoclave

cycles at 121°C for 30 minutes, with a cooling interval between cycles.

Maternal care manipulation

To examine the role of parental care on the acquisition and composition of beetle gut

bacteria, we reared larvae under three treatment conditions that modified the degree of

parental care they received [25]: 1) Full Care (FC) broods experienced complete

parental care, including pre- and post-hatch care; 2) Pre-hatch parental care (PPC)

broods were reared on a carcass that had been prepared by the female, after which she

was removed prior to the hatch/arrival of larvae; and 3) no-care (NC) broods

experienced neither pre- nor post-hatch care. Broods in all treatments were initiated

similarly. Mated females were provided with a fresh carcass and induced to lay eggs.

Eggs were collected and surface sterilized within 12-24 hours and these were then

used to generate replicate broods of 15-20 larvae each. Females remained with their

prepared carcasses in FC broods, while females were removed prior to reintroducing

larvae in the PPC broods. NC larvae were provided with a freshly thawed carcass with

a sterile incision in the abdomen to permit larval entry.

Bacterial density and composition throughout development

We examined the dynamics of *N. vespiloides* intestinal microbiota through time

by destructively sampling beetles throughout development. To quantify gut bacterial

CFU, the whole intestinal tract from each beetle from independent broods (n = 3 at

each time point) was carefully removed with fine forceps and suspended in 0.7 ml

sterile sodium phosphate buffer (PBS; 100 mM; pH 7.2). The inner contents of pupa

were examined in their entirety owing to the absence of a clear gut at this stage.

Individual gut/pupal contents were serially diluted in PBS and plated on 1/3 strength

Tryptic Soy Broth agar and incubated at 30°C. In experiments shown in Supplemental

Figure 1, we also directly compared bacterial densities determined from total

microscopic counts versus via plating. Although plating for CFU consistently

underestimates bacterial densities, this approach recovered up to 60% of total counts

and the dynamics of bacterial densities perfectly mirror those based on total counts.

The composition of the maternal microbiota was characterized from n = 3 mated

females.

At each time point from each treatment, we isolated random colonies (n ≥ 100) from

individual beetles to analyze for species identification using MALDI-TOF Mass

Spectrometry (Matrix Assisted Laser Desorption Ionization-Time of Flight) with the

Biotyper platform (Bruker Daltonic GmbH). By generating unique whole-cell protein-

based fingerprints for each colony, the Biotyper permits highly reproducible

identification of bacterial colonies to the genus or species level. Because of its

reproducibility, ease of use and cost effectiveness, the Biotyper is used extensively in

clinical and public health microbiological laboratories [41] and is finding increased

use in ecological studies [42, 43, 44]. To standardize growth prior to analysis,

individual colonies were tooth-picked onto a 1/3 TS plate and grown overnight.

Colonies were then transferred directly to a 96-well steel MALDI-TOF target plate

and coated with 1 μl of alpha-cyano-4-hydroxy cinamic acid (HCCA) matrix

comprised of Acetonitrile (50%), Trifluoroacetic acid (2.5%) and water (47.5%), and

dried at room temperature. The target plate was subsequently inserted into the

Biotyper system for analysis. Next, mass spectrometry was carried out using the

MALDI Biotyper RTC (Realtime classification) and analyzed using Biotyper 3.0

(Bruker DAltonic GmbH). Spectra were collected under the linear positive mode in

the mass range of 3 to 20 kDa and a sample rate of 0.5 GS/s (laser frequency, 60 Hz;

ion source 1 voltage, 20.08 kV; ion source 2 voltage, 18.6 kV; lens voltage, 7.83 kV).

The Bruker bacterial test standard (BTS 8255343) was measured for standardization

of MALDI calibration before the specimens were processed. Spectra were compared

to the reference library provided by Bruker which identified 62.3% of the colonies to

species level overall using a stringent cut-off of 1.699, below which indicated no

reliable identification (in the Bruker library) [45, 46]. To confirm these assignments

and to establish the identity of colonies whose spectra were not included in the Bruker

database, all unique MS spectra (including both those with positive hits and those not

present in the Biotyper database) were subsequently analyzed using 16s rDNA

sequencing. Colony PCR using primers 27F (5'AGAGTTTGATCCTGGCTCAG-3')

and 1492R (5'-GGTTACCTTGTTACGACTT-3') was used for bacterial 16s rRNA

gene amplification [47]. The PCR cycling conditions were as follow: 95 °C for 5 min,

then 34 cycles of 95°C for 45 s, 55°C for 45 s, 72°C for 1 min. PCR products were

directly sequenced via the DNA Markerpoint in Leiden and 16s sequences were

classified for bacterial taxonomy using a nucleotide BLAST against the NCBI

database. The Bruker database was manually updated to include new samples thus

obtained.

A second experiment was conducted to determine the source of bacterial re-

colonization following beetle pupation. Pupae were removed from their chambers and

both the inside of the chamber and the cuticle were swabbed with a sterile, moist,

cotton swab. The bacteria on the swab were resuspended in sterile water and serially

diluted onto 1/3 TS agar. Finally, soil from outside the pupal chamber was collected

and diluted into PBS and plated. Colonies were isolated and identified as above using

a combination of MALDI-TOF Biotyping and 16s rDNA sequencing. To exclude rare

or transient bacterial species, we established a minimum threshold frequency of 1%,

averaged over all sampling periods for each treatment set, prior to analysis of

community composition.

Statistical analysis

Bacterial CFU through time were analyzed using General Linear Models (GLM) with

time and treatment as factors. Community composition was analyzed using the Vegan

package in R [48]. Beta diversity among the different treatments was analyzed using

ANOSIM, which is based on a Bray-Curtis dissimilarity matrix [49, 50]. The

Dendrograms to examine community similarity were generated based on the matrix of

mean within-group and between-group distances and the R function *hclust* was used

for hierarchical clustering.

Results

Bacterial CFU vary through development and as a function of parental care

The CFU of intestinal bacteria was quantified throughout development for three

treatment groups corresponding to different levels of parental care. Following

hatching from sterile eggs, larvae from all treatments rapidly acquire high bacterial

densities within their guts. Bacterial densities vary significantly through time (GLM

analyses: $df = 10$, $P < 0.001$) and as a function of treatment (GLM: $df = 2$, $P = 0.006$)

and vary across nearly 6 orders of magnitude as a function of developmental stage.

These dynamics are insensitive to experimental methods, as estimates of density

based on total microscopic counts perfectly mirror those determined by plate counting

(Supplemental Figure 2). During larval feeding on the carcass, bacterial densities

increase in all treatments, reaching densities of $\sim 10^6$ / larva. By contrast, following

dispersal, bacterial populations precipitously decline until, during pupation, bacteria

were undetectable. Finally, as pupae eclose and reemerge from pupal chambers, they

reacquire a high-density bacterial population within their guts (Fig. 1). It is notable

that this recovery occurs prior to feeding and before emergence from the pupal

chamber, indicating that recolonization takes place from bacteria resident within the

pupal chamber itself.

Composition of *N. vespilloides* larval symbionts

Although bacterial densities differ across parental-care treatments there is broad

overlap in the dynamics of CFU change through time. Despite these similarities, the

composition and diversity (Table S1) of these communities may vary. To understand

these differences and to illuminate transmission dynamics from mothers to larvae, we

tracked community composition of gut bacteria within larvae throughout development

(Fig. 2) using MALDI-TOF Mass-Spectrometry and compared these to the maternal

samples. The maternal microbiota was dominated by four bacterial Genera that

together comprised > 65% of recovered CFU, including *Providencia*, *Morganella*,

Vagococcus and *Proteus*, with several other genera appearing in lower frequencies

(Fig. 2). We next examined genus level composition across the three larval treatment

groups. As anticipated if transmission occurs via parents, we observed significant

overlap in the bacterial communities of parental and larval gut communities from

larvae receiving parental care throughout development ($R_{\text{FC-g vs Mother}}$=0.277; P=0.028,

Table 1), as R values < 0.25 correspond to "barely separable" groups [51]. Equally,

although to a lesser degree there is concordance between the maternal microbiota and

those of larvae receiving pre-hatch care only ($R_{\text{PPC-g vs Mother}}$= 0.331; P = 0.066, Table

1). By contrast, larvae reared in the absence of parental care are highly diverged from

the parental microbiota ($R_{\text{NC-g vs Mother}}$= 1; P = 0.007, Table 1) (Figs 3A and 3B). In

particular, the gut community of NC larvae was shifted towards bacterial groups

likely acquired from either the soil or the carcass (Figs 2 and 3C), e.g. *Escherichia*

coli (23.5%), *Serratia* (20.4%) and *Staphylococcus* (19.2%).

In comparing the larval microbiota of the three treatment groups, ANOSIM analysis

illustrated clear differences between the treatment groups overall (Global test: R =

0.815, P = 0.001) and although there are differences between the FC and PPC larvae,

there is much greater similarity between the two groups with parental care ($R_{FC-g \, vs}$

$_{PPC-g}$ = 0.665; P = 0.001) compared to either care group and the no-care larvae ($R_{FC-g \, vs}$

$_{NC-g}$ = 0.956, P=0.001; $R_{PPC-g \, vs \, NC-g}$ = 0.994, P=0.001) (Table 1, Fig. S1). This is also

apparent in the Venn diagrams in Fig. 3A, focusing on presence/absence of specific

bacterial groups. Together, these results indicate that transmission of the beetle

microbiota occurs predominantly from parents to offspring. However, they also reveal

that continued replenishment of bacteria from parent to offspring via feeding is

unnecessary to establish the endogenous microbiota. Instead transmission can occur

indirectly via deposition of the maternal bacteria on to the carcass by the mother

during carcass preparation and subsequent colonization of larva via self-feeding.

Re-colonization of *N.vespilloides* symbionts

A striking result from these analyses is the aposymbiotic stage occurring during

pupation, followed by recolonization from within the pupal chamber. Notably, this

result based on CFU was further confirmed by direct microscopic counts

(Supplemental Figure 2). To assess the source of recolonization, we sampled bacterial

populations from the pupal cuticle and the wall of the pupal chamber, together with

samples from the bulk soil in which pupal chambers were constructed. Treatment

designations are as above, with the addition of subscripts corresponding to each

sampling site. For example, FC-g refers to samples taken from the guts of larvae

receiving Full Care, while FC-c represents samples from the cuticle and chamber wall

of these same larvae. These analyses showed that the *N. vespilloides* pupal cuticle and

chamber soil had very similar compositions (FC-g, FC-c: R = 0.32, P = 0.068; PPC-g,

PPC-c: R = 0.02, P = 0.052; NC-g, NC-c: R = 0.03, P= 0.0397 by Pairwise test of

ANOSIM, Table 1), and that these were diverged compared to the bulk soil (FC-c,

Soil: R = 0.89, P = 0.094; PPC-c, Soil: R = 1, P = 0.114; NC-c, Soil: R = 1, P = 0.099

by Pairwise test of ANOSIM, Table 1). Importantly, many bacterial genera

irrespective of treatment, were found in the pre-pupal gut and the cuticle but

infrequently or not at all in the soil. For example, the most common bacterial groups

in FC larvae contained *Providencia* (FC-g: 18.3% vs FC-c: 17.1%), *Morganella* (FC-

g: 10.0% vs FC-c: 8.7%), *Proteus* (FC-g: 14.0% vs FC-c: 3.9%), *Vagococcus* (FC-g:

7.0% vs FC-c: 6.1%), *Neisseria* (FC-g: 8.3% vs FC-c: 4.7%), and *Koukoulia* (FC-g:

13.2% vs FC-c: 8.9%), while these were absent from soil. Similarly, the most

abundant genera in NC beetles were only found in NC-g and NC-c: *Escherichia coli*

(NC-g: 23.5% vs NC-c: 23.8%), *Enterococcus* (NC-g: 18.5% vs NC-c: 18.8%) (Fig.

4A and 4B). These results indicate that the core components of previously colonized

gut bacteria can successfully recolonize the host intestinal system after the

aposymbiotic stage characteristic of pupation. Thus, although transmission and

recolonization to larvae occurs via the environment, the bacterial species that

recolonize the newly eclosing adult are highly biased towards bacterial species that

were already present in the pre-pupal gut and which were originally acquired from the

mother.

Discussion

Animal symbionts can be passed to offspring through different mechanisms that vary

in their reliability of transmission [10, 11]. While strict endosymbionts of animals are

typically transmitted vertically via eggs, other mechanisms that include an

environmental component may also reliably transmit bacteria between generations [14,

19]. Here, we examined the mechanisms of bacterial transmission from *Nicrophorus*

vespilloides mothers to offspring. *Nicrophorus* larvae are exposed to and consume

high densities of bacteria throughout their development on decomposing carrion [25].

In earlier studies we showed that parental care, including preservation of the carcass

through secretion of lysozyme and potentially other antimicrobials, is essential for

maintaining larval fitness [21]. Additionally, preliminary metagenomic analyses from

our own lab (unpublished data) and published studies from others [39] have found

that parental beetles significantly modify the bacterial composition of decomposing

carrion, thereby potentially influencing the bacteria that larvae are exposed to and

ingest.

To examine the influence of parental care on the transmission of bacteria from parents

to offspring, we manipulated the level of care parents provide to their larvae. With

full parental care, parents apply oral and anal secretions to the carcass both before

larvae hatch and throughout larval development [21, 32]; they also regurgitate food to

larvae during the first three to four days of development [27, 52]. As expected, given

the continuous direct and indirect exposure to parental bacteria, larvae in this

treatment were colonized predominantly with parental symbionts (Fig 2); importantly,

despite limitations associated with a CFU-based approach, we observed broad overlap

between the dominant bacterial species we cultured and those identified using

sequence-based approaches (e.g *Providencia, Morganella, Vagococcus, Proteus,*

Koukoulia, and *Serratia*) [53]. However, with this Full Care treatment alone, it could

not be determined if larvae require constant replenishment of the parental species for

these to be maintained in the larval gut [54]. One possibility, for example, is that the

dominant bacteria from the carcass could outcompete endogenous beetle bacteria

within the larval gut; this could be driven actively, if the bacteria on the carcass are

particularly good colonizers, or passively since larval exposure to carcass bacteria is

continuous [22, 23]. To address this question, we established broods that only

received pre-hatch care. In this treatment, parents have no direct exposure to larvae,

and can only influence larval exposure to bacteria indirectly through their influence

on the carcass. It is important to note that because eggs are sterile, transmission is also

prevented through this route [26]. As with the Full Care treatment, larvae receiving

only Pre-hatch care were also predominantly colonized by maternal bacteria (Figs 2

and 3). This was not due to an inability of bacteria from the gut to colonize larvae, as

larvae in the No-Care treatment were also colonized by a high-density bacterial

microbiota. Also, bacteria in the pre-hatch groups were partially colonized by carcass-

derived bacteria (Figs 2 and 3C), leading to higher bacterial diversity overall in this

group (Supplemental Table 1) and indicating the capacity for carcass-derived bacteria

to establish themselves within the larval gut. Rather, we interpret this result to

indicate that "endogenous" bacteria from the mother are able to outcompete the

carrion associated microbes. Furthermore, this effect is long-lasting and can persist

entirely in the absence of direct maternal feeding. Although this interpretation is

consistent with our data, this hypothesis will require experimental testing using the

culturable species we have now established in our collection of *N. vespilloides*

symbionts.

At present, we understand relatively few of the mechanisms used by parents to

manipulate the carcass bacteria. However, several factors are likely to be important.

First, when parents locate a carcass they strip it of fur, while simultaneously coating

the carcass surface with oral and anal secretions. The composition of these secretions

has only been partially characterized, but a key component is lysozyme, a broad-
spectrum antibacterial with greater specificity towards Gram-positive bacteria [32,
55]. Additionally, oral secretions contain bacteria that can serve as an inoculum to
feeding larvae (unpublished results). In addition to these behaviors, we have also
observed parents opening the carcass and removing the mouse gut, behaviors that
could potentially have a dramatic influence on larval bacterial exposure by
introducing oxygen that could bias the bacterial community towards aerobic species
or more simply by directly reducing the overall density of bacteria to which larvae are
exposed. Following gut removal, parents continue to coat the carcass in secretions and
then bury the balled up carrion underground [21, 29, 52], which could influence
moisture or temperature levels. Both behaviors could possibly bias the persisting
microbial species, and potentially in favor of species originally introduced by caring
parents. In addition, caring parents and their larvae may be exposed to different
bacterial numbers and composition as a function of carcass age, a factor that is known
to have a dramatic influence on larval fitness [25, 56]. Although much remains to be
determined of these processes, our results clarify the importance of more completely
understanding how parents influence both the bacteria on the carcass and how this, in
turn, affects larval microbiota establishment.
418
After larvae complete feeding, they migrate into the soil to pupate [29, 52]. Bacterial
numbers during this stage decline precipitously (Fig. 1 and Supplemental Figure 2), in
part due to the absence of feeding and also to the evacuation of the larval gut. In
addition, larvae in some metamorphosing insects undergo a pre-pupal molt which
would further reduce bacterial numbers [50, 57]. Regardless of the mechanisms, it
appears that *Nicrophorus* larvae become effectively sterile during pupation, an
outcome previously seen in several flies and mosquitoes [58, 59, 60]. It is possible
that host immunity facilitates pupal symbiont suppression during metamorphosis [57,
61, 62], as a decline of phagocytic haemocytes and an increasing phenoloxidase
activity were both detected in *Nicrophorus* pupa [63]. Following this aposymbiotic
state, bacterial densities are quickly recovered at eclosion with bacterial communities
that significantly overlap with those present prior to pupation (Figs 1 and 2). To
determine the source of recolonization, we sampled bacteria from the pupal molt as
well as the wall of the pupal chambers, and in both cases we observed striking
similarity to the microbiomes of earlier developmental stages. Interestingly, this was

true for all treatment groups, suggesting that there is no intrinsic bias to recolonization,

but rather that eclosing beetles are colonized by a subset of the bacterial species

present in the pupal chamber.

The larval gut of *N. vespilloides* thus appears to be colonized via a combination of

mechanisms that are dependent on the degree of parental care and the stage of

development. With complete parental care, parents transmit bacteria to larvae through

a combination of direct feeding and through an indirect effect mediated by the carcass

[36, 39]. At present, it remains unclear if this latter component is because

Nicrophorus symbionts outcompete the mouse carrion microbiota within the larval

gut, or if this occurs primarily on the carcass surface itself. However, the former

seems more likely given the vast differences in larval exposure to these two groups of

bacteria, and the fact that larvae in the pre-hatch group remained colonized by beetle

symbionts, despite lacking any direct exposure to parents (Fig. 2). It is tempting,

given the reliable mode of transmission from parents to larvae, to speculate about the

function of these sybmionts for *Nicrophorus* growth and development, particularly the

role of these bacteria in limiting infection from carrion-borne bacteria [39, 53].

However, this remains an active area of research that we will hope to address in future

publications. In addition, it will be important to supplement the present work with

more detailed analyses based upon sequencing [39, 53]. Although culture-based

methods play an essential role in unraveling the relationships between invertebrate

host sociality and their symbiont strain-level diversity [64], they are clearly

complementary to sequence-based methods that can recover bacterial groups that may

be difficult or impossible to culture in the laboratory. Our work clarifies the key links

between *Nicrophorus* social behavior and symbiont transmission. This is likely to

have parallels in other animal systems where parents invest in the care of offspring.

Acknowledgements

Funding for this work was provided by start-up funds from Leiden University to DR

and from a studentship from the China Scholarship Council to YW. We gratefully

acknowledge expert assistance with beetle maintenance from Kees Koops, and thank

Andres Arce and Chris Jacobs for their extremely helpful comments on an earlier

version of this manuscript. In addition, we gratefully acknowledge the helpful and

constructive comments from the Editor, Eric Stabb, and three anonymous reviewers

on an earlier submission of this manuscript.

References

1. Engela, P., Kwong, W. K., McFrederick, Q., Anderson, K. E., Barribeau, S. M., Chandler, J. A., Cornman, R. S., Dainat, J., de Miranda, J. R., Doublet, V., Emery, O., Evans, J. D., Farinelli, L., Flenniken, M. L., Granberg, F., Grasis, J. A., Gauthier, L., Hayer, J., Koch, H., Kocher, S., Martinson, V. G., Moran, N., Munoz-Torres, M., Newton, I., Paxton, R. J., Powell, E., Sadd, B. M., Schmid-Hempel, P., Schmid-Hempel, R., Song, S. J., Schwar, R. S., van Engelsdorp, D., and Dainat, B.. 2016 The Bee Microbiome: Impact on Bee Health and Model for Evolution and Ecology of Host-Microbe Interactions. *MBio* **7**, e02164–15. (doi:10.1128/mBio.02164-15)

2. Peterson, B. F. & Scharf, M. E. 2016 Lower Termite Associations with Microbes: Synergy, Protection, and Interplay. *Front. Microbiol.* **7**, 422. (doi:10.3389/fmicb.2016.00422)

3. Ezenwa, V. O., Gerardo, N. M., Inouye, D. W., Medina, M. & Xavier, J. B. 2012 Animal Behavior and the Microbiota. *Science (80-.).* **338**, 198–199. (doi:10.1126/science.1227412)

4. Sanders, D., Kehoe, R., van Veen, F. F., McLean, A., Godfray, H. C. J., Dicke, M., Gols, R. & Frago, E. 2016 Defensive insect symbiont leads to cascading extinctions and community collapse. *Ecol. Lett.* **19**, 789–799. (doi:10.1111/ele.12616)

5. Shigenobu, S., Watanabe, H., Hattori, M., Sakaki, Y. & Ishikawa, H. 2000 Genome sequence of the endocellular bacterial symbiont of aphids Buchnera sp. APS. *Nature* **407**, 81–86. (doi:10.1038/35024074)

6. Vallet-Gely, I., Lemaitre, B. & Boccard, F. 2008 Bacterial strategies to overcome insect defences. *Nat Rev Microbiol* **6**, 302–313. (doi:10.1038/nrmicro1870)

7. Bulla, L. a 1975 Bacteria as insect pathogens. *Annu. Rev. Microbiol.* **29**, 163–190. (doi:10.1146/annurev.mi.29.100175.001115)

8. Lund, M. B., Holmstrup, M., Lomstein, B. A., Damgaard, C. & Schramm, A. 2010 Beneficial effect of verminephrobacter nephridial symbionts on the fitness of the earthworm aporrectodea tuberculata. *Appl. Environ. Microbiol.* **76**, 4738–4743. (doi:10.1128/AEM.00108-10)

9. Douglas, A. E. 2014 Multiorganismal Insects: Diversity and Function of Resident Microorganisms. *Annu. Rev. Entomol.* , 1–18. (doi:10.1146/annurev-ento-010814-020822)

10. Bright, M. & Bulgheresi, S. 2010 A complex journey: transmission of microbial symbionts. *Nat. Rev. Microbiol.* **8**, 218–230. (doi:10.1038/nrmicro2262)

11. Baalen, M. Van 2000 Parent-to-offspring infection and the struggle for transmission. *Evol. Biol. host-parasite relationships ...*

12. Hosokawa, T., Hironaka, M., Inadomi, K., Mukai, H., Nikoh, N. & Fukatsu, T. 2013 Diverse Strategies for Vertical Symbiont Transmission among Subsocial Stinkbugs. *PLoS One* **8**, 4–11. (doi:10.1371/journal.pone.0065081)

13. Estes, A. M., Hearn, D. J., Snell-Rood, E. C., Feindler, M., Feeser, K., Abebe,

T., Dunning Hotopp, J. C. & Moczek, A. P. 2013 Brood ball-mediated
transmission of microbiome members in the dung beetle, Onthophagus taurus
(Coleoptera: Scarabaeidae). *PLoS One* **8**, 1–15.
(doi:10.1371/journal.pone.0079061)
14. Russell, J. a & Moran, N. a 2005 Horizontal Transfer of Bacterial Symbionts :
Heritability and Fitness Effects in a Novel Aphid Host Horizontal Transfer of
Bacterial Symbionts : Heritability and Fitness Effects in a Novel Aphid Host.
Appl. Environ. Microbiol. **71**, 7987–7994. (doi:10.1128/AEM.71.12.7987)
15. Drown, D. M., Zee, P. C., Brandvain, Y. & Wade, M. J. 2013 Evolution of
transmission mode in obligate symbionts. *Evol. Ecol. Res.* **15**, 43–59.
16. Dahbi, a, Hefetz, A., Cerda, X. & Lenoir, A. 1999 Trophallaxis mediates
uniformity of colony odor in Cataglyphis iberica ants (Hymenoptera,
Formicidae). *J Insect Behav* **12**, 559–567. (doi:10.1023/A:1020975009450)
17. Zimmer, M. & Topp, W. 2002 The role of coprophagy in nutrient release from
feces of phytophagous insects. *Soil Biol. Biochem.* **34**, 1093–1099.
(doi:10.1016/S0038-0717(02)00044-5)
18. Ott, B. M., Cruciger, M., Dacks, A. M. & Rio, R. V. M. 2014 Hitchhiking of
host biology by beneficial symbionts enhances transmission. *Sci. Rep.* **4**, 5825.
(doi:10.1038/srep05825)
19. Kikuchi, Y., Hosokawa, T. & Fukatsu, T. 2007 Insect-microbe mutualism
without vertical transmission: A stinkbug acquires a beneficial gut symbiont
from the environment every generation. *Appl. Environ. Microbiol.* **73**, 4308–
4316. (doi:10.1128/AEM.00067-07)
20. Salem, H., Florez, L., Gerardo, N. & Kaltenpoth, M. 2015 An out-of-body
experience: the extracellular dimension for the transmission of mutualistic
bacteria in insects. *Proc. R. Soc. B Biol. Sci.* **282**, 20142957–20142957.
(doi:10.1098/rspb.2014.2957)
21. Arce, A. N., Johnston, P. R., Smiseth, P. T. & Rozen, D. E. 2012 Mechanisms
and fitness effects of antibacterial defences in a carrion beetle. *J. Evol. Biol.* **25**,
930–937. (doi:10.1111/j.1420-9101.2012.02486.x)
22. Coyte, K. Z., Schluter, J. & Foster, K. R. 2015 The ecology of the microbiome:
Networks, competition, and stability. *Science (80-.).* **350**, 663–666.
(doi:10.1126/science.aad2602)
23. Devevey, G., Dang, T., Graves, C. J., Murray, S. & Brisson, D. 2015 First
arrived takes all: inhibitory priority effects dominate competition between co-
infecting Borrelia burgdorferi strains. *BMC Microbiol.* **15**, 381.
(doi:10.1186/s12866-015-0381-0)
24. Ahmed, M. Z., Breinholt, J. W. & Kawahara, A. Y. 2016 Evidence for
common horizontal transmission of Wolbachia among butterflies and moths.
BMC Evol. Biol. **16**, 118. (doi:10.1186/s12862-016-0660-x)
25. Rozen, D. E., Engelmoer, D. J. P. & Smiseth, P. T. 2008 Antimicrobial
strategies in burying beetles breeding on carrion. *Proc. Natl. Acad. Sci. U. S. A.*
105, 17890–17895. (doi:10.1073/pnas.0805403105)
26. Jacobs, C. G. C., Wang, Y., Vogel, H., Vilcinskas, A., van der Zee, M. &
Rozen, D. E. 2014 Egg survival is reduced by grave-soil microbes in the
carrion beetle, *Nicrophorus vespilloides*. *BMC Evol. Biol.* **14**, 208.
(doi:10.1186/s12862-014-0208-x)
27. Smiseth, P. T., Darwell, C. T. & Moore, A. J. 2003 Partial begging: an
empirical model for the early evolution of offspring signalling. *Proc. Biol. Sci.*
270, 1773–7. (doi:10.1098/rspb.2003.2444)

28.   Scott, M. P. 1989 Male parental care and reproductive success in the burying
beetle,Nicrophorus orbicollis. *J. Insect Behav.* **2**, 133–137.
(doi:10.1007/BF01053623)
29.   Scott, M. P. 1998 The ecology and behaviour of burying beetles. *Annu. Rev.*
Entomol. **43**, 595–618.
30.   Smith, R. J. 2002 Effect of larval body size on overwinter survival and
emerging adult size in the burying beetle, Nicrophorus investigator. *Can. J.*
Zool. **80**, 1588–1593. (doi:10.1139/z02-151)
31.   Janzen D.H. 1977 Why fruits rot, seeds mold, and meat spoils. *Am Nat.* **111**,
691-713.
32.   Hall, C. L., Wadsworth, N. K., Howard, D. R., Jennings, E. M., Farrell, L. D.,
Magnuson, T. S. & Smith, R. J. 2011 Inhibition of microorganisms on a carrion
breeding resource: the antimicrobial peptide activity of burying beetle
(Coleoptera: Silphidae) oral and anal secretions. *Environ. Entomol.* **40**, 669–78.
(doi:10.1603/EN10137)
33.   Anduaga, S. & Huerta, C. 2001 Effect of parental care on the duration of larval
development and offspring survival in *Nicrophorus mexicanus* Matthews
(Coleoptera: Silphidae). *Coleopts Bull.* **55**, 264–270. (doi:10.1649/0010-
065X(2001)055)
34.   Cotter, S. C. & Kilner, R. M. 2010 Sexual division of antibacterial resource
defence in breeding burying beetles, *Nicrophorus vespilloides*. *J. Anim. Ecol.*
79, 35–43. (doi:10.1111/j.1365-2656.2009.01593.x)
35.   Hoback, W. W., Bishop, A. A., Kroemer, J., Scalzitti, J. & Shaffer, J. J. 2004
Differences among antimicrobial properties of carrion beetle secretions reflect
phylogeny and ecology. *J. Chem. Ecol.* **30**, 719–729.
(doi:10.1023/B:JOEC.0000028427.53141.41)
36.   Eggert, A.-K., Reinking, M., Mu, J. K. & Ller, } 1998 Parental care improves
offspring survival and growth in burying beetles. *Anim. Behav* **55**, 97–107.
(doi:10.1006/anbe.1997.0588)
37.   Greenberg, B. 1969 Salmonella suppression by known populations of bacteria
in flies. *J. Bacteriol.* **99**, 629–635.
38.   Russell, V. & Dunn, P. E. 1996 Antibacterial Proteins in the Midgut of
Manduca sexta During Metamorphosis. *I. Insect Physiol* **42**, 65–71.
(doi:10.1016/0022-1910(95)00083-6)
39.   Duarte, A., Welch, M., Wagner, J. & Kilner, R. M. 2016 Privatization of a
breeding resource by the burying beetle *Nicrophorus vespilloides* is associated
with shifts in bacterial communities. **44**. (doi:dx.doi.org/10.1101/065326)
40.   TRUMBO, S. T. 2016 Fate of mouse carcasses in a Northern Woodland. *Ecol.*
Entomol. , 737–740. (doi:10.1111/een.12341)
41.   Spanu T, Posteraro B, Fiori B, D'Inzeo T, Campoli S, Ruggeri A, Tumbarello
M, Canu G, Trecarichi EM, Parisi G, Tronci M, Sanguinetti M, Fadda G. 2012
Direct MALDI-TOF mass spectrometry assay of blood culture broths for rapid
identification of Candida species causing bloodstream infections: An
observational study in two large microbiology laboratories. *J. Clin. Microbiol.*
50, 176–179. (doi:10.1128/JCM.05742-11)
42.   Rahi, P., Prakash, O. & Shouche, Y. S. 2016 Matrix-Assisted Laser
Desorption/Ionization Time-of-Flight Mass-Spectrometry (MALDI-TOF MS)
Based Microbial Identifications: Challenges and Scopes for Microbial
Ecologists. *Front. Microbiol.* **7**, 1359. (doi:10.3389/fmicb.2016.01359)
43.   Santos IC, Hildenbrand ZL, Schug KA, Krásný L, Hynek R, Hochel I,

Havlicek V, Lemr K, Schug KA, Fenselau C, Demirev PA, Lay JO, Dingle TC, Butler-Wu SM, Lay JO, Freiwald A, Sauer S, Carbonnelle E, Mesquita C, Bille E, Day N, Dauphin B, Beretti J-L, Ferroni A, Gutmann L, Nassif X, Fournier P-E, Drancourt M, Colson P, Rolain J-M, Scola B La, Raoult D, Murray PR, Croxatto A, Prod'hom G, Greub G, Belkum A van, Welker M, Erhard M, Chatellier S, Bedair M, Sumner LW, Lovecka P, Pacovska I, Stursa P, Vrchotova B, Kochankova L, Demnerova K, Emami K, Askari V, Ullrich M, Mohinudeen K, Anil AC, Khandeparker L, Burgess JG, Mesbahi E, Kopcakova A, Stramova Z, Kvasnova S, Godany A, Perhacova Z, Pristas P, Koubek J, Uhlik O, Jecna K, Junkova P, Vrkoslavova J, Lipov J, Kurzawowa V, Macek T, Mackova M, Ferreira L, Sanchez-Juanes F, Garcia-Fraile P, Rivas R, Mateos PF, Martinez-Molina E, Gonzalez-Buitrago JM, Velazquez E, Štursa P, Uhlík O, Kurzawová V, Koubek J, Ionescu M, Strohalm M, Lovecká P, MacEk T, MacKová M, Ruelle V, Moualij B El, Zorzi W, Ledent P, Pauw E De, Dieckmann R, Graeber I, Kaesler I, Szewzyk U, Döhren H von, Singhal N, Kumar M, Kanaujia PK, Virdi JS, Tan KE, Ellis BC, Lee R, Stamper PD, Zhang SX, Carroll KC, Singh O V., Wang Z, Russon L, Li L, Roser DC, Long SR, Deutschbauer AM, Chivian D, Arkin AP, Green BD, Keller M, Kelleher NL, Yates JR, Ruse NA, Cristian I, Chandramouli K, Qian P, Demirev PA, Fenselau C, Zhang Y, Fonslow BR, Shan B, Baek M, Yates JR, Bogdanov B, Smith RD, Johnson RS, Davis MT, Taylor JA, Patterson SD, Demirev PA, Feldman AB, Kowalski P, Lin JS, Warscheid B, Fenselau C, Maron PA, Ranjard L, Mougel C, Lemanceau P, Kurian D, Jansèn T, Mäenpää P, Kan J, Hanson TE, Ginter JM, Wang K, Chen F, Vriendt K De, Theunissen S, Carpentier W, Smet L De, Devreese B, Beeumen J Van, Heim S, Ferrer M, Heuer H, Regenhardt D, Nimtz M, Timmis KN, Lacerda CMR, Choe LH, Reardon KF, Sandrin TR, Goldstein JE, Schumaker S, Olive DM, Bean P, Emerson D, Agulto L, Liu H, Liu L, Giebel RA, Fredenberg W, Sandrin TR, Siegrist TJ, Anderson PD, Huen WH, Kleinheinz GT, McDermott CM, Sandrin TR, Angolini CFF, Pilau EJ, Lopes-Oliveira PF, Garcia INS, Gozzo FC, Oliveira VM de, Marsaioli AJ, Munoz R, López-López A, Urdiain M, Moore ERB, Rosselló-Móra R, Donohue MJ, Best JM, Smallwood AW, Kostich M, Rodgers M, Shoemaker JA, Eddabra R, Prévost G, Scheftel J-M, Stets MI, Pinto AS, Huergo LF, Souza EM de, Guimarães VF, Alves AC, Steffens MBR, Monteiro RA, Pedrosa FDO, Cruz LM, Ghyselinck J, Hoorde K Van, Hoste B, Heylen K, Vos P De, Vidali M, Uhlik O, Strejcek M, Junkova P, Sanda M, Hroudova M, Vlcek C, Mackova M, Macek T, Tomás-Gallardo L, Canosa I, Santero E, Camafeita E, Calvo E, López JA, Floriano B, Kim YH, Cho K, Yun SH, Kim JY, Kwon KH, Yoo JS, Kim S Il, Pessione E, Giuffrida MG, Prunotto L, Barello C, Mazzoli R, Fortunato D, Conti A, Giunta C, Wilmes P, Wexler M, Bond PL, Dettmer K, Aronov PA, Hammock BD, Mapelli V, Olsson L, Nielsen J, Edwards JL, Kennedy RT, Persson S, Sönksen CP, Frigaard NU, Cox RP, Roepstorff P, Miller M, Valentine N, Wunschel S, Wunschel D, Petersen C, Wahl K, Toh-boyo GM, Basile F. 2016 Applications of MALDI-TOF MS in environmental microbiology. *Analyst* **141**, 2827–2837. (doi:10.1039/C6AN00131A)

44.   Ferreira, L., Sánchez-Juanes, F., García-Fraile, P., Rivas, R., Mateos, P. F., Martínez-Molina, E., González-Buitrago, J. M. & Velázquez, E. 2011 MALDI-TOF mass spectrometry is a fast and reliable platform for identification and ecological studies of species from family rhizobiaceae. *PLoS One* **6**.

(doi:10.1371/journal.pone.0020223)

45.   He, Y., Li, H., Lu, X., Stratton, C. W. & Tang, Y. W. 2010 Mass spectrometry
biotyper system identifies enteric bacterial pathogens directly from colonies
grown on selective stool culture media. *J. Clin. Microbiol.* **48**, 3888–3892.
(doi:10.1128/JCM.01290-10)

46.   Angelakis E, Yasir M, Azhar EI, Papadioti A, Bibi F, Aburizaiza AS, Metidji
S, Memish ZA, Ashshi AM, Hassan AM, Harakeh S, Gautret P, Raoult D.
2014 MALDI-TOF mass spectrometry and identification of new bacteria
species in air samples from Makkah, Saudi Arabia. *BMC Res. Notes* **7**, 892.
(doi:10.1186/1756-0500-7-892)

47.   Hongoh, Y., Yuzawa, H., Ohkuma, M. & Kudo, T. 2003 Evaluation of primers
and PCR conditions for the analysis of 16S rRNA genes from a natural
environment. *FEMS Microbiol. Lett.* **221**, 299–304. (doi:10.1016/S0378-
1097(03)00218-0)

48.   Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB. 2015
Vegan: community ecology packageR package version 2.3-0. Available at:
https://github.com/vegandevs/vegan.

49.   Anderson, M. J. 2006 Distance-based tests for homogeneity of multivariate
dispersions. *Biometrics* **62**, 245–253. (doi:10.1111/j.1541-0420.2005.00440.x)

50.   Martinson, V. G., Moy, J. & Moran, N. A. 2012 Establishment of characteristic
gut bacteria during development of the honeybee worker. *Appl. Environ.*
Microbiol. **78**, 2830–2840. (doi:10.1128/AEM.07810-11)

51.   Ramette, A. 2007 Multivariate analyses in microbial ecology. *FEMS*
Microbiol. Ecol. **62**, 142–160. (doi:10.1111/j.1574-6941.2007.00375.x)

52.   Milne, L. J. & Milne, M. 1976 The Social Behavior of Burying Beetles. *Sci.*
Am. **235**, 84–89. (doi:10.1038/scientificamerican0876-84)

53.   Kaltenpoth, M. & Steiger, S. 2014 Unearthing carrion beetles' microbiome:
Characterization of bacterial and fungal hindgut communities across the
Silphidae. *Mol. Ecol.* **23**, 1251–1267. (doi:10.1111/mec.12469)

54.   Blum, J. E., Fischer, C. N., Miles, J. & Handelsman, J. 2013 Frequent
replenishment sustains the beneficial microbiome of Drosophila melanogaster.
MBio **4**, 1–8. (doi:10.1128/mBio.00860-13)

55.   Arce, A. N., Smiseth, P. T. & Rozen, D. E. 2013 Antimicrobial secretions and
social immunity in larval burying beetles, Nicrophorus vespilloides. *Anim.*
Behav. **86**, 741–745. (doi:10.1016/j.anbehav.2013.07.008)

56.   Trumbo, S. T., Sikes, D. S. & Philbrick, P. K. B. 2016 Parental care and
competition with microbes in carrion beetles: a study of ecological adaptation.
Anim. Behav. **118**, 47–54. (doi:10.1016/j.anbehav.2016.06.001)

57.   Kim, J. K., Han, S. H., Kim, C. H., Jo, Y. H., Futahashi, R., Kikuchi, Y.,
Fukatsu, T. & Lee, B. L. 2014 Molting-associated suppression of symbiont
population and up-regulation of antimicrobial activity in the midgut symbiotic
organ of the Riptortus-Burkholderia symbiosis. *Dev. Comp. Immunol.* **43**, 10–
14. (doi:10.1016/j.dci.2013.10.010)

58.   Bakula, M. 1969 The persistence of a microbial flora during
postembryogenesis of *Drosophila melanogaster. J. Invertebr. Pathol.* **14**, 365–
374. (doi:10.1016/0022-2011(69)90163-3)

59.   GREENBERG, B. 1959 Persistence of bacteria in the developmental stages of
the housefly. IV. Infectivity of the newly emerged adult. *Am. J. Trop. Med.*
Hyg. **8**, 618–22.

60.   Broderick, N. A. & Lemaitre, B. 2012 Gut-associated microbes of Drosophila

melanogaster. *Gut Microbes* **3**, 307–321. (doi:10.4161/gmic.19896)

61. Tryselius, Y., Samakovlis, C., Kimbrell, D. a & Hultmark, D. 1992 CecC, a
cecropin gene expressed during metamorphosis in Drosophila pupae. *Eur. J.*
Biochem. **204**, 395–399. (doi:10.1111/j.1432-1033.1992.tb16648.x)

62. Johnston, P. R. & Rolff, J. 2015 Host and Symbiont Jointly Control Gut
Microbiota during Complete Metamorphosis. *PLoS Pathog.* **11**, 1–11.
(doi:10.1371/journal.ppat.1005246)

63. Urbański, A., Czarniewska, E., Baraniak, E. & Rosiński, G. 2014
Developmental changes in cellular and humoral responses of the burying beetle
Nicrophorus vespilloides (Coleoptera, Silphidae). *J. Insect Physiol.* **60**, 98–
103. (doi:10.1016/j.jinsphys.2013.11.009)

64. Ellegaard, K. & Engel, P. 2016 Beyond 16S rRNA community profiling: intra-
species diversity in the gut microbiota. *Front. Microbiol.* **7**, 1475.
(doi:10.3389/FMICB.2016.01475)

744
745
746 **Figure 1 CFU of *Nicrophorus vespilloides* gut bacteria through development.**
747 Overview of the entire time course of beetle developmental and change in CFU of
748 host gut contents through time (means ± SD). FC corresponds to larval guts sample
749 from full parental care broods; PPC, to guts sample from pre-parental care broods;
750 and NC from larval guts sample from no-care broods.

Figure 2 Composition of *N. vespilloides* gut microbiota through development.
The maternal gut microbiota is shown at bottom, while treatment designations are the
same as in Figure 1. No CFUs were detectable between days 15-21 of larval
development, corresponding to the duration of pupation. Three individual larvae were
independently analyzed for each time point. Y-axis of day 1 to day 9 refers to larval
stage; day 12 corresponds to prepupal stages; day 24 to day 30 refers to adult
formation, respectively.

Figure 3: Frequencies of bacteria from gut communities across parental care treatments. **A)** Shared and unique genera between treatment groups. Strains with a minimum frequency of 1% were included. **B)** Hierarchical clustering on mean similarity of gut microbiota between treatment groups. **C)** Overall composition of gut communities across treatments. Strains with frequencies lower than 1% across all communities were excluded from plots.

(A)

(B)

Figure 4: Re-colonization of bacterial communities through pupations. **A)** Shared and unique genera among treatment groups. Subscripts correspond to the site of isolation; e.g. FC-g corresponds to gut samples, FC-c corresponds to the pupal carapace in the wall of the pupal chamber, and Soil corresponds to bulk soil outside the pupal chamber. **B)** Comparison of gut bacterial communities from each sample site and treatment. Strains with frequencies lower than 1% across all communities were excluded from plots.

868 **Table 1 ANOSIM analysis on bacterial community dissimilarity.** Subscripts
869 correspond to the site of isolation; e.g. FC-g corresponds to gut samples, FC-c
870 corresponds to the pupal carapace and the wall of the pupal chamber, and Soil
871 corresponds to bulk soil outside the pupal chamber.
872
873
874

Groups	R statistic	Significance (P value)	Number of permutations	Number of observed	Test model
FC-g, PPC-g, NC-g	0.8152	0.001	999	352	Global
FC-g, PPC-g, NC-g, Mother	0.741	0.001	999	264	Global
FC-c, PPC-c, NC-c, Soil	0.7493	0.001	999	814	Global
FC-g, NC-g	0.9556	0.001	999	144	Pairwise
PPC-g, NC-g	0.9939	0.001	999	64	Pairwise
FC-g, PPC-g	0.6651	0.001	999	144	Pairwise
FC-g, Mother	0.2769	0.028	999	24	Pairwise
PPC-g, Mother	0.3306	0.066	999	24	Pairwise
NC-g, Mother	1	0.007	999	24	Pairwise
FC-g, FC-c	0.3177	0.068	999	54	Pairwise
PPC-g, PPC-c	0.0242	0.052	999	54	Pairwise
NC-g, NC-c	0.0134	0.397	999	54	Pairwise
FC-c, Soil	0.8889	0.094	720	9	Pairwise
PPC-c, Soil	1	0.114	720	9	Pairwise
NC-c, Soil	1	0.099	720	9	Pairwise

887