Title: *C. elegans* detect the color of pigmented food sources to guide foraging decisions

Authors: D. Dipon Ghosh[1], Xin Jin[1], and Michael N. Nitabach[1,2,3]*

Affiliations:

[1]Department of Cellular and Molecular Physiology, Yale University, New Haven, CT

[2]Department of Genetics, Yale University, New Haven, CT

[3]Program in Cellular Neuroscience, Neurodegeneration and Repair, Yale University, New

Haven, CT

*Correspondence to: michael.nitabach@yale.edu

Abstract:

Here we establish that contrary to expectations, *Caenorhabditis elegans* nematode worms possess a color discrimination system despite lacking any opsin or other photoreceptor genes. We found that simulated daylight guides *C. elegans* foraging decisions with respect to harmful bacteria that secrete a blue pigment toxin. By absorbing yellow-orange light, this blue pigment toxin alters the color of light sensed by the worm, and thereby triggers an increase in avoidance of harmful bacteria. These studies thus establish the existence of a color detection system that is distinct from those of other animals. In addition, these studies reveal an unexpected contribution of microbial color display to visual ecology.

One-sentence summary:

Color detection contributes to *Caenorhabditis elegans* behavioral ecology by guiding foraging decisions.

 Main text:

C. elegans live in decomposing organic matter where they feed on microorganisms (*1-3*),

some of which secrete colorful pigments. While *C. elegans* lack any specialized photoreceptor

cells or opsin genes, they possess an illuminance sensing system that mediates rapid escape

responses to bright short-wavelength light (*4-6*). However, it is unknown whether *C. elegans* use

light information, potentially including color, to inform complex decisions like foraging in

environments containing colorful food sources. To address this question, we tested whether

white light alters foraging decisions on *P. aeruginosa* bacterial lawns containing the blue

pigment toxin pyocyanin, one of a number of small molecule phenazine toxins secreted by *P.*

aeruginosa (*7-10*). Moderate intensity indirect daylight is typically 10-20 kilolux intensity and

contains peaks in the blue and yellow-orange wavelength ranges (*11*). We selected an artificial

LED array white light source with color temperature of 6500 K and 8 kilolux intensity to mimic

these natural lighting conditions (**Fig. 1A,** see **fig. S1A**). Previous studies have shown that

foraging decisions to remain on or leave a bacterial lawn are guided by a variety of factors (*12,*

13). Worms remain on bacterial lawns that are easy to eat and support its growth, while leaving

lawns that are of poor nutritive quality, repulsive, or pathogenic (*12-17*). Worms are initially

attracted to *P. aeruginosa* lawns, but over a time course of hours, as the *P. aeruginosa* continues

to divide and secrete toxins, they respond to its increasingly pathogenic qualities and begin to

leave (*15, 16, 18-21*). Whether light plays a role in guiding avoidance of *P. aeruginosa* has never

been tested. Here we employed a standard lawn avoidance assay, placing worms on a *P.*

aeruginosa lawn in the center of an agar plate, and quantified avoidance as the fraction of worms

found off the lawn as a function of time (**Fig. 1A**). Consistent with prior studies, worms

gradually avoid *P. aeruginosa* strain PA14 over a span of many hours (*15, 16, 18-21*) (**Fig. 1A**).

Surprisingly, however, this avoidance is dramatically potentiated by simulated daylight (**Fig.**

1B). This light has no effect on worms on lawns of non-toxic *E. coli* OP50 food bacteria, which

they remain on without leaving (**Fig. 1B**).

Recent studies have implicated the *lite-1* gene – which encodes a non-canonical seven-

transmembrane domain protein with closest homology to insect olfactory and gustatory

chemoreceptors – in photophobic responses to bright short-wavelength light (*4-6*). To test

whether the *lite-1* pathway is involved in light-dependent potentiation of PA14 avoidance, we

compared avoidance of PA14 by *lite-1* null-mutant worms in the light and in the dark. Like wild-

type worms, *lite-1* null-mutant worms gradually avoid PA14 over time, but their avoidance is

unaffected by white light (**Fig. 1C**). These results indicate an essential role for the *lite-1* light

response pathway in light-mediated potentiation of PA14 avoidance.

In order to determine whether the secreted blue toxin pyocyanin is involved in light-

dependent potentiation of *P. aeruginosa* avoidance, we tested avoidance of a *phzM* mutant strain,

PA14ΔphzM. This strain lacks the biosynthetic enzyme necessary for pyocyanin synthesis, but

still synthesizes other phenazine toxins (*7-10, 22*), and PA14ΔphzM cultures are no longer blue

(**Fig. 1D**). Simulated daylight has only a minimal effect on avoidance of PA14ΔphzM by wild-

type worms, and no effect on avoidance by *lite-1* null-mutant worms (**Fig. 1, E and F**). The

phzM mutation has pleiotropic effects beyond absence of pyocyanin, which include, among other

things, increased production and secretion of other phenazines (*7*). The results thus far establish

light-dependent potentiation of PA14 avoidance requiring both the *lite-1* light response pathway

of the worm and the blue toxin pyocyanin secreted by *P. aeruginosa*.

To circumvent complexities associated with *P. aeruginosa* lawns, which continue to

divide and secrete phenazines over the course of the assay, we tested whether pyocyanin is

sufficient to mediate light-dependent potentiation of avoidance of otherwise non-toxic bacteria.

Accordingly, we employed lawns of non-toxic *E. coli* OP50 doped with pyocyanin (**Fig. 2A**).

Wild-type worms avoid OP50 doped with 2.5 mM pyocyanin within one hour of placing on the

lawn, but only in the presence of simulated daylight (**Fig. 2B**). This is more rapid than the

avoidance of PA14 (see **Fig. 1B**), and is already at the maximum avoidance reached during

extended exposure (**Fig. 2B**). This rapid, unchanging avoidance of OP50 lawns doped with 2.5

mM pyocyanin is likely due to the constant amount of pyocyanin present in the lawn, in contrast

to *P. aeruginosa* lawns that continue to secrete pyocyanin over the course of the assay. In light of

this unchanging avoidance of OP50 lawns doped with pyocyanin, for subsequent experiments we

relied on this one hour time point. Like avoidance of PA14, light-dependent avoidance of OP50

+ 2.5 mM pyocyanin is abolished by the *lite-1* null mutation (**Fig. 2C**). While doping with 0.25

mM pyocyanin fails to elicit avoidance, worms avoid OP50 + 5 mM pyocyanin both in the light

and dark, and independently of *lite-1* function (**Fig. 2C**). These results indicate that an

intermediate concentration of pyocyanin confers light-dependent and *lite-1*-dependent rapid

avoidance of otherwise innocuous food bacteria, with higher concentrations conferring

avoidance independent of light or *lite-1*.

lite-1 is expressed in a number of neurons in the worm nervous system, including the

primary sensory neurons ASJ, ASI, and ASK (*5, 6, 23*). ASJ and ASI have been previously

implicated in long-term avoidance of PA14 mediated by their direct responses to phenazines

with Ca^{2+} increases and gene activation (*16*). In addition, *lite-1* expression in ASJ and ASK is

sufficient for escape responses to short-wavelength light (*5, 6*). To determine the cellular locus of

lite-1 expression responsible for rapid light-dependent avoidance of OP50 + 2.5 mM pyocyanin,

we employed *lite-1* null-mutant worms with targeted re-expression of *lite-1* in neurons ASJ,

ASH/ASI/PVQ, or ASK. Restoration of *lite-1* expression in ASJ or ASH/ASI/PVQ are each

sufficient to rescue rapid light-dependent avoidance of OP50 + 2.5 mM pyocyanin, while

restoration in ASK is not (**Fig. 2D**). These results localize *lite-1* function in rapid light- and

pyocyanin-dependent bacteria avoidance to a small number of primary sensory neurons also

involved in long-term avoidance of PA14, and which only partially overlap with those

responsible for escape responses to short-wavelength light (**Fig. 2E**).

To test whether the blue color of pyocyanin independent of its toxic chemistry is

sufficient to induce rapid avoidance of non-toxic OP50 lawns, we employed OP50 lawns doped

with non-toxic inert blue food dye (**Fig. 3A**). Neither wild-type nor *lite-1* null-mutant worms

avoid OP50 lawns doped with blue food dye, whether in dark or light, indicating that it is not

solely the blue color of pyocyanin that drives light-potentiated rapid avoidance but also its

chemical reactivity (**Fig. 3B**). One of the chemical features of pyocyanin that confers toxicity is

that it enters eukaryotic cells and generates reactive oxygen species (ROS) through various

mechanisms (*7-10, 22*). To determine if pyocyanin's combination of blue color and ROS-

generating toxic chemistry underlies light-potentiated avoidance, we employed OP50 lawns

doped with the colorless ROS-generating toxin paraquat (*24*) and non-toxic inert blue food dye

(**Fig. 3C**). Wild-type worms rapidly avoid OP50 doped with 30 mM paraquat and blue dye, but

only in the presence of light (**Fig. 3D**). This light-potentiated avoidance is abolished in *lite-1*

null-mutant worms (**Fig. 3D**). As with OP50 + pyocyanin, higher concentrations of paraquat

mediate rapid avoidance independent of both incident light and *lite-1*, while lower concentrations

of paraquat with blue dye are insufficient to mediate avoidance (**Fig. 3D**). Doping OP50 lawns

with 30 mM paraquat without any dye or with inert non-toxic red dye (**Fig. 3D**) are each

insufficient to confer light-potentiated avoidance (**Fig. 3F**). These results indicate that rapid

light- and *lite-1*-dependent avoidance of pyocyanin-containing bacterial lawns relies both on its

chemical reactivity as a generator of ROS and on its blue color. They also indicate that

avoidance conferred by higher concentrations of pyocyanin relies solely on its ROS-generating

capacity and is independent of *lite-1* function. This integration of color and chemical information

to guide avoidance of food sources could enable more accurate discrimination of toxic from non-

toxic lawns.

We hypothesized that blue pigment confers rapid light-potentiated avoidance of bacterial

lawns containing ROS-generating toxins by absorbing long-wavelength light and thereby altering

the spectral composition of light sensed by the worm. To test this possibility, we employed a

series of shortpass and longpass optical filters to alter the spectra of incident light (corresponding

photographs and spectra of filtered light are shown in **fig. S2**). Consistent with the previously

determined action spectrum for *lite-1*-dependent photophobic responses (*4-6*), longpass filtered

light lacking blue light fails to potentiate avoidance of OP50 + 2.5 mM pyocyanin (**Fig. 4A**).

Surprisingly, however, light shortpass filtered with cut-offs of <500 nm or <550 nm also failed to

potentiate avoidance (**Fig. 4A**). Only light shortpass filtered with <600 nm cut-off, which still

includes the yellow-orange peak of 6500 K simulated daylight, was sufficient to potentiate

avoidance of OP50 + 2.5 mM pyocyanin (**Fig. 4A**). This establishes that light-potentiated

avoidance of bacterial lawns containing ROS-generating toxins and blue pigment requires not

only short-wavelength blue light, but also long-wavelength yellow-orange light. This suggests

the existence of a yellow-orange sensing pathway in addition to the *lite-1*-dependent blue light

pathway.

Completely eliminating yellow-orange light abolishes light-dependent potentiation of

avoidance of OP50 + 2.5 mM pyocyanin (**Fig. 4A**). Thus, we hypothesized that the presence of

blue pigment in the lawn reduces, but without eliminating, the yellow-orange content of light

sensed by the worm, and that it is this decreased ratio of yellow-orange to blue light that

potentiates avoidance of bacterial lawns containing ROS-generating toxins. It is uncertain

exactly how blue pigment in its microenvironment alters the spectrum of light sensed by the

worm. To circumvent this uncertainty, we eliminated blue pigment from the lawn and directly

modified the spectral composition of light sensed by the worm with a "blue vinyl" filter that

reduces without eliminating yellow-orange content to mimic the hypothesized effect of blue

pigment (**Fig. 4B, fig. S2, A and B**). Remarkably, simulated daylight filtered through this blue

vinyl filter recapitulates white light-potentiated rapid avoidance of OP50 + 30 mM paraquat, but

now in the absence of blue pigment in the lawn (**Fig. 4C**). These results indicate that blue

pigments confer light-potentiated avoidance of toxic bacterial lawns by absorbing yellow-orange

light and thereby altering the spectral composition of light sensed by the worm. They also imply

the existence of a yellow-orange light-sensing pathway that is required to be activated to

potentiate avoidance of toxic lawns, but with activation of this pathway beyond some threshold

preventing such potentiation.

As a further test of the hypothesis that it is the altered color of light, specifically the blue-

to-yellow-orange ratio, that potentiates avoidance and not altered illuminance, we altered the

intensity of simulated daylight (**fig. S3A**). Increasing white light intensity by ~3-fold to 25 klx

failed to increase avoidance of OP50 + 30 mM paraquat (**fig. S3B**), supporting the hypothesis

that it is the blue-to-yellow-orange ratio, and not illuminance, driving light-potentiated

avoidance. While further increases to 50 klx or 100 klx did mediate increased avoidance of OP50

+ 30 mM paraquat, these very high light intensities closer to that of direct sunlight (100-120

kilolux) identically increase avoidance of OP50 lawns without paraquat, most likely by

triggering *lite*-1-dependent escape responses (**fig. S3B**). Since *lite-1*-dependent responses to very

bright light may themselves involve ROS (*23*), we used H2DCFDA, a fluorescent ROS indicator,

to determine if intensities of light used in our experiments induce detectable ROS production in

worms. No ROS signal is detected in worms placed on OP50 lawns doped with 30 mM paraquat

in the dark or exposed to 8 klx white light for one hour (**fig. S3, C and D**). In contrast,

substantial ROS signal is detected in worms on OP50 lawns exposed to 100 klx white light for

one hour, regardless of whether paraquat is present (**fig. S3, C and D**). These results further

support the conclusion that avoidance of toxic bacterial lawns potentiated by moderate intensity

light operates by a different mechanism than photophobic responses to brighter light. They also

rule out the possibility that blue pigment potentiates avoidance of bacterial lawns containing

ROS-generating toxins by scattering blue light and thereby increasing activation of the *lite-1*

short-wavelength light-sensing pathway.

Taken together, our results thus establish that *C. elegans* worms are capable of

discriminating color in their environment by comparing activation of blue- and yellow-orange-

sensitive pathways to guide foraging decisions on colorful toxic food sources. This reveals an

unexpected mechanism by which color detection contributes to *C. elegans* behavioral ecology by

heightening vigilance to ROS and thereby guiding foraging decisions on toxic bacteria (**Fig. 4D**).

To our knowledge this is the first demonstration of the effect of microbial color display on an

animal's foraging decision. Another example of color display by potential food sources

influencing foraging decisions can be found in butterflies. Some non-toxic butterfly species

mimic the coloration pattern of toxic butterflies in the same ecological niche to ward off

predation (*25*). While selective pressure may have driven the worm's association between blue

color and toxic food sources, blue color itself is insufficient to drive avoidance (see **Fig. 3, A**

and B). This raises the possibility that other non-toxic bacterial species may have co-opted blue

coloration to ward off being eaten, forcing worms to be more vigilant in the presence of blue

pigments while simultaneously considering information beyond color alone to avoid being

tricked by mimics.

References:

1. M. A. Felix, F. Duveau, Population dynamics and habitat sharing of natural populations of Caenorhabditis elegans and C. briggsae. *BMC Biol* **10**, 59 (2012).

2. L. Frezal, M. A. Felix, C. elegans outside the Petri dish. *Elife* **4**, (2015).

3. B. S. Samuel, H. Rowedder, C. Braendle, M. A. Felix, G. Ruvkun, Caenorhabditis elegans responses to bacteria from its natural habitats. *Proceedings of the National Academy of Sciences of the United States of America*, (2016).

4. S. L. Edwards *et al.*, A novel molecular solution for ultraviolet light detection in Caenorhabditis elegans. *PLoS biology* **6**, e198 (2008).

5. J. Liu *et al.*, C. elegans phototransduction requires a G protein-dependent cGMP pathway and a taste receptor homolog. *Nature neuroscience* **13**, 715-722 (2010).

6. A. Ward, J. Liu, Z. Feng, X. Z. Xu, Light-sensitive neurons and channels mediate phototaxis in C. elegans. *Nature neuroscience* **11**, 916-922 (2008).

7. B. Cezairliyan *et al.*, Identification of Pseudomonas aeruginosa phenazines that kill Caenorhabditis elegans. *PLoS Pathog* **9**, e1003101 (2013).

8. S. Mahajan-Miklos, M. W. Tan, L. G. Rahme, F. M. Ausubel, Molecular mechanisms of bacterial virulence elucidated using a Pseudomonas aeruginosa-Caenorhabditis elegans pathogenesis model. *Cell* **96**, 47-56 (1999).

9. M. W. Tan, S. Mahajan-Miklos, F. M. Ausubel, Killing of Caenorhabditis elegans by Pseudomonas aeruginosa used to model mammalian bacterial pathogenesis. *Proceedings of the National Academy of Sciences of the United States of America* **96**, 715-720 (1999).

10. M. W. Tan, L. G. Rahme, J. A. Sternberg, R. G. Tompkins, F. M. Ausubel, Pseudomonas aeruginosa killing of Caenorhabditis elegans used to identify P. aeruginosa virulence

factors. *Proceedings of the National Academy of Sciences of the United States of America*

96, 2408-2413 (1999).

11.   L. Walmsley *et al.*, Colour as a signal for entraining the mammalian circadian clock.

PLoS biology **13**, e1002127 (2015).

12.   A. Bendesky, M. Tsunozaki, M. V. Rockman, L. Kruglyak, C. I. Bargmann,

Catecholamine receptor polymorphisms affect decision-making in C. elegans. *Nature*

472, 313-318 (2011).

13.   B. B. Shtonda, L. Avery, Dietary choice behavior in Caenorhabditis elegans. *J Exp Biol*

209, 89-102 (2006).

14.   E. Beale, G. Li, M. W. Tan, K. P. Rumbaugh, Caenorhabditis elegans senses bacterial

autoinducers. *Appl Environ Microbiol* **72**, 5135-5137 (2006).

15.   J. D. Meisel, D. H. Kim, Behavioral avoidance of pathogenic bacteria by Caenorhabditis

elegans. *Trends Immunol* **35**, 465-470 (2014).

16.   J. D. Meisel, O. Panda, P. Mahanti, F. C. Schroeder, D. H. Kim, Chemosensation of

bacterial secondary metabolites modulates neuroendocrine signaling and behavior of C.

elegans. *Cell* **159**, 267-280 (2014).

17.   E. Pradel *et al.*, Detection and avoidance of a natural product from the pathogenic

bacterium Serratia marcescens by Caenorhabditis elegans. *Proceedings of the National*

Academy of Sciences of the United States of America **104**, 2295-2300 (2007).

18.   H. C. Chang, J. Paek, D. H. Kim, Natural polymorphisms in C. elegans HECW-1 E3

ligase affect pathogen avoidance behaviour. *Nature* **480**, 525-529 (2011).

19.   X. Jin, N. Pokala, C. I. Bargmann, Distinct Circuits for the Formation and Retrieval of an

Imprinted Olfactory Memory. *Cell* **164**, 632-643 (2016).

20. K. C. Reddy, E. C. Andersen, L. Kruglyak, D. H. Kim, A polymorphism in npr-1 is a

behavioral determinant of pathogen susceptibility in C. elegans. *Science* **323**, 382-384

(2009).

21. Y. Zhang, H. Lu, C. I. Bargmann, Pathogenic bacteria induce aversive olfactory learning

in Caenorhabditis elegans. *Nature* **438**, 179-184 (2005).

22. G. W. Lau, D. J. Hassett, H. Ran, F. Kong, The role of pyocyanin in Pseudomonas

aeruginosa infection. *Trends Mol Med* **10**, 599-606 (2004).

23. N. Bhatla, H. R. Horvitz, Light and hydrogen peroxide inhibit C. elegans Feeding

through gustatory receptor orthologs and pharyngeal neurons. *Neuron* **85**, 804-818

(2015).

24. T. Blanco-Ayala, A. C. Anderica-Romero, J. Pedraza-Chaverri, New insights into

antioxidant strategies against paraquat toxicity. *Free Radic Res* **48**, 623-640 (2014).

25. T. Booker, R. W. Ness, D. Charlesworth, Molecular evolution: breakthroughs and

mysteries in Batesian mimicry. *Current biology : CB* **25**, R506-508 (2015).

Acknowledgments:

We thank S. Xu and the Caenorhabditis Genetics Center for *C. elegans* strains, F. Ausubel for *P.*

aeruginosa strains, and members of the Nitabach and Kazmierczak labs and the Yale worm

community for technical support, advice, and comments. D.D.G was supported by a National

Institute of Neurological Disease and Stroke (NINDS) National Institutes of Health (NIH)

Predoctoral Fellowship (F31NS080628). Work in the laboratory of M.N.N. was supported in part

by NINDS, NIH (R01NS055035, R01NS056443, R01NS091070) and National Institute of

General Medical Sciences, NIH (R01GM098931).

Figure 1. Light-potentiated avoidance of *P. aeruginosa* requires *lite-1* and pyocyanin blue

pigment toxin.

A. Schematic of bacterial lawn avoidance paradigm. Worms (black squiggles) are placed on a

lawn of *P. aeruginosa* strain PA14. Photographs are of *P. aeruginosa* PA14 liquid culture and 8

kilolux 6500 K white light with corresponding spectrum as measured by a CCD spectrometer.

The fraction of worms outside the lawn is counted once per hour in the absence (dark) or

presence (light) of 8 kilolux 6500 K white light simulating daylight illuminating the entire plate.

B. Timecourse of wild-type worm avoidance of lawns of PA14 or *E. coli* OP50 in the presence

of light or in the dark. Over the course of the nine hour assay, worms increase avoidance of

PA14 in the light (orange) and dark (black), but this avoidance is greatly potentiated by white

light. Worms remain on OP50 lawns (red) throughout the nine hour assay in the presence of

light. **C.** Avoidance of PA14 by *lite-1* null-mutant worms is unaffected by white light. **D.**

Schematic of lawn avoidance paradigm involving *P. aeruginosa* strain PA14ΔphzM, which is

incapable of synthesizing pyocyanin. Photographs are of *P. aeruginosa* PA14ΔphzM liquid

culture (note the change in color when pyocyanin is absent) and 8 kilolux 6500 K white light

used in these experiments. **E.** White light has a less pronounced effect on avoidance by wild-type

worms of mutant *P. aeruginosa* strain incapable of synthesizing pyocyanin. **F.** Avoidance of

PA14ΔphzM by *lite-1* null-mutant worms is unaffected by white light. (Data represent an

average of at least three assays with thirty worms per assay. All statistical comparisons are by

two-way ANOVA with time as a repeated measure and post-hoc Bonferroni tests. For

corresponding *p* and F values, see Supplemental Table 1. Error bars denote s.e.m. * indicates $p <$

0.05, ** indicates $p \leq 0.01$, and *** indicates $p \leq 0.001$.)

Figure 2. Light potentiates avoidance of non-toxic *E. coli* lawns doped with pyocyanin.

A. Schematic of experimental design for testing avoidance of *E. coli* OP50 lawns doped with pyocyanin. Photographs are of *E. coli* OP50 liquid culture and doping solutions used in these experiments. **B.** Timecourse of avoidance by wild-type worms of *E. coli* OP50 lawns doped with 2.5 mM pyocyanin in the presence or absence of light reveals white light potentiation of stable, long-term, avoidance of OP50 lawns. For corresponding p and F values, see Supplemental Table 1. **C.** White light potentiates rapid one-hour avoidance of OP50 lawns doped with 2.5 mM pyocyanin by wild-type but not *lite-1* null-mutant worms. Avoidance of OP50 lawns doped with 0.25 mM or 5 mM pyocyanin is unaffected by light. **D.** Re-expression of *lite-1* in *lite-1* null-mutant worms using the *trx-1* or *sra-6* promoters for expression in neurons ASJ and ASH/ASI/PVQ, respectively, rescues light-potentiated avoidance of OP50 lawns doped with 2.5 mM pyocyanin. Re-expression of *lite-1* using the *srg-8* promoter for expression in sensory neuron ASK has no effect. **E.** *lite-1* function in rapid light- and pyocyanin-dependent lawn avoidance localizes to a small number of primary sensory neurons previously shown to be involved in long-term PA14 avoidance, and only partially overlaps with those involved in escape responses to bright short-wavelength light. (Data in this figure and all remaining figures represent an average of at least three assays with thirty worms per assay. All statistical analyses were performed by one-way ANOVA with post-hoc Tukey-Kramer for pairwise comparisons or Bonferroni tests for comparisons with control. Error bars denote s.e.m. * indicates $p < 0.05$, ** indicates $p \leq 0.01$, and *** indicates $p \leq 0.001$.)

Figure 3. Light potentiates avoidance of *E. coli* lawns doped with paraquat colorless toxin

mixed with non-toxic blue dye.

A. Experimental design for testing avoidance of OP50 lawns doped with non-toxic blue food

dye. Photographs are of *E. coli* OP50 liquid culture and doping solutions used in these

experiments. **B.** Avoidance of OP50 lawns doped solely with blue dye is unaffected by white

light. **C.** Experimental design for testing avoidance of OP50 lawns doped with toxic paraquat and

non-toxic blue dye. **D.** White light potentiates avoidance of OP50 lawns doped with 30 mM

paraquat and blue dye by wild-type but not *lite-1* null-mutant worms. Avoidance of OP50 lawns

doped with 20 mM or 40 mM paraquat and blue dye is unaffected by light. **E.** Experimental

design for testing avoidance of OP50 lawns doped with toxic paraquat with no dye or with non-

toxic red food dye. **F.** Avoidance of OP50 doped with 30 mM paraquat alone or paired with red

dye is unaffected by white light.

Figure 4. Color of light determines avoidance of toxic bacterial lawns.

A. The spectral composition of light was modified by various optical filters. Photographs are of

E. coli OP50 liquid culture and doping solutions used in these experiments, as well as lights

modified by the indicated filter shining on a white background. Light spectra were measured

using a CCD spectrometer. Blue light or blue-green light obtained with 500 nm or 550 nm cut-

off shortpass filters each fail to potentiate avoidance of OP50 lawns doped with 2.5 mM

pyocyanin. White light without blue light obtained with a 495 nm longpass filter also fails to

potentiate avoidance. Only light containing both blue and yellow-orange light, obtained with a

600 nm cut-off shortpass filter, potentiates avoidance of OP50 doped with 2.5 mM pyocyanin. **B.**

Experimental design for testing whether blue pigment causes light-potentiated avoidance by

absorbing long-wavelength light and altering the spectral composition of light sensed by the

worm, by eliminating blue pigment from the lawn and directly controlling the spectral

composition of light with filters. **C.** Light modified by a blue vinyl filter that decreases intensity

of yellow-orange light without eliminating it potentiates avoidance of colorless OP50 lawns

doped with 30 mM paraquat, but not lawns lacking paraquat. **D.** Schematic depicting the

multisensory decision whereby integration of color and chemical information guides the foraging

decision to stay on or leave toxic or non-toxic bacterial lawns. Blue pigment absorbs long-

wavelength light and thereby alters the spectral composition of light detected by the worm. We

propose that worms increase their vigilance to ROS when they detect a decrease in yellow-

orange light content induced by blue pigments in their environment.

Supplementary Materials:

Supplementary Figure 1. Spectral composition of 6500 K 8 kilolux LED array white light

source.

A. Photograph and spectral profile of white LED backlight as measured using a CCD

spectrometer.

Supplementary Figure 2. Spectral composition of white light modified by filters.

A. Photographs of 8 klx white light modified by various filters. **B.** Spectral profiles of white light

modified by LP (longpass) or SP (shortpass) filters with cut-offs at the indicated wavelengths

were measured using a CCD spectrometer.

Supplementary Figure 3. Bright white light fails to potentiate avoidance of toxic colorless

bacterial lawns.

A. Experimental design for testing whether increasing flux of blue light by increasing intensity

of incident white light potentiates avoidance of colorless toxic food lawns. **B.** Increasing white

light intensity ~3-fold to 25 klx fails to potentiate avoidance of OP50 lawns doped with 30 mM

paraquat containing no blue dye. While further increases in white light intensity to 50 or 100 klx

potentiate avoidance of OP50 lawns, this is independent of the presence of paraquat. **C.**

Representative fluorescent images of the ROS-dependent indicator H2DCFDA in worms

exposed to various intensities of white light for one hour, with or without 30 mM paraquat. **D.**

Quantification of fluorescence from experiments shown in panel **c**. For each condition, at least 7

worms from 3 independent experiments were analyzed. Statistical analysis was performed by

one-way ANOVA with post-hoc Tukey-Kramer for pairwise comparisons.

Supplemental Table 1. F-values and *p* values from statistical analyses performed by two-

way ANOVA with time as a repeated measure.

Comparison	Relevant panel	F-value (time	*p* value (time)	F-value (condition)	*p* value (condition)
Wild-type worms on PA14 lawn, light vs. dark	Figure 1b	44.18	< 0.0001	27.57	0.0033
lite-1 null-mutant worms on PA14 lawn, light vs. dark	Figure 1c	52.93	< 0.0001	0.7630	0.4223
Wild-type worms on PA14ΔphzM lawn, light vs. dark	Figure 1e	27.04	< 0.0001	10.06	0.0248
lite-1 null-mutant worms on PA14ΔphzM lawn, light vs. dark	Figure 1f	26.31	< 0.0001	0.1715	0.6960
Wild-type worms on OP50 + 2.5 mM, light vs. dark	Figure 2b	0.4251	0.8891	462.3	0.0022

Materials and Methods:

Strains

C. elegans strains were maintained on Nematode Growth Medium (NGM) agar plates with *E.*

coli OP50 as a food source. All strains were derived from the Bristol N2 wild-type strain. The

lite-1 null-mutant strain was backcrossed to wild-type N2 at least 5 times. Strains used for each

figure are detailed below:

Figure 1: N2, KG1180 *lite-1(ce314)*

Figure 2: N2, KG1180 *lite-1(ce314),* TQ1101 *lite-1 (xu7)*, TQ1179 *lite-1(xu7)*; xuEx301 [*sra-*

6p::lite-1::sl2::yfp; sra-6p::mCherry2; unc-122p::DsRed], TQ1188 *lite-1(xu7)*; xuEx7 [*trxp-*

1::lite-1; unc-122p::gfp; trx-1p::mCherry2], TQ1203 *lite-1(xu7)*; xuEx [*srg-8p::lite-1::sl2::yfp;*

unc-122p::gfp; srg-8p::mCherry2]

Figure 3: N2, KG1180 *lite-1(ce314)*

Figure 4: N2

Supplementary Figure 3: N2

Lawn avoidance assay

P. aeruginosa lawn avoidance assays were performed on slow killing assay (SKA) plates as

described (*8-10, 16, 18, 20*). *P. aeruginosa* cultures were grown in LB liquid media at 37°C for

16-22 hours. 7 µL spots were added to the center of a 3.5 cm plate filled with 5 mL of SKA agar.

Plates were first incubated at 37°C for 22-24 hours, and then at room temperature for 16-20

hours. Worms were maintained in regular room lighting or incubator lighting conditions.

Growing worms in the presence or absence of light had no effect on assay results (data not

shown). Thirty 18-24 hours post-L4 staged adult worms were transferred to the lawn for each

assay. The fraction of worms out of those thirty that were off the lawn after the specified time

interval was recorded as fraction avoiding.

Solutions of specified concentrations of pyocyanin (Cayman Chemical) and paraquat (ULTRA

Scientific) were added to *E. coli* OP50 lawns on plates prepared exactly as described above.

Erioglaucine (Alfa Aesar) or sulfarhodamine (MP Biomedicals) were used for blue and red color,

respectively. Before adding solution, a 1.5 mL Eppendorf tube cut open at both ends was placed

on the plate such that the rim of the tube circumscribed the boundary of the lawn. Next, 300 µL

of the indicated solution was added to the tube to direct gradual application of solution to the

lawn. Plates were prepared accordingly for three hours to allow for complete drying of the

solution before transferring worms to the lawn and beginning the assay.

Statistical analysis

Sample sizes were chosen to ensure that at least two sets of experiments for each condition were

carried out. No samples or animals were excluded from statistical analysis. Animals of a

particular genotype were randomly assigned to assay plates. The order of assay plates loaded

with worms was also randomized for each experiment, ensuring that the order of conditions

tested was randomized. The investigator was blinded to genotype and condition (e.g., PA14

strain, concentration of paraquat) throughout, but not to different lighting conditions (e.g., light

vs. dark, filter vs. no filter).

At least three independent assays were performed for each condition. For time course

experiments in Figures 1 and 2, two-way ANOVA with time as a repeated measure and post-hoc

Bonferroni tests was used to analyze statistical differences.

For all other experiments, one-way ANOVA was used to analyze statistical differences, with

post-hoc Tukey-Kramer tests performed for pairwise comparisons between all conditions or

post-hoc Bonferroni tests performed for comparisons to control. * indicates $p < 0.05$, **

indicates $p \leq 0.01$, and *** indicates $p \leq 0.001$.

Illumination

For experiments shown in Figures 1, 2, and 3, assay plates were illuminated in a completely dark

room by a white LED backlight (Edmund Optics, Stock Number #83-873). After taking the lid

off the plates, the plates were inverted onto the backlight. For assays performed in the dark,

plates were positioned similarly either with the light off or on aluminum foil to block the light.

For experiments shown in Figure 4 and Extended Data Figure 3, plates were illuminated by a

white LED (Thorlabs MCWHL5 with SM2P50-A attached) clamped to a ring stand. LEDs were

positioned 18 cm above the plate, illuminating a 5 cm diameter spot covering the entire 3.5 cm

diameter assay plate. Assay plates were always centered within this spot to circumvent any

effects of imperfect collimation of the incident light. Multiple plates were illuminated

simultaneously, each by a single LED. Multiple LEDs were powered simultaneously by a 4-

Channel LED Driver (Thorlabs DC4100). In this lighting configuration, plate lids were removed,

but plates remained upright.

A calibrated optometer Gamma Scientific (UDT Instruments Model S471 with 268P illuminance

sensor) was used to measure illuminance. Irradiance at specified wavelengths was measured

using the same instrument, but connected to a 268R sensor head that was continuously calibrated

from 350 nm to 1100 nm. Irradiance at 470 nm was measured after filtering the light source

through 500 nm shortpass filters.

The following filters were used in the experiments shown in Figure 4: 495 nm longpass filter

(Thorlabs FGL495S), 500 nm shortpass filter (Edmund Optics Stock Number 64663), 550 nm

shortpass filter (Edmund Optics Stock Number 64664), 600 nm shortpass filter (Edmund Optics

Stock Number 64665), and a blue vinyl sheet (Hull's Art store). Spectral profiles were measured

using a compact CCD spectrometer (Thorlabs CCS100 with connected cosine corrector CCSB1).

Relative intensities measured by the spectrometer do not permit comparisons of absolute

intensity between the various filters. Illuminance of white light modified by each filter was

calibrated to ensure that the intensity of short-wavelength (470 nm) light passing through each

filter was identical as measured using a calibrated irradiance detector described above. This only

applied to white light modified by the blue vinyl filter, which required calibration of < 1 klx.

ROS detection

After exposure to the specified light/dark and paraquat conditions, worms were washed three

times with M9 buffer. Seven to nine worms from each treatment group were incubated with 25

μM 2'-7'-dichlorodihydrofluorescein (Biotium) in the dark for 30 min at 20 °C. After

H2DCFDA incubation, worms were washed with M9 solution and mounted in PBS buffer

containing 1 mM sodium azide. Fluorescence was measured immediately after mounting to

slides using a Leica M165FC fluorescent stereomicroscope at excitation/emission wavelengths of

488 nm and 520 nm. The mean fluorescence intensity of the whole body of the worm was

quantified using ImageJ software. Values were then compared using one-way ANOVA with

post-hoc Tukey-Kramer test. For each condition, at least 7 worms from 3 independent

experiments were analyzed.

Additional Author notes:

Author Contributions

Designed study and experiments: DDG, MNN

Performed experiments: DDG, XJ

Data analysis and statistics: DDG, XJ

Wrote the paper: DDG, MNN

Author Information

The authors declare no conflicts of interest.

Correspondence and requests for material should be addressed to michael.nitabach@yale.edu

Figure 1

Figure 2

Figure 3

Figure 4

Supplementary Figure 1

A

8 kilolux LED array white light source
(Color temp.: 6500 K)

Supplementary Figure 2

Supplementary Figure 3

A

B

C

D