1 *Clostridium difficile* colonizes alternative nutrient

2 niches during infection across distinct murine gut

3 environments

4 **Authors:** Matthew L. Jenior, Jhansi L. Leslie, Vincent B. Young, and Patrick D. Schloss[*]

5 **Abstract**

6 *Clostridium difficile* infection (CDI) has grown to be the most prevalent cause of hospital

7 acquired infection in the United States. Susceptibility to CDI is induced by recent

8 antibiotic exposure, which is known to alter the structure of the gut microbiome and to

9 affect the availability of growth nutrients in the gut. We hypothesized that *C. difficile* is a

10 generalist that adapts its physiology to the nutrients available within the gut. We orally

11 challenged C57BL/6 mice that previously received one of three antibiotics with *C.*

12 *difficile* and demonstrated that it was able to colonize the cecum within 18 hours of

13 infection. However, levels of both spore and toxin production, which are known to be

14 affected by nutrient availability, varied between each antibiotic treatment group. To

15 more closely investigate the specific responses of *C. difficile* as it colonized the cecum,

16 we performed *in vivo* transcriptional analysis of *C. difficile* from cecal content of infected

17 mice. This approach revealed variation in expression of genes that drive life-cycle

18 switches as well as metabolic pathways associated with catabolizing a variety of carbon

19 sources such as carbohydrates, amino acids, and amino sugars. To assess which

20 substrates *C. difficile* was most likely exploiting in each antibiotic-perturbed microbiome,

we developed a novel metabolite scoring algorithm within the genome-scale bipartite

metabolic network of *C. difficile* that incorporated both network topology and transcript

abundance to infer the likelihood that a given metabolite was acquired from the

environment. Applying this approach, we found that *C. difficile* indeed occupies

alternative nutrient niches across each antibiotic-perturbed microbiome and that the

highlighted metabolites support significant growth, *in vitro. In silico* findings were further

validated using *in vivo* metabolomic analysis to compare resitant and susceptible cecal

conditions. These results support the hypothesis that consumption of N-acetyl-D-

glucosamine and Stickland fermentation substrates are central components of *C.*

difficile's metabolic strategy and pathogenesis. This work has implications for

elucidating specifics of the nutrient niche of *C. difficile* during infection and may lead to

the discovery of targeted measures to prevent *C. difficile* colonization including potential

pre- or probiotic therapies.

Significance

Clostridium difficile infection has grown to be the largest single cause of hospital-

acquired infection in the United States. A compromised gut microbiota, typically through

previous antibiotic treatment, is a prerequisite feature of *C. difficile* colonization

susceptibility. This loss of resistance has also been described across classes of

antibiotics, in which many result in distinct gut communities and each presumably

presents individual metabolic challenges to *C. difficile* as it colonizes. In this study we

demonstrate that not only does *C. difficile* alter its pathogenesis between differentially

sensitized hosts, but also exploits separate available nutrient niches across these

environments. We accomplish this through an *in vivo* murine model of infection,

characterized using transcriptomic-enabled genome-scale metabolic modeling in

combination with untargeted metabolomic analysis. Our results support that *C. difficile*

possesses a highly plastic nutrient niche space, allowing it to successfully infect

numerous different hosts and ultimately cause disease. This work also provides

evidence that *C. difficile* virulence may be driven by the concentrations of specific

carbohydrate sources that are accessible for growth during each unique infection.

Introduction

Infection by the Gram-positive, spore-forming bacterium *Clostridium difficile* has

increased in both prevalence and severity across numerous countries during the last

decade (1). In the United States, *C. difficile* was estimated to have caused >500,000

infections and resulted in ~$4.8 billion worth of acute care costs in 2014 (2). *C. difficile*

infection (CDI) causes an array of toxin-mediated symptoms ranging from abdominal

pain and diarrhea to the more life-threatening conditions pseudomembraneous colitis

and toxin megacolon. Prior treatment with antibiotics is the most common risk factor

associated with susceptibility to CDI (3). It has been shown that antibiotic therapy alters

the structure and function of the gut microbiota making it susceptible to colonization by

C. difficile (4). This is referred to as colonization resistance, in which the gut microbiota

inhibits the persistence or growth of a number of pathogenic bacteria. Colonization

resistance can be achieved by multiple mechanisms including competition for physical

space or growth nutrients (5).

Mouse models have been an effective tool for studying the mechanisms of colonization

resistance. Use of distinct antibiotic classes to vary the structure of the microbiota which

have been shown to result in susceptibility *C. difficile* colonization (6–8). In this case,

the antibiotics chosen significantly impact the structure and diversity of the cecal

microbiome (Fig. S1a & S1b). It has been further demonstrated that at 18 hours after

being introduced to a cefoperazone treated mouse, *C. difficile* reached its maximum

vegetative cell density in the cecum (9). This provided a single timepoint to measure the

largest population of metabolically active *C. difficile*. Building upon these results, others

have shown that many of these antibiotic classes also alter the gut metabolome,

increasing the concentrations of known *C. difficile* growth substrates (7, 10–12). Taken

together these results are a strong indication that the healthy gut microbiota inhibits the

growth of *C. difficile* through limitation of substrates it needs to grow. The ability of an

intact gut community to exclude *C. difficile* colonization is suggestive of the nutrient-

niche hypothesis in which an organism must be able to utilize a subset of available

resources better than all competitors to colonize the intestine (13, 14).

Based on its genome sequence and *in vitro* growth characteristics, *C. difficile* appears

able to fill multiple nutrient niches. *C. difficile* has a relatively large and mosaic genome,

it is amenable to a variety of growth substrates, and is able to colonize a diverse array

of hosts suggesting that that it is a bacterial generalist (15–17). The ability to metabolize

a variety of substrates is important since these substrates affect the regulation of genes

involved in *C. difficile*'s pathogenesis. For example, *in vitro* transcriptomic analysis

suggests that high concentrations of easily metabolized carbon sources, such as

glucose or amino acids, inhibit toxin gene expression and sporulation (18, 19). These

genes are regulated by DNA-binding sigma factors, such as the pleiotropic regulator

ccpA, which are under the control of environmental nutrient concentrations, especially

carbohydrates (20, 21). Downstream effects of this regulation likely have enormous

impact on the lifestyle and metabolic strategy of *C. difficile* when colonizing across

sensitive hosts.

Previous transcriptomic studies of *C. difficile* have mainly focused on transcription of

virulence factors, *in vitro* (22, 23), with some work characterizing transcription during

colonization of germfree mice (24, 25). More relevant to nutrient acquisition, *C. difficile*

up-regulated several phosphotransferase systems (PTS) and ABC transporters in

germfree mice, alluding to metabolic adaptation to nutrient availability *in vivo* (25).

Although these analyses are informative, they are either primarily directed toward the

expression of virulence factors or lack the context of the gut microbiota which *C. difficile*

must compete against for substrates. Metabolomic analyses have also been used to

more directly assay changes in bacterial metabolism as they relate to CDI (7, 12);

however, these methods cannot focus on *C. difficile*-specific metabolites and more

closely resemble echoes of metabolism, not currently active processes. In contrast to

these approaches, *in vivo C. difficile* transcriptomic analysis from specific pathogen free

(SPF) animals may provide unique insight into its active metabolic pathways in a more

realistic model of infection. Integrating transcriptomic data with genome-scale metabolic

modeling has previously aided in identifying the most active aspects of an organism's

metabolism and which substrates are preferred by the organism (26–28). Applying

these methods to study *C. difficile* colonization would allow us to directly test the

nutrient-niche hypothesis.

Founded on the ability of *C. difficile* to grow on a diverse array of carbon sources and its

ability to colonize a variety of communities, we hypothesized that it focuses its

metabolism to fit the context of the community it is attempting colonize. To test this

hypothesis, we employed a mouse model of infection to compare the response of *C.*

difficile to the gut environment caused by different classes of antibiotics. The antibiotics

used in this study included streptomycin (Fig. 1a), cefoperazone (Fig. 1b), and

clindamycin (Fig. 1c). These antibiotics differentially affect the structure of the gut

microbiota (8). Each has also been shown to alter the gut metabolome relative to

untreated animals (7, 10, 12). As such, we predicted that *C. difficile* would encounter a

unique subset of nutrients and competitors in each environment, which would

necessitate distinct adaptive responses. To determine whether *C. difficile* is a generalist

and differentially responds to each condition, we assayed for differences in the amount

of sporulation and toxin activity phenotypes and used metabolic models built using *C.*

difficile expression data. In each of the three antibiotic conditions we challenged with *C.*

difficile, as well as in monoassociated germfree mice, we observed that *C. difficile*

adapted its nutrient utilization profile to colonize to high levels and express its virulence

factors.

Results

Levels of *C. difficile* sporulation and toxin activity vary between antibiotic-treated

specific pathogen free and germfree mice.

Due to the connection between metabolism, sporulation, and toxin production in *C.*

difficile, we measured sporulation and toxin production at 18 hours post infection in

each group. There was not a significant difference in the number of vegetative cells

between any susceptible condition tested (Fig. 2a). All antibiotic treated (Table 1)

specific pathogen free (SPF) and germfree (GF) animals were colonized to ~1×10^8

colony forming units (c.f.u.) per gram of content, while untreated SPF mice maintained

colonization resistance to *C. difficile*. Despite having the same number of vegetative *C.*

difficile cells, large differences were detected in the density of *C. difficile* spores.

Significantly more spores ($P = 0.005, 0.008, 0.003$) were detected in ex-GF mice than in

the antibiotic treated mice (Fig. 2b). The spore densities in both streptomycin and

clindamycin-treated mice were also generally higher than that in cefoperazone-treated

mice. There was significantly more toxin activity in ex-GF animals than any other

colonized group (all $P <= 0.001$), but toxin titer also varied between antibiotic treatment

groups (Fig. 2c). Although similar toxin activity was found in both the cefoperazone and

clindamycin-treated groups, toxin titer was below the limit of detection in most

streptomycin-treated animals. These results indicate that *C. difficile* was able to colonize

different communities to a consistently high level, but that the density of spores and

toxin titer varied by treatment.

***C. difficile* adapts the expression of genes for virulence and key sigma factors**

that are under the control of environmental nutrient concentrations. To more

closely investigate the responses of *C. difficile* to colonizing distinct susceptible gut

environments, we performed whole transcriptome analysis of *C. difficile* during infection

of the antibiotic treatment models. We then narrowed our analysis to focus on genes

that control or code for functions that have been linked to nutrient concentrations in the

intestines during CDI. After observing differences in spore load, we first examined

transcription of the most highly expressed genes in the *C. difficile* sporulation pathway

(29–32) (Fig. 3a). Across the four conditions where *C. difficile* colonized, we observed

transcriptional profiles consistent with observed spore levels (Fig. 2b). The mice treated

with cefoperazone had the lowest spore density and had the highest level of expression

for the anti-sigma factors *spoVG* and *spoVS*. The produts of these genes are involved in

suppressing expression of genes found later in the sporulation pathway (33).

Streptomycin-treated mice had the next highest density of spores and the highest

expression of genes associated with sporulation activation (*spoIIAB*/*spoIIE*), but they

also had relatively high levels of expression of *sspA* and *sspB*, which are genes that

code for effectors that protect DNA from damage during dormancy. Next, in mice

treated with clindamycin, *C. difficile* expressed genes associated with late stages of

sporulation, including those for spore coat components (*cdeC*, *cotD*, and *cotJB2*), spore

formation (*spoIVA*, *spoVB*, and *spoVFB*), and *sspA* and *sspB*. Finally, GF mice

harbored the highest density of spores and those *C. difficile* primarily expressed the

dormancy genes linked with the latest stages sporulation. Together these data

demonstrate that *C. difficile* differentially expressed genes associated with sporulation

that corresponded to the presence of spores in the cecum.

Expression of genes for quorum sensing and pathogenicity have been linked to

changes in the nutrients that can be found in the environment of *C. difficile*. Both the *agr*

locus and *luxS* gene are thought to be associated with inducing the expression of *C.*

difficile virulence in several strains (34, 35). Considering the link between quorum

sensing genes and toxin production, we expected the expression of genes for quorum

sensing and toxin production and toxin titer to be concordant. Based on this model, we

expected GF mice to have the highest levels of expression of genes for toxin production

(Fig. 3b) and quorum sensing (Fig. 3c); however, these transcripts were not found in the

GF mice. We also observed the highest level of expression for quorum sensing genes in

cefoperazone-treated mice, but *tcdA* expression in these animals was not the highest

among the different treatment groups. Interestingly, the levels of expression for genes

associated with toxin production did not match the toxin titers observed in the animals.

These results suggest that the relationship between toxin titer and the expression of

genes for toxin production is even more complex than current models indicate.

We next focused on the regulators of metabolic pathways. Sigma factors are master

regulators and a subset have been shown to integrate signals from intra- and

extracellular nutrient concentrations (20, 21, 31, 36). The transcription of the global

repressor *codY* is responsive to intracellular concentrations of *C. difficile* energy

sources (37). Highest transcription for this gene was found in cefoperazone-treated and

GF mice (Fig. 3d). The regulation networks of CodY and CcpA are highly

interconnected, with the expression of *ccpA* specifically linked to local concentration of

rapidly metabolizable carbon sources (38). Cefoperazone-treated mice also exhibited

increased transcription of *ccpA*, but the GF condition did not follow the same pattern.

CcpA acts directly on *spo0A* (Fig. 3d), which positively regulates initiation of the

sporulation pathway in *C. difficile.* Transcripts for *spo0A* were highly abundant in all

conditions tested except for clindamycin-treated mice, where it was still moderately

detectable. The sig-family of sigma factors is under the control of spo0A and regulate

different stages of sporulation. The genes from this family with the highest total

transcription (*sigA1*, *sigF*, *sigG*, *sigH*, and *sigK*) each demonstrated a unique pattern of

expression between conditions. These results indicate that complete expression of

sporulation likely integrates multiple levels of signaling and is more complex than a

single metabolic switch. Both CcpA and Spo0A also control pathogenicity by regulating

toxin production (Fig. 3d). We found expression of the toxin negative regulator *tcdC* in

all of the antibiotic-treated groups, but no detectable transcripts for the positive toxin A/B

regulator *tcdR* were seen in any treatment. In addition to its effects on sporulation and

virulence, CcpA also regulates the expression of other sigma factors that generally

mediate distinct forms of *C. difficile* metabolism as needed. These targets include *rex*

(general fermentation regulator) and *prdR* (Stickland fermentation regulator) (Fig. 3d).

Although the expression of both has been shown to be linked to environmental proline

concentrations, *rex* integrates additional signals from the intracellular NADH/NAD+ ratio

to also control carbohydrate fermentation. Low-level transcription of *prdR* was found

across all conditions, however *C. difficile* expression the *rex* gene highly in both

cefoperazone-treated and GF mice. Combined, the variable expression of these sigma

factors support the hypothesis that *C. difficile* adapts expression metabolism to fit its

needs between colonized environments.

Gene sets from multiple *C. difficile* metabolic pathways are differentially

expressed between colonized environments. In the context of similar colonization

between antibiotic-treated animals, differential expression of global metabolic control

mechanisms that are under the control of specific nutrient concentrations suggests that

C. difficile adapts to each environment when in competition with the resident microbiota.

To test this further, we quantified the total expression for all KEGG annotated genes in

the *C. difficile* genome (Fig. S2A). We then focused on general differences in

carbohydrate (Fig. S2b) and amino acid (Fig. S2C) metabolism in order to asses for

apparent differences in the utilization of carbon sources by *C. difficile* across

environments. However aside from overall lower expression of most gene families in GF

mice, no other clear trends were evident at this broad level of analysis so we moved

toward a more fine-scale resolution of annotation and focused on specific gene sets

known to contribute to certain forms of *C. difficile* metabolism (Table S1). Also, to more

effectively compare between colonized states, we calculated the percentage of total

expression between antibiotic-treated conditions for each gene (Fig. 4). We then

identified the condition in which each gene was most highly transcribed and adjusted

the size of the corresponding point relative to the largest transcipt abundance. This

demonstrated that genes involved in amino acid catabolism had the greatest amount of

expression overall, relative to other gene sets. This category includes those enzymes

involved in Stickland fermentation (*arg*, *fdh*, *grd*, and *prd* loci) as well as several general

peptidases (*pep* family). These results indicated that catabolizing environmental amino

acids may be important for the growth of *C. difficile* during infection.

To more clearly indentify associations of gene sets with each condition, we also

analyzed each set separately. First, we found that the expression of genes associated

with amino acid catabolism were expressed at nearly consistent levels across the

conditions. This was in agreement with the high level of overall expression associated

with these genes. Additionally, genes for the metabolism of the host-derived amino

sugars N-acetylglucosamine and N-acetylmannosamine were also expressed at

consistent levels across each treatment group (*glm*, *nan*, *mur*, and *acd* loci). Along

similar lines with related molecules, a number of genes for certain monosaccharides

entering (*gal*, *man*, *pmi*, and *tag* loci) and leading through glycolysis (*fba*, *fbp*, *gap*, and

pfk), as well as catabolism of the polysaccharides trehalose and cellibiose (*treA* and

celG) were expressed relatively evenly between each condition. Combined, these

findings suggest that catabolism of amino acids and specific carbohydrates are likely

core components of the *C. difficile* nutritional strategy during infection.

Aside from those gene sets that were equally expressed across conditions, there were

also large scale differences in expression of certain pathways between groups of mice.

We chose to assess sugar transport systems have been associated with adaptive

expression of phosphotransferase systems (PTS) and ABC transporters with many

known differences in substrate specificities (25). Among the genes classified as PTS

transporters were overrepresented in both clindamycin and streptomycin-treated mice,

while ABC sugar transporters were overrepresented in the cefoparazone-treated mice.

The most stark differences were seen in transcription for genes involved in sugar

alcohol catabolism. Expression of these genes was entirely absent from clindamycin-

treated mice and expression of genes for mannitol utilization (*mtl* operon) were

overrepresented in cefoparazone-treated mice and expression of genes for sorbitol

utilization (*srl* operon) were overrepresented in streptomycin-treated mice. Concordant

patterns also emerged in genes associated with fermentation end steps and

polysaccharide degradation. Short chain fatty acids (SCFAs) and alcohols are the end

products of both carbohydrate and amino acid fermentation in *C. difficile* through

separate pathways with shared terminal steps. Transcripts for genes involved in *C.*

difficile butyrate/butanol metabolism (*ptb*, *buk1*, *cat2*, and *adhE*) were more abundant in

clindamycin-treated mice. Additionally, alpha/beta-galactosidase genes (*aglB* and *bglA*)

were also overrepresented in clindamycin-treated mice. Together these patterns

suggested that polysaccharide fermentation occurred this condition. More subtle

differences were seen in those gene associated with glycolysis. This category includes

genes for not only the steps of glycolysis, but also several genes that mediate entry

points of monosaccharides to glycolysis. Transcripts for several genes in this group

(*eno*, *gapA*, *gpmI*, *tpi*, and *pyk*) were overrepresented in cefoparazone-treated mice,

however *fruK* was overrepresented in streptomycin-treated mice which catalyzes the

committed step of glycolysis. Overall, these results support the hypothesis that *C.*

difficile is able to adapt its metabolism to fit the nutrient availability across different

susceptible environments.

Structure of genome-scale bipartite metabolic model underscores known

bacterial metabolism. To further investigate which metabolites were differentially

utilized between conditions, we represented the metabolic network of *C. difficile* as a

directed bipartite graph using the genome annotation. Enzymes and metabolites were

represented by nodes and their interaction by the edge between the nodes (Fig. 5a). To

validate our metabolic network, we calculated betweenness centrality (BC) and overall

closeness centralization index (OCCI) for all enzyme and metabolite nodes in the

bipartite metabolic network of *C. difficile* generated for this study (Table S2). In

biological terms, BC reflects the amount of influence a given hub has on the overall flow

of metabolism through the network (39) and OCCI indicates those enzymes and

substrates that are the most central components of the organism's metabolism (40). For

both enzymes and substrates, the 18 of top 20 nodes with the highest BC values were

involved in glycolysis, fermentation, and amino acid synthesis. In agreement, almost all

nodes with the largest OCCI values were involved in glycolysis and amino acid

synthesis as well. Enzymes that scored highly in both metrics included pyruvate kinase,

aspartate aminotransferase, and formate C-acetyltransferase while substrates

consistently scoring most highly were pyruvate, acetyl-CoA, D-glyceraldehyde 3-

phosphate. This indicated to us that the topology of the network reflects established

bacterial physiology.

Metabolite importance algorithm reveals adaptive nutritional strategies of *C.*

***difficile* during infection across distinct environments.** Moving beyond a strictly

topological analysis of the *C. difficile* metabolic network, we sought to utilize

transcriptomic data to infer which metabolites *C. difficile* is most likely to obtain from its

environment in each condition. To accomplish this we mapped normalized transcript

abundances to the enzyme nodes in the network. Due to the coupling of transcription

and translation in bacteria, we were able to use this information as a proxy for enzyme

levels. The importance of each metabolite was measured as the log-transformed

difference between the average transcript levels of enzymes that use the metabolite as

a substrate and those that generate it as a product (Fig. 5b). A metabolite with a high

importance score is most likely obtained from the environment because the expression

of genes for enzymes that produce the metabolite are low. Then, using a Monte Carlo-

style simulation, we generated a random transcript abundance distribution for each

enzyme node to then calculate new metabolite importance scores for each iteration. We

then created a confidence interval of scores for each metabolite that would likely result

from random noise (41). This provided a standard of comparison for actual importance

scores from single timepoint measurments, and ultimately allow for computing the

significance level that a given score has a high probability of being excluded from its

associated null hypothesis score distribution.

Applying these methods to the *C. difficile* transcriptomic data collected from the *in vivo*

CDI models, we sought to assess differential patterns of metabolite importance. We first

ranked the importance scores to identify the most important metabolites for each

treatment group (Table S3). To identify the core metabolites that are essential to *C.*

difficile in any condition, we compared the highest 50 scoring, significant metabolites

from each treatment group ($P < 0.05$) (Fig. 6a). The host derived amino sugar N-acetyl-

D-glucosamine was found to be consistently important. Components of the Stickland

fermentation pathway were also found to be important to *C. difficile* in all conditions

tested including proline, 3–hydroxybutanoyl–CoA, and formate (42–44). This indicated

that these metabolites may be an integral component of the nutrient niche for *C. difficile*

in any infection condition. Additionally, acetate was found to be important in all

conditions, but was just below the significance cutoff in GF mice (Table S3). It has been

shown that *C. difficile* metabolizes acetate for use in glycolysis (45). We directly tested

the relative concentration of acetate in cefoperazone-treated *C. difficile*-infected mice

versus mock-infected mice. We found that *C. difficile* colonization led to a significant

decrease in the levels of acetate (Fig. S4) suggesting that *C. difficile* was utilizing

acetate in the cecum. These findings provided validation for our metabolite importance

algorithm as well as supporting known elements of *C. difficile* metabolism.

Returning to our hypothesis that *C. difficile* adapts its metabolism to fit the surrounding

community, we identified those metabolites that were uniquely important to each

condition in which *C. difficile* colonized. We cross-referenced the top 25 positively

scoring, significant substrates ($P < 0.05$) between treatment groups to uncover the most

important patterns of nutrient utilization by *C. difficile* in each (Fig. 6b). Each group of

metabolites contained at least one known carbohydrate growth substrate of *C. difficile*

(7, 46). This included sorbitol, galacititol, mannitol, N-acetylneuraminic acid, and salicin.

Furthermore, in GF mice where no other competitors are present, our model indicated

that *C. difficile* was more likely to acquire numerous amino acids from the environment

instead of expending energy to produce them itself. These data support the hypothesis

that *C. difficile* may exploit alternative nutrient sources between the susceptible

environments it colonizes.

Carbohydrates predicted to be important using network-based approach

differentially supported *C. difficile* growth *in vitro*. To validate the biological

relevance of substrates identified as uniquely important to *C. difficile* metabolism

through our network-based analysis, we tested whether *C. difficile* was able to utilize

each substrate for *in vitro* growth (Fig. 6c). This was performed using a modified defined

C. difficile minimal media (7), supplemented individually with the selected carbohydrates

implicated by high importance scores. Because *C. difficile* is auxotrophic for several

amino acids, it was necessary to include amino acids in the minimal media despite the

capability to achieve modest growth through Stickland fermentation of these substrates.

This focused our analysis on carbohydtrates and made the most effective negative

control growth in media lacking carbohydrates but containing amino acids.

Relying on network-based analysis a single carbohydrate source, N-acetyl-D-

glucosamine, was found to be important to *C. difficile* in each condition tested (Fig. 6a).

When tested for improved growth, significantly more growth (Max OD_{600} = 0.774) was

observed compared to negative controls (P < 0.001). This provided evidence that N-

acetyl-D-glucosamine, derived from the host mucus layer, may be a central component

of the *C. difficile* nutritional niche during infection. Upon analysis of carbon sources

identified as distinctly important across treatment groups, we found that at least one

substrate from cefoperazone-treated, clindamycin-treated, and GF groups provided high

levels of *C. difficile* growth relative to negative controls ($P < 0.001$). This included

mannitol (cefoperazone; Max $OD_{600} = 0.464$), salicin (clindamycin; Max $OD_{600} = 0.869$),

and N-acetylneuraminate (GF; Max $OD_{600} = 0.439$). Dinstinctly important from

streptomycin-treated mice, D-sorbitol (streptomycin; Max $OD_{600} = 0.202$) and galactitol

(streptomycin; Max $OD_{600} = 0.211$), significantly altered growth compared to negative

controls ($P < 0.001$), however did not overall growth rate (Table S4). Maximum growth

rate analysis for each carbohydrate also indicated potential hierarchy in carbohydrate

preference (Table S4) which is as follows: N-acetyl-D-glucosamine ($k = 0.085$ hr^{-1}),

salicin ($k = 0.077$ hr^{-1}), mannitol ($k = 0.044$ hr^{-1}), N-acetylneuriminate ($k = 0.024$ hr^{-1}),

sorbitol ($k = 0.022$ hr^{-1}), and finally galactitol ($k = 0.02$ hr^{-1}). Although not a

carbohydrate, we also tested acetate for the ability to support *C. difficile* growth *in vitro*,

but showed no improvement over negative controls (Fig. S5). These data suggested

that *C. difficile* was well-suited to adapt its metabolism toward nutrient sources that were

differentially available across susceptible mouse ceca.

Metabolomic analysis supported network-based results for metabolic adaptation

and uncovered additional aspects of *C. difficile*'s nutrient niche. In order to validate

results from transcriptomic and metabolic modeling analyses, we utilized untargeted

mass-specrometry (Table S5) to measure the relative *in vivo* concentrations of

metabolites highlighted by network-based importance scores. We compared cecal

content from mock *C. difficile*-infected susceptible and resistant mice at the same 18

hour time point as the previous analyses. Initially N-acetylglucosamine + N-

acetylgalactosamine (Fig. 7a) and salicylate (Fig. 7b) were found to be significantly

decreased in all susceptible treatment groups compared to resistant controls ($P < 0.05$).

Additionally, we found that galactitol (Fig. 7c), mannitol + sorbitol (Fig. 7d), and N-

acetylneuraminate (Fig. 7e) were significantly increased in both cefoperazone-treated

SPF and GF mice ($P < 0.05$). Finally, these assays also revealed that proline (Fig. 7f)

was significantly increased in all *C. difficile* susceptible conditions ($P < 0.05$), while

glycine (fig. 7g) was found to also be concordantly increased. These amino acids are

required together for *C. difficile* to perform Stickland fermentation. Together

metabolomic results further supported network-derived importance scores and our

hypothesis that *C. difficile* adapts its metabolic strategy in distinct susceptible

environments.

Discussion

Collectively, our results support the hypothesis that *C. difficile* can adapt its metabolism

to the available niche landscape across susceptible gut environments and give insight

to the adaptive strategies that *C. difficile* can use to colonize diverse human microbiota.

All of our *in vivo*, *in vitro*, and *in silico* data demonstrate the plasticity of *C. difficile* to

effectively change its metabolism to utilize alternative resources for growth. This may be

the result of increased concentration of particular metabolites as a consequence of

concordant decreases in the population of one or more competitors for those resources.

In addition to uncovering adaptive strategies of *C. difficile*, our method is also able to

identify consistent trends in metabolism across environments. The findings that N-

acetyl-D-glucosamine and Stickland fermentations substrates were consistently among

the highest scoring shared metabolite among all tested conditions strongly indicates that

these metabolites are central to the nutritional strategy of *C. difficile* and may be utilized

in and condition in which they are available. The metabolism of both substrate types

provides not only carbon and energy to *C. difficile*, but are also a source for nitrogen

which is a limited resource in the mammalian lower GI tract (47).

Apart from exploring differential patterns in known metabolism, our modeling approach

also allowed for the identification of emergent properties for the metabolic strategy of *C.*

difficile during infection. One interesting result is the appearance of CO_2, an apparent

metabolic end product, in the list of shared important metabolites (Fig. 6a). While this

may be a shortcoming of the annotation, one group has posited that *C. difficile* may

actually be autotrophic under certain conditions and could explain our results (48). This

highlights that our method does not only identify growth substrates, but also reports any

metabolites that are being removed from the environment. Although our modeling

results are consistent with previously published work on the metabolism of *C. difficile*,

there are potential limitations of this approach. Ultimately, the metabolite importance

calculation is dependent on correct and existing gene annotation. In this regard it has

been shown that the pathway annotations in KEGG are robust to missing elements (49),

however this does not completely eliminate the possibility for this type of error. Due to

the topology of the metabolic network, we were also unable to integrate stoichiometry

for each reaction which may effect rates of consumption or production. In addition to

computational limitations, our network-based approach simplifies several aspects of

bacterial physiology including the assumptions that all detectable transcript is translated

to effector protein, that each metabolite is available, and that only the neighboring

pathway is important in determining a metabolite's importance. In spite of these

assumptions, the method outlined here supports known elements of *C. difficile* biology

and our subsequent *in vitro* and *in vivo* metabolomic analysis confirmed the predictions

made here.

Based on the evidence presented, our results support the hypothesis that *C. difficile* is a

metabolic generalist and is able to catabolize alternative carbon sources across

susceptible gut environments. This may be due to an inability to outcompete a collection

of metabolic specialists in an intact community, and separate classes on antibiotics

differentially eliminate these populations and allow for *C. difficile* colonization. This may

be particularly true with competitors for proline and glycine as our data also suggests

that overall *C. difficile* colonization resistance may be driven by the availability of

fermentable amino acids. *C. difficile* may also have evolved to utilize the host-derived

aminoglycan N-acetyl-D-glucosamine in all infection conditions as it is a consistently

available nutrient source across susceptible hosts. Additionally, differential importance

scores for carbohydrates in agreement with metabolomic results across distinct

environments, and when considered with variable sporulation and toxin production, may

indicate that *C. difficile* pathogenesis is directed by which carbohydrates are available

for use. Furthermore , competition for nutrients may also potentially explain the success

rate of fecal microbial transplant (FMT), in that wholesale installation of a diverse range

of specialized metabolic strategies is enough to outcompete *C. difficile* from the majority

of perturbed gut environments. In light of these findings, our metabolic network platform

may also prove informative for generating hypotheses through reverse ecology that

could ultimately lead to uncovering new interaction between species that ultimately

impact host health (50). In conclusion, *C. difficile* is is able to optimize its nutritional

strategy for each colonized gut environment. Our results implicate that further

considerations are needed when attempting to design targeted prebiotic and probiotic

therapies for the prevention or elimination of *C. difficile* from the human gut.

Methods

Animal care and antibiotic administration Adapted from the previously described

model (51), six-to-eight week-old SPF C57BL/6 mice were obtained from a single

breeding colony maintained at the University of Michigan for all experiments. Six-to-

eight week-old GF C57BL/6 mice were obtained from a single breeding colony

maintained at the University of Michigan and fed Laboratory Rodent Diet 5001 from

LabDiet for all experiments. All animal protocols were approved by the University

Committee on Use and Care of Animals at the University of Michigan and carried out in

accordance with the approved guidelines. Specified SPF animals were administered

one of three antibiotics; cefoperazone, streptomycin, or clindamycin (Table 1).

Cefoperazone (0.5 mg/ml) and streptomycin (5.0 mg/ml) were administered in distilled

drinking water *ad libitum* for 5 days with 2 days recovery with untreated distilled drinking

water prior to infection. Clindamycin (10 mg/kg) was given via intraperitoneal injection

24 hours before time of infection.

***C. difficile* infection and necropsy** *C. difficile* strain 630 spores were prepared from a

single large batch whose concentration was determined a week prior to challenge for all

experiments. On the day of challenge, 1×10^3 *C. difficile* spores were administered to

mice via oral gavage in phosphate-buffered saline (PBS) vehicle. Subsequent

quantitative plating for c.f.u. was performed to ensure correct dosage. Infection negative

control animals were given an oral garage of 100 µl PBS at the same time as those

mice administered *C. difficile* spores. 18 hours following infection, mice were euthanized

by carbon dioxide asphyxiation. Necropsy was then performed and cecal content was

split into three small aliquots (~100 µl). Two were flash frozen immediately for later DNA

extraction and toxin titer analysis respectively. The third aliquot was quickly moved to an

anaerobic chamber for c.f.u. quantification. The remaining content in the ceca (~1 ml)

was emptied into a stainless steel mortar in a dry ice/ethanol bath using 1 ml of sterile

PBS. This process was repeated for each mouse within a treatment group to pool

content into a single large sample (9 mice across 3 cages) to compensate for cage

effects as much as possible while maximizing sequencing depth. The content was then

finely ground and stored at -80° C for subsequent RNA extraction.

C. difficile cultivation and quantification Cecal samples were weighed and serially

diluted under anaerobic conditions (6% H, 20% CO_2, 74% N_2) with anaerobic PBS.

Differential plating was performed to quantify both *C. difficile* spores and vegetative cells

by plating diluted samles on CCFAE plates (fructose agar plus cycloserine, cefoxitin,

and erythromycin) at 37° C for 24 hours under anaerobic conditions (52). It is important

to note that the germination agent taurocholate was omitted from these plates in order

to only quantify vegetative cells. In parallel, undiluted samples were heated at 60° C for

30 minutes to eliminate vegetative cells and leave only spores (53). These samples

were serially diluted under anaerobic conditions in anaerobic PBS and plated on

CCFAE with taurocholate at 37° C for 24 hours. Plating was simultaneously done for

heated samples on CCFAE to ensure all vegetative cells had been eliminated.

C. difficile **toxin titer assay** To quantify the titer of toxin in the cecum, a Vero cell

rounding assay was performed (54). Briefly, filtered-sterilized cecal content was serially

diluted 1:5 in PBS. As a control for toxin-mediated cell rounding the cecal content was

diluted a further 1:2 by the addition of an equal volume of goat anti-toxin serum (T5000;

TechLab). Vero cells were grown to a confluent monolayer in DMEM (Dulbecco's

Modified Eagle's medium), supplemented with 10% heat-inactivated fetal bovine serum

and 1% penicillin-streptomycin. The cells then were transferred to a conical tube and

centrifuged at 1,000 rpm for 5 minutes to pellet the cells. The old media was removed

and the cells were re-suspended in fresh media to a final concentration of 1×10^5 cells

per 90µL. 90µL of the cell suspension were seeded in each well of a 96-well plate and

incubated at 37° C in a 5% CO_2 humidified incubator for 4 hours. Following the

incubation, cecal samples were added to the Vero cells and the plate was incubated

overnight at 37° C. Plates were viewed after 24 hours at 10x magnification for cell

rounding. The cytotoxic titer was defined as the $\log_{10}$ transformed reciprocal of the

highest dilution that produced rounding in 80% of the cells. A more detailed protocol

with product information can be found at:

https://github.com/jlleslie/Intraspecific_Competition/blob/master/methods/Verocell_Toxin

Activity_Assay.Rmd

16S rRNA gene sequencing DNA was extracted from approximately 50 mg of cecal

content from each mouse using the PowerSoil-htp 96 Well Soil DNA isolation kit (MO

BIO Laboratories) and an epMotion 5075 automated pipetting system (Eppendorf). The

V4 region of the bacterial 16S rRNA gene was amplified using custom barcoded primers

and sequenced as described previously using an Illumina MiSeq sequencer (55). All 63

samples were sequenced on a single sequencing run.

Sequence curation The 16S rRNA gene sequences were curated using the mothur

software package (v1.36), as described previously (55). In short, paired-end reads were

merged into contigs, screened for quality, aligned to SILVA 16S rRNA sequence

database, and screened for chimeras. Sequences were classified using a naive

Bayesian classifier trained against a 16S rRNA gene training set provided by the

Ribosomal Database Project (RDP) (56). Curated sequences were clustered into

operational taxonomic units (OTUs) using a 97% similarity cutoff with the average

neighbor clustering algorithm. The number of sequences in each sample was rarefied to

2,500 per sample to minimize the effects of uneven sampling.

RNA extraction, shotgun library preparation, and sequencing To generate enough

mRNA biomass contributed by *C. difficile*, we pooled cecal content from all mouse

replicates into a single large isolation for each treatment group. Pooling was performed

in a sterile stainless steel mortar resting in dry ice and a small amount of 100% ethanol.

After all content for the given group was added, the sample was ground with a sterile

pestle to a fine powder and scraped into a sterile 50 ml polypropylene conical tube.

Samples were stored at -80° C until the time of extraction. Immediately before RNA

extraction, 3 ml of lysis buffer (2% SDS, 16 mM EDTA and 200 mM NaCl) contained in

a 50 ml polypropylene conical tube was first heated for 5 minutes in a boiling water bath

(57). The hot lysis buffer was added to the frozen and ground cecal content. The

mixture was boiled with periodic vortexing for another 5 minutes. After boiling, an equal

volume of 37° C acid phenol/chloroform was added to the cecal content lysate and

incubated at 37° C for 10 minutes with periodic vortexing. The mixture was the

centrifuged at 2,500 x g at 4° C for 15 minutes. The aqueous phase was then

transferred to a sterile tube and an equal volume of acid phenol/chloroform was added.

This mixture was vortexed and centrifuged at 2,500 x g at 4° for 5 minutes. The process

was repeated until aqueous phase was clear. The last extraction was performed with

chloroform/isoamyl alcohol to remove acid phenol. An equal volume of isopropanol was

added and the extracted nucleic acid was incubated overnight at -20° C. The following

day the sample was centrifuged at 12000 x g at 4° C for 45 minutes. The pellet was

washed with 0° C 100% ethanol and resuspended in 200 µl of RNase-free water.

Following the manufacturer's protocol, samples were then treated with 2 µl of Turbo

DNase for 30 minutes at 37° C. RNA samples were retrieved using the Zymo Quick-

RNA MiniPrep according the manufacturer's protocol. Completion of the reaction was

assessed using PCR for the V4 region of the 16S rRNA gene (Kozich, 2013). Quality

and integrity of RNA was assessed using the Agilent RNA 6000 Nano kit for total

prokaryotic RNA. The Ribo-Zero Gold rRNA Removal Kit Epidemiology was then used

to deplete prokaryotic and eukaryotic rRNA from the samples according the

manufacturer's protocol. Prior to library construction, quality and integrity as measured

again using the Agilent RNA 6000 Pico Kit. Stranded RNA-Seq libraries were made

constructed with the TruSeq Total RNA Library Preparation Kit v2, both using the

manufacturer's protocol. The Agilent DNA High Sensitivity Kit was used to measure

concentration and fragment size distribution before sequencing. High-throughput

sequencing was performed by the University of Michigan Sequencing Core in Ann

Arbor, MI. For all groups, sequencing was repeated across 4 lanes of an Illumina HiSeq

2500 using the 2x50 bp chemistry.

Sequence curation, read mapping, and normalization. Raw transcript sequencing

read curation was performed in a two step process. Residual 5' and 3' Illumina adapter

sequences were trimmed using CutAdapt (35) on a per library basis. Reads were quality

trimmed using Sickle (Joshi, 2011) on the default settings. An average of ~300,000,000

total reads (both paired and orphaned) remained after quality trimming. Mapping was

accomplished using Bowtie2 (58) and the default stringent settings. ~1,600,000 reads in

sample each mapped to the annotated nucleotide gene sequences of *PeptoClostridium*

difficile str. 630 from the KEGG: Kyoto Encyclopedia of Genes and Genomes (59).

Optical and PCR duplicates were then removed using Picard MarkDuplicates

(http://broadinstitute.github.io/picard/), leaving ~150,000 reads per sample for final

analysis. The remaining mappings were converted to idxstats format using Samtools

(60) and the read counts per gene were tabulated. Discordant pair mappings were

discarded and counts were then normalized to read length and gene length to give a per

base report of gene coverage. Unless indicated otherwise, each collection of reads was

then 1000-fold iteratively subsampled to 90% of the lowest sequence total within each

analysis, and a median expression value for each gene was calculated.

Reaction Annotation & Bipartite Network Construction. The metabolism of *C.*

difficile str. 630 was represented as a directed bipartite graph with both enzymes and

metabolites as nodes. Briefly, models were semi-automatically constructed using KEGG

ortholog (KO) gene annotations to which transcripts had been mapped. Reactions that

each KEGG ortholog mediate were extracted from ko_reaction.list located in

/kegg/genes/ko/. KOs that do not mediate simple biochemical reactions (ex. mediate

interactions of macromolecules) were omitted. Metabolites linked to each reaction were

retrieved from reaction_mapformula.lst file located in /kegg/ligand/reaction/ from the

KEGG release. Those reactions that did not have annotations for the chemical

compounds the interact with are discarded. Metabolites were then associated with each

enzyme and the directionality and reversibility of each biochemical conversion was also

saved. This process was repeated for all enzymes in the given bacterial genome, with

each enzyme and metabolite node only appearing once. The resulting data structure

was an associative array of enzymes associated with lists of both categories of

substrates (input and output), which could then be represented as a bipartite network.

The final metabolic network of C. difficile str. 630 contained a total of 1205 individual

nodes (447 enzymes and 758 substrates) with 2135 directed edges. Transcriptomic

mapping data was then re-associated with the respective enzyme nodes prior to

substrate importance calculations. Betweenness-centrality and overall closeness

centralization indices were calculated using the igraph R package found at

http://igraph.org/r/.

Metabolite Importance Calculation. The substrate importance algorithm (Fig. 5a)

favors metabolites that are more likely acquired from the environment (not produced

within the network), and will award them a higher score (Fig. 6b & 6c). The presumption

of our approach was that enzymes that were more highly transcribed were more likely to

utilize the substrates they act on due to coupled bacterial transcription and translation. If

a compound was more likely to be produced, the more negative the resulting score

would be. To calculate the importance of a given metabolite (m), we used rarefied

transcript abundances mapped to respective enzyme nodes. This was represented by t_o

and t_i to designate if an enzyme created or utilized m. The first step was to calculate the

average expression of enzymes for reactions that either created a given metabolite (i) or

consumed that metabolite (ii). For each direction, the sum of transcripts for enzymes

connecting to a metabolite were divided by the number of contributing edges (e_o or e_i) to

normalize for highly connected metabolite nodes. Next the raw metabolite importance

score was calculated by subtracting the creation value from the consumption value to

weight for metabolites that are likely acquired exogenously. The difference was $\log_2$

transformed for comparability between scores of individual metabolites. This resulted in

a final value that reflected the likelihood a metabolite was acquired from the

environment.Untransformed scores that already equaled to 0 were ignored and negative

values were accounted for by transformation of the absolute value then multiplied by -1.

These methods have been written into a single python workflow, along with supporting

reference files, and is presented as bigSMALL (BacterIal Genome-Scale Metabolic

models for AppLied reverse ecoLogy) available in a public Github repository at

https://github.com/mjenior/bigsmall.

Transcriptome Bootstrapping and Probability Distribution Comparison. As

sequencing replicates of *in vivo* transcriptomes was not feasible, we applied a Monte

Carlo style simulation to distinguish calculated metabolite importances due to distinct

transcriptional patterns for the environment measured from those metabolites that were

constitutively important. We employed a 10,000-fold bootstrapping approach of

randomly reassigning transcript abundance for enzyme nodes and recalculating

metabolite importances. This approach was chosen over fitting a simulated

transcriptome to a negative binomial distribution because it created a more relevant

standard of comparison for lower coverage sequencing efforts. Using this method, each

substrate node accumulated a random probability distribution of importance scores

which were then used to calculate the median and confidence interval in order to

ultimately generate a p-value for each metabolite. This was a superior approach to

switch randomization since the connections of the network itself was created through

natural selection and any large-scale alterations would yield biologically uninformative

comparisons(61). These calculations are also included within the standard bigSMALL

workflow presented above and examples can be found in Fig. S6.

Measuring *in vivo* concentrations of acetate. Cecal contents were flash frozen in

liquid nitrogen at the time of necropsy and subjected to short chain fatty acid

quantification analysis using GC-MS (gas chromatography–mass spectrometry) as

described in the targeted metabolomics section of Theriot et al., 2014. All assays were

performed at the Michigan Regional Comprehensive Metabolomics Resource Core in

Ann Arbor, MI.

Anaerobic *in vitro* C. difficile growth curves. The carbon-free variation of *C. difficile*

Basal Defined Medium (NCMM) was prepared as previously described (7). Individual

carbohydrate sources were added at a final concentration of 5 mg/mL and pair-wise

carbohydrate combinations were added at 2.5 mg/mL each (5 mg/mL total). A solution

of the required amino acids was made separately and added when noted at identical

concentrations to the same study. 245 µl of final media mixes were added to a 96-well

sterile clear-bottom plate. A rich media growth control was also included, consisting of

liquid Brain-Heart Infusion + 0.5% cysteine. All culturing and growth measurement were

performed anaerobically in a Coy Type B Vinyl Anaerobic Chamber (3.0% H, 5.0% CO_2,

92.0% N, 0.0% O_2). *C. difficile* str. 630 was grown for 14 hours at 37° C in 3 ml BHI +

0.5% cysteine. Cultures were then centrifuged at 2000 rpm for 5 minutes and resulting

pellets were then washed twice with sterile, anaerobic 1 × phosphate-buffered saline

(PBS). Washed pellets were resuspended in 3 ml more PBS and 5 µl of prepped culture

was added the each growth well of the plate containing aliquoted media. The plate was

then placed in a Tecan Sunrise plate reader, heated to 37° C. Plates were incubated for

24 hours with automatic optical density readings at 600 nm taken every 30 minutes.

OD_{600} values were normalized to readings from wells containing sterile media of the

same type at equal time of incubation. Growth rates and other curve metrics were

determined by differentiation analysis of the measured OD_{600} over time in R to obtain

the slope at each time point.

Quantification of *in vivo* metabolite relative concentrations. Metabolomic analysis

performed by Metabolon (Durham, NC), a brief description of their methods is as

follows. All methods utilized a Waters ACQUITY ultra-performance liquid

chromatography (UPLC) and a Thermo Scientific Q-Exactive high resolution/accurate

mass spectrometer interfaced with a heated electrospray ionization (HESI-II) source

and Orbitrap mass analyzer at 35,000 mass resolution. Samples were dried then

reconstituted in solvents compatible to each of the four methods. The first, in acidic

possitive conditions using a C18 column (Waters UPLC BEH C18-2.1x100 mm, 1.7 µm)

using water and methanol, containing 0.05% perfluoropentanoic acid (PFPA) and 0.1%

formic acid (FA). The second method was identical to the first but was

chromatographically optimized for more hydrophobic compounds. The third approach

utilized a basic negative ion optimized conditions using a separate dedicated C18

column. Basic extracts were gradient eluted from the column using methanol and water,

however with 6.5mM Ammonium Bicarbonate at pH 8. Finally, samples were analyzed

via negative ionization following elution from a hydrophilic interaction chromatography

column (Waters UPLC BEH Amide 2.1x150 mm, 1.7 μm) using a gradient consisting of

water and acetonitrile with 10mM Ammonium Formate, pH 10.8. The MS analysis

alternated between MS and data-dependent MS n scans using dynamic exclusion. The

scan range varied slighted between methods but covered 70-1000 m/z. Library matches

for each compound were checked for each sample and corrected if necessary. Peaks

were quantified using area under the curve.

Statistical methods. All statistical analyses were performed using R (v.3.2.0).

Significant differences between community structure of treatment groups from 16S

rRNA gene sequencing were determined with AMOVA in the mothur software package.

Significant differences of Inv. Simpson diversity, CFU, toxin titer, and acetate

concentration were determined by Wilcoxon rank-abundance test with Holm-Bonferroni

correction. Significant differences for growth curves compared to no carbohydrate

control (+ amino acids) were calculated using 2-way ANOVA with Holm-Bonferroni

correction. Significance for metabolite importance scores was determined as described

above. Wilcoxon rank-abundance test with Holm-Bonferroni correction was also used to

calculate significant differences is relative metabolite concentrations from mass-

spectrometry data.

Acknowledgements

The authors would like to acknowledge Charles Koumpouras for assistance with DNA extractions and metabolomic sample preparation. We would also like to acknowledge members of the University of Michigan Germfree Mouse Center, University of Michigan Sequencing Core, and Metabolon for their assistance in experimental design, execution, and data collection. This work was supported by funding from the National Institutes of Health to PDS (R01GM099514, P30DK034933, U19AI09087, and U01AI124255), VBY (P30DK034933, U19AI09087, and U01AI124255), a Translational Research Education Certificate grant to JLL (MICHR; UL1TR000433), and was partially supported by a predoctoral fellowship from the Cellular Biotechnology Training Program to MLJ (T32GM008353). Pooled and quality trimmed transcriptomic read data and experiment metadata are available through the NCBI Sequence Read Archive (SRA; PRJNA354635). Data processing steps for beginning from raw sequence data to the final manuscript are hosted at http://www.github.com/SchlossLab/Jenior_Modeling_PNAS_2016.

Author Information

Affiliations Department of Microbiology and Immunology, University of Michigan, Ann Arbor, Michigan. Matthew L. Jenior, Jhansi L. Leslie, & Patrick D. Schloss Ph.D.

Department of Internal Medicine & Infectious Diseases, University of Michigan Medical Center, Ann Arbor, Michigan. Vincent B. Young M.D. Ph.D.

Contributions M.L.J. conceived, designed and performed experiments, analyzed data, and drafted the manuscript. J.L.L. performed experiments and analyzed data. V.B.Y. contributed to the manuscript. P.D.S. interpreted data and contributed the manuscript.

727   **Competing interests** The authors declare no competing interest.

728   **Corresponding author** Correspondence to Patrick D. Schloss

729 **Table 1. Antibiotics used during *C. difficile* infection models.**
730

Antibiotic	Class	Target	Activity	Administration	Dosage
Cefoperazone	Cephalosporin (3rd generation)	Primarily Gram-positive bacteria, with increased activity against Gram-negative bacteria	Irreversibly crosslink bacterial transpeptidases to peptidoglycan and prevents cell wall synthesis	Drinking water Ad libitum	0.5 mg/ml
Streptomycin	Aminoglycoside	Active against most Gram-negative aerobic and facultative anaerobic bacilli	Protein synthesis inhibitor through binding the 30S portion of the 70S ribosomal subunit	Drinking water Ad libitum	5.0 mg/ml
Clindamycin	Lincosamide	Primarily active against Gram-positive bacteria, most anaerobic bacteria, and some mycoplasma	Protein synthesis inhibition through binding to the 23s portion of the 50S ribosomal subunit	Intraperitoneal injection	10 mg/kg body weight

Figure Legends

Figure 1 | Experimental time lines for mouse model pretreatments and *C. difficile*

infection. 9 wild-type C57BL/6 mice across 3 cages were included in each treatment

group. **(A)** Streptomycin or **(B)** cefoperazone administered *ad libitum* in drinking water

for 5 days with 2 days recovery with untreated drinking water before infection, **(C)** a

single clindamycin intraperitoneal injection one day prior to infection, or **(D)** no antibiotic

pretreatment (for both SPF control and GF mice). If no antibiotics were administered in

the drinking water, mice were given untreated drinking water for the duration of the

experiment beginning 7 days prior to infection. At the time of infection, mice were

challenged with 1×10^3 *C. difficile* str. 630 spores at the time of infection. Sacrifice and

necropsy was done 18 hours post-challenge and cecal content was then collected.

Figure 2 | *C. difficile* sporulation and toxin activity quantification after 18 hours of

infection. (A) Vegetative *C. difficile* c.f.u. per gram of cecal content. No significant

differences were observed in between any group colonized by *C. difficile*. **(B)** *C. difficile*

spore c.f.u. per gram of cecal content. Significantly more spores were detectable in GF

mice compared to any of the antibiotic-treated SPF groups ($P < 0.05$). **(C)** Toxin titer

from cecal content measured by activity in Vero cell rounding assay. GF mice also

displayed significantly more toxin activity than all other groups ($P < 0.05$). Untreated

mice in a,b,c had no detectable *C. difficile* or toxin activity and were significantly

different from all other groups in each assay ($P < 0.01$). Median values are shown for

each group with significant differences calculated using Wilcoxon rank-sum test with the

Holm-Bonferroni correction. Dotted lines denote the limit of detection for both assays,

and undetectable points are shown just below the limit of detection for clarity.

Figure 3 | Select *C. difficile* gene set expression compared between treatment

group. Relative abundances of *C. difficile* transcript for specific genes of interest,

rarefied to 90% of the total number of reads within each colonized condition shown. **(A)**

Transcription for select genes from the *C. difficile* sporulation pathway with the greatest

variation in expression between the conditions tested. **(B)** Relative abundances of

transcript for genes that encode effector proteins from the *C. difficile* pathogenicity

locus. **(C)** Transcript abundances for genes associated with quorum sensing in *C.*

difficile. **(D)** Transcript relative abundance of select sigma factors which expression or

activity is influenced by environmental metabolite concentrations. Asterisks (*) indicate

genes from which transcript was undetectable.

Figure 4 | *C. difficile* expression of gene sets for carbon metabolism pathways

across antibiotic pretreatments. Ternary plot indicating the relative abundance of

transcripts for all *C. difficile* str. 630 genes across the three colonized antibiotic-treated

conditions (gray points). Raw transcript abundances were iteratively rarefied and the

median expression of each gene was calculated (~24x coverage). Each point

represents a unique gene from the annotated genome of *C. difficile* str. 630 with

position reflecting the ratio of transcription for that gene in all three antibiotic

pretreatments. Transcripts for genes that are over-represented in a single condition are

placed more proximal to the corner associated with that treatment group. Points placed

near the center are equally expressed across all of the conditions measured at 18 hours

post-infection. Points are colored based on inclusion in specific carbon metabolic

pathways, and point sizes within groups of interest were determined based on the

highest expression value for each gene from a single condition. Gene groups from are

shown individually below without abundance information for ease of comparison. Genes

included in each group with normalized transcript abundances can be found in Table

S1, and refer to Fig. S3 for additional explanation of figure interpretation.

Figure 5 | Genome-scale bipartite metabolic modeling results using the

transcriptome of *C. difficile* str. 630 in each colonized environment. (A) Largest

component from the bipartite genome-scale metabolic model of *C. difficile* str. 630. The

complete network contains 447 enzymes and 758 metabolites, with 2135 directed

edges. Size of enzyme nodes is relative to the number of normalized reads mapped to

the corresponding gene. The sizes shown reflect the transcriptome of *C. difficile* str. 630

during infection of cefoperazone-treated mice after 18 hours of infection. Below the

representative network is the metabolite importance algorithm separated into 3

components; (i) relative transcription of reactions consuming a metabolite, (ii) relative

transcription of reactions consuming a metabolite, and (iii) difference of consumption

and creation of the given metabolite. **(B)** The expanded window displays an example of

a single metabolite importance calculation based on local enzyme gene transcription.

White values in the red nodes represent the number of normalized transcript reads

mapping to the gene sequence for each enzyme node. Average expression of input and

output reactions surrounding metabolite **m** are calculated at then the difference of these

values found to get the relative importance of **m**. Log_2 transformation is then performed

for uniform comparison between metabolites.

Figure 6 | Results from network-based metabolite importance calculation and *in*

***vitro* growth with important carbohydrates.** Prior to importance calculation, transcript

abundances for each condition were evenly rarefied for even comparison across

colonized environments (~18x coverage). **(A)** Median shared significant metabolites

among the 50 highest scoring metabolites from each condition ($P < 0.05$). Median

importance scores and pooled random distribution were recalculated per metabolite

using the scores from each condition tested. **(B)** Distinctly important significant

metabolites from each treatment group ($P < 0.05$). The top 25 scoring metabolites from

each group was cross-referenced against each other group resulting in metabolites that

are differentially important between environments. **(C)** *in vitro* growth curves validating

identified growth nutrients from network analysis. One metabolite that is consistently

important to *C. difficile* and at least one metabolite indicated as distinctly important from

each group supported growth significantly more ($P < 0.001$) than no carbohydrate

control (+ amino acids, gray line). Only those carbon sources that significantly improved

C. difficile growth over control are displayed (remainder are located in Table S4).

Significant differences were calculated using 2-Way ANOVA with Holm-Bonferroni

correction.

Figure 7 | Relative *in vivo* concentrations of metabolites highlighted through

network-based analysis Select metabolite concentrations using normalized untargeted

UPLC-MS quantification from cecal content of *C. difficile* susceptible mice compared to

resistant controls. Samples were collected at the same 18 hour timepoint as the

analyses performed in infection experiments (n = 9 mice per group). Highlighted

carbohydrates indlude: **(A)** N–Acetylglucosamine + N–Acetylgalactosamine, **(B)**

Galactitol. **(C)** Mannitol + Sorbitol (These metabolites are quantified simultaneously as

they only differ by chirality), **(D)** Salicylate, and **(E)** N–Acetylneuriminate. **(F)** Proline and

(G) Glycine were also quantified. Paired metabolites were quantified simultaneously as

the only differ by chirality making differentiation impossible. Significant differences

between experimental, susceptible groups and resistant SPF mice ($P < 0.05$) are

denoted by a single asterisk (*) and were determined using Wilcoxon rank-abundance

test with Holm-Bonferroni correction.

Supplementary Figure 1 | Analysis of bacterial community structure resulting

from antibiotic treatment. Results from 16S rRNA gene amplicon sequencing from

bacterial communities of cecal content in both mock-infected and *C. difficile* 630-

infected animals 18 hours post-infection across pretreatment models. **(A)** Non-metric

multidimensional scaling (NMDS) ordination based on Theta$_{YC}$ distances for the gut

microbiome of all conventionally-raised mice used in these experiments (n = 63). All

treatment groups are significantly different from all other groups by AMOVA ($P < 0.001$).

(B) Inverse Simpson diversity for each cecal community from the mice in (A). Cecal

communities from mice not treated with any antibiotics are significantly more diverse

than any antibiotic-treated condition by Wilcoxon test ($P < 0.001$).

Supplementary Figure 2 | Expression of specific KEGG gene families. Abundances

of normalized transcriptomic reads from *C. difficile* str. 630 in each tested condition. **(A)**

All KEGG families. **(B)** Those sub-families within Carbohydrate metabolism. **(C)** Sub-

families within Amino acid metabolism.

Supplementary Figure 3 | Additional explanation for Figure 4 interpretation.

Relative abundance of transcription for *C. difficile* 630 genes during infection across the

3 antibiotic pretreatment models used during this study. Points that are located closer to

a corner are more highly transcribed in the condition associated with that corner

compared to the others. As this shows a 3-dimensional data set in 2 dimensions, there

is an amount of distortion proximal to each corner. Simply put for points that are nearer

to an edge, a greater percentage of their total transcription was contributed by *C.*

difficile colonizing those mice. **(A)** This point represents the transcription for a gene that

is overrepresented in cefoperazone-treated mice. **(B)** This point represents a gene in

which transcripts are equally detectable in all 3 conditions. **(C)** Transcripts for this gene

are only underrepresented in only cefoperazone-treated mice, and are equally

detectable in clindamycin and streptomycin-treated animals.

Supplementary Figure 4 | *in vivo* acetate concentrations with GC-MS analysis

from cefoperazone-treated mouse cecal content. 2 groups of 5 mice each were

pretreated with cefoperazone as outlined Fig. 1b. A single cage was infected with *C.*

difficile in the same fashion as other experiments described here. Animals were

necropsied at 18 hours post-infection and cecal content was flash frozen for later GC-

MS analysis. Significance was determined using Wilcoxon signed-rank test.

Supplementary Figure 5 | Additional growth curves for additional carbon sources

and controls. Significant metabolites from network analysis that did not provide

improved growth over no carbohydrate (+ amino acids) control. Also included is the

negative control of minimal media with no amino acids as well as *C. difficile* growth in

standard Brain-Heart Infusion broth.

Supplementary Figure 6 | Example Mont-Carlo simulation results for a single

metabolite. Shown here are the distributions resulting from 10000-fold Monte-Carlo

simulation of transcript randomization for enzymes interacting with select metabolites.

These scores were derived from transcriptome of *C. difficile* colonizing streptomycin-

treated mice. The black solid line represents the median of the distribution, the red

dotted lines denote the 95% confidence interval, and the blue arrow indicates the actual

measured score for each metabolite. **(A)** Starch (Importance score = 5.492, *P* < 0.01).

(B) D-Glyceraldehyde 3-phosphate (Importance score = -4.139, *P* = n.s.).

Supplementary Table 1 | Sets of genes included in Figure 4 with normalized

abundances and citations.

Supplementary Table 2 | Topology metrics for enzyme and metabolite nodes in

the *C. difficile* str. 630 metabolic network.

Supplementary Table 3 | All metabolites with significant important scores for *C.*

***difficile* in each colonized condition.**

Supplementary Table 4 | Growth curve analysis for each tested carbon source.

Supplementary Table 5 | Normalized untargeted metabolomics results for

molecules highlighted by network analysis.

References

1. Lessa FC, Gould CV, McDonald LC (2012) Current status of *Clostridium difficile*

infection epidemiology. *Clinical infectious diseases : an official publication of the*

Infectious Diseases Society of America 55 Suppl 2(Suppl 2):S65–70.

2. Lessa FC, et al. (2015) Burden of *Clostridium difficile* Infection in the United States.

The New England journal of medicine 372(9):825–834.

3. Leffler DA, Lamont JT (2015) *Clostridium difficile* Infection. *New England Journal of*

Medicine 372(16):1539–1548.

4. Young VB, Schmidt TM (2004) Antibiotic-Associated Diarrhea Accompanied by

Large-Scale Alterations in the Composition of the Fecal Microbiota. *Journal of Clinical*

Microbiology 42(3):1203–1206.

5. Lawley TD, Walker AW (2013) Intestinal colonization resistance. *Immunology*

138(1):1–11.

6. Chen X, et al. (2008) A Mouse Model of *Clostridium difficile*-Associated Disease.

Gastroenterology 135(6):1984–1992.

7. Theriot CM, et al. (2014) Antibiotic-induced shifts in the mouse gut microbiome and

metabolome increase susceptibility to *Clostridium difficile* infection.

doi:10.1038/ncomms4114.Antibiotic-induced.

8. Schubert AM, Sinani H, Schloss PD (2015) Antibiotic-induced alterations of the

murine gut microbiota and subsequent effects on colonization resistance against

Clostridium difficile. mBio 6(4). doi:10.1128/mBio.00974-15.

9. Koenigsknecht MJ, et al. (2015) Dynamics and establishment of *Clostridium difficile*

infection in the murine gastrointestinal tract. *Infection and Immunity* 83(3):934–941.

10. Antunes LCM, et al. (2011) Effect of antibiotic treatment on the intestinal

metabolome. *Antimicrobial Agents and Chemotherapy* 55(4):1494–1503.

11. Ferreyra JA, et al. (2014) Gut microbiota-produced succinate promotes *Clostridium*

difficile infection after antibiotic treatment or motility disturbance. *Cell Host and Microbe*

16(6):770–777.

12. Jump RLP, et al. (2014) Metabolomics analysis identifies intestinal microbiota-

derived biomarkers of colonization resistance in clindamycin-treated mice. *PLoS ONE*

9(7). doi:10.1371/journal.pone.0101267.

13. Freter R, Brickner H, Botney M, Cleven D, Aranki A (1983) Mechanisms that control

bacterial populations in continuous-flow culture models of mouse large intestinal flora.

Infection and Immunity 39(2):676–685.

14. Wilson KH, Perini F (1988) Role of competition for nutrients in suppression of

Clostridium difficile by the colonic microflora. *Infection and Immunity* 56(10):2610–2614.

15. Sebaihia M, et al. (2006) The multidrug-resistant human pathogen *Clostridium*

difficile has a highly mobile, mosaic genome. *Nature genetics* 38(7):779–786.

16. Tracy BP, Jones SW, Fast AG, Indurthi DC, Papoutsakis ET (2012) Clostridia: The

importance of their exceptional substrate and metabolite diversity for biofuel and

biorefinery applications. 23:364–381.

17. Songer JG, Anderson MA (2006) *Clostridium difficile*: An important pathogen of food

animals. 12:1–4.

18. Neumann-Schaal M, Hofmann JD, Will SE, Schomburg D (2015) Time-resolved

amino acid uptake of *Clostridium difficile* 630 Delta-erm and concomitant fermentation

product and toxin formation. *BMC Microbiology*:281.

19. Nawrocki KL, Edwards AN, Daou N, Bouillaut L, McBride SM (2016) CodY-

dependent regulation of sporulation in *Clostridium difficile*. *Journal of Bacteriology*

198(15):2113–2130.

20. Antunes A, et al. (2012) Global transcriptional control by glucose and carbon

regulator CcpA in *Clostridium difficile*. 40:10701–10718.

21. Bouillaut L, Dubois T, Sonenshein AL, Dupuy B (2015) Integration of metabolism

and virulence in *Clostridium difficile*. *Research in Microbiology* 166(4):375–383.

22. Matamouros S, England P, Dupuy B (2007) *Clostridium difficile* toxin expression is

inhibited by the novel regulator TcdC. *Molecular Microbiology* 64(5):1274–1288.

23. Antunes A, Martin-Verstraete I, Dupuy B (2011) CcpA-mediated repression of

Clostridium difficile toxin gene expression. *Molecular Microbiology* 79(4):882–899.

24. Janoir C, et al. (2013) Adaptive strategies and pathogenesis of *Clostridium difficile*

from *in vivo* transcriptomics. *Infection and Immunity* 81(10):3757–3769.

25. Kansau I, et al. (2016) Deciphering adaptation strategies of the epidemic

Clostridium difficile 027 strain during infection through in vivo transcriptional analysis.

PLoS ONE 11(6). doi:10.1371/journal.pone.0158204.

26. Patil KR, Nielsen J (2005) Uncovering transcriptional regulation of metabolism by

using metabolic network topology. *Proceedings of the National Academy of the*

Sciences of the United States of America 102(8):2685–2689.

27. Borenstein E, Kupiec M, Feldman MW, Ruppin E (2008) Large-scale reconstruction

and phylogenetic analysis of metabolic environments. *Proceedings of the National*

Academy of Sciences 105(38):14482–14487.

28. Suthers PF, et al. (2009) Genome-scale metabolic reconstruction Of mycoplasma

genitalium, iPS189. *PLoS Computational Biology* 5(2).

doi:10.1371/journal.pcbi.1000285.

29. Putnam EE, Nock AM, Lawley TD, Shen A (2013) SpoIVA and sipl are *Clostridium*

difficile spore morphogenetic proteins. *Journal of Bacteriology* 195(6):1214–1225.

30. Underwood S, et al. (2009) Characterization of the sporulation initiation pathway of

Clostridium difficile and its role in toxin production. *Journal of Bacteriology*

191(23):7296–7305.

31. Fimlaid KA, et al. (2013) Global Analysis of the Sporulation Pathway of *Clostridium*

difficile. PLoS Genetics 9(8). doi:10.1371/journal.pgen.1003660.

32. Saujet L, Pereira FC, Henriques AO, Martin-Verstraete I (2014) The regulatory

network controlling spore formation in *Clostridium difficile. FEMS Microbiology Letters*

358(1):1–10.

33. Matsuno K, Sonenshein AL (1999) Role of SpoVG in asymmetric septation in

Bacillus subtilis. Journal of Bacteriology 181(11):3392–3401.

34. Lee ASY, Song KP (2005) LuxS/autoinducer-2 quorum sensing molecule regulates

transcriptional virulence gene expression in *Clostridium difficile. Biochemical and*

Biophysical Research Communications 335(3):659–666.

35. Martin MJ, et al. (2013) The *agr* locus regulates virulence and colonization genes in

Clostridium difficile 027. *Journal of Bacteriology* 195(16):3672–3681.

36. Donnelly ML, Fimlaid KA, Shen A (2016) Characterization of *Clostridium difficile*

spores lacking either SpoVAC or DPA Synthetase. *Journal of Bacteriology*

(April):JB.00986–15.

37. Dineen SS, Villapakkam AC, Nordman JT, Sonenshein AL (2007) Repression of

Clostridium difficile toxin gene expression by CodY. *Molecular Microbiology* 66(1):206–

219.

38. Martin-Verstraete I, Peltier J, Dupuy B (2016) The regulatory networks that control

Clostridium difficile toxin synthesis. 8. doi:10.3390/toxins8050153.

39. Potapov AP, Voss N, Sasse N, Wingender E (2005) Topology of mammalian

transcription networks. *Genome informatics International Conference on Genome*

Informatics 16(2):270–278.

40. Ma HW, Zeng AP (2003) The connectivity structure, giant strong component and

centrality of metabolic networks. *Bioinformatics* 19(11):1423–1430.

41. McGill R, Tukey JW, Larsen Wa (1978) Variations of Box Plots. *The American*

Statistician 32(1):12–16.

42. Aboulnaga EH, et al. (2013) Effect of an oxygen-tolerant bifurcating butyryl

coenzyme a dehydrogenase/electron-transferring flavoprotein complex from *Clostridium*

difficile on butyrate production in *Escherichia coli*. *Journal of Bacteriology*

195(16):3704–3713.

43. Fonknechten N, et al. (2010) *Clostridium sticklandii*, a specialist in amino acid

degradation:revisiting its metabolism through its genome sequence. *BMC genomics*

11:555.

44. Jackson S, Calos M, Myers A, Self WT (2006) Analysis of proline reduction in the

nosocomial pathogen *Clostridium difficile. Journal of Bacteriology* 188(24):8487–8495.

45. Karlsson S, Burman LG, Åkerlund T (2008) Induction of toxins in *Clostridium difficile*

is associated with dramatic changes of its metabolism. *Microbiology* 154(11):3430–

3436.

46. Ng KM, et al. (2013) Microbiota-liberated host sugars facilitate post-antibiotic

expansion of enteric pathogens. *Nature* 502(7469):96–9.

47. Fuller MF, Reeds PJ (1998) Nitrogen cycling in the gut. *Annual review of nutrition*

18:385–411.

48. Köpke M, Straub M, Dürre P (2013) *Clostridium difficile* Is an Autotrophic Bacterial

Pathogen. *PLoS ONE* 8(4). doi:10.1371/journal.pone.0062157.

49. Green ML, Karp PD (2006) The outcomes of pathway database computations

depend on pathway ontology. *Nucleic Acids Research* 34(13):3687–3697.

50. Levy R, Borenstein E (2012) Reverse Ecology: from systems to environments and

back. *Advances in experimental medicine and biology* 751:329–345.

51. Theriot CM, et al. (2011) Cefoperazone-treated mice as an experimental platform to

assess differential virulence of *Clostridium difficile* strains. *Gut microbes* 2(6):326–334.

52. Wilson KH, Kennedy MJ, Fekety FR (1982) Use of sodium taurocholate to enhance

spore recovery on a medium selective for *Clostridium difficile. Journal of Clinical*

Microbiology 15(3):443–446.

53. Sorg Ja, Sonenshein AL (2010) Inhibiting the initiation of *Clostridium difficile* spore

germination using analogs of chenodeoxycholic acid, a bile acid. *Journal of Bacteriology*

192(19):4983–4990.

54. Leslie JL, et al. (2015) Persistence and toxin production by *Clostridium difficile*

within human intestinal organoids result in disruption of epithelial paracellular barrier

function. *Infection and Immunity* 83(1):138–145.

55. Kozich J(of M, Schloss P (2013) 16S Sequencing with the Illumina MiSeq Personal

Sequencer. *University of Michigan Health System SOP* 3.1:1–16.

56. Wang Q, Garrity GM, Tiedje JM, Cole JR (2007) Naive Bayesian classifier for rapid

assignment of rRNA sequences into the new bacterial taxonomy. *Applied and*

Environmental Microbiology 73(16):5261–5267.

57. Lopez-Medina E, Neubauer MM, Pier GB, Koh AY (2011) RNA isolation of

Pseudomonas aeruginosa colonizing the murine gastrointestinal tract. *Journal of*

visualized experiments : JoVE (55):6–9.

58. Langmead B, Trapnell C, Pop M, Salzberg SL (2009) Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome. *Genome Biol*:1–10.

59. Ogata H, et al. (1999) KEGG: Kyoto encyclopedia of genes and genomes. 27:29–

34.

60. Li H, et al. (2009) The Sequence Alignment/Map format and SAMtools.

Bioinformatics 25(16):2078–2079.

61. Basler G, Ebenhöh O, Selbig J, Nikoloski Z (2011) Mass-balanced randomization of

metabolic networks. *Bioinformatics* 27(10):1397–1403.

Treatment: Streptomycin   Cefoperazone   Clindamycin   No Antibiotics   No Antibiotics

Mice: SPF   SPF   SPF   GF   SPF

Amino acid catabolism

Amino sugar metabolism

Glycolysis−associated

Polysaccharide metabolism

PTS transporters

ABC sugar transporters

Sugar alcohol metabolism

Fermentation product synthesis

A

(i) $\mu_i = \left(\dfrac{\Sigma t_i}{n(e_o)} \right)$

(ii) $\mu_o = \left(\dfrac{\Sigma t_o}{n(e_i)} \right)$

(iii) $\text{Importance}(m) = \log_2(\mu_i - \mu_o)$

B

● Enzyme node
● Metabolite node

dAdo Aminohydrolase

7
t_o

e_i

Deoxyadenosine (dAdo)
Importance = 6.554

m

e_o

dAdo:PO$_4^{-3}$ Ribosyltransferase

115
t_i

e_o

94
t_i

ATP:dAdo 5'–Phosphotransferase

More likely produced   More likely consumed

−   0   +

Importance Score