JNK signaling in pioneer neurons directs the architectural organization of the

CNS and coordinates the motor activity of the *Drosophila* embryo

Katerina Karkali[1,2,4], Timothy E. Saunders[2], Samuel W. Vernon[3], Richard A. Baines[3],

George Panayotou[4] and Enrique Martín-Blanco[1,2] *

1. Instituto de Biología Molecular de Barcelona (CSIC), Parc Cientific de Barcelona, Baldiri

Reixac 10-12, 08028 Barcelona, Spain

2. Mechanobiology Institute and Department of Biological Sciences, 5 Engineering Drive 1,

National University of Singapore, 117411 Singapore

3. Division of Neuroscience and Experimental Psychology, School of Biological Sciences, Faculty

of Biology, Medicine and Health, University of Manchester, Manchester Academic Health Science

Centre, Manchester, M13 9PL, UK.

4. BSRC Alexander Fleming, 34 Fleming Street, 16672 Vari, Greece

* Correspondence to: embbmc@ibmb.csic.es

ABSTRACT

The Central Nervous System (CNS) organogenesis is a complex process that obeys precise
architectural rules. Yet, both the mechanisms dictating CNS structural organization, and the impact
that this may have on its functionality remain unknown. Analyzing the Drosophila embryo Ventral
Nerve Cord (VNC) morphogenesis, we found that tight control of JNK signaling levels is essential
for its stereotyped architectural organization. The JNK pathway is active in a specific subset of
pioneer neurons, where it, autonomously, regulates the expression of cell adhesion molecules, the
integrity of axons and dendrites, and motoneurons electrophysiological properties. Interfering
during early embryogenesis in the activity of JNK signaling in these cells, it affects their pioneer
functions and leads to cumulative scaffolding defects during the establishment of axonal networks.
This global architectural disarray associates to the failure of the VNC condensation and the
coordination of the embryonic motor activities required for hatching. Altogether, our data point to
the JNK cascade in a subset of pioneer neurons as a key element underpinning VNC architecture,
and reveal critical links between the structural organization of the VNC and its functional
optimization.

INTRODUCTION

The complex nervous system of bilaterians is thought to have emerged from an ancient, diffuse nerve plexus [1], which evolved into a stereotyped segmented polarized structure [2]. Within this framework, functional neural circuits [3] [4] [5] arise as precise 3D arrangements of neurons and glial cells [6]. Several lines of evidence suggest a direct relationship between the 3D architectural organization of the nervous system and its functions. First, the spatial distribution of functional circuits in the cortex of vertebrates extensively overlaps with large-scale microstructure, connectivity, and gene expression patterns [7]. Second, in pursuit of functional efficiency, neural networks likely minimize their structural cost following the wiring optimization principle postulated by Ramón y Cajal [8]. These costs may arise from metabolic requirements, signal delay and attenuation, and complex guidance processes [9] and would increase with interneuronal distance [10]. Lastly, the spatial organization of cortical areas appears subject to evolutionary selection [11].

The nervous system's architecture is the outcome of neuro-glial interactions and axon guidance decisions. Yet, how the nervous system achieves a mechanically balanced organization throughout morphogenesis, while neuronal connectivity is being established, is poorly understood. We postulate; first, that coordinated cell communication events must be triggered to direct the acquisition of a specific 3D neural architecture; and second, that accuracy in nervous system spatial organization/topography is essential for its functionality.

To explore these hypotheses, we made use of the developing *Drosophila* embryo Central Nervous System (CNS), in particular the Ventral Nerve Cord (VNC). The mature VNC is composed of segmentally iterated ganglia with a stereotyped pattern of neurons organized in a 3D array. Neurons are surrounded by a cellular meshwork composed of different types of glia and are linked by cohesive axon bundles that establish longitudinal connectives and transverse commissures along the neuropile [12]. The neuronal cell bodies in the VNC are restricted to the cortex, while sensory axons and interneuron projections mostly accumulate in central domains. The positions of neurons and their projections have been mapped by employing segmentally repeated reference points [13]. As embryonic development proceeds, axonal and dendritic arborizations become more complex and, as a whole, the VNC condenses along the anterior-posterior (AP) axis. During condensation, embryos develop sporadic muscle contractions that initially do not correlate with

any sensory input [14] [15]. Yet, over time, these contractions evolve into peristaltic patterns,

dependent on coordinated motor circuitry activation, that allow larval hatching [16].

Here, we uncover essential functions for the JNK pathway in a subset of pioneer neurons

orchestrating the structural pattern and condensation of the VNC and enabling motor coordination.

We find that early in embryogenesis, JNK signaling modulates pioneering events, ensuring the

topographical arrangement and fasciculation of axons, and regulates cell adhesion molecules

expression. On these bases, JNK facilitates VNC condensation and muscle contractions

coordination. At late stages, JNK activity fine-tunes cell bodies allocation and dendrites

morphology and numbers. Altogether, our data uncovers a fundamental contribution from the JNK

signaling pathway in pioneer neurons to implement CNS architectural organization and to promote

functional optimization.

RESULTS

Modular Architecture of the Ventral Nerve Cord (VNC)

The 3D organization of axonal tracts and neuronal bodies dictates the size and shape of the

Drosophila embryonic VNC. Yet, the spatial distribution of these elements is irregular and noisy,

preventing the dissection of the mechanical processes outlining VNC architecture. To identify

robust structural landmarks along the VNC scaffold, we performed axial correlation analyses of

Fas 2+ axons (**Figures 1A** and **1B**, see also **Movie S1**). Fas 2 is a neural cell adhesion molecule,

which highlights a subset of the longitudinal axonal connectives, and whose loss-of-function leads

to defasciculation [17]. Correlations along the AP axis uncovered a single characteristic domain

in each segment, with a specific (robust) 3D geometrical organization (structural nodes) (**Figure**

1C). These nodes positioned slightly following each posterior commissure (**Figure S1**). At the

nodes, the longitudinal connectives were organized in three compact fasciculated units at precise

dorso-ventral (DV) and medio-lateral (ML) coordinates (**Figure 1D**). Between nodes, the

distribution of fascicles was not correlated between segments (**Figure 1E**). In a posterior direction,

noise increased very fast reaching a maximal randomness (minimal correlation) in a mid-position

amongst the posterior and the next anterior commissure. Noise then slowly decreased towards the

following node. Thus, within a segment, the 3D structural organization of the VNC is conserved

and polarized along the AP axis (**Figure 1F**).

VNC architecture is modulated by JNK signaling

Although the signaling mechanisms directing axon guidance are well understood [18], those

employed to coordinate the structural organization and topography of the nervous system remain

unexplored. Topography refers to the degree of spatial relation between cells and the cells their

axons project to.

We hypothesize that the JNK cascade, which recently emerged as a conserved regulator for axonal

growth and regeneration [19] [20], may play a role in shaping the VNC, in part by mechanically

regulating neural cells. To investigate this possibility, we interfered with the pathway modulating

the kinase activity of the JNK. In most *Drosophila* tissues, the JNK activity is restrained by a

negative feedback loop implemented by the dual-specificity phosphatase Puckered (Puc). *puc*

expression depends on the activity of Basket (Bsk), the *Drosophila* homologue of JNK, and Bsk

function is inhibited by Puc [21]. Although *puc* expression has been reported in the *Drosophila*

embryonic CNS [22], its role, if any, in CNS morphogenesis is unknown.

We found that in *puc^{E69}* and *puc^{B48}* loss-of-function alleles [21], the structural organization of the

VNC was disturbed. The Fas 2+ longitudinal connectives became highly defasciculated and the

intersegmental nerves were unevenly distributed. Axons came together irregularly into medial

positions and Fas 2 expression was enriched in cell bodies (**Figure 1G** and **S3** and **Movie S2**). In

the absence of *puc*, the level of JNK phosphorylation significantly increased in the VNC,

supporting the existence of a negative feedback on JNK activity, mediated by *puc* (**Figure S2**).

Excessive JNK activity affected axonal organization in the DV and ML axes, but not along the AP

direction (**Figure 1H**). The segmental nodes, although less robust and more widely spaced, were

still present (**Figure 1I** and **S3**). The increased internodal distance observed in *puc* mutants was

manifested as a failure of VNC condensation (on average 18% in *puc^{E69}* and 39% in *puc^{B48}* alleles)

(see direct measurements of the VNC length in **Figure S4**). Further, the segmentally iterated AP

axial structural polarity detected in wild type embryos was abolished (**Figures 1F**).

In brief, a tight control of JNK signaling appears to be essential for implementing the architectural

organization of the VNC and for its condensation. This control is exerted, at least in part, by Puc,

which restricts JNK activity and promotes cell viability (**Figure S2**).

JNK signaling is dynamically regulated in the VNC

To determine, when and where is the JNK cascade active in the VNC we studied the dynamics of

puc expression, which is itself an early readout of the pathway [21]. A viable Gal4 line under the

control of the *puc^E69* enhancer [23] highlighted a segmentally iterated subset of neurons (**Figures**

2A and **2B**). An increased number of neurons was also observed at the peripheral nervous system

(PNS) but no cross-reaction was observed with glial markers [24] (**Figures 2C** and **2D**). *puc*

expression in the VNC initiated in a few neurons at the ventral midline at stage 13 (**Figure 2E** and

Movie S3) and expanded to a minimum consensus of 19 *puc+* neurons per hemisegment at

embryonic stage 17 (**Figure 2F** and **Movie S4**).

Dual staining of *puc^E69* embryos with neuronal markers [Even-skipped (Eve) and Engrailed (En)]

enabled identification of most of the *puc+* cells (**Figure S5**). *puc* labeled the aCC, pCC, RP2, U1-

U3 (plus weak/variable expression in U4-U5) and 3 out of the 8 to 10 EL interneurons. Meanwhile,

NL1, NL2, and iVUM5 En+ neurons were *puc+*. Morphological and positional criteria further

enabled the assignment of five other neurons, AD, ADV, dMP2 and 2 VUM En- motoneurons

(**Figures 2G** and **2H**). This pattern of *puc+* neurons was largely conserved in *puc^B48* (**Figure S6**).

In summary, JNK highest activity was detected in VUMs, pCC and possibly dMP2 neurons at the

ventral midline from stage 13 and subsequently in diverse inter and motoneurons projecting

through the longitudinal connectives and the intersegmental nerves.

The level of JNK activity in *puc+* neurons directs the VNC architectural organization

To rule out the possibility that the aberrant architecture of the VNC of *puc* embryos was a

secondary consequence of the general disruption of the embryo anatomy, we restricted the loss or

gain of JNK activity to different subsets of *puc+* neurons.

The RN2-Gal4 line is specifically expressed in the aCC, pCC and RP2 neurons (**Figure 3A**).

Suppressing JNK activity in RN2+ cells by overexpressing Bsk^DN, a dominant negative form of

Bsk, resulted in the disorganization of the Fas 2+ connectives (**Figure 3B**). Spatial correlations

became noisy and irregular, internodal distances enlarged, and the condensation of the VNC failed.

We also overexpressed Hep^CA, a constitutively active form of Hemipterous (Hep), the *Drosophila*

homologue of the JNK kinase MKK7, to hyperactivate the pathway. This strongly affected the

connectives, which retained a juvenile, disorganized appearance, and resulted in internodal

distance enlargement (**Figure 3C**).

The MzVum-Gal4 line is expressed in a few lateral cells and in the 3 VUM motoneurons (**Figure**

3D). Bsk^{DN} expression in these cells altered the fasciculation of Fas 2+ connectives (**Figure 3E**),

while Hep^{CA} expression led to their disorganization, as shown by the noisier spatial correlation

profiles (**Figure 3F**). There was also an increase in the internodal distances.

The CQ2-Gal4 line drives the expression of UAS-linked genes in the U1 to U5 motoneurons

(**Figure 3G**). Neither the inhibition (**Figure 3H**), nor the hyperactivation of the pathway (**Figure**

3I) resulted in any major structural defects. The VNC architecture in both cases (e.g. nodes

positions and spatial correlations) was similar to WT.

Altogether, these data indicate that the final precise architectural organization of the VNC, requires

carefully tuned levels of JNK activity in the early-specified aCC, pCC, and RP2 pioneers and in

the VUM neurons.

JNK signaling in pioneer neurons modulates Fasciclin 2 levels

The role of JNK signaling in the overall 3D organization of the VNC suggests the existence of a

mechanism that propagates the response to JNK activity of pioneers and VUM neurons throughout

the tissue. While growth factors or neurotransmitters can be an important source of instructive

information [25], adhesion molecules themselves (e.g. Fas 2) may also act as key elements for

communication in large-scale processes [26] [27] [28] [29]. In this context, we explored whether

the global effects of *puc* could be associated with Fas 2 deregulation or subcellular mislocalization.

Fas 2 was enriched at the segmental nodes, irrespectively of the genotype (*wt*, and *puc^{E69}* and

puc^{B48} mutant embryos) (**Figures 4A-C**). Yet, at high JNK activity, in the absence of Puc, its level

was reduced by 50%, both in nodes and internodes (**Figure 4D**) and overall, in stage 16 embryo

extracts (**Figure 4E**). An equivalent decrease was also observed throughout the CNS after

hyperactivating JNK in just the RN2+ neurons (**Figure S7**). Importantly, this effect on Fas 2 levels

is autonomously restricted to the aCC, pCC and RP2 neurons and differences in expression outside

targeted cells were marginal (**Figure S8**). Further, restricting JNK hyperactivation to discrete

segments clearly supported autonomous regulation. Employing a parasegment specific PS8/13-

Gal80 transgene [30], we silenced the overexpression of Hep^{CA} by the RN2-Gal4 driver in

segments A4-A9. These embryos showed reduced Fas 2 expression anteriorly but normal levels in

posterior segments (**Figure S9**).

To explore the roles of Fas 2 in the architectural organization of the VNC and its condensation, we

employed a series of allelic combinations between hypomorphic (*Fas 2^{E76}* and *Fas 2-GFP^{CB03613}*)

and null (*Fas 2^{EB112}*) alleles. We found consistent, although mild, VNC condensation defects

associated to an extensive defasciculation of Fas 2+ longitudinal tracts. Segmental nodes were

present but the 3D correlation along the AP axis was less distinct (**Figure 4F-G)** Further, VNC

condensation was reduced, with the VNC length increased compared to WT (**Figure 4H**). These

defects partially resemble *puc* alleles phenotypes and indicate a role for Fas 2 in the structural

refinement of the VNC and its condensation. Yet, they do not support an exclusive role for Fas 2

as the single mediator of JNK activity during VNC morphogenesis. Indeed, the rescue of *puc^{E69}*

mutant embryos by pan-neural (ELAV-Gal4) overexpression of the full-length Fas 2^{PEST+} isoform

[31] fails to suppress VNC structural or condensation defects (**Figure S10**).

In summary, Fas 2 is essential for a correct VNC structural organization and condensation but it is

not the unique mediator of the JNK pathway. Yet, the cell adhesion landscape of pioneer neurons

appears to have a significant role in the relay mechanisms operating downstream of JNK during

tissue shape remodeling (see Discussion).

JNK signaling influences the axodendritic organization of *puc*-expressing neurons

Besides the contribution on modulating adhesion cues (see above), the global effects of JNK

signaling on VNC architecture may be structural. We examined the morphology of *puc*-expressing

neurons [RN2 (aCC, pCC and RP2)] upon Hep^{CA} overexpression for defective structural layouts.

aCC is a motoneuron whose axon gets incorporated into the intersegmental nerve (ISN)[32]. Its

dendritic arbor develops in a distinct region of the ISN myotopic dendritic domain and consists of

8 to 10 dendrites[33] of an average length of 4 μm (**Figure 5A**). pCC is a medially-located

interneuron whose axon extends ipsilateral in an anterior direction [34]. Finally, RP2 is a

motoneuron, whose axon proceeds through the posterior root of the ISN [35] [32] and branches

out between 7 and 9 dendrites, with an average length of 4 μm (**Figure 5A**).

As a consequence of the overexpression of Hep^{CA} in RN2+ neurons, the cell bodies of the aCC,

pCC and RP2 were frequently misplaced and came together towards the midline (compare **Figures 5A** and **5B**). The appearance of their axons was also affected; they thickened (**Figures 5B**), increased in length (for pCC and RP2) (**Figures 5C**) and developed a wiggly morphology (**Figure 5D**). JNK hyperactivation also affected the density (**Figure 5E**) and length (**Figure 5F** and **Movie S5**) of dendrites. Summarizing, increasing JNK activity in specific pioneer neurons leads to local structural and topographical disarrays that could seed for mistaken 3D arrangements and cumulative misrouting.

JNK activity in pioneer neurons is essential for embryonic motor coordination

Considering the roles of JNK signaling on promoting VNC architectural robustness, we investigated their potential functional consequences. To do so, we focused on motor circuits and evaluated late embryonic activity patterns[14, 16] (**Figure 6A** and **6B**) (see Methods, **Figure S11** and **Movie S6**). In wild type conditions, developing embryos execute forward and backward waves of peristalsis by coordinating bilaterally symmetrical muscle contractions [36] [16]. We found that these movements matured through five stages (**Figure 6C**): they began with brief, isolated, unilateral twitches and sporadic head pinching (stage A). This was followed first by a prolonged rest period occasionally interrupted by uncoordinated single contractions (B) and then by an active stage showing frequent incomplete and complete peristalsis (C). Peristalsis then ends with a short resting period (D), which precedes violent head movements and multiple complete peristalsis (E) leading to the rupture of the vitelline membrane and larval hatching (see also **Figure S11** and **Movie S7**).

In the complete absence of *puc* (*puc^{E69}*), the progression of motor patterns was interrupted, and embryos remained indefinitely in an immature stage (A) characterized by occasional unilateral twitches and incomplete peristalsis (**Figure 6D**, **Figure S11** and **Movie S8**). Muscle integrity and functionality were not affected. This systemic failure of motor coordination could be a direct consequence of the increase of JNK activity levels in the CNS in *puc* mutants or the outcome of other non-defined non-autonomous events. To discard this last possibility, we overexpressed Hep^{CA} in RN2 (**Figure 6E** and **6F**) or MzVum cells (**Figure 6G** and **6H**). In both cases, an excess of JNK activity in *puc+* neurons resulted in a phenotype equivalent to that observed in *puc^{E69}*. Only occasional unilateral twitches, head pinches and infrequent incomplete peristalsis were observed. Therefore, the motor defects in *puc* are a direct response to the loss of Puc activity in *puc+* neurons.

The misregulation of the JNK activity in the same neurons appears to be associated to both VNC
structural and embryonic motor coordination defects.

JNK signaling modulates motoneuron firing but not synaptic vesicle transport

Some *puc+* cells are motoneurons and at aberrant levels of JNK activity, their signaling capability
might be impaired. This might result in defects in the embryo's motor coordination independently
of the architectural organization of the VNC.

To test this possibility, we first studied the axonal transport of synaptic vesicles (Synaptotagmin)
in the intersegmental nerve [aCC and RP2 (RN2-Gal4) and VUM motoneurons (MzVum-Gal4)]
(see Methods, **Figure S12** and **Movie S9**). We found that synaptic vesicles in wild type embryos
moved at the same speed and had the same lifetime in aCC/RP2 and VUMs axons, although they
were at higher density in RN2+ neurons. No differences were observed either between anterograde
and retrograde transport. Upon co-expression of Hep^{CA} in RN2 and MzVUM cells, synaptotagmin
transport behavior was largely unaffected (**Figure S12)**, indicating that microtubule-mediated
transport remained fully functional.

To directly address if the JNK activity influences motoneuron function, we undertook patch clamp
recordings from the aCC and RP2 motoneurons, which receive identical cholinergic synaptic
inputs [37]. Neurons were voltage-clamped at -60mV to record endogenously produced
cholinergic synaptic currents [Spontaneously Rhythmic Currents (SRCs)]. We found that
expression of Hep^{CA} in RN2 + neurons did not influence SRC amplitude or frequency (**Figure 7A**
and **7B**), indicating that the synaptic input to aCC/RP2 remained unaffected. However, current
clamp recordings, in which no holding potential was applied (I=0), revealed that the motoneuron
Resting Membrane Potential (RMP) was significantly increased and the membrane hyperpolarized
(**Figure 7C** and **7D**). Consequently, the average number of action potential fired per Excitatory
PostSynaptic Potential (EPSP) was also notably reduced. We conclude that although the synaptic
drive was normal upon JNK hyperactivation, excitability, and thus endogenous firing, was
considerably reduced in pioneer motoneurons due to a more negative RMP.

Summarizing, the JNK signaling pathway affects pioneer neuron morphology and dendrite
integrity but does not affect synaptic vesicle transport or neuron sensory competencies. However,
it modulates their firing response.

Different temporal roles for the JNK pathway

To define if the inputs of the JNK pathway modulating the structural organization and functional optimization of the VNC are causally related or independent, we evaluated the effects of interfering in the JNK activity levels within the VNC at different times.

Programmed temperature shifts between 29 and 18ºC or *vice versa* (see Methods) allowed us to selectively overexpress Hep^{CA} early vs late during embryogenesis. The early-only (stage 10-13) hyperactivation of JNK in RN2+ neurons, before the initiation of neural activity [38], was associated with defects on the morphology of dendrites and axons, with axonal scaffold anomalies and with the collapse of RN2+ neurons towards the midline (**Figure 7E**). VNC condensation was also affected (**Figure 7F**). Furthermore, although these embryos occasionally hatched, they displayed strong muscle coordination defects with an almost complete absence of bilateral peristalsis (**Figure 7G** and **Movie S10**). Conversely, the late hyperactivation of the JNK pathway (stage 15-16) did not affect VNC condensation or peristalsis (**Figures 7F** and **7G** and **Movie S10**). However, it resulted in the misalignment of RN2+ neurons, which displayed scarce and long dendrites, and in the formation of heterogeneous axon bundles (**Figure 7E**).

Surprisingly, the level of expression of Fas 2 after early JNK hyperactivation increased by 20%, in contrast to the reduction observed upon constant JNK hyperactivation (**Figure 7F**). We hypothesize that this increment might be the consequence of the negative impact of Puc on JNK activity. Puc expression would increase in response to the early transient JNK hyperactivation and its activity sustained throughout embryogenesis.

In conclusion, the condensation of the VNC and the coordination of the peristaltic movements leading to larval hatching are both dependent on precise levels of JNK activity in pioneer neurons at the beginning of CNS morphogenesis. Early, JNK modulates the establishment of the pioneer landscape and alterations in its activity induce aberrations in the axonal scaffold and in cell bodies positioning. Late, JNK activity impinges upon the final refinement of the allocation of neurons and axonal tracts (**Figure 7H**).

DISCUSSION

The embryonic VNC has a robust architecture

During VNC morphogenesis, both neurons and glial cells grow, change shape, and rearrange over time to generate a mechanically balanced structure. Functionally, the neuropile is a well-organized structure, with dendrites of motor and sensory neurons terminating and branching in characteristic domains. Yet, anatomically, it shows few signs of organization, apart from certain regularities such as the commissures and a set of longitudinal axon bundles. This paradox may originate in a shortage of knowledge about the structural cues facilitating networking. We have identified by correlative analyses a series of anchoring positions (nodes) with a potential mechanical role. The nodes consist of sets of Fas 2+ axons organized in a stereotyped 3D conformation that are repeated segment by segment (**Figure 1** and **S1**). We speculate that in architectural terms the longitudinal and commissural connectives serve as structural beams and that nodes may act as cross-points that help to evenly distribute any load from compression, tension or shear that could arise as the VNC forms (or that it may suffer once functional). This robust organization may respond to evolutionary conserved needs in terms of mechanics and physical balance [39] [1]. Of course, the Fas 2+ nodes represent only a subset of the elements holding the tensional network of the VNC, since other Fas 2- axons could organize in equivalent structures.

The JNK pathway in pioneer neurons modulates VNC architectural organization and condensation

Several studies have shown the important function of JNK signaling in axonal plasticity, stability or regeneration [40] [41]. However, its roles in directing CNS morphogenesis or implementing its functional coordination have not been addressed.

We found that autonomously increasing JNK activity in early-specified midline VUMs and aCC, pCC and dMP2 pioneers affects the axonal scaffold as a whole (**Figure 5**). These precise neurons are specified and singled out at stage 9 (**Figure 2**) and initiate axonogenesis around stage 11 [34]. As the CNS develops, each neuron makes a number of stereotypic pathfinding decisions as they navigate to their particular targets. This is an orchestrated process that in the VNC initiates when the early ipsilateral interneurons MP1, dMP2, vMP2 and pCC, pioneer a scaffold of longitudinal fascicles (stage 12-13). Pioneer axons are subsequently joined by other axons, ipsilateral and contralateral, projecting in anterior or posterior directions. The longitudinal axons fasciculate and

defasciculate (stage 14 and 15) in a process in which glial cells intercalate between axonal tracts

helping them to distribute in three sets, internal, medial and external in the neuropile (stage 16).

At stage 17 the neuron cell bodies positionally relocate and align linearly along the AP axis [42].

This lengthy and complex process appears to be affected when JNK activity levels are increased

at early stages of development. First, the proper allocation of pioneer neurons along the neuropile

and the development of their axonal paths become abnormal. Then, their inaccurate early

axonogenesis apparently mislead the fasciculation of other followers' axons that encounter an

aberrant guidance landscape. Last, as a result, the execution of the VNC architectural plan and its

condensation are challenged (**Figure 7**).

Our own data (**Figure 4**) and some data from the literature; e.g. the role of DSCAM regulating

circuit level plasticity by inhibiting the receptive field size of retinal bipolar cells [43] point to cell

adhesion as an additional major player in the structural organization at a mesoscopic level of the

CNS. In particular, changes in the adhesion properties (Fas 2 levels) of pioneer neurons (e.g. RN2

and MzVum) upon interference in JNK activity appear to contribute to the global topographical

changes observed in the VNC. Fas 2 is involved in homophilic interactions and its deficit leads to

fasciculation failures and the disruption of the organization of the local axonal network. The shape

and compaction of individual segments is thus affected and, as a consequence, the global

architecture and condensation of the VNC perturbed (**Figure 7H**). Such a direct link between JNK

signaling and adhesion elements has been noticed in other related scenarios; e.g. in the third instar

Drosophila larvae, Bsk promotes the axonal pruning of mushroom body gamma neurons by

reducing the levels of Fas 2 at the membrane [44]. Although our genetic analyses show that Fas 2

is essential for the structural organization and condensation of the VNC, Fas 2 overexpression does

not rescue the effects caused by the loss of Puc activity (**Figure 4**).

The function of the JNK pathway on the structural organization of the VNC may also involve

short-range intercellular communication. Such mechanism has been observed in different

developmental, physiological and pathological paradigms. JNK signaling causes non-

autonomously patterning changes in imaginal discs [45]. It also induces paracrine compensatory

proliferation during regeneration through downstream pathways like Upd, Wg and Dpp [46].

Finally, JNK signaling non-autonomously influences tumor microenvironments by affecting the

transcription of various cell growth- and proliferation-promoting molecules [47]. A further

element involved could be the modulation of cell-cell or cell-matrix contacts. As many as 89 JNK

substrates have been described [48], and amongst them are extracellular effectors as Matrix
Metalloproteases (MMPs) or integrins). Indeed, in third instar larvae, altering the expression of
MMPs or integrins in glial cells results in the elongation of the VNC [49]. Further, integrin mutant
phenotypes partially recapitulate the VNC structural defects observed in *puc* mutants [50].
Whether any of these elements have any role in the structural organization of the axonal network
of the VNC remains to be investigated.

JNK signaling coordinates motor activities

In the *Drosophila* embryo, muscles become active very early and isolated twitches and unilateral
waves of contraction occur prior to the generation of propagated action potentials [38]. Do the
activity of pioneer neurons and the organization of the VNC have any functional significance in
this context? Our data strongly advocate for a close relationship between the acquisition of VNC
architectural robustness and the rhythmic coordination of motor functions. Mutant embryos of
different *puc* alleles and those overexpressing Hep^{CA} in pioneer neurons show severe structural
defects and also slower and uncoordinated muscle twitching; they never hatch. This outcome could
be explained as an indirect result of firing defects associated to the aberrant dendritic arborizations
of pioneer motoneurons (**Figures 5** and **7**) [51]. Our results suggest, however, that muscle
coordination failure is primarily linked to the altered organization of the VNC, consequence of
pioneer cells malfunction upon early hyperactivation of the JNK pathway (**Figure 7**).

Dendrites carry the vast majority of synaptic inputs and their morphology and properties are
essential for propagation of postsynaptic potentials [52]. Conceivably, in RN2 neurons, synaptic
integration will be affected by those changes in dendritic arborization and morphology associated
to high JNK activity. This in turn may alter the frequency or capability of firing action potentials.
A remarkable consequence of increasing JNK activity in pioneer motoneurons is the
hyperpolarization of their membranes (**Figure 7**) and it is well known that the membrane potential
can regulate cell differentiation as well as cortical tissue arrangement, shape and size. Indeed,
hyperpolarizing conditions lead to a reduction of the number of connections in immature neurons
in vitro [53]. Accordingly, in the embryonic VNC, hyperpolarization of RN2 cells after Hep^{CA}
overexpression may in part account for their anomalous dendritic arborization. Less probable
would be its participation in the implementation of motor coordination. If this were the case,
hyperpolarization of RN2 cells should be associated with early hyperactivation of the JNK

pathway, which is unlikely.

Wiring Optimization

It is well established that spontaneous activities that persistently stimulate postsynaptic cells result

in long-term synapse potentiation [54]. These activities propagate efficiently between nearby cells,

ensuring connections are strengthened, whereas those from distant ones are lost [55]. Bringing

neurons in proximity improves functional interconnectedness and improves wiring optimization

[8]. We suggest that these classic principles are followed during the coordination of motor

activities in the *Drosophila* embryo. Several arguments support this premise. First, both VNC

condensation and motor coordination require a precise level of JNK activity at early developmental

stages, enabling activity of pioneer neurons (**Figure 7**). Second, either activation or ablation of the

EL interneurons disrupt bilateral muscle contractions in the larvae [56]. Remarkably, these ELs

make presynaptic contacts with the RP2 motoneurons, which we found critical for the architectural

organization of the VNC and for motor coordination (**Figures 3** and **6**). Third, ELs, and the dorsal

projecting motoneurons they innervate, are nearly all *puc*-expressing neurons (**Figure 2**).

We foresee that the principles of wiring optimization may also apply at a postembryonic level. As

larvae develop, their body wall area grows by two orders of magnitude [57]. To accommodate

such a change, the mechanosensory neurons must extend their dendrites and increase their

receptive field [58], while motoneurons add more synapses to adapt to larger muscles [57] [59].

During scaling, synaptic contacts between nearby neurons are sustained [60]. In this scenario, a

post-synaptic neuron can only connect with nearby specific pre-synaptic sites. The propensity to

form stable local functional synapses appears to operate across development.

Here we uncovered an intimate causal link between the temporal control of the activation of the

JNK pathway in pioneer neurons, the structural organization of axonal networks, VNC

condensation, and embryonic motor coordination (**Figure 7H**). Correlative geometrical and

mechanical studies and further cell type-specific genetic interference analyses will be necessary to

dissect the precise mechanisms underlying this process. Yet, wiring optimization across different

circuits, by enhancing partners' proximity and topographical robustness, appears to be an essential

character associated with morphogenesis of the nervous system.

METHODS

Drosophila strains

The following stocks were used:

$w1118$; $puc^{E69}LacZ/TM3, twi-GFP$ [21]

$w1118$; $puc^{B48}LacZ/ TM3, twi-GFP$ [21]

$w1118$; $UAS-GFP; puc^{E69}I–Gal4/TM6B$ [23]

$w1118$; $UAS-GFP; puc^{B48}–Gal4/TM6B$ [23]

$w1118$; $P\{w[+mC]=UAS-bsk.DN\}2$ (BDSC #6409)

$w1118$; $P\{w[+mC]=UAS-Hep.Act\}2$ (BDSC #9306)

$P\{GawB\}elav[C155]$ (BDSC #6923)

$w1118$; $P\{w[+mC]=eve-Gal4.RN2\}T$, $P\{w[+mC]=UAS-mCD8::GFP.L\}$; $P\{w[+mC]=eve-Gal4.RN2\}G$, $P\{w[+mC]=UAS-mCD8::GFP.L\}$ (Dr. Irene Miguel-Aliaga)

$y1$ $w1118$; $P\{w[+mC]=CQ2-Gal4\}O$, $P\{w[+mC]=UAS-mCD8::GFP.L\}LL6$ (BDSC #7465)

$w1118$ $P\{w[+mW.hs]=GawB\}MzVUM$; $P\{y[+t7.7]w[+mC] = 10xUAS-IVS-mCD8::GFP\}$ $attP40$ (Dr. Irene Miguel-Aliaga)

$w1118$; $G203:UAS-TNT; ZCL2144$ [16] (Dr. Matthias Landgraf)

$w1118$; $; ZCL2144$ [16] (Dr. Matthias Landgraf)

$w1118$; $P\{UAS-syt.eGFP\}2$ (BDSC #6925)

$w1118$; $UAS-Fas 2^{PEST+}/CyO [Nuc::GFP]$ [31] (Dr. Brian McCabe)

$w1118$, $Fas 2^{E76}$ (Dr. Brian McCabe)

$w1118$, $Fas 2^{EB112}/FM7[Actin::GFP]$ (Dr. Brian McCabe)

$w1118$, $Fas 2-GFP^{CB03616}$ (Dr. Christian Klämbt)

$w1118$; $UAS-GFP; HCJ192(iab3)-Gal80 (PS8/13-Gal80)$ [30] (Dr. Welcome Bender)

In all cases, unless otherwise stated, embryos of the $w1118$ strain served as controls.

Genetics

All crosses were performed at room temperature and after 48 hours were shifted to different
temperatures as the individual experiments required.

For temperature shifts, 2 hours embryo collections were done at room temperature in agar plates
and aged for a further 2 hours. Then, they were transferred, either to a 29ºC incubator for 4 hours
and then to an 18ºC incubator for 12 additional hours [Hep^{CA} (10-13)] or to an 18ºC incubator for
12 hours and then to a 29ºC incubator for 4 hours [Hep^{CA} (15-16)]. These timing regimes were
calculated to enhance Hep^{CA} expression during the early pioneering stages or during the final cell
allocation and axonal refinement stages. After the temperature shifts the embryos were processed
for immunocytochemistry or for live imaging.

Immunohistochemistry

Immunostaining of flat-prepped stage 16 *Drosophila* embryos was performed using the following
primary antibodies: mouse anti-Fas 2 (1:100, clone 1D4, DHSB), mouse anti-axons CNS (1:100,
BP 102, DHSB), mouse anti-Futsch (1:1000, 22c10, DHSB), rabbit anti-phospho-JNK (Thr^{183} /
Tyr^{185}) (1:100, Cell Signaling #9251), rabbit anti-Dcp-1 (Asp216) (1:100, Cell Signaling #9578),
rabbit anti-GFP tag polyclonal (1:600, Thermo Fisher Scientific), rat anti-Elav (1:1000, clone
7E8A10, DHSB), mouse anti-Repo (1:100, clone 8D12 DHSB), mouse anti-Even-skipped (1:100,
clone 3C10, DHSB), and mouse anti-Engrailed (1:100, clone 4D9, DHSB).

The secondary antibodies used for detection were: Goat anti-Rabbit IgG (H+L), Alexa Fluor 488
conjugate (A-11008), Goat anti-Rabbit IgG (H+L) Alexa Fluor 555 conjugate (A-21428), Goat
anti-Mouse IgG (H+L) Alexa Fluor 488 conjugate (A-11001), Goat anti-Mouse IgG (H+L) Alexa
Fluor 555 conjugate (A-21422) and Goat anti-Rat IgG (H+L) Alexa Fluor 555 conjugate (A-
21434). All secondary antibodies were used in a dilution of 1:600 and were from Invitrogen.

Sample preparations for immunodetection and image acquisition

Drosophila embryo dissections for generating flat preparations were performed according to [32].
Briefly, flies maintained in apple juice-agar plates at 25ºC were synchronized by repetitive changes
of the juice-agar plate, with a time interval of 1.5 hours. All embryos laid within this time window
were aged for approximately 22 hours at 18ºC, or until reaching mid-stage 16 (3-part gut stage).
At this point embryos were dechorionated with bleach for 1 min, poured into a mesh and rinsed

extensively with water. For dissection, embryos were transferred with forceps on the surface of a

small piece of double-sided tape, adhered on one of the sides of a poly-L-Lysine coated coverslip.

After orienting the embryos dorsal side up and posterior end towards the center of the coverslip,

the coverslip was flooded with saline (0.075 M Phosphate Buffer, pH 7.2). Using a pulled glass

needle the embryos were manually de-vitelinized and dragged to the center of the coverslip, where

they were attached to the coated glass with their ventral side down. An incision on the dorsal side

of the embryo was performed using the glass needle from the anterior to the posterior end of the

embryo. The gut was removed by mouth suction and a blowing stream of saline was used to flatten

their lateral epidermis. Tissue fixation was done with 3.7 % formaldehyde in saline for 10 minutes

at room temperature. After this point standard immunostaining procedures were followed.

In the single case (**Figure S4**) where whole mount embryos were immunostained, embryo fixation

and de-vitelinization was done according to [61]. Both whole mount and flat-prepped

immunostained embryos were mounted in Vectashield anti-fading medium (Vector Laboratories,

USA).

Image acquisition was performed on a Zeiss LSM 700 inverted confocal microscope, using a 63 x

oil objective lens (NA 1.4). Z-stacks spanning the whole VNC thickness were acquired applying

optimal, with respect to pinhole diameter, sectioning with a step size that varied between 0.2 – 0.6

µm. Image processing was performed with Fiji [62].

Live imaging

Dechorionated stage 14 embryos were glued ventral side down on a MatTek glass bottom dish and

they were covered with S100 Halocarbon Oil (Merck) to avoid desiccation. Image acquisition was

performed on a Zeiss LSM 700 inverted confocal microscope, using a 25 X oil immersion lens

(N.A 0.8, Imm Korr DIC M27). Z-stacks spanning the whole VNC thickness, with a 2 µm step

size, were acquired every 6 minutes for a total of 8 hours. Processing of time-lapse data was done

with Fiji [62].

Image analysis

Image analysis and quantification of fluorescence intensity were performed using Fiji [62]. In

immunostainings where quantification of the fluorescent intensity was required, extra care was

taken so that the same antibody batch was used for staining embryos of different genotypes, while

identical confocal acquisition settings were applied.

For calculating the average integrated density of Fas 2 signal in an image, regions of interest

(ROIs) of fixed size were measured at the nodes and internodes positions. Average Integrated

Density values obtained by measuring ROIs in several animals (n stated in Figure legends) were

then pooled for calculating the mean value as well as standard deviations. For those cases where

the quantification of integrated density was performed on a segment basis, the measured values

represent the raw integrated density, since the area measured slightly varies in between animals.

To calculate the percentage of autonomous and non-autonomous reduction in Fas 2 signal, the

GFP signal (RN2 expression) was thresholded and 3D binary masks were created. Masks were

employed to delimit the quantification of Fas 2 levels in different domains.

Statistical analysis

528   All statistical analyses were performed using GraphPad Software (GraphPad Software Inc., La

529   Jolla, CA, USA). In all cases one sample t test was performed and probability values $p<0.05$ were

530   considered as significant.

Correlations Data analysis

Images were scaled to be isotropic in all axes. Deconvolution was found to not significantly

improve the image correlation. Viewed along the AP axis, images were cropped to include only

the VNC, and then the VNC was split into 50 bins, corresponding to 1.65 μm length each. Within

each bin, a maximum intensity projection was performed along the included (typically eleven)

planes in the z-axis. The Matlab function *imregister* was then used to perform the image

registration for both inter- and intra- embryo correlations. The optimizer for *imregister* was defined

with multimodal metric, tolerance of 10^{-6} and 500 iterations. Before image registration, the center

of mass of the two images was aligned so to maximize image overlap. The code was run on a PC

with 128 Gb memory and in the most demanding case (full embryo correlation: 50 x 50 = 2500

correlations) took around one hour per sample.

To create the image correlation traces, the regions of high correlation (e.g. as identified by arrows

in **Figure 1**) were manually identified and then plotted considering the mean correlation along ±

1 row in the correlation matrices. The diagonal values were ignored. Traces were offset along AP

axis to ensure maximum overlap of peaks in the region 25-57 μm to aid comparison between

different datasets. In all figures, individual profiles are plotted in gray, with the mean (black) and

± 1 standard deviation (red) shown.

To analyze the steepness of the spatial correlations along the AP axis, we created *in silico* data

from the real experimental data points - i.e. each *in silico* data point was drawn from a distribution

defined by the mean and standard deviation in the samples. We determined the best linear fit to

each *in silico* data set and repeated 500 times to find the average gradient fit. This was a robust

way to find the steepness as it accounts for experimental noise. To compare between datasets, we

defined the variable (where gradient corresponds to the spatial derivative of the correlation along

the AP axis):

$r = (\max(gradient) - \mathrm{abs}(\min(gradient))) / [(\max(gradient) + \mathrm{abs}(\min(gradient)))/2]$.

Muscle contraction analysis

Stage 17 embryos carrying the G203, ZCL2144 or both GFP muscle markers were dechorionated

and mounted in halocarbon oil (3:1 mix of S700 and S27) following standard procedures. Muscle

contractions were monitored at the embryo's middle plane, using a 25 X oil immersion lens (N.A

0.8, Imm Korr DIC M27) in a Zeiss LSM780 confocal microscope. For each acquisition 1000

frames were obtained with a scanning speed of 200 msec per frame with a time interval between

acquisition cycles of 15 minutes. Muscle activity was monitored up to embryo's hatching or for a

total of 12 hours at room temperature. For the experiments where Hep[CA] over-expression in RN2

and MzVum neurons was required, both control and Hep[CA] overexpressing embryos were filmed

at 29°C for equal time periods. For temperature-shift analyses of animals expressing Hep[CA] in RN2

neurons, embryos of equivalent pre-hatching age were recorded at 200 msec per frame for 500

frames. Post-imaging, movies were processed with Fiji [62] and Kymographs were generated

using the Fiji plugin KymoResliceWide.

Axonal Trafficking

To examine axonal trafficking, stage 17 embryos expressing UAS-Syt-GFP were dechorionated

and mounted laterally in halocarbon oil (3:1 mix of S700 and S27) for live imaging. Synaptic

vesicle transport was monitored in the medial part of the Inter-Segmental Nerve (ISN) using a 40

X oil immersion lens (N.A 1.3, Oil DIC M27) in a Zeiss LSM780 confocal microscope. A constant

frame size of 512x100 pixels, with an XY zoom of 6 and a recording line spacing of 2 were

routinely employed. Single plane movies were acquired with a scanning speed of 1.11 sec, with 1

sec interval for a total of 60 cycles. All movies were processed with Fiji and Kymographs were

generated using the Fiji plugin KymoResliceWide.

The Mean Velocity and the Lifetime parameters for each particle were calculated from the

Kymographs as follows. The particles trajectories were traced and saved as independent ROIs. The

XY coordinates for each ROI were then extracted using the getSelectionCoordinates function in

Fiji of the custom made "Macro coordinates.ijm" shown below:

run("Fit Spline", "straighten");

getSelectionCoordinates(x, y);

x2=0; y2=0; distance=0;

for (i=0; i<x.length; i++) {

if (i>0) {

590   dx = x[i] - x[i-1];

591   dy = y[i] - y[i-1];

592   distance = sqrt(dx*dx+dy*dy);

}

print(i, x[i], y[i], distance);

}

These XY coordinates represent the particles displacement in both space (X values) and time (Y

values), thus permitting the calculation of Mean Velocity and Lifetime.

Analysis of Axonal and dendrite morphology by Segmentation

To describe the axonal and dendrite morphology of the RN2+ neurons, we analyzed images of flat-

prepped embryos immunostained against GFP and Fasciclin 2 using the Simple Neurite Tracer Fiji

plugin [63]. Image segmentation by this tool allows a semi-automated tracing of axons and

dendrites in 3D, providing measurements of total axonal and dendrite length, as well as of dendrite

number directly from skeletonized images.

Western Blotting

Dechorionated stage 16 embryos of the *puc^{E69}LacZ / TM3, twi-GFP* and *puc^{B48}LacZ / TM3, twi-*

GFP lines were used for generating whole embryo extracts for western blot analysis. In detail,

homozygous embryos were separated from their balanced siblings under a fluorescent dissecting

scope based on the *twi-GFP* expression. Approximately 100 embryos of each type were fully

homogenized in 100 μL of iced cold RIPA buffer, supplemented with Protease Inhibitors (Protease

Inhibitor Cocktail Tablets, Roche Life Sciences). Subsequently, the extracts were centrifuged at

3500 g for 15 minutes at 4ºC and the supernatants were collected. Total protein concentration in

the lysates was determined by Bradford assay (Bio-Rad) and 30 μg of sample were analyzed in a

10 % SDS-PAGE. Proteins were blotted on Protran nitrocellulose membrane (Whatman) while the

antibodies employed were: anti-Fas 2 (1:1000, clone 1D4, DSHB) and Pan-actin (1:1000, Cell

Signaling). The ECL Plus chemiluminescent substrate (Pierce, Thermo Fisher Scientific) was used

for antibody detection. Western Blot signal quantification was performed with Fiji by measuring

the average density of each band.

Embryo whole-cell patch-clamp recordings

Dissections and recordings were performed as described[38] at room temperature (20-22°C). Late

stage 17 *Drosophila* embryos were dissected in external saline (in mM: 135 NaCl, 5 KCl, 4

MgCl2·6H2O, 2 CaCl2·2H2O, 5 N-Tris[hydroxymethyl]methyl-2-aminoethanesulfonic acid, and

36 sucrose, pH 7.15). The CNS was exposed through a dorsal-longitudinal cut of the dechorionated

embryonic body wall musculature. Embryonic cuticle was secured to a Sylgard (Dow-Corning,

Midland, Michigan, USA)-coated cover slip using tissue glue (GLUture; WPI, Hitchin, UK) and

larval gut and fat body removed. The glia surrounding the CNS was partially removed using

protease (1% type XIV; Sigma, Dorset, UK) contained in a wide-bore (15 μm) patch pipette.

Whole cell recordings were carried out using borosilicate glass electrodes (GC100TF-10; Harvard

Apparatus, Edenbridge, UK), fire-polished to resistances of between 16-20 MΩ. Cell identity was

confirmed through RN2-GAL4 driven expression of UAS-CD8-GFP. Recordings were made

using a MultiClamp 700B amplifier. Endogenous excitatory synaptic cholinergic input to the

aCC/RP2 motoneurons were recorded in voltage clamp and current clamp sampled at 20 kHz and

low pass filtered at 10 kHz, using pClamp 10.6 (Molecular Devices, Sunnyvale, CA). Only neurons

with an input resistance of $\geq 1G\Omega$ were accepted for analysis.

ACKNOWLEDGEMENTS

We are extremely grateful to all colleagues that have provided us materials, training and guidance

throughout this work as well as for commenting on the manuscript. Amongst them we like to

specially thank Elisabeth Knust, Christian Klambt, Claude Desplan, Matthias Landgraf, Chris Doe,

Andreas Prokop, Natalia Sanchez-Soriano, Alicia Hidalgo, Marta Zlatic, Albert Cardona, Brian

McCabe and Irene Miguel-Aliaga. We also thank our colleagues at the Parc Cientific de Barcelona,

the Mechanobiology Institute of Singapore and the BSRC "Alexander Fleming" of Athens for their

constant encouragement and support. The Martín-Blanco laboratory was supported by funds from

Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia (BFU2014-

57019-P and BFU2017-82876-P) and from Fundación Ramón Areces. TES was supported by a

Singapore National Research Foundation Fellowship (NRF2012NRF-NRFF001-094) and core

support from the Mechanobiology Institute. RAB was supported by a Biotechnology and

Biological Sciences Research Council grant (BBSRC BB/L027690/1) and SWV by a BBSRC

CASE studentship. Work on this project benefited from the Manchester Fly Facility, established

through funds from University and the Wellcome Trust (087742/Z/08/Z).

REFERENCES

1.   Arendt D, Tosches MA, Marlow H. From nerve net to nerve ring, nerve cord and brain--
evolution of the nervous system. Nature reviews Neuroscience. 2016;17(1):61-72. doi:
10.1038/nrn.2015.15.

2.   Qing Yu ZL, Chuanzi OuYang, Lingjun Li. Biologically Active Peptides in Invertebrates:
Discovery and Functional Studies. Colloqium Series on neuropeptides: Morgan & Claypool
Publishers; 2015. p. 182.

3.   Leise EM. Modular construction of nervous systems: a basic principle of design for
invertebrates and vertebrates. Brain research Brain research reviews. 1990;15(1):1-23.

4.   Chen BL, Hall DH, Chklovskii DB. Wiring optimization can relate neuronal structure and
function. Proceedings of the National Academy of Sciences of the United States of America.
2006;103(12):4723-8. doi: 10.1073/pnas.0506806103.

5.   Redies C, Puelles L. Modularity in vertebrate brain development and evolution. BioEssays
: news and reviews in molecular, cellular and developmental biology. 2001;23(12):1100-11. doi:
10.1002/bies.10014.

6.   Sporns O, Betzel RF. Modular Brain Networks. Annu Rev Psychol. 2016;67:613-40. doi:
10.1146/annurev-psych-122414-033634.

7.   Huntenburg JM. A core organizing axis of the human cerebral cortex: PhD Thesis. Freie
Universität Berlin; 2018.

8.   Cajal SRy. Textura del Sistema Nervioso del Hombre y de los Vertebrados. Madrid, Spain:
Nicolas Moya; 1899.

9.   Tessier-Lavigne M, Goodman CS. The molecular biology of axon guidance. Science.
1996;274(5290):1123-33.

10.   Chklovskii DB, Koulakov AA. Maps in the brain: what can we learn from them? Annual
review of neuroscience. 2004;27:369-92. doi: 10.1146/annurev.neuro.27.070203.144226.

11.   Huntenburg JM, Bazin PL, Margulies DS. Large-Scale Gradients in Human Cortical
Organization. Trends Cogn Sci. 2018;22(1):21-31. doi: 10.1016/j.tics.2017.11.002.

12.   Technau GM. Brain development in Drosophila melanogaster: Springer Science+Business
Media, LLC - Landes Bioscience; 2008.

13. Landgraf M, Sanchez-Soriano N, Technau GM, Urban J, Prokop A. Charting the
Drosophila neuropile: a strategy for the standardised characterisation of genetically amenable
neurites. Developmental biology. 2003;260(1):207-25.

14. Crisp SJ, Evers JF, Bate M. Endogenous patterns of activity are required for the maturation
of a motor network. J Neurosci. 2011;31(29):10445-50. doi: 10.1523/JNEUROSCI.0346-11.2011.

15. Suster ML, Bate M. Embryonic assembly of a central pattern generator without sensory
input. Nature. 2002;416(6877):174-8. doi: 10.1038/416174a.

16. Crisp S, Evers JF, Fiala A, Bate M. The development of motor coordination in Drosophila
embryos. Development. 2008;135(22):3707-17. doi: 10.1242/dev.026773.

17. Lin DM, Fetter RD, Kopczynski C, Grenningloh G, Goodman CS. Genetic analysis of
Fasciclin II in Drosophila: defasciculation, refasciculation, and altered fasciculation. Neuron.
1994;13(5):1055-69.

18. Kolodkin AL, Tessier-Lavigne M. Mechanisms and molecules of neuronal wiring: a
primer. Cold Spring Harbor perspectives in biology. 2011;3(6):a001727.

19. Srahna M, Leyssen M, Choi CM, Fradkin LG, Noordermeer JN, Hassan BA. A signaling
network for patterning of neuronal connectivity in the Drosophila brain. PLoS biology.
2006;4(11):e348. doi: 10.1371/journal.pbio.0040348.

20. Koch M, Nicolas M, Zschaetzsch M, de Geest N, Claeys A, Yan J, et al. A Fat-Facets-
Dscam1-JNK Pathway Enhances Axonal Growth in Development and after Injury. Front Cell
Neurosci. 2017;11:416. doi: 10.3389/fncel.2017.00416.

21. Martin-Blanco E, Gampel A, Ring J, Virdee K, Kirov N, Tolkovsky AM, et al. puckered
encodes a phosphatase that mediates a feedback loop regulating JNK activity during dorsal closure
in Drosophila. Genes & development. 1998;12(4):557-70.

22. Martin-Blanco E. Regulatory control of signal transduction during morphogenesis in
Drosophila. The International journal of developmental biology. 1998;42(3):363-8.

23. Pastor-Pareja JC, Grawe F, Martin-Blanco E, Garcia-Bellido A. Invasive cell behavior
during Drosophila imaginal disc eversion is mediated by the JNK signaling cascade. Dev Cell.
2004;7(3):387-99. doi: 10.1016/j.devcel.2004.07.022.

24. Shklover J, Mishnaevski K, Levy-Adam F, Kurant E. JNK pathway activation is able to
synchronize neuronal death and glial phagocytosis in Drosophila. Cell death & disease.
2015;6:e1649. doi: 10.1038/cddis.2015.27.

25.   Foty RA, Steinberg MS. The differential adhesion hypothesis: a direct evaluation.
Developmental biology. 2005;278(1):255-63. doi: 10.1016/j.ydbio.2004.11.012.

26.   Amack JD, Manning ML. Knowing the boundaries: extending the differential adhesion
hypothesis   in   embryonic   cell   sorting.   Science.   2012;338(6104):212-5.   doi:
10.1126/science.1223953.

27.   Grenningloh G, Rehm EJ, Goodman CS. Genetic analysis of growth cone guidance in
Drosophila: fasciclin II functions as a neuronal recognition molecule. Cell. 1991;67(1):45-57.

28.   Neuert H, Deing P, Krukkert K, Naffin E, Steffes G, Risse B, et al. The Drosophila NCAM
homolog Fas 2 signals independently of adhesion. Development. 2020;147(2). Epub 2019/12/22.
doi: 10.1242/dev.181479.

29.   Zhang S, Amourda C, Garfield D, Saunders TE. Selective Filopodia Adhesion Ensures
Robust Cell Matching in the Drosophila Heart. Dev Cell. 2018;46(2):189-203 e4. Epub
2018/07/18. doi: 10.1016/j.devcel.2018.06.015.

30.   Bowman SK, Deaton AM, Domingues H, Wang PI, Sadreyev RI, Kingston RE, et al.
H3K27 modifications define segmental regulatory domains in the Drosophila bithorax complex.
Elife. 2014;3:e02833. Epub 2014/08/02. doi: 10.7554/eLife.02833.

31.   Beck ES, Gasque G, Imlach WL, Jiao W, Jiwon Choi B, Wu PS, et al. Regulation of
Fasciclin II and synaptic terminal development by the splicing factor beag. J Neurosci.
2012;32(20):7058-73. Epub 2012/05/18. doi: 10.1523/JNEUROSCI.3717-11.2012.

32.   Landgraf M, Bossing T, Technau GM, Bate M. The origin, location, and projections of the
embryonic abdominal motorneurons of Drosophila. J Neurosci. 1997;17(24):9642-55.

33.   Furrer MP, Vasenkova I, Kamiyama D, Rosado Y, Chiba A. Slit and Robo control the
development of dendrites in Drosophila CNS. Development. 2007;134(21):3795-804. doi:
10.1242/dev.02882.

34.   Jacobs JR, Goodman CS. Embryonic development of axon pathways in the Drosophila
CNS. II. Behavior of pioneer growth cones. J Neurosci. 1989;9(7):2412-22.

35.   Sink H, Whitington PM. Pathfinding in the central nervous system and periphery by
identified embryonic Drosophila motor axons. Development. 1991;112(1):307-16.

36.   Pereanu W, Spindler S, Im E, Buu N, Hartenstein V. The emergence of patterned
movement during late embryogenesis of Drosophila. Dev Neurobiol. 2007;67(12):1669-85. doi:
10.1002/dneu.20538.

37. Baines RA, Robinson SG, Fujioka M, Jaynes JB, Bate M. Postsynaptic expression of tetanus toxin light chain blocks synaptogenesis in Drosophila. Curr Biol. 1999;9(21):1267-70.

38. Baines RA, Bate M. Electrophysiological development of central neurons in the Drosophila embryo. J Neurosci. 1998;18(12):4673-83.

39. Matheson T. Invertebrate Nervous Systems. Encyclopedia of Life Sciences. 2002:1-6.

40. Hartwig CL, Worrell J, Levine RB, Ramaswami M, Sanyal S. Normal dendrite growth in Drosophila motor neurons requires the AP-1 transcription factor. Dev Neurobiol. 2008;68(10):1225-42. doi: 10.1002/dneu.20655.

41. Vonhoff F, Kuehn C, Blumenstock S, Sanyal S, Duch C. Temporal coherency between receptor expression, neural activity and AP-1-dependent transcription regulates Drosophila motoneuron dendrite development. Development. 2013;140(3):606-16. doi: 10.1242/dev.089235.

42. Hartenstein V. Atlas of Drosophila development: Cold Spring Harbor Laboratory Press; 1993.

43. Simmons AB, Bloomsburg SJ, Sukeena JM, Miller CJ, Ortega-Burgos Y, Borghuis BG, et al. DSCAM-mediated control of dendritic and axonal arbor outgrowth enforces tiling and inhibits synaptic plasticity. Proceedings of the National Academy of Sciences of the United States of America. 2017;114(47):E10224-E33. Epub 2017/11/09. doi: 10.1073/pnas.1713548114.

44. Bornstein B, Zahavi EE, Gelley S, Zoosman M, Yaniv SP, Fuchs O, et al. Developmental Axon Pruning Requires Destabilization of Cell Adhesion by JNK Signaling. Neuron. 2015;88(5):926-40. doi: 10.1016/j.neuron.2015.10.023.

45. Ahmed-de-Prado S, Diaz-Garcia S, Baonza A. JNK and JAK/STAT signalling are required for inducing loss of cell fate specification during imaginal wing discs regeneration in Drosophila melanogaster. Developmental biology. 2018;441(1):31-41. Epub 2018/06/06. doi: 10.1016/j.ydbio.2018.05.021.

46. Pinal N, Calleja M, Morata G. Pro-apoptotic and pro-proliferation functions of the JNK pathway of Drosophila: roles in cell competition, tumorigenesis and regeneration. Open Biol. 2019;9(3):180256. Epub 2019/03/07. doi: 10.1098/rsob.180256.

47. La Marca JE, Richardson HE. Two-Faced: Roles of JNK Signalling During Tumourigenesis in the Drosophila Model. Front Cell Dev Biol. 2020;8:42. Epub 2020/03/03. doi: 10.3389/fcell.2020.00042.

48.   Bogoyevitch MA, Kobe B. Uses for JNK: the many and varied substrates of the c-Jun N-781 terminal kinases. Microbiol Mol Biol Rev. 2006;70(4):1061-95. Epub 2006/12/13. doi: 782 10.1128/MMBR.00025-06.

49.   Meyer S, Schmidt I, Klambt C. Glia ECM interactions are required to shape the Drosophila 784 nervous system. Mech Dev. 2014;133:105-16. Epub 2014/05/27. doi: 10.1016/j.mod.2014.05.003.

50.   Stevens A, Jacobs JR. Integrins regulate responsiveness to slit repellent signals. J Neurosci. 786 2002;22(11):4448-55. Epub 2002/06/01. doi: 20026413.

51.   Katz LC, Shatz CJ. Synaptic activity and the construction of cortical circuits. Science. 788 1996;274(5290):1133-8.

52.   Spruston N, Stuart G, Häusser M. Principles of dendritic integration. In: Spruston N, Stuart 790 G, Häusser M, editors. Dendrites. Oxford: Oxford University Press; 2016.

53.   Ozkucur N, Quinn KP, Pang JC, Du C, Georgakoudi I, Miller E, et al. Membrane potential 792 depolarization causes alterations in neuron arrangement and connectivity in cocultures. Brain 793 Behav. 2015;5(1):24-38. doi: 10.1002/brb3.295.

54.   Hebb D. The Organization of Behavior: A Neuropsychological Theory. John Wiley And 795 Sons Inc; New York; Chapman And Hall Limited; London; 1949.

55.   Eglen SJ, Demas J, Wong RO. Mapping by waves. Patterned spontaneous activity regulates 797 retinotopic map refinement. Neuron. 2003;40(6):1053-5.

56.   Heckscher ES, Zarin AA, Faumont S, Clark MQ, Manning L, Fushiki A, et al. Even-799 Skipped(+) Interneurons Are Core Components of a Sensorimotor Circuit that Maintains Left-800 Right Symmetric Muscle Contraction Amplitude. Neuron. 2015;88(2):314-29. doi: 801 10.1016/j.neuron.2015.09.009.

57.   Keshishian H, Chiba A. Neuromuscular development in Drosophila: insights from single 803 neurons and single genes. Trends Neurosci. 1993;16(7):278-83.

58.   Grueber WB, Jan LY, Jan YN. Tiling of the Drosophila epidermis by multidendritic 805 sensory neurons. Development. 2002;129(12):2867-78.

59.   Rasse TM, Fouquet W, Schmid A, Kittel RJ, Mertel S, Sigrist CB, et al. Glutamate receptor 807 dynamics organizing synapse formation in vivo. Nat Neurosci. 2005;8(7):898-905.

60.   Gerhard S, Andrade I, Fetter RD, Cardona A, Schneider-Mizell CM. Conserved neural 809 circuit structure across Drosophila larval development revealed by comparative connectomics. 810 Elife. 2017;6. doi: 10.7554/eLife.29089.

61.   Patel NH. Imaging neuronal subsets and other cell types in whole-mount Drosophila

embryos and larvae using antibody probes. Methods in cell biology. 1994;44:445-87.

62.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an

open-source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82. doi:

10.1038/nmeth.2019.

63.   Longair MH, Baker DA, Armstrong JD. Simple Neurite Tracer: open source software for

reconstruction, visualization and analysis of neuronal processes. Bioinformatics.

2011;27(17):2453-4. doi: 10.1093/bioinformatics/btr390.

FIGURES

Figure 1

Figure 1. Structural Architecture of the Ventral Nerve Cord (VNC) of the *Drosophila* embryo

A) Maximum projection of the VNC of a stage 16 *wt* embryo stained with Fas 2. See **Movie S1**. Scale Bar is 10 μm.

B) Self cross-correlation matrix of the Z sections of the image in **A** along the AP axis. The color-coded representation, in all figures from now on, shows the correlation level for each possible comparison at each position along the AP axis. In this way, it was possible to identify an oscillatory pattern of hot spots (3D nodes - purple double headed arrows) of high correlation. One conspicuous node of robust correlation per segment was detected (white arrowheads).

C) Image cross-correlation score along the AP axis for **B** (n=6). The cross-correlation profile shows that segmental Fas 2 nodes fell 28 μm apart in *wt* animals and that the internodal maximum noise/minimum correlation maps 9 μm posterior to the node. The grey lines show individual profiles. The black line corresponds to Average values and the red lines to Standard Deviations. The green bars indicate the positions of maximum correlation (nodes).

D) Z sections at two different maximal (green and magenta) correlation positions (from **A**) showing coalescence of Fas 2 connectives at the nodes. DV and ML [Left (L) and R (Right)] axes are indicated by arrows.

E) Z sections at two different minimal (green and magenta) correlation positions (from **A**) showing dispersion and stochastic distribution associated to internodal noise. DV and ML [Left (L) and R (Right)] axes are indicated by arrows.

F) Box plots of the gradient of the spatial correlation along the AP axis, normalized by the difference between the maximum and minimum gradients for *wt, puc^{E69}* and *puc^{B48}* individuals. Boxes denote 95% confidence interval. The mutant values were not distinguishable from zero.

G) Equivalent image to **A** for a *puc^{E69}* embryo. Note the defasciculation and collapse of the longitudinal connectives, specially the most lateral ones. See **Movie S2**. Scale Bar is 10 μm.

H) Self cross-correlation matrix of the Z sections of the image in **D**. In *puc^{E69}* mutants, while correlation matrices are noisier, a single node per segment can still be observed (white arrowheads).

I) Image cross-correlation score along the AP axis for **E** (n=9). Segmental Fas 2 nodes spread 33 μm away in *puc^{E69}* and the internodal minima gets diffuse. Lines and bars are coded as in **C**.

Figure 2

Figure 2. JNK signaling is dynamically regulated during development

A to **D)** Dorsal and ventral sections highlighting the expression of *puc* (green) in the VNC of stage 17 *puc^{E69}I-Gal4 > UAS-GFP* embryo double-stained for the pan-neuronal marker Elav (**A** and **B** - red) or the glial marker Repo (**C** and **D** - red). *puc* is expressed in a subset of neurons in each segment but not in glia. Scale Bar is 10 μm.

E to **E'''')** four different snapshots from **Movie S3** at the indicated time points from the UAS-GFP; *puc^{E69}I*-Gal4 viable line. *puc* expression can be observed from early stage 13 in a few neurons at midline. As development progressed, more and more neurons express *puc*. Scale Bar is 10 μm.

F to **F'''')** Diagram displaying the cumulative expression of *puc* at the same time points as in **E** to **E'''**. A minimum consensus of 19 neurons per hemisegment was defined. At each time point the intensity (defined by visual inspection) was displayed as low (red), medium (blue) and high (green) (see also **Movie S4**).

G) Composition summarizing the expression of *puc* in the CNS [*puc* expressing cells in the VNC at stage 16 including those identified by coexpressing Even-skipped (magenta) or Engrailed (cyan) plus those putatively identified by morphological and positional criteria (yellow - AD, ADV, dMP2 and 2 VUM En-negative motor neurons)]. To facilitate understanding, a scaled Fas 2 expression image from an embryo of the same stage was overimposed to the diagram. Positions of the robustness nodes are highlighted as dark blue lines.

H) Diagram representing the whole set of *puc* expressing cells (compare to G) and their expected axonal contributions to connectives and exiting nerves (each individual cell carried a single color-code). This diagram was over-imposed over a grey mask covering the scaled width and shape of the VNC.

Figure 3

Figure 3. Targeted activation and inhibition of the JNK pathway in the VNC

A) Cartoon representing the full set of *puc* expressing neurons per segment in the VNC
highlighting the cells expressing RN2-Gal4 (turquoise).

B and **C)** From left to right, pattern of Fas 2 expression, self-cross-correlation matrix and image
cross-correlation profiles for VNCs of 16-stage embryos upon expression of UAS-Bsk^{DN} **(B)** and
UAS-Hep^{CA} **(C)** transgenes under the control of the RN2-Gal4 respectively, inhibiting and
hyperactivating JNK activity in the aCC, pCC and RP2 cells. These conditions affect connectives
defasciculation, nodes profiles and VNC condensation. Scale Bar is 10 μm.

D) Cartoon highlighting the *puc+* neurons expressing MzVum-Gal4 (purple).

E and **F)** As **B** and **C** but employing MzVum-Gal4 for interfering in JNK activity in the VUM
neurons. Affecting JNK activity levels in VUMs also affects defasciculation, nodes profiles and
condensation. Scale Bar is 10 μm.

G) Cartoon highlighting the *puc+* cells expressing CQ2-Gal4 (yellow).

H and **I)** As **B** and **C** but employing CQ2-Gal4 for interfering in JNK activity in the U
motoneurons. None of these conditions result in noticeable defects. Scale Bar is 10 μm.

Figure 4

Figure 4. Fasciclin 2 is necessary for VNC condensation and its expression is modulated by
the JNK pathway

A) *wt*, **B)** *puc^{E69}* and **C)** *puc^{B48}* stage 16 embryos Fas 2 immunoreactivity. Maximum projection of
ventral views across three VNC abdominal segments. Overlaid circles represent the fixed area
ROIs used for measuring the Average Integrated Density of Fas 2 signal, either at the node (red)
or at the internode (yellow) positions. Scale Bar is 10 μm.

D) Quantification of the Average Integrated Density of Fas 2 signal at node and internode positions
as indicated. Statistically significant differences in Fas 2 levels ($p < 0.0001$) were detected between
wt and *puc* embryos, both at the node and internode positions.

E) Immunodetection of Fas 2 in extracts from stage 16 *puc* embryos. Extracts of balanced siblings
were used as controls. In control extracts, the antibody against Fas 2 detected a strong band at 95
kD (red arrow), the intensity of which was reduced by half in extracts from *puc* homozygous
embryos. Detection of actin (42 kD) served as loading control for the quantification of the Fas 2
signal. The graph represents the averaged Fas 2 band intensities normalized with respect to actin.

F) Fas 2-GFP expression pattern of a stage 16 embryo mutant for Fas 2 (*Fas 2^{CB03613} / Fas 2^{EB112}*).
Maximum projection of ventral views across three VNC abdominal segments. A general
derangement of the axonal network was observed [compare to the Fas 2-GFP expression pattern
of a stage 16 heterozygous embryo (*Fas 2^{CB03613} / +*) (inset)]. Scale Bar is 10 μm.

G) Self cross-correlation matrix along the A/P axis of the Z sections of the image in **F**. The color-
coded representation shows the correlation level at each axial position. One spread node per
segment was detected.

H) Quantification of the VNC length in μm (average and standard deviation) for each condition.
Statistically significant differences in length ($p < 0.01$ and $p < 0.001$) were detected between the
wt and the different *Fas 2* allelic combinations.

Figure 5

Figure 5. The axonal and dendritic landscape of pioneer neurons is altered upon JNK hyperactivation

A and **B**) Flat-prepped, stage 16 embryos expressing mCD8-GFP (**A**) and mCD8-GFP and Hep^{CA} (**B**) under the control of the RN2-Gal4 line immunostained for GFP. Traces of the aCC (red) and RP2 (green) proximal dendrites are over-imposed. Yellow arrowheads point to differences in RP2 axon morphology. Anterior is up. Scale bar is 10 μm.

C) Histograms showing the average axonal length of the aCC, RP2 and pCC neurons. Statistically significant differences ($p<0.001$) in the axonal length of the RP2 and pCC neurons were detected between control and Hep^{CA} expressing neurons (n= 4 embryos). Control embryos are represented in grey and Hep^{CA} expressing embryos in green.

D) Histograms showing the average axonal wiggliness of the aCC, RP2 and pCC neurons. RP2 axonal wiggliness was found significantly increased after Hep^{CA} expression ($p<0.001$, n= 4 embryos).

E and **F**) Graphs representing the average Dendrite Number (**E**) and average Dendrite Length (**F**) of the aCC and RP2 neurons. In all cases significant differences were scored in the comparison of control and Hep^{CA} expressing embryos ($p<0.001$, n= 4 embryos).

Figure 6

Figure 6. Defective motor coordination results in incomplete peristalsis in *puc^{E69}* embryos

A) Representative mid-plane image of a stage 17 embryo carrying two GFP protein traps expressed at muscle Z-lines (*w; G203; ZCL2144*) (see **Movie S7**). A selected segmented-line (highlighted in yellow) was employed to create kymographs spanning muscle units along the A/P axis (see below) at both sides of the embryo, right (R) and left (L). The evolution overtime of these kymographs was monitored live. Scale bar is 25 μm.

B) Intensity Profile plots generated along the segmented-line selection in **A**. Gray values as a function of distance highlight the distribution of the segmental muscle units along the axis. Muscle contraction is detected by the differences in the distance between consecutive peaks at sequential time points.

C) Kymograph displaying muscle profiles from 17 to 25 hours AEL for the left side of the embryo in **A**. Anterior (A) and posterior (P). Motor coordination maturation occurs in 5 stages of approximated stereotyped length (color coded): A, Intense unilateral twitching; B, general inactivity with occasional uncoordinated contractions; C, initiation of backward and forward peristalsis; D, pre-hatching inactivity period and E, complete peristalsis and specialized hatching head movements.

D) Kymograph displaying muscle profiles from 18 to 23 hours AEL, when the embryo dies, for the left side of a *puc^{E69}* embryo (see **Movie S8**). Anterior (A) and posterior (P). Motor coordination fails and embryos do not progress beyond the unilateral contraction period (stage A).

E and G) Ventral view of stage 17 embryos expressing GFP under the control of the RN2-Gal4 (**E**) and MzVum-Gal4 (**G**) lines co-expressing Hep^{CA} and carrying the muscle Z-lines specific GFP protein trap ZCL2144. Muscles are pseudo-colored in red and neurons in green. Scale bar is 25 μm.

F and H) Kymographs (ZCL2144 marker) revealing occasional unilateral contractions in embryos co-expressing Hep^{CA} in the RN2-Gal4 (**F**) and MzVum-Gal4 (**H**) neurons. For each case, the kymographs correspond to three periods of 1 hour. These embryos never manage to coordinate muscle movements. Both sides, right (R) and left (L) are shown.

Figure 7

Figure 7. JNK activity modulates VNC condensation, neuronal firing and embryo motor coordination

A) Representative traces of endogenously produced cholinergic SRCs from aCC/RP2 motoneurons in late stage 17 embryos in control (*RN2-Gal4 / CyO*) and experimental lines (*RN2-Gal4 > UAS-Hep^{CA}*). Scale Bar: 20 pA / 2s.

B) No significant difference in SRC amplitude (21.7 ± 3.0 vs. 17.0 ± 3.1 pA / pF, control vs. *RN2-Gal4 > UAS-Hep^{CA}* respectively, P = 0.28) or frequency (16.3 ± 3.3 vs. 15.4 ± 2.4 per min, P = 0.83) was observed.

C) Representative traces of endogenously produced cholinergic EPSPs from aCC/RP2 motoneurons in late stage 17 embryos in control (*RN2-Gal4 / CyO*) and experimental lines (*RN2-Gal4 > UAS-Hep^{CA}*). Action potentials are indicated by asterisks. The amplitude of the action potentials is shunted because of the underlying depolarization. Scale Bar: 10 mV / 100ms.

D) Motoneuron resting membrane potential (RMP) is significantly hyperpolarized in *RN2-Gal4 > UAS-Hep^{CA}* (-57.0 ± 7.6 vs. -81.5 ± 2.9 mV, control vs. *RN2-Gal4 > UAS-Hep^{CA}* respectively, P = 0.009). Consequently, the average number of action potentials fired per EPSP was also significantly reduced (20.6 ± 2.4 vs. 9.3 ± 1.7 spikes per EPSP, P = 0.003). All data points are shown on graphs, bars represent mean ± sem.

E) Fas 2 (red) and GFP (green) immunoreactivity of RN2-Gal4 embryos at late stage 16 subjected to different temperature shift conditions and expressing, from left to right: GFP alone (WT) (maintained at 29ºC continuously from stage 10 to 16); or GFP in combination with a *Hep^{CA}* transgene; Hep^{CA} (10-16) (maintained at 29ºC continuously from stage 10 to 16); Hep^{CA} (10-13) (kept at 29ºC between stages 10 and 13) and Hep^{CA} (15-16) (kept at 29ºC during stages 15 and 16). Maximum projection of ventral views across three VNC abdominal segments. Scale Bar is 10 μm. Overexpressing *Hep^{CA}* at early stages affects pioneer functions and provoke the collapse of neurons cell bodies into the midline. At late stages, dendrite arborizations, axons fasciculation and cell bodies alignment are disturbed.

F) Quantification of the VNC length in μm (average and standard deviation) for each condition (**E**) (top). Statistically significant differences in length (p < 0.0001) were detected between WT and (Hep^{CA} 10-16) and (Hep^{CA} 10-13) embryos. Quantification of the Average Integrated Density of Fas 2 signal by segment (bottom). Statistically significant differences in Fas 2 levels (p <

0.0001) were detected between WT and (Hep^{CA} 10-16) and (Hep^{CA} 10-13) embryos. Note the

increased expression of Fas 2 in (Hep^{CA} 10-13) (see Discussion).

G) Kymographs (bright field) displaying muscle profiles at 25 hours AEL (100 sec) for the left

(L) and right (R) sides of embryos expressing Hep^{CA} in RN2+ cells at early (10-13) or late (15-16)

developmental stages. Anterior (A) and posterior (P). Embryos overexpressing *Hep^{CA}* at early

stages in RN2+ neurons never managed to coordinate muscle movements but those overexpressing

Hep^{CA} late, developed bilaterally symmetrical peristalsis (red arrows) and hatched in all occasions

(**Movie S10**).

H) Regulatory network modulating VNC architectural robustness and condensation and

coordinating motor activities. Precise JNK activity levels in early-specified neurons (aCC, pCC,

RP2 and VUMs) are regulated by a negative feedback loop mediated by Puc. Excessive JNK

activity in *puc* mutants or upon early overexpression of Hep^{CA} in aCC, pCC, RP2 and/or VUMs

neurons leads to a general downregulation of Fas 2 expression (autonomous and non-autonomous),

defects in axonal paths and a general affectation of the architectural robustness of the VNC,

preventing its condensation. The coordination of embryonic motor activities and peristaltic

movements is also inhibited, which could be in part consequence of a failure in wiring optimization

arising from the VNC incomplete condensation. Late interference in JNK activity prevents cell

bodies allocation, dendrites correct arborization and axonal late defasciculation. Images show an

otherwise WT embryo at stages 13 and 16 carrying ELAV-GFP [pan-neural expression (green)]

and Hh-RFP [posterior compartment cells (red)] transgenes.

SUPPLEMENTAL FIGURES

Figure S1

Figure S1. Fine localization of robustness nodes

A) Self cross-correlation matrix of the Z sections of the image in **B**. A conspicuous node of robust correlation is highlighted (white arrowhead).

B to **D)** Images for Fas 2 (**B** - green in **D**), BP102 (**C** - red in **D**) and their superimposed image (**D**) of a double-stained 16 stage *wt* embryo. Discontinued red lines indicate the localization of the robustness nodes in relation to the VNC pattern. Scale Bar is 10 μm.

E) Self cross-correlation matrix of the Z sections of the image in **F**. A conspicuous node of robust correlation is highlighted (white arrowhead).

F to **H)** Images for Fas 2 (**F** - green in **H**), 22C10 (**G** - red in **H**) and their superimposed image **(H)** of a double-stained 16 stage *wt* embryo. Discontinued red lines indicate the localization of the robustness nodes in relation to the VNC pattern. Scale Bar is 10 μm.

Figure S2

Figure S2. JNK activity and cell death in *puc* mutants

A) Single (left and right) and combined (center) channels highlighting the expression of Fas 2 (left - green) and anti-P-JNK (right- red) of a stage 17 *wt* embryo. Scale Bar is 10 μm.

B) Equivalent as A for a stage 17 *puc^E69^* embryo. Scale Bar is 10 μm.

C) Equivalent as A for a stage 17 *puc^B48^* embryo. Scale Bar is 10 μm.

In mutant embryos (**B** and **C**) a significant fraction of the P-JNK signal remained in or around cell bodies (yellow arrowheads) and is not found just in axons as in *wt*.

D) Graph displaying the averaged Integrated Density per segment for the P-JNK signal. Statistically significant differences in P-JNK levels ($p < 0.001$) were detected between the *wt* and both mutant conditions.

E) From left to right, stage 17 heterozygous and homozygous *puc^E69^* embryo stained for Fas 2 (green) and Dcp1 (red), which highlights those cells in the process of apoptosis. Scale Bar is 10 μm.

F) From left to right, equivalent images as in **E** for the *puc^B48^* allele. Scale Bar is 10 μm.

G) Graph displaying the averaged Integrated Density per segment for the Dpc1 signal. Statistically significant differences in Dcp1 levels ($p < 0.001$) were detected between *wt* and *puc^B48^* embryos. The Dpc1 levels of *puc^E69^* embryos showed a tendency to be consistently larger than in the *wt*, although they were not statistically significant.

Figure S3

Figure S3. Robustness and modularity of the VNC architecture in *puc^B48* embryos

A) Maximum projection of the VNC of a *Drosophila* 16-stage *puc^B48* embryo stained with Fas 2.

Defasciculation and collapse of longitudinal connectives is widely spread. Scale Bar is 10 μm.

B) Self cross-correlation matrix of **A**. In *puc^B48*, single nodes per segment are consistently

developed (purple double headed arrows).

C) Image cross-correlation score along the AP axis for **B** (n=7). Segmental Fas 2 nodes are on

average at 39 μm from each other in *puc^B48* and the internodal profile flattens. The grey lines show

individual profiles. The black line corresponds to the Average values and the red lines to the

Standard Deviations. The green bars indicate the positions of maximum correlation (nodes).

Figure S4

1055

Figure S4. Loss of VNC condensation in *puc* mutants

A) *wt*, **B)** *puc^{E69}* and **C)** *puc^{B48}* embryos Fas 2 immunoreactivity. Lateral view of whole mounts of

stage 17 embryos. Direct measurements of the VNC length were performed along the outline of

the abdominal segments of the VNC as shown in A (dotted red line). Scale Bar is 10 μm.

D) Quantification of the VNC length in μm (average and standard deviation) for each condition.

Statistically significant differences in length ($p < 0.001$) were detected between the *wt* and *puc^{B48}*

embryos. The length of the VNC of *puc^{E69}* embryos showed a tendency to be consistently longer

than the *wt*, although it was not statistically significant.

Figure S5

Figure S5. Identities of *puc*-expressing neurons

A and **B)** Dorsal and ventral sections highlighting the expression of *puc* (green) in the VNC of

stage 17 *puc^{E69}I-Gal4 > UAS-GFP* embryo double-stained for Even-Skipped (red). Scale Bar is

10 μm.

C) Diagram indicating those *puc* expressing cells also expressing Even-Skipped [magenta - aCC,

pCC, RP2, U1-U3 (plus weak/variable expression in U4-U5) and 3 out of the 8 to 10 EL

interneurons].

D and **E)** Dorsal and ventral sections highlighting the expression of *puc* (green) in the VNC of

stage 17 *puc^{E69}I-Gal4 > UAS-GFP* embryo double-stained for Engrailed (red). Scale Bar is 10 μm.

F) Diagram indicating those *puc*-expressing cells also expressing Engrailed (cyan - NL1, NL2,

and iVUM5 neurons).

Figure S6

1076

Figure S6. *puc^{B48}* expressing neurons

A to D) Dorsal and ventral sections highlighting the expression of *puc* (green) in the VNC of stage

17 *puc^{B48}-Gal4 > UAS-GFP* embryo double-stained for the pan-neuronal marker Elav (**A** and **B** -

red) or the glial marker Repo (**C** and **D** - red). *puc* is expressed in a subset of neurons in each

segment but not in glia. Scale Bar is 10 μm.

E to H) Dorsal and ventral sections highlighting the expression of *puc* (green) in the VNC of stage

17 *puc^{B48}-Gal4 > UAS-GFP* embryo double-stained for Even-Skipped (**E** and **F** - red) or Engrailed

(**G** and **H** - red). Scale Bar is 10 μm.

I) Cartoon representing the full set of *puc^{B48}* expressing neurons per segment in the VNC.

J) Cartoon representing the *puc^{E69}* neurons highlighting the cells expressing *puc^{B48}* (deep blue).

Figure S7

Figure S7. Fasciclin 2 expression under the control of Hep^{CA}

A) *RN2-Gal4* and **B)** *RN2-Gal4 > UAS-Hep^{CA}* stage 16 embryos Fas 2 immunoreactivity.
Maximum projection of ventral views across two-three VNC abdominal segments. Overlaid circles
represent the fixed area ROIs used for measuring the Average Integrated Density of Fas 2 signal,
either at the node (red) or at the internode (yellow) positions. Scale Bar is 10 μm.

C and **D)** Quantification of the Average Integrated Density of Fas 2 signal at node (**C**) and
internode (**D**) positions as indicated. Statistically significant differences in Fas 2 levels (p < 0.001)
were detected between embryos expressing and non-expressing Hep^{CA}, both at the node and
internode positions.

Figure S8

Figure S8. JNK signaling modulates Fasciclin 2 expression

A) *RN2-Gal4 / +* and **B)** *RN2-Gal4 > UAS-Hep^{CA}* stage 16 embryos Fas 2 immunoreactivity. Sum
projection of ventral views across two-three VNC abdominal segments. Panels from left to right
display: 1) Maximum projection of GFP (RN2-Gal4) (green) and Fas 2 (red) expression; 2) Sum
projection of GFP binary masks; 3) Fas 2 expression associated to RN2^{+} cells; and 4) Fas 2
expression associated to RN2^{-} cells. Scale Bar is 10 μm. **C)** Graphic representation of the reduction
of Fas 2 Average Integrated Density per segment between conditions **B** (HepCA overexpressing
embryos; n=13) and **A** (WT controls; n=6). Statistically significant differences in Fas 2 levels were
detected for whole segments (0.13; p< 0.01) and for RN2+ cells (0.25; P<0.001) but not for RN2-
cells (0.07; P>0.05). JNK hyperactivation in RN2+ cells leads to the autonomous reduction of Fas
2 expression by 25%.

Figure S9

1109

Figure S9. JNK signaling autonomously modulates Fasciclin 2 expression

A) Cartoon showing a schematic view of the embryonic CNS highlighting the territory of expression of a Gal80 transgene (PS8-Gal80) under the control of the *iab3* promoter (parasegments 8 to 13) (pale yellow box). Thoracic (pale green) and abdominal (dark green) segments are indicated. **B)** Maximum projection of ventral views across three VNC segments of *RN2-Gal4 > UAS-GFP* (i and ii) and *RN2-Gal4 > UAS-GFP; UAS-Hep^{CA}* (iii and iv) carrying PS8-Gal80. GFP expression is shown in green and Fas 2 in red. GFP is highly expressed in RN2 cells in the thoracic and anterior abdominal segments (i and iii), while its expression is strongly reduced in the posterior abdominal segments (ii and iv). Scale Bar is 10 μm. **C)** Quantification of the Average Integrated Density of Fas 2 signal per segment. Statistically significant differences in Fas 2 levels (p < 0.01) were detected between embryos expressing and non-expressing Hep^{CA}, only in areas anterior to the parasegment 8. Restricting JNK hyperactivation to discrete segments confirm that Fas 2 expression was autonomously regulated by JNK.

Figure S10

Figure S10. Fasciclin 2 does not rescue Hep^{CA} overexpression phenotypes

A) Superimposed images for Fas 2 (green) and HRP-TRITC (red) of double-stained 16 stage *ELAV/+, puc^{E69}, ELAV-Gal4 > UAS-Fas 2^{PEST+},* and *ELAV-Gal4 > UAS-Fas 2^{PEST+}; puc^{E69}* embryos. Neural expression of Fas 2^{PEST+} in a *puc^{E69}* background exacerbates axonal network defects. Scale Bar is 10 μm. **B)** Quantification of the VNC length in μm (average and standard deviation) for each condition. Statistically significant differences in length (p < 0.001) were detected between *wt* and *puc^{E69}* embryos regardless of the presence of Fas 2^{PEST+}. Pan-neural overexpression of the full length Fas 2^{PEST+} isoform fails to suppress VNC structural or condensation defects.

Figure S11

A

Unilateral Constrictions

B

Head Movements

C

Forward Peristalsis

D

Unilateral Constrictions

Incomplete Peristalsis

Complete Peristalsis

Head Movements

E

1133

Figure S11. Types of embryonic muscle movements

Snapshots from **Movie S6** displaying mid-plane images of a stage 17 embryo carrying two GFP

protein traps expressed at muscle Z-lines (*w; G203, ZCL2144*). Three main types of movement

were observed during the muscle coordination process. Red arrows point to the contracting muscle

units. Anterior is to the left. Scale bar is 50 µm.

A) Unilateral muscle contractions. They can occur at any embryo side and in different segments.

B) Head turning and mouth hook pinching.

C) Peristalsis. They can be incomplete or complete and develop forward (in the image) or

backward displacements.

D) Histograms showing the progression of the muscle coordination process in wild type embryos.

Over-imposed colors follow the 5 stages of motor coordination maturation described in **C**. Muscle

movements were distributed in four classes: Unilateral contractions [sum of the contractions

occurring at the right (green) and the left (purple) side of the embryo]; Incomplete peristalsis [sum

of the incomplete forward (blue) and incomplete backward (red) peristaltic attempts]; Complete

peristalsis [sum of the full forward (blue) and backward (red) peristalsis]; and Head movements

[sum of head turning (red) and mouth hook pinching events (blue)].

E) Histograms equivalent to **D** showing the unsuccessful muscle coordination of puc^{E69} embryos.

Figure S12

Figure S12. JNK activity does not affect synaptic vesicle transport

A and **B**) Flat-prepped, stage 16 embryos expressing Synaptotagmin-GFP under the control of the RN2-Gal4 (**A**) and MzVum-Gal4 (**B**) lines immunostained for GFP (gray scale - top panels and green - bottom panels) and Fas 2 (red - bottom panel). The grey rectangle marks the medial part of the ISN, where synaptic vesicles axonal transport was monitored (see **Movie S9**). Scale bar is 25 μm.

C and **D**) Histograms showing synaptotagmin vesicles density and directional motility in control (grey) versus Hep^{CA} expressing (green) aCC/ RP2 (**C**) and VUM motoneurons (**D**). No statistically significant differences were observed between control and JNK over-active neurons.

E and **F**) Lifetime range distribution of motile synaptotagmin vesicles in aCC/RP2 (**E**) and VUM (**F**) motoneurons. No differences were observed in lifetime between control (grey) and Hep^{CA} expressing (green) embryos.

G and **H**) Mean velocity range distribution comparison of motile synaptotagmin vesicles in aCC/RP2 (**G**) and VUM (**H**) motoneurons between control (grey) and JNK gain-of function (green) conditions. For both sets of motoneurons the distribution was unaltered.

SUPPLEMENTAL MOVIES LEGENDS

Movie S1. Fas 2 positive connectives

Animated 3D reconstruction of the Fas 2 positive connectives of a section of the VNC of a late embryo (Stage 17).

Movie S2. Fas 2 positive connectives of a *puc^{E69}* embryo

Animated 3D reconstruction of the Fas 2 positive connectives of a section of the VNC of a late *puc^{E69}* mutant embryo (Stage 17).

Movie S3. Dynamics of *puc^{E69}* expression

Time course of puc^{E69} expression from a $puc^{E69}I$-$Gal4 > UAS$-GFP viable line. puc expression can be observed from early stage 13 in a few neurons at the midline. As development progressed, more and more neurons express puc, both within the VNC and at the peripheral nervous system (PNS). Scale bar is 50 μm. Anterior is to the left.

Movie S4. Minimum consensus for the segmental puc^{E69} expression

Animated 3D cartoon representing a single segment of a late puc^{E69} mutant embryo (Stage 17). The 19 puc expressing cells per hemisegment represented in the minimum consensus are shown in different colors.

Movie S5. Axonal and dendritic landscape of pioneer neurons upon JNK hyper-activation

Animated 3D reconstruction of the axonal and dendritic pattern of stage 16 embryos expressing mCD8-GFP (**A**) and mCD8-GFP and Hep^{CA} (**B**) under the control of the RN2-Gal4 line. Traces of the dendrites (purple) are over-imposed. Scale bar is 10 μm.

Movie S6. Embryonic motor patterns

Representative mid-plane movies of a stage 17 embryo carrying two GFP protein traps expressed at muscle Z-lines (*w; G203; ZCL2144*). **A)** Forward peristalsis. **B)** Backward Peristalsis. **C)** Left twitch. **D)** Right twitch. **E)** Head tumble. **F)** Mandible pinch. Scale bar is 50 μm. Anterior is to the left.

Movie S7. Time course of motor maturation

Representative mid-plane movies (250 time points per stage) of an embryo carrying two GFP protein traps expressed at muscle Z-lines (*w; G203; ZCL2144*). From top to bottom stages **A** (Unilateral twitches and incomplete peristalsis), **B** (Resting period), **C** (Complete peristalsis), **D** (Late resting period) and **E** (Hatching movements). Color coded as in **Figure 6**. Scale bar is 50 μm. Anterior is to the left.

Movie S8. Motor activity of puc^{E69} embryos

1209 Representative mid-plane movie (1000 time points) of a late stage 17 *puc^{E69}* embryo carrying two

1210 GFP protein traps expressed at muscle Z-lines (*w; G203; ZCL2144*). Scale bar is 50 µm. Anterior

1211 is to the left.

1212

1213 Movie S9. Synaptic vesicles axonal transport in RN2 neurons

1214 Stage 16 embryos expressing Synaptotagmin-GFP under the control of the RN2-Gal4 in the

1215 absence **(A)** or presence **(B)** of Hep^{CA}. Left panels present representative single plane movies (Fire

1216 Luts) with 1 sec interval for a total of 60 cycles. Right panels display in different colors the traces

1217 of the different particles analyzed. Proximal is to the left. Scale bar is 25 µm.

1218

1219 Movie S10. Motor coordination responses to early and late JNK hyperactivation

1220 Representative mid-plane movies of stage 17 embryos expressing Hep^{CA} in RN2+ cells at early

1221 (13 - top) or late (16 - bottom) developmental stages. Embryos over-expressing Hep^{CA} at early

1222 stages never coordinate muscle movements but those overexpressing Hep^{CA} late, developed

1223 multiple bilaterally symmetrical peristalsis and hatched. Scale bar is 50 µm

1224