**The JNK signaling links the CNS architectural organization to motor**

**coordination in the *Drosophila* embryo**

**Katerina Karkali[1] [2] [3],George Panayotou[2], Timothy E. Saunders[3] and Enrique Martín-**

**Blanco[1] [3] [*]**

1. Instituto de Biología Molecular de Barcelona (CSIC), Parc Cientific de Barcelona, Baldiri

Reixac 10-12, 08028 Barcelona, Spain

2. BSRC Alexander Fleming, 34 Fleming Street, 16672 Vari, Greece

3. Mechanobiology Institute, 5 Engineering Drive 1, National University of Singapore, 117411

Singapore

* Correspondence to: embbmc@ibmb.csic.es

**ABSTRACT**

In any organism, the construction of the Central Nervous System (CNS) is a complex process
that must comply with precise architectural rules. How it is attained and how it affects neural
functionality are key issues to solve. In the Drosophila Ventral Nerve Cord (VNC), we
uncovered a set of novel robust architectural elements (3D nodes) arranged in a precise pattern
responding to a delicate balance on Jun-N-terminal kinase (JNK) signaling activity in specific
pioneer neurons. This cascade controls axons and dendrites integrity and, non-autonomously,
Fasciclin 2 (Fas 2) expression. A deficit in Fas 2 impairs the VNC architecture and condensation.
Remarkably, the coordination of the embryonic motor activity essential for embryo hatching also
responds to JNK activity in the same cells. Altogether, our data point to the JNK pathway as an
essential element underpinning VNC robustness and reveal critical links between the architecture
of the VNC and its functional optimization.

**INTRODUCTION**

In higher organisms, behavioral and cognitive functions are directed and coordinated by neural networks [1-3]. Networks interconnect single neural modules, which are comprised of sets of neurons and glial cells with a specific operative design that are arranged in stereotyped structures conserved during evolution [4]. It is still poorly understood: (1) how neural networks acquire their architecturally stable 3D structure; (2) how their spatial organization influences neural functions.

To explore these problems in depth we focused on the *Drosophila* embryonic Central Nervous System (CNS), and in particular in the Ventral Nerve Cord (VNC). The VNC of the *Drosophila* embryo is composed of segmentally iterated ganglia with a stereotyped pattern of neurons organized in a particular 3D array. The neuronal cell bodies are restricted to the cortex, while sensory axons and interneuron projections mostly accumulate in the central neuropile. The positions of these neurons and their projections can be mapped employing segmentally repeated reference points [5]. Cohesive axon bundles build longitudinal connectives and establish transverse commissures [6].

As development proceeds, the VNC autonomously condenses along the anterior/posterior (A/P) axis into a compact bilaterally symmetrical structure. In parallel, the nervous system functionally matures and the sporadic activities of individual motor neurons, interneurons and sensory neurons integrate into specific patterns to direct the motor activities responsible for larval hatching [7]. This maturation into coordinated crawling-like behaviours proceeds autonomously, independent of any sensory input, [8, 9].

Many considerations suggest a direct relationship between the 3D structural organization of the nervous system and its function. Classical neuroanatomy points to evolutionary selection as a key element guiding the spatial organization of cortical areas [10]. Further, in the cortex of vertebrates, large-scale trends in microstructure, connectivity, and gene expression extensively overlap with the spatial distribution of functional processing hierarchies [11]. Last, in pursuit of functional efficiency, neural modules should minimize their wiring cost following the wiring optimization principle postulated by Ramón y Cajal [12]. These costs may arise from metabolic requirements, signal delay and attenuation, and potential guidance defects [13] and would

increase with interneuronal distance [14].

The Jun-N-terminal (JNK) pathway has recently emerged as a conserved regulator for axonal

growth and regeneration in the CNS and PNS in mammals, flies and worms [15, 16]. Further, it

affects radial cell migration via microtubule-based events [17] and neural tube closure and

cortical development in the brain of mouse embryos [18, 19]. Yet, a direct assesment of the role

of JNK signaling in the morphogenesis of the CNS has not been performed.

Here we describe the consequences of deregulating the activity of the JNK cascade on the

architecture of the VNC and the motor coordination of the *Drosophila* embryo. We first

employed a correlative analysis and uncovered an iterated robust pattern of structural elements

(3D nodes or cross-pieces). Then we found that the JNK pathway - by directing axon integrity,

axonal fasciculation and dendrites formation and non-autonomously modulating Fasciclin 2 (Fas

2) expression - controls the robustness of the VNC architectural organization and its

condensation. The attainment of a correct geometry is the result of a delicate balance of JNK

signaling activity in specific pioneer neurons. We further found that the precise levels of JNK

activity necessary for correct VNC condensation are instrumental in the integration of embryonic

motor circuitries. Altering JNK activity levels resulted in a failure in the coordination of muscle

contractions and the inability to build the harmonic peristalsis essential for embryo hatching.

Altogether, our data elucidates some of the essential molecular and cellular mechanisms

underpinning robustness in CNS architecture and highlights the links between its architectural

organization and functional optimization.

**RESULTS**

**Modular Architecture of the Ventral Nerve Cord (VNC)**

The *Drosophila* embryonic VNC is a complex structure with a stereotypical 3D organization and

a defined distribution of axonal tracts and neuronal cell bodies. Yet, the elements sustaining its

architectural organization are unknown. To shed light on their identity we analyzed the spatial

correlations of Fas 2 patterns (**Figures 1A** and **1B**, see also **Movie S1**). Fas 2 is a neural cell

adhesion molecule that highlights a subset of the longitudinal axonal connectives, and which

loss-of-function leads to complete or partial defasciculation [20]. We uncovered an iterated

domain per segment along the A/P axis with a robust 3D geometrical organization (3D node) at

an average internodal distance of 28 μm (**Figure 1C**). These nodes positioned slightly anterior to

the anterior commissure (see **Figure S1**). At the nodes, the longitudinal connectives organized in

three compactly fasciculated units at precise dorso-ventral/medio-lateral positions (**Figure 1D**).

Outside the nodes (**Figure 1E**), the fascicles showed an imprecise distribution with a maximal

randomness (minimal correlation) at a level in-between the anterior and the posterior

commissures. The correlation profiles of the Fas 2 connectives showed not just segmental

periodicity (see Experimental Procedures) but a strong structural bias within individual segments

(relative difference in the maximum and minimum spatial derivatives of aligned profiles)

(**Figure 1F**). The internodal noise increased steeply posteriorly and shallowly anteriorly

uncovering a segmental structural polarization (**Figure 1F**).

In summary, nodes define topographical domains where the axonal arrangements are sustained

from segment to segment in the 3D space. We speculate that nodes correspond to conserved

crosspieces holding the architectural organization of the VNC.

**VNC architecture is modulated by JNK signaling**

In *Drosophila*, the JNK activity is moderated by a negative feedback loop implemented by the

dual-specificity phosphatase Puckered (Puc). *puc* expression depends on Basket (Bsk), the

*Drosophila* homologue of JNK, activity, and Bsk function is inhibited by Puc [21]. Although *puc*

expression has been reported in the CNS [22], its function during CNS morphogenesis is

unknown. We found that in *puc*$^{E69}$ loss-of-function mutants, the Fas 2 longitudinal connectives

were highly defasciculated and that the intersegmental nerves were unevenly positioned. Axons

became irregularly recruited to more medial positions and the Fas 2 signal was frequently

relocated to the cell bodies (aCC and pCC neurons) (**Figure 1G** and **Movie S2**). Equivalent

defects were also observed for the unrelated *puc*$^{B48}$ allele (**Figure S2**) [21].

Although in *puc*, the Fas 2 fascicles became disorganized on the dorso-ventral/medio-lateral

plane (**Figure 1H**), their topographical organization in the A/P axis was robustly conserved and

segmental 3D nodes, although more spaced, were still present (**Figure 1I** and **S2**). This increased

internodal distance reflected a failure on VNC condensation (on average 18% in *puc*$^{E69}$ and 39%

in $puc^{B48}$) (see also direct measurements of the VNC length in **Figure S3**). Further, the A/P

structural polarity within segments was also abolished (**Figures 1F**). Puc activity thus appears

not to be participating in the establishment of the segmental body plan but into the structural

reinforcement of the VNC and its condensation. We further found that in the absence of the

negative feedback loop mediated by *puc*, the level of JNK phosphorylation significantly rose in

the VNC (**Figure S4**). This increase in JNK activity also correlated with an increment in

apoptosis above the VNC characteristic levels (**Figure S4**). In brief, a tight control of JNK

signaling appears to be essential for the acquisition of the final architectural organization of the

VNC. This control is exerted, at least in part, by Puc, which limits JNK activity and promotes

viability.

**JNK signaling is dynamically regulated in the VNC**

When and where is the JNK cascade active in the VNC? To answer this question, we studied the

dynamics of *puc* expression, an early readout of JNK activity [21]. A viable Gal4 line under the

control of the $puc^{E69}$ enhancer [23] identified a segmentally iterated subset of neurons at

embryonic stage 17 (**Figures 2A** and **2B**). This pattern originated at stage 13 in a few neurons at

the ventral midline (**Figure 2E** and **Movie S3**). No cross-reaction was observed with glial

markers (**Figures 2C** and **2D** and [24]). As development progressed, an increased number of

neurons within the VNC and at the peripheral nervous system (PNS) expressed *puc*. We

identified a minimum consensus of 19 *puc* positive neurons of varying intensity per

hemisegment (see diagrams in **Figure 2F** and **Movie S4**).

Dual staining of $puc^{E69}$ embryos with neuronal markers [Even-skipped (Eve) and Engrailed (En)]

identified most of the *puc*-expressing cells (**Figure S5**). *puc* labeled the aCC, pCC, RP2, U1-U3

(plus weak/variable expression in U4-U5) and 3 out of the 8/10 EL interneurons. Meanwhile,

NL1, NL2, and iVUM5 En+ neurons were *puc* positive. Morphological and positional criteria

further enabled the assignment of five other neurons, AD, ADV, dMP2 and 2 VUM En-

motoneurons (**Figures 2G** and **2H**). This pattern of *puc*-expressing neurons was largely

conserved in $puc^{B48}$ (**Figure S6**). Summarizing, *puc* expression was detected from stage 13 in

VUMs, pCC and possibly dMP2 neurons at the ventral midline. This expression pattern

subsequently expands to diverse inter and motoneurons projecting through the longitudinal

connectives and segmental and intersegmental nerves.

**The level of JNK activity in *puc*-positive neurons controls the VNC architectural organization**

The altered architecture of the VNC observed in *puc* mutants could be the result of an autonomous malfunction in the CNS, or just a structural consequence of the general disruption of embryo development. To clarify this possibility, we generated JNK loss- or gain-of-function conditions overexpressing Bsk$^{DN}$, a dominant negative form of Basket (Bsk), or Hep$^{CA}$, a constitutively active form of Hemipterous (Hep), the *Drosophila* homologue of the JNK kinase MKK7, in *puc*-expressing cells with different Gal4 lines.

The CQ2-Gal4 line drives expression of UAS-linked genes in the U1 to U5 motoneurons (**Figure 3A**). Neither the inhibition (**Figure 3B**), nor the hyperactivation of the pathway (**Figure 3C**) resulted in any major structural defects. The VNC architecture in both cases (e.g. nodes positions and spatial correlation) was similar to *wt* (**Figures 1A** to **1C**).

The RN2-Gal4 line is specifically expressed in the aCC, pCC and RP2 neurons (**Figure 3D**). Interference with JNK signaling in RN2 cells resulted in the disorganization of the Fas 2 connectives (**Figure 3E**). The spatial correlations became noisy and irregular, the internodal distances expanded and the condensation of the VNC failed. On the other hand, the hyperactivation of the pathway, mimicking reduced *puc* levels, also strongly affected Fas 2 connectives, which kept a juvenile appearance (**Figure 3F**). Although 3D nodes were observed, the internodal distance enlarged and the VNC lengthened.

The MzVum-Gal4 line is expressed in a few lateral cells and in the 3 VUM motoneurons (**Figure 3G**). Bsk$^{DN}$ expression in these cells just slightly affected Fas 2 connectives fasciculation (**Figure 3H**). On the contrary, Hep$^{CA}$ expression resulted in disorganization of Fas 2 connectives and on an increased noise in spatial correlations (**Figure 3I**). Internodal distances also became larger and the VNC does not condense.

Altogether these data suggest that the level of JNK activity must be carefully tuned in the early-specified aCC, pCC, and RP2 pioneers and VUM neurons to facilitate the robust architectural organization of the VNC.

**JNK signaling modulates Fasciclin 2 levels in the developing VNC**

The non-autonomous impact of JNK signaling from local sources on global VNC morphogenesis requires the propagation of the cellular responses to JNK. While direct instructive communication by growth factors or neurotransmitters can be an important determinant of structural changes [25], adhesion molecules also stand as key elements for large-scale processes [26].

We explored whether the global effects of *puc* on structural robustness could be associated to Fas 2 deregulation or subcellular mislocalization. Fas 2 has been implicated in fasciculation by promoting adherence between growing axons [27]. By quantifying Fas 2 integrated densities in *wt*, *puc^E69^*, and *puc^B48^* embryos (**Figures 4A** to **4C**) we found that in all conditions it was enriched at the 3D nodes, which was consistent with their high axonal density. Fas 2 expression was reduced by 50% in the absence of Puc (at high JNK activity) (**Figure 4D**). This reduction was systemic and could not be assigned to individual domains within the VNC. In agreement, the quantification of protein levels in stage 16 embryo extracts showed a global reduction of Fas 2 expression in mutant conditions (**Figure 4E**).

These analyses support a model in which the global architectural organization of the VNC is influenced by the proper deployment of Fas 2 in response to the controlled activity of the JNK pathway in, at least, the aCC, pCC, RP2 and VUMs neurons.

**JNK signaling influences the axodendritic organization of *puc*-expressing neurons**

To investigate the structural origins of the effects of JNK signaling on the overall architectural organization of the VNC we monitored the axonal integrity and dendritic arborization of *puc*-expressing neurons. We observed that the overexpression of Hep^CA^ in RN2 (aCC, pCC and RP2) cells led to the local disorganization of the tissue affecting axonal morphologies and dendrites number and organization.

The aCC is a motoneuron whose axon gets incorporated into the intersegmental nerve (ISN) and proceeds laterally to innervate the dorsal muscle DA1 [28]. Its dendritic arbor develops in a distinct region of the ISN myotopic dendritic domain and consists of 8-10 dendrites [29] of an average length of 4 μm (**Figure 5A**). The pCC is a medially-located interneuron whose axon extends ipsilaterally in an anterior direction [30]. Finally, the RP2 motoneuron axonal trajectory

proceeds through the posterior root of the ISN and forms endings on all dorsolateral and dorsal internal muscles [28, 31]. Accordingly, the RP2 develop its dendrites into the region of the neuropile that represent its targets in the myotopic map [32]. Between 7 and 9 dendrites of an average length of 4 μm branching out of the RP2 axon can be regularly found (**Figure 5A**). Upon overexpression of Hep$^{CA}$, the cell bodies of the aCC, pCC and RP2 neurons were frequently misplaced and came together towards the midline (compare **Figures 5A** and **5B**). The appearance of their axons was affected; they thicken and became slack, while their length increased (for pCC and the RP2) (**Figures 5C** and **5D**). Hep$^{CA}$ overexpression also affected the density and length of dendrites. Both for aCC and RP2, dendrites numbers were reduced by half (**Figure 5E**) and they became longer reaching up to 15 μm of length (**Figure 5F** and **Movie S5**).

Our data show that altering JNK activity levels led autonomously to structural and, potentially, transmission defects in motoneurons. These faults along the loss of the adhesive capabilities of the tissue, consequence of the reduction of Fas 2 levels, may be responsible in certain degree of the architectural failures of the VNC as a whole, including its lack of condensation.

**The VNC architectural organization is essential for embryonic motor coordination**

To determine the functional relevance of the robust architectural organization of the VNC, we analyzed the process of maturation of embryonic motor patterns [7, 8] (**Figure 6A** and **6B**) (see Experimental Procedures, **Figure S7** and **Movie S6**). The *Drosophila* larva moves by peristaltic crawling and even before hatching, embryos execute forward and backward waves of peristalsis. These waves require the coordinated, bilaterally symmetrical, contraction of the thoracic and abdominal muscles [7, 33]. We found that embryonic movements began with isolated unilateral brief muscle twitches, sporadic head pinching events and sequences resembling incomplete peristalsis (stage A). This stage was followed by a prolonged resting period (stage B) occasionally interrupted by uncoordinated single contractions involving one or few segments. As development proceeds multi-segmental events arose with incomplete and complete peristalsis (stage C). These events then became less and less frequent until they ceased completely into a short resting period (stage D). Last, violent head movements and multiple peristaltic waves took place (stage E) leading to the rupture of the viteline membrane and larva hatching (**Figure 6C** and **6D** and **Movie S7**). These stages of stereotyped length closely recapitulate previous reports (Crisp et al., 2008). In the absence of *puc* (*puc$^{E69}$*), the developmental progression of motor

patterns was interrupted and embryos remained indefinitely in an immature stage (A)

characterized by occasional unilateral twitches and incomplete peristalsis, (**Figure 6E** and **6F**

and **Movie S8**).

Muscle integrity and functionality were not affected in *puc* embryos. Yet, the systemic failure of

motor coordination could still be the indirect consequence of non-defined events resulting from

an overall increase of JNK activity in *puc* mutants. To evaluate this possibility we overexpressed

Hep$^{CA}$ in RN2 (**Figure 7A** and **7B**) or MzVum (**Figure 7C** and **7D**) cells. In both cases, an

excess of JNK activity in just the *puc*-expressing neurons resulted in an equivalent phenotype to

that observed in *puc$^{E69}$* mutants. Only occasional unilateral twitches and head pinches and

infrequent incomplete peristalsis were observed in either condition. Therefore, the motor defects

in *puc* appear to be a direct response to loss of Puc activity in *puc*-expressing neurons.

Would the loss of motile synchrony be related to the aberrant architectural organization of the

VNC or to a signaling defects consequence of altering JNK activity in *puc*-expressing neurons?

Some of these *puc+* cells are motoneurons and in the absence of a correct level of JNK activity,

their signaling capability might be impaired. To test this possibility, we studied the axonal

transport of synaptotagmin, a key component of synaptic vesicles in the intersegmental nerve

[aCC and RP2 (RN2-Gal4) (**Figure 7E**) and VUM motoneurons (MzVum-Gal4) (**Figure 7F**)]

(see Experimental Procedures and **Movie S9**). Both in RN2 and MzVum axons, synaptotagmin

(synaptic vesicles) moved at a speed of around 0,7 μm/s with a tracking lifetime of 50 s per

event. Particles densities (frequency of events) were larger for aCC and RP2 (0,3 N/μm) than for

VUMs (0,2 N/μm). No differences were observed between anterograde and retrograde transport.

Next, particle transport was analyzed upon co-expression of Hep$^{CA}$ in RN2 and MzVUM cells.

Synaptotagmin transport behavior was largely unaffected (**Figures 7G to 7L)** which indicate that

microtubules, motor systems and synaptic vesicles transport remained fully functional. As for the

wild type, we could not distinguish differences between anterograde and retrograde transport.

In conclusion, synaptotagmin and, potentially, neurotransmitters transport appeared not to be

disturbed by the deregulation of JNK activity. These results suggest that the main function of the

attenuation of the JNK pathway in the VNC by Puc is mainly structural ensuring its robust

architectural organization. The embryonic motor coordination would be a foreseeable

consequence.

## DISCUSSION

### The VNC attains a robust Architectural Organization

The complex nervous system of bilaterians is thought to have emerged from an ancient, diffuse nerve plexus [34] that evolved into a segmented structure polarized along the Anterior/Posterior (A/P) axis [35]. 'Homologous' neurons functioning in different body segments are serially repeated and conserved across species in other arthropods. We have found that this conservation at the cellular level is reflected in a unique architectural pattern. Structurally, the VNC of the *Drosophila* embryo is composed of segmentally iterated ganglia with well-structured commissures and well-defined segmental and intersegmental nerves. Fasciculated axons (those expressing Fas 2) become robustly organized in 3D at one anchoring position (nodes) per segment slightly anterior to the connecting anterior commissure (**Figure 1** and **S1**). These commissures contain bundles of primary neurites where many synapses are made. A likely reason for this robust organization is that it responds to conserved architectural needs in terms of mechanics and physical balance [34, 36]. We speculate that in architectural terms the connectives may work as structural beams and that nodes help to distribute evenly any load from compression, tension or shear arising as the VNC is built (or that it may suffer once functional). This backbone will support the accumulated cargo of neuronal cell bodies, assuring that the elements required to fulfill neural functions will be robustly established. Yet, the Fas 2 connectives may not be the only element holding the tensional network of the VNC as the global axonal landscape of the embryo includes many other elements not expressing Fas 2.

### The JNK signaling modulates the 3D structure of the Nervous System

During the morphogenesis of the nervous system, both neurons and glia grow, change shape and accommodate overtime to establish a mechanically equilibrated and precisely organized 3D structure. Doing so, they generate - and also become the target of - forces and tensional stresses. It has been suggested that the compaction and clustering of cells in neural modules could be the result of the spatio-temporal regulation of mechanical tension in and along axons [37]. Indeed, tension forces contribute to neuronal morphology and connectivity [38], neurite connections

stabilization [37] and remodeling of neuronal processes and functional wiring [39]. Yet, the

information on how the tensional and topographical landscape of the nervous system is

implemented is scarce. A candidate to modulate neural structural organization and function and,

eventually, tension within the CNS is the stress activated JNK cascade [40].

The robust structural organization and the condensation of the VNC appear to be associated

events and they are both regulated by JNK activity. In *puc* mutants, Fas 2 bundles defasciculate,

the overall shape of the neuropile is affected, and the VNC fails to condense (**Figures 1** and **S3**).

3D nodes are still present at the right position within segments but the internodal distances are

enlarged. Remarkably, the comparison of spatial correlations profiles indicate that in each

segment the shrinkage of the domain immediately posterior to the nodes (roughly covering the

anterior commisure) is key for VNC condensation. If these iterated narrowings are temporally or

spatially (segmentally) regulated or if they are the only ones or just major participants in the

condensation process remains to be studied. In any case the regulated attenuation of the JNK

signaling cascade is essential.

In *Drosophila*, the pioneer neurons are specified and singled out at stage 9 and initiate

axonogenesis around stage 11 [30]. JNK activity, in parallel, builds up in the CNS and is

detected (*puc* expression) in the midline VUMs, and the pCC and dMP2 early pioneers (**Figure**

**2**). This temporal correlation argues that the major aberrations in fasciculation observed in *puc*

embryos may result from pioneering defects. Potentially, inaccurate axonogenesis of pioneers

could impact non-autonomously in the fasciculation of axons from late neurons that encounter an

altered guidance landscape. Indeed, we observed that autonomously increasing JNK activity in a

few early-specified neurons or in midline pioneers (RN2 and MzVum positive neurons) affects

their axons lengths and paths and their dendritic arborizations (**Figure 5**) preceeding widely

spread fasciculation defects. The failure of the VNC condensation may have the same origin.

VNC defects are remarkably similar upon reduction or increase in JNK function, indicating that

precision in the control of its contribution is essential. The negative feedback loop mediated by

*puc* would thus be a main factor in this tight spatio-temporal regulation.

How JNK activity levels would impinge in the architectural organization of the VNC? Fas 2 is a

neural cell adhesion molecule that highlights a subset of the longitudinal axonal connectives, and

which loss-of-function leads to complete or partial defasciculation [20]. The regular segmental

arrangement of Fas 2 nodes suggests they are important elements for the global stabilization of

the VNC structure. Defective node formation or alterations in neurons adhesion properties (e.g.

by Fas 2 downregulation) would weaken the structural scaffold. It has been shown that tension,

alongside cell adhesion and neurotransmitter-dependent signaling is necessary for axonal growth

and transport, it also promotes neurites maturation and supports dendrites development in

different models [41]. The disassembly of Fas 2 connectives and architectural rearrangements in

*puc* mutants are associated with a strong reduction of Fas 2 levels. Remarkably, The Fas 2

reduction is not a consequence of its relocation (e.g. node to internode) but instead to its systemic

downregulation (**Figure 4**). Interestingly, an equivalent link between JNK signaling and Fas 2

has been observed in the pruning of mushroom body gamma neurons axons. Bsk promotes

pruning by reducing the membrane levels of Fas 2 [42].

Altogether, our data suggest that the activation of *puc* expression by JNK in early-specified

pioneer neurons (aCC, pCC, RP2, dMP2 and VUMs) results in the downregulation of the

pathway, and an overall increase in Fas 2 levels. Fas 2 would be essential for proper structural

development [20]. A deficit in Fas 2 would affect the fasciculation of the axons of primary

neurons, leading to secondary bundling defects resulting in a general reduction in spatial

correlations across the whole axonal network of the neuropile. Failure to properly fasciculate, the

misrouting of axonal paths and alterations in dendritic arborizations would eventually affect both

the architectural organization and the tensional balance of the VNC, ultimately impeding its

condensation (**Figure 8**).

**Coordination of motor activities by JNK signaling**

In the *Drosophila* embryo, motor networks become active very early and isolated twitches and

unilateral waves of contraction occur prior to the generation of propagated action potentials by

neurons [43]. The early muscle contractions are thus myogenic in origin [7]. The elaborated

rhythmic coordination of motor function and its optimization are, however, the consequence of a

gradual maturation dependent on a functional Central Pattern Generator (CPG) [44]. In other

words, the invariant pattern of muscle contraction that constitutes a full peristaltic wave depends,

besides specific motoneuron activities, on the development of excitatory and inhibitory

interactions accross segments.

Does the geometrical and architectural organization of the VNC have any functional

significance? Which are the consequences of acquiring its precise shape and size? Our data
strongly advocate for a close relationship between the VNC axial condensation and the
development of a functional CPG and the acquisition of coordinated motor functions. Both, in
different *puc* alleles or after overexpressing Hep$^{CA}$ in the aCC, pCC, RP2 and VUM neurons,
when the robustness of the VNC organization is compromised, the embryo's motor coordination
becomes interrupted. Mutant embryos shake their muscles at low pace continuously and stay
alive for up to 24 h but do not hatch. Their muscles motility patterns can be best described as
uncoordinated individual segments contractions, where left and right hemisegments are often out
of phase and peristaltic waves are absent. These animals do not proceed beyond the early stage
of spontaneous twitches (A) observed in wild type embryos. While this outcome could be
explained as an indirect result of signaling defects associated to odd dendritic arborizations [45]
(**Figure 5**), we postulate that they are most probably linked to the incomplete compaction of the
VNC (**Figures 1** and **3**).

It is well established that correlated spontaneous activities [46] repeatedly and persistently
stimulating postsynaptic cells will result in long-term synapse strengthening (long-term
potentiation [LTP]) [47]. These spontaneous activity patterns are known to propagate, ensuring
that the connections between neighboring cells are strengthened, whereas those from more
distant ones are lost [48]. Thus, functional interconnectedness will only be reached by bringing
cells in close proximity leading to wiring optimization [12]. We hypothesize that these principles
will be followed during the condensation of the VNC ensuring the coordination of motor
activities. Several arguments support this premise. The muscle activity phenotype observed for
*puc* closely resembles that of *glial cells missing* (*gcm*). In *gcm*, neurons are unaffected but most
classes of glial cells are lost and VNC condensation does not take place. Peristaltic waves are
absent and uncoordinated movements are continuous [33]. Further, the activation or ablation of
the Even-skipped+ EL interneurons (Eve/Evx) disrupt the bilaterally symmetrical muscle
contractions in the larvae [49]. These ELs make direct presynaptic contacts to the contralateral
dorsal-projecting motoneurons RP2, U1, and U2. Amongst these, we found the RP2 to be key in
directing the structural organization of the VNC (**Figure 3**) and for motor coordination (**Figure
6**). The ELs and the dorsal projecting motoneurons they innervate are nearly all *puc* expressing
neurons (**Figure 2**). It will be of interest in the future the analysis of the EL sensorimotor circuit
and its signaling capabilities upon perturbation of JNK activity.

The principle of wiring optimization also likely applies to postembryonic development. As a

*Drosophila* larva develops, its body wall surface area grows by two orders of magnitude [50]. To

accommodate such physiological changes, mechanosensory neurons must extend their dendrites

to maintain receptive fields [51], while larval motor neurons add more synapses to maintain

functional responses in much larger muscles [50, 52]. Interestingly, during scaling, the likelihood

of synaptic contact between nearby neurons is stereotyped and cell type-specific [53]. In this

scenario, a post-synaptic neuron can only connect to spatially nearby pre-synaptic sites. This

local propensity to form stable synapses with a nearby cell type is preserved across development.

The application of correlative geometrical and mechanical analyses in combination with cell

type-specific genetic interference will allow wiring optimization to be tested across different

circuits and to better elucidate the detailed mechanisms underlying the CNS structural

development.

**EXPERIMENTAL PROCEDURES**

**Drosophila strains**

The following stocks were used:

w; *puc$^{E69}$LacZ/TM3, twi-GFP* [21]

*w; puc$^{B48}$LacZ/ TM3, twi-GFP* [21]

w; UAS-GFP; *puc$^{E69}$I–Gal4/TM6B* [23]

w; UAS-GFP; *puc$^{B48}$–Gal4/TM6B* [23]

*w[1118], P{w[+mC]=UAS-bsk.DN}2* (BDSC #6409)

*w[*]; P{w[+mC]=UAS-Hep.Act}2* (BDSC #9306)

*P{GawB}elav[C155]* (BDSC #6923)

*w[*]; P{w[+mC]=eve-GAL4.RN2}T, P{w[+mC]=UAS-mCD8::GFP.L}; P{w[+mC]=eve-*

*GAL4.RN2}G, P{w[+mC]=UAS-mCD8::GFP.L} (*Dr. Irene Miguel-Aliaga)

*y[1] w[*];P{w[+mC]=CQ2-GAL4}O, P{w[+mC]=UAS-mCD8::GFP.L}LL6* (BDSC #7465)

*w[\*] P{w[+mW.hs]=GawB}MzVUM; P{y[+t7.7] w[+mC] = 10xUAS-IVS-mCD8::GFP}attP40*

*(*Dr. Irene Miguel-Aliaga)

*w[\*]; G203:UAS-TNT; ZCL2144* (Dr. Matthias Landgraf)

*w[\*]; ; ZCL2144* (Dr. Matthias Landgraf)

*w\*; P{UAS-syt.eGFP}2* (BDSC #6925)

In all cases, unless otherwise stated, embryos of the *w1118* strain served as controls.

**Immunohistochemistry**

Immunostaining of flat-prepped stage 16 *Drosophila* embryos was performed using the

following primary antibodies: mouse anti-Fas 2 (1:100, clone 1D4, DHSB), mouse anti-axons

CNS (1:100, BP 102, DHSB), mouse anti-Futsch (1:1000, 22c10, DHSB), rabbit anti-phospho-

JNK (Thr$^{183}$ / Tyr$^{185}$) (1:100, Cell Signaling #9251), rabbit anti-Dcp-1 (Asp216) (1:100, Cell

Signaling #9578), rabbit anti-GFP tag polyclonal (1:600, Thermo Fisher Scientific), rat anti-Elav

(1:1000, clone 7E8A10, DHSB), mouse anti-Repo (1:100, clone 8D12 DHSB), mouse anti-Even-

skipped (1:100, clone 3C10, DHSB), and mouse anti-Engrailed (1:100, clone 4D9, DHSB).

The secondary antibodies used for detection were: Goat anti-Rabbit IgG (H+L), Alexa Fluor 488

conjugate (A-11008), Goat anti-Rabbit IgG (H+L) Alexa Fluor 555 conjugate (A-21428), Goat

anti-Mouse IgG (H+L) Alexa Fluor 488 conjugate (A-11001), Goat anti-Mouse IgG (H+L)

Alexa Fluor 555 conjugate (A-21422) and Goat anti-Rat IgG (H+L) Alexa Fluor 555 conjugate

(A-21434). All secondary antibodies were used in a dilution of 1:600 and were from Invitrogen.

**Sample preparations for immunodetection and image acquisition**

*Drosophila* embryo dissections for generating flat preparations were performed according to

[28]. Briefly, flies maintained in apple juice-agar plates at 25ºC were synchronized by repetitive

changes of the juice-agar plate, with a time interval of 1.5 hours. All embryos laid within this

time window were aged for approximately 22 hours at 18ºC, or until reaching mid-stage 16 (3-

part gut stage). At this point embryos were dechorionated with bleach for 1 min, poured into a

mesh and rinsed extensively with water. For dissection, embryos were transferred with forceps

on the surface of a small piece of double sided tape, adhered on one of the sides of a poly-L-

Lysine coated coverslip. After orienting the embryos dorsal side up and posterior end towards

the center of the coverslip, the coverslip was flooded with saline (0.075 M Phosphate Buffer, pH

7.2). Using a pulled glass needle the embryos were manually de-vitellinized and dragged to the

center of the coverslip, where they were attached to the coated glass with their ventral side down.

An incision on the dorsal side of the embryo was performed using the glass needle from the

anterior to the posterior end of the embryo. The gut was removed by mouth suction and blowing

stream of saline was used to flatten their lateral epidermis. Tissue fixation was done with 3.7 %

formaldehyde in saline for 10 minutes at room temperature. After this point standard immuno-

staining procedures were followed.

In the single case (**Figure S3**) where whole mount embryos were immunostained, embryo

fixation and de-vitalinization was done according to [54]. Both whole mount and flat-prepped

immunostained embryos were mounted in Vectashield anti-fading medium (Vector Laboratories,

USA).

Image acquisition was performed on a Zeiss LSM 700 inverted confocal microscope, using a 63

x oil objective lens (NA 1.4).  Z-stacks spanning the whole VNC thickness were acquired

applying optimal, with respect to pinhole diameter, sectioning with a step size that varied

between $0.2 - 0.6$ μm. Image processing was performed with Fiji [55].

**Live imaging**

Dechorionated stage 14 embryos were glued ventral side down on a MatTek glass bottom dish

and they were covered with S100 Halocarbon Oil (Merck) to avoid desiccation. Image

acquisition was performed on a Zeiss LSM 700 inverted confocal microscope, using a 25 x oil

objective lens (NA 0.8).  Z-stacks spanning the whole VNC thickness and with a 2 μm step size

were acquired every 6 minutes for a total of 8 hours. Processing of time-lapse data was done with

Fiji [55].

**Image analysis**

Image analysis and quantification of fluorescence intensity were performed using Fiji [55]. In

immunostainings where quantification of the fluorescent intensity was required, extra care was

taken so that the same antibody batch was used for staining embryos of different genotypes,

while identical confocal acquisition settings were applied.

For calculating the average integrated density of Fas 2 signal in an image, regions of interest
(ROIs) of fixed size were measured at the nodes and internodes positions. Average Integrated
Density values obtained by measuring ROIs in several animals (n stated in Figure legends) were
then pooled for calculating the mean value as well as standard deviations. For those cases where
the quantification of integrated density was performed on a segment basis, the measured values
represent the raw integrated density, since the area measured slightly varies in between animals.

**Statistical analysis**

All statistical analyses were performed using GraphPad Software (GraphPad Software Inc., La
Jolla, CA, USA). In all cases one sample t test was performed and probability values $p<0.05$
were considered as significant.

**Correlations Data analysis**

Images were scaled to be isotropic in all axes. Deconvolution was found to not significantly
improve the image correlation. Viewed along the antero-posterior axis, images were cropped to
include only the VNC, and then the VNC was differentiated into 50 bins, corresponding to 1,65
μm length each. Within each bin, a maximum intensity projection was performed along the
included (typically eleven) planes. The Matlab function *imregister* was then used to perform the
image registration for both inter- and intra- embryo correlations. The optimizer for *imregister*
was defined with multimodal metric, tolerance of $10^{-6}$ and 500 iterations. Before image
registration, the center of mass of the two images was aligned so to maximize image overlap.
The code was run on a PC with 128 Gb memory and in the most demanding case (full embryo
correlation: $50 \times 50 = 2500$ correlations) took around one hour per sample.

To create the image correlation traces, the regions of high correlation (e.g. as identified by
arrows in **Figure 1**) were manually identified and then plotted considering the mean correlation
along ± 1 row in the correlation matrices. The diagonal values were ignored. Traces were offset
along AP axis to ensure maximum overlap of peaks in the region 25-57 μm to aid comparison
between different datasets. In all figures, individual profiles are plotted in gray, with the mean
(black) and ± 1 standard deviation (red) shown. For each maximum in the spatial correlation
profile, and a subset of data was sliced that lay between the two minima at either side. Then, we

created *in silico* data from the real experimental data points - i.e. each *in silico* data point was

drawn from a distribution defined by the mean and standard deviation in the samples. We

determined the best linear fit to each *in silico* data set and repeated 500 times to find the average

gradient fit. This was a robust way to find the steepness as it accounts for experimental noise. To

compare between datasets, we defined the variable

r = (max(gradient) - abs(min(gradient))) / [(max(gradient)+abs(min(gradient)))/2]

*i.e.* the difference / mean.

**Muscle contraction analysis**

Stage 17 embryos carrying the G203, ZCL2144 or both GFP muscle markers were dechorionated

and mounted in halocarbon oil (3:1 mix of S700 and S27) following standard procedures. Muscle

contractions were monitored at the embryo's middle plane, using a 25 X oil immersion lens (N.A

0.8, Imm Korr DIC M27) in a Zeiss LSM780 confocal microscope. For each acquisition 1000

frames were obtained with a scanning speed of 200 msec per frame with a time interval between

acquisition cycles of 15 minutes. Muscle activity was monitored up to embryo's hatching or for a

total of 12 hours at room temperature. For the experiments where $Hep^{CA}$ over-expression in RN2

and MzVum neurons was required, both control and $Hep^{CA}$ overexpressing embryos were filmed

at $29^{o}C$ for equal time periods. Post-imaging, movies were processed with Fiji [55] and

Kymographs were generated using the Fiji plugin KymoResliceWide.

**Axonal Trafficking**

To examine axonal trafficking, stage 17 embryos expressing UAS-Syt-GFP were dechorionated

and mounted laterally in halocarbon oil (3:1 mix of S700 and S27) for live imaging. Synaptic

vesicle transport was monitored in the medial part of the Inter-Segmental Nerve (ISN) using a 40

X oil immersion lens (N.A 1.3, Oil DIC M27) in a Zeiss LSM780 confocal microscope. A

constant frame size of 512x100 pixels, with an XY zoom of 6 and a recording line spacing of 2

were routinely employed. Single plane movies were acquired with a scanning speed of 1.11 sec,

with 1 sec interval for a total of 60 cycles. All movies were processed with Fiji and Kymographs

were generated using the Fiji plugin KymoResliceWide.

The Mean Velocity and the Lifetime parameters for each particle were calculated from the

Kymographs as follows. The particles trajectories were traced and saved as independent ROIs.

The XY coordinates for each ROI were then extracted using the getSelectionCoordinates

function in Fiji of the custom made "Macro coordinates.ijm" shown below:

run("Fit Spline", "straighten");

getSelectionCoordinates(x, y);

x2=0; y2=0; distance=0;

for (i=0; i<x.length; i++) {

if (i>0) {

dx = x[i] - x[i-1];

dy = y[i] - y[i-1];

distance = sqrt(dx*dx+dy*dy);

}

print(i, x[i], y[i], distance);

}

These XY coordinates represent the particles displacement in both space (X values) and time (Y

values), thus permitting the calculation of Mean Velocity and Lifetime.

**Analysis of Axonal and dendrite morphology by Segmentation**

To describe the axonal and dendrite morphology of the RN2-Gal4 positive neurons, we analyzed

images of flat-prepped embryos immunostained against GFP and Fasciclin II using the Simple

Neurite Tracer Fiji plugin [56]. Image segmentation by this tool allows a semi-automated tracing

of axons and dendrites in 3D, providing measurements of total axonal and dendrite length, as

well as of dendrite number directly from skeletonized images.

**Western Blotting**

Dechorionated stage 16 embryos of the $puc^{E69}LacZ$ / *TM3, twi-GFP* and $puc^{B48}LacZ$ / *TM3, twi-GFP* lines were used for generating whole embryo extracts for western blot analysis. In detail, homozygous embryos were separated from their balanced siblings under a fluorescent dissecting scope based on the *twi-GFP* expression. Approximately 100 embryos of each type were fully homogenized in 100 μL of iced cold RIPA buffer, supplemented with Protease Inhibitors (Protease Inhibitor Cocktail Tablets, Roche Life Sciences). Subsequently, the extracts were centrifuged at 3500 g for 15 minutes at 4ºC and the supernatants were collected. Total protein concentration in the lysates was determined by Bradford assay (Bio-Rad) and 30 μg of sample were analyzed in a 10 % SDS-PAGE. Proteins were blotted on Protran nitrocellulose membrane (Whatman) while the antibodies employed were: anti-Fas 2 (1:1000, clone 1D4, DSHB) and Pan-actin (1:1000, Cell Signaling). The ECL Plus chemiluminescent substrate (Pierce, Thermo Fisher Scientific) was used for antibody detection. Western Blot signal quantification was performed with Fiji by measuring the average density of each band.

**REFERENCES**

1.      Leise EM. Modular construction of nervous systems: a basic principle of design for invertebrates and vertebrates. Brain research reviews. 1990;15(1):1-23.

2.      Chen BL, Hall DH, Chklovskii DB. Wiring optimization can relate neuronal structure and function. Proceedings of the National Academy of Sciences of the United States of America. 2006;103(12):4723-8.

3.      Redies C, Puelles L. Modularity in vertebrate brain development and evolution. BioEssays. 2001;23(12):1100-11.

4.      Sporns O, Betzel RF. Modular Brain Networks. Annu Rev Psychol. 2016;67:613-40.

5.      Landgraf M, Sanchez-Soriano N, Technau GM, Urban J, Prokop A. Charting the Drosophila neuropile: a strategy for the standardised characterisation of genetically amenable neurites. Developmental biology. 2003;260(1):207-25.

6.     Technau GM. Brain development in Drosophila melanogaster: Springer
Science+Business Media, LLC  - Landes Bioscience; 2008.

7.     Crisp S, Evers JF, Fiala A, Bate M. The development of motor coordination in
Drosophila embryos. Development. 2008;135(22):3707-17.

8.     Crisp SJ, Evers JF, Bate M. Endogenous patterns of activity are required for the
maturation of a motor network. The Journal of neuroscience. 2011;31(29):10445-50.

9.     Suster ML, Bate M. Embryonic assembly of a central pattern generator without
sensory input. Nature. 2002;416(6877):174-8.

10.    Huntenburg JM, Bazin PL, Margulies DS. Large-Scale Gradients in Human Cortical
Organization. Trends Cogn Sci. 2018;22(1):21-31.

11.    Huntenburg JM. A core organizing axis of the human cerebral cortex: Freie
Universität Berlin; 2018.

12.    Cajal SRy. Textura del Sistema Nervioso del Hombre y de los Vertebrados. Madrid,
Spain: Nicolas Moya; 1899.

13.    Tessier-Lavigne M, Goodman CS. The molecular biology of axon guidance. Science.
1996;274(5290):1123-33.

14.    Chklovskii DB, Koulakov AA. Maps in the brain: what can we learn from them?
Annual review of neuroscience. 2004;27:369-92.

15.    Srahna M, Leyssen M, Choi CM, Fradkin LG, Noordermeer JN, Hassan BA. A signaling
network for patterning of neuronal connectivity in the Drosophila brain. PLoS biology.
2006;4(11):e348.

16.    Koch M, Nicolas M, Zschaetzsch M, de Geest N, Claeys A, Yan J, et al. A Fat-Facets-
Dscam1-JNK Pathway Enhances Axonal Growth in Development and after Injury. Front Cell
Neurosci. 2017;11:416.

17.    Hirai S, Banba Y, Satake T, Ohno S. Axon formation in neocortical neurons depends
on stage-specific regulation of microtubule stability by the dual leucine zipper kinase-c-Jun
N-terminal kinase pathway. The Journal of neuroscience. 2011;31(17):6468-80.

18.    Kuan CY, Yang DD, Samanta Roy DR, Davis RJ, Rakic P, Flavell RA. The Jnk1 and Jnk2
protein kinases are required for regional specific apoptosis during early brain
development. Neuron. 1999;22(4):667-76.

19.    Waetzig V, Herdegen T. Neurodegenerative and physiological actions of c-Jun N-
terminal kinases in the mammalian brain. Neuroscience letters. 2004;361(1-3):64-7.

20.    Lin DM, Fetter RD, Kopczynski C, Grenningloh G, Goodman CS. Genetic analysis of
Fasciclin II in Drosophila: defasciculation, refasciculation, and altered fasciculation. Neuron.
1994;13(5):1055-69.

21.    Martin-Blanco E, Gampel A, Ring J, Virdee K, Kirov N, Tolkovsky AM, et al. puckered
encodes a phosphatase that mediates a feedback loop regulating JNK activity during dorsal
closure in Drosophila. Genes & development. 1998;12(4):557-70.

22.    Martin-Blanco E. Regulatory control of signal transduction during morphogenesis in
Drosophila. The International journal of developmental biology. 1998;42(3):363-8.

23.    Pastor-Pareja JC, Grawe F, Martin-Blanco E, Garcia-Bellido A. Invasive cell behavior
during Drosophila imaginal disc eversion is mediated by the JNK signaling cascade. Dev
Cell. 2004;7(3):387-99.

24.    Shklover J, Mishnaevski K, Levy-Adam F, Kurant E. JNK pathway activation is able to
synchronize neuronal death and glial phagocytosis in Drosophila. Cell death & disease.
2015;6:e1649.

25.    Foty RA, Steinberg MS. The differential adhesion hypothesis: a direct evaluation.
Developmental biology. 2005;278(1):255-63.

26.    Amack JD, Manning ML. Knowing the boundaries: extending the differential
adhesion hypothesis in embryonic cell sorting. Science. 2012;338(6104):212-5.

27.    Grenningloh G, Rehm EJ, Goodman CS. Genetic analysis of growth cone guidance in
Drosophila: fasciclin II functions as a neuronal recognition molecule. Cell. 1991;67(1):45-
57.

28.    Landgraf M, Bossing T, Technau GM, Bate M. The origin, location, and projections of
the embryonic abdominal motorneurons of Drosophila. The Journal of neuroscience.
1997;17(24):9642-55.

29.    Furrer MP, Vasenkova I, Kamiyama D, Rosado Y, Chiba A. Slit and Robo control the
development of dendrites in Drosophila CNS. Development. 2007;134(21):3795-804.

30.    Jacobs JR, Goodman CS. Embryonic development of axon pathways in the Drosophila
CNS. II. Behavior of pioneer growth cones. The Journal of neuroscience. 1989;9(7):2412-22.

31.    Sink H, Whitington PM. Pathfinding in the central nervous system and periphery by
identified embryonic Drosophila motor axons. Development. 1991;112(1):307-16.

32.    Landgraf M, Jeffrey V, Fujioka M, Jaynes JB, Bate M. Embryonic origins of a motor
system: motor dendrites form a myotopic map in Drosophila. PLoS biology. 2003;1(2):E41.

33.    Pereanu W, Spindler S, Im E, Buu N, Hartenstein V. The emergence of patterned
movement during late embryogenesis of Drosophila. Dev Neurobiol. 2007;67(12):1669-85.

34.    Arendt D, Tosches MA, Marlow H. From nerve net to nerve ring, nerve cord and
brain--evolution of the nervous system. Nature reviews Neuroscience. 2016;17(1):61-72.

35.    Qing Yu ZL, Chuanzi OuYang, Lingjun Li. Biologically Active Peptides in
Invertebrates: Discovery and Functional Studies. Colloqium Series on neuropeptides:
Morgan & Claypool Publishers; 2015. p. 182.

36.    Matheson T. Invertebrate Nervous Systems. Encyclopedia of Life Sciences. 2002:1-6.

37.    Siechen S, Yang S, Chiba A, Saif T. Mechanical tension contributes to clustering of
neurotransmitter vesicles at presynaptic terminals. Proceedings of the National Academy
of Sciences of the United States of America. 2009;106(31):12611-6.

38.    Franze K. The mechanical control of nervous system development. Development.
2013;140(15):3069-77.

39.    Van Essen DC. A tension-based theory of morphogenesis and compact wiring in the
central nervous system. Nature. 1997;385(6614):313-8.

40.    Hsieh MH, Nguyen HT. Molecular mechanism of apoptosis induced by mechanical
forces. International review of cytology. 2005;245:45-90.

41.    Heiman MG, Shaham S. Twigs into branches: how a filopodium becomes a dendrite.
Curr Opin Neurobiol. 2010;20(1):86-91.

42. Bornstein B, Zahavi EE, Gelley S, Zoosman M, Yaniv SP, Fuchs O, et al. Developmental
Axon Pruning Requires Destabilization of Cell Adhesion by JNK Signaling. Neuron.
2015;88(5):926-40.

43. Baines RA, Bate M. Electrophysiological development of central neurons in the
Drosophila embryo. The Journal of neuroscience. 1998;18(12):4673-83.

44. Marder E, Bucher D. Central pattern generators and the control of rhythmic
movements. Curr Biol. 2001;11(23):R986-96.

45. Katz LC, Shatz CJ. Synaptic activity and the construction of cortical circuits. Science.
1996;274(5290):1133-8.

46. Kirkby LA, Sack GS, Firl A, Feller MB. A role for correlated spontaneous activity in
the assembly of neural circuits. Neuron. 2013;80(5):1129-44.

47. Hebb D. The Organization of Behavior: A Neuropsychological Theory. John Wiley
and Sons Inc; New York; Chapman and Hall Limited; London; 1949.

48. Eglen SJ, Demas J, Wong RO. Mapping by waves. Patterned spontaneous activity
regulates retinotopic map refinement. Neuron. 2003;40(6):1053-5.

49. Heckscher ES, Zarin AA, Faumont S, Clark MQ, Manning L, Fushiki A, et al. Even-
Skipped(+) Interneurons Are Core Components of a Sensorimotor Circuit that Maintains
Left-Right Symmetric Muscle Contraction Amplitude. Neuron. 2015;88(2):314-29.

50. Keshishian H, Chiba A. Neuromuscular development in Drosophila: insights from
single neurons and single genes. Trends Neurosci. 1993;16(7):278-83.

51. Grueber WB, Jan LY, Jan YN. Tiling of the Drosophila epidermis by multidendritic
sensory neurons. Development. 2002;129(12):2867-78.

52. Rasse TM, Fouquet W, Schmid A, Kittel RJ, Mertel S, Sigrist CB, et al. Glutamate
receptor dynamics organizing synapse formation in vivo. Nat Neurosci. 2005;8(7):898-905.

53. Gerhard S, Andrade I, Fetter RD, Cardona A, Schneider-Mizell CM. Conserved neural
circuit structure across Drosophila larval development revealed by comparative
connectomics. Elife. 2017;6.

54. Patel NH. Imaging neuronal subsets and other cell types in whole-mount Drosophila
embryos and larvae using antibody probes. Methods in cell biology. 1994;44:445-87.

55.    Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji:
an open-source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82.

56.    Longair MH, Baker DA, Armstrong JD. Simple Neurite Tracer: open source software
for reconstruction, visualization and analysis of neuronal processes. Bioinformatics.
2011;27(17):2453-4.

**ACKNOWLEDGEMENTS**

We are extremely grateful to all colleagues that have provided us materials, training and
guidance throughout this work as well as for commenting on the manuscript. Amongst them we
like to specially thank Claude Desplan, Matthias Landgraf, Chris Doe, Andreas Prokop, Natalia
Sanchez-Soriano, Marta Zlatic, Albert Cardona and Irene Miguel-Aliaga. We also have to thank
our colleagues at the Parc Cientific de Barcelona, the Mechanobiology Institute of Singapore and
the Fleming Institute of Athens for their constant encouragement and support.

 **FIGURE LEGENDS**

# Figure 1

**Figure 1. Structural Architecture of the Ventral Nerve Cord (VNC) of the *Drosophila* embryo**

**A)** Maximum projection of the VNC of a *Drosophila* 16-stage *wt* embryo stained with Fas 2. See **Movie S1**. Scale Bar is 10 μm.

**B)** Self cross-correlation matrix of the Z sections of the image in **A** along the A/P axis. The color-coded representation, in all figures from now on, shows the correlation level for each possible comparison at each position along the A/P axis. In this way it was possible to identify an oscillatory pattern of hot spots (3D nodes - purple double headed arrows) of high correlation. One conspicuous node of robust correlation per segment is detected (white arrowheads).

**C)** Image cross-correlation score along the A/P axis for **B**. The cross-correlation profile shows that segmental Fas 2 nodes fell 28 μm apart in *wt* animals and that the internodal maximum noise/minimum correlation maps 9 μm posterior to the node.

**D)** Z sections at two different maximal (green and magenta) correlation positions (from **A**) showing coalescence of Fas 2 connectives at the nodes.

**E)** Z sections at two different minimal (green and magenta) correlation positions (from **A**) showing dispersion and stochastic distribution associated to internodal noise.

**F)** Box plots of the differences vs. the mean for the variable r = [max (grad) - abs (min (grad))] / [(max (grad) + abs (min (grad))) / 2] between datasets for *wt, puc$^{E69}$* and *puc$^{B48}$* individuals. The median from *wt* was clearly less than zero with a 95% confidence interval. For the mutants this was lost, with the values not distinguishable from zero. The bias in *wt* values indicates that the decay in robustness from nodes in the posterior direction is more pronounced than its recovery towards subsequent nodes. This behavior is abolished in *puc$^{E69}$* and *puc$^{B48}$*.

**G)** Equivalent image to **A** for a *puc$^{E69}$* embryo. Note the defasciculation and collapse of the longitudinal connectives, specially the most lateral ones. See **Movie S2**. Scale Bar is 10 μm.

**H)** Self cross-correlation matrix of the Z sections of the image in **D**. In *puc$^{E69}$* mutants, while correlation matrices are noisier, a single node per segment can still be observed (white arrowheads).

**I)** Image cross-correlation score along the A/P axis for **E**. Segmental Fas 2 nodes spread 33 μm away in *puc$^{E69}$* and the internodal minima gets diffuse.

# Figure 2

**Figure 2. JNK signaling is dynamically regulated during development**

**A** to **D)** Dorsal and ventral sections highlighting the expression of *puc* (green) in the VNC of stage 17 UAS-GFP; *puc*$^{E69}$*I*-Gal4 embryo double-stained for the pan-neuronal marker Elav (**A** and **B** - red) or the glial marker Repo (**C** and **D** - red). *puc* is expressed in a subset of neurons in each segments but not in glia. Scale Bar is 10 μm.

**E** to **E''')** four different snapshots from **Movie S3** at the indicated time points from the UAS-GFP; *puc*$^{E69}$*I*-Gal4 viable line. *puc* expression can be observed from early stage 13 in a few neurons at midline. As development progressed, more and more neurons express *puc*. Scale Bar is 10 μm.

**F** to **F''')** Diagram displaying the cumulative expression of *puc* at the same time points as in **E** to **E'''**. A minimum consensus of 19 neurons per hemisegment was defined. At each time point the intensity (defined by visual inspection) was displayed as low (red), medium (blue) and high (green) (see also **Movie S4**).

**G)** Composition summarizing the expression of *puc* in the CNS [*puc* expressing cells in the VNC at stage 16 including those identified by coexpressing Even-skipped (magenta) or Engrailed (cyan) plus those putatively identified by morphological and positional criteria (yellow - AD, ADV, dMP2 and 2 VUM En-negative motor neurons)]. To facilitate understanding, a scaled Fas 2 expression image from an embryo of the same stage was overimposed to the diagram. Positions of the robustness nodes are highlighted as dark blue lines.

**H)** Diagram representing the whole set of *puc* expressing cells (compare to G) and their expected axonal contributions to connectives and exiting nerves (each individual cell carried a single color-code). This diagram was over-imposed over a grey mask covering the scaled width and shape of the VNC.

# Figure 3

**Figure 3. Targeted activation and inhibition of the JNK pathway in the VNC**

**A)** Cartoon representing the full set of *puc* expressing neurons per segment in the VNC highlighting the cells expressing CQ2-Gal4 (yellow).

**B** and **C)** From left to right, pattern of Fas 2 expression, self cross-correlation matrix and image cross-correlation profiles for VNCs of 16-stage embryos upon expression of UAS-Bsk$^{DN}$ **(B)** and UAS-Hep$^{CA}$ **(C)** transgenes under the control of the CQ2-Gal4 respectively inhibiting and hyperactivating JNK activity in the U motoneurons. None of these conditions result in noticeable defects. Scale Bar is 10 μm.

**D)** Cartoon highlighting the *puc* positive cells expressing RN2-Gal4 (turquoise).

**E** and **F)** As **B** and **C** but employing RN2-Gal4 for interfering in JNK activity in the aCC, pCC and RP2 cells. Affecting JNK activity levels in aCC, pCC and RP2 cells affect connectives defasciculation, nodes profiles and VNC condensation. Scale Bar is 10 μm.

**G)** Cartoon highlighting the *puc* positive neurons expressing MzVum-Gal4 (purple).

**H** and **I)** As **B** and **C** but employing MzVum-Gal4 for interfering in JNK activity in the VUM neurons. Affecting JNK activity levels in VUMs also affects defasciculation, profiles and condensation. Scale Bar is 10 μm.

# Figure 4

**Figure 4. Fasciclin 2 expression is reduced in *puc* mutants**

**A)** *wt*, **B)** *puc^{E69}* and **C)** *puc^{B48}* stage 16 embryos Fas 2 immunoreactivity. Maximum projection of ventral views across three VNC abdominal segments. Overlaid squares represent the fixed area ROIs used for measuring the Average Integrated Density of Fas 2 signal, either at the node (red) or at the internode (yellow) positions. Scale Bar is 10 μm.

**D)** Quantification of the Average Integrated Density of Fas 2 signal at node and internode positions as indicated. Statistically significant differences in Fas 2 levels ($p < 0.0001$) were detected between *wt* and *puc* embryos, both at the node and internode positions.

**E)** Immunodetection of Fas 2 in extracts from stage 16 *puc* embryos. Extracts of balanced siblings were used as controls. In control extracts, the antibody against Fas 2 detected a strong band at 95 kD (red arrow), the intensity of which was reduced by half in extracts from *puc* homozygous embryos. Detection of actin (42 kD) served as loading control for the quantification of the Fas 2 signal. The graph represents the averaged Fas 2 band intensities normalized with respect to actin.

# Figure 5

**Figure 5. Pioneer neurons axonal and dendritic landscape is altered upon JNK hyper-**

**activation**

**A** and **B**) Flat-prepped, stage 16 embryos expressing mCD8-GFP (**A**) and mCD8-GFP and

Hep$^{CA}$ (**B**) under the control of the RN2-Gal4 line immunostained for GFP. Traces of the aCC

(red) and RP2 (green) proximal dendrites are over-imposed. Yellow arrowheads point to

differences in RP2 axon morphology. Anterior is up. Scale bar is 10 μm.

**C**) Histograms showing the average axonal length of the aCC, RP2 and pCC neurons.

Statistically significant differences ($p<0.001$) in the axonal length of the RP2 and pCC neurons

were detected between control and Hep$^{CA}$ expressing neurons (n= 4 embryos). Control embryos

are represented in light green and Hep$^{CA}$ expressing embryos in dark green.

**D**) Histograms showing the average axonal wiggliness of the aCC, RP2 and pCC neurons. RP2

axonal wiggliness was found significantly increased after Hep$^{CA}$ expression ($p<0.001$, n= 4

embryos).

**E** and **F**) Graphs representing the average Dendrite Number (**E**) and average Dendrite Length (**F**)

of the aCC and RP2 neurons. In all cases significant differences were scored in the comparison

of control and Hep$^{CA}$ expressing embryos ($p<0.001$, n= 4 embryos).

# Figure 6

**Figure 6. Defective motor coordination results in the absence of complete peristalsis in *puc^E69* embryos**

**A)** Representative mid-plane image of a stage 17 embryo carrying two GFP protein traps expressed at muscle Z-lines (w; G203, ZCL2144) (see **Movie S7**). A selected segmented-line (highlighted in yellow) was employed to create kymographs spanning muscle units along the antero-posterior (A/P) axis (see below) at both sides of the embryo, right (R) and left (L). The evolution overtime of these kymographs was monitored live. Scale bar is 50 μm.

**B)** Intensity Profile plots generated along the segmented-line selection in **A**. Gray values as a function of distance highlight the distribution of the segmental muscle units along the axis. Muscle contraction is detected by the differences in the distance between consecutive peaks at sequential time points.

**C)** Kymograph displaying muscle units profiles from 17 to 25 hours AEL for the left side of the embryo in **A**. Anterior (A) and posterior (P). Motor coordination maturation occurs in 5 stages of approximated stereotyped length (color coded): A, Intense unilateral twitching; B, general inactivity with occasional uncoordinated contractions; C, initiation of backward and forward peristalsis; D, pre-hatching inactivity period and E, complete peristalsis and specialized hatching head movements.

**D)** Histograms showing the progression of the muscle coordination process in wild type embryos. Over-imposed colors follow the 5 stages of motor coordination maturation described in **C**. Muscle movements were distributed in four classes: Unilateral contractions [sum of the contractions occurring at the right (green) and the left (purple) side of the embryo]; Incomplete peristalsis [sum of the incomplete forward (blue) and incomplete backward (red) peristaltic attempts]; Complete peristalsis [sum of the full forward (blue) and backward (red) peristalsis]; and Head movements [sum of head turning (red) and mouth hook pinching events (blue)].

**E)** Kymograph displaying muscle units profiles from 18 to 23 hours AEL, when the embryo dies, for the left side of a *puc^E69* embryo (see **Movie S8**). Anterior (A) and posterior (P). Motor coordination fails and embryos do not progress beyond the unilateral contraction period (stage A).

**F)** Histograms equivalent to **D** showing the unsuccessful muscle coordination of *puc^E69* embryos.

# Figure 7

**Figure 7. JNK activity modulates VNC condensation and embryo motor coordination but does not affect synaptic vesicles transport**

**A** and **C**) Ventral view of stage 17 embryos expressing GFP under the control of the RN2-Gal4 (**A**) and MzVum-Gal4 (**C**) lines carrying the muscle Z-lines specific GFP protein trap ZCL2144. Muscles are pseudo-colored in red and neurons in green. Scale bar is 50 µm.

**B** and **D**) Kymographs (ZCL2144 marker) revealing occasional unilateral contractions in embryos co-expressing Hep$^{CA}$ in the RN2-Gal4 (**B**) and MzVum-Gal4 (**D**) neurons. For each case, the kymographs correspond to three periods of 1 hour. These embryos never manage to coordinate muscle movements. Both sides, right (R) and left (L) are shown.

**E** and **F**) Flat-prepped, stage 16 embryos expressing syt-GFP under the control of the RN2-Gal4 (**E**) and MzVum-Gal4 (**F**) lines immunostained against GFP (grayscale - top panels and green - bottom panels) and Fas 2 (red - bottom panel). The grey rectangle marks the medial part of the ISN, where synaptic vesicles axonal transport was monitored (see **Movie S9**). Scale bar is 25 µm.

**G** and **H**) Histograms showing synaptotagmin vesicles density and directional motility in control (light green) versus Hep$^{CA}$ expressing (dark green) aCC/ RP2 (**G**) and VUM motoneurons (**H**). No statistically significant differences were observed between control and JNK over-active neurons.

**I** and **J**) Lifetime range distribution of motile synaptotagmin vesicles in aCC/RP2 (**I**) and VUM (**J**) motoneurons. No differences were observed in lifetime between control (light green) and Hep$^{CA}$ expressing (dark green) embryos.

**K** and **L**) Mean velocity range distribution comparison of motile synaptotagmin vesicles in aCC/RP2 (**K**) and VUM (**L**) motoneurons between control (light green) and JNK gain-of function (dark green) conditions. For both sets of motoneurons the distribution was unaltered.

**M)** Regulatory network modulating VNC architectural robustness and condensation and coordinating motor activities. Precise JNK activity levels in early-specified neurons (at least aCC, pCC, RP2 and VUMs) are regulated by a negative feedback loop mediated by Puc. Excessive JNK activity in *puc* mutants or upon overexpressing HepCA in aCC, pCC, RP2 and/or VUMs leads to autonomous defects in axonal paths and dendrites numbers and shape. It also results in a general downregulation of Fas 2 expression (autonomous and non-autonomous)

(discontinuous red inhibitory sign). As a result, the architectural robustness of the VNC will be

affected and its condensation prevented. Further, the coordination of embryonic motor activities

and peristaltic movements are also inhibited. This could be in part consequence of a failure in

wiring optimization arising from the incomplete condensation of the VNC (discontinuous green

arrow).

**SUPPLEMENTAL FIGURES LEGENDS**

# Figure S1

**Figure S1. Fine localization of robustness nodes**

**A)** Self cross-correlation matrix of the Z sections of the image in **B**. A conspicuous node of robust correlation is highlighted (white arrowhead).

**B to D)** Images for Fas 2 (**B** - green in **D**), BP102 (**C** - red in **D**) and their superimposed image (**D**) of a double-stained 16 stage *wt* embryo. Discontinued red lines indicate the localization of the robustness nodes in relation to the VNC pattern. Scale Bar is 10 μm.

**E)** Self cross-correlation matrix of the Z sections of the image in **F**. A conspicuous node of robust correlation is highlighted (white arrowhead).

**F to H)** Images for Fas 2 (**F** - green in **H**), 22C10 (**G** - red in **H**) and their superimposed image (**H**) of a double-stained 16 stage *wt* embryo. Discontinued red lines indicate the localization of the robustness nodes in relation to the VNC pattern. Scale Bar is 10 μm.

# Figure S2

**Figure S2. Robustness and modularity of the VNC architecture in *puc^B48* embryos**

**A)** Maximum projection of the VNC of a *Drosophila* 16-stage *puc^B48* embryo stained with Fas 2.

Defasciculation and collapse of longitudinal connectives is widely spread. Scale Bar is 10 μm.

**B)** Self cross-correlation matrix of **A**. In *puc^B48*, single nodes per segment are consistently

developed (white arrowheads).

**C)** Image cross-correlation score along the A/P axis for **B**. Segmental Fas 2 nodes are on average

at 39 μm from each other in *puc^B48* and the internodal profile flattens.

# Figure S3

**Figure S3. Loss of VNC condensation in *puc* mutants**

**A)** *wt*, **B)** *puc^{E69}* and **C)** *puc^{B48}* embryos Fas 2 immunoreactivity. Lateral view of whole mounts

of stage 17 embryos. Direct measurements of the VNC length were performed along the outline

of the abdominal segments of the VNC as shown in A (dotted red line). Scale Bar is 10 μm.

**D)** Quantification of the VNC in μm (average and standard deviation) for each condition.

Statistically significant differences in length ($p < 0.001$) were detected between the *wt* and *puc^{B48}*

embryos. The length of the VNC of *puc^{E69}* embryos showed a tendency to be consistently longer

than the *wt*, although it was not statistically significant.

# Figure S4

**Figure S4. JNK activity and Cell Death in *puc* mutants**

**A)** Single (left and right) and combined (centre) channels highlighting the expression of Fas 2

(left - green) and anti-P-JNK (right- red) of a stage 17 *wt* embryo. Scale Bar is 10 μm.

**B)** Equivalent as A for a stage 17 *puc$^{E69}$* embryo. Scale Bar is 10 μm.

**C)** Equivalent as A for a stage 17 *puc$^{B48}$* embryo. Scale Bar is 10 μm.

In mutant embryos (**B** and **C**) a significant fraction of the P-JNK signal remained in or around

cell bodies (yellow arrowheads) and is not strictly found as in *wt* only in axons.

**D)** Graph displaying the averaged Integrated Density per segment for the P-JNK signal.

Statistically significant differences in P-JNK levels ($p < 0.001$) were detected between the *wt* and

both mutant conditions.

**E)** From left to right, stage 17 heterozygous and homozygous *puc$^{E69}$* embryo stained for Fas 2

(green) and Dcp1 (red), which highlights those cells in the process of apoptosis. Scale Bar is 10

μm.

**F)** From left to right, equivalent images as in **E** for the *puc$^{B48}$* allele. Scale Bar is 10 μm.

**G)** Graph displaying the averaged Integrated Density per segment for the Dpc1 signal.

Statistically significant differences in Dcp1 levels ($p < 0.001$) were detected between *wt* and

*puc$^{B48}$* embryos. The Dpc1 levels of *puc$^{E69}$* embryos showed a tendency to be consistently larger

than in the *wt*, although they were not statistically significant.

# Figure S5

**Figure S5. Identities of *puc*-expressing neurons**

**A** and **B)** Dorsal and ventral sections highlighting the expression of *puc* (green) in the VNC of
stage 17 UAS-GFP; *puc^E69^I*-Gal4 embryo double-stained for Even-Skipped (red). Scale Bar is 10
µm.

**C)** Diagram indicating those *puc* expressing cells also expressing Even-Skipped [magenta - aCC,
pCC, RP2, U1-U3 (plus weak/variable expression in U4-U5) and 3 out of the 8 to 10 EL
interneurons].

**D** and **E)** Dorsal and ventral sections highlighting the expression of *puc* (green) in the VNC of
stage 17 UAS-GFP; *puc^E69^I*-Gal4 embryo double-stained for Engrailed (red). Scale Bar is 10 µm.

**F)** Diagram indicating those *puc*-expressing cells also expressing Engrailed (cyan - NL1, NL2,
and iVUM5 neurons).

# Figure S6

**Figure S6.** *puc*$^{B48}$ **expressing neurons**

**A** to **D)** Dorsal and ventral sections highlighting the expression of *puc* (green) in the VNC of

stage 17 UAS-GFP; *puc*$^{B48}$-Gal4 embryo double-stained for the pan-neuronal marker Elav (**A**

and **B** - red) or the glial marker Repo (**C** and **D** - red). *puc* is expressed in a subset of neurons in

each segments but not in glia. Scale Bar is 10 μm.

**E** to **H)** Dorsal and ventral sections highlighting the expression of *puc* (green) in the VNC of

stage 17 UAS-GFP; *puc*$^{B48}$-Gal4 embryo double-stained for Even-Skipped (**E** and **F** - red) or

Engrailed (**G** and **H** - red). Scale Bar is 10 μm.

**I)** Cartoon representing the full set of *puc*$^{B48}$ expressing neurons per segment in the VNC.

**J)** Cartoon representing the *puc*$^{E69}$ neurons highlighting the cells expressing *puc*$^{B48}$ (deep blue).

# Figure S6

**A**

**Unilateral Constrictions**

**B**

**Head Movements**

**C**

**Forward Peristalsis**

**Figure S7. Types of embryonic muscles movements**

Snapshots from **Movie S6** displaying mid-plane images of a stage 17 embryo carrying two GFP protein traps expressed at muscle Z-lines (w; G203, ZCL2144). Three main types of movement were observed during the muscle coordination process. Red arrows point to the contracting muscle units. Anterior is to the left. Scale bar is 50 µm.

**A)** Unilateral muscle contractions. They can occured at any embryo side and in different segments.

**B)** Head turning and mouth hook pinching.

**C)** Peristalsis. They can be incomplete or complete and develop forward (in the image) or backward displacements.

**SUPPLEMENTAL MOVIES LEGENDS**

**Movie S1. *Drosophila* Fas 2 positive connectives**

Animated 3D reconstruction of the Fas 2 positive connectives of a section of the VNC of a late *Drosophila* embryo (Stage 17).

**Movie S2. *Drosophila* Fas 2 positive connectives of a *puc^{E69}* embryo**

Animated 3D reconstruction of the Fas 2 positive connectives of a section of the VNC of a late *Drosophila puc^{E69}* mutant embryo (Stage 17).

**Movie S3. Dynamics of *puc^{E69}* expression**

Time course of *puc^{E69}* expression from a UAS-GFP; *puc^{E69}I*-Gal4 viable line. *puc* expression can be observed from early stage 13 in a few neurons at the midline. As development progressed, more and more neurons express *puc*, both within the VNC and at the peripheral nervous system (PNS). Scale bar is 50 µm. Anterior is to the left.

1016 **Movie S4. Minimum consensus for the segmental $puc^{E69}$ expression**

1017 Animated 3D cartoon representing a single segment of a late *Drosophila* $puc^{E69}$ mutant embryo

1018 (Stage 17). The 19 *puc* expressing cells per hemisegment represented in the minimum consensus

1019 are shown in different colors.

1020

1021 **Movie S5. Axonal and dendritic landscape of pioneer neurons upon JNK hyper-activation**

1022 Animated 3D reconstruction of the axonal and dendritic pattern of stage 16 embryos expressing

1023 mCD8-GFP (**A**) and mCD8-GFP and Hep$^{CA}$ (**B**) under the control of the RN2-Gal4 line. Traces

1024 of the dendrites (green) are over-imposed. Scale bar is 10 μm.

1025

1026 **Movie S6. Embryonic motor patterns**

1027 Representative mid-plane movies of a stage 17 embryo carrying two GFP protein traps expressed

1028 at muscle Z-lines (w; G203, ZCL2144). **A)** Forward peristalsis. **B)** Backward Peristalsis. **C)** Left

1029 twitch. **D)** Right twitch. **E)** Head tumble. **F)** Mandible pinch. Scale bar is 50 μm. Anterior is to

1030 the left.

1031

**Movie S7. Time course of motor maturation**

Representative mid-plane movies (250 time points per stage) of an embryo carrying two GFP

protein traps expressed at muscle Z-lines (w; G203, ZCL2144). From top to bottom stages **A**

(Unilateral twitches and incomplete peristalsis), **B** (Resting period), **C** (Complete peristalsis), **D**

(Late resting period) and **E** (Hatching movements). Color coded as in **Figure 6**. Scale bar is 50

μm. Anterior is to the left.

1038

1039 **Movie S8. Motor activity of $puc^{E69}$ embryos**

1040 Representative mid-plane movie (1000 time points) of a late stage 17 $puc^{E69}$ embryo carrying

1041 two GFP protein traps expressed at muscle Z-lines (w; G203, ZCL2144). Scale bar is 50 μm.

1042 Anterior is to the left.

1043

1044

1045 **Movie S9. Synaptic vesicles axonal transport in RN2 neurons**

1046 Stage 16 embryos expressing Synaptotagmin-GFP under the control of the RN2-Gal4 in the

1047 absence **(A)** or presence **(B)** of Hep$^{CA}$. Left panels present representative single plane movies

1048 (Fire Luts) with 1 sec interval for a total of 60 cycles. Right panels display in different colors the

1049 traces of the different particles analyzed. Proximal is to the left. Scale bar is 25 μm.

1050

1051