1   **The SEK-1 p38 MAP kinase pathway modulates Gq signaling in *C. elegans***

4   Jill M. Hoyt, Samuel K. Wilson, Madhuri Kasa, Jeremy S. Rise,

5   Irini Topalidou, Michael Ailion

Department of Biochemistry, University of Washington, Seattle, WA, 98195

Running Title: SEK-1 modulates Gq signaling

key words: p38 MAPK pathway, Gq signaling, animal behavior, *C. elegans*

Corresponding author:

Michael Ailion

Department of Biochemistry

University of Washington

Box 357350

1705 NE Pacific St

Seattle, WA 98195

Phone: 206-685-0111

email: mailion@uw.edu

Abstract

Gq is a heterotrimeric G protein that is widely expressed in neurons and regulates neuronal activity. To identify pathways regulating neuronal Gq signaling we performed a forward genetic screen in *Caenorhabditis elegans* for suppressors of activated Gq. One of the suppressors is an allele of *sek-1*, which encodes a mitogen-activated protein kinase kinase (MAPKK) in the p38 MAPK pathway. Here we show that *sek-1* mutants have a slow locomotion rate and that *sek-1* acts in acetylcholine neurons to modulate both locomotion rate and Gq signaling. Furthermore, we find that *sek-1* acts in mature neurons to modulate locomotion. Using genetic and behavioral approaches we demonstrate that other components of the p38 MAPK pathway also play a positive role in modulating locomotion and Gq signaling. Finally, we find that mutants in the SEK-1 p38 MAPK pathway partially suppress an activated mutant of the sodium leak channel NCA-1/NALCN, a downstream target of Gq signaling. Our results suggest that the SEK-1 p38 pathway may modulate the output of Gq signaling through NCA-1.

Introduction

Gq is a widely expressed heterotrimeric G protein that regulates a variety of biological processes ranging from neurotransmission to cardiovascular pathophysiology (Sánchez-Fernández *et al.* 2014). In the canonical Gq pathway, Gq activates phospholipase Cβ (PLCβ), which cleaves phosphatidylinositol 4,5-bisphosphate (PIP$_2$) into the second messengers diacylglycerol (DAG) and inositol trisphosphate (IP$_3$) (Rhee 2001). In addition to PLCβ, other Gq effectors have been identified including kinases, such as protein kinase Cζ (PKCζ) and Bruton's tyrosine kinase (Btk) (Bence *et al.* 1997; García-Hoz *et al.* 2010; Vaqué *et al.* 2013), and guanine nucleotide exchange factors (GEFs) for the small GTPase Rho, such as Trio (Williams *et al.* 2007; Vaqué *et al.* 2013). These noncanonical effectors bridge the activation of Gq to other cellular signaling cascades.

In order to study noncanonical pathways downstream of Gq, we used the nematode *C. elegans* which has a single Gαq homolog (EGL-30) and conservation of the other components of the Gq signaling pathway (Koelle 2016). In neurons, EGL-30 signals through EGL-8 (PLCβ) (Lackner *et al.* 1999) and UNC-73 (ortholog of Trio RhoGEF) (Williams *et al.* 2007). UNC-73 activates RHO-1 (ortholog of RhoA), which has been shown to enhance neurotransmitter release through both diacylglycerol kinase (DGK-1)-dependent and DGK-1-independent pathways (McMullan *et al.* 2006).

To identify additional signaling pathways that modulate Gq signaling, we screened for suppressors of the activated Gq mutant *egl-30(tg26)* (Doi and Iwasaki 2002). *egl-30(tg26)* mutant animals exhibit hyperactive locomotion and a "loopy" posture in which worms have exaggerated, deep body bends and loop onto themselves (Bastiani *et al.*

2003; Topalidou *et al.* 2017). Here we identify one of the suppressors as a deletion

allele in the gene *sek-1*. SEK-1 is a mitogen-activated protein kinase kinase (MAPKK),

the *C. elegans* ortholog of mammalian MKK3/6 in the p38 MAPK pathway (Tanaka-Hino

et al. 2002). The p38 MAPK pathway has been best characterized as a pathway

activated by a variety of cellular stresses and inflammatory cytokines (Kyriakis and

Avruch 2012). However, the p38 MAPK pathway has also been shown to be activated

downstream of a G protein-coupled receptor in rat neurons (Huang *et al.* 2004). Btk, a

member of the Tec family of tyrosine kinases, has been shown to act downstream of Gq

to activate the p38 MAPK pathway (Bence *et al.* 1997), but *C. elegans* lacks Btk and

other Tec family members (Plowman *et al.* 1999).

SEK-1 is activated by the MAPKKK NSY-1 (ortholog of ASK1) and activates the p38

MAPKs PMK-1 and PMK-2 (Andrusiak and Jin 2016). The p38 MAPK pathway

consisting of NSY-1, SEK-1, and PMK-1 is required for innate immunity in *C. elegans*

(Kim *et al.* 2002). NSY-1 and SEK-1 are also required for the specification of the

asymmetric AWC olfactory neurons (Sagasti *et al.* 2001; Tanaka-Hino *et al.* 2002); the

p38 orthologs PMK-1 and PMK-2 function redundantly in AWC specification (Pagano *et*

al. 2015). For both innate immunity and AWC specification, the p38 MAPK pathway acts

downstream of the adaptor protein TIR-1 (an ortholog of SARM) (Couillault *et al.* 2004;

Chuang and Bargmann 2005). Here we show that the pathway consisting of TIR-1,

NSY-1, SEK-1, PMK-1 and PMK-2 also acts to modulate locomotion downstream of Gq

signaling.

Materials and Methods

C. elegans strains and maintenance

All strains were cultured using standard methods and maintained at 20°C

(Brenner 1974). The *sek-1(yak42)* mutant was isolated from an ENU mutagenesis

suppressor screen of the activated Gq mutant *egl-30(tg26)* (Ailion *et al.* 2014). *sek-*

1(yak42) was outcrossed away from *egl-30(tg26)* before further analysis. Double mutant

strains were constructed using standard methods (Fay 2006), often with linked

fluorescent markers (Frokjaer-Jensen *et al.* 2014) to balance mutations with subtle

visible phenotypes. Table S1 contains all the strains used in this study.

Mapping

yak42 was mapped using its slow locomotion phenotype and its *egl-30(tg26)*

suppression phenotype. *yak42* was initially mapped to the X chromosome using strains

EG1000 and EG1020, which carry visible marker mutations. These experiments

showed that *yak42* was linked to *lon-2*, but at least several map units away. *yak42* was

further mapped to about one map unit (m.u.) away from the red fluorescent insertion

marker *oxTi668* which is located at +0.19 m.u. on the X chromosome.

Whole-genome sequencing

Strain XZ1233 *egl-30(tg26); yak42* was used for whole-genome sequencing to

identify candidate *yak42* mutations. XZ1233 was constructed by crossing a 2X

outcrossed *yak42* strain back to *egl-30(tg26)*. Thus, in XZ1233, *yak42* has been

outcrossed 3X from its original isolate. DNA was isolated from XZ1233 and purified

according to the Hobert Lab protocol (http://hobertlab.org/whole-genome-sequencing/).

Ion Torrent sequencing was performed at the University of Utah DNA Sequencing Core

Facility. The resulting data contained 10,063,209 reads of a mean read length of 144

bases, resulting in about 14X average coverage of the *C. elegans* genome. The

sequencing data were uploaded to the Galaxy web platform and we used the public

server at usegalaxy.org to analyze the data (Afgan *et al.* 2016). We identified and

annotated variants with the Unified Genotyper and SnpEff tools, respectively (DePristo

et al. 2011; Cingolani *et al.* 2012). We filtered out variants found in other strains we

sequenced, leaving us with 605 homozygous mutations. The X chromosome contained

94 mutations: 55 SNPs and 39 indels. Of these, four SNPs were non-synonymous

mutations in protein-coding genes, but only two were within 5 m.u. of *oxTi668.* However,

we were unable to identify *yak42* from the candidate polymorphisms located near

oxTi668. Transgenic expression of the most promising candidate *pcyt-1* did not rescue

yak42. Instead, to identify possible deletions, we scrolled through 2 MB of aligned reads

on the UCSC Genome Browser starting at -4.38 m.u. and moving towards the middle of

the chromosome (0 m.u.), looking for regions that lacked sequence coverage. We found

a 3713 bp deletion that was subsequently confirmed to be the *yak42* causal mutation,

affecting the gene *sek-1* located at -1.14 m.u.

Locomotion assays

Locomotion assay plates were made by seeding 10 cm nematode growth

medium plates with 150 µl of an *E. coli* OP50 stock culture, spread with sterile glass

beads to cover the entire plate. Bacterial lawns were grown at room temperature

(22.5°C -24.5°C) for 24 hrs and then stored at 4°C until needed. All locomotion assays

were performed on first-day adults at room temperature (22.5°C -24.5°C). L4 stage

larvae were picked the day before the assay and the experimenter was blind to the

genotypes of the strains assayed. For experiments on strains carrying

extrachromosomal arrays, the *sek-1(km4)* control worms were animals from the same

plate that had lost the array.

Body bend assays were performed as described (Miller *et al.* 1999). A single

animal was picked to the assay plate, the plate lid was returned, and the animal allowed

to recover for 30 s. Body bends were then counted for one minute, counting each time

the worm's tail reached the minimum or maximum amplitude of the sine wave. All

strains in an experiment were assayed on the same assay plate. For experiments with

egl-8, *unc-73*, and *rund-1* mutants, worms were allowed a minimal recovery period (until

the worms started moving forward, 5 sec maximum) prior to counting body bends.

For the heat shock experiment, plates of first-day adults were parafilmed and

heat-shocked in a 34°C water bath for 1 hr. Plates were then un-parafilmed and

incubated at 20°C for five hours before performing body bend assays.

Radial locomotion assays were performed by picking animals to the middle of an

assay plate. Assay plates were incubated at 20°C for 20 hr and the distances of the

worms from the starting point were measured.

Quantitative analysis of the waveform of worm tracks was performed as

described (Topalidou *et al.* 2017). Briefly, worm tracks were photographed and Image J

was used to measure the period and amplitude. The value for each animal was the

average of five period/amplitude ratios.

C. elegans pictures

Pictures of worms were taken at 60X on a Nikon SMZ18 microscope with the DS-

L3 camera control system. The worms were age-matched as first-day adults and each

experiment set was photographed on the same locomotion assay plate prepared as

described above. The images were processed using ImageJ and were rotated, cropped,

and converted to grayscale.

Molecular biology

Plasmids were constructed using the Gateway cloning system (Invitrogen).

Plasmids and primers used are found in Table S2. The *sek-1* cDNA was amplified by

RT-PCR from worm RNA and cloned into a Gateway entry vector. To ensure proper

expression of *sek-1*, an operon GFP was included in expression constructs with the

following template: (promoter)p::*sek-1*(cDNA)::*tbb-2utr::gpd-2 operon::GFP::H2B:cye-*

1utr (Frøkjær-Jensen *et al.* 2012). This resulted in untagged SEK-1, but expression

could be monitored by GFP expression.

Injections

C. elegans strains with extrachromosomal arrays were generated by standard

methods (Mello *et al.* 1991). Injection mixes were made with a final total concentration

of 100 ng/µL DNA. Constructs were injected at 5 ng/µL, injection markers at 5 ng/µL,

and the carrier DNA Litmus 38i at 90 ng/µL. Multiple lines of animals carrying

extrachromosomal arrays were isolated and had similar behaviors as observed by eye.

The line with the highest transmittance of the array was assayed.

Statistical analysis

At the beginning of the project, a power study was conducted on pilot body bend

assays using wild type and *sek-1(yak42)* worms. To achieve a power of 0.95, it was

calculated that 17 animals should be assayed per experiment. Data were analyzed to

check if normally distributed (using the D'Agostino-Pearson and Shapiro-Wilk normality

tests) and then subjected to the appropriate analysis using GraphPad Prism 5. For data

sets with three or more groups, if the data were normal they were analyzed with a one-

way ANOVA; if not, with a Kruskal-Wallis test. Post-hoc tests were used to compare

data sets within an experiment. Reported p-values are corrected. Table S3 contains the

statistical tests for each experiment. $p<0.05$ = *; $p<0.01$ = **; $p<0.001$ = ***.

Reagent and Data Availability

190   Strains and plasmids are shown in Table S1 and Table S2 and are available from

191   the *Caenorhabditis* Genetics Center (CGC) or upon request. The authors state that all

192   data necessary for confirming the conclusions presented in the article are represented

193   fully within the article and Supplemental Material.

Results

sek-1 suppresses activated Gq

To identify genes acting downstream of Gαq, we performed a forward genetic

screen for suppressors of the activated Gq mutant, *egl-30(tg26)* (Doi and Iwasaki 2002).

egl-30(tg26) worms are hyperactive and have a "loopy" posture characterized by an

exaggerated waveform (Figure 1B-1E). Thus, we screened for worms that are less

hyperactive and less loopy. We isolated a recessive suppressor, *yak42*, and mapped it

to the middle of the X chromosome (see Materials and Methods). Whole-genome

sequencing revealed that *yak42* carries a large deletion of the *sek-1* gene from

upstream of the start codon into exon 4 (Figure 1A). *yak42* also failed to complement

sek-1(km4), a previously published *sek-1* deletion allele, for the Gq suppression

phenotype (Figure 1A) (Tanaka-Hino *et al.* 2002).

egl-30(tg26) double mutants with either *sek-1(yak42)* or *sek-1(km4)* are not loopy

(Figure 1B-1D) and are not hyperactive (Figure 1E and S1A). *sek-1(yak42)* was

outcrossed from *egl-30(tg26)* and assayed for locomotion defects. Both the *sek-*

1(yak42) and *sek-1(km4)* mutants are coordinated but move more slowly than wild-type

(Figure 1F). The *sek-1(ag1)* point mutation (Kim *et al.* 2002) also causes a similar slow

locomotion phenotype (Figure S1B). To test whether the *egl-30(tg26)* suppression

phenotype might be an indirect effect of the slow locomotion of a *sek-1* mutant, we built

an *egl-30(tg26)* double mutant with a mutation in *unc-82*, a gene required for normal

muscle structure. *unc-82* mutants are coordinated but move slowly, similar to a *sek-1*

mutant (Hoppe *et al.* 2010). However, although an *egl-30(tg26) unc-82(e1220)* double

mutant moves more slowly than *egl-30(tg26)* (Figure S1C), it is still loopy (Figure 1B-

1D). Thus, *sek-1* appears to be a specific suppressor of activated *egl-30*.

The *egl-30(tg26)* allele causes an R243Q missense mutation in the Gα switch III

region that has been shown to reduce both the intrinsic GTPase activity of the G protein

and render it insensitive to GTPase-activation by a regulator of G protein signaling

(RGS) protein, thus leading to increased G protein activation (Natochin and Artemyev

2003). To test whether the suppression of *egl-30(tg26)* by *sek-1* is specific for this *egl-*

30 allele, we built a double mutant between *sek-1(km4)* and the weaker activating

mutation *egl30(js126)*. *egl-30(js126)* causes a V180M missense mutation in the Gα

switch I region immediately adjacent to one of the key residues required for GTPase

catalysis (Hawasli *et al.* 2004). Thus, the *tg26* and *js126* alleles activate EGL-30

through different mechanisms. The *sek-1(km4)* mutant also suppresses the

hyperactivity and loopy waveform of *egl-30(js126)* (Figure 1G, 1H), demonstrating that

sek-1 suppression of activated *egl-30* is not allele-specific.

EGL-30/Gαq is negatively regulated by GOA-1, the worm Gαo/i ortholog, and the

RGS protein EAT-16 (Hajdu-Cronin *et al.* 1999). We tested whether *sek-1* also

suppresses the *goa-1* and *eat-16* loss-of-function mutants that cause a hyperactive and

loopy phenotype similar to activated *egl-30* mutants. *sek-1(km4)* suppresses the

hyperactivity and loopy waveform of *goa-1(sa734)* (Figure S1D, S1E). However, though

sek-1(km4) suppresses the hyperactivity of *eat-16(tm775)*, it did not significantly

suppress the loopy waveform (Figure S1F, S1G). One possible downstream effector of

GOA-1 is the DAG kinase DGK-1 that inhibits DAG-dependent functions such as

synaptic vesicle release (Nurrish *et al.* 1999; Miller *et al.* 1999). *dgk-1(sy428)* animals

are hyperactive, but the *sek-1 dgk-1* double mutant is uncoordinated and looks like

neither *sek-1* nor *dgk-1* mutants, confounding the interpretation of how *sek-1* genetically

interacts with *dgk-1*.

sek-1 acts in mature acetylcholine neurons

egl-30 is widely expressed and acts in neurons to modulate locomotion (Lackner

et al. 1999), so it is possible that *sek-1* also acts in neurons to modulate Gq signaling.

sek-1 is expressed in neurons, intestine, and several other tissues (Tanaka-Hino *et al.*

2002) and has been shown to function in GABA neurons to promote synaptic

transmission (Vashlishan *et al.* 2008).

To identify the cell type responsible for the *sek-1* locomotion phenotypes, we

expressed the wild-type *sek-1* cDNA under different cell-specific promoters and tested

for transgenic rescue of a *sek-1* null mutant. Expression of *sek-1* in all neurons (using

the *unc-119* promoter) or in acetylcholine neurons (*unc-17* promoter) was sufficient to

rescue the *sek-1* mutant slow locomotion phenotype, but expression in GABA neurons

(*unc-47* promoter) was not sufficient to rescue (Figure 2A, B). These results indicate

that *sek-1* acts in acetylcholine neurons to modulate locomotion rate.

We next tested whether *sek-1* acts in neurons to suppress *egl-30(tg26)*.

Expression of *sek-1* under pan-neuronal and acetylcholine neuron promoters reversed

the *sek-1* suppression of *egl-30(tg26)*. Specifically, *egl-30(tg26) sek-1* double mutants

expressing wild-type *sek-1* in all neurons or acetylcholine neurons resembled the *egl-*

30(tg26) single mutant (Figure 2C-E). However, expression of *sek-1* in GABA neurons

did not reverse the suppression phenotype (Figure 2C-E). Together, these data show

that *sek-1* acts in acetylcholine and not GABA neurons to modulate both wild-type

locomotion rate and to modulate Gq signaling.

To narrow down the site of *sek-1* action, we expressed *sek-1* in head (*unc-17H*

promoter) and motorneuron (*unc-17β* promoter) acetylcholine neuron subclasses

(Topalidou *et al.* 2017). Expression of *sek-1* in acetylcholine motorneurons rescued the

sek-1 slow locomotion phenotype (Figure S2A), suggesting that the slow locomotion of

sek-1 mutants is due to a loss of *sek-1* in acetylcholine motorneurons. However,

expression of *sek-1* in either the head acetylcholine neurons or motorneurons partially

reversed the *sek-1* suppression of *egl-30(tg26)* hyperactivity (Figure S2B), suggesting

that the hyperactivity of activated Gq mutants may result from excessive Gq signaling in

both head acetylcholine neurons and acetylcholine motorneurons; *sek-1* may act in Gq

signaling in both neuronal cell types. By contrast, expression of *sek-1* in head

acetylcholine neurons but not motorneurons reversed the *sek-1* suppression of the *egl-*

30(tg26) loopy waveform (Figure S2C), suggesting that the loopy posture of activated

Gq mutants may result from excessive Gq signaling in head acetylcholine neurons, and

sek-1 may act in those neurons to control body posture.

Because *sek-1* acts in the development of the AWC asymmetric neurons, we

asked whether *sek-1* also has a developmental role in modulating locomotion by testing

whether adult-specific *sek-1* expression (driven by a heat-shock promoter) is sufficient

to rescue the *sek-1* mutant. We found that *sek-1* expression in adults rescues the *sek-1*

slow locomotion phenotype (Figure 2F). This result indicates that *sek-1* is not required

for development of the locomotion circuit and instead acts in mature neurons to

modulate locomotion.

The p38 MAPK pathway is a positive regulator of Gq signaling

SEK-1 is the MAPKK in the p38 MAPK pathway consisting of the adaptor protein

TIR-1, NSY-1 (MAPKKK), SEK-1 (MAPKK), and PMK-1 or PMK-2 (MAPKs) (Tanaka-

Hino *et al.* 2002; Andrusiak and Jin 2016). We tested whether the entire p38 MAPK

signaling module also modulates locomotion rate and suppression of activated Gq. Both

tir-1(tm3036) and *nsy-1(ok593)* mutant animals have slow locomotion on their own and

also suppress the hyperactivity and loopy waveform of *egl-30(tg26)* (Figure 3A-D, G and

H). We also tested single mutants in each of the three worm p38 MAPK genes (*pmk-1,*

pmk-2 and *pmk-3)* and a *pmk-2 pmk-1* double mutant. Although we found that the *pmk-*

2 and *pmk-3* single mutants were slightly slow on their own, only the *pmk-2 pmk-1*

double mutant phenocopied *sek-1* and suppressed both the hyperactivity and loopy

waveform of *egl-30(tg26)* (Figure 3E-H). Thus, *pmk-2* and *pmk-1* act redundantly

downstream of *sek-1* to suppress *egl-30(tg26).* These data suggest that the p38 MAPK

pathway modulates locomotion rate in *C. elegans* and acts genetically downstream of

egl-30.

The JNK MAPK pathway, related to the p38 MAPK family, also modulates

locomotion in *C. elegans*. Specifically, the JNK pathway members *jkk-1* (JNK MAPKK)

and *jnk-1* (JNK MAPK) have been shown to act in GABA neurons to modulate

locomotion (Kawasaki 1999). We found that the *jkk-1* and *jnk-1* single mutants had slow

locomotion and that the double mutants with p38 MAPK pathway members exhibited an

additive slow locomotion phenotype (Figure S3A). Moreover, neither *jkk-1* nor *jnk-1*

suppressed the loopy phenotype of *egl-30(tg26)* (Figure S3B). Thus, the JNK and p38

MAPK pathways modulate locomotion independently and the JNK pathway is not

involved in Gq signaling.

We also tested the involvement of possible p38 MAPK pathway effectors. One of

the targets of PMK-1 is the transcription factor ATF-7 (Shivers *et al.* 2010). Both the *atf-*

7(qd22 qd130) loss-of-function mutant and the *atf-7(qd22)* gain-of-function mutant

moved slowly compared to wild-type animals (Figure S3C). However, *atf-7(qd22 qd130)*

did not suppress the loopiness of *egl-30(tg26)* (Figure S3B), suggesting that *atf-7* is not

a target of this pathway, or else it acts redundantly with other downstream p38 MAPK

targets. We also tested *gap-2*, the closest *C. elegans* homolog of ASK1-interacting

Protein (AIP1) which activates ASK1 (the ortholog of *C. elegans* NSY-1) in mammalian

systems (Zhang *et al.* 2003). A *C. elegans gap-2* mutant has no locomotion defect

(Figure S3D). Finally, we tested VHP-1, a phosphatase for p38 and JNK MAPKs that

inhibits p38 MAPK signaling (Kim *et al.* 2004). However, the *vhp-1(sa366)* mutant also

has no locomotion defect (Figure S3D).

egl-30(tg26) animals are loopy and hyperactive so we tested whether increased

activation of the TIR-1/p38 MAPK signaling module causes similar phenotypes. The *tir-*

1(ky648tg26) allele leads to a gain-of-function phenotype in the AWC neuron

specification (Chang *et al.* 2011), but does not cause loopy or hyperactive locomotion

(Figure S3E, F).

Genetic interactions of *sek-1* with pathways acting downstream of Gq

Our forward genetic screen for suppressors of *egl-30(tg26)* identified mutants

that fall into three different categories: mutants in the canonical Gq pathway such as the

PLC *egl-8* ((Lackner *et al.* 1999), mutants in the RhoGEF Trio pathway such as *unc-73*

(Williams et al., 2007), and mutants that affect dense-core vesicle biogenesis and

release (Ailion *et al.* 2014; Topalidou *et al.* 2016).

To test if *sek-1* acts in any of these pathways we built double mutants between

sek-1 and members of each pathway. Loss-of-function alleles of *egl-8(sa47)*, *unc-*

73(ox317), and *rund-1(tm3622)* have slow locomotion (Figure 4A-C). We found that

sek-1 enhances the slow locomotion phenotype of *egl-8* and *rund-1* single mutants,

suggesting that *sek-1* does not act in the same pathway as *egl-8* or *rund-1* (Figure 4A,

B). By contrast, *sek-1* does not enhance the slow locomotion phenotype of *unc-73*

mutants (Figure 4C), suggesting that *sek-1* may act in the same genetic pathway as the

Trio RhoGEF *unc-73.*

We next tested whether *sek-1* interacts with *rho-1*, encoding the small G protein

Rho that is activated by Trio. Because *rho-1* is required for viability (Jantsch-Plunger *et*

al. 2000), we used an integrated transgene overexpressing an activated *rho-1* mutant

allele specifically in acetylcholine neurons. Animals carrying this activated RHO-1

transgene, referred to here as *rho-1(gf)*, have a loopy posture reminiscent of *egl-*

30(tg26) (McMullan *et al.* 2006), and a decreased locomotion rate (Figure 4D-F). *rho-*

1(gf) sek-1(km4) double mutants had a loopy body posture like *rho-1(gf)* and an even

slower locomotion rate (Figure 4D-F), suggesting that *sek-1* and *rho-1(gf)* mutants have

additive locomotion phenotypes. However, both *sek-1(km4)* and *sek-1(yak42)* weakly

suppress the slow growth rate of the *rho-1(gf)* mutant (data not shown). Because *sek-1*

does not enhance *unc-73* mutants and suppresses some aspects of the *rho-1(gf)*

mutant, *sek-1* may modulate output of the Rho pathway, though it probably is not a

direct transducer of Rho signaling.

sek-1 and nsy-1 partially suppress activated NCA

To clarify the relationship of the SEK-1 p38 MAPK pathway to the Rho pathway

acting downstream of Gq, we examined interactions with *nca-1*, a downstream target of

the Gq-Rho pathway (Topalidou *et al.* 2017). NCA-1 and its orthologs are sodium leak

channels associated with rhythmic behaviors in several organisms (Nash *et al.* 2002; Lu

et al. 2007; Shi *et al.* 2016). In *C. elegans*, NCA-1 potentiates persistent motor circuit

activity and sustains locomotion (Gao *et al.* 2015).

364   We tested whether *sek-1* and *nsy-1* mutants suppress the activated NCA-1

mutant *ox352*, referred to as *nca-1(gf)*. The *nca-1(gf)* animals are coiled and

uncoordinated; thus, it is difficult to measure their locomotion rate by the body bend

assay because they do not reliably propagate sinusoidal waves down the entire length

of their body. Instead, we used a radial locomotion assay in which we measured the

distance animals moved from the center of a plate. *nca-1(gf)* double mutants with either

sek-1(km4) or *nsy-1(ok593)* uncoil a bit but still exhibit uncoordinated locomotion

(Figure 5A). In fact, though these double mutants show more movement in the anterior

half of their bodies than *nca-1(gf)*, they propagate body waves to their posterior half

even more poorly than the *nca-1(gf)* mutant. However, both *sek-1* and *nsy-1* partially

suppress the loopy waveform of the *nca-1(gf)* mutant (Figure 5A, B) and in radial

locomotion assays, *sek-1* and *nsy-1* weakly suppressed the *nca-1(gf)* locomotion defect

(Figure 5C). Additionally, both *sek-1* and *nsy-1* partially suppress the small body size of

nca-1(gf) (Figure S4A). Together these data suggest that mutations in the SEK-1 p38

MAPK pathway suppress some aspects of the *nca-1(gf)* mutant.

379   Given that *sek-1* acts in acetylcholine neurons to modulate wild-type and *egl-*

30(tg26) locomotion, we tested whether *sek-1* also acts in these neurons to suppress

nca-1(gf). Expression of *sek-1* in all neurons or in acetylcholine neurons of *nca-1(gf)*

sek-1(km4) animals restored the *nca-1(gf)* loopy phenotype (Figure 5D, E). By contrast,

expression of *sek-1* in GABA neurons did not affect the loopy posture of the *nca-1(gf)*

sek-1 double mutant (Figure 5D, E). These data suggest that *sek-1* acts in acetylcholine

neurons to modulate the body posture of *nca-1(gf)* as well. However, in radial

locomotion assays, expression of *sek-1* in none of these neuron classes significantly

altered the movement of the *nca-1(gf) sek-1* double mutant (Figure S4B), though the

weak suppression of *nca-1(gf)* by *sek-1* in this assay makes it difficult to interpret these

negative results. To further narrow down the site of action of *sek-1* for its NCA

suppression phenotypes, we expressed it in subclasses of acetylcholine neurons.

Surprisingly, expression of *sek-1* in acetylcholine motorneurons but not head

acetylcholine neurons was sufficient to restore the loopy posture of the *nca-1(gf)* mutant

(Figure 5E), the opposite of what we found for *sek-1* modulation of the loopy posture of

the activated Gq mutant, suggesting that the loopy posture of *nca-1(gf)* mutants may

result from excessive NCA-1 activity in acetylcholine motorneurons. Additionally,

expression of *sek-1* in either the head acetylcholine neurons or the motorneurons

restored the *nca-1(gf)* small body size phenotype (Figure S4C). We make the tentative

conclusion that *sek-1* acts in acetylcholine neurons to modulate *nca-1(gf)* body posture

and size, but we were not able to conclusively narrow down its site of action further,

possibly due to the uncoordinated phenotype of *nca-1(gf)* and the weaker suppression

of *nca-1(gf)* by *sek-1* .

Discussion

The p38 MAPK pathway has been best characterized as a pathway activated by

a variety of cellular stresses and inflammatory cytokines (Kyriakis and Avruch 2012), but

it has also been implicated in neuronal function, including some forms of mammalian

synaptic plasticity (Bolshakov *et al.* 2000; Rush *et al.* 2002; Huang *et al.* 2004). In this

study we identified a new neuronal role for the mitogen-activated protein kinase kinase

SEK-1 and the p38 MAPK pathway as a positive modulator of locomotion rate and Gq

signaling. The physiological importance of this pathway is clear under conditions of

elevated Gq signaling but is less obvious during normal wild-type locomotion, consistent

with the observation that *sek-1* mutations have a relatively weak effect on synaptic

transmission in a wild-type background (Vashlishan *et al.* 2008). Thus, the SEK-1 p38

MAPK pathway may be more important for modulation of Gq signaling and synaptic

strength than for synaptic transmission per se.

In addition to SEK-1, we identified other p38 pathway components that modulate

Gq signaling. Specifically, we found that *tir-1, nsy-1* and *pmk-1 pmk-2* mutants exhibit

locomotion defects identical to *sek-1* and suppress activated Gq, suggesting that they

act in a single p38 pathway to modulate signaling downstream of Gq. These results

indicate a redundant function for PMK-1 and PMK-2 in modulating locomotion rate and

Gq signaling. PMK-1 and PMK-2 also act redundantly for some other neuronal roles of

the p38 pathway, such as the development of the asymmetric AWC neurons and to

regulate induction of serotonin biosynthesis in the ADF neurons in response to

pathogenic bacteria (Shivers *et al.* 2009; Pagano *et al.* 2015). By contrast, PMK-1 acts

alone in the intestine to regulate innate immunity and in interneurons to regulate

trafficking of the GLR-1 glutamate receptor (Pagano *et al.* 2015; Park and Rongo 2016).

What are the downstream effectors of the SEK-1 p38 MAPK pathway that

modulates locomotion? There are several known downstream effectors of p38 MAPK

signaling in *C. elegans*, including the transcription factor ATF-7 (Shivers *et al.* 2010).

Our data indicate that ATF-7 is not required for the p38 MAPK-dependent modulation of

Gq signaling. The p38 MAPK pathway may activate molecules other than transcription

factors or may activate multiple downstream effectors.

How does the SEK-1 p38 pathway modulate the output of Gq signaling? One of

the pathways that transduces signals from Gq includes the RhoGEF Trio/UNC-73, the

small GTPase Rho, and the cation channel NALCN/NCA-1 (Williams *et al.* 2007;

Topalidou *et al.* 2017). Compared to other pathways downstream of Gq, mutants in the

Rho-Nca pathway are particularly strong suppressors of the loopy waveform phenotype

of the activated Gq mutant (Topalidou *et al.* 2017). Similarly, we found that mutations in

the SEK-1 p38 MAPK pathway strongly suppress the loopy waveform of the activated

Gq mutant, suggesting that the SEK-1 pathway might modulate Gq signal output

through the Rho-Nca branch. Consistent with this, we found that mutations in the SEK-1

p38 MAPK pathway partially suppress an activated NCA-1 mutant. Given the

precedence for direct phosphorylation of sodium channels by p38 to regulate channel

properties (Wittmack *et al.* 2005; Hudmon *et al.* 2008), it is possible that PMK-1 and

PMK-2 phosphorylate NCA-1 to regulate its expression, localization, or activity.

Consistent with the observation that Gq acts in acetylcholine neurons to stimulate

synaptic transmission (Lackner *et al.* 1999), we found that *sek-1* acts in acetylcholine

neurons to modulate the locomotion rate in both wild-type and activated Gq mutants.

sek-1 also acts in acetylcholine neurons to modulate the loopy waveform of both

activated Gq and activated *nca-1* mutants, and the size of activated *nca-1* mutants.

However, our data attempting to narrow down the site of action of *sek-1* suggest that it

may act in both head acetylcholine neurons and acetylcholine motorneurons, and that

the waveform is probably controlled by at least partially distinct neurons from those that

control locomotion rate. Further work will be required to identify the specific neurons

where Gq, NCA-1 and the SEK-1 pathway act to modulate locomotion rate and

waveform, and determine whether they all act together in the same cell.

Acknowledgements

We thank Dennis Kim and Chiou-Fen Chuang for strains, Pin-An Chen and Erik

Jorgensen for the *nca-1(gf)* mutant *ox352*, Chris Johnson for the fine mapping of *yak42*,

Jordan Hoyt for help with Galaxy to analyze WGS data, and Dana Miller for providing

access to her microscope camera. Some strains were provided by the CGC, which is

funded by NIH Office of Research Infrastructure Programs (P40 OD010440). J.M.H was

supported in part by Public Health Service, National Research Service Award

T32GM007270, from the National Institute of General Medical Sciences. M.A. is an

Ellison Medical Foundation New Scholar. This work was supported by NIH grant R00

MH082109 to M.A.

Figure Legends

Figure 1. *sek-1* acts downstream of Gαq to modulate locomotion behavior

(A) Gene structure of *sek-1*. White boxes depict the 5' and 3' untranslated regions,

black boxes depict exons, and lines show introns. The positions of the *yak42* and *km4*

deletions are shown. *yak42* is a 3713 bp deletion that extends to 1926 bp upstream of

the start codon. Drawn with Exon-Intron Graphic Maker

(http://www.wormweb.org/exonintron). Scale bar is 100 bp.

(B-D) *sek-1(yak42)* and *sek-1(km4)* suppress the loopy waveform of the activated Gq

mutant *egl-30(tg26). unc-82(e1220)* does not suppress *egl-30(tg26).* (B) Photos of first-

day adult worms. WT: wild type. (C) Quantification of the waveform phenotype. ***,

p<0.001; ns, p>0.05 compared to *egl-30(tg26)*. Error bars = SEM, n=5. (D) Photos of

worm tracks.

(E) The activated Gq mutant *egl-30(tg26)* is hyperactive and is suppressed by *sek-*

1(yak42) and *sek-1(km4)*. ***, p<0.001, error bars = SEM, n=20.

(F) *sek-1* mutant worms have slow locomotion. ***, p<0.001 compared to wild-type.

Error bars = SEM, n=20.

(G) *sek-1(km4)* suppresses the hyperactive locomotion of the activated Gq mutant *egl-*

30(js126). ***, p<0.001, error bars = SEM, n=20.

(H) *sek-1(km4)* suppresses the loopy waveform of the activated Gq mutant *egl-*

30(js126). ***, p<0.001, error bars = SEM, n=5.

Figure 2. *sek-1* acts in mature acetylcholine neurons to modulate locomotion

(A) *sek-1* acts in neurons to modulate locomotion rate. The *sek-1* wild-type cDNA driven

by the *unc-119* pan-neuronal promoter *[unc-119p::sek-1(+)]* rescues the slow

locomotion phenotype of *sek-1(km4)* worms. ***, p< 0.001, error bars = SEM, n=20.

(B) *sek-1* acts in acetylcholine neurons to modulate locomotion rate. *sek-1* WT cDNA

driven by the *unc-17* acetylcholine neuron promoter *[unc-17p::sek-1(+)]* rescues the

slow locomotion phenotype of *sek-1(km4)* worms but *sek-1* expression in GABA

neurons using the *unc-47* promoter *[unc-47p::sek-1(+)]* does not. ***, p< 0.001, error

bars = SEM, n=20.

(C-D) *sek-1* acts in acetylcholine neurons to modulate the loopy waveform of *egl-*

30(tg26). *egl-30(tg26) sek-1(km4)* worms expressing *unc-119p::sek-1(+)* or *unc-*

17p::sek-1(+) are loopy like *egl-30(tg26)*, but *egl-30(tg26) sek-1(km4)* worms expressing

unc-47p::sek-1(+) are similar to *egl-30(tg26) sek-1*. (C) Photos of worms. (D)

Quantification of waveform phenotype. ***, p<0.001; ns, p>0.05. Error bars = SEM, n=5.

(E) *sek-1* acts in acetylcholine neurons to modulate the locomotion rate of *egl-30(tg26)*.

egl-30(tg26) sek-1(km4) worms expressing *unc-119p::sek-1(+)* or *unc-17p::sek-1(+)*

have an increased locomotion rate compared to *egl-30(tg26) sek-1*, but *egl-30(tg26)*

sek-1(km4) worms expressing *unc-47p::sek-1(+)* are similar to *egl-30(tg26) sek-1*. ***,

p< 0.001, **, p<0.01; ns, p>0.05. Error bars = SEM, n=17-20.

(F) *sek-1* acts in mature neurons to modulate locomotion rate. Heat-shock induced

expression of *sek-1* in adults (*hsp-16.2p::sek-1(+)*) rescues the slow locomotion

phenotype of *sek-1(km4)*. ***, p< 0.001, error bars = SEM, n=20.

Figure 3. The p38 MAPK pathway modulates locomotion downstream of *egl-30*

(A) *tir-1(tm3036)* mutant animals have slow locomotion. ***, p<0.001, error bars = SEM,

n=20.

(B) *tir-1(tm3036)* suppresses *egl-30(tg26)*. *egl-30(tg26) tir-1* animals move more slowly

than the hyperactive *egl-30(tg26)* animals. ***, p< 0.001, error bars = SEM, n=20.

(C) *nsy-1(ok593)* mutant animals have slow locomotion. ***, p<0.001, error bars = SEM,

n=20.

(D) *nsy-1(ok593)* suppresses *egl-30(tg26)*. *egl-30(tg26) nsy-1* animals move more

slowly than the hyperactive *egl-30(tg26)* animals. ***, p< 0.001, error bars = SEM, n=20.

(E) *pmk-2, pmk-2 pmk-1,* and *pmk-3* mutant animals have slow locomotion. ***, p<

0.001; *, p<0.05, compared to WT. Error bars = SEM, n=20.

(F) A *pmk-2 pmk-1* double mutant suppresses the hyperactivity of *egl-30(tg26)*. ***, p<

0.001 compared to *egl-30(tg26)*. Error bars = SEM, n=20.

(G, H) *tir-1(tm3036)*, *nsy-1(ok593)*, and the *pmk-2 pmk-1* double mutant suppress the

loopy waveform of *egl-30(tg26)*. *egl-30(tg26)* animals with mutations in either *pmk-1,*

pmk-2, or *pmk-3* are still loopy. (G) Worm photos. (H) Quantification. ***, p<0.001

compared to *egl-30(tg26)*. Error bars = SEM, n=5.

Figure 4. *sek-1* acts in the same genetic pathway as *unc-73*

(A) *sek-1* does not act in the same genetic pathway as *egl-8.* The *sek-1(yak42)*

mutation enhances the slow locomotion of the *egl-8(sa47)* mutant. ***, p< 0.001, error

bars = SEM, n=20.

(B) *sek-1* does not act in the same genetic pathway as *rund-1*. The *sek-1(yak42)*

mutation enhances the slow locomotion of the *rund-1(tm3622)* mutant. ***, p< 0.001,

error bars = SEM, n=20.

(C) *sek-1* may act in the same genetic pathway as *unc-73*. The *sek-1(yak42)* mutation

does not enhance the slow locomotion phenotype of the *unc-73(ox317)* mutant. ns,

p>0.05, error bars = SEM, n=20.

(D-E) *sek-1(km4)* does not suppress the loopy waveform of *nzIs29 rho-1(gf)* animals.

(D) Worm photos. (E) Quantification. ns, p>0.05, error bars = SEM, n=5.

(F) *sek-1(km4)* does not suppress the slow locomotion of *rho-1(gf)* animals. ***, p<

0.001, error bars = SEM, n=20.

Figure 5. *sek-1* and *nsy-1* weakly suppress *nca-1(gf)*

(A) *nca-1(gf)* mutants are small, loopy, and uncoordinated. The phenotypes of *nca-*

1(ox352) animals are partially suppressed by *sek-1(km4)* and *nsy-1(ok593)*. Photos of

first-day adults.

(B) *sek-1(km4)* and *nsy-1(ok593)* suppress the loopy waveform of *nca-1(gf)*. ***,

p<0.001; **, p<0.01. Error bars = SEM, n=5.

(C) *sek-1* and *nsy-1* suppress *nca-1(gf)* locomotion. *nca-1(gf)* animals travel a small

distance from the center of the plate in the radial locomotion assay. *nca-1(gf) nsy-*

1(ok593) and *nca-1(gf) sek-1(km4)* worms move further than *nca-1(gf)* worms. **,

p<0.01; *, p<0.05. Error bars = SEM, n=30.

(D) *sek-1* acts in acetylcholine neurons to modulate the size and loopy waveform of

nca-1(gf). *nca-1(ox352) sek-1(km4)* animals expressing *sek-1* in all neurons (*unc-*

119p::sek-1(+)) or in acetylcholine neurons (*unc-17p::sek-1(+)*) are loopy and small like

nca-1(gf), but *nca-1(ox352) sek-1(km4)* animals expressing *sek-1* in GABA neurons

(*unc-47p::sek-1(+)*) resemble *nca-1(gf) sek-1*. White arrowheads depict food piles

created by *nca-1(gf) sek-1(km4)* animals due to their uncoordinated locomotion. Such

food piles are not made by *nca-1(gf)* animals.

(E) *sek-1* acts in acetylcholine motorneurons to modulate the loopy waveform of *nca-*

1(gf). *nca-1(ox352) sek-1(km4)* worms expressing *sek-1* in all neurons (*unc-119p::sek-*

1(+)), acetylcholine neurons (*unc-17p::sek-1(+)*), or acetylcholine motorneurons (*unc-*

17βp::sek-1(+)) are loopy like *nca-1(gf)*, but *nca-1(ox352) sek-1(km4)* worms expressing

sek-1 in GABA neurons (*unc-47p::sek-1(+)*) or head acetylcholine neurons (*unc-*

17Hp::sek-1(+)) are similar to *nca-1(gf) sek-1*. ***, $p < 0.001$, **, $p < 0.01$; ns, $p > 0.05$.

Error bars = SEM, n=5.

References

Afgan, E., D. Baker, M. van den Beek, D. Blankenberg, D. Bouvier *et al.*, 2016 The

573   Galaxy platform for accessible, reproducible and collaborative biomedical

574   analyses: 2016 update. Nucleic Acids Res. 44: W3–W10.

575   Ailion, M., M. Hannemann, S. Dalton, A. Pappas, S. Watanabe *et al.*, 2014 Two Rab2

576   interactors regulate dense-core vesicle maturation. Neuron 82: 167–180.

577   Andrusiak, M. G., and Y. Jin, 2016 Context Specificity of Stress-activated Mitogen-

578   activated Protein (MAP) Kinase Signaling: The Story as Told by Caenorhabditis

579   elegans. J. Biol. Chem. 291: 7796–7804.

580   Bastiani, C. A., S. Gharib, M. I. Simon, and P. W. Sternberg, 2003 Caenorhabditis

581   elegans Gαq Regulates Egg-Laying Behavior via a PLCβ-Independent and

582   Serotonin-Dependent Signaling Pathway and Likely Functions Both in the

583   Nervous System and in Muscle. Genetics 165: 1805–1822.

584   Bence, K., W. Ma, T. Kozasa, and X.-Y. Huang, 1997 Direct stimulation of Bruton's

585   tyrosine kinase by Gq-protein α-subunit. Nature 389: 296–299.

Bolshakov, V. Y., L. Carboni, M. H. Cobb, S. A. Siegelbaum, and F. Belardetti, 2000

Dual MAP kinase pathways mediate opposing forms of long-term plasticity at

CA3-CA1 synapses. Nat. Neurosci. 3: 1107–1112.

Brenner, S., 1974 The genetics of Caenorhabditis elegans. Genetics 77: 71–94.

Chang, C., Y.-W. Hsieh, B. J. Lesch, C. I. Bargmann, and C.-F. Chuang, 2011

Microtubule-based localization of a synaptic calcium-signaling complex is

required for left-right neuronal asymmetry in C. elegans. Development 138:

3509–3518.

Chuang, C.-F., and C. I. Bargmann, 2005 A Toll-interleukin 1 repeat protein at the

synapse specifies asymmetric odorant receptor expression via ASK1 MAPKKK

signaling. Genes Dev. 19: 270–281.

Cingolani, P., A. Platts, L. L. Wang, M. Coon, T. Nguyen *et al.*, 2012 A program for

annotating and predicting the effects of single nucleotide polymorphisms, SnpEff:

SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. Fly

6: 80–92.

Couillault, C., N. Pujol, J. Reboul, L. Sabatier, J.-F. Guichou *et al.*, 2004 TLR-

independent control of innate immunity in Caenorhabditis elegans by the TIR

domain adaptor protein TIR-1, an ortholog of human SARM. Nat. Immunol. 5:

488–494.

DePristo, M. A., E. Banks, R. Poplin, K. V. Garimella, J. R. Maguire *et al.*, 2011 A

framework for variation discovery and genotyping using next-generation DNA

sequencing data. Nat. Genet. 43: 491–498.

Doi, M., and K. Iwasaki, 2002 Regulation of retrograde signaling at neuromuscular

junctions by the novel C2 domain protein AEX-1. Neuron 33: 249–259.

Fay, D., 2006 Genetic mapping and manipulation: Chapter 7-Making compound

mutants. *WormBook*, ed. The *C. elegans* Research Community, WormBook,

doi/10.1895/wormbook.1.96.2, http://www.wormbook.org

Frøkjær-Jensen, C., M. W. Davis, M. Ailion, and E. M. Jorgensen, 2012 Improved

Mos1-mediated transgenesis in C. elegans. Nat. Methods 9: 117–118.

Frokjaer-Jensen, C., M. W. Davis, M. Sarov, J. Taylor, S. Flibotte *et al.*, 2014 Random

and targeted transgene insertion in C. elegans using a modified Mosl transposon.

Nat. Methods 11: 529–534.

Gao, S., L. Xie, T. Kawano, M. D. Po, J. K. Pirri *et al.*, 2015 The NCA sodium leak

channel is required for persistent motor circuit activity that sustains locomotion.

Nat. Commun. 6: 6323.

García-Hoz, C., G. Sánchez-Fernández, M. T. Díaz-Meco, J. Moscat, F. Mayor *et al.*,

2010 Gαq Acts as an Adaptor Protein in Protein Kinase Cζ (PKCζ)-mediated

ERK5 Activation by G Protein-coupled Receptors (GPCR). J. Biol. Chem. 285:

13480–13489.

Hajdu-Cronin, Y. M., W. J. Chen, G. Patikoglou, M. R. Koelle, and P. W. Sternberg,

1999 Antagonism between Goα and Gqα in Caenorhabditis elegans: the RGS

protein EAT-16 is necessary for Goα signaling and regulates Gqα activity. Genes

Dev. 13: 1780–1793.

Hawasli, A. H., O. Saifee, C. Liu, M. L. Nonet, and C. M. Crowder, 2004 Resistance to

volatile anesthetics by mutations enhancing excitatory neurotransmitter release

in Caenorhabditis elegans. Genetics 168: 831–843.

Hoppe, P. E., J. Chau, K. A. Flanagan, A. R. Reedy, and L. A. Schriefer, 2010

Caenorhabditis elegans unc-82 Encodes a Serine/Threonine Kinase Important

for Myosin Filament Organization in Muscle During Growth. Genetics 184: 79–90.

Huang, C.-C., J.-L. You, M.-Y. Wu, and K.-S. Hsu, 2004 Rap1-induced p38 mitogen-

activated protein kinase activation facilitates AMPA receptor trafficking via the

GDI.Rab5 complex. Potential role in (S)-3,5-dihydroxyphenylglycene-induced

long term depression. J. Biol. Chem. 279: 12286–12292.

Hudmon, A., J.-S. Choi, L. Tyrrell, J. A. Black, A. M. Rush *et al.*, 2008 Phosphorylation

of Sodium Channel Nav1.8 by p38 Mitogen-Activated Protein Kinase Increases

Current Density in Dorsal Root Ganglion Neurons. J. Neurosci. 28: 3190–3201.

Jantsch-Plunger, V., P. Gönczy, A. Romano, H. Schnabel, D. Hamill *et al.*, 2000 CYK-4:

A Rho family gtpase activating protein (GAP) required for central spindle

formation and cytokinesis. J. Cell Biol. 149: 1391–1404.

Kawasaki, M., 1999 A Caenorhabditis elegans JNK signal transduction pathway

regulates coordinated movement via type-D GABAergic motor neurons. EMBO J.

18: 3604–3615.

Kim, D. H., R. Feinbaum, G. Alloing, F. E. Emerson, D. A. Garsin *et al.*, 2002 A

Conserved p38 MAP Kinase Pathway in Caenorhabditis elegans Innate

Immunity. Science 297: 623–626.

Kim, D. H., N. T. Liberati, T. Mizuno, H. Inoue, N. Hisamoto *et al.*, 2004 Integration of

Caenorhabditis elegans MAPK pathways mediating immunity and stress

resistance by MEK-1 MAPK kinase and VHP-1 MAPK phosphatase. Proc. Natl.

Acad. Sci. U. S. A. 101: 10990–10994.

Koelle, M. R., 2016 Neurotransmitter signaling through heterotrimeric G proteins:

insights from studies in C. elegans. WormBook, doi:10.1895/wormbook.1.75.2.

Kyriakis, J. M., and J. Avruch, 2012 Mammalian MAPK signal transduction pathways

activated by stress and inflammation: a 10-year update. Physiol. Rev. 92: 689–

737.

Lackner, M. R., S. J. Nurrish, and J. M. Kaplan, 1999 Facilitation of synaptic

transmission by EGL-30 Gqalpha and EGL-8 PLCbeta: DAG binding to UNC-13

is required to stimulate acetylcholine release. Neuron 24: 335–346.

Lu, B., Y. Su, S. Das, J. Liu, J. Xia *et al.*, 2007 The Neuronal Channel NALCN

Contributes Resting Sodium Permeability and Is Required for Normal Respiratory

Rhythm. Cell 129: 371–383.

McMullan, R., E. Hiley, P. Morrison, and S. J. Nurrish, 2006 Rho is a presynaptic

activator of neurotransmitter release at pre-existing synapses in C. elegans.

Genes Dev. 20: 65–76.

Mello, C. C., J. M. Kramer, D. Stinchcomb, and V. Ambros, 1991 Efficient gene transfer

in C.elegans: extrachromosomal maintenance and integration of transforming

sequences. EMBO J. 10: 3959–3970.

Miller, K. G., M. D. Emerson, and J. B. Rand, 1999 Goalpha and diacylglycerol kinase

negatively regulate the Gqalpha pathway in C. elegans. Neuron 24: 323–333.

Nash, H. A., R. L. Scott, B. C. Lear, and R. Allada, 2002 An unusual cation channel

mediates photic control of locomotion in Drosophila. Curr. Biol. 12: 2152–2158.

Natochin, M., and N. O. Artemyev, 2003 A point mutation uncouples transducin-alpha

from the photoreceptor RGS and effector proteins. J. Neurochem. 87: 1262–

1271.

Nurrish, S., L. Ségalat, and J. M. Kaplan, 1999 Serotonin inhibition of synaptic

transmission: Galpha(0) decreases the abundance of UNC-13 at release sites.

Neuron 24: 231–242.

Pagano, D. J., E. R. Kingston, and D. H. Kim, 2015 Tissue Expression Pattern of PMK-2

p38 MAPK Is Established by the miR-58 Family in C. elegans. PLoS Genet 11:

e1004997.

Park, E. C., and C. Rongo, 2016 The p38 MAP kinase pathway modulates the hypoxia

response and glutamate receptor trafficking in aging neurons. eLife 5: e12010.

Plowman, G. D., S. Sudarsanam, J. Bingham, D. Whyte, and T. Hunter, 1999 The

protein kinases of Caenorhabditis elegans: a model for signal transduction in

multicellular organisms. Proc. Natl. Acad. Sci. U. S. A. 96: 13603–13610.

Rhee, S. G., 2001 Regulation of Phosphoinositide-Specific Phospholipase C. Annu.

Rev. Biochem. 70: 281–312.

Rush, A. M., J. Wu, M. J. Rowan, and R. Anwyl, 2002 Group I metabotropic glutamate

receptor (mGluR)-dependent long-term depression mediated via p38 mitogen-

activated protein kinase is inhibited by previous high-frequency stimulation and

activation of mGluRs and protein kinase C in the rat dentate gyrus in vitro. J.

Neurosci. 22: 6121–6128.

Sagasti, A., N. Hisamoto, J. Hyodo, M. Tanaka-Hino, K. Matsumoto *et al.*, 2001 The

CaMKII UNC-43 Activates the MAPKKK NSY-1 to Execute a Lateral Signaling

Decision Required for Asymmetric Olfactory Neuron Fates. Cell 105: 221–232.

Sánchez-Fernández, G., S. Cabezudo, C. García-Hoz, C. Benincá, A. M. Aragay *et al.*,

2014 Gαq signalling: The new and the old. Cell. Signal. 26: 833–848.

Shi, Y., C. Abe, B. B. Holloway, S. Shu, N. N. Kumar *et al.*, 2016 Nalcn Is a "Leak"

Sodium Channel That Regulates Excitability of Brainstem Chemosensory

Neurons and Breathing. J. Neurosci. 36: 8174–8187.

Shivers, R. P., T. Kooistra, S. W. Chu, D. J. Pagano, and D. H. Kim, 2009 Tissue-

specific activities of an immune signaling module regulate physiological

responses to pathogenic and nutritional bacteria in C. elegans. Cell Host Microbe

6: 321–330.

Shivers, R. P., D. J. Pagano, T. Kooistra, C. E. Richardson, K. C. Reddy *et al.*, 2010

Phosphorylation of the conserved transcription factor ATF-7 by PMK-1 p38

MAPK regulates innate immunity in Caenorhabditis elegans. PLoS Genet. 6:

e1000892.

Tanaka-Hino, M., A. Sagasti, N. Hisamoto, M. Kawasaki, S. Nakano *et al.*, 2002 SEK-1

MAPKK mediates Ca2+ signaling to determine neuronal asymmetric

development in Caenorhabditis elegans. EMBO Rep. 3: 56–62.

Topalidou, I., J. Cattin-Ortolá, A. L. Pappas, K. Cooper, G. E. Merrihew *et al.*, 2016 The

EARP Complex and Its Interactor EIPR-1 Are Required for Cargo Sorting to

Dense-Core Vesicles. PLOS Genet 12: e1006074.

Topalidou, I., P.-A. Chen, K. Cooper, S. Watanabe, E. M. Jorgensen *et al.*, 2017 The

NCA-1 and NCA-2 Ion Channels Function Downstream of Gq and Rho To

Regulate Locomotion in Caenorhabditis elegans. Genetics 206: 265–282.

Vaqué, J. P., R. T. Dorsam, X. Feng, R. Iglesias-Bartolome, D. J. Forsthoefel *et al.*,

2013 A genome-wide RNAi screen reveals a Trio-regulated Rho GTPase circuitry

transducing mitogenic signals initiated by G protein-coupled receptors. Mol. Cell

49: 94–108.

Vashlishan, A. B., J. M. Madison, M. Dybbs, J. Bai, D. Sieburth *et al.*, 2008 An RNAi

Screen Identifies Genes that Regulate GABA Synapses. Neuron 58: 346–361.

Williams, S. L., S. Lutz, N. K. Charlie, C. Vettel, M. Ailion *et al.*, 2007 Trio's Rho-specific

GEF domain is the missing Galpha q effector in C. elegans. Genes Dev. 21:

2731–2746.

Wittmack, E. K., A. M. Rush, A. Hudmon, S. G. Waxman, and S. D. Dib-Hajj, 2005

Voltage-Gated Sodium Channel Nav1.6 Is Modulated by p38 Mitogen-Activated

Protein Kinase. J. Neurosci. 25: 6621–6630.

Zhang, R., X. He, W. Liu, M. Lu, J.-T. Hsieh *et al.*, 2003 AIP1 mediates TNF-α–induced

ASK1 activation by facilitating dissociation of ASK1 from its inhibitor 14-3-3. J.

Clin. Invest. 111: 1933–1943.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

WT

nca-1(gf)

nca-1(gf); sek-1(km4)

nca-1(gf); nsy-1

B

period/amplitude

WT sek-1(km4) nca-1(gf) nca-1(gf); sek-1(km4) nca-1(gf); nsy-1

** ***

C

radial distance at 20hr (mm)

WT nca-1(gf) nca-1(gf); sek-1(km4) nca-1(gf); nsy-1

* **

D

WT nca-1(gf) nca-1(gf); sek-1(km4)

nca-1(gf); sek-1(km4); unc-119p::sek-1(+) nca-1(gf); sek-1(km4); unc-17p::sek-1(+) nca-1(gf); sek-1(km4); unc-47p::sek-1(+)

E

period/amplitude

WT nca-1(gf) nca-1(gf) ;sek-1 unc-119p::sek-1 unc-17p::sek-1 unc-47p::sek-1 unc-17Hp::sek-1 unc-17βp::sek-1

*** *** ** ns ns ***

nca-1(gf); sek-1(km4)

Supplementary Information

Table S1. Strain List

Strain	Genotype
AU1	*sek-1(ag1)* X
BS3383	*pmk-3(ok169)* IV
CX3695	*kyIs140[str-2p::gfp, lin-15(+)]* I
CX5959	*kyIs140[str-2p::gfp, lin-15(+)]* I; *tir-1(ky648gf)* III
EG317	*unc-73(ox317)* I
EG1000	*dpy-5(e61)* I; *rol-6(e187)* II; *lon-1(e1820)* III
EG1020	*bli-6(sc16)* IV; *dpy-11(e224)* V; *lon-2(e678)* X
EG3745	*eat-16(tm775)* I ; *him-5(e1490)* V
EG4782	*nzIs29[unc-17p::rho-1(G14V), unc-122::gfp]* II
EG5505	*rund-1(tm3622)* X
EG7989	*unc-119(ed3)* III; *oxTi668[eft-3p::TdTomato::H2B, Cb-unc-119(+)]* X
IG685	*tir-1(tm3036)* III
JN147	*gap-2(tm748)* X
JT47	*egl-8(sa47)* V
JT366	*vhp-1(sa366)* II
JT734	*goa-1(sa734)* I
KU2	*jkk-1(km2)* X
KU4	*sek-1(km4)* X
KU25	*pmk-1(km25)* IV
N2	Bristol wild isolate, standard lab wild-type
NM1380	*egl-30(js126gf)* I

VC8	*jnk-1(gk7)* IV
VC390	*nsy-1(ok593)* IV
XZ42	*sek-1(yak42)* X
XZ1233	*egl-30(tg26)* I*; sek-1(yak42)* X
XZ1151	*egl-30(tg26)* I
XZ1566	*egl-8(sa47)* V*; sek-1(yak42)* X
XZ1567	*unc-73(ox317)* I*; sek-1(yak42)* X
XZ1574	*rund-1(tm3622) sek-1(yak42)* X
XZ1575	*egl-30(tg26)* I*; sek-1(km4)* X
XZ1588	*egl-30(tg26)* I*; nsy-1(ok593)* IV
XZ1589	*egl-30(tg26)* I*; sek-1(km4)* X*; qdEx8[unc-119p::sek-1::GFP, myo-2p::mStrawberry::unc-54-3'UTR]*
XZ1590	*egl-30(tg26)* I *; jkk-1(km2)* X
XZ1593	*egl-30(tg26)* I*; pmk-1(km25)* IV
XZ1597	*egl-30(tg26)* I *; jnk-1(gk7)* IV
XZ1642	*sek-1(km4)* X*; yakEx72[unc-17p::sek-1::tbb-2utr-operon-GFP::H2B::cye-1utr, myo-2p::mCherry]*
XZ1643	*sek-1(km4)* X*; yakEx73[unc-47p::sek-1::tbb-2utr-operon-GFP::H2B::cye-1utr, myo-2p::mCherry]*
XZ1717	*nzIs29[unc-17p::rho-1(G14V) unc-122::gfp]* II*; sek-1(km4)* X
XZ1720	*sek-1(km4)* X *; yakEx82[unc-17Hp::sek-1:: tbb-2utr-operon-GFP::H2B::cye-1utr, myo-2p::mCherry]*
XZ1721	*sek-1(km4)* X *; yakEx83[unc-17βp::sek-1::tbb-2utr-operon-GFP::H2B::cye-1utr, myo-2p::mCherry]*
XZ1770	*egl-30(tg26)* I*; pmk-2(qd279 qd171) pmk-1(km25)* IV

XZ1771	*egl-30(tg26) I; pmk-2(qd287) IV*
XZ1772	*egl-30(tg26) I; pmk-3(ok169) IV*
XZ1815	*egl-30(tg26) I; tir-1(tm3036) III*
XZ1816	*nca-1(ox352) IV; sek-1(km4) X; qdEx8[unc-119p::sek-1::GFP::unc-54-3' UTR, myo-2p::mStrawberry::unc-54-3'UTR]*
XZ1820	*nca-1(ox352) IV ; sek-1(km4) X ; yakEx83[unc-17βp::sek-1::tbb-2utr-operon-GFP::H2B::cye-1utr, myo-2p::mCherry]*
XZ1830	*egl-30(tg26) I ; sek-1(km4) X ; yakEx83[unc-17βp::sek-1::tbb-2utr-operon-GFP::H2B::cye-1utr, myo-2p::mCherry]*
XZ1834	*egl-30(tg26) I; sek-1(km4) X; yakEx72[unc-17p::sek-1::tbb-2utr-operon-GFP::H2B::cye-1utr, myo-2p::mCherry]*
XZ1835	*nca-1(ox352) IV; sek-1(km4) X; yakEx72[unc-17p::sek-1::tbb-2utr-operon-GFP::H2B::cye-1utr, myo-2p::mCherry]*
XZ1861	*nca-1(ox352) IV; sek-1(km4) X; yakEx73[unc-47p::sek-1::tbb-2utr-operon-GFP::H2B::cye-1utr, myo-2p::mCherry]*
XZ1862	*nca-1(ox352) IV ; sek-1(km4) X ; yakEx82[unc-17Hp::sek-1::tbb-2utr-operon-GFP::H2B::cye-1utr, myo-2p::mCherry]*
XZ1863	*egl-30(tg26) I; sek-1(km4) X; yakEx73[unc-47p::sek-1::tbb-2utr-operon-GFP::H2B::cye-1utr, myo-2p::mCherry]*
XZ1872	*jnk-1(gk7) I; sek-1(km4) X*
XZ1873	*pmk-2(qd279 qd171) pmk-1(km25) IV; jkk-1(km2) X*
XZ1879	*egl-30(tg26) I ; sek-1(km4) X ; yakEx82[unc-17Hp:: sek-1::tbb-2utr-operon-GFP::H2B::cye-1utr, myo-2p::mCherry]*
XZ1902	*egl-30(tg26) I ; unc-82(e1220) IV*
XZ1937	*sek-1(km4) X; yakEx121[hsp-16.2p::sek-1::tbb-2-3' UTR::gld-1 operon*

	linker::gfp::h2b, myo-2p::mCherry]
XZ1938	*egl-30(tg26)* I ; *agIs219[T24B8.5p::GFP::unc-54-3'UTR + ttx-3p::GFP::unc-54-3'UTR]* III *atf-7(qd22 qd130)* III
XZ1939	*goa-1(sa734)* I; *sek-1(km4)* X
XZ1942	*tir-1(ky648gf)* III
XZ2054	*eat-16(tm775)* I ; *sek-1(km4)* X
XZ2062	*egl-30(js126gf)* I ; *sek-1(km4)* X
ZD202	*sek-1(km4)* X; *qdEx8[unc-119p::sek-1::GFP::unc-54-3' UTR + myo-2p::mStrawberry::unc-54-3'UTR]*
ZD318	*agIs29 atf-7(qd22 qd130)* III
ZD442	*agIs29 atf-7(qd22)* III
ZD934	*pmk-2(qd279 qd171) pmk-1(km25)* IV
ZD1020	*pmk-2(qd287)* IV

Table S2. Plasmids and Primers

Gateway entry clones

Plasmid	Details
pADA180	*unc-17Hp* [4-1]
pJH21	*sek-1* cDNA [1-2]
pCFJ150	pDEST5605[4-3]
pCFJ326	*tbb-2utr-operon-GFP::H2B::cye-1utr* [2-3]
pMA23	*unc-17βp* [4-1]
pMH522	*unc-47p* [4-1]
pGH1	*unc-17p* [4-1]
pCM1.56	*hsp-16.2p* [4-1]

Gateway Expression Constructs

Plasmid	Details	Used to make
pJH23	*unc-17p::sek-1::tbb-2utr-operon-GFP::H2B::cye-1utr*	*yakEx72*
pJH24	*unc-47p::sek-1::tbb-2utr-operon-GFP::H2B::cye-1utr*	*yakEx73*
pJH28	*unc-17Hp::sek-1::tbb-2utr-operon-GFP::H2B::cye-1utr*	*yakEx82*
pJH29	*unc-17βp::sek-1::tbb-2utr-operon-GFP::H2B::cye-1utr*	*yakEx83*
pJH46	*hsp-16.2p::sek-1::tbb-2utr-operon-GFP::H2B::cye-1utr*	*yakEx121*

Primers

oJH114	GGGGACAAGTTTGTACAAAAAAGCAGGCTcaATGGAGCGAAAAGGACGTG	F to clone *sek-1* cDNA into [1-2]
oJH115	GGGGACCACTTTGTACAAGAAAGCTGGGTgTCATCGTCGCCAAACAGTG	R to clone *sek-1* cDNA into [1-2]

Table S3. Statistical Tests

Figure	Test	p value
1C	One-way ANOVA and Bonferroni's Multiple Comparison Test WT vs *sek-1(km4)* (ns)	< 0.001

	WT vs *egl-30(tg26)* (p<0.001) *egl-30(tg26)* vs *egl-30(tg26); sek-1(km4)* (p<0.001) *egl-30(tg26)* vs *egl-30(tg26); sek-1(yak42)* (p<0.001) *egl-30(tg26)* vs *egl-30(tg26); unc-82(e1220)* (ns)	
1E	One-way ANOVA and Bonferroni's Multiple Comparison Test WT vs *egl-30(tg26)* (p<0.001) *egl-30(tg26)* vs *egl-30(tg26); sek-1(yak42)* (p<0.001) *egl-30(tg26)* vs *egl-30(tg26); sek-1(km4)* (p<0.001)	< 0.001
1F	One-way ANOVA and Dunnett's Multiple Comparison Test WT vs *sek-1(yak42)* (p<0.001) WT vs *sek-1(km4)* (p<0.001)	< 0.001
1G	One-way ANOVA and Bonferroni's Multiple Comparison Test WT vs *egl-30(js126)* (p<0.001) WT vs *sek-1(km4)* (p<0.001) *egl-30(js126)* vs *egl-30(js126); sek-1(km4)* (p<0.001)	< 0.001
1H	One-way ANOVA and Bonferroni's Multiple Comparison Test WT vs *egl-30(js126)* (p<0.001) *egl-30(js126)* vs *egl-30(js126); sek-1(km4)* (p<0.001)	<0.001
2A	One-way ANOVA and Bonferroni's Multiple Comparison Test *sek-1(km4)* vs *sek-1(km4); qdEx8[unc-119::sek-1(+)]* (p<0.001)	< 0.001
2B	One-way ANOVA and Bonferroni's Multiple Comparison Test *sek-1(km4)* vs *sek-1(km4); yakEx72[unc-17p::sek-1(+)]* (p<0.001) *sek-1(km4)* vs *sek-1(km4); yakEx73[unc-47p::sek-1(+)]* (ns)	<0.001
2D	One-way ANOVA and Bonferroni's Multiple Comparison Test WT vs *egl-30(tg26)* (p<0.001)	<0.001

	egl-30(tg26); sek-1(km4) vs *egl-30(tg26); sek-1(km4); qdEx8[unc-119p::sek-1(+)]* (p<0.001) *egl-30(tg26); sek-1(km4)* vs *egl-30(tg26); sek-1(km4); yakEx72[unc-17p::sek-1(+)]* (p<0.001) *egl-30(tg26); sek-1(km4)* vs *egl-30(tg26); sek-1(km4); yakEx73[unc-47p::sek-1(+)]* (ns)	
2E	Kruskal-Wallis Test and Dunn's Multiple Comparison Test *egl-30(tg26); sek-1(km4)* vs *egl-30(tg26); sek-1(km4); qdEx8[unc-119p::sek-1(+)]* (p<0.001) *egl-30(tg26); sek-1(km4)* vs *egl-30(tg26); sek-1(km4); yakEx72[unc-17p::sek-1(+)]* (p<0.01) *egl-30(tg26); sek-1(km4)* vs *egl-30(tg26); sek-1(km4); yakEx73[unc-47p::sek-1(+)]* (ns)	< 0.001
2F	One-way ANOVA and Bonferroni's Multiple Comparison Test *sek-1(km4)* vs *sek-1(km4); yakEx121[hsp-16.2p::sek-1(+)]* (p<0.001)	< 0.001
3A	Unpaired t test, two-tailed	< 0.001
3B	One-way ANOVA and Bonferroni's Multiple Comparison Test *egl-30(tg26)* vs *egl-30(tg26); tir-1(tm3036)* (p<0.001)	< 0.001
3C	Unpaired t test, two-tailed	< 0.001
3D	One-way ANOVA and Bonferroni's Multiple Comparison Test *egl-30(tg26)* vs *egl-30(tg26); nsy-1(ok593)* (p<0.001)	< 0.001
3E	One-way ANOVA and Dunnett's Multiple Comparison Test	< 0.001

	WT vs *pmk-1(km25)* (ns)	
	WT vs *pmk-2(qd287)* (p<0.05)	
	WT vs *pmk-2(qd279 qd171) pmk-1 (km25)* (p<0.001)	
	WT vs *pmk-3(ok169)* (p<0.001)	
3F	One-way ANOVA and Dunnett's Multiple Comparison Test *egl-30(tg26)* vs *egl-30(tg26); pmk-1(km25)* (p<0.001) *egl-30(tg26)* vs *egl-30(tg26); pmk-2(qd287)* (p<0.001) *egl-30(tg26)* vs *egl-30(tg26); pmk-2(qd279 qd171) pmk-1 (km25)* (p<0.001) *egl-30(tg26)* vs *egl-30(tg26); pmk-3(ok169)* (p<0.001)	< 0.001
3H	One-way ANOVA and Bonferroni's Multiple Comparison Test *egl-30(tg26)* vs *egl-30(tg26); tir-1(tm3036)* (p<0.001) *egl-30(tg26)* vs *egl-30(tg26); nsy-1(ok593)* (p<0.001) *egl-30(tg26)* vs *egl-30(tg26); sek-1(km4)* (p<0.001) *egl-30(tg26)* vs *egl-30(tg26); pmk-2(qd279 qd171)* *pmk-1(km25)* (p<0.001)	< 0.001
4A	One-way ANOVA and Bonferroni's Multiple Comparison Test *egl-8(sa47)* vs *egl-8(sa47); sek-1(yak42)* (p<0.001)	< 0.001
4B	One-way ANOVA and Bonferroni's Multiple Comparison Test *sek-1(yak42)* vs *rund-1(tm3622); sek-1(yak42)* (p<0.001)	< 0.001
4C	One-way ANOVA and Bonferroni's Multiple Comparison Test *unc-73(ox317)* vs *unc-73(ox317); sek-1(yak42)* (ns)	< 0.001
4E	One-way ANOVA and Bonferroni's Multiple Comparison Test WT vs *nzIs29[unc-17p::rho-1(G14V)]* (p<0.001) WT vs *nzIs29[unc-17p::rho-1(G14V)]; sek-1(km4)* (p<0.001)	< 0.001

	nzIs29[unc-17p::rho-1(G14V)] vs *nzIs29[unc-17p::rho-1(G14V)]; sek-1(km4)* (ns)	
4F	One-way ANOVA and Bonferroni's Multiple Comparison Test *nzIs29[unc-17p::rho-1(G14V)]* vs *nzIs29[unc-17p::rho-1(G14V)]; sek-1(km4)* (p<0.001)	< 0.001
5B	One-way ANOVA and Bonferroni's Multiple Comparison Test WT vs *sek-1(km4)* (ns) WT vs *nca-1(ox352)* (p<0.001) *nca-1(ox352)* vs *nca-1(ox352); sek-1(km4)* (p<0.01) *nca-1(ox352)* vs *nca-1(ox352); nsy-1(ok593)* (p<0.001)	<0.001
5C	One-way ANOVA and Bonferroni's Multiple Comparison Test WT vs *nca-1(ox352)* (p<0.001) *nca-1(ox352)* vs *nca-1(ox352); nsy-1(ok593)* (p<0.01) *nca-1(ox352)* vs *nca-1(ox352); sek-1(km4)* (p<0.05)	< 0.001
5E	One-way ANOVA and Bonferroni's Multiple Comparison Test *nca-1(ox352)* vs *nca-1(ox352); sek-1(km4)* (p<0.001) *nca-1(ox352); sek-1(km4)* vs *nca-1(ox352); sek-1(km4); qdEx8[unc-119p::sek-1(+)]* (p<0.001) *nca-1(ox352); sek-1(km4)* vs *nca-1(ox352); sek-1(km4); yakEx72[unc-17p::sek-1(+)]* (p<0.01) *nca-1(ox352); sek-1(km4)* vs *nca-1(ox352); sek-1(km4); yakEx73[unc-47p::sek-1(+)]* (ns) *nca-1(ox352); sek-1(km4)* vs	< 0.001

	nca-1(ox352); sek-1(km4); yakEx82[unc-17Hp::sek-1(+)] (ns) *nca-1(ox352); sek-1(km4)* vs *nca-1(ox352); sek-1(km4); yakEx83[unc-17βp::sek-1(+)]* (p<0.001)	
S1A	One-way ANOVA and Bonferroni's Multiple Comparison Test WT vs *egl-30(tg26)* (p<0.01) WT vs *sek-1(km4)* (p<0.001) *egl-30(tg26)* vs *egl-30(tg26); sek-1(km4)* (p<0.001) *sek-1(km4)* vs *egl-30(tg26); sek-1(km4)* (p<0.01)	< 0.001
S1B	One-way ANOVA and Newman-Keuls Multiple Comparison Test WT vs *sek-1(yak42)* (p<0.001) WT vs *sek-1(ag1)* (p<0.001) *sek-1(yak42)* vs *sek-1(ag1)* (ns)	< 0.001
S1C	One-way ANOVA and Bonferroni's Multiple Comparison Test WT vs *egl-30(tg26)* (ns) *egl-30(tg26)* vs *egl-30(tg26); unc-82(e1220)* (p<0.001)	< 0.001
S1D	One-way ANOVA and Bonferroni's Multiple Comparison Test WT vs *sek-1(km4)* (p<0.001) *goa-1(sa734)* vs *sek-1(km4)* (p<0.001) *goa-1(sa734)* vs *goa-1(sa734); sek-1(km4)* (p<0.001)	< 0.001
S1E	One-way ANOVA and Bonferroni's Multiple Comparison Test WT vs *sek-1(km4)* (ns) WT vs *goa-1(sa734)* (p<0.001) *goa-1(sa734)* vs *goa-1(sa734); sek-1(km4)* (p<0.001)	< 0.001
S1F	One-way ANOVA and Bonferroni's Multiple Comparison Test	< 0.001

	WT vs *sek-1(km4)* (p<0.001) WT vs *eat-16(tm775)* (p<0.001) *eat-16(tm775)* vs *eat-16(tm775); sek-1(km4)* (p<0.001) *sek-1(km4)* vs *eat-16(tm775); sek-1(km4)* (ns)	
S1G	One-way ANOVA and Bonferroni's Multiple Comparison Test WT vs *eat-16(tm775)* (p<0.001) *eat-16(tm775)* vs *eat-16(tm775); sek-1(km4)* (ns)	< 0.001
S2A	One-way ANOVA and Bonferroni's Multiple Comparison Test WT vs *sek-1(km4)* (p<0.001) *sek-1(km4)* vs *sek-1(km4); yakEx82[unc-17Hp::sek-1(+)]* (p<0.05) *sek-1(km4)* vs *sek-1(km4); yakEx83[unc-17βp::sek-1(+)]* (p<0.01)	< 0.001
S2B	One-way ANOVA and Bonferroni's Multiple Comparison Test *egl-30(tg26); sek-1(km4)* vs *egl-30(tg26); sek-1(km4); yakEx82[unc-17Hp::sek-1(+)]* (p<0.001) *egl-30(tg26); sek-1(km4)* vs *egl-30(tg26); sek-1(km4); yakEx83[unc-17βp::sek-1(+)]* (p<0.001)	< 0.001
S2C	One-way ANOVA and Bonferroni's Multiple Comparison Test WT vs *egl-30(tg26)* (p<0.001) *egl-30(tg26); sek-1(km4)* vs *egl-30(tg26); sek-1(km4); qdEx8[unc-119p::sek-1(+)]* (p<0.001) *egl-30(tg26); sek-1(km4)* vs *egl-30(tg26); sek-1(km4); yakEx72[unc-17p::sek-1(+)]* (p<0.001) *egl-30(tg26); sek-1(km4)* vs	< 0.001

	egl-30(tg26); sek-1(km4); yakEx73[unc-47p::sek-1(+)] (ns) *egl-30(tg26); sek-1(km4)* vs *egl-30(tg26); sek-1(km4);* *yakEx82[unc-17Hp::sek-1(+)]* (p<0.001) *egl-30(tg26); sek-1(km4)* vs *egl-30(tg26); sek-1(km4);* *yakEx83[unc-17βp::sek-1(+)]* (ns)	
S3A	One-way ANOVA and Bonferroni's Multiple Comparison Test *jnk-1(gk7)* vs *jnk-1(gk7); sek-1(km4)* (p<0.01) *jkk-1(km2)* vs *pmk-2(qd279 qd171) pmk-1 (km25); jkk-1(km2)* (p<0.05)	< 0.05
S3B	One-way ANOVA and Bonferroni's Multiple Comparison Test WT vs *egl-30(tg26)* (P<0.001) *egl-30(tg26)* vs *egl-30(tg26); jkk-1* (ns) *egl-30(tg26)* vs *egl-30(tg26); jnk-1* (ns) *egl-30(tg26)* vs *egl-30(tg26); atf-7* (ns)	
S3C	One-way ANOVA and Bonferroni's Multiple Comparison Test WT vs *atf-7(qd22)* (p<0.001) WT vs *atf-7(qd22 qd130)* (p<0.001)	< 0.001
S3D	One-way ANOVA	p=0.806
S3F	Unpaired t test, two-tailed	< 0.001
S4A	One-way ANOVA and Bonferroni's Multiple Comparison Test WT vs *nca-1(ox352)* (p<0.001) *nca-1(ox352)* vs *nca-1(ox352); sek-1(km4)* (p<0.001) *nca-1(ox352)* vs *nca-1(ox352); nsy-1(ok593)* (p<0.001)	<0.001
S4B	Kruskal-Wallis Test and Dunn's Multiple Comparison Test	0.001

	nca-1(ox352);sek-1(km4) vs *nca-1(ox352);sek-1(km4); qdEx8[unc-119p::sek-1(+)]* (ns) *nca-1(ox352);sek-1(km4)* vs *nca-1(ox352);sek-1(km4); yakEx72[unc-17p::sek-1(+)]* (ns) *nca-1(ox352);sek-1(km4)* vs *nca-1(ox352);sek-1(km4); yakEx73[unc-47p::sek-1(+)]* (ns)	
S4C	One-way ANOVA and Bonferroni's Multiple Comparison Test *nca-1(ox352)* vs *nca-1(ox352); sek-1(km4)* (p<0.001) *nca-1(ox352); sek-1(km4)* vs *nca-1(ox352); sek-1(km4); qdEx8[unc-119p::sek-1(+)]* (p<0.001) *nca-1(ox352); sek-1(km4)* vs *nca-1(ox352); sek-1(km4); yakEx72[unc-17p::sek-1(+)]* (p<0.001) *nca-1(ox352); sek-1(km4)* vs *nca-1(ox352); sek-1(km4); yakEx73[unc-47p::sek-1(+)]* (ns) *nca-1(ox352); sek-1(km4)* vs *nca-1(ox352); sek-1(km4); yakEx82[unc-17Hp::sek-1(+)]* (p<0.001) *nca-1(ox352); sek-1(km4)* vs *nca-1(ox352); sek-1(km4); yakEx83[unc-17βp::sek-1(+)]* (p<0.001)	<0.001

Figure S1

Figure S1. *sek-1* interacts with Gq and Go mutants

(A) The *sek-1(km4)* mutation suppresses the hyperactive locomotion of the activated Gq mutant *egl-30(tg26)*. ***, p<0.001, error bars = SEM, n=20.

(B) *sek-1(ag1)* mutant animals have slow locomotion. ***, p<0.001, error bars = SEM, n=10.

(C) The *unc-82(e1220)* mutation reduces the locomotion rate of the activated Gq mutant *egl-30(tg26)*. ***, p<0.001, error bars = SEM, n= 20.

(D) *sek-1(km4)* suppresses the hyperactivity of *goa-1(sa734)*. ***, p<0.001, error bars = SEM, n=20.

(E) *sek-1(km4)* suppresses the loopy waveform of *goa-1(sa734)*. ***, p<0.001, error bars = SEM, n=5.

(F) *sek-1(km4)* suppresses the hyperactivity of *eat-16(tm775)*. ***, p<0.001, error bars = SEM, n=20.

(G) *sek-1(km4)* does not suppress the loopy waveform of *eat-16(tm775)*. ns, p>0.05, error bars = SEM, n=5.

Figure S2

Figure S2. *sek-1* **acts in both head acetylcholine neurons and acetylcholine motorneurons**
(A) *sek-1* acts in acetylcholine motorneurons to modulate locomotion rate. The *sek-1* WT cDNA driven by the *unc-17β* acetylcholine motorneuron promoter [*unc-17βp::sek-1(+)*] rescues the slow locomotion phenotype of *sek-1(km4)* worms, but *sek-1* expression in head acetylcholine neurons using the *unc-17H* promoter [*unc-17Hp::sek-1(+)*] does not rescue. **, p< 0.01; *, p<0.05 compared to *sek-1*. Error bars = SEM, n=20.
(B) *sek-1* acts in both head acetylcholine neurons and acetylcholine motorneurons to modulate the locomotion rate of the activated Gq mutant *egl-30(tg26)*. *egl-30(tg26) sek-1(km4)* worms expressing either *unc-17Hp::sek-1(+)* or *unc-17βp::sek-1(+)* have an increased locomotion rate compared to *egl-30(tg26) sek-1*. ***, p<0.001 compared to *egl-30(tg26) sek-1*. Error bars = SEM, n=20.
(C) *sek-1* acts in head acetylcholine neurons to modulate the loopy waveform of the activated Gq mutant *egl-30(tg26)*. *egl-30(tg26) sek-1(km4)* worms expressing *unc-17Hp::sek-1(+)* are loopy like *egl-30(tg26)*, but *egl-30(tg26) sek-1(km4)* worms expressing *unc-17βp::sek-1(+)* are similar to *egl-30(tg26) sek-1*. ***, p<0.001; ns, p>0.05 compared to *egl-30(tg26) sek-1*. Error bars = SEM, n=5.

Figure S3

Figure S3. Locomotion of p38 and JNK MAPK pathway mutants

(A) *jkk-1* and *jnk-1* act in parallel to *sek-1* and *pmk-2 pmk-1*. The *jnk-1(gk7) sek-1(km4)* double mutant and *pmk-2(qd279 qd171) pmk-1(km25) jkk-1(km2)* triple mutants move more slowly than the respective individual mutants. **, p< 0.01, *, p<0.05. Error bars = SEM, n=20.

(B) Mutations in *jkk-1, jnk-1,* and *atf-7* do not suppress the loopy waveform of the activated Gq mutant *egl-30(tg26)*. ns, p>0.05, error bars = SEM, n=5.

(C) Worms with gain-of-function or loss-of-function alleles of *atf-7* are slower than wild-type worms. ***, p< 0.001, error bars = SEM, n=20.

(D) Worms lacking *gap-2* and *vhp-1* move like wild-type worms. Neither *gap-2(tm478)* nor *vhp-1(sa366)* confers a slow locomotion phenotype. ns, p>0.05 compared to WT. Error bars = SEM, n=20.

(E-F) *tir-1(ky648gf)* animals do not have loopy or hyperactive locomotion. *tir-1(ky648gf)* worms have wild-type posture and are slower than wild-type animals. ***, p< 0.001, error bars = SEM, n=20.

Figure S4

A

B

C

Figure S4. *sek-1* and *nsy-1* weakly suppress *nca-1(gf)*
(A) Mutations in *sek-1* and *nsy-1* suppress the small body size of *nca-1(ox352)* mutant worms.
***, p<0.001, error bars = SEM, n=10.
(B) None of the neuronal *sek-1* rescuing constructs reverse the radial locomotion phenotype of
nca-1(gf) sek-1(km4) animals. ns, p>0.05. Error bars = SEM, n=19-24.
(C) *sek-1* acts in both head acetylcholine neurons and acetylcholine motorneurons to control the
body size of *nca-1(gf)*. *nca-1(ox352) sek-1(km4)* worms expressing *sek-1* in all neurons (*unc-119p::sek-1(+)*),
acetylcholine neurons (*unc-17p::sek-1(+)*), head acetylcholine neurons (*unc-17Hp::sek-1(+)*), or acetylcholine
motorneurons (*unc-17βp::sek-1(+)*) have a similar size to *nca-1(gf),* but *nca-1(ox352) sek-1(km4)* worms
expressing sek-1 in GABA neurons (*unc-47p::sek-1(+)*) are similar to *nca-1(gf) sek-1.*
***, p< 0.001; ns, p>0.05. Error bars = SEM, n=7-10.