1 **Cultivation and genomics of the first freshwater SAR11 (LD12) isolate**
2

3 Michael W. Henson[1], V. Celeste Lanclos[1], Brant C. Faircloth[1,2], and J. Cameron Thrash[1,3]
4
5 1. Department of Biological Sciences, Louisiana State University, Baton Rouge, LA 70803,
6 U.S.A.
7
8 2. Museum of Natural History, Louisiana State University, Baton Rouge, LA 70803, U.S.A.
9
3. Materials and Correspondence:
J. Cameron Thrash
Department of Biological Sciences
202 Life Sciences Bldg.
Louisiana State University
Baton Rouge, LA 70803
thrashc@lsu.edu
225-578-8210

18
19
20
21 Key Words: LD12, SAR11, freshwater microbiology, genomics, physiology

Abstract

Abstract
Evolutionary transitions between fresh and salt water happen infrequently among
bacterioplankton. Within the ubiquitous and highly abundant heterotrophic
Alphaproteobacteria order *Pelagibacterales* (SAR11), most members live in marine
habitats but the LD12 subclade has evolved as a unique freshwater lineage. LD12 cells
occur as some of the most dominant freshwater bacterioplankton, yet this group has
remained elusive to cultivation, hampering a more thorough understanding of its biology.
Here we report the first successful isolation of an LD12 representative, strain LSUCC0530,
using high throughput dilution to extinction cultivation methods, and its complete genome
sequence. Growth experiments corroborate ecological data indicating active populations of
LD12 in brackish water up to salinities of ~5. LSUCC0530 has the smallest closed genome
thus far reported for a SAR11 strain (1.16 Mbp). The genome affirms many previous
metabolic predictions from cultivation-independent analyses, like a complete Embden-
Meyerhof-Parnas glycolysis pathway, but also provides novel insights, such as the first
isocitrate dehydrogenase in LD12, a likely homologous recombination of malate synthase
from outside of the SAR11 clade, and analogous substitutions of ion transporters with
others that occur throughout the rest of the SAR11 clade. Growth data support
metagenomic recruitment results suggesting temperature-based ecotype diversification
within LD12, and key gene losses for osmolyte uptake provide a succinct hypothesis for the
evolutionary transition of LD12 from salt to fresh water. For strain LSUCC0530 we propose
the provisional nomenclature *Candidatus* Fonsibacter ubiquis.
43

Introduction

Bacterioplankton in the SAR11 clade of *Alphaproteobacteria* are dominant heterotrophs in both marine and freshwater systems. In the oceans, SAR11 can represent 25-50% of total planktonic cells[1,2]. Numerous subclades with unique spatio-temporal distributions comprise SAR11[1,3-5]. At least nine subclades defined via 16S rRNA gene sequences occupy marine niches[4], and more likely exist[6]. However, in spite of its global distribution[1], massive predicted population size of 10^{28} cells[1], and an estimated divergence time from the last common ancestor of 1.1 billion years ago[7], the bulk of existing evidence suggests that SAR11 has only successfully colonized freshwater environments once in its natural history[8-10]. All known freshwater SAR11 belong to subclade IIIb, a.k.a. LD12, although a very recent report challenges this assertion[11]. Regardless, the limited evolutionary diversification into less saline habitats has not prevented LD12 from achieving ecosystem prominence where it lives. In many lotic and lentic environments, LD12 occupies similar relative abundances as its marine cousins[9,12-15]. Study of LD12 is important to understand SAR11 evolution specifically, and how successful transitions between marine and freshwater environments occur in bacterioplankton[16] more generally.

Ecological, functional, and sequence-based inference from single amplified genomes (SAGs) and metagenomes support the hypothesis that LD12 bacterioplankton evolved from a genome-streamlined marine ancestor[7,9,17,18]. As such they share many of the same characteristics as marine SAR11, such as small cell volumes; adaptation to oligotrophic habitats; small, streamlined genomes; an obligate aerobic chemoorganoheterotrophic lifestyle with limited metabolic flexibility; preference for small molecular weight compounds like carboxylic and amino acids as carbon/energy sources; and auxotrophies for some amino acids and vitamins[5,9,10,15,17-24]. Previous research suggests that where LD12 differ from their marine counterparts lies in specific elements of metabolic potential that suggest a greater emphasis on production, rather than uptake, of osmolytes, and important metabolic changes related to energy production[15,17]. For example, metagenomic population data showed a correlation between decreased salinity and greater proportion of the Embden-Meyerhof-Parnass (EMP) vs. Entner-Doudoroff (ED) glycolysis pathways[15]. Comparative genomic analyses of SAGs from different SAR11 strains concurred: LD12 genomes contained the EMP pathway which is not found in most marine SAR11[17,19]. SAGs also suggested that LD12 lacks the glyoxylate shunt and some single carbon (C1) metabolism[17].

In spite of what has been learned from cultivation-independent methods, the lack of cultured LD12 representatives has hampered a more detailed exploration of the group. Potential ecotypes within LD12 have been identified[18], but we cannot delineate what distinguishes them without better physiological and genomic data. Interpreting data on the ecological distribution of LD12 remains challenging without information on growth tolerances and optima for salinity and temperature. We also don't understand whether a connection exists between more efficient energy production through glycolysis and the

freshwater lifestyle, or what other adaptations explain LD12 evolution away from salt
water.
The next steps in translating 'omics-based predictions into measured data for
integration with ecosystem models require living experimental subjects. For example,
cultures of marine SAR11, such as HTCC1062 and HTCC7211, have facilitated testing of
metabolism and growth[21-23,25-29], structural organization[30], and virus-host interactions[31].
We need cultivated representatives to provide this kind of understanding of other
important bacterioplankton like LD12. In service of this goal, we pursued a systematic high
throughput cultivation effort from coastal regions in the northern Gulf of Mexico. Here we
report the first isolation of a member of the LD12 clade, strain LSUCC0530, its complete
genome sequence, preliminary physiological data, and an examination of the ecological
distribution of this organism compared with other LD12 clade members. Our results
provide evidence for temperature-dependent ecotype diversification within LD12, and a
hypothesis about evolutionary process that led to the unique SAR11 colonization of fresh
water.
99
Results
Isolation and growth of LSUCC0530
We isolated strain LSUCC0530 as part of a series of high throughput dilution-to-extinction
cultivation experiments (HTC) conducted with inoculation water obtained across the
southern Louisiana coast. The particular experiment that yielded LSUCC0530 utilized
surface water from the coastal lagoon of Lake Borgne, 39 km southeast of New Orleans (Fig.
S1). At the time of collection, the surface water had a salinity of 2.39 and was 30.5°C. Later
community analysis demonstrated that LD12-type organisms represented 8.7% of the
bacterioplankton population in the inoculum (see below). We diluted Lake Borgne surface
water for inoculation to an estimated 2 cells well^{-1} into our JW5 medium (salinity 1.45,
Table S1), and incubated cultures for 30 days at 23°C (room temperature in the Thrash
Lab). LSUCC0530 grew slowly, reaching 8.6 x 10^5 cells mL^{-1} after 24 days. It had a low
fluorescence and low side scatter flow cytometry signature characteristic of other SAR11
cells (Fig. S1).
114
Taxonomy and morphology
The LSUCC0530 16S rRNA gene sequence had > 99% identity with eight sequences from
members of LD12, and only 91% and 92% sequence identity with sequences from the
sister subclade IIIa representatives IMCC9063 and HIMB114, respectively. LSUCC0530 also
had average amino acid identities (AAI) of 88.7% and 89.8% with HIMB114 and IMCC9063,
respectively, corroborating LSUCC0530 as belonging to separate species from subclade
IIIa[32]. Phylogenetic inference using 16S rRNA genes placed LSUCC0530 within the Order
Pelagibacteriales (a.k.a. SAR11[5]), in a monophyletic group with the original LD12 clone
library sequence (Fig. S2). Phylogenomic inference using 83 single copy marker genes from

49 SAR11 genomes also strongly supported placement of LSUCC0530 within LD12 as an
early diverging member with only the SAG AAA280-B11 from Lake Damariscotta diverging
earlier (Fig. 1A). This tree also recovered the three microclusters within LD12 (A, B, C)
observed previously[18] (Fig. 1A). Cells of strain LSUCC0530 were small, curved rods,
approximately 1 µm x 0.1 µm (Fig. 1B, C), showing conserved morphology with the
distantly related SAR11 strain HTCC1062 in subclade Ia[30,33].
130
Genome characteristics
Genome assembly resulted in a single, circularized scaffold that we tested for completeness
using multiple tools (Supplemental Information). The completed genome is 1,160,202 bp
with a GC content of 29.02% and 1271 predicted genes (Fig. 2). There are 1231 putative
protein coding genes and 40 RNA genes, including one each of the 5S, 16S, and 23S rRNA
genes. We found no genomic evidence of lysogenic bacteriophage or a CRISPR-*cas* system.
LSUCC0530 has the characteristics of genome streamlining previously reported for other
SAR11 genomes, with an estimated coding percentage of 96.36% and the smallest complete
SAR11 genome reported thus far[19,20]. However, in spite of its unusually small overall size,
intergenic spacer regions are not significantly smaller than those of the sister subclade IIIa
(Fig. S3). All LD12 genomes had much higher AAI and synteny with each other than with
the subclade IIIa (Table S1), mirroring their phylogenomic relationships (Fig. 1A).
However, within the LD12 clade, AAI and synteny percentages did not clearly delineate the
same patterns seen in the phylogenomic tree, which may have resulted from the LD12 SAGs
being fragmented and incomplete.
The LSUCC0530 genome has the conserved architecture for hypervariable region
(HVR) 2, observed in most other complete SAR11 genomes[19,34,35] and other LD12 SAGs[18],
between the 16S rRNA-tRNA-tRNA-23S rRNA operon and the 5S rRNA gene operon. This
region is 54,755 bp and has a clear variation in GC content from neighboring parts of the
genome (Fig. 2). Metagenomic recruitment also verified this region as hypervariable based
on the lack of matching sequences in community data (Fig. S4). The predicted gene
annotations in HVR2 resemble those in the same location in other SAR11 genomes[18,19],
namely genes likely functioning in membrane modification such as glycosyltransferases,
epimerases, transaminases, and methyltransferases; as well as hypothetical proteins.
155
Metabolic reconstruction
Like most other SAR11s[5,17,19], the LSUCC0530 genome encodes for an obligate aerobic
lifestyle, with a complete oxidative phosphorylation pathway including a heme/copper-
type cytochrome c oxidase, a proton-translocating NADH dehydrogenase, and a proton-
translocating ATP synthase (Fig. 3). As previously detected in LD12 SAGs[17], the LSUCC0530
genome encodes a complete EMP glycolysis pathway, including phosphofructokinase and
pyruvate dehydrogenase, and complete gluconeogenesis through nucleotide sugar
production (Fig. 3). We recovered genes for the pentose phosphate pathway and a

predicted fructokinase gene for conversion of D-fructose to ß-D-fructose-6-P. As expected
based on other SAR11 genomes, we did not find genes for utilization of allose, fucose,
fuculose, galactose, maltose, mannose, rhamnose, sucrose, starch, trehalose, or xylose.
LSUCC0530 has a complete TCA cycle and glycolate oxidase, and like other SAR11s, the
non-mevalonate pathway for isopentenyl diphosphate (Fig. 3).
We recovered the first isocitrate lyase in an LD12 genome, conferring a complete
glyoxylate bypass. The isocitrate lyase shares its ancestry with the majority of the SAR11
clade except subclade IIIa, where it appears to have been lost[17,19] (Fig. S5). However, the
LSUCC0530 malate synthase shared with other LD12 SAGs belongs to isoform A, whereas
all other SAR11 clades contain isoform G[36] (Figs. 3; S6). Comparison of gene neighborhoods
in a variety of SAR11 genomes revealed that both isoforms occur in the same location (Fig.
S7).
We predict production of eighteen amino acids (Fig. 3) and partial synthesis
pathways for three more (histidine, cysteine, and methionine), as well as the ability to
interconvert, but not synthesize *de novo*, phenylalanine and tyrosine. The LSUCC0530
genome has lost some of the C1 and methylated compound pathways found in other
SAR11s[24]. For example, as in the LD12 SAGs[17], LSUCC0530 does not have DMSP,
methylamine, or glycine-betaine metabolism. However, it retains tetrahydrafolate
metabolism, formate oxidation, and the glycine cleavage pathway (Fig. 3).
The genome has the same PII nitrogen response system and *amtB* ammonium
transporter found in other SAR11s, and is predicted to utilize ammonia as a nitrogen
source in amino acid synthesis (Fig. 3). The genome also encodes a putative thiosulfate/3-
mercaptopyruvate sulfurtransferase that may be involved in sulfur and thiocyanate
reactions (Fig. 3). A possible product of this enzyme is sulfite, which could be further
oxdized to sulfate in the periplasm via putative *sorAB* genes. This gene pair occurs
infrequently in the SAR11 clade, but sulfite oxidation may serve in some strains as a means
to generate additional proton motive force[37,38], similarly to proteorhodopsin[21,29,39], which
the LSUCC0530 genome also contains. Also like other SAR11s, a complete assimilatory
sulfate reduction pathway is missing, making LSUCC0530 dependent on reduced sulfur
compounds for cellular sulfur requirements[22]. In our media, this was supplied in the form
of sulfur-containing amino acids (Table S1). The genome encodes partial pathways for
incorporation of sulfide into cysteine and methionine, but is missing predicted serine O-
acetyltransferase and homoserine O-succinyltransferase genes (Fig. 3).
In addition to the PII nitrogen sensor, LSUCC0530 also has putative two component
systems handling phosphate limitation, pH, osmotic upshift via potassium, and redox state
(Fig. 3). The LSUCC genome encodes ABC transporters for phosphate (*ptsABCS*),
lipoprotein, as well as branched-chain and L amino acid uptake. We identified tripartite
ATP-independent periplasmic (TRAP) transporter genes for C4-dicarboxylate and
mannitol/chloroaromatic compounds. However, the gene cluster appears to be missing the
substrate binding subunit, and we did not find predicted genes for mannitol utilization.

In general, LSUCC0530 and the LD12 SAGs share many of the same pathways for
osmolyte synthesis as the subclade IIIa and other SAR11 organisms, such as N-acetyl-
ornithine, glutamate, glutamine, and proline (Table 1). In some cases, there appear to be
analogous substitutions in osmolyte transport systems. For example, LD12 genomes have
potassium antiporters, but not the *trkAH* genes for potassium transport in other SAR11s.
Similarly, a sodium/proton antiporter ortholgus cluster comprised genes exclusively from
LD12 genomes, but a separate sodium/proton antiporter occurred in all other SAR11
genomes (Table 1). Perhaps the most important difference in genomic content concerning
osmolytes occurs in two key transporter losses: LSUCC0530 and the other LD12s have no
genes for the glycine betaine/proline or ectoine/hydroxyectoine ABC transporters that are
present in HIMB114 and IMCC9063 and other subclades of SAR11 (Table 1).
215
Physiology
Under isolation conditions (23°C, JW5 medium), strain LSUCC0530 grew to a density of 2 x
10^7 cells mL^{-1} with an average growth rate of 0.53 day^{-1}. Our physiological experiments to
determine the salinity tolerance and growth optima revealed that LSUCC0530 could grow
at salinities between 0.36 and 4.7 (Fig. 4A). The optimal salinity for LSUCC0530 was at or
below the isolation medium of 1.45. Growth rates decreased between salinities of 2.9 and
4.7 (Fig. 4A), and cells died (determined by rapid loss of flow cytometric signal) in salinities
above 8 (Fig. S8). We observed growth after a very extended lag phase in a subset of
cultures at salinity 5.8, but could not reproduce this growth in repeated experiments (Fig.
S8). Among the temperatures tested, LSUCC0530 grew optimally at 23°C, and we observed
growth up to 30°C, but not at 12°C or below, nor at 35°C or above (Fig. 4B). However, we
did not observe a clear loss of signal at 4, 12, 35, or even 40°C (Fig. S8), even after over 33
days, suggesting that LSUCC0530 can endure extended periods at these temperatures in a
non-growth state. Notably during growth experiments, one of the five replicates during any
given experiment frequently grew at a reduced rate than the others (Fig. S8).
231
Ecology
We quantified the abundance of LD12 taxa in the coastal Louisiana ecosystems using 16S
rRNA gene sequences clustered into operational taxonomic units (OTUs). These sequences
were obtained as part of the combined sampling and HTC campaign that included the
successful isolation of LSUCC0530. Based on BLASTn sequence similarity, the
representative sequence for OTU7 and LSUCC0530 shared 100% identity over the 250bp
v4 region of the 16S rRNA gene, confirming the Mothur classification of OTU7 as belonging
to the SAR11 LD12 clade. Over the three year sampling period, OTU7 was the fifth most
abundant OTU based on total read abundance and sixth ranked OTU based on median read
abundance. At the time of sampling for the experiment that resulted in the isolation of
LSUCC0530, OTU7 had a relative abundance of 8.7% in Lake Borgne. In some sites, such as
the Atchafalaya River Delta (salinity 0.18), LD12 represented as much as 16% of total

bacterioplankton (Fig. S9). Generally, the LD12 OTU7 relative abundance correlated
negatively with salinity. At seven sites with salinities below 6, OTU7 had a relative
abundance of 4.0% or greater with a median relative abundance of 8.8% (Fig. S9). At the
remaining ten sites above a salinity of 6 (median 19.44), OTU7 had relative abundances
around 2% or lower (Fig. S9).
The phylogenetic separation of LSUCC0530 from B11 and the IIIb A-C microclusters
raised the question of whether LSUCC0530 represents a novel ecotype with unique
spatiotemporal distribution compared to other LD12 taxa. Therefore, we explored the
distribution of LD12 genomes in 90 fresh and brackish water environments through
competitive recruitment of metagnomic sequences to all the SAR11 genomes in the
phylogenomic tree and the outgroup sequence HIMB59. We quantified relative abundance
via recruited reads (per kpb of metagenomic sequence per Mbp of genome sequence-
RPKM), and aggregated LD12 data based on the A, B, and C microclusters (Fig. 1A), the B11
SAG, and LSUCC0530. Figure 5A displays RPKM values for reads with ≥ 95% identity,
roughly corresponding to a species delineation via average nucleotide identity (ANI)[40], but
additional comparisons at ≥ 85, 90, 92, 98, and 100% identity are provided in Figure S10.
All LD12 subgroups had their highest RPKM values in very low salinity sites,
although we could detect LD12 across various brackish salinities (Fig. 5A). In general, IIIb.A
and LSUCC0530 averaged the highest relative abundances across all sites with salinities
between 0 and 5.75. All subgroups showed site-specific variation in abundances (Fig. 5A),
indicating that the available genomes likely do represent specific ecotypes within LD12.
Microcluster IIIb.A clearly dominated recruitment at Lake Michigan, whereas LSUCC0530
represented the most abundant genome at Feitsui Reservoir and Lake Gatun (Fig. 5A).
By examining recruitment density to the different microclusters across a range of
percent identities, we also observed patterns indicating additional LD12 diversity beyond
that represented in the current genomes. For example, whereas the LSUCC0530 genome
recruited the majority of high percent identity reads in the Feitsui Reservoir and Lake
Gatun samples (Fig. 5B), in some locations the majority of recruitment to this genome
centered around lower percent identity hits (Fig. 5C). We observed the same kind of
alternative patterns for the other microclusters as well (Fig. 5D). Since reads were
competitively recruited against all close relative genomes, these cases where a
microcluster primarily recruited reads ~90-92% identity likely mean that we lack a
representative genome for that fraction of the metagenomic data.
277
Discussion
The successful cultivation of and LD12 strain has demonstrated for the first time that this
organism shares many physiological traits with cultured marine SAR11 representatives.
LSUCC0530 doubles roughly once every two days, similarly to taxa in subclade Ia[33] and
Ib[41]. It has an aerobic, chemoorganoheterotrophic lifestyle, thriving in oligotrophic media
with simple carbon compounds, and likely depends on reduced organosulfur compounds

and ammonium as its sulfur and nitrogen sources, respectively. Cell size and shape
demonstrated considerable morphological conservation (small, curved rods roughly 1 x 0.1
μm) across large evolutionary distances (LD12 vs. subclade Ia ~89% 16S rRNA gene
identity).

Cultivation of LSUCC0530 has also helped to constrain important physiological
differences between LD12 and marine SAR11. Although a systematic survey of salinity
tolerance has not been conducted for other cultured SAR11 representatives, we have
shown that LD12 cannot survive in media with salinities over eight, whereas all other
cultured SAR11 grow in seawater salinities[25,33,41-43]. LSUCC0530 thrives in relatively high
temperatures (23-30°C), and we could not confirm growth at low temperatures where
other SAR11 have been isolated[42].

LSUCC0530 shares many genomic characteristics with marine SAR11, such as a
small, streamlined genome with few paralogs, short intergenic spaces, and a conserved
hypervariable region, confirming predictions from LD12 SAGs and metagenomes[12,17,18]. The
LSUCC0530 genome also encodes much of the same metabolic potential as what has been
reported through LD12 SAGs and metagenomic data. For example, we confirm the EMP
glycolysis pathway, TCA cycle, non-mevalonate terpenoid biosynthesis, numerous amino
acid synthesis pathways, TRAP transporters for C4 dicarboxylate compounds, and ABC
transporters for various amino acids. We can confirm the lack of a phosphoenolpyruvate
carboxylase (ppc) and missing pathways for DMSP, glycine-betaine, and methylamine
metabolism.

We have also recovered important genetic information previously unknown for
LD12. At 1.16 Mbp, LSUCC0530 has a smaller genome, with fewer genes, than any other
SAR11. LSUCC0530 has a complete glyoxylate bypass, which differs from previous
inferences using SAGs[17]. The unusual LD12 malate synthase isoform A gene occupies the
same genomic location in LSUCC0530 as the isoform G copy in most subclade Ia, Ic, and IIIa
genomes, with the exception of HIMB114 (Fig. S7). Therefore, the most parsimonious
explanation for the presence of isoform A in LSUCC0530 is a homologous recombination
event that replaced the isoform G homolog. The LSUCC0530 genome also contains putative
pathways for oxidation of sulfide, thiosulfate, and sulfite, which have not yet been explored
in any cultured SAR11.

The relative abundance data from coastal Louisiana, the isolation of an LD12 strain
from a coastal environment, and the ability of LSUCC0530 to grow in slightly brackish
water supports previous work indicating that LD12 occupies a much wider range of aquatic
habitats than inland freshwater systems[9,15,44,45]. However, metagenomic recruitment
provides supporting evidence that LD12 has diversified into multiple ecotypes with unique
distributions (Fig. 5A), and new evidence that in many environments, additional LD12
diversity exists beyond that found in current genomes (Fig. 5C-D). In addition, our data
suggests that LSUCC0530 likely represents a novel LD12 ecotype since it preferentially

recruits metagenomic sequences at high percent identity in different environments than
for the other microclusters of LD12 (Fig. 5).
What distinguishes the distribution of these ecotypes remains unclear. Small
variations in salinity optima for different ecotypes may drive diversification. However,
given the paucity of low brackish sites in available datasets, we cannot establish whether
salinity plays a role in ecotype diversification. Furthermore, without complete genomes
from other LD12 ecotypes, we cannot rule out whether alternative ionic strength
management strategies, or some other metabolic characteristic, help define evolutionary
differences associated with ecotypes. For example, the ability to utilize alternative sulfur
sources may distinguish LSUCC0530 from other LD12. The *sorAB* sulfite dehydrogenase
and the thiosulfate/3-mercaptopyruvate sulfurtransferase genes have uneven distribution
among SAR11 genomes, including within the LD12 subclade.
An alternative hypothesis is that temperature has driven diversification between
ecotypes. A striking feature of LSUCC0530 is its growth preference for relatively high
temperatures and inability to grow at low temperatures. While robust growth at 30°C may
seem intuitive given that this represents typical Lake Borgne surface water temperature,
the inability to grow at 12°C or below would likely place LSUCC0530-type LD12 cells at a
fitness disadvantage in places like Lake Michigan, where annual temperatures rarely get
above 22°C and average temperatures remain below 15°C for the majority of the year
(https://coastwatch.glerl.noaa.gov/). The metagenomic recruitment data support this
hypothesis: the samples in which LSUCC0530 dominated recruitment (Lake Gatun, Feitsui
Reservoir) were all above 19°C. Conversely, other microclusters dominated recruitment of
samples from colder environments (e.g. IIIb.A in Lake Michigan)(Fig. 5). More sequence
data may strengthen the relationship between temperature and ecotypes, but since
temperature optima is extremely hard to infer from genomes alone, additional cultivars
will be needed to test the temperature hypothesis for ecotype diversification.
A critical question remains for LD12. What has made these cells uniquely adapted to
low salinity waters? Genomic analyses here and elsewhere have helped define the
differences in LD12 metabolic potential with that of other SAR11, but very few adaptations
appear related to salinity. It has been noted that the EMP pathway provides more ATP and
NADH than the ED pathway, thus conferring a greater energy conservation through
glycolysis for LD12 cells than other SAR11s[15,17]. If or how this energy conservation relates
to salinity adaptation has not been pursued. The presence of unique LD12 cation-proton
antiporters (Fig. 3) could provide a connection. LD12 cells may benefit from additional
NADH for the production of a proton motive force that drives the potassium-proton
antiporters, thus improving active transport to lower internal ionic strength. However,
while this adaptation may improve the ability of LD12 cells to live in fresh water, it doesn't
explain why they have lost the ability to survive in salt water where marine SAR11 cousins
thrive.

The most likely explanation comes in the distribution of genes for osmolyte uptake.
Critically, LSUCC0530 and LD12 in general have lost the *proVWX* glycine-betaine/proline
ABC transporter shared by all other marine SAR11 (Table 1). Although a relatively minimal
number of genes, this uptake capability likely represents a significant loss of function for
LD12 salinity tolerance. In *Escherichia coli*, the *proVWX* operon has a strong promoter that
upregulates these genes two orders of magnitude in response to increased salinity[46].
Indeed, rapid proline uptake is a common response to osmotic upshift in many bacteria[47].
Furthermore, LSUCC0530 and other LD12 genomes have lost the ABC transporters for
ectoine/hydroxyectoine present in the other SAR11 genomes. Given the slow growth of
LD12, rapid production of new intracellular osmolytes in response to increased salinity
seems unlikely. Thus, the loss of these transport systems for quickly equilibrating
intracellular osmolarity probably prevents successful dispersal of this organism into higher
salinity waters.
This finding allows us to speculate further about the evolutionary trajectory that led
to successful colonization of freshwater environments by LD12. All evidence suggests that
the last common SAR11 ancestor was a marine organism with a streamlined genome[7,8,18].
During further diversification into subclade III, additional genes and pathways were lost,
for example, ppc and genes for DMSP and some C1 compound metabolisms[17,48]. Therefore,
the common ancestor of subclade IIIa and LD12 would have had to adapt to freshwater
environments starting with an already reduced genome and limited metabolic flexibility. A
simple way to inhabit lower salinity environments would have been to decrease expression
of, or altogether lose, genes for uptake and/or production of osmolytes. If the adaptation
were reversible, we would expect to see examples of LD12 organisms growing in marine
systems. However, existing evidence suggests that while LD12 organisms can grow
(established herein) and be active[44] in low brackish conditions, they are not successful in
higher salinity brackish or marine water[8,9,15,45]. Thus, we propose that LD12 adaptation to
freshwater involved irreversible loss of function, decreasing osmotic response capability in
LD12 cells, and preventing recolonization of higher salinity environments. The missing ABC
transporters for proline/glycine betaine and ectoine/hydroxyectoine in LD12 are likely
candidates for this loss of function. Experimental tests on the differences in production and
uptake of osmolytes among subclade IIIa and LD12 representatives will also provide
critical insight into how these different evolutionary paths have resulted in unique salinity
ranges.
395
396   For the first cultured representative of the LD12 clade, we propose the provisional
397   taxonomic assignment for strain LSUCC0530 as '*Candidatus* Fonsibacter ubiquis',
398
Description of *Fonsibacter* gen. nov.
Fonsibacter (*Fons* L. noun; fresh water, spring water, -bacter, Gr. adj.; rod, bacterium.
Fonsibacter referring to the preferred low salinity habitat of this organism).

Aerobic, chemoorganoheterotrophic, and oligotrophic. Cells are small, curved rods roughly

1 x 0.1μm. Non-motile. Based on phylogenomics and 16S rRNA gene phylogenetics,

subclade IIIb/LD12 occurs on a separate branch within the *Pelagibacteraceae* (SAR11),

sister to subclade IIIa containing strains HIMB114 and IMCC9063. Due to the depth of

branching between these clades using concatenated protein-coding genes, and the 92%

and 91% 16S rRNA gene sequence identity with HIMB114 and IMCC9063, respectively, we

propose LSUCC0530 and subclade IIIb/LD12 as a novel genus in the *Pelagibacteraceae*.

Description of *Fonsibacter ubiquis* sp. nov.

Fonsibacter ubiquis (L. adv. ubiquitous).

In addition to the properties given in the genus description, the species is described as

follows. Growth occurs at temperatures between 23°C and 30°C, but not at 11°C or below,

nor at 35°C or above. Optimal salinity is 1.45 and below, and growth occurs between 0.36

and 4.68. At optimal temperature and salinity, cells divide at an average rate of 0.53 day^{-1}.

Genome size is 1.16 Mbp, with 1271 predicted genes, and a GC content of 29.02%

(calculated). LSUCC0530 had an ANI of 75.3% and 75.9% and an AAI of 88.7% and 89.8%

with HIMB114 and IMCC9063, respectively. The genome is available on GenBank under

accession number CP024034.

The type strain, LSUCC0530^T, was isolated from brackish water from the coastal lagoon

Lake Borgne off the coast of southeastern Louisiana, USA.

Materials and Methods

Isolation and Identification of LSUCC0530

Isolation, propagation, and identification of strain LSUCC0530 (subclade IIIb) was

completed as previously reported[14]. Briefly, water was collected from surface water in the

coastal lagoon Lake Borgne (Shell Beach, Louisiana) (29.872035 N, -89.672819 W) on 1

July, 2016. We recorded salinity on site using a YSI 556 MPS (YSI, Solon, OH, USA) (Table

S1). Whole water was filtered through a 2.7 μm filter (Whatman), stained with 1X SYBR

Green (Lonza, Basal, Switzerland), and enumerated using the Guava EasyCyte 5HT flow

cytometer (Millipore, Massachusetts, USA). After serial dilution, water was inoculated into

five 96 x 2.1 mL well PTFE plates (Radleys, Essex, UK) containing 1.7 mL of JW5 medium at

an estimated 2 cells well^{-1}. Cultures were incubated at room temperature in the dark for

three weeks and evaluated for growth using flow cytometry. Positive wells (> 10^4 cells mL^{-1}

^{1}) were transferred in duplicate to capped, 125 mL polycarbonate flasks (Corning, New

York, USA) containing 50 mL of JW5 medium. Upon reaching a density of > 5.0 x 10^5 cells

mL^{-1}, cells were filtered onto 25 mm 0.22 μm polycarbonate filters (Millipore,

Massachusetts, USA), and DNA extractions were performed using the MoBio PowerWater
DNA kit (QIAGEN, Massachusetts, USA) following the manufacturer's instructions. DNA was
amplified as previously reported[14] and sequenced at Michigan State University RTSF
Genomics Core. Sanger sequences were evaluated using the freely available software
4Peaks (v. 1.7.1)(http://nucleobytes.com/4peaks/). Forward and reverse sequences were
assembled using the CAP3 webserver (http://doua.prabi.fr/software/cap3), after reverse
reads were converted to their reverse complement at
http://www.bioinformatics.org/sms/rev_comp.html.
450
Community iTag analysis
Community DNA from 17 sites of various salinities was filtered, extracted, and analyzed
following our previously reported protocol[14]. The sites sampled were Lake Borgne (LKB;
Shell Beach, LA), Bay Pomme D'or (JLB; Buras, LA), Terrebonne Bay (TBon; Cocodrie, LA),
Atchafalaya River Delta (ARD; Franklin, LA), Freshwater City (FWC; Kaplan, LA), Calcasieu
Jetties (CJ; Cameron, LA). Each site was visited three times with the exception of TBon,
which was visited only twice. Briefly, extracted DNA was sequenced at the 16S rRNA gene
V4 region (515F, 806R)[49], with Illumina MiSeq 2x250bp paired-end sequencing at Argonne
National Laboratories. 16S rRNA gene amplicon data was analyzed and classified (OTU$_{0.03}$)
with Mothur v.1.33.3[50] using the Silva v119 database[51]. To determine the best matching
OTU for LSUCC0530, sequences from the OTU representative fasta files, provided by
mothur using *get.oturep*(), were used to create a blast database (*formatdb*) against which
the LSUCC isolate 16S rRNA genes could be searched via blastn (BLAST v 2.2.26). All best
hits ranked at 100% identity. OTU abundance analysis was done within the R statistical
environment v3.2.3, within the package PhyloSeq (v1.14.0)[52]. Sequences were rarefied
using the command *rarefy_even_depth*() and then trimmed for OTUs with at least 2 reads in
at least 90% of the samples. Abundances of an OTU were averaged across biological
duplicates and a rank abundance matrix was calculated using the command
transform_sample_counts() with the argument function of function(x) x / sum(x).
Completed rarefied rank abundance OTU tables were plotted using the graphing program
GGPLOT2 (v. 2.0.0)[53]. All iTag sequences are available at the Short Read Archive with
accession number SRR6235382-SRR6235415.
473
Growth experiments
Salinity tolerance was tested by growing LSUCC0530 in different media distinguished by
proportional changes in the concentration of major ions. All other media elements (carbon,
iron, phosphate, nitrogen, vitamins, and trace metals) were kept consistent. Media included
JW1, JW2, JW3, and JW4, previously published[14], and JW3.5, JW4.5, JW5, and JW6. Recipes
for all media used in this study are provided in Table S1. Salinity tolerance experiments
were conducted in quintuplicate at room temperature. Temperature range and sole carbon
substrate use were tested using the isolation medium, JW5. Temperature range

experiments were also conducted in quintuplicate. For all experiments, growth was
measured using flow cytometry as described above.
484
Microscopy
Strain LSUCC0530 cells were grown to near max density ($1x10^7$ cells mL^{-1}) in JW5 medium.
Cells for transmission electron microscopy were prepared as previously described[33] with
one minor change: centrifuged cells were concentrated in JW5 medium with no added
fixative. Cells were visualized under a JEOL JSM-2011 High Resolution transmission
electron microscope at the LSU Socolofsky Microscopy Center, Baton Rouge, LA. For
scanning electron microscopy, cultured cells were fixed for four hours in 4%
Glutaraldehyde buffered with 0.2M Cacodylate buffer and filtered onto a 0.2 µm filter (Pall,
Florida, USA). Samples were dehydrated in an ethanol series (30%, 50%, 70%, 96 and
100%) for 15 minutes each, followed by critical point drying. Cells were visualized under a
JEOL JSM-6610 scanning electron microscope at the LSU Socolofsky Microscopy Center,
Baton Rouge, LA.
497
Genome sequencing, assembly, and annotation
Cells were grown in 1 L JW5 medium (Table S1) and filtered onto 0.2 µm nylon filters
(Manuf, STATE). DNA was obtained via phenol-chloroform extraction and sequenced using
both Illumina HiSeq and MiSeq approaches. *HiSeq*. 350 ng of DNA was sheared to ~500 bp
with a Qsonica Q800R (QSonica) using 16 cycles with a amplitude of 25% and pulse rate of
20 seconds on, 20 seconds off. DNA was visualized on a 1.5% agrose gel stained with Gel
Red (Biotum) to determine average sheared length and quantified using Qubit high
sensitivity dsDNA assay kit (Invitrogen). Library prep was conducted following a modified
Kapa Hyper Prep Kit protocol (Kapa Biosystems, Inc.). Briefly, we ligated end-repaired and
adenylated DNA to universal Y-yoke oligonucleotide adapters and custom Illumina iTru
dual-indexed primers[54]. Prior to index amplification, post ligated LSUCC0530 DNA was
cleaned using a 1.2x SPRI cleanup to increase ligated DNA retention. Following a brief, 8
cycle PCR amplification of the index library, the genomic library was quantified using a
Qubit broad range dsDNA assay kit (Invitrogen). Illumina HiSeq 3000 sequencing was
performed at Oklahoma Medical Research Facility using SBS kit chemistry to generate 150
bp paired-end reads, insert size 400 bp. *MiSeq*. DNA was sent to the Argonne National
Laboratory Environmental Sample Preparation and Sequencing Facility which performed
library prep and sequencing, generating 250 bp paired-end reads, insert size 550 bp. Due
to the large number of reads from the HiSeq sequencing, we created a random subset using
14% of the total reads by using the program *seqtk* (v. 1.0-r75-dirty) and the flag –s100.
Initial assembly with the subset of reads was done with SPAdes[55]. This generated a large
single scaffold with overlapping ends. We performed a quality assessment of the single
scaffold via iterations of Reapr[56]-advised scaffold breaks and SSPACE[57] extensions using
both the full set of HiSeq reads and the MiSeq reads. Final assessment of the scaffold with

overlaps removed from the ends was performed with Pilon[58] using all *HiSeq* reads mapped
to the scaffold with BWA[59]. No issues were reported. An analysis of the GC skew was
performed on the final scaffold using the gc_skew script provided via Brown et al. 2015[60].
Contamination and completion were also evaluated with CheckM[61]. Detailed commands
and outputs for the assembly and quality assessment are provided in Supplementary
Information. The final circular chromosome scaffold was annotated by IMG[62] and is
publicly available with IMG Taxon ID number 2728369501. It is also publicly available in
GenBank with accession number CP024034.
530
Comparative genomics
Orthologus clusters were determined using the Get Homologues pipeline[63] using the
LSUCC0530 genome and 48 publicly available SAR11 genomes from IMG, including the 10
LD12 SAGs analyzed previously[17,18] (Table S1). Clusters were determined from amino acid
sequences using the OrthoMCL option in Get Homologues. All clusters are provided in
cluster_list.txt as part of Supplemental Information. Average amino acid identity (AAI) and
synteny were calculated as previously reported[64]. Average nucleotide identity (ANI) was
calculated with the Konstantinidis Lab website ANI calculator (http://enve-
omics.ce.gatech.edu/ani/index) as reported[65]. Compatible solutes identified in Table S1
were identified in SAR11 genomes using KO numbers, annotations, and cross-referenced
with orthologous clusters.
542
543   *Bacteriophage searches*
544   We looked for signatures of bacteriophage in the LSUCC0530 genome using two methods,
545   VirSorter[66] and PHASTER[67] with default settings. CRISPR regions are identified as part of
546   the standard IMG annotation process, and none were found in LSUCC0530.
547
Phylogenomics
We selected single-copy genes present in at least 39 of the 49 genomes as determined via
Get Homologues (above). Amino acid sequences from these 83 markers were separately
aligned with MUSCLE[68], culled with Gblocks[69], and concatenated into a superalignment
totaling 23,356 positions. Inference was performed with RAxML[70] using the PROTCATLG
model and 100 bootstrapping runs. All scripts (with settings) used in the alignment, culling,
concatenation, and inference processes are available in Supplemental Information. Node
labels were renamed with Newick Utilities[71] *nw_rename*, and the tree was visualized with
Archaeopteryx[72].
557
Single gene phylogeny
The 16S rRNA gene sequence from the LSUCC0530 genome was aligned with 284
alphaproteobacterial sequences and seven outgroup sequences. We completed
phylogenetic inference for *aceA* and *malA* in an analogous manner. For each, the

LSUCC0530 gene was searched against the NCBI nr database using blastp (v. 2.2.28). The
top 100 sequence hits were combined with all homologus SAR11 sequences identified with
Get Homologues for the *aceA* phylogeny, and both the *malA* and *malG* sequences in all
SAR11s for the *malA* phylogeny. Sequences from redundant taxa were removed. For all
trees, alignment with MUSCLE[68], culling with Gblocks[69], and tree inference with
FastTree2[73] was completed with the FT_pipe script, provided in Supplemental Information,
along with the initial fasta file containing all sequences and the newick tree output file.
Node labels were changed with Newick Utilities[71] *nw_rename* and visualization was
performed with Archaeopteryx[72].
571
Metagenomic recruitment
Scaffolds for the 49 SAR11 genomes, including strain LSUCC0530, used in the
phylogenomic analyses were used for competitive recruitment of metagenomic sequences
from 89 different samples. Prior to recruitment, multi-scaffold assemblies were condensed
into single scaffolds, and all single scaffolds were used to generate a BLAST database with
the command *makeblastdb* (BLAST v. 2.2.28). Metagenomic datasets from 90 different
environmental samples of various salinities from the Baltic Sea[15], Amazon River,
freshwater lakes[10,74], Lagoon Albufera[75], Columbia River [76,77], Lake Lanier[78], Lake
Michigan[79], Feitsui Reservoir[80], and GOS[81] were downloaded from the Short Read Archive
(SRA) and European Nucleotide Archive (ENA) using *ftp* and SRA toolkit (v.2.8.2-1),
respectively. Reads from pyrosequencing were dereplicated with the program CD-HIT(v
4.6)[82] using the *cd-hit-454* command and flag -c 0.95. Illumina sequence data was converted
from fastq to fasta files with the FastX toolkit (v. 0.0.13.2)
(http://hannonlab.cshl.edu/fastx_toolkit) using the command *fastq_to_fa*sta with the flag –
Q33. Metagenomic sequences were separated into 1 million sequence files and recruited
against the SAR11 database with blastn (v. 2.2.28) with –max_target_seqs 1. Best hits for
each metagenomic sequence were kept if the alignment length was > 90% of the median
length of all metagenomic sequences for a given sample. Metagenomic recruitment was
expressed in read density per genome using ggplot2 *geom_density*() and reads per kilobase
of genome per million mapped reads (RPKM) as previously reported[83].
592
593   *Script availability*
594   All scripts used in this work can be found on the Thrash Lab website with the manuscript
595   link at http://thethrashlab.com/publications.
596

Acknowledgements

Funding for this work was provided in part by a Louisiana Board of Regents (grant
LEQSF(2014-17)-RD-A-06) grant to JCT and the LSU Department of Biological Sciences. The
authors thank the LSU Socolofsky Microscopy Center for assistance with electron

microscopy imaging. Portions of this research were conducted with high performance
computing resources provided by Louisiana State University (http://www.hpc.lsu.edu).
603
Author Contributions
JCT designed the study; MWH conducted cultivation experiments, nucleic acid extraction,
microscopy, and iTag and metagenomics analyses; JCT and BCF assembled the genome and
conducted quality assessment; MWH, VCL, and JCT conducted comparative genomics
analyses, metabolic reconstruction, phylogenetic, and phylogenomic analyses; JCT lead
manuscript writing and all authors contributed to and reviewed the text.
610
611 **Competing Financial Interests**
612 The authors declare no competing financial interests.
613
614
615

Tables and Figures

618   **Table 1.** Compatible solute production and transport in SAR11

Osmolyte system	LSUCC0530	LD12 SAGs	HIMB114	IMCC9063	Other SAR11
Alanine synthesis	+	+	+	+	+
Di-myo-inositol phosphate synthesis	+/-	+/-	+/-	+/-	+/-
Ectoine/hydroxyectoine transport	-	-	+	+	+
Glutamate synthesis	+	+	+	+	+
Glutamate transport	-	+*	-	-	-
Glutamine synthesis	+	+	+	+	+
Glycine synthesis	+	+	+	+	+
Glycine betaine synthesis	-	+/-*	+	+	+
Glycine betaine/proline ABC transporter	-	-	+	+	+
N-∂-Acetyl-ornithine synthesis	+	+	+	+	+
Potassium transport (antiporters K03549, K03455)	+	+	-	-	-
Potassium transport (trkAH + K16052)	-	-	+	+	+
Proline synthesis	+	+	+	+	+
Proline transport (symporter)	+	+	+	+	+
Sodium transport (antiporter K03313)	-	-	+	+	+
Sodium transport (other antiporter)	+	+	-	-	-
Sorbitol synthesis	-	-	+	-	+
Taurine transport	-	-	-	-	+
TMAO synthesis	-	-	-	-	+

+ genes present; - genes absent; +/- partial pathway; *B11 LD12 SAG only. Table shows

only systems with predicted genes in at least one SAR11 genome. For a complete list of all

systems searched, see Table S1.

Figure Legends

Figure 1. Phylogeny and morphology of LSUCC0530. A. Phylogenetic position of

LSUCC0530 within the SAR11 clade using a concatenation of 83 single copy marker genes

and 23,356 amino acid positions. The tree was inferred using RAxML with 100 bootstrap

runs. Values at the nodes indicate 100% bootstrap support unless otherwise noted. The

tree was rooted on HIMB59. Subclades are indicated on the right, and LD12 microclusters

are indicated A-C. Asterisk- we have included AAA028-C07 in microcluster A, even though

it was excluded previously[18]. B. Scanning electron micrograph of LSUCC0530 cells at

35,000x magnification. Scale bar represents 0.5 μm. C. Transmission electron micrograph

of a LSUCC0530 cell thin section. Scale bar represents 0.5 μm.

Figure 2. Circular diagram of the LSUCC0530 genome. Rings indicate, from outside to the
center: Position relative to the replication start site; Forward strand genes, colored by COG
category; Reverse strand genes, colored by COG category; RNA genes (tRNAs green, rRNAs
red, other RNAs black); GC content; GC skew.
641
Figure 3. Metabolic reconstruction of LSUCC0530. Solid boxes indicate recovered genes,
dashed boxes indicate missing genes. ABC transporters are depicted with circles indicating
subunits. Symporters and antiporters are depicted with ovals. Two component systems are
depicted with large and small rectangles. The C4 TRAP transporter is depicted separately.
Major subsystems are colored for easier identification. Multiple arrows indicate all genes
present in a given pathway. Numbers identify genes according to the key in Table S1.
Question marks indicate a single missing gene in an otherwise complete pathway. Blue fill
indicates genes with no orthologs outside of the LD12 subclade.
650
651   **Figure 4.** Growth rates for LSUCC0530 according to salinity (A) and temperature (B).
652   Points indicate individual biological replicates, calculated from corresponding growth
653   curves in Figure S8.
654
Figure 5. Results of competitive metagenomic recruitment to all genomes depicted in Fig.
1. A. 95% identity RPKM values for LD12 genomes only, by site, with data aggregated for all
genomes in microclusters A-C, according to the key. Colors indicate broad environmental
categories. B-D. Density plots of metagenomic sequence recruitment according to percent
identity of the best hit. Data was aggregated according to the key, with LD12 separated as
in A, HIMB59 separately depicted as "V", and all other SAR11 genomes together as "Other."
B. Two sites where LSUCC0530 dominated recruitment with high percent identity. C. Two
sites where the LSUCC0530 only recruited sequences at ~90% identity and no other group
had high recruitment at higher identity. D. Two sites where other LD12 microclusters
showed similar patterns as LSUCC0530 in B and C.
665
666

References Cited

1   Morris, R. M. *et al.* SAR11 clade dominates ocean surface bacterioplankton
communities. *Nature* **420**, 806-810, (2002).
2   Schattenhofer, M. *et al.* Latitudinal distribution of prokaryotic picoplankton
populations in the Atlantic Ocean. *Environmental Microbiology* **11**, 2078-2093,
(2009).
3   Carlson, C. A. *et al.* Seasonal dynamics of SAR11 populations in the euphotic and
mesopelagic zones of the northwestern Sargasso Sea. *ISME J.* **3**, 283-295, (2009).
4   Giovannoni, S. J. & Vergin, K. L. Seasonality in ocean microbial communities. *Science*
335, 671-676, (2012).
5   Thrash, J. C. *et al.* Single-cell enabled comparative genomics of a deep ocean SAR11
bathytype. *ISME J.* **8**, 1440-1451, (2014).
6   Tsementzi, D. *et al.* SAR11 bacteria linked to ocean anoxia and nitrogen loss. *Nature*
536, 179-183, (2016).
7   Luo, H., Csűros, M., Hughes, A. L. & Moran, M. Evolution of Divergent Life History
Strategies in Marine Alphaproteobacteria. *mBio* **4**, e00373-00313, (2013).
8   Logares, R., Brate, J., Heinrich, F., Shalchian-Tabrizi, K. & Bertilsson, S. Infrequent
Transitions between Saline and Fresh Waters in One of the Most Abundant
Microbial Lineages (SAR11). *Mol. Biol. Evol.* **27**, 347-357, (2010).
9   Salcher, M. M., Pernthaler, J. & Posch, T. Seasonal bloom dynamics and
ecophysiology of the freshwater sister clade of SAR11 bacteria 'that rule the waves'
(LD12). *ISME J.* **5**, 1242-1252, (2011).
10   Eiler, A. *et al.* Productivity and salinity structuring of the microplankton revealed by
comparative freshwater metagenomics. *Environmental Microbiology* **16**, 2682-2698,
(2014).
11   Cabello-Yeves, P. J. *et al.* Genomes of novel microbial lineages assembled from the
sub-ice waters of Lake Baikal. *Appl. Environ. Microbiol.* **AOP**, (2017).
12   Newton, R. J., Jones, S. E., Eiler, A., McMahon, K. D. & Bertilsson, S. A Guide to the
Natural History of Freshwater Lake Bacteria. *Microbiol. Mol. Biol. Rev.* **75**, 14-49,
(2011).
13   Garcia, S. L. *et al.* Contrasting patterns of genome-level diversity across distinct co-
occurring bacterial populations. *bioRxiv*, (2017).
14   Henson, M. W. *et al.* Artificial Seawater Media Facilitate Cultivating Members of the
Microbial Majority from the Gulf of Mexico. *mSphere* **1**, e00028-00016.
doi:00010.01128/mSphere.00028-00016., (2016).
15   Dupont, C. L. *et al.* Functional Tradeoffs Underpin Salinity-Driven Divergence in
Microbial Community Composition. *PLOS ONE* **9**, e89549, (2014).
16   Logares, R. *et al.* Infrequent marine–freshwater transitions in the microbial world.
Trends in Microbiology **17**, 414-422, (2009).
17   Eiler, A. *et al.* Tuning fresh: radiation through rewiring of central metabolism in
streamlined bacteria. *ISME J.* **10**, 1902-1914, (2016).
18   Zaremba-Niedzwiedzka, K. *et al.* Single-cell genomics reveal low recombination
frequencies in freshwater bacteria of the SAR11 clade. *Genome Biol* **14**, 1-14, (2013).

19 Grote, J. *et al.* Streamlining and Core Genome Conservation among Highly Divergent
Members of the SAR11 Clade. *mBio* **3**, e00252-00212, (2012).
20 Giovannoni, S. J. *et al.* Genome streamlining in a cosmopolitan oceanic bacterium.
Science **309**, 1242-1245, (2005).
21 Giovannoni, S. J. *et al.* Proteorhodopsin in the ubiquitous marine bacterium SAR11.
Nature **438**, 82-85, (2005).
22 Tripp, H. J. *et al.* SAR11 marine bacteria require exogenous reduced sulphur for
growth. *Nature* **452**, 741-744, (2008).
23 Schwalbach, M. S., Tripp, H. J., Steindler, L., Smith, D. P. & Giovannoni, S. J. The
presence of the glycolysis operon in SAR11 genomes is positively correlated with
ocean productivity. *Environmental Microbiology* **12**, 490-500, (2010).
24 Sun, J. *et al.* One Carbon Metabolism in SAR11 Pelagic Marine Bacteria. *PLOS ONE* **6**,
e23973, (2011).
25 Carini, P., Steindler, L., Beszteri, S. & Giovannoni, S. J. Nutrient requirements for
growth of the extreme oligotroph 'Candidatus Pelagibacter ubique' HTCC1062 on a
defined medium. *ISME J.* **7**, 592-602, (2013).
26 Carini, P. *et al.* Discovery of a SAR11 growth requirement for thiamin's pyrimidine
precursor and its distribution in the Sargasso Sea. *ISME J.* **8**, 1727-1738, (2014).
27 Smith, D. P. *et al.* Proteome Remodeling in Response to Sulfur Limitation in
"Candidatus Pelagibacter ubique". *mSystems* **1**, 16, (2016).
28 Smith, D. P. *et al.* Proteomic and Transcriptomic Analyses of "Candidatus
Pelagibacter ubique" Describe the First PII-Independent Response to Nitrogen
Limitation in a Free-Living Alphaproteobacterium. *mBio* **4**, 12, (2013).
29 Steindler, L., Schwalbach, M. S., Smith, D. P., Chan, F. & Giovannoni, S. J. Energy
Starved Candidatus Pelagibacter Ubique Substitutes Light-Mediated ATP Production
for Endogenous Carbon Respiration. *PLOS ONE* **6**, e19725, (2011).
30 Zhao, X. *et al.* Three-Dimensional Structure of the Ultraoligotrophic Marine
Bacterium "Candidatus Pelagibacter ubique". *Appl. Environ. Microbiol.* **83**, 16,
(2017).
31 Zhao, Y. *et al.* Abundant SAR11 viruses in the ocean. *Nature* **494**, 357-360, (2013).
32 Konstantinidis, K. T. & Tiedje, J. M. Prokaryotic taxonomy and phylogeny in the
genomic era: advancements and challenges ahead. *Current Opinion in Microbiology*
10, 504-509, (2007).
33 Rappé, M. S., Connon, S. A., Vergin, K. L. & Giovannoni, S. J. Cultivation of the
ubiquitous SAR11 marine bacterioplankton clade. *Nature* **418**, 630-633, (2002).
34 Wilhelm, L. J., Tripp, J. H., Givan, S. A., Smith, D. P. & Giovannoni, S. J. Natural
variation in SAR11 marine bacterioplankton genomes inferred from metagenomic
data. *Biology Direct* **2**, 1-19, (2007).
35 Rodriguez-Valera, F. *et al.* Explaining microbial population genomics through phage
predation. *Nat Rev Micro* **7**, 828-836, (2009).
36 Anstrom, D. M., Kallio, K. & Remington, J. S. Structure of the Escherichia coli malate
synthase G:pyruvate:acetyl‐coenzyme A abortive ternary complex at 1.95 Å
resolution. *Protein Science* **12**, 1822-1832, (2003).
37 Kappler, U. Bacterial sulfite-oxidizing enzymes. *Biochimica et Biophysica Acta (BBA)
- Bioenergetics* **1807**, 1-10, (2011).

38   Denger, K., Weinitschke, S., Smits, T. H. M., Schleheck, D. & Cook, A. M. Bacterial
sulfite dehydrogenases in organotrophic metabolism: separation and identification
in Cupriavidus necator H16 and in Delftia acidovorans SPH-1. *Microbiology* **154**,
256-263, (2008).
39   Béjà, O., Spudich, E. N., Spudich, J. L., Leclerc, M. & DeLong, E. F. Proteorhodopsin
phototrophy in the ocean. *Nature* **411**, 786-789, (2001).
40   Konstantinidis, K. T. & Tiedje, J. M. Genomic insights that advance the species
definition for prokaryotes. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 2567-2572, (2005).
41   Jimenez-Infante, F. *et al.* Genomic Characterization of Two Novel SAR11 Isolates
From the Red Sea, Including the First Strain of the SAR11 Ib clade. *FEMS Microbiol*
Ecol, (2017).
42   Song, J., Oh, H.-M. & Cho, J.-C. Improved culturability of SAR11 strains in dilution-to-
extinction culturing from the East Sea, West Pacific Ocean. *FEMS Microbiology*
Letters **295**, 141-147, (2009).
43   Stingl, U., Tripp, H. J. & Giovannoni, S. J. Improvements of high-throughput culturing
yielded novel SAR11 strains and other abundant marine bacteria from the Oregon
coast and the Bermuda Atlantic Time Series study site. *ISME J.* **1**, 361-371, (2007).
44   Piwosz, K., Salcher, M. M. & and ..., Z.-M. Seasonal dynamics and activity of typical
freshwater bacteria in brackish waters of the Gulf of Gdańsk. *Limnology and ...*,
(2013).
45   Herlemann, D., Woelk, J., Labrenz, M. & Jürgens, K. Diversity and abundance of
"Pelagibacterales" (SAR11) in the Baltic Sea salinity gradient. *Syst. Appl. Microbiol.*
37, 601-604, (2014).
46   Dattananda, C. S. & Gowrishankar, J. Osmoregulation in Escherichia coli:
complementation analysis and gene-protein relationships in the proU locus. *J.*
Bacteriol. **171**, 1915-1922, (1989).
47   Empadinhas, N. & da Costa, M. Osmoadaptation mechanisms in prokaryotes:
distribution of compatible solutes. *International Microbiology*, (2008).
48   Sun, J. *et al.* The abundant marine bacterium Pelagibacter simultaneously
catabolizes dimethylsulfoniopropionate to the gases dimethylsulfide and
methanethiol. *Nature Microbiology* **1**, (2016).
49   Parada, A. E., Needham, D. M. & Fuhrman, J. A. Every base matters: assessing small
subunit rRNA primers for marine microbiomes with mock communities, time series
and global field samples. *Environmental Microbiology* **18**, (2015).
50   Schloss, P. D. *et al.* Introducing mothur: Open-Source, Platform-Independent,
Community-Supported Software for Describing and Comparing Microbial
Communities. *Appl. Environ. Microbiol.* **75**, 7537-7541, (2009).
51   Pruesse, E. *et al.* SILVA: a comprehensive online resource for quality checked and
aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Research*
35, 7188-7196, (2007).
52   McMurdie, P. J. & Holmes, S. phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. *PLOS ONE* **8**, e61217, (2013).
53   Wickham, H. ggplot2. *Wiley Interdisciplinary Reviews: Computational Statistics* **3**,
180-185, (2011).
54   Glenn, T. C. *et al.* Adapterama I: Universal stubs and primers for thousands of dual-
indexed Illumina libraries (iTru & iNext). *bioRxiv*, (2016).

55   Bankevich, A. *et al.* SPAdes: A New Genome Assembly Algorithm and Its Applications
803   to Single-Cell Sequencing. *Journal of Computational Biology* **19**, 455-477, (2012).

804   56   Hunt, M. *et al.* REAPR: a universal tool for genome assembly evaluation. *Genome Biol*
805   **14**, 1-10, (2013).

806   57   Boetzer, M., Henkel, C. V., Jansen, H. J., Butler, D. & Pirovano, W. Scaffolding pre-
807   assembled contigs using SSPACE. *Bioinformatics* **27**, 578-579, (2011).

808   58   Walker, B. J. *et al.* Pilon: An Integrated Tool for Comprehensive Microbial Variant
809   Detection and Genome Assembly Improvement. *PLOS ONE* **9**, (2014).

810   59   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler
811   transform. *Bioinformatics* **25**, 1754-1760, (2009).

812   60   Brown, C. T. *et al.* Unusual biology across a group comprising more than 15% of
813   domain Bacteria. *Nature* **523**, 208-211, (2015).

814   61   Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:
815   assessing the quality of microbial genomes recovered from isolates, single cells, and
816   metagenomes. *Genome Research* **25**, 1043-1055, (2015).

817   62   Markowitz, V. M. *et al.* IMG 4 version of the integrated microbial genomes
818   comparative analysis system. *Nucleic Acids Research* **42**, D560-567, (2014).

819   63   Contreras-Moreira, B. & Vinuesa, P. GET_HOMOLOGUES, a versatile software
820   package for scalable and robust microbial pangenome analysis. *Appl. Environ.*
821   *Microbiol.* **79**, 7696-7701, (2013).

822   64   Dick, G. J. *et al.* Community-wide analysis of microbial genome sequence signatures.
823   *Genome Biol* **10**, R85, (2009).

824   65   Goris, J. *et al.* DNA–DNA hybridization values and their relationship to whole-
825   genome sequence similarities. *IJSEM* **57**, 81-91, (2007).

826   66   Roux, S., Enault, F., Hurwitz, B. L. & Sullivan, M. B. VirSorter: mining viral signal from
827   microbial genomic data. *PeerJ* **3**, (2015).

828   67   Arndt, D. *et al.* PHASTER: a better, faster version of the PHAST phage search tool.
829   *Nucleic Acids Research* **44**, (2016).

830   68   Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high
831   throughput. *Nucleic Acids Research* **32**, 1792-1797, (2004).

832   69   Castresana, J. Selection of conserved blocks from multiple alignments for their use in
833   phylogenetic analysis. *Mol. Biol. Evol.* **17**, 540-552, (2000).

834   70   Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
835   large phylogenies. *Bioinformatics* **30**, 1312-1313, (2014).

836   71   Junier, T. & Zdobnov, E. M. The Newick utilities: high-throughput phylogenetic tree
837   processing in the Unix shell. *Bioinformatics* **26**, 1669-1670, (2010).

838   72   Han, M. V. & Zmasek, C. M. phyloXML: XML for evolutionary biology and comparative
839   genomics. *BMC Bioinformatics* **10**, 1-6, (2009).

840   73   Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2--approximately maximum-
841   likelihood trees for large alignments. *PLOS ONE* **5**, e9490, (2010).

842   74   Martinez-Garcia, M. *et al.* High-throughput single-cell sequencing identifies
843   photoheterotrophs and chemoautotrophs in freshwater bacterioplankton. *ISME J.* **6**,
844   113-123, (2011).

75   Ghai, R. *et al.* Metagenomes of Mediterranean Coastal Lagoons. *Sci. Rep.* **2**, (2012).

76   Smith, M. W., Allen, L., Allen, A. E., Herfort, L. & Simon, H. M. Contrasting genomic
properties of free-living and particle-attached microbial assemblages within a
coastal ecosystem. *Front. Microbiol.* **4**, (2013).
77   Fortunato, C. S. & Crump, B. C. Microbial Gene Abundance and Expression Patterns
across a River to Ocean Salinity Gradient. *PLOS ONE* **10**, (2015).
78   Oh, S. *et al.* Metagenomic Insights into the Evolution, Function, and Complexity of
the Planktonic Microbial Community of Lake Lanier, a Temperate Freshwater
Ecosystem. *Appl. Environ. Microbiol.* **77**, 6000-6011, (2011).
79   Denef, V. J., Mueller, R. S., Chiang, E., Liebig, J. R. & Vanderploeg, H. A. Chloroflexi
CL500-11 Populations That Predominate Deep-Lake Hypolimnion Bacterioplankton
Rely on Nitrogen-Rich Dissolved Organic Matter Metabolism and C1 Compound
Oxidation. *Appl. Environ. Microbiol.* **82**, 1423-1432, (2016).
80   Tseng, C.-H. *et al.* Microbial and viral metagenomes of a subtropical freshwater
reservoir subject to climatic disturbances. *ISME J.* **7**, 2374-2386, (2013).
81   Rusch, D. B. *et al.* The Sorcerer II Global Ocean Sampling Expedition: Northwest
Atlantic through Eastern Tropical Pacific. *Plos Biol* **5**, e77, (2007).
82   Fu, L., Niu, B., Zhu, Z., Wu, S. & Li, W. CD-HIT: accelerated for clustering the next-
generation sequencing data. *Bioinformatics* **28**, 3150-3152, (2012).
83   Thrash, J. *et al.* Metabolic Roles of Uncultivated Bacterioplankton Lineages in the
Northern Gulf of Mexico "Dead Zone". *mBio* **8**, (2017).
866

A.

Candidatus Pelagibacter sp IMCC9063 — IIIa
Rickettsiales sp HIMB114

alpha proteobacterium SCGC AAA280-B11
Candidatus Fonsibacter ubiquis LSUCC0530
alpha proteobacterium SCGC AAA028-C07
alpha proteobacterium SCGC AAA028-D10 — A*
alpha proteobacterium SCGC AAA023-L09
alpha proteobacterium SCGC AAA024-N17
alpha proteobacterium SCGC AAA280-P20 — C
alpha proteobacterium SCGC AAA487-M09
alpha proteobacterium SCGC AAA027-J10 — B
alpha proteobacterium SCGC AAA027-C06
alpha proteobacterium SCGC AAA027-L15

LD12/IIIb

Candidatus Pelagibacter ubique HIMB058 — II
SAR11 cluster bacterium JGI GoM 1470m 185 D04
SAR11 cluster bacterium JGI GoM 375m 188 H08
alpha proteobacterium SCGC AAA240-E13
alpha proteobacterium SCGC AAA288-E13
alpha proteobacterium SCGC AAA288-N07
SAR11 cluster bacterium JGI GoM 375m 188 D08
alpha proteobacterium SCGC AAA288-G21
SAR11 cluster bacterium JGI ETNP 125m 186 C10
SAR11 cluster bacterium JGI ETNP 125m 186 G05

Ic

SAR11 cluster bacterium JGI GoM 1470m 185 E04
SAR11 cluster bacterium JGI GoM 375m 188 A02
SAR11 cluster bacterium JGI GoM 375m 188 E04
Candidatus Pelagibacter ubique SAR11 HTCC9565
Candidatus Pelagibacter ubique HTCC1040
Candidatus Pelagibacter ubique HTCC1016
Candidatus Pelagibacter ubique HTCC1002
Candidatus Pelagibacter ubique HTCC1062
Candidatus Pelagibacter ubique SAR11 HTCC1013
Alphaproteobacteria sp SAR11 HIMB5
SAR11 cluster bacterium JGI GoM 1m 182 G01
SAR11 cluster bacterium JGI GoM 1m 182 D02
SAR11 cluster bacterium JGI GoM 1m 182 F07
Candidatus Pelagibacter sp HIMB1321
Candidatus Pelagibacter-like SAR11 HIMB4
Candidatus Pelagibacter ubique SAR11 HTCC8051
Candidatus Pelagibacter ubique SAR11 HTCC9022
Candidatus Pelagibacter ubique HTCC7214
Candidatus Pelagibacter sp HTCC7211
Candidatus Pelagibacter ubique HTCC7217
Candidatus Pelagibacter sp HIMB122
Candidatus Pelagibacter-like SAR11 HIMB140
Candidatus Pelagibacter-like SAR11 HIMB083
SAR11 cluster bacterium JGI ETNP 60m 184 G07
SAR11 cluster bacterium JGI GoM 1m 182 A11

Ia

0.1

B.

SEI 10kV WD8mm SS33 x35,000 0.5µm
530 2

C.

500 nm

Candidatus
Fonsibacter ubiquis
1,160,202 bp