Cultivation and genomics of the first freshwater SAR11 (LD12) isolate

Michael W. Henson[1], V. Celeste Lanclos[1], Brant C. Faircloth[1,2], and J. Cameron Thrash[1,3]

1. Department of Biological Sciences, Louisiana State University, Baton Rouge, LA 70803,
U.S.A.

2. Museum of Natural History, Louisiana State University, Baton Rouge, LA 70803, U.S.A.

3. Materials and Correspondence:
J. Cameron Thrash
Department of Biological Sciences
202 Life Sciences Bldg.
Louisiana State University
Baton Rouge, LA 70803
thrashc@lsu.edu
225-578-8210

Key Words: LD12, SAR11, freshwater microbiology, genomics, physiology

Abstract
Evolutionary transitions between fresh and salt water happen infrequently among
bacterioplankton. Within the ubiquitous and highly abundant heterotrophic
Alphaproteobacteria order *Pelagibacterales* (SAR11), most members live in marine
habitats, but the LD12 subclade has evolved as a unique freshwater lineage. LD12 cells
occur as some of the most dominant freshwater bacterioplankton, yet this group has
remained elusive to cultivation, hampering a more thorough understanding of its biology.
Here, we report the first successful isolation of an LD12 representative, strain LSUCC0530,
using high throughput dilution to extinction cultivation methods, and its complete genome
sequence. Growth experiments corroborate ecological data suggesting active populations
of LD12 in brackish water up to salinities of ~5. LSUCC0530 has the smallest closed
genome thus far reported for a SAR11 strain (1.16 Mbp). The genome affirms many
previous metabolic predictions from cultivation-independent analyses, like a complete
Embden-Meyerhof-Parnas glycolysis pathway, but also provides novel insights, such as the
first isocitrate dehydrogenase in LD12, a likely homologous recombination of malate
synthase from outside of the SAR11 clade, and analogous substitutions of ion transporters
with others that occur throughout the rest of the SAR11 clade. Growth data support
metagenomic recruitment results suggesting temperature-based ecotype diversification
within LD12. Key gene losses for osmolyte uptake provide a succinct hypothesis for the
evolutionary transition of LD12 from salt to fresh water. For strain LSUCC0530, we
propose the provisional nomenclature *Candidatus* Fonsibacter ubiquis.
43

Introduction

Bacterioplankton in the SAR11 clade of *Alphaproteobacteria* are dominant heterotrophs in marine and freshwater systems. In the oceans, SAR11 can represent 25-50% of total planktonic cells (Morris *et al.*, 2002, Schattenhofer *et al.*, 2009). Numerous subclades with unique spatio-temporal distributions comprise SAR11 (Carlson *et al.*, 2009, Giovannoni and Vergin, 2012, Morris *et al.*, 2002, Thrash *et al.*, 2014). At least nine subclades defined via 16S rRNA gene sequences occupy marine niches (Giovannoni and Vergin, 2012), and more likely exist (Tsementzi *et al.*, 2016). However, in spite of its global distribution (Morris *et al.*, 2002), massive predicted population size of 10^{28} cells (Morris *et al.*, 2002), and an estimated divergence time from its last common ancestor of 1.1 billion years ago (Luo *et al.*), the bulk of existing evidence suggests that SAR11 has only successfully colonized freshwater environments once in its natural history (Eiler *et al.*, 2014, Logares *et al.*, 2010, Salcher *et al.*, 2011). Traditionally, all known freshwater SAR11 belong to subclade IIIb, a.k.a. LD12. However, a recent report challenges this assertion: a genome sister to subclade I was recovered in Lake Baikal metagenomic data (Cabello-Yeves *et al.*, 2018). Regardless, the limited evolutionary diversification into less saline habitats has not prevented LD12 from achieving prominence in the ecosystems it inhabits. In many lotic and lentic environments, LD12 occupies similar relative abundances as its marine cousins (Dupont *et al.*, 2014, Garcia *et al.*, 2017, Henson *et al.*, 2016, Newton *et al.*, 2011, Salcher *et al.*, 2011). Study of LD12 is important to understand SAR11 evolution, specifically, and how successful transitions between marine and freshwater environments occur in bacterioplankton (Logares *et al.*, 2009), more generally.

Ecological, functional, and sequence-based inference from single amplified genomes (SAGs) and metagenomes support the hypothesis that LD12 bacterioplankton evolved from a genome-streamlined marine ancestor (Eiler *et al.*, 2016, Luo *et al.*, 2013, Salcher *et al.*, 2011, Zaremba-Niedzwiedzka *et al.*, 2013). As such, they share many of the same characteristics as marine SAR11, such as small cell volumes; adaptation to oligotrophic habitats; small, streamlined genomes; an obligate aerobic chemoorganoheterotrophic lifestyle with limited metabolic flexibility; preference for small molecular weight compounds like carboxylic and amino acids as carbon/energy sources; and auxotrophies for some amino acids and vitamins (Dupont *et al.*, 2014, Eiler *et al.*, 2014, Eiler *et al.*, 2016, Giovannoni *et al.*, 2005a, Giovannoni *et al.*, 2005b, Grote *et al.*, 2012, Salcher *et al.*, 2011, Schwalbach *et al.*, 2010, Sun *et al.*, 2011, Thrash *et al.*, 2014, Tripp *et al.*, 2008, Zaremba-Niedzwiedzka *et al.*, 2013). Previous research suggests that LD12 differ from their marine counterparts in specific elements of metabolic potential that indicate a greater emphasis on production, rather than uptake, of osmolytes, and important metabolic changes related to energy production (Dupont *et al.*, 2014, Eiler *et al.*, 2016). For example, metagenomic population data showed a correlation between decreased salinity and greater proportion of the Embden-Meyerhof-Parnass (EMP) vs. Entner-Doudoroff (ED) glycolysis pathways (Dupont *et al.*, 2014). Comparative genomic analyses of SAGs from different SAR11 strains

concurred: LD12 genomes contained the EMP pathway that is not found in most marine
SAR11 (Eiler *et al.*, 2016, Grote *et al.*, 2012). SAG data also suggested that LD12 lacks the
glyoxylate shunt and some single carbon (C1) metabolism (Eiler *et al.*, 2016).
Despite what has been learned from cultivation-independent methods, the lack of
cultured LD12 representatives has hampered a more detailed exploration of the group.
Potential ecotypes within LD12 have been identified (Zaremba-Niedzwiedzka *et al.*, 2013),
and their population dynamics recently described with five-year time series data in
freshwater lakes (Garcia *et al.*, 2017). However, we cannot delineate what distinguishes
ecotypes without better physiological and genomic data. Similarly, interpreting data on the
ecological distribution of LD12 remains challenging without information on growth
tolerances and optima for salinity and temperature. We also don't understand whether a
connection exists between more efficient energy production through EMP-based glycolysis
and the freshwater lifestyle, or what other adaptations might explain LD12 evolution away
from salt water.
The next steps in translating 'omics-based predictions into measured data for
integration with ecosystem models require living experimental subjects. For example,
cultures of marine SAR11, such as HTCC1062 and HTCC7211, have facilitated testing of
metabolism and growth (Carini *et al.*, Carini *et al.*, Giovannoni *et al.*, 2005a, Schwalbach *et
al.*, 2010, Smith *et al.*, Smith *et al.*, Steindler *et al.*, 2011, Tripp *et al.*, 2008), structural
organization (Zhao *et al.*, 2017), and virus-host interactions (Zhao *et al.*, 2013). We need
cultivated representatives to provide this kind of understanding of other important
bacterioplankton like LD12. In service of this goal, we pursued a systematic high
throughput cultivation effort from coastal regions in the northern Gulf of Mexico. Here, we
report the first isolation of a member of the LD12 clade, strain LSUCC0530, its complete
genome sequence, preliminary physiological data, and an examination of the ecological
distribution of this organism compared with other LD12 clade members. Our results
provide evidence for temperature-dependent ecotype diversification within LD12 and a
hypothesis about the evolutionary trajectory that led to LD12 colonization of fresh water.
112

Results
Isolation and growth of LSUCC0530
We isolated strain LSUCC0530 as part of a series of high throughput dilution-to-extinction
cultivation experiments (HTC) conducted with inoculation water obtained across the
southern Louisiana coast. The particular experiment that yielded LSUCC0530 utilized
surface water from the coastal lagoon of Lake Borgne, 39 km southeast of New Orleans. At
the time of collection, the surface water had a salinity of 2.39 and was 30.5°C. Subsequent
community analysis demonstrated that LD12-type organisms represented 8.7% of the
bacterioplankton population in the inoculum (see below). We diluted 2.7 μm-filtered Lake
Borgne surface water for inoculation to an estimated 2 cells well^{-1} into our JW5 medium
(salinity 1.45, Table S1), and incubated cultures for 30 days at 23°C (room temperature in

the Thrash Lab). LSUCC0530 grew slowly, reaching 8.6 x 10^5 cells mL^{-1} after 24 days. It had
a low fluorescence/low side scatter flow cytometric signature characteristic of other
SAR11 cells (Fig. S1).
127
Taxonomy and morphology
The LSUCC0530 16S rRNA gene sequence had > 99% identity with eight SAG sequences
from members of LD12 (Zaremba-Niedzwiedzka *et al.*, 2013) and only 91% and 92%
sequence identity with sequences from the sister subclade IIIa representatives IMCC9063
and HIMB114, respectively. Of note for taxonomic incorporation of genomic data (Chun *et
al.*, 2018), the genome-generated 16S rRNA gene sequence matched the PCR-generated 16*S*
rRNA gene sequence (GenBank accession number KY290650.1) at 1316 of 1320 sites. Two
of these mismatches were single bp gaps the genome sequence, and three of the four
mismatches occurred in the first 20 bp of the PCR-generated sequence. LSUCC0530 also
had average amino acid identities (AAI) of 58.1% and 59.7% with HIMB114 and IMCC9063,
respectively (Table S1). In contrast, AAI between LSUCC0530 and the LD12 SAGs ranged
from 81.9% to 88.0%, corroborating that LSUCC0530 belongs to separate species from
subclade IIIa (Konstantinidis and Tiedje, 2007). Phylogenetic inference using 16S rRNA
genes placed LSUCC0530 within the Order *Pelagibacteriales* (a.k.a. SAR11 (Thrash *et al.*,
2014)), in monophyly with the original LD12 clone library sequence (Fig. S2).
Phylogenomic inference using 83 single copy marker genes from 49 SAR11 genomes also
strongly supported placement of LSUCC0530 within LD12 as an early diverging member,
with only the SAG AAA280-B11 from Lake Damariscotta diverging earlier (Fig. 1A). This
tree also recovered the three microclusters within LD12 (A, B, C) observed previously
(Zaremba-Niedzwiedzka *et al.*, 2013) (Fig. 1A). Cells of strain LSUCC0530 were small,
curved rods, approximately 1 μm x 0.1 μm (Fig. 1B, C), showing conserved morphology
with the distantly related SAR11 strain HTCC1062 in subclade Ia (Rappé *et al.*, 2002, Zhao
et al., 2017).
151
Genome characteristics
Genome assembly resulted in a single, circularized scaffold that we tested for completeness
using multiple tools (Supplemental Information). The completed genome is 1,160,202 bp
with a GC content of 29.02% and 1271 predicted genes (Fig. 2). There are 1231 putative
protein coding genes and 40 RNA genes, including one each of the 5S, 16S, and 23S rRNA
genes. We found no genomic evidence of lysogenic bacteriophage or a CRISPR-*cas* system.
LSUCC0530 has the characteristics of genome streamlining previously reported for other
SAR11 genomes, with an estimated coding percentage of 96.36% and the smallest complete
SAR11 genome reported thus far (Giovannoni *et al.*, 2005b, Grote *et al.*, 2012). Despite its
smaller overall genome size, intergenic spacer regions are not significantly smaller than
those of the sister subclade IIIa (Fig. S3). All LD12 genomes had much higher AAI and
synteny with each other than with the subclade IIIa (Table S1), mirroring their

phylogenomic relationships (Fig. 1A). However, within the LD12 clade, AAI and synteny
percentages did not clearly delineate the same patterns we observed in the phylogenomic
tree, which may have resulted from the LD12 SAGs being fragmented and incomplete.
The LSUCC0530 genome has the conserved architecture for hypervariable region 2
(HVR2) that has been observed in most other complete SAR11 genomes (Grote *et al.*, 2012,
Rodriguez-Valera *et al.*, 2009, Wilhelm *et al.*, 2007) and other LD12 SAGs (Zaremba-
Niedzwiedzka *et al.*, 2013), between the 16S rRNA-tRNA-tRNA-23S rRNA operon and the
5S rRNA gene operon. This region is 54,755 bp and has a strong variation in GC content
from neighboring parts of the genome (Fig. 2). Metagenomic recruitment also verified this
region as hypervariable based on the lack of matching sequences in community data (Fig.
S4). The predicted gene annotations in HVR2 resemble those in the same location in other
SAR11 genomes (Grote *et al.*, 2012, Zaremba-Niedzwiedzka *et al.*, 2013), namely genes
likely functioning in membrane modification such as glycosyltransferases, epimerases,
transaminases, and methyltransferases; as well as hypothetical proteins.
178
Metabolic reconstruction
Like most other SAR11s (Eiler *et al.*, 2016, Grote *et al.*, 2012, Thrash *et al.*, 2014), the
LSUCC0530 genome encodes for an obligate aerobic lifestyle, with a complete oxidative
phosphorylation pathway including a heme/copper-type cytochrome c oxidase, a proton-
translocating NADH dehydrogenase, and a proton-translocating ATP synthase (Fig. 3). As
previously detected in LD12 SAGs (Eiler *et al.*, 2016), the LSUCC0530 genome encodes a
complete EMP glycolysis pathway, including phosphofructokinase and pyruvate
dehydrogenase, and complete gluconeogenesis through nucleotide sugar production (Fig.
3). We recovered genes for the pentose phosphate pathway and a predicted fructokinase
gene for conversion of D-fructose to ß-D-fructose-6-P. As expected based on other SAR11
genomes, we did not find annotated genes for utilization of allose, fucose, fuculose,
galactose, maltose, mannose, rhamnose, sucrose, starch, trehalose, or xylose. LSUCC0530
has a complete TCA cycle and glycolate oxidase, and like other SAR11s, the non-mevalonate
pathway for isopentenyl diphosphate (Fig. 3).
We recovered the first isocitrate lyase in an LD12 genome, conferring a complete
glyoxylate bypass in combination with malate synthase. The isocitrate lyase shares its
ancestry with the majority of the SAR11 clade except subclade IIIa, where it appears to
have been lost (Eiler *et al.*, 2016, Grote *et al.*, 2012) (Fig. S5). However, the LSUCC0530
malate synthase shared with other LD12 SAGs belongs to isoform A, whereas all other
SAR11 clades contain isoform G (Anstrom *et al.*, 2003) (Figs. 3; S6). Comparison of gene
neighborhoods in a variety of SAR11 genomes revealed that both isoforms occur in the
same location (Fig. S7).
We predict production of eighteen amino acids (Fig. 3) and partial synthesis
pathways for three more (histidine, cysteine, and methionine), as well as the ability to
interconvert, but not synthesize *de novo*, phenylalanine and tyrosine. The LSUCC0530

genome has lost some of the C1 and methylated compound pathways found in other
SAR11s (Sun *et al.*, 2011). For example, as in the LD12 SAGs (Eiler *et al.*, 2016), LSUCC0530
does not have DMSP, methylamine, or glycine-betaine metabolism. However, it retains
tetrahydrafolate metabolism, formate oxidation, and the glycine cleavage pathway (Fig. 3).
The genome has the same PII nitrogen response system and *amtB* ammonium
transporter found in other SAR11s and is predicted to utilize ammonia as a nitrogen source
in amino acid synthesis (Fig. 3). The genome also encodes a putative thiosulfate/3-
mercaptopyruvate sulfurtransferase that may be involved in sulfur and thiocyanate
reactions (Fig. 3). A possible product of this enzyme is sulfite, which could be further
oxidized to sulfate in the periplasm via putative *sorAB* genes. This gene pair occurs
infrequently in the SAR11 clade, but sulfite oxidation may serve, in some strains, as a
means to generate additional proton motive force (Denger *et al.*, 2008, Kappler, 2011). This
is similar to the function conferred by proteorhodopsin (Béjà *et al.*, 2001, Giovannoni *et al.*,
2005a, Steindler *et al.*, 2011), which the LSUCC0530 genome also contains. Also like other
SAR11s, a complete assimilatory sulfate reduction pathway is missing, making LSUCC0530
dependent on reduced sulfur compounds for cellular sulfur requirements (Tripp *et al.*,
2008). In our media, this was supplied in the form of sulfur-containing amino acids (Table
S1). The genome encodes partial pathways for incorporation of sulfide into cysteine and
methionine but is missing predicted serine O-acetyltransferase and homoserine O-
succinyltransferase genes (Fig. 3).
In addition to the PII nitrogen sensor, LSUCC0530 also has putative two component
systems handling phosphate limitation, pH, osmotic upshift via potassium, and redox state
(Fig. 3). The LSUCC genome encodes ABC transporters for phosphate (*ptsABCS*),
lipoprotein, and branched-chain and L amino acid uptake. We identified tripartite ATP-
independent periplasmic (TRAP) transporter genes for C4-dicarboxylate and
mannitol/chloroaromatic compounds. However, the gene cluster appears to be missing the
substrate binding subunit, and we did not find predicted genes for mannitol utilization.
In general, LSUCC0530 and the LD12 SAGs share many of the same pathways for
osmolyte synthesis as the subclade IIIa and other SAR11 organisms, such as N-acetyl-
ornithine, glutamate, glutamine, and proline (Table 1). In some cases, there appear to be
analogous substitutions in osmolyte transport systems. For example, LD12 genomes have
potassium antiporters, but not the *trkAH* genes for potassium transport in other SAR11s.
Similarly, a sodium/proton antiporter ortholouges cluster comprised genes exclusively from
LD12 genomes, but a separate sodium/proton antiporter occurred in all other SAR11
genomes (Table 1). Perhaps the most important difference in genomic content concerning
osmolytes occurs in two key transporter losses: LSUCC0530 and the other LD12s have no
genes for the glycine betaine/proline or ectoine/hydroxyectoine ABC transporters that are
present in HIMB114 and IMCC9063 and other subclades of SAR11 (Table 1).
242
Physiology

Under isolation conditions (23°C, JW5 medium), strain LSUCC0530 grew to a density of 2 x
10^7 cells mL^{-1} with an average growth rate of 0.52 day^{-1}. Our physiological experiments to
determine the salinity tolerance and growth optima revealed that LSUCC0530 could grow
at salinities between 0.36 and 4.7 (Fig. 4A). The optimal salinity for LSUCC0530 was at or
below the isolation medium of 1.45. Growth rates decreased between salinities of 2.9 and
4.7 (Fig. 4A), and cells died (determined by rapid loss of flow cytometric signal) in salinities
above 8 (Fig. S8). We observed growth after a very extended lag phase in a subset of
cultures at salinity 5.8, but could not always reproduce this growth in repeated
experiments (Fig. S8). Among the temperatures tested, LSUCC0530 grew optimally at 23°C,
and we observed growth up to 30°C, but not at 12°C or below, nor at 35°C or above (Fig.
4B). However, we did not observe a clear loss of signal at 4, 12, 35, or even 40°C (Fig. S8),
even after over 33 days, suggesting that LSUCC0530 can endure extended periods at these
temperatures in a non-growth state. Notably during growth experiments, one of the five
replicates during any given experiment frequently grew at a reduced rate than the others
(Fig. S8).
259
Ecology
We quantified the abundance of LD12 taxa in the coastal Louisiana ecosystems using 16S
rRNA gene sequences clustered into operational taxonomic units (OTUs). These sequences
were obtained as part of the combined sampling and HTC campaign that included the
successful isolation of LSUCC0530. Based on BLASTn sequence similarity, the
representative sequence for OTU7 and LSUCC0530 shared 100% identity over the 250bp
v4 region of the 16S rRNA gene, confirming the Mothur classification of OTU7 as belonging
to the SAR11 LD12 clade. Over the three-year sampling period, OTU7 was the fifth most
abundant OTU based on total read abundance and sixth ranked OTU based on median read
abundance. At the time of sampling for the experiment that resulted in the isolation of
LSUCC0530, OTU7 had a relative abundance of 8.7% in Lake Borgne. In some sites, such as
the Atchafalaya River Delta (salinity 0.18), LD12 represented as much as 16% of total
bacterioplankton (Fig. S9). Generally, the LD12 OTU7 relative abundance correlated
negatively with salinity. At seven sites with salinities below 6, OTU7 had a relative
abundance of 4.0% or greater with a median relative abundance of 8.8% (Fig. S9). At the
remaining ten sites above a salinity of 6 (median 19.44), OTU7 had relative abundances
around 2% or lower (Fig. S9).
The phylogenetic separation of LSUCC0530 from B11 and the IIIb A-C microclusters
raised the question of whether LSUCC0530 represents a novel ecotype with unique
spatiotemporal distribution compared to other LD12 taxa. Therefore, we explored the
distribution of LD12 genomes in 90 fresh and brackish water environments through
competitive recruitment of metagenomic sequences to all the SAR11 genomes in the
phylogenomic tree and the outgroup sequence HIMB59. We quantified relative abundance
via recruited reads (per kpb of metagenomic sequence per Mbp of genome sequence-

RPKM), and aggregated LD12 data based on the A, B, and C microclusters (Fig. 1A), the B11
SAG, and LSUCC0530. Figure 5A displays RPKM values for reads with ≥ 95% identity,
roughly corresponding to a species delineation via average nucleotide identity
(ANI)(Konstantinidis and Tiedje, 2005), but we also provide additional comparisons at ≥
85, 90, 92, 98, and 100% identity (Fig. S10).
All LD12 subgroups had their highest RPKM values in very low salinity sites,
although we could detect LD12 across various brackish salinities (Fig. 5A). In general, IIIb.A
and LSUCC0530 averaged the highest relative abundances across all sites with salinities
between 0 and 5.75. All subgroups showed site-specific variation in abundances (Fig. 5A).
Microcluster IIIb.A clearly dominated recruitment at Lake Michigan, whereas LSUCC0530
represented the most abundant genome at Feitsui Reservoir and Lake Gatun (Fig. 5A).
We repeated the previous examination of recruitment density to the different
microclusters across a range of percent identities, now including LSUCC0530 (Garcia *et al.*,
2017). We corroborate observed patterns indicating additional LD12 diversity beyond that
represented in the current genomes, despite the inclusion of our new representative. For
example, whereas the LSUCC0530 genome recruited the majority of high percent identity
reads in the Feitsui Reservoir and Lake Gatun samples (Fig. 5B), in some locations the
majority of recruitment to this genome centered around lower percent identity hits (Fig.
5C). We observed the same kind of alternative patterns for the other microclusters as well
(Fig. 5D). As noted our colleagues, since reads were competitively recruited against all
close relative genomes, these cases where a microcluster primarily recruited reads ~90-
92% identity likely mean that we lack a representative genome for that fraction of the
metagenomic data (Garcia *et al.*, 2017).
307
Discussion
Our successful cultivation of an LD12 strain for the first time has demonstrated that this
organism shares many physiological traits with cultured marine SAR11 representatives,
and also has some important distinctions. LSUCC0530 doubles roughly once every two
days, similar to taxa in subclade Ia (Rappé *et al.*, 2002) and Ib (Jimenez-Infante *et al.*, 2017).
It has an aerobic, chemoorganoheterotrophic lifestyle, thriving in oligotrophic media with
simple carbon compounds and likely depends on reduced organosulfur compounds and
ammonium as its sulfur and nitrogen sources, respectively. Cell size and shape showed
considerable morphological conservation (small, curved rods roughly 1 x 0.1 μm) across
large evolutionary distances (LD12 vs. subclade Ia ~89% 16S rRNA gene identity).
Although a systematic survey of salinity tolerance has not been conducted for other
cultured SAR11 representatives, we have shown that LD12 cannot survive in media with
salinities over eight, whereas all other cultured SAR11 grow in seawater salinities (Carini *et*
al., 2013, Jimenez-Infante *et al.*, 2017, Rappé *et al.*, 2002, Song *et al.*, 2009, Stingl *et al.*,
2007). LSUCC0530 thrives in relatively high temperatures (23-30°C), and we could not

confirm growth at low temperatures in which other SAR11 have been isolated (Song *et al.*,
2009).

LSUCC0530 shares many genomic characteristics with marine SAR11, such as a
small, streamlined genome with few paralogs, short intergenic spaces, and a conserved
hypervariable region, confirming predictions from LD12 SAGs and metagenomes (Eiler *et*
al., 2016, Newton *et al.*, 2011, Zaremba-Niedzwiedzka *et al.*, 2013). The LSUCC0530
genome also encodes much of the same metabolic potential as what has been reported
through LD12 SAGs and metagenomic data. For example, we confirm the EMP glycolysis
pathway, TCA cycle, non-mevalonate terpenoid biosynthesis, numerous amino acid
synthesis pathways, TRAP transporters for C4 dicarboxylate compounds, and ABC
transporters for various amino acids. We can confirm the lack of a phosphoenolpyruvate
carboxylase (ppc) and missing pathways for DMSP, glycine-betaine, and methylamine
metabolism.

We have also recovered important genetic information previously unknown for
LD12. At 1.16 Mbp, LSUCC0530 has a smaller genome, with fewer genes, than any other
SAR11. LSUCC0530 has a complete glyoxylate bypass, which differs from previous
inferences using SAGs (Eiler *et al.*, 2016). The unusual LD12 malate synthase isoform A
gene occupies the same genomic location in LSUCC0530 as the isoform G copy in most
subclade Ia, Ic, and IIIa genomes, with the exception of HIMB114 (Fig. S7). Therefore, the
most parsimonious explanation for the presence of isoform A in LSUCC0530 is a
homologous recombination event that replaced the isoform G homolog. The LSUCC0530
genome also contains putative pathways for oxidation of sulfide, thiosulfate, and sulfite,
which have not yet been explored in any cultured SAR11.

The relative abundance data from coastal Louisiana and the isolation of an LD12
strain from a coastal environment supports previous work indicating that LD12 occupies a
much wider range of aquatic habitats than inland freshwater systems (Dupont *et al.*, 2014,
Herlemann *et al.*, 2014, Piwosz *et al.*, 2013, Salcher *et al.*, 2011). Questions have been raised
as to whether these organisms occur in brackish waters as active and growing community
members, or whether there are present simply as a result of freshwater input. LD12 cells
have been shown to uptake thymidine in brackish water (Piwosz *et al.*, 2013). Our
observation that LSUCC0530 can grow in slightly brackish water provides further evidence
that such habitats may support LD12 cells as active community members. However, the
relative competitiveness of LD12 within brackish microbial assemblages requires further
research.

Metagenomic recruitment provides evidence that LD12 has diversified into multiple
ecotypes with unique distributions (Fig. 5A) (Garcia *et al.*, 2017, Zaremba-Niedzwiedzka *et*
al., 2013). Furthermore, our data suggests that LSUCC0530 likely represents a novel LD12
ecotype because it preferentially recruits metagenomic sequences at high percent identity
in different locations than the other microclusters of LD12 (Fig. 5). What distinguishes the
distribution of these ecotypes remains unclear. The observed patterns may represent

temporal variation captured in these sampling snapshots (Garcia *et al.*, 2017). Another
possibility is that small variations in salinity optima for different ecotypes may drive
diversification. However, given the paucity of low-brackish sites in available datasets, we
cannot establish whether salinity plays a role in ecotype diversification. Furthermore,
without complete genomes from other LD12 ecotypes, we cannot rule out whether variant
ionic strength management strategies, or some other metabolic characteristic, help define
evolutionary differences associated with ecotypes. For example, the ability to utilize
different sulfur sources may distinguish LSUCC0530 from other LD12. The *sorAB* sulfite
dehydrogenase and the thiosulfate/3-mercaptopyruvate sulfurtransferase genes have
uneven distribution among SAR11 genomes, including within the LD12 subclade.
An alternative hypothesis is that temperature has driven diversification between
ecotypes. A striking feature of LSUCC0530 is its growth preference for relatively high
temperatures and inability to grow at low temperatures. While robust growth at 30°C may
seem intuitive given that this represents typical Lake Borgne surface water temperature,
the inability to grow at 12°C or below would likely place LSUCC0530-type LD12 cells at a
fitness disadvantage in places like Lake Michigan, where annual water temperatures rarely
get above 22°C and average temperatures remain below 15°C for the majority of the year
(https://coastwatch.glerl.noaa.gov/). The metagenomic recruitment data support the
temperature hypothesis: the samples in which LSUCC0530 dominated recruitment (Lake
Gatun, Feitsui Reservoir) were all collected from locations having average temperatures
above 19°C. Conversely, other microclusters dominated recruitment of samples from colder
environments (e.g. IIIb.A in Lake Michigan)(Fig. 5). More sequence data may strengthen the
relationship between temperature and ecotypes, but since temperature optima are
extremely hard to infer from genomes alone, additional cultivars will be needed to test the
temperature hypothesis for ecotype diversification.
A critical question remains for LD12. What has made these cells uniquely adapted to
low salinity waters? Genomic analyses here and elsewhere (Dupont *et al.*, 2014, Eiler *et al.*,
2016) have helped define the differences in LD12 metabolic potential compared to other
SAR11, but very few adaptations appear related to salinity. It has been noted that the EMP
pathway provides more ATP and NADH than the ED pathway, thus conferring greater
energy conservation through glycolysis for LD12 cells than other SAR11s (Dupont *et al.*,
2014, Eiler *et al.*, 2016). If or how this energy conservation relates to salinity adaptation
has not been pursued. The presence of unique LD12 cation-proton antiporters (Fig. 3)
could provide a connection. LD12 cells may benefit from additional NADH for the
production of a proton motive force that drives the potassium-proton antiporters, thus
improving active transport to lower internal ionic strength. However, while this adaptation
may improve the ability of LD12 cells to live in fresh water, it doesn't explain why they
have lost the ability to survive in salt water where their marine SAR11 cousins thrive.
The most likely explanation comes in the distribution of genes for osmolyte uptake.
Critically, LSUCC0530, and LD12 in general, have lost the *proVWX* glycine-betaine/proline

ABC transporter shared by all other marine SAR11 (Table 1). Although a relatively minimal number of genes, this uptake capability likely represents a significant loss of function for LD12 salinity tolerance. In *Escherichia coli*, the *proVWX* operon has a strong promoter that upregulates these genes two orders of magnitude in response to increased salinity (Dattananda and Gowrishankar, 1989). Indeed, rapid proline uptake is a common response to osmotic upshift in many bacteria (Empadinhas and da Costa, 2008). Furthermore, LSUCC0530 and other LD12 genomes have lost the ABC transporters for ectoine/hydroxyectoine present in the other SAR11 genomes. Given the slow growth of LD12, rapid production of new intracellular osmolytes in response to increased salinity seems unlikely. Thus, the loss of these transport systems for quickly equilibrating intracellular osmolarity probably prevents successful dispersal of this organism into higher salinity waters.

This finding allows us to speculate further about the evolutionary trajectory that led to successful colonization of freshwater environments by LD12. All evidence suggests that the last common SAR11 ancestor was a marine organism with a streamlined genome (Logares *et al.*, 2010, Luo *et al.*, 2013, Zaremba-Niedzwiedzka *et al.*, 2013). During further diversification into subclade III, additional genes and pathways were lost, for example, ppc and genes for DMSP and some C1 compound metabolisms (Eiler *et al.*, 2016, Sun *et al.*, 2016). Therefore, the common ancestor of subclade IIIa and LD12 would have had to adapt to freshwater environments starting with an already reduced genome and limited metabolic flexibility. A simple way to inhabit lower salinity environments would have been to decrease expression of, or altogether lose, genes for uptake and/or production of osmolytes. If the adaptation were reversible, we would expect to see examples of LD12 organisms growing in marine systems. However, existing evidence suggests that while LD12 organisms can grow (established herein) and be active (Piwosz *et al.*, 2013) in low brackish conditions, they are not successful in higher salinity brackish or marine water (Dupont *et al.*, 2014, Herlemann *et al.*, 2014, Logares *et al.*, 2010, Salcher *et al.*, 2011). Thus, we propose that LD12 adaptation to fresh water involved irreversible loss of function, which decreased osmotic response capability in LD12 cells, thereby preventing recolonization of higher salinity environments. The missing ABC transporters for proline/glycine betaine and ectoine/hydroxyectoine in LD12 are likely candidates for this loss of function. Experimental tests on the differences in production and uptake of osmolytes among subclade IIIa and LD12 representatives will provide critical insight into how these different evolutionary paths have resulted in unique salinity ranges.

For the first cultured representative of the LD12 clade, we propose the provisional taxonomic assignment for strain LSUCC0530 as '*Candidatus* Fonsibacter ubiquis',

Description of *Fonsibacter* gen. nov.

442   *Fonsibacter* (*Fons* L. noun; fresh water, spring water, -bacter, Gr. adj.; rod, bacterium.
443   Fonsibacter referring to the preferred low salinity habitat of this organism).
444
Aerobic, chemoorganoheterotrophic, and oligotrophic. Cells are small, curved rods roughly
1 x 0.1μm. Non-motile. Based on phylogenomics and 16S rRNA gene phylogenetics,
subclade IIIb/LD12 occurs on a separate branch within the *Pelagibacteraceae* (SAR11),
sister to subclade IIIa containing strains HIMB114 and IMCC9063. Due to the depth of
branching between these clades using concatenated protein-coding genes, and the 92%
and 91% 16S rRNA gene sequence identity with HIMB114 and IMCC9063, respectively, we
propose LSUCC0530 and subclade IIIb/LD12 as a novel genus in the *Pelagibacteraceae*.
452

Description of *Fonsibacter ubiquis* sp. nov.
453
454   *Fonsibacter ubiquis* (L. adv. ubiquitous).
455
In addition to the properties given in the genus description, the species is described as
follows. Growth occurs at temperatures between 23°C and 30°C, but not at 11°C or below,
nor at 35°C or above. Optimal salinity is 1.45 and below, and growth occurs between 0.36
and 4.68. At optimal temperature and salinity, cells divide at an average rate of 0.52 day^{-1}.
Genome size is 1.16 Mbp, with 1271 predicted genes, and a GC content of 29.02%
(calculated). LSUCC0530 had an ANI of 75.3% and 75.9% and an AAI of 58.1% and 59.7%
with HIMB114 and IMCC9063, respectively. The genome is available on GenBank under
accession number CP024034.
464
465   The type strain, LSUCC0530^T, was isolated from brackish water from the coastal lagoon
466   Lake Borgne off the coast of southeastern Louisiana, USA.
467
468

Materials and Methods
469
Isolation and Identification of LSUCC0530
Isolation, propagation, and identification of strain LSUCC0530 (subclade IIIb) was
completed as previously reported (Henson *et al.*, 2016). Briefly, water was collected from
the surface of the coastal lagoon Lake Borgne (Shell Beach, Louisiana) (29.872035 N, -
89.672819 W) on 1 July, 2016. We recorded salinity on site using a YSI 556 MPS (YSI, Solon,
OH, USA) (Table S1). Whole water was filtered through a 2.7 μm filter (Whatman), stained
with 1X SYBR Green (Lonza, Basal, Switzerland), and enumerated using the Guava EasyCyte
5HT flow cytometer (Millipore, Massachusetts, USA). After serial dilution, water was
inoculated into five 96 x 2.1 mL well PTFE plates (Radleys, Essex, UK) containing 1.7 mL of
JW5 medium at an estimated 2 cells well^{-1}. Cultures were incubated at room temperature in
the dark for three weeks and evaluated for growth using flow cytometry. Positive wells (>
10^4 cells mL^{-1}) were transferred in duplicate to capped, 125 mL polycarbonate flasks

(Corning, New York, USA) containing 50 mL of JW5 medium. Upon reaching a density of >
5.0×10^5 cells mL^{-1}, cells were filtered onto 25 mm 0.22 μm polycarbonate filters (Millipore,
Massachusetts, USA), and DNA extractions were performed using the MoBio PowerWater
DNA kit (QIAGEN, Massachusetts, USA) following the manufacturer's instructions. DNA was
amplified as previously reported(Henson *et al.*, 2016) and sequenced at Michigan State
University RTSF Genomics Core. Sanger sequences were evaluated using the freely
available software 4Peaks (v. 1.7.1)(http://nucleobytes.com/4peaks/). Forward and
reverse sequences were assembled using the CAP3 webserver
(http://doua.prabi.fr/software/cap3), after reverse reads were converted to their reverse
complement at http://www.bioinformatics.org/sms/rev_comp.html. The PCR-generated
16S rRNA gene sequence for strain LSUCC0530 is accessible on NCBI GenBank under the
accession number KY290650.
494
Community iTag analysis
Community DNA from 17 sites of various salinities was filtered, extracted, and analyzed
following our previously reported protocol (Henson *et al.*, 2016). The sites sampled were
Lake Borgne (LKB; Shell Beach, LA), Bay Pomme D'or (JLB; Buras, LA), Terrebonne Bay
(TBon; Cocodrie, LA), Atchafalaya River Delta (ARD; Franklin, LA), Freshwater City (FWC;
Kaplan, LA), Calcasieu Jetties (CJ; Cameron, LA). Each site was visited three times with the
exception of TBon, which was visited only twice. Briefly, extracted DNA was sequenced at
the 16S rRNA gene V4 region (515F, 806R) (Parada *et al.*, 2015), with Illumina MiSeq
2x250bp paired-end sequencing at Argonne National Laboratories. 16S rRNA gene
amplicon data was analyzed and classified (OTU$_{0.03}$) with Mothur v.1.33.3 (Schloss *et al.*,
2009) using the Silva v119 database (Pruesse *et al.*, 2007). To determine the best matching
OTU for LSUCC0530, sequences from the OTU representative fasta files, provided by
mothur using *get.oturep*(), were used to create a blast database (*formatdb*) against which
the LSUCC isolate 16S rRNA genes could be searched via blastn (BLAST v 2.2.26). All best
hits ranked at 100% identity. OTU abundance analysis was conducted within the R
statistical environment v3.2.3, within the package PhyloSeq (v1.14.0) (McMurdie and
Holmes, 2013). Sequences were rarefied using the command *rarefy_even_depth*() and then
trimmed for OTUs with at least 2 reads in at least 90% of the samples. Abundances of an
OTU were averaged across biological duplicates and a rank abundance matrix was
calculated using the command *transform_sample_counts*() with the argument function of
function(x) x / sum(x). Completed rarefied rank abundance OTU tables were plotted using
the graphing program GGPLOT2 (v. 2.0.0) (Wickham, 2011). All iTag sequences are
available at the Short Read Archive with accession number SRR6235382-SRR6235415.
518
Growth experiments
Salinity tolerance was tested by growing LSUCC0530 in different media distinguished by
proportional changes in the concentration of major ions. All other media elements (carbon,

iron, phosphate, nitrogen, vitamins, and trace metals) were kept consistent. Media included
JW1, JW2, JW3, and JW4, previously published (Henson *et al.*, 2016), and JW3.5, JW4.5, JW5,
and JW6. Recipes for all media used in this study are provided in Table S1. Salinity
tolerance experiments were conducted in quintuplicate at room temperature. Temperature
range and sole carbon substrate use were tested using the isolation medium, JW5.
Temperature range experiments were also conducted in quintuplicate. For all experiments,
growth was measured using flow cytometry as described above.
529
Microscopy
Strain LSUCC0530 cells were grown to near max density (1×10^7 cells mL^{-1}) in JW5 medium.
Cells for transmission electron microscopy were prepared as previously described (Rappé
et al., 2002) with one minor change: centrifuged cells were concentrated in JW5 medium
with no added fixative. Cells were visualized under a JEOL JSM-2011 High Resolution
transmission electron microscope at the LSU Socolofsky Microscopy Center, Baton Rouge,
LA. For scanning electron microscopy, cultured cells were fixed for four hours in 4%
Glutaraldehyde buffered with 0.2M Cacodylate buffer and filtered onto a 0.2 μm nylon filter
(Pall, Michigan, USA). Samples were dehydrated in an ethanol series (30%, 50%, 70%, 96
and 100%) for 15 minutes each, followed by critical point drying. Cells were visualized
under a JEOL JSM-6610 scanning electron microscope at the LSU Socolofsky Microscopy
Center, Baton Rouge, LA.
542
Genome sequencing, assembly, and annotation
Cells were grown in 1 L JW5 medium (Table S1) and filtered onto 0.2 μm nylon filters (Pall,
Michigan, USA). DNA was obtained via phenol-chloroform extraction and sequenced using
both Illumina HiSeq and MiSeq approaches. *HiSeq*. 350 ng of DNA was sheared to ~500 bp
with a Qsonica Q800R (QSonica) using 16 cycles with a amplitude of 25% and pulse rate of
20 seconds on, 20 seconds off. DNA was visualized on a 1.5% agrose gel stained with Gel
Red (Biotum) to determine average sheared length and quantified using Qubit high
sensitivity dsDNA assay kit (Invitrogen). Library prep was conducted following a modified
Kapa Hyper Prep Kit protocol (Kapa Biosystems, Inc.). Briefly, we ligated end-repaired and
adenylated DNA to universal Y-yoke oligonucleotide adapters and custom Illumina iTru
dual-indexed primers (Glenn *et al.*, 2016). Prior to index amplification, post-ligation
LSUCC0530 DNA was cleaned using 1.2x SPRI (Roland and Reich, 2012). Following a brief,
8 cycle PCR amplification of the library to add indexes, the genomic library was quantified
using a Qubit broad range dsDNA assay kit (Invitrogen). Illumina HiSeq 3000 sequencing
was performed at Oklahoma Medical Research Facility using SBS kit chemistry to generate
150 bp paired-end reads, insert size 400 bp. *MiSeq*. DNA was sent to the Argonne National
Laboratory Environmental Sample Preparation and Sequencing Facility which performed
library prep and sequencing, generating 250 bp paired-end reads, insert size 550 bp.

Due to the large number of reads from the HiSeq sequencing, we created a random
subset using 14% of the total reads by using the program *seqtk* (v. 1.0-r75-dirty) and the
flag –s100. We trimmed reads for adapter contamination and quality using Trimmomatic
(Bolger *et al.*). Initial assembly with the subset of reads was done with SPAdes (Bankevich
et al., 2012). This generated a large single scaffold with overlapping ends. We performed a
quality assessment of the single scaffold via iterations of Reapr (Hunt *et al.*, 2013)-advised
scaffold breaks and SSPACE (Boetzer *et al.*, 2011) extensions using both the full set of HiSeq
reads and the MiSeq reads. Final assessment of the scaffold with overlaps removed from
the ends was performed with Pilon (Walker *et al.*, 2014) using all *HiSeq* reads mapped to
the scaffold with BWA (Li and Durbin, 2009). No issues were reported. An analysis of the
GC skew was performed on the final scaffold using the gc_skew script provided via Brown
et al. 2015 (Brown *et al.*, 2015). Contamination and completion were also evaluated with
CheckM (Parks *et al.*, 2015). Detailed commands and outputs for the assembly and quality
assessment are provided in Supplementary Information. The final circular chromosome
scaffold was annotated by IMG (Markowitz *et al.*, 2014) and is publicly available with IMG
Taxon ID number 2728369501. It is also publicly available in GenBank with accession
number CP024034. Genome sequencing reads are available in SRA with accession numbers
SRR6257553 (MiSeq) and SRR6257608 (HiSeq).
579
Comparative genomics
Orthologus clusters were determined using the Get Homologues pipeline (Contreras-
Moreira and Vinuesa, 2013) using the LSUCC0530 genome and 48 publicly available SAR11
genomes from IMG, including the 10 LD12 SAGs analyzed previously (Eiler *et al.*, 2016,
Zaremba-Niedzwiedzka *et al.*, 2013) (Table S1). Clusters were determined from amino acid
sequences using the OrthoMCL option in Get Homologues. All clusters are provided in
cluster_list.txt as part of Supplemental Information. Average amino acid identity (AAI) and
synteny were calculated as previously reported (Dick *et al.*, 2009). Average nucleotide
identity (ANI) was calculated with the Konstantinidis Lab website ANI calculator
(http://enve-omics.ce.gatech.edu/ani/index) as reported (Goris *et al.*, 2007). Compatible
solutes identified in Table S1 were identified in SAR11 genomes using KO numbers,
annotations, and cross-referenced with orthologous clusters.
592
593 *Bacteriophage searches*
594 We looked for signatures of bacteriophage in the LSUCC0530 genome using two methods,
595 VirSorter (Roux *et al.*, 2015) and PHASTER (Arndt *et al.*, 2016) with default settings.
596 CRISPR regions are identified as part of the standard IMG annotation process, and none
597 were found in LSUCC0530.
598
Phylogenomics

We selected single-copy genes present in at least 39 of the 49 genomes as determined via
Get Homologues (above). Amino acid sequences from these 83 markers were separately
aligned with MUSCLE (Edgar, 2004), culled with Gblocks (Castresana, 2000), and
concatenated into a superalignment totaling 23,356 positions. Inference was performed
with RAxML (Stamatakis *et al.*, 2008) using the PROTCATLG model and 100 bootstrapping
runs. All scripts (with settings) used in the alignment, culling, concatenation, and inference
processes are available in Supplemental Information. Node labels were renamed with
Newick Utilities (Junier and Zdobnov, 2010) *nw_rename*, and the tree was visualized with
Archaeopteryx (Han and Zmasek, 2009).
609
Single gene phylogeny
The 16S rRNA gene sequence from the LSUCC0530 genome was aligned with 284
alphaproteobacterial sequences and seven outgroup sequences. We completed
phylogenetic inference for *aceA* and *malA* in an analogous manner. For each, the
LSUCC0530 gene was searched against the NCBI nr database using blastp (v. 2.2.28). The
top 100 sequence hits were combined with all homologus SAR11 sequences identified with
Get Homologues for the *aceA* phylogeny, and both the *malA* and *malG* sequences in all
SAR11s for the *malA* phylogeny. Sequences from redundant taxa were removed. For all
trees, alignment with MUSCLE (Edgar, 2004), culling with Gblocks (Castresana, 2000), and
tree inference with FastTree2 (Price *et al.*, 2010) was completed with the FT_pipe script,
provided in Supplemental Information, along with the initial fasta file containing all
sequences and the newick tree output file. Node labels were changed with Newick Utilities
(Junier and Zdobnov, 2010) *nw_rename* and visualization was performed with
Archaeopteryx (Han and Zmasek, 2009).
624
Metagenomic recruitment
Scaffolds for the 49 SAR11 genomes, including strain LSUCC0530, used in the
phylogenomic analyses were used for competitive recruitment of metagenomic sequences
from 89 different samples. Prior to recruitment, multi-scaffold assemblies were condensed
into single scaffolds, and all single scaffolds were used to generate a BLAST database with
the command *makeblastdb* (BLAST v. 2.2.28). Metagenomic datasets from 90 different
environmental samples of various salinities from the Baltic Sea (Dupont *et al.*, 2014),
Amazon River, freshwater lakes (Eiler *et al.*, 2014, Martinez-Garcia *et al.*, 2011), Lagoon
Albufera (Ghai *et al.*, 2012), Columbia River (Fortunato and Crump, 2015, Smith *et al.*,
2013b), Lake Lanier (Oh *et al.*, 2011), Lake Michigan (Denef *et al.*, 2016), Feitsui Reservoir
(Tseng *et al.*), and GOS (Rusch *et al.*, 2007) were downloaded from the Short Read Archive
(SRA) and European Nucleotide Archive (ENA) using *ftp* and SRA toolkit (v.2.8.2-1),
respectively. Reads from pyrosequencing were dereplicated with the program CD-HIT (v
4.6) (Fu *et al.*, 2012) using the *cd-hit-454* command and flag -c 0.95. Illumina sequence data
was converted from fastq to fasta files with the FastX toolkit (v. 0.0.13.2)

(http://hannonlab.cshl.edu/fastx_toolkit) using the command *fastq_to_fa*sta with the flag –
Q33. Metagenomic sequences were separated into 1 million sequence files and recruited
against the SAR11 database with blastn (v. 2.2.28) with –max_target_seqs 1. Best hits for
each metagenomic sequence were kept if the alignment length was > 90% of the median
length of all metagenomic sequences for a given sample. Metagenomic recruitment was
expressed in read density per genome using ggplot2 *geom_density*() and reads per kilobase
of genome per million mapped reads (RPKM) as previously reported (Thrash *et al.*, 2017).
647
648   *Script availability*
649   All scripts used in this work can be found on the Thrash Lab website with the manuscript
650   link at http://thethrashlab.com/publications.
651

Acknowledgements
Funding for this work was provided in part by a Louisiana Board of Regents (grant
LEQSF(2014-17)-RD-A-06) grant to JCT and the LSU Department of Biological Sciences
(startup funds to JCT and BCF). The authors thank the LSU Socolofsky Microscopy Center
for assistance with electron microscopy imaging. Portions of this research were conducted
with high performance computing resources provided by Louisiana State University
(http://www.hpc.lsu.edu).

Author Contributions
JCT designed the study; MWH conducted cultivation experiments, nucleic acid extraction,
microscopy, and iTag and metagenomics analyses; JCT and BCF assembled the genome and
conducted quality assessment; MWH, VCL, and JCT conducted comparative genomics
analyses, metabolic reconstruction, phylogenetic, and phylogenomic analyses; JCT lead
manuscript writing and all authors contributed to and reviewed the text.

Competing Financial Interests
The authors declare no competing financial interests.
669
670
671

 Tables and Figures

674 **Table 1.** Compatible solute production and transport in SAR11

Osmolyte system	LSUCC0530	LD12 SAGs	HIMB114	IMCC9063	Other SAR11
Alanine synthesis	+	+	+	+	+
Di-myo-inositol phosphate synthesis	+/-	+/-	+/-	+/-	+/-
Ectoine/hydroxyectoine transport	-	-	+	+	+
Glutamate synthesis	+	+	+	+	+
Glutamate transport	-	+*	-	-	-
Glutamine synthesis	+	+	+	+	+
Glycine synthesis	+	+	+	+	+
Glycine betaine synthesis	-	+/-*	+	+	+
Glycine betaine/proline ABC transporter	-	-	+	+	+
N-∂-Acetyl-ornithine synthesis	+	+	+	+	+
Potassium transport (antiporters K03549, K03455)	+	+	-	-	-
Potassium transport (trkAH + K16052)	-	-	+	+	+
Proline synthesis	+	+	+	+	+
Proline transport (symporter)	+	+	+	+	+
Sodium transport (antiporter K03313)	-	-	+	+	+
Sodium transport (other antiporter)	+	+	-	-	-
Sorbitol synthesis	-	-	+	-	+
Taurine transport	-	-	-	-	+
TMAO synthesis	-	-	-	-	+

+ genes present; - genes absent; +/- partial pathway; *B11 LD12 SAG only. Table shows

only systems with predicted genes in at least one SAR11 genome. Table only shows

systems for which genes could be identified in the SAR11 genomes examined in this study.

For a complete list of all systems searched, see Table S1.

Figure Legends

Figure 1. Phylogeny and morphology of LSUCC0530. **A)** Phylogenetic position of

LSUCC0530 within the SAR11 clade using a concatenation of 83 single copy marker genes

and 23,356 amino acid positions. The tree was inferred using RAxML with 100 bootstrap

runs. Values at the nodes indicate 100% bootstrap support unless otherwise noted. The

tree was rooted on HIMB59. Subclades are indicated on the right, and LD12 microclusters

are indicated A-C. Asterisk- we have included AAA028-C07 in microcluster A, even though

it was excluded previously (Zaremba-Niedzwiedzka *et al.*, 2013). **B)** Scanning electron

micrograph of LSUCC0530 cells at 35,000x magnification. Scale bar represents 0.5 μm. **C)**

Transmission electron micrograph of a LSUCC0530 cell thin section. Scale bar represents

0.5 μm.

Figure 2. Circular diagram of the LSUCC0530 genome. Rings indicate, from outside to the
center: Position relative to the replication start site; Forward strand genes, colored by COG
category; Reverse strand genes, colored by COG category; RNA genes (tRNAs green, rRNAs
red, other RNAs black); GC content; GC skew.

Figure 3. Metabolic reconstruction of LSUCC0530. Solid boxes indicate recovered genes,
dashed boxes indicate missing genes. Color shading distinguishes major metabolic sub-
systems, which are also labeled in bold text. ABC transporters are depicted with circles
indicating subunits. Symporters and antiporters are depicted with ovals. Two component
systems are depicted with large and small rectangles. The C4 TRAP transporter is depicted
separately. Major subsystems are colored for easier identification. Multiple arrows indicate
all genes present in a given pathway. Numbers identify genes according to the key in Table
S1. Question marks indicate a single missing gene in an otherwise complete pathway (e.g.,
PRPP -> His). Light blue fill indicates genes with no SAR11 orthologs outside of the LD12
subclade.

Figure 4. Growth rates for LSUCC0530 according to **A)** salinity and **B)** temperature. The
boxes indicate the interquartile range (IQR) of the data, with vertical lines indicating the
upper and lower extremes according to 1.5 x IQR. Horizontal lines within each box indicate
the median. The underlying data points of individual biological replicates, calculated from
corresponding growth curves in Figure S8, are plotted on top of each box.

Figure 5. Results of competitive metagenomic recruitment to all genomes used in the
phylogenomic tree. **A)** 95% identity RPKM values for LD12 genomes only, by site, with data
aggregated for all genomes in microclusters A-C, according to the key. Colors indicate broad
environmental categories. **B-D)** Density plots of metagenomic sequence recruitment
according to percent identity of the best hit. Data was aggregated according to the key, with
LD12 separated as in A, HIMB59 separately depicted as "V", and all other SAR11 genomes
together as "Other." **B)** Two sites where LSUCC0530 dominated recruitment with high
percent identity. **C)** Two sites where the LSUCC0530 only recruited sequences at ~90%
identity and no other group had high recruitment at higher identity. **D)** Two sites where
other LD12 microclusters showed similar patterns as LSUCC0530 in B and C.

References Cited

Anstrom DM, Kallio K, Remington JS (2003). Structure of the Escherichia coli malate synthase G:pyruvate:acetyl‐coenzyme A abortive ternary complex at 1.95 Å resolution. *Protein Science* **12:** 1822-1832.

Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y *et al.* (2016). PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Research* **44**.

Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS *et al.* (2012). SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. *Journal of Computational Biology* **19:** 455-477.

Béjà O, Spudich EN, Spudich JL, Leclerc M, DeLong EF (2001). Proteorhodopsin phototrophy in the ocean. *Nature* **411:** 786-789.

Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W (2011). Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics* **27:** 578-579.

Bolger AM, Lohse M, Usadel B (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30:** 2114-2120.

Brown CT, Hug LA, Thomas BC, Sharon I, Castelle CJ, Singh A *et al.* (2015). Unusual biology across a group comprising more than 15% of domain Bacteria. *Nature* **523:** 208-211.

Cabello-Yeves PJ, Zemskaya TI, Rosselli R, Coutinho FH, Zakharenko AS, Blinov VV *et al.* (2018). Genomes of novel microbial lineages assembled from the sub-ice waters of Lake Baikal. *Appl Environ Microbiol* **84:** e02132-02117.

Carini P, Steindler L, Beszteri S, Giovannoni SJ (2013). Nutrient requirements for growth of the extreme oligotroph 'Candidatus Pelagibacter ubique' HTCC1062 on a defined medium. *ISME J* **7:** 592-602.

Carini P, Campbell EO, Morré J, Sañudo-Wilhelmy SA, Thrash CJ, Bennett SE *et al.* (2014). Discovery of a SAR11 growth requirement for thiamin's pyrimidine precursor and its distribution in the Sargasso Sea. *ISME J* **8:** 1727-1738.

Carlson CA, Morris R, Parsons R, Treusch AH, Giovannoni SJ, Vergin KL (2009). Seasonal dynamics of SAR11 populations in the euphotic and mesopelagic zones of the northwestern Sargasso Sea. *ISME J* **3:** 283-295.

Castresana J (2000). Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol* **17:** 540-552.

773 Chun J, Oren A, Ventosa A, Christensen H, Arahal D, da Costa MS *et al.* (2018). Proposed
774 minimal standards for the use of genome data for the taxonomy of prokaryotes. *IJSEM* **68:**
775 461-466.
776
777 Contreras-Moreira B, Vinuesa P (2013). GET_HOMOLOGUES, a versatile software package
778 for scalable and robust microbial pangenome analysis. *Appl Environ Microbiol* **79:** 7696-
779 7701.
780
781 Dattananda CS, Gowrishankar J (1989). Osmoregulation in Escherichia coli:
782 complementation analysis and gene-protein relationships in the proU locus. *J Bacteriol*
783 **171:** 1915-1922.
784
785 Denef VJ, Mueller RS, Chiang E, Liebig JR, Vanderploeg HA (2016). Chloroflexi CL500-11
786 Populations That Predominate Deep-Lake Hypolimnion Bacterioplankton Rely on
787 Nitrogen-Rich Dissolved Organic Matter Metabolism and C1 Compound Oxidation. *Appl*
788 *Environ Microbiol* **82:** 1423-1432.
789
790 Denger K, Weinitschke S, Smits THM, Schleheck D, Cook AM (2008). Bacterial sulfite
791 dehydrogenases in organotrophic metabolism: separation and identification in Cupriavidus
792 necator H16 and in Delftia acidovorans SPH-1. *Microbiology* **154:** 256-263.
793
794 Dick GJ, Andersson AF, Baker BJ, Simmons SL, Thomas BC, Yelton AP *et al.* (2009).
795 Community-wide analysis of microbial genome sequence signatures. *Genome Biol* **10:** R85.
796
797 Dupont CL, Larsson J, Yooseph S, Ininbergs K, Goll J, Asplund-Samuelsson J *et al.* (2014).
798 Functional Tradeoffs Underpin Salinity-Driven Divergence in Microbial Community
799 Composition. *PLOS ONE* **9:** e89549.
800
801 Edgar RC (2004). MUSCLE: multiple sequence alignment with high accuracy and high
802 throughput. *Nucleic Acids Research* **32:** 1792-1797.
803
804 Eiler A, Zaremba-Niedźwiedzka K, Martinez-Garcia M, McMahon KD, Stepanauskas R,
805 Andersson SGE *et al.* (2014). Productivity and salinity structuring of the microplankton
806 revealed by comparative freshwater metagenomics. *Environmental Microbiology* **16:** 2682-
807 2698.
808
809 Eiler A, Mondav R, Sinclair L, Fernandez-Vidal L, Scofield DG, Schwientek P *et al.* (2016).
810 Tuning fresh: radiation through rewiring of central metabolism in streamlined bacteria.
811 *ISME J* **10:** 1902-1914.
812
813 Empadinhas N, da Costa M (2008). Osmoadaptation mechanisms in prokaryotes:
814 distribution of compatible solutes. *International Microbiology*.
815
816 Fortunato CS, Crump BC (2015). Microbial Gene Abundance and Expression Patterns across
817 a River to Ocean Salinity Gradient. *PLOS ONE* **10**.
818

819   Fu L, Niu B, Zhu Z, Wu S, Li W (2012). CD-HIT: accelerated for clustering the next-
820   generation sequencing data. *Bioinformatics* **28:** 3150-3152.
821
822   Garcia SL, Stevens SLR, Crary B, Martinez-Garcia M, Stepanauskas R, Woyke T *et al.* (2017).
823   Contrasting patterns of genome-level diversity across distinct co-occurring bacterial
824   populations. *ISME J* **AOP doi:10.1038/s41396-017-0001-0**.
825
826   Ghai R, Hernandez C, Picazo A, Mizuno C, Ininbergs K, Díez B *et al.* (2012). Metagenomes of
827   Mediterranean Coastal Lagoons. *Sci Rep* **2**.
828
829   Giovannoni SJ, Bibbs L, Cho J-C, Stapels MD, Desiderio R, Vergin KL *et al.* (2005a).
830   Proteorhodopsin in the ubiquitous marine bacterium SAR11. *Nature* **438:** 82-85.
831
832   Giovannoni SJ, Tripp HJ, Givan S, Podar M, Vergin KL, Baptista D *et al.* (2005b). Genome
833   streamlining in a cosmopolitan oceanic bacterium. *Science* **309:** 1242-1245.
834
835   Giovannoni SJ, Vergin KL (2012). Seasonality in ocean microbial communities. *Science* **335:**
836   671-676.
837
838   Glenn TC, Nilsen R, Kieran TJ, Finger JW, Pierson TW, Bentley KE *et al.* (2016). Adapterama
839   I: Universal stubs and primers for thousands of dual-indexed Illumina libraries (iTru &
840   iNext). *bioRxiv*.
841
842   Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM (2007).
843   DNA–DNA hybridization values and their relationship to whole-genome sequence
844   similarities. *IJSEM* **57:** 81-91.
845
846   Grote J, Thrash JC, Huggett MJ, Landry ZC, Carini P, Giovannoni SJ *et al.* (2012). Streamlining
847   and Core Genome Conservation among Highly Divergent Members of the SAR11 Clade.
848   *mBio* **3:** e00252-00212.
849
850   Han MV, Zmasek CM (2009). phyloXML: XML for evolutionary biology and comparative
851   genomics. *BMC Bioinformatics* **10:** 1-6.
852
853   Henson MW, Pitre DM, Weckhorst J, Lanclos VC, Webber AT, Thrash JC (2016). Artificial
854   Seawater Media Facilitate Cultivating Members of the Microbial Majority from the Gulf of
855   Mexico. *mSphere* **1:** e00028-00016. doi:00010.01128/mSphere.00028-00016.
856
857   Herlemann D, Woelk J, Labrenz M, Jürgens K (2014). Diversity and abundance of
858   "Pelagibacterales" (SAR11) in the Baltic Sea salinity gradient. *Syst Appl Microbiol* **37:** 601-
859   604.
860
861   Hunt M, Kikuchi T, Sanders M, Newbold C, Berriman M, Otto TD (2013). REAPR: a universal
862   tool for genome assembly evaluation. *Genome Biol* **14:** 1-10.
863

Jimenez-Infante F, Ngugi D, Vinu M, Blom J, Alam I, Bajic VB *et al.* (2017). Genomic
Characterization of Two Novel SAR11 Isolates From the Red Sea, Including the First Strain
of the SAR11 Ib clade. *FEMS Microbiol Ecol*.

Junier T, Zdobnov EM (2010). The Newick utilities: high-throughput phylogenetic tree
processing in the Unix shell. *Bioinformatics* **26:** 1669-1670.

Kappler U (2011). Bacterial sulfite-oxidizing enzymes. *Biochimica et Biophysica Acta (BBA)
- Bioenergetics* **1807:** 1-10.

Konstantinidis KT, Tiedje JM (2005). Genomic insights that advance the species definition
for prokaryotes. *Proc Natl Acad Sci USA* **102:** 2567-2572.

Konstantinidis KT, Tiedje JM (2007). Prokaryotic taxonomy and phylogeny in the genomic
era: advancements and challenges ahead. *Current Opinion in Microbiology* **10:** 504-509.

Li H, Durbin R (2009). Fast and accurate short read alignment with Burrows–Wheeler
transform. *Bioinformatics* **25:** 1754-1760.

Logares R, Bråte J, Bertilsson S, Clasen JL, Shalchian-Tabrizi K, Rengefors K (2009).
Infrequent marine–freshwater transitions in the microbial world. *Trends in Microbiology*
17: 414-422.

Logares R, Brate J, Heinrich F, Shalchian-Tabrizi K, Bertilsson S (2010). Infrequent
Transitions between Saline and Fresh Waters in One of the Most Abundant Microbial
Lineages (SAR11). *Mol Biol Evol* **27:** 347-357.

Luo H, Csűrös M, Hughes AL, Moran M (2013). Evolution of Divergent Life History
Strategies in Marine Alphaproteobacteria. *mBio* **4:** e00373-00313.

Markowitz VM, Chen IMA, Palaniappan K, Chu K, Szeto E, Pillay M *et al.* (2014). IMG 4
version of the integrated microbial genomes comparative analysis system. *Nucleic Acids
Research* **42:** D560-567.

Martinez-Garcia M, Swan BK, Poulton NJ, Gomez M, Masland D, Sieracki ME *et al.* (2011).
High-throughput single-cell sequencing identifies photoheterotrophs and
chemoautotrophs in freshwater bacterioplankton. *ISME J* **6:** 113-123.

McMurdie PJ, Holmes S (2013). phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. *PLOS ONE* **8:** e61217.

Morris RM, Rappé MS, Connon SA, Vergin KL, Siebold WA, Carlson CA *et al.* (2002). SAR11
clade dominates ocean surface bacterioplankton communities. *Nature* **420:** 806-810.

Newton RJ, Jones SE, Eiler A, McMahon KD, Bertilsson S (2011). A Guide to the Natural
History of Freshwater Lake Bacteria. *Microbiol Mol Biol Rev* **75:** 14-49.

Oh S, Caro-Quintero A, Tsementzi D, DeLeon-Rodriguez N, Luo C, Poretsky R *et al.* (2011).
Metagenomic Insights into the Evolution, Function, and Complexity of the Planktonic
Microbial Community of Lake Lanier, a Temperate Freshwater Ecosystem. *Appl Environ*
Microbiol **77:** 6000-6011.

Parada AE, Needham DM, Fuhrman JA (2015). Every base matters: assessing small subunit
rRNA primers for marine microbiomes with mock communities, time series and global field
samples. *Environmental Microbiology* **18**.

Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW (2015). CheckM: assessing
the quality of microbial genomes recovered from isolates, single cells, and metagenomes.
Genome Research **25:** 1043-1055.

Piwosz K, Salcher MM, Zeder M, Ameryk A, Pernthaler J (2013). Seasonal dynamics and
activity of typical freshwater bacteria in brackish waters of the Gulf of Gdańsk. *Limnology*
and Oceanography **58:** 817-826.

Price MN, Dehal PS, Arkin AP (2010). FastTree 2--approximately maximum-likelihood trees
for large alignments. *PLOS ONE* **5:** e9490.

Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig W, Peplies J *et al.* (2007). SILVA: a
comprehensive online resource for quality checked and aligned ribosomal RNA sequence
data compatible with ARB. *Nucleic Acids Research* **35:** 7188-7196.

Rappé MS, Connon SA, Vergin KL, Giovannoni SJ (2002). Cultivation of the ubiquitous
SAR11 marine bacterioplankton clade. *Nature* **418:** 630-633.

Rodriguez-Valera F, Martin-Cuadrado A-B, Rodriguez-Brito B, Pašić L, Thingstad FT,
Rohwer F *et al.* (2009). Explaining microbial population genomics through phage
predation. *Nat Rev Micro* **7:** 828-836.

Roland N, Reich D (2012). Cost-effective, high-throughput DNA sequencing libraries for
multiplexed target capture. *Genome Research* **22:** 939-946.

Roux S, Enault F, Hurwitz BL, Sullivan MB (2015). VirSorter: mining viral signal from
microbial genomic data. *PeerJ* **3**.

Rusch DB, Halpern AL, Sutton G, Heidelberg KB, Williamson S, Yooseph S *et al.* (2007). The
Sorcerer II Global Ocean Sampling Expedition: Northwest Atlantic through Eastern Tropical
Pacific. *Plos Biol* **5:** e77.

Salcher MM, Pernthaler J, Posch T (2011). Seasonal bloom dynamics and ecophysiology of
the freshwater sister clade of SAR11 bacteria 'that rule the waves' (LD12). *ISME J* **5:** 1242-
1252.

Schattenhofer M, Fuchs BM, Amann R, Zubkov MV, Tarran GA, Pernthaler J (2009).
Latitudinal distribution of prokaryotic picoplankton populations in the Atlantic Ocean.
Environmental Microbiology **11:** 2078-2093.
959
960   Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB *et al.* (2009).
961   Introducing mothur: Open-Source, Platform-Independent, Community-Supported Software
962   for Describing and Comparing Microbial Communities. *Appl Environ Microbiol* **75:** 7537-
963   7541.
964
965   Schwalbach MS, Tripp HJ, Steindler L, Smith DP, Giovannoni SJ (2010). The presence of the
966   glycolysis operon in SAR11 genomes is positively correlated with ocean productivity.
967   *Environmental Microbiology* **12:** 490-500.
968
969   Smith DP, Thrash JC, Nicora CD, Lipton MS, Burnum-Johnson KE, Carini P *et al.* (2013a).
970   Proteomic and Transcriptomic Analyses of "Candidatus Pelagibacter ubique" Describe the
971   First PII-Independent Response to Nitrogen Limitation in a Free-Living
972   Alphaproteobacterium. *mBio* **4:** 12.
973
974   Smith DP, Nicora CD, Carini P, Lipton MS, Norbeck AD, Smith RD *et al.* (2016). Proteome
975   Remodeling in Response to Sulfur Limitation in "Candidatus Pelagibacter ubique".
976   *mSystems* **1:** 16.
977
978   Smith MW, Allen L, Allen AE, Herfort L, Simon HM (2013b). Contrasting genomic properties
979   of free-living and particle-attached microbial assemblages within a coastal ecosystem.
980   *Front Microbiol* **4**.
981
982   Song J, Oh H-M, Cho J-C (2009). Improved culturability of SAR11 strains in dilution-to-
983   extinction culturing from the East Sea, West Pacific Ocean. *FEMS Microbiology Letters* **295:**
984   141-147.
985
986   Stamatakis A, Hoover P, Rougemont J (2008). A Rapid Bootstrap Algorithm for the RAxML
987   Web Servers. *Systematic Biology* **57:** 758-771.
988
989   Steindler L, Schwalbach MS, Smith DP, Chan F, Giovannoni SJ (2011). Energy Starved
990   Candidatus Pelagibacter Ubique Substitutes Light-Mediated ATP Production for
991   Endogenous Carbon Respiration. *PLOS ONE* **6:** e19725.
992
993   Stingl U, Tripp HJ, Giovannoni SJ (2007). Improvements of high-throughput culturing
994   yielded novel SAR11 strains and other abundant marine bacteria from the Oregon coast
995   and the Bermuda Atlantic Time Series study site. *ISME J* **1:** 361-371.
996
997   Sun J, Steindler L, Thrash JC, Halsey KH, Smith DP, Carter AE *et al.* (2011). One Carbon
998   Metabolism in SAR11 Pelagic Marine Bacteria. *PLOS ONE* **6:** e23973.
999

1000   Sun J, Todd JD, Thrash CJ, Qian Y, Qian MC, Temperton B *et al.* (2016). The abundant marine
1001   bacterium Pelagibacter simultaneously catabolizes dimethylsulfoniopropionate to the
1002   gases dimethylsulfide and methanethiol. *Nature Microbiology* **1**.
1003
1004   Thrash J, Seitz KW, Baker BJ, Temperton B, Gillies LE, Rabalais NN *et al.* (2017). Metabolic
1005   Roles of Uncultivated Bacterioplankton Lineages in the Northern Gulf of Mexico "Dead
1006   Zone". *mBio* **8**.
1007
1008   Thrash JC, Temperton B, Swan BK, Landry ZC, Woyke T, DeLong EF *et al.* (2014). Single-cell
1009   enabled comparative genomics of a deep ocean SAR11 bathytype. *ISME J* **8:** 1440-1451.
1010
1011   Tripp HJ, Kitner JB, Schwalbach MS, Dacey JWH, Wilhelm LJ, Giovannoni SJ (2008). SAR11
1012   marine bacteria require exogenous reduced sulphur for growth. *Nature* **452:** 741-744.
1013
1014   Tsementzi D, Wu J, Deutsch S, Nath S, Rodriguez-R LM, Burns AS *et al.* (2016). SAR11
1015   bacteria linked to ocean anoxia and nitrogen loss. *Nature* **536:** 179-183.
1016
1017   Tseng C-H, Chiang P-W, Shiah F-K, Chen Y-L, Liou J-R, Hsu T-C *et al.* (2013). Microbial and
1018   viral metagenomes of a subtropical freshwater reservoir subject to climatic disturbances.
1019   *ISME J* **7:** 2374-2386.
1020
1021   Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S *et al.* (2014). Pilon: An
1022   Integrated Tool for Comprehensive Microbial Variant Detection and Genome Assembly
1023   Improvement. *PLOS ONE* **9**.
1024
1025   Wickham H (2011). ggplot2. *Wiley Interdisciplinary Reviews: Computational Statistics* **3:**
1026   180-185.
1027
1028   Wilhelm LJ, Tripp JH, Givan SA, Smith DP, Giovannoni SJ (2007). Natural variation in SAR11
1029   marine bacterioplankton genomes inferred from metagenomic data. *Biology Direct* **2:** 1-19.
1030
1031   Zaremba-Niedzwiedzka K, Viklund J, Zhao W, Ast J, Sczyrba A, Woyke T *et al.* (2013). Single-
1032   cell genomics reveal low recombination frequencies in freshwater bacteria of the SAR11
1033   clade. *Genome Biol* **14:** 1-14.
1034
1035   Zhao X, Schwartz CL, Pierson J, Giovannoni SJ, McIntosh RJ, Nicastro D (2017). Three-
1036   Dimensional Structure of the Ultraoligotrophic Marine Bacterium "Candidatus Pelagibacter
1037   ubique". *Appl Environ Microbiol* **83:** 16.
1038
1039   Zhao Y, Ben T, Thrash JC, Schwalbach MS, Vergin KL, Landry ZC *et al.* (2013). Abundant
1040   SAR11 viruses in the ocean. *Nature* **494:** 357-360.
1041
1042

Candidatus
Fonsibacter ubiquis
1,160,202 bp

HVR2

100001

200001

300001

400001

500001

600001

700001

800001

900001

1000001

1100001