**Genome-wide analysis of ivermectin response by *Onchocerca volvulus***
**reveals that genetic drift and soft selective sweeps contribute to loss of**
**drug sensitivity**

Stephen R. Doyle [1,2*+], Catherine Bourguinat [3+], Hugues C. Nana-Djeunga [4,5],
Jonas A. Kengne-Ouafo [6], Sébastien D.S. Pion [7], Jean Bopda [8], Joseph
Kamgno [5,8], Samuel Wanji [6], Hua Che [3], Annette C. Kuesel [9], Martin Walker
[10], Maria-Gloria Basáñez [10], Daniel A. Boakye [11], Mike Y. Osei-Atweneboana
[12], Michel Boussinesq [7], Roger K. Prichard [3*], Warwick N. Grant [1*]

1. Department of Animal, Plant and Soil Sciences, La Trobe University, Bundoora, 3086, Australia
2. Wellcome Trust Sanger Institute, Hinxton, Cambridge, CB10 1SA, United Kingdom
3. Institute of Parasitology, McGill University, 21111 Lakeshore Road, Sainte Anne-de-Bellevue, Québec, H9X3V9 Canada.
4. Parasitology and Ecology Laboratory, Department of Animal Biology and Physiology, Faculty of Science, P.O. Box 812, University of Yaoundé 1, Yaoundé, Cameroon
5. Centre for Research on Filariasis and other Tropical Diseases (CRFilMT), P.O. Box 5797, Yaoundé, Cameroon
6. Research Foundation in Tropical Diseases and the Environment (REFOTDE), P.O. Box 474, Buea, Cameroon
7. Institut de Recherche pour le Développement (IRD), IRD UMI 233 TransVIHMI – Université Montpellier – INSERM U1175, 911 avenue Agropolis, P.O. Box 64501, 34394 Montpellier Cedex 5, France
8. Faculty of Medicine and Biomedical Sciences, University of Yaoundé 1, P.O. Box 1364, Yaoundé, Cameroon
9. UNICEF/UNDP/World Bank/World Health Organization Special Programme for Research and Training in Tropical Diseases (WHO/TDR), World Health Organization, 20, avenue Appia, 1211, Geneva, Switzerland
10. London Centre for Neglected Tropical Disease Research, Department of Infectious Disease Epidemiology, Faculty of Medicine, School of Public Health, Imperial College London, W2 1PG, United Kingdom
11. Noguchi Memorial Institute for Medical Research, University of Ghana, P.O. Box LG 581, Legon, Ghana
12. Council for Scientific and Industrial Research (CSIR), Department of Environmental Biology and Health Water Research Institute, P.O. Box M 32, Accra, Ghana

38  *Corresponding authors

39  Stephen R Doyle: sd21@sanger.ac.uk

40  Roger K Prichard: roger.prichard@mcgill.ca

41  Warwick N Grant: w.grant@latrobe.edu.au

44  + These authors contributed equally to this work
45

**Running title:**

*Population genomics of sub-optimal ivermectin response by Onchocerca*

*volvulus*

**Keywords**:

*Onchocerca volvulus*; ivermectin; Illumina genomics; Pool-seq; sub-optimal

drug response; population genetics; genetic drift; soft selection

**ABSTRACT**

*Background*

Treatment of onchocerciasis using mass ivermectin administration has reduced morbidity and transmission throughout Africa and Central/South America. Mass drug administration is likely to exert selection pressure on parasites, and phenotypic and genetic changes in several *Onchocerca volvulus* populations from Cameroon and Ghana - exposed to more than a decade of regular ivermectin treatment - have raised concern that sub-optimal responses to ivermectin's anti-fecundity effect are becoming more frequent and may spread.

*Methodology/Principal Findings*

Pooled next generation sequencing (Pool-seq) was used to characterise genetic diversity within and between 108 adult female worms differing in ivermectin treatment history and response. Genome-wide analyses revealed genetic variation that significantly differentiated good responder (GR) and sub-optimal responder (SOR) parasites. These variants were not randomly distributed but clustered in ~31 quantitative trait loci (QTLs), with little overlap in putative QTL position and gene content between countries. Published candidate ivermectin SOR genes were largely absent in these regions; QTLs differentiating GR and SOR worms were enriched for genes in molecular pathways associated with neurotransmission, development, and stress responses. Finally, single worm genotyping demonstrated that geographic isolation and genetic change over time (in the presence of drug exposure) had a significantly greater role in shaping genetic diversity than the evolution of SOR.

*Conclusions/Significance*

This study is one of the first genome-wide association analyses in a parasitic nematode, and provides insight into the genomics of ivermectin response and population structure of *O. volvulus*. We argue that ivermectin response is a polygenically-determined quantitative trait in which identical or related molecular pathways but not necessarily individual genes likely determine the extent of ivermectin response in different parasite populations. Furthermore,

90    we propose that genetic drift rather than genetic selection of SOR is the

91    underlying driver of population differentiation, which has significant

92    implications for the emergence and potential spread of SOR within and

93    between these parasite populations.

**Author summary**

Onchocerciasis is a human parasitic disease endemic across large areas of Sub-Saharan Africa, where more that 99% of the estimated 100 million people globally at-risk live. The microfilarial stage of *Onchocerca volvulus* causes pathologies ranging from mild itching to visual impairment and ultimately, irreversible blindness. Mass administration of ivermectin kills microfilariae and has an anti-fecundity effect on adult worms by temporarily inhibiting the development *in utero* and/or release into the skin of new microfilariae, thereby reducing morbidity and transmission. Phenotypic and genetic changes in some parasite populations that have undergone multiple ivermectin treatments in Cameroon and Ghana have raised concern that sub-optimal response to ivermectin's anti-fecundity effect may increase in frequency, reducing the impact of ivermectin-based control measures. We used next generation sequencing of small pools of parasites to define genome-wide genetic differences between phenotypically characterised good and sub-optimal responder parasites from Cameroon and Ghana, and identified multiple genomic regions differentiating the response types. These regions were largely different between parasites from both countries but revealed common molecular pathways that might be involved in determining the extent of response to ivermectin's anti-fecundity effect. These data reveal a more complex than previously described pattern of genetic diversity among *O. volvulus* populations that differ in their geography and response to ivermectin treatment.

## INTRODUCTION

*Onchocerca volvulus* is a filarial nematode pathogen responsible for causing human onchocerciasis. The infection is associated with significant morbidity, ranging from itching to severe dermatitis and from visual impairment to blindness. This morbidity and its economic impact have motivated large scale disease control programmes in the foci located in South and Central America, Yemen and throughout Sub-Saharan Africa, where more than 99% of the global at-risk population, estimated at 100 million people, live. Currently, onchocerciasis control is based primarily on annual community directed treatment with the macrocyclic lactone, ivermectin (CDTI).

Ivermectin has a least two pronounced effects on the parasites: (i) an acute "microfilaricidal effect" that results in the rapid and almost complete removal of microfilariae—the larval progeny of adult worms—from the skin within days to weeks after treatment, and (ii) a sustained "anti-fecundity effect" that results in prolonged but temporary inhibition of the release of new microfilariae from adult female worms into the skin for approximately three to six months [1]. Although some reports describe an increased proportion of adult dead worms following multiple ivermectin treatment rounds [2, 3], ivermectin is generally considered not to be a macrofilaricide [4, 5]. Ivermectin mass treatment therefore aims to: (i) prevent, and to the extent possible revert, pathology by removing microfilariae from the skin and eyes and delaying repopulation of these tissues with new microfilariae, and (ii) reduce transmission of *O. volvulus* by reducing the number of microfilariae that can be ingested by the blackfly (*Simulium*) vectors. Biannual (6-monthly) mass administration of ivermectin in hypo- and meso-endemic areas, and three-monthly ivermectin administration in hyperendemic areas of Central and South America have or are likely to have permanently eliminated transmission in most foci of the Americas [6-12]. Annual CDTI (expanded to biannual CDTI in some cases) and/or vector control, have or are likely to have also eliminated onchocerciasis in a number of endemic areas in Africa [13-19]. The World Health Organization Roadmap on Neglected Tropical Diseases has set the ambitious target of achieving onchocerciasis elimination where feasible in Africa by 2020 [20], and the African Programme for Onchocerciasis Control proposed expanding this goal to 80% of the countries by 2025 [21].

In some areas of Africa, however, persistent microfilardermia
(microfilaria in the skin) and transmission have been reported after 15-20
years of ivermectin treatment [22-33]. Already in 2004, an *O. volvulus*
ivermectin response phenotype termed "sub-optimal response" (SOR),
characterized by the presence of live stretched microfilariae in the uteri of the
adult worms 90 days after treatment, and consequent repopulation of the skin
with microfilariae earlier/more extensively than expected based on prior data,
was described in Ghana [29, 30]. Additional investigations have reported this
phenotype also in other areas in Ghana [27, 34, 35] and in Cameroon [36,
37]. Allele frequency change in a number of "candidate" ivermectin response
genes (chosen for analysis based on specific hypotheses concerning
mechanisms of resistance to the acute effects of ivermectin in *O. volvulus*)
has also been demonstrated in these populations when sampled before and
after several rounds of ivemectin treatment [38-44], which is consistent with
population genetic changes associated with drug selection pressure. While
genetic selection for SOR was not demonstrated, these studies suggest that
these populations are being influenced at the genetic level by ivermectin
treatment and, if these "candidate" genes mediate the phenotypic changes in
ivermectin response, that SOR has a genetic basis that may involve selection
on several genes.
The reproduction biology and transmission dynamics of the parasite
after ivermectin treatment have been described [5, 36, 37, 45]. However,
further work is required to understand: (i) the variability in response to
ivermectin that has been observed in ivermectin naïve parasite populations
[46-48], (ii) the biological mechanism(s) by which parasites may tolerate
and/or actively mitigate the inhibitory effects of ivermectin, and (iii) the
corresponding genetic architecture underpinning these biological
mechanisms, the influence of genetic selection, and the potential for SOR
genotypes to be transmitted preferentially within and between parasite
populations and thus increase in frequency. Although there has been some
debate regarding the existence of SOR to ivermectin in *O. volvulus* [49-53],
modeling of SOR – using individual-patient data on the rate of skin
repopulation by microfilariae following treatment in communities with different
histories of control [27, 54] – has provided support for the conclusion that the

early reappearance of microfilariae in the skin that defines SOR is most likely

due to a decreased susceptibility of the parasite to ivermectin's anti-fecundity

effect. Onchocerciasis morbidity reduction is driven primarily by ivermectin's

microfilaricidal effect but its anti-fecundity effect, delaying repopulation of the

skin with microfilariae, also contributes. More importantly, since both of these

effects are critical for achieving interruption of parasite transmission, an

increase in the frequency of SORs could jeopardise onchocerciasis

elimination goals [55].

Genome-wide approaches are increasingly being employed to

investigate the effects of drug selection in human pathogens, including but not

limited to, *Plasmodium falciparum* [56-60], *Schistosoma mansoni* [61],

*Leishmania donovani* [62] and *Salmonella enterica* Typhi [63]. Such

approaches have been instrumental in both confirming known and identifying

novel drug resistance-conferring loci in experimental and natural populations,

and in clarifying the roles played by selection, parasite transmission and

genetic drift in drug responses. An important feature of these studies is that

they do not rely on assumptions concerning mechanisms of resistance or

candidate resistance genes. The unbiased whole-genome approach has

proven particularly useful where there is a plausible hypothesis of polygenic

inheritance of a quantitative trait (QT, i.e. a trait that is determined by

interactions between multiple genes and the environment, and therefore will

have a continuous distribution of phenotypic values in a population) and when

the analysis is based on natural, outbreeding field populations in a non-model

species in which the population genetic structure is unknown but may

confound simple candidate gene analysis. Given that a number of different

candidate genes have been proposed to be associated with genetic response

to ivermectin treatment in *O. volvulus* and other parasitic nematode species

(see reviews from Gilleard [64] and Kotze *et al.* [65] for discussion of success

and limitations of candidate gene approaches), the hypothesis that variation in

response by the parasite to ivermectin is a polygenic quantitative trait is both

plausible and likely. Analyses to understand SOR have been limited by the

fact that *O. volvulus* is genetically diverse [66], populations are likely to be

genetically structured at an unknown spatial scale [67, 68], and the parasite is

a non-model outbreeding organism that is not amenable to more direct

controlled genetic crosses and quantitative trait loci (QTL) mapping methods.
These challenges are further exacerbated by limitations in providing an
accurate estimate of the degree of SOR in a given individual or population;
microfilarial density in the skin determined by skin snip is an indirect estimate
of SOR and is not precise, especially when the parasite density in the skin is
low due to ivermectin treatment, and analysis of the reproductive status of
adult worms by embryogram is typically (but not exclusively) qualitatively
measured although it is a quantitative trait. However, recent advances in our
understanding of the genome [69] and transcriptome/proteome of *O. volvulus*
[70], and more broadly, investment in publicly available helminth biology
resources such as WormBase Parasite (http://parasite.wormbase.org/, [71]),
have provided the means by which a comprehensive evaluation of ivermectin-
mediated selection of drug response in *O. volvulus* may now take place.
In this study, we present a genome-wide genetic analysis of drug
response by comparing genetic diversity within and between pools of adult *O.*
*volvulus* from Cameroon and Ghana that have been classified as "ivermectin-
naïve or little treated" [NLT]), "good responder" (GR), and ivermectin SOR
based on host population and/or individual host treatment history, microfilarial
repopulation in the host skin after ivermectin treatment and embryogram
analysis of female worms. This analysis has provided new insight into the
putative genetic and biological mechanism(s) of response by *O. volvulus* to
ivermectin. Underlying population structure, low susceptibility to ivermectin's
anti-fecundity effect, and potential for increase in the frequency of such
phenotypes are discussed in the context of efforts to control and eliminate
onchocerciasis in Africa.

**MATERIALS & METHODS**

*Sample history and phenotyping of ivermectin response*

**MATERIALS & METHODS**

*Sample history and phenotyping of ivermectin response*

*O. volvulus* samples used in this study were acquired from previously

described studies conducted in Cameroon [2, 36, 37] and Ghana [34, 35].

Ethical clearances were obtained for the original sampling of parasites from

the National Ethics Committee of Cameroon (041/CNE/MP/06), the Cameroon

Ministry of Public Health (D30-65/NHA/MINSANTE/SG/DROS/CRC and

D31/L/MSP/SG/DMPR/DAMPR/SDE/SLE), the Ethics Committee of Noguchi

Memorial Institute for Medical Research, (NMIMR-IRB CPN 006/01-04) and

McGill University (FWA 00004545).

Two separate experiments utilizing adult female worms collected from

people with known individual and/or community ivermectin treatment history

and response phenotype are reported here (see **Additional file 2: Figure S1**

for maps of sampling regions in each country). Phenotypic classification of

ivermectin response has been described in detail previously [34-37], and for

this study was determined based on a combination of host level (skin

microfilariae density) and female worm level (presence or absence of

stretched microfilariae *in utero*) characteristics (**Additional file 2: Table S1**).

A total of 108 adult female parasites were used in the Pool-seq

analyses; a description of the samples used, phenotypic characterization, host

and host-community ivermectin treatment history is presented in **Additional**

**file 2: Table S2**. Briefly, Pool-seq samples from Cameroon consisted of 3

pools of parasites composed of: (i) ~40 "drug-naïve or little treated" (NLT)

female worms from the Nkam valley, (ii) 22 ivermectin GR worms, and (iii) 16

SOR worms. The Cameroon GR and SOR parasites were collected from

people living in the Mbam valley who live in communities that have

participated in mass drug administration of ivermectin since 1994 (at least 13

years prior to sampling). In addition to these treatments, these people from

which parasites were collected also received between 4 to 13 doses of

ivermectin between 1994 and 1997 in a controlled clinical trial [2]. Similarly,

the Pool-seq samples from Ghana consisted of 3 pools of parasites made up

from: (i) 10 worms categorized as NLT, having been exposed to ivermectin

455 and 90 days prior to the time of sampling, (ii) 7 GR worms exposed to 11-

16 annual doses of ivermectin prior to sampling, and (iii) 13 SOR worms who
had been exposed to 9-16 annual doses of ivermectin.
A total of 592 adult female worms were used in the single worm
genotyping experiments; a summary of the worms selected for Sequenom
genotyping as well as the treatment history of the hosts and the area in which
the hosts live are described in **Additional file 2: Table S3**. Parasites from
Cameroon (n = 436) are categorized in three groups of samples: (i) the Nkam
Valley (NKA07; n = 140), from individuals who had been exposed to only a
single ivermectin treatment 80 days prior to sampling, i.e., considered NLT
[37], (ii) Mbam Valley (MBM07; n = 112), from people who had been exposed
to multiple rounds of ivermectin treatment prior to sampling as described
above [37], and (iii) parasites from individuals in the Mbam Valley who were
truly ivermectin naïve (MBM94; n = 184), sampled in 1994 prior to introduction
of CDTI [2]. Response phenotype data are not available for the MBM94
parasites. In total, 184 ivermectin naïve with unknown phenotype, 225 GR,
and 27 SOR parasites were genotyped. Worms from Ghana (n= 156) used for
Sequenom genotyping were sampled from 6 communities with different
ivermectin exposure histories (range: 2-17 treatments; **Additional file 2:**
**Table S3**), and were composed of 105 GR and 43 SOR parasites. An
additional 8 parasites that had been exposed to multiple annual ivermectin
treatments but whose phenotype is not available were also included.
300
301
***DNA preparation and genome resequencing***
DNA extraction from worms from Cameroon was performed in the REFOTDE
laboratory in Cameroon using an EZNA tissue DNA kit (Omega Bio-Tek Inc.,
Norcross, GA, USA). DNA extraction from worms from Ghana was performed
using the DNeasy® Blood and Tissue kit (Qiagen®, Toronto, ON, Canada) at
McGill University in Montreal, Canada.
The sequence data were generated from pools of worm DNA that were
prepared by combining the DNA of individual worms that shared drug
phenotype and geographic origin (i.e. NLT, GR or SOR from Ghana in 3
pools; NLT, GR and SOR from Cameroon in 3 pools; **Additional file 1: Table**
**S1**). DNA was pooled from individual worms at McGill University and

sequencing was carried out at the McGill University and Génome Québec

Innovation Centre (Montreal, Canada) across 8 Illumina GAII lanes

(Cameroon: GR and Ghana: SOR pools had sufficient DNA to allow two

sequencing lanes to increase sequencing depth). Overall, ~270-million 76-bp

single-end sequences were generated, resulting in an estimated 35.18-fold

(standard deviation [SD] ± 13.30) unmapped coverage per pool (or ~0.65–4

fold per worm if equal amounts of DNA/worm are assumed). Sequence data

are archived at the European Nucleotide Archive (ENA) under the study

accession PRJEB17785.

***Read mapping and variant calling***

Reads from each pool were aligned to the draft genome sequence

O_volvulus_Cameroon_v3 (WormBase Parasite; [69]) using BWA-MEM [72],

after which reads were aligned around putative indels and duplicate reads

removed using GATK (v3.3-0) [73]. Approximately 70% of the raw data were

mapped, resulting in an average mapped coverage of 24.47 ± 9.07 per pool.

Three 'pooled-sequencing' aware variant calling approaches were used to

analyse the pooled mapping data: CRISP [74], FreeBayes (using the "pooled-

continuous" parameter) (https://github.com/ekg/freebayes) and PoPoolation2

[75](**Additional file 1: Table S2**). To ensure that comparisons could be made

between all sequencing pools, the raw sequence nucleotide polymorphism

(SNP) data were filtered using the following criteria: (i) the mapped read depth

across a variant site was at least 8 reads but not greater than 3 SD from the

genome-wide mean read depth for the given pool; (ii) there was no evidence

of significant strand bias between forward and reverse reads; (iii) variants

were bi-allelic SNPs, and (iv) variants associated with reads mapped to

*Wolbachia* or mtDNA sequence were removed. The intersection of these

three approaches post-filtering yielded 248,102 shared variable sites relative

to the reference sequence.

***Genome-wide analyses of differentiation***

To perform the genome-wide analyses of genetic differentiation of the

sequencing pools, we used only variants found in the intersection of the three

Pool-seq-aware SNP calling approaches described above for all following

analyses. Given the statistical uncertainty associated with interpreting low

coverage read frequencies as a proxy for allele frequency for any given SNP

locus, and in turn, the need to use heavily-filtered variant call sets to enable

comparison between groups, we focused on analyses of relative genetic

diversity between groups by calculating $F_{ST}$ in 10-kb sliding windows across

the genome, and $F_{ST}$ of gene features based on the

O_volvulus_Cameroon_v3 genome build from WormBase Parasite

(http://parasite.wormbase.org/;

onchocerca_volvulus.PRJEB513.WBPS5.annotations). Both datasets were

generated using PoPoolation2 with the following parameters: --min-count 2, --

min-coverage 8, --max-coverage 2%, with corresponding individual haploid

pool sizes specified). Analyses of significance between different groups for

individual SNPs were performed using Fisher's exact tests to explore shared

variants present in both Ghana and Cameroon. Analyses of variant frequency

were performed using the variant read frequency generated from the CRISP

output.

One of the primary aims of this analysis was to define the physical map

positions of regions of the genome that differed significantly between

phenotypic classes in each country, on the assumption that such a region

would contain a locus or loci that contributed to the phenotypic difference

between the pools in question. We defined a genomic location as significant if

it consisted of two or more adjacent 10-kb windows (from the $F_{ST}$ sliding

window analysis) in which the GR *vs* SOR $F_{ST}$ values were greater than 3

SDs from the genome-wide mean $F_{ST}$ within a 50-kb window (unless

otherwise stated).

***Single worm genotyping by Sequenom® MassARRAY***

A subset of genome-wide SNPs (160 in total) were chosen to explore

ivermectin association and population structure further by Sequenom

genotyping of individual adult female worms phenotyped for ivermectin

response. DNA from 592 individual female worms (described in **Additional**

**file 2: Table S3**) was prepared for Sequenom® MassARRAY genotyping

(Sequenom, Inc., San Diego, CA, USA)[76]. Due to the DNA quantity

requirements for Sequenom analysis (600 ng per sample), many individual

worm DNA samples (401 of 436 samples from Cameroon, 96 of 156 samples

from Ghana) required whole genome amplification to increase the DNA

concentration. This was performed using the REPLI-g screening kit

(Qiagen®,Toronto, ON, Canada). DNA concentrations of all samples were

quantified using Quant-iT™ Pico Green dsDNA Assay Kit (Life Technologies

Inc, ON, Canada), before sending to the McGill University and Génome

Québec Innovation Centre for genotyping.

***Population- and ivermectin association analyses***

Sequenom data were analysed using PLINK [77]. Raw data were filtered

based on allele frequency (loci with <0.05 minor allele frequency were

removed; PLINK --maf 0.05), genotype quality (samples with <80% assay

success rate were removed: PLINK --mind 0.2; SNPs with <80% genotype

call frequency were removed: PLINK --geno 0.2). Filtered Sequenom data

(130 SNPs in 446 samples [**Additional file 2: Table S3**; sample numbers in

analysis are indicated in parentheses]; 81.25% of total SNPs and 75.34% of

total samples, respectively) were analysed by multidimensional scaling to

assess geographic vs phenotypic influence on genetic diversity. Hardy-

Weinberg equilibrium (HWE) was analysed using the same filtering conditions

described above (PLINK --hardy), using the "--keep" function to analyse

samples from each country separately. Discriminant analysis of principal

components (DAPC) and population assignment based on membership

probabilities were performed using the R package *adegenet* [78].

**RESULTS and DISCUSSION**

***Genome resequencing of pooled, phenotyped O. volvulus from Ghana***
***and Cameroon***

A genome-wide approach was used to detect genetic signatures associated
with the "sub-optimal responder" (SOR) phenotype of *O. volvulus* when
exposed to ivermectin. The criteria used to differentiate the SOR phenotype
from the "good responder" (GR) phenotype were based on criteria described
previously [29, 30]. In the present study, GR and SOR differ in two (deemed
to be related) ways: (i) at the host level, the number of microfilariae in the skin
determined by diagnostic skin snips is >7% of the pre-treatment value in SOR
but microfilariae are largely undetectable in GR around 3 months post
treatment (80 or 90 days for the samples from Cameroon and Ghana,
respectively), and (ii) at the parasite level, SOR macrofilariae contain
"stretched" microfilariae (ready to be released) *in utero* around 3 months post
treatment (80 or 90 days for the samples from Cameroon and Ghana,
respectively), as determined by embryogram, while GR do not. A SOR
parasite, therefore, causes earlier repopulation of the skin with microfilariae
than a GR parasite due to an earlier resumption of microfilarial release after
temporary inhibition of fecundity.

  To investigate the genetic differences between GR and SOR adult
*O. volvulus*, whole genome sequencing was performed on pools of adult
female worms from Cameroon and Ghana that were classified as "naïve or
little treated" [NTL], or multiply treated GR and SOR groups based on the prior
ivermectin exposure of the hosts and/or community from whom they were
collected, and the host and parasite level response to ivermectin as described
above (see **Additional file 2: Table S1** for response criteria and **Additional
file 2: Table S2** for characteristics of the worm pools). We identified 248,102
variable positions that were shared among all groups and passed our
stringent filtering criteria for inclusion in a genome-wide scan of genetic
differentiation within and between pools (**Additional file 1: Table S2**), at an
average marker density of 1 variant per ~389-bp (of the 96,457,494-bp
nuclear genome). This represented only 32.7% of the total putative variants
identified using the three variant analysis pipelines, primarily as a result of the
relatively relaxed variant calling conditions of PoPoolation2 (**Additional file 1:**

439    **Table S2;** 34.4% of SNPs called by PoPoolation2 were unique to this tool

440    under the conditions used compared to 2.6% and 6.2% of unique SNPs called

441    by CRISP and FreeBayes, respectively) and in part due to the stochastic

442    variation in allele detection in low sequence coverage Pool-seq data.

443            Pool-seq has been used to estimate population genetic diversity in a

number of different species [75, 79-82] on the assumption that individual read

frequencies at a variant site are a proxy for allele frequency. This approach

relies on sampling sufficient reads at any given position to be confident in

detecting the diversity present; the more reads sequenced at a given position,

the closer the read frequency is to the "true" allele frequency. Given that

approximately 24.47-fold mapped coverage per pool (range: 0.53-3.05-fold

coverage per worm) was obtained overall, it is unlikely that any of the pools

was sequenced at sufficient depth to sample each genome present. Analyses

of variation in read frequency between pools at low sequence coverage for

any given nucleotide variant should, therefore, be treated with caution as they

are confounded by significant statistical variation in coverage per genome

[83]. In addition to strict filtering of the variants (e.g. to remove putative variant

sites that were not detected by all three variant callers in all pools), we have

attempted to account for this uncertainty by focusing on genetic variation

calculated from multiple SNPs, either in sliding windows across the genome

or on whole genes, rather than on individual nucleotide variants. A significant

finding of this study was that genetic variation that differentiated GR and SOR

pools was not randomly distributed but was strongly clustered in multiple

discrete regions of genome. This observation establishes clearly, for the first

time, both the genetic architecture and likely mode of selection of ivermectin

response in *O. volvulus*, which is described in detail below.

***Genome-wide genetic differentiation between ivermectin response***

***phenotypes***

The two important questions with respect to the evolution of SOR are (i) which

locus or loci are under selection, and (ii) whether the same loci are under

selection in different populations, i.e., whether variation that differentiated

SOR and GR worms in Ghana also differentiated SOR and GR worms in

Cameroon. Shared variation between diverse geographic regions that

differentiated SOR from GR would provide candidate markers that may

predict ivermectin response in previously uncharacterised populations, and

thus form the basis for the development of a ubiquitously applicable tool for

monitoring the relative frequency of SOR and GR before and during CDTI.

Pairwise analyses of individual SNPs (p-values from Fisher's exact test;

**Figure 1A**) and $F_{ST}$ calculated from 10-kb windows (**Figure 1B**) both revealed

a higher degree of differentiation between SOR and GR pools from Cameroon

than between SOR and GR from Ghana, i.e., there were more loci or regions

above significance thresholds in the Cameroonian pools. This difference is

likely to be due to the unequal sample size between the two countries

(Cameroon = 22 GR and 16 SOR worms; Ghana = 7 GR and 13 SOR

worms); a greater proportion of total genetic diversity will be present in the

Cameroon dataset as more worms are present, however, at the same time,

the sequencing depth per Cameroon genome is lower than for the Ghana

samples, and hence, will increase the stochastic variation in the Cameroon

variant frequency. Although many single SNP loci from each country showed

significant variation between SOR and GR pools when each country was

analysed independently, only a single intergenic SNP (OM1b_7179218) was

significantly different between pools and common to both countries (**Figure**

**1A**; red dot) after a Bonferroni genome-wide correction (**Figure 1A**; dashed

lines) was applied. Furthermore, only six 10-kb regions that showed

significant deviation above a genome-wide mean $F_{ST}$ threshold of +3 SDs

between GR and SOR were shared between Ghana and Cameroon (**Figure**

**1B**; dashed lines; red dots; **Table S3**). Relaxing the threshold to +2 SDs

yielded 22 additional 10-kb regions that were able differentiate GR from SOR

and were shared between countries (**Figure 1B**, orange dots; **Table S3**). A

total of 28 $F_{ST}$ windows at > + 2SDs is only marginally more than the number

of windows that is expected to exceed +2 SDs by chance alone ($0.05^2$ x 9893

10-kb loci = 24.73), and may in fact be inflated considering: (i) 11 of the 28

10-kb windows were immediately adjacent to at least one other 10-kb window

in the genome and, therefore, are unlikely to all be segregating independently,

and (ii) four of these 28 10-kb windows contained sequences associated with

Pao retrotransposon peptidase- and integrase-related proteins and a further

two windows contained ribosomal subunits (**Additional file 1: Table S3**).

Given the multi-copy nature of these sequences, the high $F_{ST}$ value for these

6 windows is likely to be a technical error associated with poor read mapping

of multicopy sequences rather than true biological differentiation. These

results suggest that, even at a reduced genome-wide level of significance

threshold (i.e., > + 2 SDs), little variation that similarity discriminated GR from

SOR parasites was shared between the two countries.

To investigate the distribution of genome-wide genetic variation

between ivermectin response groups, we analysed the relative positions of

SOR vs GR 10 kb $F_{ST}$ windows > +3SD in the *O. volvulus* genome. A striking

finding was that the 10-kb regions that provided the most genetic

differentiation between SOR and GR were not randomly distributed but were

found in discrete clusters (**Figure 1C & D**); we interpret these clusters of

significantly differentiated 10-kb windows (defined as 2 or more adjacent 10-

kb windows within a 50-kb window > +3 SDs from the genome-wide mean $F_{ST}$

calculated separately for each country) as putative QTLs. These QTLs are

composed of a variable number of 10-kb windows and contain multiple genes.

In total, 18 putative QTLs that mapped to well-assembled regions of the

genome were found in the Cameroon data (**Figure 1C**; mean QTL size of

66,389-bp ± 55,157-bp SD) and 14 putative QTLs in the Ghana data (**Figure**

**1D**; mean size of 102,143-bp ± 95,690-bp SD), representing 1.23% and

1.48% of the ~96.4-Mb nuclear genome, respectively (**Additional file 1:**

**Table S4**). These data provide strong evidence that the SOR phenotype is a

polygenic quantitative trait (QT) and, further, given that only a single putative

QTL shared between Ghana and Cameroon was detected, suggest that

different putative QTLs may be under selection in these two geographically

separated parasite populations.

The apparent lack of concordance between Ghana and Cameroon

SOR QTLs demonstrates that these two populations are genetically distinct.

This observation, coupled with the number and location of putative QTLs,

suggests that soft selection on pre-treatment genetic variation in the two

parasite populations has acted on different loci in each country, and in turn,

resulted in a different signature of selection in each country. Soft selection

also implies that it may therefore be difficult to separate differences between

Ghana and Cameroon SOR populations that are the result of selection from

those that are the result of drift. However, the pattern of genetic variation

within each country (see below) is consistent with the conclusion that the

differentiation between GR and SOR worms (within the same country) is

associated with phenotypic differences in ivermectin response (for

background information on hard vs. soft selection, see **Additional File 2:**

**Section 2**).

**Analysis of between-country genetic variation, and genetic diversity**

**between responder phenotypes and "drug-naïve" worms**

To test more explicitly for population structure between the two countries and

thus, better understand the extent to which the standing genetic variation (see

**Additional File 2: Section 2**) in these populations may have been shaped by

the combination of selection and genetic drift, we analysed genetic

differentiation between and within the two countries across all three treatment

history/response categories (i.e. NLT, GR and SOR). A comparison of

genome-wide allele frequency correlation within and between countries

demonstrated that there was a significantly higher degree of shared variation

(i.e., less population structure) within each country than there was between

countries, where allele frequency correlation was low **(Figure 2A).** This

supports the conclusion that there is significant genetic differentiation between

the parasite populations in the two countries and that any genetic signal

associated with SOR that might have been common to both countries would

likely be masked by the presence of significant population genetic structure.

Somewhat unexpectedly, a comparison of allele frequency correlations

between response groups within each country suggested that, for both

Cameroon and Ghana, the NLT worms were genetically more similar to the

SOR worms than to the GR worms **(Figure 2A).** A direct pairwise comparison

of the three treatment history/response categories within each country using

genome-averaged $F_{ST}$ values (calculated from 9,893 10-kb windows) was

consistent with this observation: NLT *vs* SOR $F_{ST}$ genomic medians were

significantly smaller (Cameroon: 0.059; Ghana: 0.068) than either NLT *vs* GR

medians (Cameroon: median = 0.095; two-sample Kolmogorov-Smirnov [KS]

test: D = 0.0438, p < 0.001; Ghana: median = 0.110; KS D = 0.575, p <

0.001), or GR *vs* SOR medians (Cameroon: median = 0.112, KS D = 0.544, p

< 0.001; Ghana: median = 0.104, KS D = 0.495, p < 0.001) (**Figure 2B**). The

closer relationship between NLT and SOR than between NLT and GR is

particularly surprising for Cameroon when one considers that the NLT and

GR/SOR populations are from geographically distinct areas, i.e., the sampling

sites within Nkam and Mbam Valleys were >100-km apart in two separate

river basins separated by the Western High Plateau of Cameroon (see

**Additional File 2: Figure S1 for map of sampling sites)**. However,

seasonal dispersal of the local vector species, *Simulium squamosum*, has

been observed in Cameroon over greater distances than the distance

between the two sampling regions here [84], and therefore, some seasonal

transmission between the two river basins may occur and requires further

investigation.

The observation of higher genetic similarity between the NLT and SOR

populations supports the hypothesis of soft selection for SOR because it

suggests that selection for SOR from an ivermectin naive population led to

relatively little genome wide reduction in genetic variation. A characteristic

prediction of soft as opposed to hard selection (see **Additional file 2: Section**

**2** for an extended discussion of soft- versus hard-selection) is that unlike hard

selection, soft selection should not markedly reduce the genome-wide genetic

diversity following selection. Since the SOR phenotype is associated with an

early resumption of reproduction and, consequently, with the early availability

of SOR offspring in the skin for vectors to ingest, ivermectin treatment likely

interrupts the transmission of SOR genotypes for only a relatively short period

of time between CDTI rounds compared to GR transmission interruption. If

SOR alleles are already present on many different genetic backgrounds in the

NLT starting population, this continued transmission of SOR genotypes under

drug treatment will maintain genetic diversity in the SOR population, which is

consistent with what we have observed in the present study.

The strong, genome-wide, between country genetic differentiation in

the NLT populations implies that the standing genetic variation from which

SOR is selected varies significantly between Ghana and Cameroon, and

explains why the genetic outcome of selection of SOR differs between

countries (as shown by the lack of concordance between Ghana SOR and

Cameroon SOR populations). The presence of strong population structure

between Cameroon and Ghana is not unexpected. The *O. volvulus* lifecycle is

characterised by significant population bottlenecks at each transmission

event: only a miniscule proportion of microfilariae in the skin are ingested by

blackflies, and very few are subsequently transmitted to humans and establish

as adult worms [85]. Repeated bottlenecks increase the severity of genetic

drift by strongly enhancing the stochastic processes that generate genetic

diversity between populations, independent of drug treatment, suggesting that

genetic drift had created genetic differentiation between different parasite

populations before initiation of CDTI, and that subsequent soft selection of

SOR genotypes from these genetically distinct populations has led to SOR

populations that are genetically distinct despite their phenotypic similarity.

In contrast, CDTI interrupts transmission of GR genotypes for a longer

period of time. If this reduces the proportion of GR worms that contributes to

the next generation, it will lead to a loss of genetic diversity and increase in

genetic drift in subsequent GR populations relative to both the population

before CDTI was initiated and the SOR population. This expectation is

consistent with genetic change described in single-gene studies following

treatment over time in *O. volvulus* [38-44].

To investigate the impact of soft selection on GR and SOR genetic

diversity further, we estimated the genetic diversity within the group of GR

worms and the group of SOR worms from each country in two complementary

ways: (i) by calculating the variant read spectrum per group (as a proxy for the

allele frequency spectrum, **Figure 2C**), and (ii) from the relative proportion of

"invariant" SNP loci, which we have defined as SNP loci with variant read

frequencies <0.05 or > 0.95 (**Figure 2D**). These complementary analyses

provide insight into the degree of genetic variation present within each drug

response parasite pool, and will detect loss of genetic diversity. The GR

worms from both countries were less diverse and had a greater proportion of

invariant loci than the SOR and NLT parasites, particularly in Cameroon,

whereas the SOR and NLT parasites had similar levels of genetic diversity.

In conclusion, comparisons of genetic diversity and genetic similarity

between NLT, GR and SOR support our hypothesis of soft selection for SOR,

and explain both the relatively subtle signature of selection in the SOR

parasites and the strong population structure observed between the SOR

parasites from Cameroon and Ghana.

***Characterisation of molecular pathways identified from genes within***

***QTLs that differentiate ivermectin response***

In light of evidence suggesting that genetically separated parasite populations

contain different standing genetic variation (likely to be the result of genetic

drift) before introduction of CDTI, and thus that ivermectin-mediated soft

selection may produce different genetic outcomes in SOR parasites between

Cameroon and Ghana, it was of interest to compare genes within the putative

QTLs identified in the SOR worms from Ghana and Cameroon. While we

identified only a single putative QTL that was common to the SOR parasites

from the two countries, the putative QTLs from each country contained genes

encoding proteins that act in a limited number of molecular pathways. This

implies that although different genes may be under selection in Ghana and

Cameroon, there may be a common biological mechanism that confers SOR

in both countries (see **Table 1** for a summary of genes with common

functional characteristics and/or shared pathways identified in the QTLs, and

**Additional file 1: Table S5** for characterization of all genes within each QTL).

The most prominent group of genes under the putative QTLs defined

above were associated with neurotransmission (17 genes in 14 QTLs), which

was encouraging considering that the primary target of ivermectin is a ligand-

gated channel at neuromuscular junctions [86]. Given that the duration of the

anti-fecundity effect of ivermectin distinguishes the GR and SOR phenotypes,

the fact that nine of those genes (in eight QTLs) were associated with

acetylcholine signaling is of particular interest because acetylcholine signaling

plays an important role in the regulation of egg laying in *C. elegans,* and

therefore, may be relevant to ivermectin's anti-fecundity effect in *O. volvulus*.

These genes include a number of ion channels (*Ovo-acc-1, Ovo-lgc-46, Ovo-*

*lgc-47*), and components of acetylcholine synthesis (*Ovo-cha-1*), transport

(*Ovo-unc-17, Ovo-aex-3*), and regulation (*Ovo-pha-2, Ovo-snb-1, Ovo-emc-6,*

*Ovo-nrfl-1*). The *unc-17* and *cha-1* mutants in *C. elegans* exhibit hyperactive

egg-laying phenotypes associated with defects in laying inhibition [87, 88],

and acetylcholine activation of egg laying in *C. elegans* is regulated by

neuropeptides, serotonin and glutamate [89]. Furthermore, acetylcholine

receptors have recently been proposed to be involved in the development of

the nervous system during embryo- and spermatogenesis in the filarial

parasite, *Brugia malayi* [90]. In addition, two recent studies demonstrated

inhibition of L-AChR receptors in *C. elegans* by ivermectin [91] and

antagonistic effects of abamectin on nicotinic acetylcholine receptors [92],

implying that under some circumstances ivermectin may act directly on

acetylcholine signaling. We therefore hypothesise that modification of

neurotransmission in general, and acetylcholine signaling pathways in

particular, may contribute to ivermectin SOR and reflects an overall

adjustment in neuromuscular signaling that mitigates the effects of ivermectin.

This variation may also contribute to the changes in fecundity that have been

associated with the SOR phenotype [93] by changing the way in which

neurotransmission might influence the release of microfilariae in an analogous

fashion to neuronal control of egg laying in *C. elegans*. Given the enrichment

of neuronal genes that may play a role in regulating reproduction, it is of

interest to note that the single QTL that is shared between Cameroon and

Ghana SOR parasites includes *Ovo-aex-3,* a neuronal protein and regulator of

synaptic transmission. This gene may be of particular interest because the *C.*

*elegans* orthologue plays a role in reproduction and also in regulation of

pharyngeal pumping [80] (a phenotype that is also consistent with reduced

sensitivity to ivermectin in *C. elegans*). In addition, this QTL also includes

*Ovo-stg-1*, which may be of interest due to its putative chaperone-like role in

regulating ionotropic glutamate receptor (iGluR) function and a hypothesised

role in protecting neurons from excitotoxicity or inappropriate depolarisation in

*C. elegans* [81].

Nine putative QTLs contained genes associated with stress responses,

including heat-shock proteins (*Ovo-hsp-17, Ovo-hsb-1*), and genes required

for the synthesis (*Ovo-acs-16, Ovo-fat-4, Ovo-spl-1, Ovo-tat-2*), regulation

and storage (*Ovo-obr-2, Ovo-sms-1, Ovo-math-46, Ovo-mtd-15, Ovo-cuc-1*)

of lipids. Variation in gene expression in lipid metabolism-encoding genes

following ivermectin treatment in *C. elegans* has been described previously

[94], which was interpreted as a metabolic adaptation to starvation due to

ivermectin inhibition of pharyngeal pumping but may also be a more general

indicator of organismal stress. Genes associated in general with stress

responses, including lipid metabolism, are often reported in genome-wide

analyses for loci under drug selection pressure [56].

Other genes under the putative QTLs include those acting in pathways

involved in developmental processes, including muscle assembly and myosin

organization (three QTLs contained 6 loci; *Ovo-mup-4, Ovo-mua-3, Ovo-mlc-*

*5, Ovo-unc-82, Ovo-nmy-1, Ovo-sca-1*), and germline and larval development

signals associated with notch signaling (three QTLs), specifically with the

suppression (*Ovo-bre-5*) or cleavage (*Ovo-pen-2,Ovo- crb-1*) of the LIN-12

receptor. The relevance, if any, of these developmental genes to ivermectin

response is not known.

This list of genes that fall under the QTLs leads to a hypothesis of SOR

as tolerance to ivermectin's anti-fecundity effect brought about by a "re-tuning"

of neuronal function in combination with a stress response. We acknowledge

the tentative nature of this hypothesis given: (i) the limitations in the resolving

power of genetic-association analysis based on modest sequencing depth of

pooled samples, and only a single comparison between GR and SOR

parasites from each country, (ii) the inability to carry out more specific

analysis of coding sequence variation imposed by limited genetic resolution,

(iii) the large number of predicted genes that lack functional annotation in the

*O. volvulus* genome (~67.4% of the coding sequences within the QTLs are

unannotated or hypothetical), and (iv) the methods available for phenotype

classification, which has limited sensitivity for the determination of the

presence and extent of microfilariae density in the skin [95] and is qualitative

with respect to the presence or absence of live stretched microfilariae *in utero*,

which in turn decreases the power of genetic association. However, the

enrichment of neurotransmission and stress response genes in the QTLs that

differentiate GR and SOR parasites in two geographically independent

populations is striking, and does provide support for our hypothesis of a

polygenic quantitative trait characterised by earlier recovery from the acute

effects of ivermectin on fecundity.

If correct, this hypothesis implies that the SOR adult worms remain

sensitive to the acute effects of ivermectin on fecundity but recover more

quickly than GR worms. Early recovery from the acute effect of ivermectin on

fecundity does not require a mechanism that is specific to the mode of action

of that acute effect, so a polygenic "retuning" or "buffering" of the neuronal

regulation of reproduction that leads to earlier recovery of fertility is

biologically plausible. It may also explain why our analyses have failed to

validate "candidate genes" that would protect worms against ivermectin's

acute effects (see below).

**Analysis of "candidate" ivermectin-resistance genes**

Given the extensive literature focused on "candidate" genes (i.e. genes

chosen for analysis based on specific hypotheses concerning mechanisms of

resistance to the acute effects of ivermectin in *O. volvulus*), and their apparent

absence from the QTLs described here, it was important to investigate these

genes, which included glutamate-gated chloride channels (*Ovo-avr-14, Ovo-*

*glc-2, Ovo-avr-15, Ovo-glc-4*), p-glycoproteins (*Ovo-pgp-1, Ovo-pgp-10, Ovo-*

*pgp-11, Ovo-plp-1*), ABC transporters (*Ovo-abcf-1, Ovo-abcf-2, Ovo-abcf-3*)

and other candidates (Ovo-*ben-1* [beta tubulin], *Ovo-lgc-37, Ovo-mrp-7, Ovo-*

*dyf-7*). *Ovo-abcf-1* was the only "candidate" gene found in a QTL. For the

majority of the other genes, no significant SOR *vs* GR $F_{ST}$ differences were

observed (**Additional file 1: Table S6).** *Ovo-abcf-1* (in QTL-5), *Ovo-abce-1,*

*Ovo-dyf-7, and Ovo-pgp-10* did show moderate levels of differentiation, but

none were statistically significant. Given that our data suggests that parasite

populations are structured and that alleles associated with response may vary

between populations, we cannot exclude that these loci contribute to

variability in ivermectin susceptibility in other *O. volvulus* populations.

However, considering the evidence of multiple putative QTLs and strong

geographic population structure, it is possible that single "candidate" gene

comparisons may have been confounded by population structure and the

polygenic nature of the trait. We conclude that there was no evidence of

significant genetic differentiation in these candidate genes between the SOR

and GR populations compared here.

Three of the 31 putative QTLs did contain "neuronal function" genes

that have been linked previously to ivermectin sensitivity in *C. elegans* (**Table**

**1; Additional file 1: Table S5**): *Ovo-unc-44*, a likely orthologue of a *C.*

*elegans* gene that influences ivermectin sensitivity and is involved in neuronal

development [96], *Ovo-inx-5*, which in *C. elegans* encodes an innexin gap

junction protein that is associated with the pharyngeal motor neurons and is

related to *unc-9* (a known ivermectin-resistance allele [97]), and *Ovo-klp-11*, a

kinesin motor protein found in the cilia of chemosensory neurons in

*C. elegans*, which has significant homology to, and is a likely binding partner

of, *Ovo-osm-3*, alleles of which have been described to decrease sensitivity to

ivermectin in *C. elegans* [96].

***Geographic and genetic distribution of ivermectin susceptibility***

Given the importance of the distinction between population structure as

a result of genetic drift and phenotypic differentiation as a result of selection

for SOR, we sought to investigate ivermectin response genetics in a larger

cohort of individual female worms (most having been characterized for their

ivermectin response phenotypes via embryograms [34-37]; **Additional file 2:**

**Table S3**) by genotyping individual worms at 160 SNP loci by Sequenom

(**Additional file 1: Table S7**). The loci were chosen based on the original

Pool-seq analyses to determine the degree of genetic association with

response phenotype, and to characterize genetic structure within these

populations. Multidimensional scaling analysis (MDS) was used to interrogate

the Sequenom genotyping data from 446 female worms at 130 SNPs that

passed filtering criteria. Three aspects of the data are highlighted (in different

panels, **Figure 3**): (i) between country population structure, (ii) within country

population structure, and (iii) differentiation between response phenotypes.

Consistent with the Pool-Seq analysis, we observed clear

differentiation between worms from Ghana and worms from Cameroon: there

were very few [blue] Ghanaian worms mixed with the [red] Cameroon worms

(**Figure 3A**), regardless of response phenotype. Analysis of worms by their

community of origin to allow within as well as between country comparisons

did not reveal any internal genetic structure in the Ghanaian populations or

between the Cameroonian NKA07 and MBM07 [pink and red, respectively]

populations (**Figure 3B**).

Interpretation of the Cameroon data is more complex than for Ghana

due to the likely existence of underlying population structure that is both

temporal and spatial in origin: within Cameroon, MBM94 [light orange] was

genetically distinct from both NKA07 and MBM07 populations but NKA07 and

MBM07 were indistinguishable. MBM94 and MBM07 are two samples drawn

from the same geographical location in Cameroon (the Mbam valley) at

different times: MBM94 was sampled in 1994 before ivermectin distribution

began, and is drug-naïve, whereas MBM07 was sampled in the Mbam Valley

after 13 years of annual CDTI and contains approximately 16% SOR and 84%

GR parasites (n = 112). Given that MBM07 is still largely GR, we interpret this

signal of strong population differentiation as the result of genetic drift in

MBM94, i.e., of largely stochastic (rather than deterministic, i.e., selection)

genetic change likely brought about by multiple population bottlenecks

imposed by the impact of 13 years of ivermectin treatment on parasite

transmission and hence on effective population size. If this interpretation is

correct, the striking similarity between MBM07 and NKA07 is coincidental:

genetic drift is a stochastic process, and previously distinct populations may

become more similar by chance alone.

Alternatively, the similarity between NKA07 and MBM07 may indicate

that there is transmission between NKA07 and MBM07, such that as

ivermectin shrank the originally naïve MBM94 population over 13 years,

exchange of parasites between the Nkam Valley and the Mbam valley

mediated by vector or human migration brought these two initially dissimilar

populations closer together genetically. While the Nkam valley had not

received CDTI at the time the NKA07 samples were collected (the samples

genotyped are classified as NLT, having only been exposed to a single

experimentally controlled round of ivermectin treatment [37]), some

communities in the Nkam valley are only around 10 km from communities

across the Nkam river which have received annual CDTI for >10 years.

Consequently, the NKA07 parasites may not be entirely "naïve". However,

transmission of multiply-exposed parasites from these villages into NKA07

patients is unlikely to account for the similarities between NKA07 and MBM07,

as the NKA07 population was largely composed of GR parasites (94.3%, n =

140).

Our data do not allow discrimination between these two hypotheses. In

either case, however, the Cameroon data underline how little is known about

the determinants of *O. volvulus* population structure in general, and the

genetic impact of population bottlenecks imposed by ivermectin-mediated

temporary interruption and long-term reduction of transmission in *O. volvulus.*

These data also illustrate the confounding effect of underlying population

structure on our ability to distinguish between selection and drift as drivers of

genetic differentiation between two population samples, regardless of the

origin of that structure, whether due to a drug induced population bottleneck,

natural drift over time or parasite immigration from a drug naïve into a drug

treated population.

Analysis of the distribution of response phenotypes revealed no clear

separation of GR (green) from SOR (red) in either Ghana or Cameroon

samples (**Figure 3C**). **Additional File 2 Table S3** and careful comparison of

**Figures 3B** and **3C** show that the Cameroon NKA07 parasite group in **Figure**

**3B** (pink dots) is composed almost entirely of GR worms (SOR frequency of

approximately 5%, n = 140), whereas the MBM07 parasite group (**Figure 3B**,

red dots) is composed of a mix of GR worms (**Figure 3C**, green dots) and

SOR worms (**Figure 3C**, red dots; SOR frequency approximately 16% [n =

112]), but that the GR and SOR worms in both NKA07 and MBM07 are

intermingled, i.e. the differentiation between (NKA07 + MBM07) and MBM94

in **Figure 3B** is unrelated to the distribution of drug phenotypes in **Figure 3C**.

Thus, these data suggest that genetic differentiation between the Cameroon

populations is not correlated with response phenotype.

To further characterise the differences between GR and SOR parasites

within each country, we analysed genotype frequencies for Hardy-Weinberg

equilibrium (HWE). A greater number of SNPs were significantly out of

equilibrium in the GR populations compared to the SOR populations from both

countries (significance threshold of $p < 0.05$; 111 *vs* 20 (of 131) SNPs in the

Cameroon GR *vs* SOR parasites, **Additional file 1: Table S8;** and 80 *vs* 68

(of 121) SNPs in the Ghana GR *vs* SOR parasites, **Additional file 1: Table**

**S9**). In this respect, the Sequenom genotype and the whole genome PoolSeq

data are concordant: greater deviation from HWE in the GR populations is

best explained by ivermectin treatment reducing effective population size and

genetic diversity and increasing genetic drift in the GR worms, but not in the

SOR worms or naïve populations.

Collectively, these Sequenom data suggest that spatial (between

Ghana and Cameroon) and temporal (pre- and post 13 years of drug

exposure within Cameroon) genetic differentiation were readily detectable by

genotyping at these 130 SNP loci, but that this genotyping failed to detect the

relatively weaker signal of genetic differentiation between GR and SOR

phenotypes. This is illustrated most clearly by comparison of the Ghanaian

and Cameroon SOR populations (**Figure 3C**), where the MDS coordinates of

Ghanaian SOR individuals cluster with the Ghanaian GR individuals, and

Cameroon SOR individuals cluster with Cameroon GR, evidence that SOR

alleles exist in genetic backgrounds determined by their population of origin,

and that the genetic signature of soft selection for SOR in different naïve

populations is weak compared to the pre-existing population structure that is

the product of genetic drift. This is further supported by an analysis of

genotype-based assignment of individual parasites to their respective

populations (**Additional file 2: Section 3**). Greater that 99.5% of individuals

were correctly assigned to their country of origin by their genotype, and

between 40-92.4% of individuals from Cameroon and 0-97% of individuals

from Ghana to their respective communities or regions. In contrast, 0% of

Cameroon worms and only 8.1% of Ghanaian worms were correctly assigned

as SOR worms on the basis of their genotypes, i.e., there is strong correlation

between Sequenom genotype and place of origin but little or no correlation

between Sequenom genotype and ivermectin response phenotype.

Furthermore, the MDS analysis of the three Cameroon groups of samples

(**Figure 3C**) supports the view that not only do past histories of geographic or

ecological separation result in genetic drift and hence differentiation between

*O. volvulus* populations, but that reduction or interruption of transmission and

reduction in population size by ivermectin, and/or migration from untreated

into treated populations also drives drift and masks selection.

***Implications for control and elimination***

The success of CDTI in the majority of areas in Africa where CDTI has been

implemented with good coverage [17], suggests that the alleles responsible

for SOR are not common across Africa and that CDTI can interrupt

transmission in many populations, depending on epidemiological and

programmatic factors such as the intensity of blackfly biting on humans and

the frequency and achieved coverage of CDTI [28, 33, 98-100].

However, it is also clear that drug response by *O. volvulus* is not

uniform in all ivermectin-naïve populations [5, 27, 37, 46, 47], and that in

some naïve populations (such as those sampled here), parasites with sub-

optimal response to ivermectin's anti-fecundity effect are present prior to

ivermectin treatment. The data presented here provide an explanation for the

presence of SOR worms in ivermectin-naïve populations, and support for the

view that sub-optimal response to ivermectin in those populations is a

genetically determined trait that can increase in frequency as a result of

selection. Selection for SOR has progressed to an extent in the populations

sampled for analysis here that the genetic signal of that selection can be

detected in the Pool-seq data. The genetic signal is, however, weak because

it is based on soft selection acting on many genes that contribute to a

quantitative trait. The implications of these conclusions for control and

elimination hinge on three crucial questions.

The first question is whether, and under what circumstances, SOR can

be detected genetically. The whole genome Pool-seq data demonstrate that it

is possible to detect SOR genotypes in an analysis of 248,102 SNPs but our

first attempt to validate SOR alleles in individual worms by reducing the

number of SNP's to a panel 130 SNPs in Sequenom genotyping failed. This

failure is likely because soft selection for SOR has left a faint genetic

signature in the SOR populations compared to the very strong, pre-existing

population structure signals and likely ivermectin-induced genetic drift in

treated populations. More careful marker selection based on better quality

whole genome sequence data from individual- rather than pooled-worms may

solve this problem and permit sensitive detection of SOR genotypes in

samples drawn from a single parasite population or transmission zone.

The second crucial question is whether it is possible to predict the

likelihood that SOR frequency will increase to a level that prevents elimination

of *O. volvulus* transmission in a given population. Our data suggest that

ivermectin reduces the population size of GR worms, as one would expect,

but that the population of SOR worms is stable and/or increasing as the GR

population shrinks. In very simple terms it is, therefore, a race: will the rate of

population contraction (driven by the temporary interruption and long term

reduction of transmission of GR worms) outpace the rate at which the SOR

population stabilises and expands (because they continue to be transmitted).

If GR decline is faster than SOR expansion and the population threshold for

maintenance of transmission is reached, local elimination will occur. If

expansion of the SOR population prevents that threshold from being reached,

transmission will persist, prevalence of infection will rebound and elimination

will not occur. The outcome of the race will be determined by: (i) the starting

frequency of SOR in a naïve population when ivermectin treatment begins, (ii)

the relative rates of GR population contraction and SOR population

expansion, and (iii) the population size at which transmission is no longer

sustainable (the threshold for maintenance of transmission). Developing

genotyping assays that can measure the pre-treatment (or current) SOR

frequency and monitor the relative rates of GR contraction and SOR

stabilization/expansion (the two critical parameters identified above) in

populations undergoing CDTI is therefore essential to detect populations in

which elimination may prove problematic. Furthermore, the response

phenotypes in terms of dynamics of microfilariae production [36] and thus

availability of skin microfilariae for vectors to ingest need to be better

understood. They could then be taken into account for timing CDTI relative to

vectors abundance so that the ratio of progeny of SOR/progeny of GR is

minimal when the availability of vectors is maximal.

The third crucial question is whether SOR is likely to spread from one

location to another. The analysis of population structure suggests that all of

the communities sampled in Ghana are in a single transmission zone (there is

no internal population structure between the Ghanaian communities), so that

SOR could be transmitted between communities in this region of Ghana. This

is also likely true for Cameroon. The tentative conclusion from analysis of the

Sequenom data for MBM94, MBM07 and NKA07 from Cameroon suggests

that immigration from an area not under CDTI (Nkam Valley) into a population

under CDTI may occur (suggested by the similarity between MBM07 and

NKA07), even if the pre-treatment level of transmission between the two

populations is low (as is suggested by comparison of NKA07 and MBM94,

which are genetically distinct populations). While the concept of

recrudescence in areas undergoing control as a result of immigration from

neighbouring regions with less or no control is not new, these data provide the

first genetic evidence that such immigration has likely occurred and may have

had an impact on the success of ivermectin distribution in the Mbam valley

parasite populations. In contrast, Ghana and Cameroon are clearly separated

genetically, which implies there is no transmission of *O. volvulus* at this scale

and SOR will not spread from Ghana to Cameroon (or *vice versa*). While this

is not surprising given the location of the two countries, the data show that

genetic markers were able to detect transmission between endemic areas

within Cameroon (i.e. are transmission zone markers) and thus could be a

useful tool for onchocerciasis control programs. Such a tool could support

decisions on whether to stop CDTI when criteria for stopping CDTI have been

met in one area but not in others in a geographic context that makes

transmission between these two areas possible. These data also suggest that

not only can genetic markers sensitively detect transmission between

geographic locations, but also that QLTs that are associated with SOR could

increase in frequency in neighbouring regions having little drug selection

pressure as a result of that transmission.

With respect to SORs, genotyping assays will benefit control programs

in multiple ways by (i) providing diagnostic tools to monitor changes in the

frequency of SORs (e.g. genotyping infective larvae in the vector), (ii)

discriminating between genetic explanations for persistence of transmission

(i.e., selection for SOR) and other factors that determine CDTI success (such

as host-related factors, treatment coverage and compliance, pre-control

prevalence and intensity of infection and vector biting rates [48-53, 99]), and

(iii) suggesting a trigger for the initiation of alternative treatment strategies,

such as anti-Wolbachia treatment [101, 102], local vector control, or new

treatments that may become available sooner or later like moxidectin [46,

103], or combination treatments, flubendazole or emodepside (as reviewed in

[104]) in populations where persistent transmission is observed in spite of

CDTI [105].

1016

1017

***Summary***

The data presented suggest that the evolution of SOR to ivermectin in

*O. volvulus* is via soft selective sweeps of pre-existing quantitative trait loci

(QTLs) rather than via a hard selective sweep of a relatively rare resistance-

conferring mutation. The outcome of this soft selection is the accumulation of

many alleles in a limited number of functional pathways that facilitate the

recovery of adult female worm fecundity from the inhibitory effects of

ivermectin. This is consistent with the observation that the acute

microfilaricidal and macrofilarial anti-fecundity effects of ivermectin remain

unaltered in SOR populations, and that the difference between SOR and GR

parasites is quantitative variation in the rate and extent to which fertility

recovers after ivermectin treatment. This conclusion (of a soft selective sweep

of quantitative variation in the rate of recovery from ivermectin toxicity) is

based on the presence of multiple, geographically independent genetic

signals throughout the genome that differentiate GR and SOR pools and the

apparent preservation of genetic diversity within the SOR populations

following selection. Furthermore, when these data are considered together

with the population bottlenecks that characterise the transmission of

*O. volvulus* through its lifecycle and the likelihood of some degree of

inbreeding, we conclude that *O. volvulus* populations are variable and can be

structured due to allele frequency change in the absence of selection, i.e., as

a consequence of genetic drift. What is unclear is the extent to which each

putative QTL contributes to the development of SOR. The failure to detect

differences between SOR and GR worms using a panel of 130 SNPs does

suggest that the signature of selection in these populations is subtle and that

no single putative QTL dominates the response; further validation of the

putative QTLs is clearly required. A more thorough association analysis of

each genomic region with SOR may be achieved by additional whole genome

sequencing using single adult worms (as opposed to pool sequencing as

described here) together with a more precise estimation of the ivermectin

response of the individual worm. This would allow fine mapping of QTLs, and

estimation of their effect size and penetrance, and ultimately, of the heritability

of SOR. Such analyses should take transmission zones and population

stratification into account to correctly determine the extent of gene flow and

therefore, the ability of SOR alleles to be transmitted within and between

populations. Additional sequencing of individual parasites may also provide

greater confidence in assigning genetic associations to individual SNPs,

opening the way to investigation of the likely function of individual SNPs as

putative SOR-conferring variants. Given the increasing accessibility to

genomic resources, the reduced cost of next-generation sequencing, and the

ability to look at the whole genome in an unbiased way, we propose that

genome-wide analyses such as those applied here should replace candidate

gene approaches for future work concerned with the genetics and diagnosis

of drug resistance in helminth parasites.

1062

**ABBREVIATIONS**

**CDTI,** community-directed treatment with ivermectin; **GR,** good responders; **HWE,** Hardy-Weinberg equilibrium; **KS**, Kolmogorov-Smirnov; **MDS**, multidimensional scaling; **NLT**, naïve or little treated; **QT,** quantitative trait; **QTL,** quantitative trait loci, **SD,** standard deviation; **SNP,** single nucleotide polymorphism; **SOR,** sub-optimal responders.

**ACKNOWLEDGEMENTS**

We would like to thank all of the patients and collaborators from Cameroon and Ghana who participated in the studies, as well as the Ministry of Public Health from Cameroon and the Ministry of Health from Ghana. We would like to acknowledge Pierre Lepage, Rosalie Frechette, Audrey Ann Kustec, Patrice Charbonneau-Larose and Valérie Catudal from the Genome Quebec Innovation Centre, Andrew Robinson and Nathan Hall from La Trobe University for computational support, James Cotton and Shannon Hedtke for critical reading of the manuscript, and Matthew Berriman and James Cotton from the Wellcome Trust Sanger Institute for pre-publication access to *O. volvulus* genomic resources.

**FUNDING STATEMENT**

This investigation received financial support from UNICEF/UNDP/World Bank/World Health Organization Special Programme for Research and Training in Tropical Diseases (WHO/TDR); Agence Nationale pour la Recherche, France (Grant: ANR-06-SEST-32); the Wellcome Trust, UK (Grant: 092677/Z/10/Z); the Natural Sciences and Engineering Research Council of Canada (Grant: RGPIN/2777-2012), Fonds de Recherche du Québec – Nature et technologies (FQRNT), Canadian Institutes of Health Research (CIHR grants: 85021 and 94593); Centre of Host-Parasite interaction (CHPI), Quebec; an Early Career Development Fellowship from La Trobe University (SRD); and computational resources from the Victorian Life Sciences Computation Initiative.

1097

**AUTHOR CONTRIBUTIONS**

Conceived and designed the experiments: WNG, RKP, MB, CB

Conducted the fieldwork, collected and processed parasite material for

genetic analysis: MB, MYOA, SW, SDSP, CB, HCND, JAKO, JB, HC

Contributed reagents and materials: MYOA, SW, MB, SDSP, RKP

Led the parasitology and molecular biology: CB

Led the bioinformatics and population genetics analyses: SRD

Provided intellectual input, participated in meetings and discussions on the

interpretation and presentation of the results and contributed to the writing of

the manuscript: SRD, CB, HCND, JAKO, SDSP, JK, SW, ACK, MW, MGB,

DAB, MYOA, MB, RKP, WNG

Drafted the paper: SRD, WNG

Read and approved the final submitted manuscript: SRD, CB, HND, JAKO,

SDSP, JB, JK, SW, HC, ACK, MW, MGB, DAB, MYOA, MB, RKP, WNG

## REFERENCES

1.      Remme JHF, Boatin B, Boussinesq M. Helminthic Diseases: Onchocerciasis and Loiasis. In: SR Q, WC C, editors. The International Encyclopedia of Public Health. Vol. 3. 2nd ed. Oxford: Elsevier; 2016. p. 576-87.

2.      Gardon J, Boussinesq M, Kamgno J, Gardon-Wendel N, Demanga N, Duke BO. Effects of standard and high doses of ivermectin on adult worms of *Onchocerca volvulus*: a randomised controlled trial. Lancet. 2002;360(9328):203-10.

3.      Duke BO, Zea-Flores G, Castro J, Cupp EW, Munoz B. Effects of multiple monthly doses of ivermectin on adult *Onchocerca volvulus*. Am J Trop Med Hyg. 1990;43(6):657-64.

4.      Klager SL, Whitworth JA, Downham MD. Viability and fertility of adult *Onchocerca volvulus* after 6 years of treatment with ivermectin. Trop Med Int Health. 1996;1(5):581-9.

5.      Basáñez MG, Pion SDS, Boakes E, Filipe JA, Churcher TS, Boussinesq M. Effect of single-dose ivermectin on *Onchocerca volvulus*: a systematic review and meta-analysis. Lancet Infect Dis. 2008;8(5):310-22.

6.      Rodriguez-Perez MA, Unnasch TR, Dominguez-Vazquez A, Morales-Castro AL, Pena-Flores GP, Orozco-Algarra ME, et al. Interruption of transmission of *Onchocerca volvulus* in the Oaxaca focus, Mexico. Am J Trop Med Hyg. 2010;83(1):21-7.

7.      Rodriguez-Perez MA, Dominguez-Vazquez A, Unnasch TR, Hassan HK, Arredondo-Jimenez JI, Orozco-Algarra ME, et al. Interruption of transmission of *Onchocerca volvulus* in the Southern Chiapas Focus, Mexico. PLoS Negl Trop Dis. 2013;7(3):e2133.

8.      Gonzalez RJ, Cruz-Ortiz N, Rizzo N, Richards J, Zea-Flores G, Dominguez A, et al. Successful interruption of transmission of *Onchocerca volvulus* in the Escuintla-Guatemala focus, Guatemala. PLoS Negl Trop Dis. 2009;3(3):e404.

9.      Lovato R, Guevara A, Guderian R, Proano R, Unnasch T, Criollo H, et al. Interruption of infection transmission in the onchocerciasis focus of Ecuador leading to the cessation of ivermectin distribution. PLoS Negl Trop Dis. 2014;8(5):e2821.

10.     Convit J, Schuler H, Borges R, Olivero V, Dominguez-Vazquez A, Frontado H, et al. Interruption of *Onchocerca volvulus* transmission in Northern Venezuela. Parasit Vectors. 2013;6(1):289.

11.     Richards F, Jr., Rizzo N, Diaz Espinoza CE, Monroy ZM, Crovella Valdez CG, de Cabrera RM, et al. One Hundred Years After Its Discovery in Guatemala by Rodolfo Robles, *Onchocerca volvulus* Transmission Has Been Eliminated from the Central Endemic Zone. Am J Trop Med Hyg. 2015;93(6):1295-304.

12.     Botto C, Basáñez MG, Escalona M, Villamizar NJ, Noya-Alarcón O, Cortez J, et al. Evidence of suppression of onchocerciasis transmission in the Venezuelan Amazonian focus. Parasit Vectors. 2016;9(1):40.

13.     Higazi TB, Zarroug IM, Mohamed HA, Elmubark WA, Deran TC, Aziz N, et al. Interruption of *Onchocerca volvulus* transmission in the Abu Hamed focus, Sudan. Am J Trop Med Hyg. 2013;89(1):51-7.

14.     Katabarwa MN, Walsh F, Habomugisha P, Lakwo TL, Agunyo S, Oguttu DW,
et al. Transmission of onchocerciasis in wadelai focus of northwestern Uganda has
been interrupted and the disease eliminated. J Parasitol Res. 2012;2012:748540.

15.     Lakwo TL, Garms R, Rubaale T, Katabarwa M, Walsh F, Habomugisha P, et
al. The disappearance of onchocerciasis from the Itwara focus, western Uganda after
elimination of the vector *Simulium neavei* and 19 years of annual ivermectin
treatments. Acta Trop. 2013;126(3):218-21.

16.     Tekle AH, Elhassan E, Isiyaku S, Amazigo UV, Bush S, Noma M, et al.
Impact of long-term treatment of onchocerciasis with ivermectin in Kaduna State,
Nigeria: first evidence of the potential for elimination in the operational area of the
African Programme for Onchocerciasis Control. Parasit Vectors. 2012;5:28.

17.     Tekle AH, Zoure HG, Noma M, Boussinesq M, Coffeng LE, Stolk WA, et al.
Progress towards onchocerciasis elimination in the participating countries of the
African Programme for Onchocerciasis Control: epidemiological evaluation results.
Infect Dis Poverty. 2016;5(1):66.

18.     Traore MO, Sarr MD, Badji A, Bissan Y, Diawara L, Doumbia K, et al. Proof-
of-principle of onchocerciasis elimination with ivermectin treatment in endemic foci in
Africa: final results of a study in Mali and Senegal. PLoS Negl Trop Dis.
2012;6(9):e1825.

19.     Zarroug IM, Hashim K, ElMubark WA, Shumo ZA, Salih KA, ElNojomi NA, et
al. The First Confirmed Elimination of an Onchocerciasis Focus in Africa: Abu
Hamed, Sudan. Am J Trop Med Hyg. 2016.

20.     Accelerating work to overcome the global impact of neglected tropical
diseases. A roadmap for implementation. Geneva: World Health Organization; 2012.

21.     Eighteenth Session of the Joint Action Forum (JAF) of the African Programme
for Onchocerciasis Control, Bujumbura, Burundi, 11-13 December 2012. Final
Communiqué. Africa Programme for Onchocerciasis Control (APOC); 2012.

22.     Borsboom GJ, Boatin BA, Nagelkerke NJ, Agoua H, Akpoboua KL, Alley EW,
et al. Impact of ivermectin on onchocerciasis transmission: assessing the empirical
evidence that repeated ivermectin mass treatments may lead to
elimination/eradication in West-Africa. Filaria J. 2003;2(1):8.

23.     Remme J, Baker RH, De Sole G, Dadzie KY, Walsh JF, Adams MA, et al. A
community trial of ivermectin in the onchocerciasis focus of Asubende, Ghana. I.
Effect on the microfilarial reservoir and the transmission of *Onchocerca volvulus*.
Trop Med Parasitol. 1989;40(3):367-74.

24.    Lakwo TL. Preliminary studies on the biting activity and transmission of
*Onchocerca volvulus* by *Simulium neavei* (diptera: simuliidae) in Kashoya-Kitomi
focus, Western Uganda. East Afr Med J. 2004;81(5):244-7.

25.     Katabarwa M, Eyamba A, Habomugisha P, Lakwo T, Ekobo S, Kamgno J, et
al. After a decade of annual dose mass ivermectin treatment in Cameroon and
Uganda, onchocerciasis transmission continues. Trop Med Int Health.
2008;13(9):1196-203.

26.      Katabarwa MN, Eyamba A, Nwane P, Enyong P, Yaya S, Baldiagai J, et al.
Seventeen years of annual distribution of ivermectin has not interrupted
onchocerciasis transmission in North Region, Cameroon. Am J Trop Med Hyg.
2011;85(6):1041-9.

27.      Frempong KK, Walker M, Cheke RA, Tetevi EJ, Gyan ET, Owusu EO, et al.
Does Increasing Treatment Frequency Address Suboptimal Responses to Ivermectin
for the Control and Elimination of River Blindness? Clin Infect Dis. 2016;62(11):1338-
47.

28.      Eisenbarth A, Achukwi MD, Renz A. Ongoing Transmission of *Onchocerca*
*volvulus* after 25 Years of Annual Ivermectin Mass Treatments in the Vina du Nord
River Valley, in North Cameroon. PLoS Negl Trop Dis. 2016;10(2):e0004392.

29.      Awadzi K, Attah SK, Addy ET, Opoku NO, Quartey BT, Lazdins-Helds JK, et
al. Thirty-month follow-up of sub-optimal responders to multiple treatments with
ivermectin, in two onchocerciasis-endemic foci in Ghana. Ann Trop Med Parasitol.
2004;98(4):359-70.

30.      Awadzi K, Boakye DA, Edwards G, Opoku NO, Attah SK, Osei-Atweneboana
MY, et al. An investigation of persistent microfilaridermias despite multiple treatments
with ivermectin, in two onchocerciasis-endemic foci in Ghana. Ann Trop Med
Parasitol. 2004;98(3):231-49.

31.      Lamberton PHL, Cheke RA, Winskill P, Tirados I, Walker M, Osei-
Atweneboana MY, et al. Onchocerciasis Transmission in Ghana: Persistence under
Different Control Strategies and the Role of the Simuliid Vectors. PLoS Negl Trop
Dis. 2015;9(4):e0003688.

32.      Kamga G-R, Dissak-Delon FN, Nana-Djeunga HC, Biholong BD, Mbigha-
Ghogomu S, Souopgui J, et al. Still mesoendemic onchocerciasis in two
Cameroonian community-directed treatment with ivermectin projects despite more
than 15 years of mass treatment. Parasit Vectors. 2016;9(1):581.

33.      Koala L, Nikiema A, Post RJ, Paré AB, Kafando CM, Drabo F, et al.
Recrudescence of onchocerciasis in the Comoé valley in Southwest Burkina Faso.
Acta Tropica. 2017;166:96-105.

34.      Osei-Atweneboana MY, Awadzi K, Attah SK, Boakye DA, Gyapong JO,
Prichard RK. Phenotypic evidence of emerging ivermectin resistance in *Onchocerca*
*volvulus.* PLoS Negl Trop Dis. 2011;5(3):e998.

35.      Osei-Atweneboana MY, Eng JK, Boakye DA, Gyapong JO, Prichard RK.
Prevalence and intensity of *Onchocerca volvulus* infection and efficacy of ivermectin
in endemic communities in Ghana: a two-phase epidemiological study. Lancet.
2007;369(9578):2021-9.

36.      Pion SDS, Nana-Djeunga HC, Kamgno J, Tendongfor N, Wanji S, Njiokou F,
et al. Dynamics of *Onchocerca volvulus* microfilarial densities after ivermectin
treatment in an ivermectin-naive and a multiply treated population from Cameroon.
PLoS Negl Trop Dis. 2013;7(2):e2084.

37.      Nana-Djeunga HC, Bourguinat C, Pion SDS, Bopda J, Kengne-Ouafo JA,
Njiokou F, et al. Reproductive Status of *Onchocerca volvulus* after Ivermectin

Treatment in an Ivermectin-Naive and a Frequently Treated Population from
Cameroon. PLoS Negl Trop Dis. 2014;8(4):e2824.

38.     Osei-Atweneboana MY, Boakye DA, Awadzi K, Gyapong JO, Prichard RK.
Genotypic analysis of beta-tubulin in *Onchocerca volvulus* from communities and
individuals showing poor parasitological response to ivermectin treatment. Int J
Parasitol Drugs Drug Resist. 2012;2.

39.     Ardelli BF, Guerriero SB, Prichard RK. Genomic organization and effects of
ivermectin selection on *Onchocerca volvulus* P-glycoprotein. Mol Biochem Parasitol.
2005;143(1):58-66.

40.     Ardelli BF, Guerriero SB, Prichard RK. Ivermectin imposes selection pressure
on P-glycoprotein from *Onchocerca volvulus*: linkage disequilibrium and genotype
diversity. Parasitology. 2006;132(Pt 3):375-86.

41.     Ardelli BF, Guerriero SB, Prichard RK. Characterization of a half-size ATP-
binding cassette transporter gene which may be a useful marker for ivermectin
selection in *Onchocerca volvulus*. Mol Biochem Parasitol. 2006;145(1):94-100.

42.     Ardelli BF, Prichard RK. Identification of variant ABC-transporter genes
among *Onchocerca volvulus* collected from ivermectin-treated and untreated patients
in Ghana, West Africa. Ann Trop Med Parasitol. 2004;98(4):371-84.

43.     Ardelli BF, Prichard RK. Reduced genetic variation of an *Onchocerca*
*volvulus* ABC transporter gene following treatment with ivermectin. Trans R Soc Trop
Med Hyg. 2007;101(12):1223-32.

44.     Bourguinat C, Ardelli BF, Pion SD, Kamgno J, Gardon J, Duke BO. P-
glycoprotein-like protein, a possible genetic marker for ivermectin resistance
selection in *Onchocerca volvulus*. Mol Biochem Parasitol. 2008;158.

45.     Bottomley C, Isham V, Collins RC, Basáñez MG. Rates of microfilarial
production by *Onchocerca volvulus* are not cumulatively reduced by multiple
ivermectin treatments. Parasitology. 2008;135(13):1571-81.

46.     Awadzi K, Opoku NO, Attah SK, Lazdins-Helds J, Kuesel AC. A randomized,
single-ascending-dose, ivermectin-controlled, double-blind study of moxidectin in
*Onchocerca volvulus* infection. PLoS Negl Trop Dis. 2014;8(6):e2953.

47.     Bakajika DK, Kanza E, Howard H, Opoku N, Tchatchu J-PL, Kataliko K, et al.,
editors. Is *O. volvulus* suboptimal response to ivermectin a result of selection under
ivermectin pressure? Insights from a study comparing ivermectin and moxidectin in
areas without prior ivermectin mass treatment. Am J Trop Med Hyg; 2013;
Washington, DC.

48.     Pion SDS, Grout L, Kamgno J, Nana-Djeunga H, Boussinesq M. Individual
host factors associated with *Onchocerca volvulus* microfilarial densities 15, 80 and
180 days after a first dose of ivermectin. Acta Trop. 2011;120 Suppl 1:S91-9.

49.     Burnham G. Efficacy of ivermectin against *Onchocerca volvulus* in Ghana.
Lancet. 2007;370(9593):1125.

50.     Cupp E, Richards F, Lammie P, Eberhard M. Efficacy of ivermectin against
*Onchocerca volvulus* in Ghana. Lancet. 2007;370(9593):1123; author reply 4-5.

51.    Mackenzie CD. Efficacy of ivermectin against *Onchocerca volvulus* in Ghana.
Lancet. 2007;370(9593):1123; author reply 4-5.

52.    Remme JH, Amazigo U, Engels D, Barryson A, Yameogo L. Efficacy of
ivermectin against *Onchocerca volvulus* in Ghana. Lancet. 2007;370(9593):1123-4;
author reply 4-5.

53.    Hotez PJ. Control of onchocerciasis--the next generation. Lancet.
2007;369(9578):1979-80.

54.    Churcher TS, Pion SDS, Osei-Atweneboana MY, Prichard RK, Awadzi K,
Boussinesq M, et al. Identifying sub-optimal responses to ivermectin in the treatment
of River Blindness. Proc Natl Acad Sci U S A. 2009;106(39):16716-21.

55.    Turner HC, Churcher TS, Walker M, Osei-Atweneboana MY, Prichard RK,
Basáñez MG. Uncertainty surrounding projections of the long-term impact of
ivermectin treatment on human onchocerciasis. PLoS Negl Trop Dis.
2013;7(4):e2169.

56.    Park DJ, Lukens AK, Neafsey DE, Schaffner SF, Chang HH, Valim C, et al.
Sequence-based association and selection scans identify drug resistance loci in the
*Plasmodium falciparum* malaria parasite. Proc Natl Acad Sci U S A.
2012;109(32):13052-7.

57.    Cheeseman IH, Miller BA, Nair S, Nkhoma S, Tan A, Tan JC, et al. A major
genome region underlying artemisinin resistance in malaria. Science.
2012;336(6077):79-82.

58.    Miotto O, Almagro-Garcia J, Manske M, Macinnis B, Campino S, Rockett KA,
et al. Multiple populations of artemisinin-resistant *Plasmodium falciparum* in
Cambodia. Nat Genet. 2013;45(6):648-55.

59.    Cheeseman IH, McDew-White M, Phyo AP, Sriprawat K, Nosten F, Anderson
TJ. Pooled sequencing and rare variant association tests for identifying the
determinants of emerging drug resistance in malaria parasites. Mol Biol Evol.
2015;32(4):1080-90.

60.    Chang H-H, Moss EL, Park DJ, Ndiaye D, Mboup S, Volkman SK, et al.
Malaria life cycle intensifies both natural selection and random genetic drift.
Proceedings of the National Academy of Sciences. 2013;110(50):20129-34.

61.    Valentim CL, Cioli D, Chevalier FD, Cao X, Taylor AB, Holloway SP, et al.
Genetic and molecular basis of drug resistance and species-specific drug action in
schistosome parasites. Science. 2013;342(6164):1385-9.

62.    Downing T, Imamura H, Decuypere S, Clark TG, Coombs GH, Cotton JA, et
al. Whole genome sequencing of multiple *Leishmania donovani* clinical isolates
provides insights into population structure and mechanisms of drug resistance.
Genome Res. 2011;21(12):2143-56.

63.    Holt KE, Parkhill J, Mazzoni CJ, Roumagnac P, Weill FX, Goodhead I, et al.
High-throughput sequencing provides insights into genome variation and evolution in
*Salmonella Typhi*. Nat Genet. 2008;40(8):987-93.

64.     Gilleard JS. Understanding anthelmintic resistance: The need for genomics
and genetics. Int J Parasitol. 2006;36(12):1227-39.

65.     Kotze AC, Hunt PW, Skuce P, von Samson-Himmelstjerna G, Martin RJ,
Sager H, et al. Recent advances in candidate-gene and whole-genome approaches
to the discovery of anthelmintic resistance markers and the description of
drug/receptor interactions. Int J Parasitol Drugs Drug Resist. 2014;4(3):164-84.

66.     Choi Y-J, Tyagi R, McNulty SN, Rosa BA, Ozersky P, Martin J, et al. Genomic
diversity in *Onchocerca volvulus* and its Wolbachia endosymbiont. Nat Microbiol.
2016;2:16207.

67.     Meredith SEO, Lando G, Gbakima AA, Zimmerman PA, Unnasch TR.
*Onchocerca volvulus* - Application of the Polymerase Chain-Reaction to Identification
and Strain Differentiation of the Parasite. Exp Parasitol. 1991;73(3):335-44.

68.     Zimmerman PA, Dadzie KY, Desole G, Remme J, Alley ES, Unnasch TR.
*Onchocerca volvulus* DNA Probe Classification Correlates with Epidemiologic
Patterns of Blindness. J Infect Dis. 1992;165(5):964-8.

69.     Cotton JA, Bennuru S, Grote A, Harsha B, Tracey A, Beech R, et al. The
genome of *Onchocerca volvulus*, agent of river blindness. Nat Microbiol.
2016;2:16216.

70.     Bennuru S, Cotton JA, Ribeiro JM, Grote A, Harsha B, Holroyd N, et al.
Stage-Specific Transcriptome and Proteome Analyses of the Filarial Parasite
*Onchocerca volvulus* and Its Wolbachia Endosymbiont. MBio. 2016;7(6).

71.     Howe KL, Bolt BJ, Shafie M, Kersey P, Berriman M. WormBase ParaSite – a
comprehensive resource for helminth genomics. Mol Biochem Parasitol. 2016;In
press: Available online 27 November 2016.

72.     Li H. Aligning sequence reads, clone sequences and assembly contigs with
BWA-MEM. arXiv preprint arXiv:13033997. 2013.

73.     McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, et
al. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-
generation DNA sequencing data. Genome Res. 2010;20(9):1297-303.

74.     Bansal V. A statistical method for the detection of variants from next-
generation resequencing of DNA pools. Bioinformatics. 2010;26(12):i318-24.

75.     Kofler R, Pandey RV, Schlötterer C. PoPoolation2: identifying differentiation
between populations using sequencing of pooled DNA samples (Pool-Seq).
Bioinformatics. 2011;27(24):3435-6.

76.     Storm N, Darnhofer-Patel B, van den Boom D, Rodi CP. MALDI-TOF mass
spectrometry-based SNP genotyping. Methods Mol Biol. 2003;212:241-62.

77.     Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, et al.
PLINK: a tool set for whole-genome association and population-based linkage
analyses. Am J Hum Genet. 2007;81(3):559-75.

78.     Jombart T. adegenet: a R package for the multivariate analysis of genetic
markers. Bioinformatics. 2008;24(11):1403-5.

79.  Rellstab C, Zoller S, Tedder A, Gugerli F, Fischer MC. Validation of SNP
allele frequencies determined by pooled next-generation sequencing in natural
populations of a non-model plant species. PLoS One. 2013;8(11):e80422.

80.  Kim SY, Li Y, Guo Y, Li R, Holmkvist J, Hansen T, et al. Design of association
studies with pooled or un-pooled next-generation sequencing data. Genet Epidemiol.
2010;34(5):479-91.

81.  Futschik A, Schlötterer C. The next generation of molecular markers from
massively parallel sequencing of pooled DNA samples. Genetics. 2010;186(1):207-
18.

82.  Schlötterer C, Tobler R, Kofler R, Nolte V. Sequencing pools of individuals -
mining genome-wide polymorphism data without big funding. Nat Rev Genet.
2014;15(11):749-63.

83.  Anderson EC, Skaug HJ, Barshis DJ. Next-generation sequencing for
molecular ecology: a caveat regarding pooled samples. Mol Ecol. 2014;23(3):502-12.

84.  Traore-Lamizana M, Lemasson J. Participation à une étude de faisabilité
d'une campagne de lutte contre l'onchocercose dans la région du bassin du Logone:
répartition des espèces du complexe *Simulium damnosum* dans la zone
camerounaise du projet. Cahiers-ORSTOM Entomologie médicale et parasitologie.
1988;25(3-4):171-86.

85.  Basáñez MG, Churcher TS, Grillet ME. *Onchocerca-Simulium* interactions
and the population and evolutionary biology of *Onchocerca volvulus*. Adv Parasitol.
2009;68.

86.  Cully DF, Vassilatis DK, Liu KK, Paress PS, Vanderploeg LHT, Schaeffer JM,
et al. Cloning of an Avermectin-Sensitive Glutamate-Gated Chloride Channel from
*Caenorhabditis elegans*. Nature. 1994;371(6499):707-11.

87.  Bany IA, Dong MQ, Koelle MR. Genetic and cellular basis for acetylcholine
inhibition of *Caenorhabditis elegans* egg-laying behavior. J Neurosci.
2003;23(22):8060-9.

88.  Ringstad N, Horvitz HR. FMRFamide neuropeptides and acetylcholine
synergistically inhibit egg-laying by *C. elegans*. Nat Neurosci. 2008;11(10):1168-76.

89.  Waggoner LE, Zhou GT, Schafer RW, Schafer WR. Control of alternative
behavioral states by serotonin in *Caenorhabditis elegans*. Neuron. 1998;21(1):203-
14.

90.  Li B-W, Rush AC, Weil GJ. Expression of five acetylcholine receptor subunit
genes in *Brugia malayi* adult worms. Int J Parasitol Drugs Drug Resist.
2015;5(3):100-9.

91.  Hernando G, Bouzat C. *Caenorhabditis elegans* neuromuscular junction:
GABA receptors and ivermectin action. PLoS One. 2014;9(4):e95072.

92.  Abongwa M, Buxton SK, Robertson AP, Martin RJ. Curiouser and Curiouser:
The Macrocyclic Lactone, Abamectin, Is also a Potent Inhibitor of
Pyrantel/Tribendimidine Nicotinic Acetylcholine Receptors of Gastro-Intestinal
Worms. PLoS One. 2016;11(1).

93.      Bourguinat C, Pion SDS, Kamgno J, Gardon J, Duke BOL, Boussinesq M, et
al. Genetic Selection of Low Fertile *Onchocerca volvulus* by Ivermectin Treatment.
PLoS Negl Trop Dis. 2007;1(1).

94.      Laing ST, Ivens A, Butler V, Ravikumar SP, Laing R, Woods DJ, et al. The
transcriptional response of *Caenorhabditis elegans* to Ivermectin exposure identifies
novel genes involved in the response to reduced food intake. PLoS One.
2012;7(2):e31367.

95.      Bottomley C, Isham V, Vivas-Martínez S, Kuesel AC, Attah SK, Opoku NO, et
al. Modelling Neglected Tropical Diseases diagnostics: the sensitivity of skin snips for
*Onchocerca volvulus* in near elimination and surveillance settings. Parasit Vectors.
2016;9(1):343.

96.      Hunt PW. Identification and characterisation of dominant ivermectin resistant
genes in *Caenorhabditis elegans* [PhD thesis]: University of New England; 1995.

97.      Barnes TM, Hekimi S. The *Caenorhabditis elegans* avermectin resistance
and anesthetic response gene unc-9 encodes a member of a protein family
implicated in electrical coupling of excitable cells. J Neurochem. 1997;69(6):2251-60.

98.      Turner HC, Walker M, Churcher TS, Osei-Atweneboana MY, Biritwum NK,
Hopkins A, et al. Reaching the London Declaration on Neglected Tropical Diseases
Goals for Onchocerciasis: An Economic Evaluation of Increasing the Frequency of
Ivermectin Treatment in Africa. Clin Infect Dis. 2014;59(7):923-32.

99.      Stolk WA, Walker M, Coffeng LE, Basáñez MG, de Vlas SJ. Required
duration of mass ivermectin treatment for onchocerciasis elimination in Africa: a
comparative modelling analysis. Parasit Vectors. 2015;8(1):552.

100.    Coffeng LE, Stolk WA, Hoerauf A, Habbema D, Bakker R, Hopkins AD, et al.
Elimination of African Onchocerciasis: Modeling the Impact of Increasing the
Frequency of Ivermectin Mass Treatment. PLoS One. 2015;9(12):e115886.

101.    Taylor MJ, Hoerauf A, Townson S, Slatko BE, Ward SA. Anti-*Wolbachia* drug
discovery and development: safe macrofilaricides for onchocerciasis and lymphatic
filariasis. Parasitology. 2014;141(1):119-27.

102.    Walker M, Specht S, Churcher TS, Hoerauf A, Taylor MJ, Basáñez MG.
Therapeutic Efficacy and Macrofilaricidal Activity of Doxycycline for the Treatment of
River Blindness. Clinical Infectious Diseases. 2015;60(8):1199-207.

103.    Turner HC, Walker M, Attah SK, Opoku NO, Awadzi K, Kuesel AC, et al. The
potential impact of moxidectin on onchocerciasis elimination in Africa: an economic
evaluation based on the Phase II clinical trial data. Parasit Vectors. 2015;8(1):167.

104.    Kuesel AC. Research for new drugs for elimination of onchocerciasis in
Africa. Int J Parasitol Drugs Drug Resist. 2016;In Press: Available online 19 May
2016.

105.    African Programme for Onchocerciasis Control (APOC): Strategic Options
and Alternative Treatment Strategies for Accelerating Onchocerciasis Elimination in
Africa. Ouagadougou: World Health Organization 2015.

1447

1448

**Figure Legends**

**Figure 1.** Analysis of shared variation that differentiates ivermectin good

responder (GR) from sub-optimally responding (SOR) *Onchocerca volvulus*

adult worms in both the Cameroon and Ghana population samples, measured

using individual single nucleotide polymorphisms (SNPs) (Fisher's exact test;

**A**) or 10-kb windows ($F_{ST}$; **B**). Dotted lines represent statistical cutoff applying

the Bonferroni correction for SNPs and genome-wide mean $F_{ST}$ + 3 standard

deviations (SDs) (Cam: 0.355; Gha: 0.255) for 10-kb windows. Red dots

highlight differentiation above genome-wide cutoffs that is shared by both

groups. Orange dots represent additional shared differentiation at 2 SDs in

the $F_{ST}$ analysis (**B**). Manhattan plots of genome-wide $F_{ST}$ describing spatial

genetic differentiation between GR and SOR pools for both Cameroon (**C**)

and Ghana (**D**). Each point represents $F_{ST}$ calculated for a non-overlapping

10-kb window. Plots are colored to differentiate the main genomic scaffolds

from unplaced scaffolds and contigs. Dotted lines represents genome-wide

mean $F_{ST}$ + 3 SDs (dark grey; Cam: 0.355; Gha: 0.255) and $F_{ST}$ + 5 SDs (light

grey; Cam: 0.508; Gha: 0.351).

1466

**Figure 2.** Analysis of genetic diversity between ivermectin responder

phenotypes and drug-"naïve" (NTL) worms. (**A**) Spearman rank correlation

analysis of variant read frequencies from 248,102 SNPs. Values within

squares represents correlation coefficients for each pairwise analysis. (**B**)

Degree of shared variation determined by pairwise comparisons of $F_{ST}$

between treatment and response groups, summarized from 9893 10-kb

windows throughout the genome. $F_{ST}$ distributions were compared using a

two-sample Kolmogorov-Smirnov [KS] test. (**C**) Variant read frequency

spectrum from treatment and response subgroups for Cameroon and Ghana.

The variant read frequency was calculated at each of the 248,102 SNP

positions, from which the proportion of total variants in 0.05 frequency bins is

presented. (**D**) Analysis of "invariant" loci per group as a proportion of the total

number of variants observed, defined as variant read frequencies < 0.05

(blue) and >0.95 (red).

1481

**Figure 3.** Analysis of genetic differentiation among 446 *O. volvulus* female

worms from Ghana and Cameroon individually genotyped at 130 loci

distributed throughout the genome. Multi-dimensional scaling analysis was

used to determine the relative genetic similarity between worms. Plots contain

the same data, but have been presented to emphasise the degree of genetic

differences between countries (**A**), sampling communities within each country

(**B**), and their treatment exposure and phenotypic response to ivermectin if

known (**C**). Ghanaian sampling sites: Asubende (ASU), Begbomdo (BEG),

Jagbenbendo (JAG), Kyingakrom (KYG), New Longoro (NLG), and Wiae

(WIA). Cameroonian sampling sites: Nkam (NKA07), Mbam valley sampled in

1994 (MBM94) before introduction of annual CDTI in 1994 and sampled in

2007 (MBM07).

**Table 1:** Summary of genes with shared functional characteristics / pathways from QTL peaks from Ghana and Cameroon [a,b]

| Country | Ivermectin-associated | Neuro-transmission [c] | LIN-12/Notch signaling | Muscle assembly / myosin organization | Lipid synthesis and storage / stress |
|---|---|---|---|---|---|
| Cameroon | | *Ovo-emc-6* (4; ACh) <br> *Ovo-cha-1* (9; ACh) <br> *Ovo-unc-17* (9; ACh) <br> *Ovo-stg-1* (10.1) <br> *Ovo-aex-3* (10.1; ACh) <br> *Ovo-nrfl-1* (16; ACh) <br> *Ovo-ncx-5* (24) <br> *Ovo-sca-1* (25) <br> *Ovo-lgc-46* (27; ACh) | *Ovo-crb-1* (4) | *Ovo-mup-4* (25) <br> *Ovo-mua-3* (25) <br> *Ovo-sca-1* (25) <br> *Ovo-mlc-5* (26) | *Ovo-acs-16* (3) <br> *Ovo-obr-2* (4) <br> *Ovo-math-46* (7) <br> Ovo-hsp-17 (22) <br> *Ovo-mtd-15* (25) <br> *Ovo-cuc-1* (25) |
| Ghana | *Ovo-inx-5*[b] (5) <br> *Ovo-klp-11*[b] (13) <br> *Ovo-unc-44* (14) | *Ovo-nud-1* (1) <br> *Ovo-inx-5* (5) <br> *Ovo-stg-1* (10.2) <br> *Ovo-aex-3* (10.2; ACh) <br> *Ovo-unc-31* (13) <br> *Ovo-lgc-47* (17; ACh) <br> *Ovo-kin-2* (17) <br> *Ovo-unc-26* (19) <br> *Ovo-acc-1* (21; ACh) <br> *Ovo-snb-1* (23; ACh) | *Ovo-bre-5* (5) <br> *Ovo-pen-2* (29) | *Ovo-unc-82* (21) <br> *Ovo-nmy-1* (21) | *Ovo-hsb-1* (1) <br> *Ovo-sms-1* (13) <br> *Ovo-spl-1* (17) <br> *Ovo-fat-4* (19) <br> *Ovo-tat-2* (19) |

[a] Quantitative trait loci (QTL) identification is provided in parentheses (refer to **Additional file 1: Table S4** for a description of each QTL and **Additional file 1:**

**Table S5** for the genes within).

[b] putative ivermectin association as described in text

[c] loci associated with acetylcholine neurotransmission are abbreviated with ACh.

**SUPPLEMENTARY DATA**

**Additional file 1 (xls).**

**Table S1.** Sequencing library composition, sequencing data and mapping
statistics**.**

**Table S2.** SNP calling statistics derived from CRISP, FreeBayes and
PoPoolation2 Pool-seq variant analysis, highlighting total SNPs called and the
intersection between different SNP callers.

**Table S3.** Analysis of shared regions of genetic differentiation between SOR
and GR of both Cameroon and Ghana. Table provides details of variants
presented in Figure 1 A and B.

**Table S4**. Characterisation of QTL clusters that differentiate SOR and GR in
either Cameroon or Ghana. Table provides details of regions of differentiation
presented in Figure 1 C&D.

**Table S5.** Characterisation of genes within QTL clusters that differentiate
SOR and GR in either Cameroon or Ghana.

**Table S6**. Analysis of genetic differentiation between GR and SOR pools in
ivermectin-resistance associated genes from the literature and in clusters.

**Table S7**. Names and coordinates of the 160 SNPs genotyped by Sequenom.

**Table S8**. HWE analysis of GR and SOR populations from Cameroon.

**Table S9**. HWE analysis of GR and SOR populations from Ghana

1519

**Additional file 2**: Supplementary information (pdf)

*Section 1. Extended information regarding sample history*

**Table S1.** Criteria for phenotypic classification of *Onchocerca volvulus*

**Table S2.** Overview of samples selected for Pool-seq

**Table S3.** Overview of samples selected for single worm genotyping

**Figure S1.** Maps of sampling sites in Ghana and Cameroon

**Table S4.** Sampling sites and mapping coordinates in Ghana and Cameroon.

*Section 2. Extended discussion of hard- and soft-selective sweeps*

**Figure S2.** Demonstrating the consequences of hard- versus soft-selection on
genetic diversity

**Table S5**. A simple multilocus quantitative trait model demonstrating multiple
genotypes conferring the same quantitative phenotype

**Table S6.** Summary of features differentiating hard and soft selection

***Section 3. Population assignment based on Sequenom genotyping of***

***individual worms***

**Table S7.** Population assignment of individual worms based on their genotype

profile from 130 SNPs analysed by Sequenom

**Figure S3.** DAPC analysis of genetic diversity to compare predicted versus

known population assignment of individuals based on their genotype.

**A**

**B**

**C**

**D**