Integrating proteomics and selected reaction monitoring to develop a non-invasive assay

for geoduck reproductive maturation

Emma B. Timmins-Schiffman[1], Grace A. Crandall[2], Brent Vadopalas[2], Michael E. Riffle[3], Brook

L. Nunn[1], Steven B. Roberts[2],*

[1] Department of Genome Sciences, University of Washington, Seattle, WA 98105

[2] Department of Aquatic and Fishery Sciences, University of Washington, Seattle, WA 98105

[3] Department of Biochemistry, University of Washington, Seattle, WA 98105

* Corresponding author:

Steven B. Roberts, Department of Aquatic and Fishery Sciences, University of Washington,

Seattle WA 98105. Email: sr320@uw.edu

Abstract

Geoduck clams (*Panopea generosa*) are an increasingly important fishery and

aquaculture product in the United States while also holding essential ecological roles. These

long-lived clams are highly fecund, but sustainable hatchery production of genetically diverse

larvae is hindered by the lack of sexual dimorphism, resulting in asynchronous spawning of

broodstock, unequal sex ratios, and low numbers of breeders. Assays of gonad physiology could

indicate both sex and maturation stage. Proteomic profiles were determined for three

reproductive maturation stages in both male and female clams using data dependent acquisition

(DDA), or whole proteome profiling, of gonad proteins. Gonad proteomes became increasingly

divergent between males and females as maturation progressed. The DDA data was used to

develop targets analyzed with selected reaction monitoring (SRM) in gonad tissue as well as

hemolymph. The SRM assay yielded a suite of indicator peptides that can be used as an efficient

assay to non-lethally determine geoduck sex and maturation stage pre-spawning. These results

lay the foundation for the development of field-based tools to assess reproductive status in

marine mollusks.

Key words: reproduction, gonad, geoduck, hemolymph, mollusk, proteomics, SRM,

targeted proteomics

Introduction

Wild Pacific geoduck clams (*Panopea generosa*), like other suspension feeders, provide

vital ecosystem services as both primary consumers of phytoplankton and biodepositors.[1]

Additionally, commercial geoduck fisheries provide significant social benefits as the most

economically important clam fishery in North America.[2] Commercial farming of geoduck now

generates over 20 million dollars in annual sales and geoduck are among the most valuable

farmed shellfish on a per acre basis.[3] The rapid development of geoduck aquaculture close to

wild populations of conspecifics has raised concerns over farmed- wild interbreeding and the

subsequent loss of genetic diversity. To address these concerns, geoduck hatcheries are currently

encouraged to use large numbers of wild broodstock to maximize genetic diversity in farmed

geoducks. An ongoing impediment to fulfilling this conservation goal is the absence of nonlethal

identification methods for sex and maturation stage determination in hatchery broodstock.

Geoducks are not sexually dimorphic, so these determinations require hatcheries to

increase the temperature of seawater and introduce large rations of microalgae to induce the

broodstock to spawn. Further, within a hatchery incubator only a few of the broodstock mature

and spawn during a given spawning attempt. This thwarts the sustainability goal to breed large

numbers of broodstock to maintain genetic diversity. Highly skewed sex ratios in some

populations and the inability to determine when to induce spawning based on maturation state

presents additional substantial economic challenges for shellfish hatcheries. Since sexual

differentiation and maturation processes in marine mollusks cannot be well characterized with

visual inspection, a low-cost assay for sex and maturation stage would be advantageous to these

efforts.

To date, only a few studies have investigated molecular-level analyses to identify sex or

reproductive stages in marine invertebrates. For example, contentious studies on vertebrate

steroids explored these organic molecules as an analyte for tracking mollusk reproduction

processes and stages.[4] Given the lack of functional receptors for androgens and estrogens in

marine mollusks, vertebrate-specific steroids may not be critical analytes that play a role in

reproduction; they have yet to yield viable assays for mollusk reproductive stages.[5] Further

research on gonad lipids [6], specific genetic transcripts [7,8], and RNA/DNA ratios [9] have also not

provided definitive, reproducible assays for the determination of mollusk sexual

maturation. Arcos et al. [10] explored the use of enzyme-linked immunoassay assays (ELISA) to

quantify proteins associated with vitellogenesis in order to determine oyster oocyte

maturation. Although the two proteins targeted by Arcos et al.[10] allowed distinction of males

and females, multiple mechanisms for maturation processing may be at play and, to date,

unidentified.[4] Supporting this hypothesis, Li et al. [11] used an alternative protein-profile approach

to simultaneously examine 139 peptides in the Eastern oyster, *Crassostrea virginica*, and

revealed that numerous unidentified proteins change in abundance with relation to maturation

and the overall protein profile could predict gender differences.

Since male and female geoduck gonads are morphologically and functionally distinct at

reproductive maturity, molecular assays to track these early biochemical changes could help

identify sex and indicate when gametes are mature. Significant changes in both gene and protein

expression are expected as gonad tissue differentiates. Proteomic assays designed to detect these

expression changes could lead to non-lethal, informative, and efficient tools for selecting

broodstock. Here, we apply mass spectrometry (MS)-based proteomics technology to provide a

large-scale, unbiased approach for examining relative abundances of all proteins present in

gonad tissue across several maturation stages of both female and male geoducks. Whole

proteome characterization via data dependent acquisition (DDA) provided a framework for

understanding molecular processes behind reproductive differentiation and directed peptide

selection for the determination of sex and maturation stage in subsequent downstream MS

analyses. Selected peptides underwent targeted proteomic analyses via selected reaction

monitoring (SRM) to more accurately quantify geoduck gonad proteins. These analyses were

used to increase our fundamental understanding of sex and stage determination in marine

mollusks and to develop non-invasive peptide assays on the circulating hemolymph.

Methods

Tissue Sampling

Geoduck clams were collected in November 2014 from Nisqually Reach, Washington

(latitude:47 08.89, longitude:122 47.439 WGS84). Clams were collected at depths between 9 to

14 meters from a sandy substrate. Gonad tissue and hemolymph from geoduck clams at early-,

mid-, and late-stage gonad maturation, from both males and females, were characterized

histologically. Female reproductive maturation stages were categorized as follows: early-stage

(no secondary oocytes, or oocytes that measure ~5-15μ), middle-stage (secondary oocytes ~50-

70μ), and late-stage (secondary oocytes ~65-85μ). Male reproductive maturation stages were

characterized as follows: early-stage (mostly somatic cells and ~5% spermatid composition per

acinus), middle-stage (about equal parts somatic cells and reproductive tissue and ~50%

spermatid composition per acinus), and late-stage (very little somatic cells and ~75-90%

spermatid composition per acinus). Details of gonadal maturation classification and histological

details have been previously described.[12] Gonad tissue samples were taken from 3 early-stage

females (fG03, fG04, fG08), 3 mid-stage females (fG34, fG35, fG38), 3 late-stage females

(fG51, fG69, fG70), 3 early-stage males (mG02, mG07, mG09), 3 mid-stage males (mG41,

mG42, mG46), and 3 late-stage males (mG65, mG67, mG68) with identification codes

corresponding to those reported in Crandall et al. (2015).[12] Hemolymph tissue samples were

taken from 3 early-stage females (fG18, fG29, fG30), 2 mid-stage females (fG25, fG35), 3 late-

stage females (fG51, fG69, fG70), 3 early-stage males (mG17, mG20, mG028), 2 mid-stage

males (mG42, mG46), and 3 late-stage males (mG65, mG67, mG68).

Protein Preparation

Gonad tissue from each of the eighteen geoduck clams was sonicated in lysis buffer (50

mM NH4HCO3, 6 M urea). Gonad tissue protein homogenate content and hemolymph protein

content were quantified using Pierce BCA Protein Assay Kit (Thermo Fisher Scientific,

Waltham, MA; catalog #23225). Protein (100 ug - gonad samples, 50 μg - hemolymph) was

evaporated to ~20 µl and resuspended in 100 µl of lysis buffer followed by sonication. Protein

digestion for gonad tissue samples and hemolymph followed the protocol outlined in Timmins-

Schiffman et al. [13] Briefly, each sample was incubated with TCEP buffered at pH 8.8 (1 hr,

37°C). Samples were alkylated with iodoacetamide (IAM; 1hr, 20°C) followed by a 1 hr

incubation with dithiothreitol to absorb any remaining IAM. To each sample, NH_4HCO_3 and

HPLC grade methanol were added to dilute urea and to increase solubilization of membrane

proteins. Samples were digested overnight with trypsin at 37°C. Digested samples were

evaporated and reconstituted in 5% acetonitrile (ACN) + 0.1% trifluoroacetic acid (TFA) (100

µl) and pH was decreased to < 2. Desalting of the samples was done using Macrospin columns

(sample capacity 0.03-300 µg; The Nest Group, Southborough, MA, USA) following the

manufacturer's specifications. Dried peptides were reconstituted in 100 µl of 5% ACN + 0.1%

formic acid.

Global Proteome Analysis: Data-Dependent Acquisition (DDA) LC-MS/MS

Data-dependent acquisition (DDA), also referred to as shotgun proteomics, was

performed to examine gonad proteomic profiles across samples (Figure 1). Liquid

chromatography coupled with tandem mass spectrometry (LC-MS/MS) was accomplished on a

Q-Exactive-HF (Thermo) on technical triplicates for each sample. The analytical column (20 cm

long) was packed in-house with C18 beads (Dr. Maisch, 0.3 µm) with a flow rate of 0.3 µl/min.

Chromatography was accomplished with an increasing ratio of solvent A (ACN + 0.1% formic

acid): solvent B (water + 0.1% formic acid). The solvent gradient consisted of: 0-1 minutes 2%

A; 1-60 minutes 5% A; 60-61 minutes 35% A; 61-71 minutes 80% A; 71-90 minutes 2%

A. Quality control standards (Pierce Peptide Retention Time Calibration mixture (PRTC) +

bovine serum albumin peptides (BSA)) were analyzed throughout the experiment to ensure

consistency of peptide detection and elution times.

Figure 1. Illustration of experimental setup and workflow for mass spectrometry data

acquisition and analysis. MS experimental workflow: 1. Data dependent acquisition (DDA) was

performed on the Q-Exactive-HF (QE) to assess the global proteomic differences in gonad (G)

tissue between sexes and maturation stages. 2. Targeted assay development followed the steps of

(a) data independent acquisition (DIA) on gonad tissue was also completed on the QE to create

spectral libraries for selected reaction monitoring (SRM) method development; (b&c) spectral

libraries were analyzed in PECAN using Skyline to select optimal transitions and design an

instrument method for SRM analyses; (d) SRM was completed on the TSQ Vantage for geoduck

peptide transitions in gonad and hemolymph (G & H); (e) Peptide transition detection and

quantification was performed in Skyline. The chromatograms of peptide KEEELIDYMVKQ

(from protein 130261_c0_seq1|m.17926) were collected using the 3 different MS approaches

(DDA, DIA, and SRM) from the same late-stage female. Black vertical lines indicate peak

integration boundaries, and colored peaks represent the different transitions (i.e. peptide

fragments) collected.

DDA protein identification and quantification

159   Gonad peptides were identified and proteins inferred using a proteome derived from a *de*

novo assembled transcriptome of a male and female geoduck clam gonad tissue libraries (NCBI

Bioproject Accession #PRJNA316216) (Del Rio-Portilla et al unpublished). Briefly, reads were

assembled using Trinity [14] and deduced protein sequences determined using the Transdecoder

algorithm within Trinity. Raw mass spectrometry data (PRIDE Accession #PXD003127) was

searched against the protein database (Supporting Information 1) using Comet v 2015.01 rev.2

specifying trypsin as the cleaving enzyme, two missed cleavages allowed, peptide mass tolerance

of 20 ppm, cysteine modification of 57 Da (from the IAM) and methionine modification of

15.999 Da (from oxidation) (Supporting information 2) to find peptide spectral

matches.[15,16] Protein inference and match probability were found using the Trans-Proteomic

Pipeline with a probability cut-off of 0.9.[17,18] Protein identifications were considered true

matches when the probability of the match was at least 0.9 and at least 2 independent spectra

were associated with the protein across all samples.

Non-metric multidimensional scaling analysis (NMDS) was used to determine the

similarity of technical replicates (Supporting Information 3) using the vegan package [19] in R v.

3.2.3 [20]. NMDS was performed on log-transformed data using a Bray-Curtis dissimilarity

matrix. As technical replicates clustered closely together and showed less variability than

biological replicates (Supporting Information 4), spectral counts were averaged across each

sample (n=18). Normalized spectral abundance factors (NSAF) were calculated on the averaged

spectral counts for all high confidence, detected proteins as a proxy for protein expression.[21]

Proteomic differences between sexes and maturation stages were explored with two methods: 1)

NMDS and analysis of similarity (ANOSIM) were used to compare the entire proteomic profiles

in multivariate space and 2) QSpec [22] was used to determine significant differences at the

individual protein level. NMDS was performed on NSAF data followed by ANOSIM in the

vegan package in R to determine the differences between sexes and maturation stages.

Differentially abundant proteins among the six conditions were identified using QSpec.[22]

Spectral counts were summed across technical replicates to create input files for QSpec

(http://www.nesvilab.org/qspec.php/) with data normalized by abundance. Nine comparisons

were analyzed for differentially abundant proteins: Early-stage male (EM) vs. early-stage female

(EF); mid-stage male (MM) vs. mid-stage female (MF); late-stage male (LM) vs. late-stage

female (LF); EM vs. MM; EM vs. LM; MM vs. LM; EF vs. MF; EF vs. LF; MF vs. LF. Proteins

were considered significantly different if the absolute value of the log fold change was ≥ 0.5 and

the absolute value of the z-statistic was ≥ 2.

Enrichment Analysis

We used enrichment analysis to determine which biological processes (represented by

detected proteins) were predominant in each gonad sample. GO enrichment analysis was

performed for each maturation stage (for each sex) on 1) unique proteins and 2) all detected

proteins. In both cases, the entire detected proteome was used as the background. The uniquely

detected proteins were the basis for targeted proteomics assay development (see below).

Enrichment analysis was also performed on sets of differentially abundant proteins from QSpec

analyses. Code used to perform enrichment analysis is available in a corresponding GitHub

repository (https://github.com/yeastrc/compgo-geoduck-public) as is the underlying code for the

end user web-interface (http://yeastrc.org/compgo_geoduck/pages/goAnalysisForm.jsp).

Briefly, a p-value was calculated representing the likelihood that a GO term would be as

represented as it was (or more) by chance among the set of tested proteins, given the annotation

of the geoduck gonad proteome (3,627 proteins). A p-value cutoff of 0.01 was used to ascribe

statistical significance to GO terms representing biological process.

GO terms were first assigned directly to the geoduck gonad protein names resulting from

the *de novo*-assembled transcriptome. This was done by assigning GO terms associated with

Uniprot-KB Swiss-Prot BLASTp hits. A p-value representing the statistical significance of the

representation of a GO term in a set of proteins was calculated using the hypergeometric

distribution using the following formula:

$$P(I) = \frac{\binom{A}{I} * \binom{T-A}{B-I}}{\binom{T}{B}}$$

Where A = total number of proteins submitted that have a GO annotation, B = the total number

of proteins in the background proteome annotated with the given GO term (or any of its

descendants), I (intersection of A and B) = the total number of submitted proteins annotated with

the given GO term (or any of its descendants), and T = the total number of annotated proteins in

the proteome background.

Then, the p-value describing the chance of having an intersection of size I or larger by

chance may be computed as:

P-value = $\displaystyle\sum_{i=I}^{\min(A,B)} P(i)$, where min (A,B) is the minimum of values A and B.

The p-value is then corrected for multiple hypothesis testing using the Bonferroni method

by multiplying the p-value by the number of GO terms tested (setting resulting values over 1 to

1). The number of GO terms tested equals the number of GO terms found (and all ancestors) for

the submitted set of proteins.

Complete directed acyclic graphs (DAG) that represent a subset of the whole GO DAG

were generated to visualize enriched biological processes. These DAGs were then filtered by

removing all childless terms that had an associated p-value > 0.01. The resulting DAGs were

then filtered using the same method, and this process repeated until no childless terms remained

with a p-value above the cutoff. The final result is a filtered subset of the GO DAG that contains

no leaf nodes with a p-value greater than the cutoff, but where a given term is guaranteed to have

all of its ancestor terms, even if those terms have a p-value greater than the cutoff. Having the

ancestor terms present is critical to visualization as they provide context for interpreting the

results. Visualizations of DAGs for combined full and unique proteomes for each sex and stage

are in Supporting Information 7 and proteins that contribute to specific enriched processes for all

enrichment analyses are included in Supporting Information 6.

Targeted Assay Development: Selected Reaction Monitoring (SRM)

A subset of proteins was chosen for development of a suite of targeted assays using

selected reaction monitoring (SRM) on the mass spectrometer. Based on the data dependent

acquisition, proteins that were detected only in one of these stages; early-stage males (EM),

early-stage females (EF), late-stage males (LM), or late-stage females (LF) were screened for

usable peptide transitions in SRM in Skyline daily v. 3.5.1.9706 [23] in gonad tissue.

Data independent acquisition (DIA) was used to generate spectral libraries for biomarker

development in the gonad tissue (Figure 1). Equal amounts of isolated peptides from the three

biological replicates for EM, EF, LM, and LF used in the DDA experiment (described above)

were pooled in equal quantities for DIA on the Q-Exactive HF (Thermo). Each sample included

a spiked-in internal quality control peptide standard (375 fmol PRTC + BSA; Pierce, hereafter

referred to as "QC"). Sample injections for all DIA experiments included protein (1 ug) plus the

QC (375 fmol) in a 2 μl injection. An analytical column (27cm) packed with C18 beads (3 μm;

Dr. Maisch) and a trap (3 cm) with C12 beads (3 μm; Dr. Maisch) were used for

chromatography. Technical replicate DIA spectra were collected in 4 m/z isolation width

windows spanning 125 m/z ranges each [24] (400-525, 525-650, 650-775, 775-900). For each

method, a gradient of 5-80% ACN over 90 minutes was applied for peptide spectra acquisition.

Raw data can be accessed via ProteomeXchange under identifier PXD004921. MSConvert [25]

was used to generate mzML files from the raw DIA files.

In order to generate spectral libraries for targeted method development, Peptide Centric

Analysis was completed with the software program PECAN (Ting et al., in review; Ting et al.,

2015[26]). Input files included the list of peptides generated for SRM (n=212), as described above,

and the mzML files generated from the raw DIA files. PECAN correlates a list of peptide

sequences with the acquired DIA spectra in order to locate the peptide-specific spectra within the

acquired DIA dataset. A background proteome of the *in silico* digested geoduck gonad proteome

was used.

The PECAN output file (.blib) was imported into Skyline to select peptide transitions and

create MS methods that would target specific peptides and transitions. Peptide transitions are the

reproducible fragments of peptides that are generated during the MS2 scan. Peptides reliably

fragment in the same way in the mass spectrometer, therefore transitions are a robust and

consistent signal of a peptide's presence.[27] Peptide transitions were selected if peak morphology

was uniform and consistent across the MS2 scans for technical replicates. Peptides were selected

for targeted analysis if they had $\geq$ 3 good quality transitions and there were $\geq$ 2 peptides per

protein. A maximum of 4 transitions per peptide were selected for targeted analysis and no more

than 3 peptides per protein were selected. The final list included 25 transitions for EM

biomarkers, 22 for EF, 133 for LM, and 52 for LF. This transition list was divided between two

method files for the final SRM analyses to provide adequate dwell time on individual transitions

in order to accurately detect and measure all peptides desired.[28]

Selected reaction monitoring (SRM) was carried out on a Thermo Vantage for all

eighteen geoduck gonad samples used in the original DDA analysis. Samples were prepared as

described above for DIA (1 µg of protein per 3 µl injection). A new C18 trap (2 cm) and C18

analytical column (27.5 cm) were used and each sample was analyzed in triplicate across two

MS experiments to cover the entire peptide transition list (n=212). Raw data can be accessed in

the PeptideAtlas under accession PASS00943.

Hemolymph from early-, mid-, and late-stage males and females was also assayed for

sex- and stage-specific biomarkers. An augmented SRM assay was used on the hemolymph that

included all the peptide transitions analyzed in the gonad with an additional 40 transitions from

five new proteins. These proteins were selected based on gonad proteome annotation to include

1) proteins that would likely be circulating in the hemolymph and 2) gonad proteins that had

homology with the mussel hemolymph proteome [29]. The additional peptides for SRM analysis of

hemolymph were from vitellogenin (2 proteins), glycogen synthase (2 proteins), and glycogenin-

1. PECAN generated spectral libraries, as described above, and Skyline software selected

hemolymph protein transitions to target during SRM analyses. For the new hemolymph proteins,

the minimums for peptides and transitions could not be met for every protein, but were still

included in the analysis. These transitions, and the previous gonad transitions, were analyzed

across sixteen geoduck hemolymph samples in two technical replicates, as described above for

the gonad. Some of these peptides yielded no data in the hemolymph, resulting in a dataset of

171 peptide transitions that were reliably detected. Raw data can be accessed in PeptideAtlas

under accession PASS00942.

Acquired SRM data in gonad and hemolymph were analyzed in Skyline for peptide

transition quantification. Skyline documents that were used to analyze the monitored peptide

transitions can be found on Panorama for the gonad dataset and the hemolymph datasets

(panoramaweb.org/labkey/geoduckrepro.url). Peak presence was determined based on

consistency of retention time (verified by spiked in QC peptides) and peak morphology.

All peptide transition peak intensities were exported from Skyline for automated peak

selection and peak integration analysis. QC internal standard transitions were monitored for

consistency across runs by calculating the coefficient of variation (CV) of transition peak area

across injections. Peak intensities for the geoduck peptide transitions were normalized by

dividing by the averaged intensities for the 6 QC peptide transitions that had the lowest CV in

the gonad data (CV < 13) and the 6 with the lowest CV in the hemolymph data (CV < 10).

NMDS and ANOSIM were performed on the QC-normalized SRM dataset as described

above for the DDA dataset. An initial NMDS showed that technical replicates clustered together

well and that variation was lower within biological replicates compared to between replicates,

therefore technical replicate peak intensities were averaged for the rest of the analysis

(Supporting Information 5). ANOSIM was performed using grouping by sex and reproductive

stage alone, as well as by a combined sex-stage factor. Coefficients of variation were calculated

for combined technical replicates using the raster package in R.

Eigenvector loadings were calculated for the gonad and hemolymph data using the vegan

package in R. For each dataset, the top 20 transitions with the combination of lowest p-value and

highest loading value were selected as the most "significant" biomarkers. Heatmaps of the log-

normalized transition intensities for these biomarker transitions were made using pheatmap[30] in

R, clustering rows (peptide transitions) and columns (samples) using euclidean distance and the

average clustering method.

Results

DDA Proteomics

The mass spectrometry data (PRIDE Accession #PXD003127) interpreted with a species-

and tissue-specific transcriptome yielded 3,627 proteins inferred with high confidence across all

gonad tissue samples (total spectral count across all replicates > 1) (Supporting Information

6). Female and male gonad proteomic profiles were more similar in early-stage maturation, with

proteomes diverging as reproductive maturity advanced (**Figure 2**). Proteomic profiles were

significantly different between sexes (R = 0.4122, p = 0.002) and maturation stages (R = 0.3025,

p = 0.004).

Unique proteomic profiles include the set of proteins that were detected in a specified sex

or specific stage within sex in the DDA analysis. The female unique proteomic profile across all

stages contained 305 proteins whereas 522 proteins were unique to males. The number of

proteins unique to a specific maturation stage increased with maturity from 24 and 20 in early-

stage females and males, respectively, to 161 and 145 in late-stage females and males (Figure 3).

These unique proteomes were used to develop the downstream SRM assays. Enriched GO

biological processes for the unique and full proteomes for each sex and stage are summarized in

Table 1. Only the most specific GO terms (i.e., farthest down on the DAG) are listed; parent

terms were frequently enriched as well and can be seen in the DAGs (Supporting Information 7).

345 *Table 1. Enriched biological process GO terms for the detected and unique gonad proteomes. If*

346 *no GO terms are listed, no enriched processes were identified*

Gonad Stage	Detected Proteome GO terms	Unique Proteome GO terms
Early-stage Female	Actin filament organization; Translation	Protein phosphorylation
Mid-stage Female	Hexose metabolic process; Translation; Translational initiation	Regulation of hydrolase activity
Late-stage Female	DNA replication; Translation; Translational initiation	Protein glycosylation; Signal peptide processing; Homophilic cell adhesion via plasma membrane adhesion molecules
Early-stage Male	Translation; Oxidation-reduction process; Small molecule metabolic process	
Mid-stage Male	Translation; ATP metabolic process; Pyruvate metabolic process; Oxidation-reduction process; Nucleotide phosphorylation; Single organism catabolic process	
Late-stage Male	Translation; Cellular macromolecular complex assembly; Nucleotide metabolic process	Microtubule-based movement

Figure 2. Non-metric multidimensional scaling plot (NMDS) of geoduck gonad whole proteomic

profiles generated by data dependent acquisition. Gonad proteomes differ among clams by both

sex (male = orange, female = blue) and stage (early-stage = circles, mid-stage = squares, late-

stage = triangles; p<0.05).

Numerous proteins were differentially abundant in comparisons between reproductive

stage and sex based on analysis with QSpec. The number of differentially abundant proteins

within a stage and between sexes increased from 387 between early-stage males and females to

1035 between late-stage males and females (Figure 4, Supporting Information 6). There were

more differentially abundant proteins between the stages within males than females (Supporting

Information 6). Enrichment analysis revealed which processes were over-represented in the sets

of differentially abundant proteins, when compared to the entire proteome (Figure 4 and Table

2).

Table 2. GO biological processes enriched in proteins that were differentially abundant between

stages within a sex (e.g. early- vs. mid-stage female) or between sexes within a stage (e.g LF vs.

LM). Only the most specific GO terms in the GO hierarchy are listed.

	Mid-stage Female	Late-stage Female	Early-stage Male	Mid-stage Male	Late-stage Male
Early-stage Female	Translation; DNA duplex unwinding	Translation; DNA duplex unwinding	Translation		
Mid-stage Female		DNA replication; DNA replication initiation; DNA duplex unwinding		Translation	
Late-stage Female	DNA replication; DNA				Translation; Microtubule-based process;

	replication initiation; DNA duplex unwinding				Translational initiation; Regulation of translational initiation
Early-stage Male				Ribonucleotide biosynthetic process; Purine ribonucleotide metabolic process; Purine ribonucleotide triphosphate biosynthetic process; Purine ribonucleoside metabolic process; Cytoskeletal anchoring at plasma membrane	Microtubule-based process; Oxidation-reduction process; Purine ribonucleoside triphosphate biosynthetic process; Purine ribonucleotide metabolic process
Mid-stage Male	Translation		Ribonucleotide biosynthetic process; Purine ribonucleotide metabolic process; Purine ribonucleotide triphosphate biosynthetic process; Purine ribonucleoside metabolic process; Cytoskeletal anchoring at plasma membrane		Monocarboxylic acid metabolic process; Microtubule-based movement

Figure 3. Venn diagram with corresponding enriched biological processes for sex-stage full

proteomes of early-stage female (blue), early-stage male (yellow), late-stage female (green), and

late-stage male (red). Venn diagrams produced in Venny. [31]

Figure 4. Venn diagram with corresponding enriched biological processes for differentially

abundant proteins from QSpec comparisons between early-stage males (EM) and mid-stage

males (MM); EM and late-stage males (LM); early-stage females (EF) and mid-stage females

(MF); EF and late-stage females (LF). Venn diagrams produced in Venny.[31]

Targeted Assay Development: Selected Reaction Monitoring (SRM)

SRM was applied to create peptide assays that could resolve geoduck sex and maturation

stages. Peptide transitions in the gonad (n=212) and hemolymph (n=171) derived from proteins

detected in a single sex-maturation stage from one of the following; early-stage male (EM),

early-stage female (EF), late-stage male (LM), and late-stage female (LF) proteomes were

measured across 18 geoduck gonad samples and 16 hemolymph samples that included the 3

maturation stages and both sexes (Supporting Information 8). Many transitions had stable

coefficients of variation across all biological replicates, although more than 6% of the peptide

transitions had CVs > 100, indicating a need for assay optimization (Supporting Information 9).

In an ordination plot, the gonad SRM data resolved males and females better in late-stage

samples than the early- or mid-stage (Figure 5A). There was significant separation based on

gonad SRM data for sex (R = 0.2373, p=0.003), stage (R=0.2189, p=0.007), and a combined sex-

stage factor (R=0.4132, p=0.001). The hemolymph SRM data only resolved the late-stage female

group from the rest of the geoduck (Figure 6A). There was significant separation of the

hemolymph proteomic profiles based on sex (R=0.1384, p=0.043) and sex-stage (R=0.4892,

p=0.001), but not by stage alone (R=0.1435, p=0.065).

Peptide transitions that directed the observed spread of the geoduck sex-stage data along

Axis 1 and Axis 2 in the NMDS plots (determined by eigenvector analysis) were considered key

analytes for differentiating groups. These peptide transitions represent a starting point for

biomarker assay optimization. Peptides that drove the separation of the male maturation stages

along Axis 1 in the gonad NMDS plot (Figure 5A) are from a set of 9 proteins: flap

endonuclease, IQ domain-containing protein K, WD repeat-containing protein on Y

chromosome, tetratricopeptide repeat protein 18, spectrin alpha chain, and four uncharacterized

proteins (Figure 5B). Proteins that played the most significant role in the separation of the female

stages along Axis 2 for the gonad data included WD repeat-containing protein on Y

chromosome, tetratripcopeptide repeat protein 18, spectrin alpha chain, centrosomal protein of

70 kDa, and four uncharacterized proteins. Analysis of the hemolymph SRM NMDS (Figure 6A)

reveals that the main peptide drivers along Axis 1 (separating late-stage females from all other

geoduck) were from three proteins: flap endonuclease, vitellogenin, and one uncharacterized

protein (Figure 6B). A cluster analysis of the 20 transitions that were primary drivers of sex

differentiation demonstrates that sex and stage groups from the gonad data clustered tightly, but

more specifically exposed a significant separation of males and females in the hemolymph data

(Figure 6B).

Figure 5. A) NMDS of geoduck gonad peptide transition abundance generated by selected

reaction monitoring. Gonad proteomes differ among clams by both sex (male = orange, female

= blue) and stage (early = circles, mid = squares, late = triangles; p<0.05). B) Heatmap of log-

normalized peptide transition intensities for the top 20 most significant transitions (based on

NMDS analysis (A)) across all samples (early-stage/circles, mid-stage/squares, and late-

stage/triangles males/orange and females/blue) gonad. Each row (peptide transition) is labeled

by its corresponding protein annotation in gray. Superscripts indicate for which NMDS axis a

transition is significant.

Figure 6. A) NMDS of geoduck hemolymph peptide transition abundance generated by selected

reaction monitoring. The main plot contains a point for each technical and biological replicate;

the inset (lower right-hand corner) shows the same data in a NMDS with the technical replicates

averaged. Hemolymph proteomes differ among clams by sex (male = orange, female = blue;

p<0.05). B) Heatmap of log-normalized peptide transition intensities for the top 20 most

significant transitions (based on NMDS analysis (A)) across all samples (early-stage/circles,

mid-stage/squares, and late-stage/triangles males/orange and females/blue) hemolymph. Each

row (peptide transition) is labeled by its corresponding protein annotation in gray. Superscripts

indicate for which NMDS axis a transition is significant.

Discussion

      Proteomics has the capacity to uncover molecular physiological processes underlying

functional phenotypic change, such as the maturation of reproductive tissue. We have

characterized the geoduck clam gonad proteome throughout reproductive maturation for both

males and females. Our data dependent analysis (DDA) approach yielded 3,627 detected proteins

across both sexes and three maturation stages. This is a significant escalation in our

understanding of proteomic responses in maturation stages of marine mollusks compared to the

study by Li et al. [32] that focused on protein-chip assays of 139 peptides. Based on the DDA data,

27 proteins from early- and late-stage male and female clams were chosen for selected reaction

monitoring (SRM), or targeted proteomics, to initiate development of a biomarker assay to

identify geoduck reproductive stages based on 85 peptides. Peptide transitions (n=212) were

targeted from the selected peptides as candidates for biomarker development in gonad and 171

transitions were targeted in hemolymph.

DDA

There are clear proteomic profile differences in the geoduck gonad by both sex and

maturation stage and these differences reveal the biochemical pathways underlying tissue

reproductive specialization. More proteins are differentially abundant between early and late

stages of reproductive maturity, compared to early and mid-stages, for both males and females,

reflecting that, as maturation progresses, gonad tissue becomes highly specialized to male or

female reproduction. Throughout geoduck reproductive maturation, the gonad goes from

undifferentiated connective tissue to a highly specialized reproductive tissue that can release

fully mature sperm and eggs. These dramatic phenotypic changes required for reproductive

maturity are visible with histology [12] and are realized via changes at the molecular level that

prompt the biochemical shifts necessary to create mature gonad. For example, the zona pellucida

(ZP) containing protein (cds.comp134923_c0_seq3|m.23445) and vitellogenin

(cds.comp144315_c0_seq1|m.50928) became significantly more abundant in the female gonad as

geoduck matured. The zona pellucida surrounds vertebrate oocytes and its invertebrate homolog

is the vitelline envelope, which is likely the source of this protein annotation and is the site of

sperm-oocyte recognition and binding. Vitellogenin is an egg yolk protein that is essential in

reproduction and development across taxa.

The sex- and stage-specific proteomes were enriched for specific biological processes,

further demonstrating tissue specialization. In females, the early-stage gonad proteome was

enriched in proteins involved in protein phosphorylation, the mid-stage was enriched in

regulation of hydrolase activity and carbohydrate metabolism, and the late-stage was enriched in

protein glycosylation, signal peptide processing, DNA replication, and cell adhesion. These

shifts in biological pathways clarify the mechanisms behind what has already been observed in

bivalve tissue morphology during maturation. The enrichment in the process "protein

phosphorylation" likely represents the activation of translated proteins required to effect gonad

structural and functional changes. The enrichment of carbohydrate metabolism in mid-stage

females reflects the increased mobilization of carbohydrates to support the energy-intensive

process of gonad maturation. Previous biochemical work has highlighted the importance of

glycogen metabolism during bivalve reproductive maturation [33,34] and carbohydrate metabolism

during vertebrate oocyte maturation [35,36], however, a more complex overview of gonad

carbohydrate metabolism is revealed in the proteomic data. The proteins contributing to the

enrichment of carbohydrate metabolism processes in mid-stage females are involved in glycan

metabolism (2 enzymes), glycolysis (3 enzymes), gluconeogenesis (2 enzymes), and the pentose

phosphate pathway (2 enzymes). These enzymes were detected in the geoduck gonad, but were

not differentially abundant. It is likely that, similar to other marine invertebrates, there is a rapid

upregulation of glycolysis in geoduck eggs upon fertilization.[37]

As geoduck females entered the late stages of maturation, more proteins were detected

that are related to cell replication as well as cell adhesion, which is an essential biological

process in maturing oocytes in vertebrates.[38,39] The strong cell adhesion signal in late-stage

females is driven in large part by the many cadherin and protocadherin proteins detected in a

single female. However, seven protocadherins were detected at significantly higher abundance in

more than a single late-stage female - compared to early- and mid-stages. Further, laminin, talin,

and vinculin were significantly less abundant in late-stage females compared to other females.

Evidence for the importance of DNA replication was more consistent: DNA helicase, DNA

ligase, proliferating cell nuclear antigen, and ribonucleoside-diphosphate reductase were all more

abundant in late-stage females than early- or mid-stages. This surge in DNA replication-

associated proteins is likely a reflection of an up-regulation of meiosis and/or mitosis. These

changes represent the diversity in function of female gonad tissue as it undergoes reproductive

maturation.

The male tissue similarly undergoes increased complexity and specialization: the early-

stage male gonad proteome was enriched in translation processes, oxidation-reduction processes,

and small molecule metabolic process; the mid-stage male proteome was enriched in ATP

metabolic process, pyruvate metabolism, oxidation-reduction process, and nucleotide

phorphorylation; and the late-stage for nucleotide metabolic process, microtubule-based

movement, and cellular macromolecular complex assembly. Translation, which was enriched in

every stage in both sexes, signifies the physiological need in the gonad for a larger suite of

proteins as it changes form and function. In mid-stage males, increases in proteins involved in

ATP metabolic processes reflected a greater metabolic demand in the tissue as it matures. This

was particularly reflected in the increased abundance of ATP synthase proteins in mid-stage

males relative to early-stage males and the decreased abundance of V-type proton ATPase

relative to early-stage males (this trend continued, although not significantly, in late-stage

males). ATP synthase produces ATP from a proton gradient while V-type proton ATPase

consumes ATP to fuel a proton gradient. These pieces of evidence suggest a physiological need

for increased ATP production as male maturation proceeds. However, glycolysis pathway

enzymes trended towards decreased abundance in mid- versus early-stage males, suggesting a

decrease in ATP generated via this pathway, which is responsible for the proton gradient that

drives ATP synthase. There may be a trend of ATP consumption over production during male

geoduck gametogenesis. Adenosine phosphate molecule consumption was observed in mussel

gonad tissue, where amounts of AMP, ADP, and ATP decreased as gametogenesis

progressed.[40] In late-stage males, the enriched processes reflected an increase in cell replication

during the last steps of sperm formation. Several proteins implicated in mitosis and meiosis were

detected at significantly increased abundance in late-stage males compared to early- and mid-

stage: centrosomal proteins, dynein, intraflagellar transport protein, kinesin, and alpha and beta

tubulin. All of these proteins are instrumental in the cell division and replication and their

importance at this maturation stage suggests the presence of rapidly dividing cells. This could be

a sign of both meiosis and mitosis. In mussels off the Atlantic coast of Spain, meiotic division

begins approximately four months before the onset of spawning, while mitosis reconstructs the

gonad during and after spawning.[41] These pieces of proteomic evidence point towards increased

cellular energy metabolism and cell division during the development of mature, motile sperm.

Even with a highly specific protein identification database, the DDA method excludes

many important, relatively low abundance proteins from analysis.[42] The ten most abundant

proteins (as measured by total spectral count) for each sex-maturation stage accounted for 14-

26% of the total spectral counts across all proteins for a given sex-stage. Many of these proteins

are "housekeeping" proteins, such as actin and myosin, and their dominance in the peptide

mixture was likely masking many informative, low abundance peptides. Similarly, in zebrafish,

highly abundant vitellogenin (also highly abundant in the mid- and late-stage females in this

study) made it difficult to detect lower abundant proteins using DDA.[43] Vitellogenin can be an

informative biomarker, but by masking the detection of other proteins that may have a more

nuanced expression profile, it and other highly abundant proteins can obscure the finer-scale

changes that occur.

SRM

DDA analysis yields biologically relevant results, but for practical purposes of biomarker

development we do not need the abundance profiles of >3000 proteins. Selected reaction

monitoring allows for the selection and accurate detection of only the most informative proteins'

peptides to address a specific hypothesis. We leveraged the DDA dataset to create informative

SRM assays of geoduck sex and maturation stage in both gonad and hemolymph. Both of these

tissues have the potential for being the basis of non-lethal assays for broodstock assessment in

aquaculture. Reproductive maturation is a complex process and involves suites of interacting

proteins to achieve the end goal of mature gametes. By applying an unbiased method in the first

step of our biomarker development (DDA) we were able to detect and include these informative

peptides in our assay.

In the gonad tissue, we measured 212 peptide transitions that could differentiate males

and females as well as early and late maturation stages. Our assay is based on proteins that are

involved in calcium ion binding, phosphorylation, meiosis, DNA replication and repair, and

membrane structure. Additionally, seven of these informative proteins were unannotated, so we

cannot know their function based on homology. These unknown proteins may represent taxon-

specific reproductive proteins and warrant further characterization. Reproductive proteins are

some of the fastest-evolving, resulting in many species-specific proteins.[44] A subset of the

peptide transitions from these proteins (n=29) were deemed highly informative based on our

analysis and would be good candidates for a more streamlined SRM assay. These transitions

correspond to six annotated proteins and five unannotated proteins. The annotated proteins are

implicated in cytoskeleton structure (spectrin alpha chain), DNA replication and repair

(centrosomal protein of 70 kDa, flap endonuclease), and proteins that were identified by motifs

and thus yield little functional information (WD repeat-containing protein on Y chromosome, IQ

domain containing protein K, and tetracopeptide repeat protein 18). The 29 transitions were

almost uniformly at higher abundance in males than females, which is likely because we

included more proteins derived from the late-stage male proteome in our assay since these

yielded transitions with higher MS-based ion intensities.

Hemolymph is a more reliable non-lethal option for sampling and possesses differentially

abundant peptides based on sex and reproductive stage. The peptide-based assay developed from

the evidence found in the gonad tissue that revealed sex and stage was applied to circulating

hemolymph. To the gonad tissue SRM transition list we added 40 more transitions

that were related to reproductive maturation. Proteome profiles of bivalve hemocytes can reveal

patterns of reproductive status, such as differing oocyte quality in Pacific oysters, as measured by

larval production.[45] However, biomarkers of tissue-specific changes can be difficult to detect in

circulating fluid since hemolymph (or blood) is not the primary site of physiological change. The

sensitivity of targeted proteomics has a lower limit of detection than DDA and is sensitive

enough to detect some biomarkers of interest.[46] But even if the same proteins were circulating in

the hemolymph as were detected in the gonad, they may not be informative because 1) the

proteins could be fragmented in ways that make them undetectable with our assay or 2) the

protein abundance could be changing out of sync with reproductive changes. We were able to

collect data on 171 peptide transitions that were derived from gonad tissue but detectable in the

hemolymph. Peptide transitions from 16 proteins were reliably detected in both gonad and

hemolymph, however, only two proteins had transitions that were highly informative in both

tissues: flap endonuclease and an uncharacterized protein. The inclusion of vitellogenin in our

assay confirmed its presence in geoduck hemolymph during later stages of reproductive maturity

in females, but since it was such a strong physiological signal it dominated our statistical

analysis. It was similarly the strongest signal in hemocytes of maturing Pacific oysters [45] and the

basis of a SRM assay developed in sea turtles [47]. Since we were limited to the proteins from the

gonad DDA dataset for our assay development, there are likely additional circulating proteins

that would expand our hemolymph assay and make it informative beyond distinguishing late-

stage females.

Conclusions

Through characterization of the geoduck gonad proteome and development of proteomic

assays for reproductive maturity, we not only exponentially increased the molecular resources

available for this species, but we also developed effective tools for non-lethal detection of sex

and maturation stage in this species. The gonad peptide transitions could be further decreased to

a set of 29 transitions that would accurately predict a) geoduck sex and b) reproductive

maturation status. In the hemolymph, 20 peptide transitions can accurately differentiate late-

stage female geoduck from others. Together these molecular tools can directly address the

geoduck production problem of asynchronous spawning due to different maturation stages and

unknown sexes.

Acknowledgements

This work was supported by and funded by a grant from the University of Washington

Royalty Research Fund (to SR), a Training Grant from the National Institutes of Health (to ETS)

(T32 HG00035), and the University of Washington's Proteomics Resource (UWPR95794). We

would like to thank Jimmy Eng and Priska von Haller for their help in data acquisition and

analysis; Sonia Ting, Jarrett Egertson, Lindsay Pino, Yuval Boss, and Nick Shulman for

assistance with data analysis; and the Genome Sciences Information and Technology group for

their technical support. ETS and BLN would like to thank IJE and TAN for their ongoing

inspiration.

Supporting Information. The following files are available free of charge via the internet

goo.gl/uoi9H1

S1-Pgenerosa-gonad-protein-v3.fasta Deduced protein sequences derived from gonad

transcriptome sequencing

S2-comet.params Parameter file used in Comet searches for the DDA data

S3-technical-reps.pdf NMDS of the whole proteomic profiles (DDA data) for the three

biological replicates and corresponding three technical replicates for each sex and maturation

stage. Blue points represent female proteomes and orange represent male. Shapes and shade

(light to dark) represent the different stages: circles for early-stage, squares for mid-stage, and

triangles for late-stage

S4-technical-reps-box-plots.pdf Box plot of the coefficients of variation for protein spectral

counts from DDA data A) across technical replicates for each geoduck biological replicate and

B) across biological replicates for each sex-maturation stage. CVs were calculated in R using the

raster package. X-axis labels correspond to the geoduck sex and maturation stage, indicated by

e.g. "EF3" represents early-stage female sample 3. The boxes represent the upper and lower

quartiles of the data distribution; horizontal black line represents median value; "whiskers"

extend to the greatest and least values, excluding outliers; open circles represent outliers (more

or less than 3/2 times the upper or lower quartiles

S5-NMDS-SRM-tech-reps.pdf NMDS of gonad SRM data for the 3 biological replicates and

corresponding three technical replicates for each sex and maturation stage. Blue points represent

female proteomes and orange represent male. Shapes and shade (light to dark) represent the

different stages: circles for early-stage, squares for mid-stage, and triangles for late-stage

S6-spc-nsaf-prot-names.txt All identified proteins from the DDA experiment with Uniprot

annotations (e-value cut-off of 1E-10), total spectral counts for each technical replicate,

calculated normalized spectral abundance factor (NSAF) for combined technical replicates, and

indication of significantly differentially abundant proteins by pairwise comparison. Columns

containing spectral count data have headers "SpC" followed by the biological replicate number,

sex, maturation stage, and technical replicate (for example, "SpC 3FE" is technical replicate 1

from early-stage female 3). Notation for NSAF is similar to SpC. The 9 columns in the sheet

have headers such as "EFvLF" (comparison between early- and late-stage females) have

asterisks in the cells that correspond to proteins that were differentially abundant for each given

comparison. The last two columns ("NMDS Gonad" and "NMDS Hemolymph") have asterisks

in cells that correspond to the proteins with peptide transitions that contribute significantly to the

SRM NMDS plot distributions

S7_DAGs.pdf Enrichment plots for early- and mid-maturation stage geoduck (EM, EF, MM,

MF). Processes colored in blue were enriched from the set of all proteins detected in the given

sex-stage, while red processes are from the protein set that was uniquely detected in a sex-stage.

Darker colors represent higher significance of enrichment

S8-SRM-data.xlsx Raw Skyline output (in the tab "Skyline output) and peak intensities

normalized by QC peptide abundance ("Normalized Intensities") for the gonad and hemolymph

SRM data.

S9-SRM-cv-boxplots.pdf Integrated peak area coefficient of variation values across all gonad

peptide transitions (n=212) and hemolymph peptide transitions (n=171) A) across technical

replicates for each geoduck biological replicate in gonad tissue, B) across biological replicates

for each sex-maturation stage in gonad tissue, C) across technical replicates for each geoduck

biological replicate in hemolymph, and D) across biological replicates for each sex-maturation

stage in hemolymph. CVs were calculated in R using the raster package. X-axis labels

correspond to the geoduck sex and maturation stage, indicated by e.g. "EF3" represents early-

stage female sample 3. The boxes represent the upper and lower quartiles of the data distribution;

horizontal black line represents median value; "whiskers" extend to the greatest and least values,

excluding outliers; open circles represent outliers (more or less than 3/2 times the upper or lower

quartiles)

References

(1) Newell, R. I. Ecosystem influences of natural and cultivated populations of suspension-
feeding bivalve mollusks: A review. *J. Shellfish Res.* **2004**, *23* (1), 51–62.

(2) Hoffmann, A.; Bradbury, A.; Goodwin, C. L. Modeling geoduck, Panopea abrupta (Conrad,
1849) population dynamics. I. Growth. *J. Shellfish Res.* **2000**, *19* (1), 57–62.

(3) Northern Economics, I. *The Economic Impact of Shellfish Aquaculture in Washington,*
Oregon and California. Prepared for Pacific Shellfish Institute; 2013.

(4) Scott, A. P. Do mollusks use vertebrate sex steroids as reproductive hormones? II. Critical
review of the evidence that steroids have biological effects. *Steroids* **2013**, *78* (2), 268–281.

(5) Fernandes, D.; Denise, F.; Barbara, L.; Cinta, P. Biosynthesis and metabolism of steroids in
molluscs. *J. Steroid Biochem. Mol. Biol.* **2011**, *127* (3-5), 189–195.

(6) Palacios, E.; Racotta, I. S.; Arjona, O.; Marty, Y.; Le Coz, J. R.; Moal, J.; Samain, J. F.
Lipid composition of the pacific lion-paw scallop, Nodipecten subnodosus, in relation to
gametogenesis. *Aquaculture* **2007**, *266* (1-4), 266–273.

(7) Anantharaman, S.; Craft, J. A. Annual variation in the levels of transcripts of sex-specific
genes in the mantle of the common mussel, Mytilus edulis. *PLoS One* **2012**, *7* (11), e50861.

(8) Ciocan, C. M.; Cubero-Leon, E.; Minier, C.; Rotchell, J. M. Identification of reproduction-
specific genes associated with maturation and estrogen exposure in a marine bivalve
Mytilus edulis. *PLoS One* **2011**, *6* (7), e22326.

(9) Li, Q.; Qi, L.; Makoto, O.; Katsuyoshi, M. Seasonal biochemical variations in Pacific oyster
gonadal tissue during sexual maturation. *Fish. Sci.* **2000**, *66* (3), 502–508.

(10) Arcos, F. G.; Ibarra, A. M.; del Carmen Rodríguez-Jaramillo, M.; García-Latorre, E. A.;
Celia, V.-B. Quantification of vitellin/vitellogenin-like proteins in the oysterCrassostrea

corteziensis(Hertlein 1951) as a tool to predict the degree of gonad maturity. *Aquac. Res.*

2009, *40* (6), 644–655.

(11) Li, Y.; Yaqin, L.; Ghazala, S.; Wikfors, G. H. Non-lethal determination of sex and

reproductive condition of Eastern oysters Crassostrea virginica Gmelin using protein

profiles of hemolymph by Proteinchip® and SELDI-TOF-MS technology. *Aquaculture*

2010, *309* (1-4), 258–264.

(12) Crandall, G.; Roberts, S. Reproductive Maturation in Geoduck clams (Panopea generosa)

https://figshare.com/articles/Reproductive_Maturation_in_Geoduck_clams_Panopea_gener

osa_/3205975/1. DOI: https://dx.doi.org/10.6084/m9.figshare.3205975.v1

(13) Timmins-Schiffman, E.; Nunn, B. L.; Goodlett, D. R.; Roberts, S. B. Shotgun proteomics as

a viable approach for biological discovery in the Pacific oyster. *Conserv Physiol* **2013**, *1*

(1), cot009.

(14) Grabherr, M. G.; Haas, B. J.; Yassour, M.; Levin, J. Z.; Thompson, D. A.; Amit, I.;

Adiconis, X.; Fan, L.; Raychowdhury, R.; Zeng, Q.; et al. *Full-length transcriptome*

assembly from RNA-seq data without a reference genome; 2011.

(15) Eng, J. K.; Jahan, T. A.; Hoopmann, M. R. Comet: an open-source MS/MS sequence

database search tool. *Proteomics* **2013**, *13* (1), 22–24.

(16) Eng, J. K.; Hoopmann, M. R.; Jahan, T. A.; Egertson, J. D.; Noble, W. S.; MacCoss, M. J.

A deeper look into Comet--implementation and features. *J. Am. Soc. Mass Spectrom.* **2015**,

26 (11), 1865–1874.

(17) Pedrioli, P. G. A. Trans-Proteomic Pipeline: A Pipeline for Proteomic Analysis. In *Methods*

in Molecular Biology; 2009; pp 213–238.

(18) Deutsch, E. W.; Mendoza, L.; Shteynberg, D.; Slagel, J.; Sun, Z.; Moritz, R. L. Trans-

Proteomic Pipeline, a standardized data processing pipeline for large-scale reproducible

proteomics informatics. *Proteomics Clin. Appl.* **2015**, *9* (7-8), 745–754.

(19) Oksanen, J.; Guillaume Blanchet, F.; Friendly, M.; Kindt, R.; Legendre, P.; McGlinn, D.;

Minchin, P. R.; O'Hara, R. B.; Simpson, G. L.; Solymos, P.; et al. *vegan: Community*

Ecology Package; 2016.

(20) R Core Team. *R: A language and environment for statistical computing*; R Foundation for

Statistical Computing, Vienna, Austria, 2015.

(21) Florens, L.; Carozza, M. J.; Swanson, S. K.; Fournier, M.; Coleman, M. K.; Workman, J.

L.; Washburn, M. P. Analyzing chromatin remodeling complexes using shotgun proteomics

and normalized spectral abundance factors. *Methods* **2006**, *40* (4), 303–311.

(22) Choi, H.; Fermin, D.; Nesvizhskii, A. I. Significance analysis of spectral count data in label-

free shotgun proteomics. *Mol. Cell. Proteomics* **2008**, *7* (12), 2373–2385.

(23) MacLean, B.; Tomazela, D. M.; Shulman, N.; Chambers, M.; Finney, G. L.; Frewen, B.;

Kern, R.; Tabb, D. L.; Liebler, D. C.; MacCoss, M. J. Skyline: an open source document

editor for creating and analyzing targeted proteomics experiments. *Bioinformatics* **2010**, *26*

(7), 966–968.

(24) Panchaud, A.; Scherl, A.; Shaffer, S. A.; von Haller, P. D.; Kulasekara, H. D.; Miller, S. I.;

Goodlett, D. R. Precursor acquisition independent from ion count: how to dive deeper into

the proteomics ocean. *Anal. Chem.* **2009**, *81* (15), 6481–6488.

(25) Chambers, M. C.; Maclean, B.; Burke, R.; Amodei, D.; Ruderman, D. L.; Neumann, S.;

Gatto, L.; Fischer, B.; Pratt, B.; Egertson, J.; et al. A cross-platform toolkit for mass

spectrometry and proteomics. *Nat. Biotechnol.* **2012**, *30* (10), 918–920.

(26) Ting, Y. S.; Egertson, J. D.; Payne, S. H.; Kim, S.; MacLean, B.; Käll, L.; Aebersold, R.;

Smith, R. D.; Noble, W. S.; MacCoss, M. J. Peptide-Centric Proteome Analysis: An

Alternative Strategy for the Analysis of Tandem Mass Spectrometry Data. *Mol. Cell.*

Proteomics **2015**, *14* (9), 2301–2307.

(27) Wolf-Yadlin, A.; Hautaniemi, S.; Lauffenburger, D. A.; White, F. M. Multiple reaction

monitoring for robust quantitative proteomic analysis of cellular signaling networks. *Proc.*

Natl. Acad. Sci. U. S. A. **2007**, *104* (14), 5860–5865.

(28) Picotti, P.; Aebersold, R. Selected reaction monitoring-based proteomics: workflows,

potential, pitfalls and future directions. *Nat. Methods* **2012**, *9* (6), 555–566.

(29) Moreira, R.; Pereiro, P.; Canchaya, C.; Posada, D.; Figueras, A.; Novoa, B. RNA-Seq in

Mytilus galloprovincialis: comparative transcriptomics and expression profiles among

different tissues. *BMC Genomics* **2015**, *16* (1), 728.

(30) Kolde, R. *pheatmap: Pretty Heatmaps*; 2015.

(31) Oliveros, J. C. *Venny. An interactive tool for comparing lists with Venn's diagrams*; 2007-

2015.

(32) Li, Y.; Yaqin, L.; Ghazala, S.; Wikfors, G. H. Non-lethal determination of sex and

reproductive condition of Eastern oysters Crassostrea virginica Gmelin using protein

profiles of hemolymph by Proteinchip® and SELDI-TOF-MS technology. *Aquaculture*

2010, *309* (1-4), 258–264.

(33) Beukema, J. J.; De Bruin, W. Seasonal changes in dry weight and chemical composition of

the soft parts of the tellinid bivalve Macoma balthica in the Dutch Wadden Sea. *Neth. J. Sea*

Res. **1977**, *11* (1), 42–55.

(34) Li, Q.; Qi, L.; Makoto, O.; Katsuyoshi, M. Seasonal biochemical variations in Pacific oyster

gonadal tissue during sexual maturation. *Fish. Sci.* **2000**, *66* (3), 502–508.

(35) Dworkin, M. B.; Eva, D.-R. Metabolic regulation during early frog development:

Glycogenic flux in Xenopus oocytes, eggs, and embryos. *Dev. Biol.* **1989**, *132* (2), 512–

523.

(36) Harris, S. E.; Adriaens, I.; Leese, H. J.; Gosden, R. G.; Picton, H. M. Carbohydrate

metabolism by murine ovarian follicles and oocytes grown in vitro. *Reproduction* **2007**, *134*

(3), 415–424.

(37) Yasumasu, I.; Tazawa, E.; Fujiwara, A. Glycolysis in the eggs of the echiuroid, Urechis

unicinctus and the oyster, Crassostrea gigas. *Exp. Cell Res.* **1975**, *93* (1), 166–174.

(38) Fusi, F. M.; Vignali, M.; Busacca, M.; Bronson, R. A. Evidence for the presence of an

integrin cell adhesion receptor on the oolemma of unfertilized human oocytes. *Mol. Reprod.*

Dev. **1992**, *31* (3), 215–222.

(39) Campbell, S.; Swann, H. R.; Seif, M. W.; Kimber, S. J.; Aplin, J. D. Integrins and adhesion

mlecules: Cell adhesion molecules on the oocyte and preimplantation human embryo. *Hum.*

Reprod. **1995**, *10* (6), 1571–1578.

(40) Díaz Enrich, M. J.; Ibarguren, I. Implication of adenosine 3',5'-cyclic monophosphate,

guanosine 3',5'-cyclic monophosphate, adenosine 5'-mono-, di-, and triphosphate and

fructose-2,6-bisphosphate in the regulation of the glycolytic pathway in relation to the

gametogenic cycle in the mussel mytilus galloprovincialis Lmk. *Mol. Cell. Biochem.* **2003**,

252 (1-2), 339–345.

(41) Suárez, M. P.; Alvarez, C.; Molist, P.; Fs, A. J. Particular aspects of gonadal cycle and

seasonal distribution of gametogenic stages of Mytilus galloprovincialis cultured in the

estuary of Vigo. *J. Shellfish Res.* **2005**, *24* (2), 531–540.

(42) Pedersen, S. K.; Harry, J. L.; Sebastian, L.; Baker, J.; Traini, M. D.; McCarthy, J. T.;

Manoharan, A.; Wilkins, M. R.; Gooley, A. A.; Righetti, P. G.; et al. Unseen proteome:

mining below the tip of the iceberg to find low abundance and membrane proteins. *J.*

Proteome Res. **2003**, *2* (3), 303–311.

(43) Groh, K. J.; Nesatyy, V. J.; Segner, H.; Eggen, R. I. L.; Suter, M. J.-F. Global proteomics

analysis of testis and ovary in adult zebrafish (Danio rerio). *Fish Physiol. Biochem.* **2011**,

37 (3), 619–647.

(44) Swanson, W. J.; Vacquier, V. D. THE RAPID EVOLUTION OF REPRODUCTIVE

PROTEINS. *Nat. Rev. Genet.* **2002**, *3* (2), 137–144.

(45) Corporeau, C.; Vanderplancke, G.; Boulais, M.; Suquet, M.; Quéré, C.; Boudry, P.; Huvet,

A.; Madec, S. Proteomic identification of quality factors for oocytes in the Pacific oyster

Crassostrea gigas. *J. Proteomics* **2012**, *75* (18), 5554–5563.

(46) Klee, E. W.; Bondar, O. P.; Goodmanson, M. K.; Dyer, R. B.; Erdogan, S.; Bergstralh, E. J.;

Bergen H., R.; Sebo, T. J.; Klee, G. G. Candidate Serum Biomarkers for Prostate

Adenocarcinoma Identified by mRNA Differences in Prostate Tissue and Verified with

Protein Measurements in Tissue and Blood. *Clin. Chem.* **2012**, *58* (3), 599–609.

(47) Plumel, M. I.; Wasselin, T.; Plot, V.; Strub, J.-M.; Van Dorsselaer, A.; Carapito, C.;

Georges, J.-Y.; Bertile, F. Mass spectrometry-based sequencing and SRM-based

quantitation of two novel vitellogenin isoforms in the leatherback sea turtle (Dermochelys

coriacea). *J. Proteome Res.* **2013**, *12* (9), 4122–4135.