Title 1

Genome-wide association study in Collaborative Cross mice reveals a role for 2

Rhbdf2 **in skeletal homeostasis** 3

Short title 4

GWAS in mice reveals a new skeletal role for *Rhbdf2* 5

Authors 6

Roei Levy[1,2], Clemence Levet[3], Keren Cohen[1], Matthew Freeman[3], Richard Mott[4], 7

Fuad Iraqi[5], Yankel Gabet[1] 8

Affiliations 9

[1] Department of Anatomy and Anthropology, [2] Department of Human Molecular 10

Genetics and Biochemistry, and [5] Department of Clinical Microbiology and 11

Immunology, Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv 69978, 12

Israel. [3] Dunn School of Pathology, South Parks Road, Oxford OX1 3RE, UK. [4] UCL 13

Genetics Institute, University College London, Gower St., London, WC1E 6BT, UK. 14

Keywords Bone; Collaborative Cross; Rhbdf2; GWAS; iRhom2; cancellous bone; 15
bone mass; causal genes; femur; RIL 16

Abstract   17

Osteoporosis, the most common bone disease, is characterized by a low bone mass and   18

increased risk of fractures. Importantly, individuals with the same bone mineral density   19

(BMD), as measured on two dimensional (2D) radiographs, have different risks for   20

fracture, suggesting that microstructural architecture is an important determinant of   21

skeletal strength. Here we took advantage of the rich phenotypic and genetic diversity   22

of the Collaborative Cross (CC) mice. Using microcomputed tomography, we   23

examined key structural parameters in the femoral cortical and trabecular compartments   24

of male and female mice from 34 CC lines. These traits included the trabecular bone   25

volume fraction, number, thickness, connectivity, and spacing, as well as structural   26

morphometric index. In the mid-diaphyseal cortex, we recorded cortical thickness and   27

volumetric BMD.   28

The broad-sense heritability of these traits ranged between 50 to 60%. We   29

conducted a genome-wide association study to unravel 5 quantitative trait loci (QTL)   30

significantly associated with 6 of the traits. We refined each locus by combining   31

information obtained from the known ancestry of the mice and RNA-Seq data from   32

publicly available sources, to shortlist potential candidate genes. We found strong   33

evidence for new candidate genes, including *Rhbdf2*, which association to trabecular   34

bone volume fraction and number was strongly suggested by our analyses. We then   35

examined knockout mice, and validated the causal action of *Rhbdf2* on bone mass   36

accrual and microarchitecture.   37

Our approach revealed new genome-wide QTLs and a series of genes that have   38

never been associated with bone microarchitecture. This study demonstrates for the first   39

time the skeletal role of *Rhbdf2* on the physiological remodeling of both the cortical   40

and trabecular bone. This newly assigned function for *Rhbdf2* can prove useful in   41

deciphering the predisposing factors of osteoporosis and propose new investigative   42

avenues toward targeted therapeutic solutions.   43

Author summary   44

In this study, we used the novel mouse reference population, the Collaborative Cross   45

(CC), to identify new causal genes in the regulation of bone microarchitecture, a   46

critical determinant of bone strength. This approach provides a clear advantage in   47

terms of resolution and dimensionality of the morphometric features (versus humans)   48

and rich allelic diversity (versus classical mouse populations), over current practices 49

of bone-related genome-wide association studies.   50

Our genome-wide study revealed 5 loci significantly associated with microstructural   51

traits in the cortical and trabecular bone. We found strong evidence for new candidate 52

genes, in particular, *Rhbdf2*. We then validated the specific role of *Rhbdf2* on bone 53

mass accrual and microarchitecture using knockout mice. Importantly, this study is   54

the first demonstration of a physiological role for *Rhbdf2*.   55

Introduction   57

Osteoporosis is the most common bone disease in humans, affecting nearly half the US   58

and European population over the age of 50 years. With the globally increasing life   59

expectancy, osteoporosis and related bone fractures are becoming a pandemic health   60

and economic concern. By 2050, the world-wide incidence of hip fractures is expected   61

to increase by 2.5 to 6 fold [1,2]. Importantly, the mortality rate in the 12 months   62

following bone fracture is as high as 20% [3]. Risk of fracture is determined largely by 63

bone density and quality/strength, which are the end result of peak values achieved at 64

skeletal maturity and subsequent age and menopause-related bone loss. Genetic factors   65

have a major role in determining the wide range in the so-called "normal" peak bone   66

mass. Measures of bone status are inherently complex traits, as opposed to Mendelian   67

traits; i.e. they are controlled by the cumulative effect and interactions of numerous   68

genetic loci and environmental factors.   69

Genome-wide association studies (GWAS), including a large meta-analysis,   70

have identified more than 50 loci associated with bone mineral density (BMD) [4–10].   71

However, many other genes that were experimentally associated with bone mass were   72

not confirmed by GWAS in human cohorts [11–13]. This suggests that the BMD   73

phenotype does not capture the structural complexity of the bone; there may be other   74

relevant bone phenotypes not yet studied in human GWAS [12], which hitherto have   75

generally relied on areal bone mineral density (aBMD) as the sole bone feature. aBMD   76

measured by dual energy x-ray absorptiometry (DXA) is a two-dimensional projection   77

that cannot measure bone size, individual bone compartments' shape (whether   78

trabecular or cortical) or underlying microstructure, and thus likely conceals important   79

features which are assumed to be controlled by unique genetic determinants. Indeed,   80

there is a growing body of evidence that argues for distinct genetic influences of the   81

cortical and trabecular bone and thus they should be accordingly distinguished [8,9]. A   82

recent GWAS in the Collaborative Cross (CC) mice based on DXA failed to find any   83

heritability of BMD [14], whereas another report based on the same mouse panel   84

showed highly significant heritability levels in most of the cortical and trabecular   85

microstructural parameters measured by micro-computed tomography (μCT) [15]   86

Traditional peripheral quantitative CT (pQCT) has the capacity to distinguish   87

between the cortical and trabecular bone compartments, but it lacks the required   88

resolution to detect microstructural differences. A recent report based on high   89

resolution pQCT (HR-pQCT) data in humans, identified two novel bone-related loci,   90

thus far undetected by DXA and pQCT-based GWAS [9]. Another [16], found strong   91

genetic correlations between 1047 adult participants of the Framingham heart study,   92

therefore indicating that the heritability of bone microstructure constitutes a phenotypic   93

layer which is at least partially independent of DXA-derived BMD. Like HR-pQCT   94

studies in humans, understanding the genetic regulation of bone microstructural   95

parameters using μCT in small animals is likely to identify genetic factors distinct from   96

those previously identified for DXA-derived traits.   97

The CC mouse panel is designed to provide high resolution analysis of complex   98

traits, with particular emphasis on traits relevant to human health [17,18]. This unique   99

resource currently consists of a growing number of recombinant inbred lines (RIL)   100

generated from full reciprocal breeding of eight divergent strains of mice [19]. In   101

contrast to commonly used laboratory mouse strains, the ancestry of the CC lines   102

includes wild-derived inbred strains that encompass genetic variations accumulated   103

over ~1 million years [20]; more than 50 million single nucleotide polymorphisms   104

segregate in founders of the CC. The high genetic diversity means that QTLs can be   105

mapped using this panel that would have been invisible in a population that involved   106

only classical strains [21,22].   107

This claim is substantiated in a recent study that identified a genome-wide   108

significant association between *Oxt* (oxytocin) and *Avp* (vasopressin) and skeletal   109

microarchitecture in CC mice (Levy et al. 2015). Here, our GWAS in the CC mouse   110

panel identified a novel gene, *Rhbdf2,* associated with bone traits and using a specific   111

knockout model we validated its role in the regulation of cortical and trabecular bone   112

structure. This exemplifies the effectiveness and relative ease by which a GWAS with   113

a small CC population can associate a bone-related function to novel genes, and to   114

reveal overlooked key players in skeletal biology.   115

Results   116

CC lines widely differ in bone microarchitecture traits   117

We examined the variation in femoral cortical and trabecular microstructure between   118

34 unique CC lines totaling in 174 mice (71 females and 103 males, with an average of   119

4.25 mice per line). In the trabecular bone compartment we measured bone volume   120

fraction (BV/TV), trabecular number (Tb.N), thickness (Tb.Th), connectivity   121

(Conn.D), and spacing (or separation; Tb.Sp), as well as structural morphometric index   122

(SMI) of the trabecular framework. In the mid-diaphyseal cortex, we recorded cortical   123

thickness (Ct.Th) and volumetric bone mineral density (vBMD). These traits were   124

approximately normally distributed; BV/TV ranged from 0.017 to 0.26 (i.e. 1.7% to   125

26%; mean = 10.2%); Tb.N from 0.52 to 6.11 mm^{-1} (mean = 2.7 mm^{-1}); Tb.Th from 31   126

to 69 μm (mean = 47 μm); Conn.D from 10.9 to 268.3 mm^{-3} (mean = 104.2 mm^{-3}); SMI   127

from 0.6 to 3.3 (mean = 2.3); Tb.Sp from 0.16 to 0.7 mm (mean = 0.33 mm); Ct.Th   128

from 0.14 to 0.29 mm (mean = 0.2 mm); and vBMD from 402.5 to 809.2 $mgHA/cm^3$ 129

(mean = 581.1 $mgHA/cm^3$). μCT images taken from two mice with distinct cortical   130

(Fig. 1A) and trabecular (Fig. 1B) characteristics demonstrate the great variation in   131

bone traits due solely to the genetic background. Color-codes on the graphs in Fig. 2   132

indicate Duncan's least significance range (LSR), which dictates whether the mean   133

value of a line, or a group of lines, for a given trait differs to a degree of at least *P*-value   134

< 0.001 from any other group. LSR allows for a visual representation of the   135

heterogeneity amongst the lines.   136

With 11 distinct groups, vBMD (Fig. 2B) is the most heterogeneous trait, while 137

SMI and Conn.D are the least, with only 3 significantly distinct groups (Fig. 2A). 138

Notably, the heterogeneity of females is greater than that of males for cortical traits but 139

milder for trabecular traits (Figs. S1A1, 2 for males and S1B1, 2 for females). 140

To examine the inter-dependency between the traits, we assessed the correlation 142

between all the measured parameters, in a pairwise fashion using Pearson's correlation 143

test. The strongest correlation was between BV/TV and Tb.N (Pearson's $r = 0.94$), in 144

line with our previous findings [15], while the weakest was between Tb.N and vBMD 145

($r < 0.01$). There was also a moderately high correlation between Ct.Th and Tb.Th ($r =$ 146

0.61; and see Table S1). The correlation between sexes for each trait (Table 1) ranged 147

from $r = 0.75$ (Tb.Sp) to $r = 0.20$ (Ct.Th). Body weight (range = 17.4 - 35.0 gr) did not 148

significantly correlate with any of the traits ($r = 0.01$ for Conn.D to $r = 0.19$ for Ct.Th; 149

data not shown). After separating males from females the correlation slightly increased, 150

yet remained low. Weak correlation was found between weight and Tb.N, SMI, and 151

Ct.Th for females (Pearson's $r = -0.20, 0.23$, and 0.25 respectively), and between weight 152

and Tb.Th and Tb.Sp (Pearson's $r = 0.25$ and -0.25) for males. 153

Table 1 Heritability, sex correlations, and covariate interactions for trabecular and cortical traits

Trait	H^2	logP	H^2n	Sex Cor.	Interactions %
BV/TV	0.61	12.43	0.87	0.6695754	-
Tb.N	0.63	13.76	0.88	0.7628464	-
Tb.Th	0.54	9.08	0.83	0.7039887	34.70
Conn.D	0.56	9.90	0.84	0.5188039	-
SMI	0.55	9.43	0.84	0.2984166	26.02
Tb.Sp	0.63	13.45	0.88	0.754563	-
vBMD	0.62	12.15	0.87	0.6268818	53.92
Ct.Th	0.51	7.63	0.82	0.2055684	41.07

H^2 is the broad-sense heritability (which includes epistatic and environmental influences); logP is the negative 10-base logarithm of the P-value; H^2n is the line-mean heritability; Sex Cor. Is the sex correlation of each trait; and interactions % refers to the relative contribution of the cumulative covariate-interactions, which include sex, age, batch, month, season, year, and experimenter (see table S2).

QTL	Trait	Chr	logP	99th % thresh	H^2_r	Simulation					
						50% CI (Mb)		90% CI (Mb)		95% CI (Mb)	
						Position	Width	Position	Width	Position	Width
Trl7	BV/TV	11	7.60	4.90	0.69	116.6-116.7	0.12	113.6-118.1	4.50	112.1-118.3	6.40
Trl7	Tb.N	11	6.80	4.84	0.71	116.6-116.87	0.29	114.2-118.35	4.15	112.41-118.68	6.27
Trl8	Tb.Th	4	8.00	4.50	0.61	117.2-117.58	0.32	113.05-125.54	12.49	110.87-126.52	15.65
Trl9	Tb.Sp	5	9.40	6.01	0.78	105.78-106.14	0.35	101.6-109.11	7.54	99.8-110.39	10.62
Crl1	Ct.Th	4	8.20	4.80	0.80	9.29-9.72	0.43	4.0-11.7	7.70	3.4-11.8	8.49
Crl2	vBMD	3	9.80	4.93	0.86	97.2-97.4	0.20	94.4-103.1	8.50	93.22-104.3	10.80

Table 2 Positions of QTLs associated with trabecular and cortical traits

Chr = chromosome; logP = negative 10-base logarithm of P value; Sig = genome-wide significance level; 99th % threshold logP = threshold used to define cut-off for QTL peaks (Fig. 4); H^2_r = regional heritability (the proportion to which the locus explains the phenotypic variability). Positions and widths of the simulation-based 50, 90, and 95% CIs are given.

While in most lines the traits' correlations were predictive of a given line's rank, in others a less expected pattern was observed; e.g., IL-1513 displayed unusually extreme phenotypes for all trabecular traits and was at the higher end for BV/TV, Tb.N, Tb.Th, and Conn.D and at the lower end for SMI and Tb.Sp, but IL-188 was more discordant between these same traits (Fig. 2 and Table S3), illustrating unexpected co-variation of the traits in the CC.

Heritability and confounder-control

We quantitated the effects of the covariates sex, age, batch, month, season, year, and experimenter on each trait. Age ranged from 9 (n=6) to 13 (n=9) weeks and the mice

were dissected in 20 batches over a three-year course across 8 months during winter,   179

spring and summer, by two experimenters. Whereas age alone had no effect on any   180

trait, sex affected only Ct.Th; batch affected Tb.Th, vBMD, and Ct.Th; month affected   181

Tb.Sp, vBMD, and Ct.Th; season and year affected vBMD and Ct.Th; and Tb.Th and   182

Ct.Th were affected by experimenter. The cumulative effect of the covariates' pairwise   183

interactions was noted for Tb.Th, SMI, vBMD, and Ct.Th. (Table S2).   184

We then estimated the broad-sense heritability (H^2) of each trait among the CC   185

lines, which includes additive and non-additive epistatic effects and gene-environment   186

interactions. The greatest H^2 is seen for Tb.N (0.63, logP = 13.76; where logP stands   187

for the negative 10-base logarithm of the P value and tests the null hypothesis that the   188

heritability is zero), and the smallest for Ct.Th (0.51, logP = 7.63).   189

We calculated the heritability for the mean values in each line to get a better   190

representation of the percentage of genetic contribution to the phenotypic heterogeneity   191

by incorporating H^2 and the average number of lines [23]. This defines H^2n, which is   192

directly proportional to H^2 (Methods; Table 1) and ranges between 82 (Ct.Th) and 88%   193

(Tb.N and Tb.Sp).   194

Overall the cortical traits seemed more prone to covariate variation; they were   195

particularly sensitive to sex, batch, and season. This stands in contrast to our previous   196

results [15] where BV/TV, Tb.N, and Conn.D displayed a profound sex effect, although   197

there cortical traits were not measured. This means there is a deeper, complex layer of   198

sex effect dependent upon cooperative environmental and genetical factors which   199

requires further work to fully comprehend.   200

Association analysis for microarchitectural traits highlights 5 QTLs   201

We first measured statistical association between each trait and the founder haplotype 202

at each locus in the genome. Association analyses of the cortical and trabecular traits to 203

the haplotypes segregating in the CC (as defined by the ~70 K MegaMuga SNPs) 204

yielded 5 distinct QTLs. For BV/TV and Tb.N we recognized a marked peak at a locus 205

of length ~0.45 Mb between 116.5 and 116.9 Mb on chromosome 11, with peak logP 206

values of 7.6 and 6.8, which extended above the 99[th] percentile permutation-threshold 207

by 2.7 and 1.94 logP units, respectively. In Tb.Th, Tb.Sp, Ct.Th, and vBMD we 208

identified different QTLs on chromosomes 4, 5, 4 and 3, with logPs of 8.0, 9.4, 8.2, and 209

9.8, respectively, above threshold (Fig. 3 and Table 2). To account for false positive 210

results, we kept the false discovery rate (FDR) at 1% for each scan, by employing the 211

Benjamini–Hochberg multiple testing procedure on the logPs of the haplotype 212

associations to the traits. Conn.D and SMI lacked significant peaks above the stringent 213

permutated threshold and thus were not further analyzed (Fig. S2), but Conn.D 214

displayed a borderline peak in a region that matches the peak identified for BV/TV and 215

Tb.N. The 5 QTLs we describe are hereafter referred to as *Trl* (trabecular related locus) 216

7-9, and *Crl* (cortical related locus) 1-2 (respectively for BV/TV and Tb.N, Tb.Th, 217

Tb.Sp, Ct.Th, and vBMD, and in keeping with our previous report [15] that introduced 218

Trl 1-6). The 95% widths of the confidence intervals ranged from 6.4 to 15.6 Mb for 219

the *Trl*s, and were between 8.5-10.8 Mb for the *Crl*s (Table 2 and Fig. S3). 220

We measured the contribution of each CC founder to the QTLs, relative to the 221

wild-derived strain WSB/EiJ (Fig. 4). *Trl*7 is mostly affected by the classic laboratory 222

strains 129S1/SvImJ, NOD/LtJ, and NZO/HiLtJ; notably, the other traits were more 223

strongly driven by the following wild strains: *Trl*8 and *Crl*2 by PWK/PhJ; *Trl*9 by 224

WSB/EiJ; and *Crl*1 by CAST/Ei. 225

For *Trl*7, at the SNP most adjacent to the QTL peak UNC20471277, we found 226

that the majority of lines with a TT allelic variant (where T refers to the nucleic acid 227

Thymine) mostly congregate at the higher end of the BV/TV and Tb.N values (Mean 228

BV/TV = 17%); lines with a CC variant (where C refers to the nucleic acid Cytosine) 229

are at the lower end (mean BV/TV = 10%); and those with a CT variant are at the 230

intermediate range (Fig. 5). Largely, the more the trait examined is distantly correlated 231

with BV/TV, the less differentiated the CC and TT variants are, at the SNP 232

UNC20471277. This is accentuated in vBMD where there is a weak correlation with 233

BV/TV (Table S1) and leveled CC and TT groups (*P* value = 0.8 Welch's two sample 234

t-test). 235

Candidate genes identified by merge analysis and RNA-seq 237

To identify the gene most likely driving the skeletal trait, we next performed a merge 238

analysis and RNA-seq analysis and used a scoring system to rank the potential 239

candidates. 240

Merge analysis uses the catalogue of variants segregating in the eight CC 241

founders to impute the genotype dosage of each SNP in each CC line, based on the 242

haplotype reconstruction used for haplotype association [24]. Candidate causal 243

variants, if they exist, would be expected to be more significant (have higher logP 244

values) than the haplotype-based test in the flanking region. We found that *Trl*7 had the 245

highest density of polymorphisms (grey and crimson dots in Fig. 6) with merge-logP 246

values above the haplotype logPs (continuous black line in Fig. 6), while *Trl*8 and *Crl*2 247

had very few. The two latter loci congregated more upstream, in accordance with the 248

left-skewness of their respective CI simulations (Fig. S3). By calculating the relative   249

density of merge logP values which are considerably higher than the haplotype merge   250

logPs -and above the 99[th] percentile threshold for each scan - at intervals defined by   251

each gene within the QTLs (in meaningful regions; usually between the 50[th] and 90[th]   252

CI percentile) we devised a gene ranking scale (Table S4). For example, while the   253

proportion of merge analysis SNPs for BV/TV and Tb.N with logPs greater than that   254

of the haplotype scan is 1.4% at the genome-wide scale (as well as at the region   255

spanning the 95% CI, between ~112 − 118 Mb), it is 9.4% and ranked 5/36 (for BV/TV)   256

or 14.07% and ranked 3/36 (for Tb.N) at the region in which the gene *Rhbdf2* is situated   257

(~116.5 − 116.6 Mb) (See Table S4 and further discussion below).   258

To strengthen the criteria that classifies potential putative candidate gene as true   259

positives, we analyzed RNA-seq datasets of osteoclasts (Fig. S4) and osteocytes (Fig.   260

S5) made publicly available by Kim *et al* [25] (Gene Expression Omnibus accession   261

number GSE72846) and St John *et al* [26] (Gene Expression Omnibus accession   262

number GSE54784). We focused on local maximas that span ~0.5 Mb in and around   263

the peaks suggested by the merge analysis, for each QTL. From the raw count reads we   264

found that *Trl7* had the strongest gene expression differential; e.g while *Mxra7* in the   265

osteoclasts was expressed to a negligible degree (Fig. S4A), it had a strong presence in   266

osteocytes (Fig. S5A), whereas the genes of the other loci had much less prominent   267

differences. This suggests that genes at Trl7 are differentially expressed between   268

osteocytes and osteoclasts more prominently than in the other loci.   269

For each of the genes with the highest merge analysis scores in *Trl7*, we   270

attributed an *osteoclast* and *osteocyte* RNA-seq ($S_{RNA-seq1}$ and $S_{RNA-seq2}$, respectively)   271

based on the following formula: $S_{RNA-seq} = 1 - \frac{r}{n}$; where r is the local gene rank, n is   272

the total number of genes at the locus where if n>10 then n=10. For example, *Ube2o* 273

had an *osteoclast* RNA-seq score of 0.8 (ranked 2nd out of n>10 genes) and *osteocyte* 274

RNA-seq score of 0.8. *Rhbdf2* had an *osteoclast* RNA-seq score of 0.9 (ranked 1st) and 275

osteocyte RNA-seq score of 0.7. 276

We then summed up the cumulative *MS* (**M**erge, **S**equencing) score for each 277

gene defined as $MS = ln(merge) + (S_{RNA-seq1} + S_{RNA-seq2})$. Our analytical approach and 278

ranking strategy enabled us to shortlist the most plausible causal genes at the QTLs. In 279

Tlr7, *Ube2o* had an $MS_{Ube2o} = 2.4 + (0.8 + 0.8) = 4.0$; while *Aanat* scored $MS_{Aanat} = 2.7$ 280

$+ (0 + 0) = 2.7$, and *Rhbdf2* scored $MS_{Rhbdf2} = 2.6 + (0.9 + 0.7) = 4.2$. In the other loci, 281

the highest ranked genes were *Klf17* and *Kdm4a* for *Trl8*; *Barhl2* and *Zfp644* for *Trl9*; 282

Asph and *Gdf6* for *Crl1*; and *Hfe2*, *Acp6*, *Bcl9*, and *Notch2* for *Crl2*. 283

Because BV/TV is a predominant parameter in bone biology, we first focused 284

on *Trl7*. *Rhbdf2* had a merge strength of 14% (the 3rd strongest at the QTL and 2nd at 285

the 50% CI), and a local maxima at the RNA-seq of the osteoclasts. It was located near 286

the haplotype mapping peak, and because it received the highest cumulative *MS* score, 287

Rhbdf2 was retained for validation. The comprehensive list of the genes under the 50, 288

90, and 95% CI of the QTL, is supplied in table S4. 289

Validation of the skeletal role of *Rhbdf2* in knock-out mice 290

Femora of male mice (n=14) null at *Rhbdf2* (on a C57Bl/6J background) were collected, 291

on which we measured the same morphometric traits as above, including BV/TV, Tb.N, 292

Tb.Th, SMI, Tb.SP, Ct.Th, and vBMD. These were compared to their wild-type (WT) 293

counterparts (n=13), after adjusting for batch, age and weight. 294

Strikingly, we found that *Rhbdf2$^{-/-}$* mice had a significant bone phenotype. In 295

line with our GWAS data, *Rhbdf2$^{-/-}$* mice displayed a highly significant increase in 296

BV/TV and Tb.N (Fig. 7, 8). As expected, *Rhbdf2* KO also affected other 297

microstructural parameters, partly due to the high correlation between the trabecular 298

traits. After adjusting for confounders, we observed a significant difference between 299

KO and WT animals in Tb.Sp (P value < 0.001; uncorrected P value = 0.017), SMI (P 300

value = 0.03; uncorrected P value = 0.156) and Conn.D (P value = 0.008; uncorrected 301

P value = 0.046). Tb.Th and vBMD were not affected by the knockout. Although the 302

cortical compartment did not display a haplotype peak at the vicinity of *Trl*7 in the CC 303

animals. However, after adjusting for confounders, we observed a significant difference 304

in Ct.Th between KO and WT bones (P value = 0. 01), suggesting that the role of *Rhbdf2* 305

is not limited to the trabecular compartment (Fig. 7). 306

Discussion 307

Genetic reference population (GRP) are very efficient for the study of complex traits 308

and biological systems, because (i) genotyping is only required once ("genotype once, 309

phenotype many times", see below), and (ii) replicate individuals with the same 310

genotype can be generated at will allowing for optimal experimental designs [27]. 311

This article is the second to present the results of an ongoing quest to delineate 312

the genetic determinants that govern microstructural bone traits [15]. Here we 313

characterize several key microstructural properties of the mouse femoral bone to assess 314

the extent to which they are heritable; to what environmental perturbations they are 315

prone; and to identify candidate genes by which they are controlled. Our approach 316

narrowed down a small number of putative candidate genes for 6 of the 8 examined 317

phenotypes. Following merge analysis and RNA-seq, we validated our leading 318

candidate gene, *Rhbdf2*, using a knockout model, which confirmed the critical role of   319

Rhbdf2 on bone mass accrual and homeostasis.   320

While the heritability rates assessed here - determined to be over 60% for all   321

traits - confirmed our previous findings, the degree to which sex explains the   322

phenotypic variation was very subtle, and appeared only for the cortical traits; this   323

discrepancy may be due to the specific cohort composition used in this study (Table   324

S3), which includes a sex bias due to smaller number of females than males. We found   325

a total of five QTLs in six traits; BV/TV and Tb.N shared one QTL, and Tb.Th, Tb.Sp,   326

vBMD, and Ct.Th yielded one each. Importantly, although bone microarchitecture   327

factors are complex traits, our analyses highlighted no more than two loci for each trait;   328

it is likely that analyzing a larger number of CC lines would result in the identification   329

of further loci.   330

Our analyses yielded three QTLs for the trabecular traits and 2 QTLs for the   331

cortical traits. These are referred to as *Trl*7-9, and *Crl*1-2, respectively. In addition, and   332

in close proximity, to *Rhbdf2* (Rhomboid 5 Homolog 2; elaborated below), *Trl*7   333

includes *Ube2o* (Ubiquitin Conjugating Enzyme E2 O), which encodes an enzyme that   334

is an important interactant of SMAD6. Ube2o monoubiquitinates SMAD6, and thereby   335

facilitates the latter to bind BMP 1 receptors [28]. The signal transduction of BMP 1 is   336

in turn limited [29,30], and endochondral bone formation, instead of ossification, is   337

favored. Importantly, 4 week-old SMAD6-overexpressed mice have significantly lower   338

humeral and vertebral BV/TV ratios than their controls [29].   339

At *Trl*8, *Klf17, Kdm4a*, and *Dmap1* are likely putative candidate genes. Since   340

Klf17 (Kruppel-Like Factor 17) is part of a network that includes BMPs [31] it is more   341

likely than a nearby gene, *St3gal3* (ST3 Beta-Galactoside Alpha-2,3-Sialyltransferase   342

3), to affect bone traits, although the latter has a greater merge strength. *Kdm4a* (Lysine Demethylase 4A) encodes a histone demethylase that promotes the differentiation of embryonal stem cells (ESCs) to an endothelial fate [32]; endothelial cells are implied in regulation of bone formation [33]. *Dmap1* (DNA Methyltransferase 1 Associated Protein) which encodes a DNA-methyl transferase known to regulate obesity complications, and is differentially methylated in women with polycystic ovary syndrome [34,35] had the highest meaningful merge density, and it might epigenetically regulate bone formation as well. *Trl*9 includes two genes of interest to bone biology: *Barhl2* and *Zfp644*. By interacting with caspase3, which is essential for ossification [36], *Barhl2* (BarH Like Homeobox 2) can inhibit β-catenin activation [37], and regulate the expression of chordin, a BMP signaling-detrimental protein [38]. *Zfp644* (Zinc Finger Protein 644), which encodes a transcription repressor zinc-finger protein, is upregulated in eight week-old ovariectomized mice following treatment with estradiol [39], a treatment associated with reduced bone loss [40]. Further support for the candidacy of *Barhl2* and *Zfp644* is given by the role of *Barhl2* in the development of amacrine cells [41,42] and the association of *Zfp644* with myopia [43], a condition speculated to propagate from amacrine cell signaling [44]; interestingly myopia was linked to reduced postnatal bone mineral content in humans [45] and decreased expression of BMP 2 and 5 in guinea pigs [46].

The first of two cortical loci, *Crl*1 contains as likely candidates the genes *Asph* and *Gdf6*. *Asph* (Aspartate Beta-Hydroxylase) encodes a protein that has a role in regulating calcium homeostasis, which may affect bone metabolism [47]. *Gdf6* (Growth Differentiation Factor 6) is bone morphogenetic protein 13: mice with mutated *Gdf6* exhibit deformed bone formation in various skeletal sites; it is among the earliest known markers of limb joint formation [48], expressed in joints of ankle and knee. In *Gdf6*

homozygous mutant mice, bones fuse at the joints early at the segmentation stage [49]. 368

For *Crl2*, we found *Hfe2*, *Bcl9*, *Notch2,* and *Prkab2* as potential candidate genes. *Hfe2* 369
(Hemochromatosis Type 2 (Juvenile)) encodes the BMP co-receptor hemojuvelin 370
which is expressed in skeletal muscles [50] and is responsible for juvenile 371
hemochromatosis, a condition linked to sex hormones depletion and osteoporosis [51]. 372
Bcl9 (B-Cell CLL/Lymphoma 9), the mammalian ortholog of the gene *Legless*, encodes 373
a protein essential to the Wnt/beta-catenin signaling which is important for bone 374
metabolism [52], without which the nuclear localization of β-catenin and myocyte 375
differentiation are compromised [53]. Of note, there are mutual effects between bone 376
and muscle, and accumulating evidence suggest many genes show pleiotropism with 377
respect to muscle strength and bone parameters [54]. *Notch2* encodes a member of the 378
notch protein family, which influence both osteoblasts and osteoclasts [55]; 379
specifically, *Notch2* is associated with the rare Hajdu-Cheney syndrome, that includes 380
severe osteoporosis as one of its main symptoms [56,57]. For this gene, we did not find 381
any significant merge logPs included within its limits. Interestingly, *Sec22b*, an 382
adjacent gene, had the strongest merge logP marks in this locus but no documented link 383
to bone biology. The third-strong gene in terms of merge values was *Prkab2* (Protein 384
Kinase AMP-Activated Non-Catalytic Subunit Beta 2). It encodes an enzyme which is 385
the regulatory subunit of mitogen-activated protein kinase (AMPK). AMPK widely 386
affects bone metabolism [58]. 387

Rhbdf2 is not yet supported by peer-reviewed reports as bearing a relation to 388
bone. Based on its closeness to the *Tlr7* peak (within the 50% CI), its merge strength 389
and RNA-seq in bone cells, we identified *Rhbdf2* as a likely causal gene associated with 390
BV/TV and Tb.N. We therefore analyzed the bone phenotype of *Rhbdf2^{-/-}* mice, to 391
validate the role of this gene in the modeling of the femoral cortex and trabeculae. 392

Rhbdf2 deletion affected all the examined trabecular traits as well as Ct.Th. While the   393

effects on BV/TV and Tb.N were in line with the haplotype mapping, the *Rhbdf2* locus   394

did not appear in any of the other traits. This however is expected, because the genetic   395

architecture of the working cohort is such that the assumed contributing variant of   396

Rhbdf2 is diluted and compensated, resulting in a QTL detected only for the most   397

affected traits. Noticeably, Tb.Sp differed greatly between the *Rhbdf2$^{-/-}$* and control   398

mice but did not show up at the haplotype mapping; this might be due to (i) the great   399

diversity of the wild-type mice in Tb.Sp, and/or (ii) the need for complete knockout   400

rather than a mere SNP to detect significant changes in Tb.Sp, and/or (iii) the SNPs   401

giving rise to *Trl7* are functioning variants, with differential behavior affecting only   402

BV/TV and Tb.N. A similar interpretation may be valid for the cortical phenotype of   403

the *Rhbdf2$^{-/-}$* mice. Importantly, the significant QTL peak we found in our GWAS for   404

BV/TV and Tb.N ended up revealing a gene that has an important skeletal function in   405

both the trabecular and cortical bone compartments.   406

Rhbdf2 encodes the iRhom2 protein, a polytopic membrane protein that is a   407

catalytically inactive member of the rhomboid intramembrane serine proteases   408

superfamily [59]. iRhom2 is necessary in macrophages for the maturation and release   409

of the inflammatory cytokine tumor necrosis factor α (TNFα): it acts in the trafficking   410

of TACE, the protease that releases active TNFα from its membrane-tethered precursor   411

[60,61]. iRhom2 is also implicated in EGF-family growth factor signaling [62–64].   412

With a recent report of its role in trafficking of another protein, STING, it appears that   413

iRhom2 may have a wider role in regulating membrane trafficking [65]. iRhom2 was   414

also implicated in the regulation of CSF1R (macrophage stimulating factor 1 receptor),   415

a critical regulator of osteoclasts differentiation and survival [61,66–68]. *In vivo,*   416

Rhbdf2 has been implicated in esophageal cancer, wound healing, bone marrow 417

repopulation by monocytic cells, and inflammatory arthritis [63,69–71]. 418

Further work will be needed to identify the mechanism by which iRhom2 419

controls bone homeostasis; a possible direction could involve a positive feedback loop 420

that leads to differentiation of macrophages to osteoclasts. Indeed, iRhom2 stimulates 421

the secretion of TNFα by macrophages [68,72]; hyperactivates EGFR [73,74]; and 422

regulates CSF1R [75,76]. Although *Rhbdf2* is expressed in both the osteocyte and 423

osteoclast lineages, one cannot rule out the possibility that this gene regulates bone 424

remodeling by virtue of its expression in non-skeletal cells. 425

In summary, our analyses disclose several putative genes, several of which are 426

newly linked to a role in bone biology. A confirmation of one such gene, *Rhbdf2*, 427

provides the first conclusive evidence for its effects on bone microstructure. 428

Importantly, this study is the first demonstration of a physiological role of *Rhbdf2*. This 429

finding prompts future investigations to elucidate the exact mechanism of action of 430

Rhbdf2 and its contribution to osteoporosis in humans. 431

Materials and Methods 432

<u>Mice</u> 433

Mice aged 10 to 13 weeks (male $n = 103$; female $n = 71$), from 34 different CC lines 434

(average of 5 mice per line) were used in this study. The mice were at inbreeding 435

generations of 11 to 37, which correspond to 80-99.9% genetic homozygosity, 436

respectively. The mice were bred and maintained at the small animal facility of the 437

Sackler Faculty of Medicine, Tel Aviv University (TAU), Israel. They were housed on 438

hardwood chip bedding in open-top cages, with food and distilled water available *ad* 439

libitum, in an identical controlled environment (temperature = 25 ± 2°C; 60% ≤   440

humidity ≤ 85%) and a 12-hour light/dark cycle. All experiments protocols were   441

approved by the Institutional Animal Care and Use Committee (IACUC M-13-014) at   442

TAU, which follows the NIH/USA animal care and use protocols. The *Rhbdf2* knock   443

out mice and their WT counterparts were bred and maintained at the University of   444

Oxford as approved by license PPL80/2584 of the UK Home Office.   445

Specimen collection   446

Mice were intraperitoneally euthanized with cervical dislocation performed   447

approximately one minute after breathing stops owing to 5% Isoflurane inhalation. The   448

Rhbdf2 knock out mice and their WT counterparts were euthanized by inhalation of a   449

rising concentration of carbon dioxide followed by dislocation of the neck. Left femora   450

were harvested and fixed for 24 hours in 4% paraformaldehyde solution, and then stored   451

in 70% ethanol.   452

μCT evaluation   453

Whole left femora from each mouse were examined as described previously [77] by a   454

μCT system (μCT 50, Scanco Medical AG, Switzerland). Briefly, scans were   455

performed at a 10-μm resolution in all three spatial dimensions. The mineralized tissues   456

were differentially segmented by a global thresholding procedure [78]. All   457

morphometric parameters were determined by a direct 3D approach [79]. Parameters   458

analyzed were determined in the metaphyseal trabecular bone, which included   459

trabecular bone volume fraction (BV/TV; %), trabecular thickness (Tb.Th; μm),   460

trabecular number (Tb.N; mm^{-1}), trabecular connectivity density (Conn.D; mm^{-3}),   461

trabecular structure model index (SMI), and trabecular separation (Tb.Sp; mm). Two   462

additional parameters are characteristics of the mid-shaft diaphysis section, and include   463

volumetric bone mineral density (vBMD; mgHA/cm^3 [mg Hydroxy-Apatite per cm^3]) 464

and cortical thickness (Ct.Th; mm). All parameters were generated and termed 465

according to the Guidelines for assessment of bone microstructure in rodents using 466

micro–computed tomography [80]. 467

Genotyping 468

A representative male mouse from each line was initially genotyped with a high mouse 469

diversity array (MDA), which consists of 620,000 SNPs (Durrant et al., 2011). After 470

about two intervals of 4 generations of inbreeding, all the CC lines were regenotyped 471

by mouse universal genotype array (MUGA, 7,500 markers) and finally with the 472

MegaMuga (77,800 markers) SNP array to confirm their genotype status [19]. The 473

founder-based mosaic of each CC line was reconstructed using a hidden Markov model 474

(HMM) in which the hidden states are the founder haplotypes and the observed states 475

are the CC lines, to produce a probability matrix of descent from each founder. This 476

matrix was then pruned to about 11,000 SNPs by averaging across a window of 20 477

consecutive markers for faster analyses and reduction of genotyping errors [81]. 478

Statistical analyses and data acquisition 479

All statistical analyses were performed with the statistical software R (R core 480

development team 2009), including the package happy.hbrem [82]. 481

Heritability and covariate effects. Broad-sense heritability (H^2) was obtained for each 482

trait by fitting the trait (the independent variable) to the CC line label in a linear 483

regression model that incorporates relevant covariates (sex, age, batch, month, season, 484

year, and experimenter). ANOVA test was used to compare a null model (in which all 485

dependent variables are set to 0) with linear models that fit the covariates and the CC 486

line labels to the examined trait. Practically, the difference between the residual sum of 487

squares (RSS; $\sum_1^n(\mu_i - \hat{\mu}_i)^2$) of the covariates model and that of the CC-line labels 488

can be seen as the net genetic contribution to the trait. Thus, this difference divided by 489

that of the covariate model gives an estimation of the heritability. Each covariate was 490

calculated separately, by dividing the RSS difference between the null and full model 491

with that of the null model. Let F_0 be the model that fits the trait to the covariates; F_1 492

the model that fits the trait to the covariates and the CC line label; and F_{00} the null 493

model. Then, employing ANOVA, heritability is: 494

$$H^2 = (RSS(F_0) - RSS(F_1)) / RSS(F_{00}).$$ 495

Similarly, the effects for each covariate were computed separately, by fitting each in 496

F_0. The covariate effect is thus: 497

$$(RSS(F_{00}) - RSS(F_1)) / RSS(F_{00}).$$ 498

H^2n was derived from H^2 according to Atamni *et al* [23]. 499

Haplotype mapping. Each trait was fitted in a multiple linear regression model to the 500

probability matrix of descent from each founder, including sex and age as covariates. 501

The expected trait value from two ancestors, termed the genetic fit, is: 502

$$\mu_i = \mu + \sum_{s,t} F_{Li}(s,t)(\beta_s + \beta_t) = \mu + \sum_s \sum_t F_{Li}(s,t)\beta_s$$ 503

where μ is a normally distributed trait mean, with sex and age incorporated; $F_{Li}(s,t)$ is 504

the probability of descent from founders s and t; and $\beta_s + \beta_t$ is the additive effect of 505

founders s and t. Because $\sum_s \sum_t F_{Li}(s,t) = 2$ for a diploid organism, the maximum 506

likelihood estimates $\hat{\beta}_s$ are not independent. Thus, they are expressed here as 507

differences from the WSB/EiJ founder effect, so that $\hat{\beta}_{WSB} = 0$. Number of members 508

per line was weighted and integrated in the linear model. ANOVA was then used to compare this model with a null model where the founder effects are all set to 0; the resulting *F*-statistic yielded the significance of the genetic model *vs.* the null model and the negative 10-base logarithms of the P values (logP) were recorded.

Permutations of the CC lines between the phenotypes were used to set significance thresholds levels. Founder effects are the estimates derived from the multiple linear regression fit above.

Regional heritability (H_r^2) was hereafter computed by ANOVA as in the broad-sense heritability computation, except that here null linear regression fit was compared with a genetic linear regression fit with the probability matrix of the founder descent at the peak QTL as the explanatory variable.

False discovery rate (FDR) was calculated using the p.adjust function in R, with the method "BH" (Benjamini-Hochberg [83]).

Confidence intervals. Confidence intervals (CIs) were obtained both by simulations and by the quick method of Li, 2011 [84]. In the simulations, we resampled the residuals of the original linear regression fit at the peak of each QTL and rescanned 100 intervals within 7-10 Mb of the original loci to find the highest logP. Accordingly, following Durrant *et al.* [22], 1000 QTLs were simulated: if $\hat{t}_i$ is a random permutation of the residuals of fitted genetic model at the QTL peak, and K is a marker interval in a neighborhood of 3.5 to 5 Mb of the QTL peak L, a set of values for each trait, Z_{iK} is provided by:

$$Z_{iK} = \hat{t}_i \exp(\hat{\mu} + \sum_s X_{Kis}\hat{\beta}_s).$$

In the merge analysis the eight founder strains are partitioned and merged according to 532

the strain distribution pattern (SDP) of the alleles at the quantitative trait nucleotides 533

(QTN) within a given QTL (formerly obtained by the initial mapping). If we denote the 534

polymorphism as p, then $X_p = 1$ if s has allele a at p, and $X_p = 0$ otherwise [24]. Then, 535

at p, the probability of i to inherit alleles a and b from s and t, *respectively,* within L is 536

$$G_{pi}(a,b) = \sum_{s,t} X_p(a,s)X_p(b,t)F_{Li}(s,t).$$

This merges the founder strains by p. The expected trait value in the merged strains 538

can now be inferred by 539

$$\sum_{a,b} G_{pi}(a,b)(\beta_a + \beta_b).$$

Because this is a sub-model of the QTL model, it is expected to yield higher logP values 541

due to a reduction in the degrees of freedom. Significance was obtained by comparing 542

the merge model with the QTL model. Individual genes were extracted from the Sanger 543

mouse SNP repository (http://www.sanger.ac.uk/sanger/Mouse_SnpViewer). 544

Merge strength. We ranked the list of genes under each QTL according to the density 545

of merge logPs associated with them: only genes that had merge logPs above the 546

haplotype mapping reading, and above the threshold, plus logP=1 were included. We 547

then computed the relative density according to the density of a given gene's merge 548

logPs versus the locus' merge logP density. Let g be the region encompassed by a gene; 549

l the region encompassed by a QTL; and mp the merge logP values above the haplotype 550

P values plus 1. Then $g_i(mp) = 1$ if at SNP i there exists a mp and 0 otherwise. 551

Similarly $l_i(mp) = 1$ if at SNP j there exists a mp and 0 otherwise. The merge strength (MS) is therefore:

$$MS\ [\%] = 100 * \frac{\sum_i g_i(mp)}{\sum_j l_j(mp)}\ \%.$$

RNA-seq data

RNA-seq data from osteoclasts and osteocytes was obtained from gene expression omnibus (GEO) database (accession numbers GSE72846 and GSE54784) and mapped to the *mus musculus* assembly mm10 using tophat v. 2 [85]. Read counts were then casted on the loci of interest using the R (R Core Team 2015) package GenomicAlignments and raw read counts were taken. For the osteocytes, the data of basal level day 3 was averaged.

Acknowledgments

This study was supported by Tel Aviv University starter funds and by Israel Science Foundation (ISF) grant 1822/12 to YG, by Wellcome Trust grants 085906/Z/08/Z, 075491/Z/04, and 090532/Z/09/Z to RM, core funding by Tel-Aviv University to FI, Wellcome Trust grant 101035/Z/13/Z and the Medical Research Council (programme number U105178780) to MF, and by a fellowship from the Edmond J. Safra Center for Bioinformatics at Tel-Aviv University to RL.

References

1.   Burge R, Dawson-Hughes B, Solomon DH, Wong JB, King A, Tosteson A. Incidence and economic burden of osteoporosis-related fractures in the United States, 2005-2025. J Bone Miner Res. 2007;22: 465–75.

doi:10.1359/jbmr.061113 573

2. Dhanwal DK, Dennison EM, Harvey NC, Cooper C. Epidemiology of hip 574
fracture: Worldwide geographic variation. Indian J Orthop. 2011;45: 15–22. 575
doi:10.4103/0019-5413.73656 576

3. Center JR, Nguyen T V, Schneider D, Sambrook PN, Eisman JA. Mortality 577
after all major types of osteoporotic fracture in men and women: an 578
observational study. Lancet (London, England). 1999;353: 878–82. 579
doi:10.1016/S0140-6736(98)09075-8 580

4. Mizuguchi T, Furuta I, Watanabe Y, Tsukamoto K, Tomita H, Tsujihata M, et 581
al. LRP5, low-density-lipoprotein-receptor-related protein 5, is a determinant 582
for bone mineral density. J Hum Genet. 2004;49: 80–6. doi:10.1007/s10038- 583
003-0111-6 584

5. Richards JB, Rivadeneira F, Inouye M, Pastinen TM, Soranzo N, Wilson SG, et 585
al. Bone mineral density, osteoporosis, and osteoporotic fractures: a genome- 586
wide association study. Lancet. 2008;371: 1505–12. doi:10.1016/S0140- 587
6736(08)60599-1 588

6. Styrkarsdottir U, Halldorsson B V, Gretarsdottir S, Gudbjartsson DF, Walters 589
GB, Ingvarsson T, et al. Multiple genetic loci for bone mineral density and 590
fractures. N Engl J Med. 2008;358: 2355–65. doi:10.1056/NEJMoa0801197 591

7. Trikalinos TA, Salanti G, Zintzaras E, Ioannidis JPA. Meta-analysis methods. 592
Adv Genet. 2008;60: 311–34. doi:10.1016/S0065-2660(07)00413-0 593

8. Estrada K, Styrkarsdottir U, Evangelou E, Hsu Y-H, Duncan EL, Ntzani EE, et 594
al. Genome-wide meta-analysis identifies 56 bone mineral density loci and 595
reveals 14 loci associated with risk of fracture. Nat Genet. 2012;44: 491–501. 596

doi:10.1038/ng.2249   597

9.   Paternoster L, Lorentzon M, Vandenput L, Karlsson MK, Ljunggren O,   598

Kindmark A, et al. Genome-wide association meta-analysis of cortical bone   599

mineral density unravels allelic heterogeneity at the RANKL locus and   600

potential pleiotropic effects on bone. PLoS Genet. 2010;6: e1001217.   601

doi:10.1371/journal.pgen.1001217   602

10.   Paternoster L, Lorentzon M, Lehtimäki T, Eriksson J, Kähönen M, Raitakari O,   603

et al. Genetic determinants of trabecular and cortical volumetric bone mineral   604

densities and bone microstructure. PLoS Genet. 2013;9: e1003247.   605

doi:10.1371/journal.pgen.1003247   606

11.   Jovanovich A, Bùzková P, Chonchol M, Robbins J, Fink HA, de Boer IH, et al.   607

Fibroblast growth factor 23, bone mineral density, and risk of hip fracture   608

among older adults: the cardiovascular health study. J Clin Endocrinol Metab.   609

2013;98: 3323–31. doi:10.1210/jc.2013-1152   610

12.   Hsu Y-H, Kiel DP. Clinical review: Genome-wide association studies of   611

skeletal phenotypes: what we have learned and where we are headed. J Clin   612

Endocrinol Metab. 2012;97: E1958-77. doi:10.1210/jc.2012-1890   613

13.   Styrkarsdottir U, Thorleifsson G, Sulem P, Gudbjartsson DF, Sigurdsson A,   614

Jonasdottir A, et al. Nonsense mutation in the LGR4 gene is associated with   615

several human diseases and other traits. Nature. Nature Publishing Group, a   616

division of Macmillan Publishers Limited. All Rights Reserved.; 2013;497:   617

517–20. doi:10.1038/nature12124   618

14.   Iraqi F a, Athamni H, Dorman A, Salymah Y, Tomlinson I, Nashif A, et al.   619

Heritability and coefficient of genetic variation analyses of phenotypic traits   620

provide strong basis for high-resolution QTL mapping in the Collaborative   621

Cross mouse genetic reference population. Mamm Genome. 2014;25: 109–19.   622

doi:10.1007/s00335-014-9503-5   623

15. Levy R, Mott RF, Iraqi FA, Gabet Y. Collaborative cross mice in a genetic   624

association study reveal new candidate genes for bone microarchitecture. BMC   625

Genomics. 2015;16: 1013. doi:10.1186/s12864-015-2213-x   626

16. Karasik D, Demissie S, Zhou Y, Lu D, Broe KE, Bouxsein ML, et al.   627

Heritability and Genetic Correlations for Bone Microarchitecture: The   628

Framingham Study Families. J Bone Miner Res. 2016; doi:10.1002/jbmr.2915   629

17. Threadgill DW, Hunter KW, Williams RW. Genetic dissection of complex and   630

quantitative traits: from fantasy to reality via a community effort. Mamm   631

Genome. 2002;13: 175–8. doi:10.1007/s00335-001-4001-Y   632

18. Churchill G a, Airey DC, Allayee H, Angel JM, Attie AD, Beatty J, et al. The   633

Collaborative Cross, a community resource for the genetic analysis of complex   634

traits. Nat Genet. 2004;36: 1133–7. doi:10.1038/ng1104-1133   635

19. Collaborative Cross Consortium. The genome architecture of the Collaborative   636

Cross mouse genetic reference population. Genetics. 2012;190: 389–401.   637

doi:10.1534/genetics.111.132639   638

20. Keane TM, Goodstadt L, Danecek P, White M a, Wong K, Yalcin B, et al.   639

Mouse genomic variation and its effect on phenotypes and gene regulation.   640

Nature. 2011;477: 289–94. doi:10.1038/nature10413   641

21. Roberts A, Pardo-Manuel de Villena F, Wang W, McMillan L, Threadgill DW.   642

The polymorphism architecture of mouse genetic resources elucidated using   643

genome-wide resequencing data: implications for QTL discovery and systems   644

genetics. Mamm Genome. 2007;18: 473–81. doi:10.1007/s00335-007-9045-1   645

22.   Durrant C, Tayem H, Yalcin B, Cleak J, Goodstadt L, Villena FP De, et al.   646

        Collaborative Cross mice and their power to map host susceptibility to   647

        Aspergillus fumigatus infection Collaborative Cross mice and their power to   648

        map host susceptibility to Aspergillus fumigatus infection. 2011;   649

        doi:10.1101/gr.118786.110   650

23.   Atamni HJA-T, Mott R, Soller M, Iraqi FA. High-fat-diet induced development   651

        of increased fasting glucose levels and impaired response to intraperitoneal   652

        glucose challenge in the collaborative cross mouse genetic reference   653

        population. BMC Genet. BioMed Central; 2016;17: 10. doi:10.1186/s12863-   654

        015-0321-x   655

24.   Yalcin B, Flint J, Mott R. Using progenitor strain information to identify   656

        quantitative trait nucleotides in outbred mice. Genetics. 2005;171: 673–81.   657

        doi:10.1534/genetics.104.028902   658

25.   Kim K, Punj V, Kim J-M, Lee S, Ulmer TS, Lu W, et al. MMP-9 facilitates   659

        selective proteolysis of the histone H3 tail at genes necessary for proficient   660

        osteoclastogenesis. Genes Dev. 2016;30: 208–19. doi:10.1101/gad.268714.115   661

26.   St. John HC, Bishop KA, Meyer MB, Benkusky NA, Leng N, Kendziorski C,   662

        et al. The Osteoblast to Osteocyte Transition: Epigenetic Changes and   663

        Response to the Vitamin D $_3$ Hormone. Mol Endocrinol. 2014;28: 1150–1165.   664

        doi:10.1210/me.2014-1091   665

27.   Broman KW. The genomes of recombinant inbred lines. Genetics. 2005;169:   666

        1133–46. doi:10.1534/genetics.104.035212   667

28.   Zhang X, Zhang J, Bauer A, Zhang L, Selinger DW, Lu CX, et al. Fine-tuning   668

BMP7 signalling in adipogenesis by UBE2O/E2-230K-mediated 669

monoubiquitination of SMAD6. EMBO J. 2013;32: 996–1007. 670

doi:10.1038/emboj.2013.38 671

29. Horiki M, Imamura T, Okamoto M, Hayashi M, Murai J, Myoui A, et al. 672

Smad6/Smurf1 overexpression in cartilage delays chondrocyte hypertrophy and 673

causes dwarfism with osteopenia. J Cell Biol. 2004;165: 433–45. 674

doi:10.1083/jcb.200311015 675

30. Estrada KD, Retting KN, Chin AM, Lyons KM. Smad6 is essential to limit 676

BMP signaling during cartilage development. J Bone Miner Res. 2011;26: 677

2498–510. doi:10.1002/jbmr.443 678

31. Kotkamp K, Mössner R, Allen A, Onichtchouk D, Driever W. A Pou5f1/Oct4 679

dependent Klf2a, Klf2b, and Klf17 regulatory sub-network contributes to EVL 680

and ectoderm development during zebrafish embryogenesis. Dev Biol. 681

2014;385: 433–47. doi:10.1016/j.ydbio.2013.10.025 682

32. Wu L, Wary KK, Revskoy S, Gao X, Tsang K, Komarova YA, et al. Histone 683

Demethylases KDM4A and KDM4C Regulate Differentiation of Embryonic 684

Stem Cells to Endothelial Cells. Stem cell reports. 2015;5: 10–21. 685

doi:10.1016/j.stemcr.2015.05.016 686

33. Collin-Osdoby P. Role of vascular endothelial cells in bone biology. J Cell 687

Biochem. 1994;55: 304–9. doi:10.1002/jcb.240550306 688

34. Kokosar M, Benrick A, Perfilyev A, Fornes R, Nilsson E, Maliqueo M, et al. 689

Epigenetic and Transcriptional Alterations in Human Adipose Tissue of 690

Polycystic Ovary Syndrome. Sci Rep. 2016;6: 22883. doi:10.1038/srep22883 691

35. Kamei Y, Suganami T, Ehara T, Kanai S, Hayashi K, Yamamoto Y, et al. 692

Increased expression of DNA methyltransferase 3a in obese adipose tissue: 693
studies with transgenic mice. Obesity (Silver Spring). 2010;18: 314–21. 694
doi:10.1038/oby.2009.246 695

36. Miura M, Chen X-D, Allen MR, Bi Y, Gronthos S, Seo B-M, et al. A crucial 696
role of caspase-3 in osteogenic differentiation of bone marrow stromal stem 697
cells. J Clin Invest. 2004;114: 1704–13. doi:10.1172/JCI20427 698

37. Juraver-Geslin HA, Ausseil JJ, Wassef M, Durand BC. Barhl2 limits growth of 699
the diencephalic primordium through Caspase3 inhibition of beta-catenin 700
activation. Proc Natl Acad Sci U S A. 2011;108: 2288–93. 701
doi:10.1073/pnas.1014017108 702

38. Larraín J, Bachiller D, Lu B, Agius E, Piccolo S, De Robertis EM. BMP- 703
binding modules in chordin: a model for signalling regulation in the 704
extracellular space. Development. 2000;127: 821–30. Available: 705
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2280033&tool=pm 706
centrez&rendertype=abstract 707

39. Davis AM, Mao J, Naz B, Kohl JA, Rosenfeld CS. Comparative effects of 708
estradiol, methyl-piperidino-pyrazole, raloxifene, and ICI 182 780 on gene 709
expression in the murine uterus. J Mol Endocrinol. 2008;41: 205–17. 710
doi:10.1677/JME-08-0029 711

40. Kameda T, Mano H, Yuasa T, Mori Y, Miyazawa K, Shiokawa M, et al. 712
Estrogen inhibits bone resorption by directly inducing apoptosis of the bone- 713
resorbing osteoclasts. J Exp Med. 1997;186: 489–95. Available: 714
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2199029&tool=pm 715
centrez&rendertype=abstract 716

41. Ding Q, Chen H, Xie X, Libby RT, Tian N, Gan L. BARHL2 differentially regulates the development of retinal amacrine and ganglion neurons. J Neurosci. 2009;29: 3992–4003. doi:10.1523/JNEUROSCI.5237-08.2009

42. Mo Z, Li S, Yang X, Xiang M. Role of the Barhl2 homeobox gene in the specification of glycinergic amacrine cells. Development. 2004;131: 1607–18. doi:10.1242/dev.01071

43. Shi Y, Li Y, Zhang D, Zhang H, Li Y, Lu F, et al. Exome sequencing identifies ZNF644 mutations in high myopia. PLoS Genet. 2011;7: e1002084. doi:10.1371/journal.pgen.1002084

44. Chen JC, Brown B, Schmid KL. Evaluation of inner retinal function in myopia using oscillatory potentials of the multifocal electroretinogram. Vision Res. 2006;46: 4096–103. doi:10.1016/j.visres.2006.07.033

45. Pohlandt F. Hypothesis: myopia of prematurity is caused by postnatal bone mineral deficiency. Eur J Pediatr. 1994;153: 234–6. Available: http://www.ncbi.nlm.nih.gov/pubmed/8194552

46. Wang Q, Xue M-L, Zhao G-Q, Liu M-G, Ma Y-N, Ma Y. Form-deprivation myopia induces decreased expression of bone morphogenetic protein-2, 5 in guinea pig sclera. Int J Ophthalmol. 2015;8: 39–45. doi:10.3980/j.issn.2222-3959.2015.01.07

47. Pruitt KD, Brown GR, Hiatt SM, Thibaud-Nissen F, Astashyn A, Ermolaeva O, et al. RefSeq: an update on mammalian reference sequences. Nucleic Acids Res. 2014;42: D756-63. doi:10.1093/nar/gkt1114

48. Chang SC, Hoang B, Thomas JT, Vukicevic S, Luyten FP, Ryba NJ, et al. Cartilage-derived morphogenetic proteins. New members of the transforming

growth factor-beta superfamily predominantly expressed in long bones during
human embryonic development. J Biol Chem. 1994;269: 28227–34. Available:
http://www.ncbi.nlm.nih.gov/pubmed/7961761

49.   Settle SH, Rountree RB, Sinha A, Thacker A, Higgins K, Kingsley DM.
        Multiple joint and skeletal patterning defects caused by single and double
        mutations in the mouse Gdf6 and Gdf5 genes. Dev Biol. 2003;254: 116–30.
        Available: http://www.ncbi.nlm.nih.gov/pubmed/12606286

50.   Verga Falzacappa MV, Casanovas G, Hentze MW, Muckenthaler MU. A bone
        morphogenetic protein (BMP)-responsive element in the hepcidin promoter
        controls HFE2-mediated hepatic hepcidin expression and its response to IL-6 in
        cultured cells. J Mol Med (Berl). 2008;86: 531–40. doi:10.1007/s00109-008-
        0313-7

51.   Angelopoulos NG, Goula AK, Papanikolaou G, Tolis G. Osteoporosis in HFE2
        juvenile hemochromatosis. A case report and review of the literature.
        Osteoporos Int. 2006;17: 150–5. doi:10.1007/s00198-005-1920-6

52.   Baron R, Kneissel M. WNT signaling in bone homeostasis and disease: from
        human mutations to treatments. Nat Med. Nature Publishing Group, a division
        of Macmillan Publishers Limited. All Rights Reserved.; 2013;19: 179–92.
        doi:10.1038/nm.3074

53.   Brack AS, Murphy-Seiler F, Hanifi J, Deka J, Eyckerman S, Keller C, et al.
        BCL9 is an essential component of canonical Wnt signaling that mediates the
        differentiation of myogenic progenitors during muscle regeneration. Dev Biol.
        2009;335: 93–105. doi:10.1016/j.ydbio.2009.08.014

54.   Karasik D, Kiel DP. Genetics of the musculoskeletal system: a pleiotropic

approach. J Bone Miner Res. 2008;23: 788–802. doi:10.1359/jbmr.080218   765

55.   Bai S, Kopan R, Zou W, Hilton MJ, Ong C, Long F, et al. NOTCH1 regulates   766

       osteoclastogenesis directly in osteoclast precursors and indirectly via osteoblast   767

       lineage cells. J Biol Chem. 2008;283: 6509–18. doi:10.1074/jbc.M707000200   768

56.   Regan J, Long F. Notch signaling and bone remodeling. Curr Osteoporos Rep.   769

       2013;11: 126–9. doi:10.1007/s11914-013-0145-4   770

57.   Canalis E, Zanotti S. Hajdu-Cheney syndrome: a review. Orphanet J Rare Dis.   771

       BioMed Central; 2014;9: 200. doi:10.1186/s13023-014-0200-y   772

58.   Jeyabalan J, Shah M, Viollet B, Chenu C. AMP-activated protein kinase   773

       pathway and bone metabolism. J Endocrinol. 2012;212: 277–90.   774

       doi:10.1530/JOE-11-0306   775

59.   Lemberg MK, Freeman M. Functional and evolutionary implications of   776

       enhanced genomic analysis of rhomboid intramembrane proteases. Genome   777

       Res. 2007;17: 1634–1646. doi:10.1101/gr.6425307   778

60.   Adrain C, Zettl M, Christova Y, Taylor N, Freeman M, References A.   779

       Supporting Online Material for. 2012;225. doi:10.1126/science.1214400   780

61.   McIlwain DR, Lang PA, Maretzky T, Hamada K, Ohishi K, Maney SK, et al.   781

       iRhom2 Regulation of TACE Controls TNF-Mediated Protection Against   782

       Listeria and Responses to LPS. Science (80-). 2012;335: 229–232.   783

       doi:10.1126/science.1214448   784

62.   Siggs OM, Grieve A, Xu H, Bambrough P, Christova Y, Freeman M. Genetic   785

       interaction implicates iRhom2 in the regulation of EGF receptor signalling in   786

       mice. Biol Open. 2014;3.   787

63.   Hosur V, Johnson KR, Burzenski LM, Stearns TM, Maser RS, Shultz LD.   788

Rhbdf2 mutations increase its protein stability and drive EGFR hyperactivation   789

through enhanced secretion of amphiregulin. Proc Natl Acad Sci. 2014;111:   790

E2200–E2209. doi:10.1073/pnas.1323908111   791

64.   Li X, Maretzky T, Weskamp G, Monette S, Qing X, Issuree PDA, et al.   792

iRhoms 1 and 2 are essential upstream regulators of ADAM17-dependent   793

EGFR signaling. Proc Natl Acad Sci U S A. National Academy of Sciences;   794

2015;112: 6080–5. doi:10.1073/pnas.1505649112   795

65.   Luo W-W, Li S, Li C, Lian H, Yang Q, Zhong B, et al. iRhom2 is essential for   796

innate immunity to DNA viruses by mediating trafficking and stability of the   797

adaptor STING. Nat Immunol. 2016;17: 1057–1066. doi:10.1038/ni.3510   798

66.   Siggs OM, Xiao N, Wang Y, Shi H, Tomisato W, Li X, et al. iRhom2 is   799

required for the secretion of mouse TNFα. Blood. American Society of   800

Hematology; 2012;119: 5769–71. doi:10.1182/blood-2012-03-417949   801

67.   Adrain C, Zettl M, Christova Y, Taylor N, Freeman M. Tumor Necrosis Factor   802

Signaling Requires iRhom2 to Promote Trafficking and Activation of TACE.   803

Science (80-). 2012;335: 225–228. doi:10.1126/science.1214400   804

68.   Udagawa N, Takahashi N, Akatsu T, Tanaka H, Sasaki T, Nishihara T, et al.   805

Origin of osteoclasts: mature monocytes and macrophages are capable of   806

differentiating into osteoclasts under a suitable microenvironment prepared by   807

bone marrow-derived stromal cells. Proc Natl Acad Sci U S A. National   808

Academy of Sciences; 1990;87: 7260–4. Available: 809

http://www.ncbi.nlm.nih.gov/pubmed/2169622   810

69.   Issuree PDA, Maretzky T, McIlwain DR, Monette S, Qing X, Lang PA, et al.   811

iRHOM2 is a critical pathogenic mediator of inflammatory arthritis. J Clin   812

Invest. American Society for Clinical Investigation; 2013;123: 928–32.   813

doi:10.1172/JCI66168   814

70.   Qing X, D. Rogers L, Mortha A, Lavin Y, Redecha P, Issuree PD, et al.   815

iRhom2 regulates CSF1R cell surface expression and non-steady state   816

myelopoiesis in mice. Eur J Immunol. 2016;46: 2737–2748.   817

doi:10.1002/eji.201646482   818

71.   Blaydon DC, Etheridge SL, Risk JM, Hennies H-C, Gay LJ, Carroll R, et al.   819

RHBDF2 mutations are associated with tylosis, a familial esophageal cancer   820

syndrome. Am J Hum Genet. Elsevier; 2012;90: 340–6.   821

doi:10.1016/j.ajhg.2011.12.008   822

72.   Kobayashi K, Takahashi N, Jimi E, Udagawa N, Takami M, Kotake S, et al.   823

Tumor necrosis factor alpha stimulates osteoclast differentiation by a   824

mechanism independent of the ODF/RANKL-RANK interaction. J Exp Med.   825

2000;191: 275–86. Available: 826

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2195746&tool=pm   827

centrez&rendertype=abstract   828

73.   Yi T, Lee H-L, Cha J-H, Ko S-I, Kim H-J, Shin H-I, et al. Epidermal growth   829

factor receptor regulates osteoclast differentiation and survival through cross-   830

talking with RANK signaling. J Cell Physiol. 2008;217: 409–422.   831

doi:10.1002/jcp.21511   832

74.   Hosur V, Johnson KR, Burzenski LM, Stearns TM, Maser RS, Shultz LD.   833

Rhbdf2 mutations increase its protein stability and drive EGFR hyperactivation   834

through enhanced secretion of amphiregulin. Proc Natl Acad Sci. 2014;111:   835

E2200–E2209. doi:10.1073/pnas.1323908111   836

75. Hung JY, Horn D, Woodruff K, Prihoda T, LeSaux C, Peters J, et al. Colony-stimulating factor 1 potentiates lung cancer bone metastasis. Lab Investig. 2014;94: 371–381. doi:10.1038/labinvest.2014.1

76. Qing X, D. Rogers L, Mortha A, Lavin Y, Redecha P, Issuree PD, et al. iRhom2 regulates CSF1R cell surface expression and non-steady state myelopoiesis in mice. Eur J Immunol. 2016;46: 2737–2748. doi:10.1002/eji.201646482

77. Hiram-Bab S, Liron T, Deshet-Unger N, Mittelman M, Gassmann M, Rauner M, et al. Erythropoietin directly stimulates osteoclast precursors and induces bone loss. FASEB J. 2015;29: 1890–900. doi:10.1096/fj.14-259085

78. Rüegsegger P, Koller B, Müller R. A microtomographic system for the nondestructive evaluation of bone architecture. Calcif Tissue Int. 1996;58: 24–29. doi:10.1007/BF02509542

79. Hildebrand T, Laib A, Müller R, Dequeker J, Rüegsegger P. Direct three-dimensional morphometric analysis of human cancellous bone: microstructural data from spine, femur, iliac crest, and calcaneus. J Bone Miner Res. 1999;14: 1167–74. doi:10.1359/jbmr.1999.14.7.1167

80. Bouxsein ML, Boyd SK, Christiansen BA, Guldberg RE, Jepsen KJ, Müller R. Guidelines for assessment of bone microstructure in rodents using micro-computed tomography. J Bone Miner Res. 2010;25: 1468–86. doi:10.1002/jbmr.141

81. Hall M, Manship G, Morahan G, Pettit K, Scholten J, Tweedie K, et al. The genome architecture of the Collaborative Cross mouse genetic reference population. Genetics. 2012;190: 389–401. doi:10.1534/genetics.111.132639

82. Mott R, Talbot CJ, Turri MG, Collins a C, Flint J. A method for fine mapping   861

      quantitative trait loci in outbred animal stocks. Proc Natl Acad Sci U S A.   862

      2000;97: 12649–54. doi:10.1073/pnas.230304397   863

83. Hochberg Y. Controlling the False Discovery Rate : A Practical and Powerful   864

      Approach to Multiple Testing Author (s): Yoav Benjamini and Yosef   865

      Hochberg Source : Journal of the Royal Statistical Society . Series B (866

      Methodological), Vol . 57 , No . 1 (1995), Publi. 2016;57: 289–300.   867

84. Li H. A quick method to calculate QTL confidence interval. J Genet. 2011;90:   868

      355–60. Available: http://www.ncbi.nlm.nih.gov/pubmed/21869489   869

85. Durrant C, Tayem H, Yalcin B, Cleak J, Goodstadt L, de Villena FP-M, et al.   870

      Collaborative Cross mice and their power to map host susceptibility to   871

      Aspergillus fumigatus infection. Genome Res. 2011;21: 1239–48.   872

      doi:10.1101/gr.118786.110   873

86. Trapnell C, Pachter L, Salzberg SL. TopHat: discovering splice junctions with   874

      RNA-Seq. Bioinformatics. Oxford University Press; 2009;25: 1105–11.   875

      doi:10.1093/bioinformatics/btp120   876

Figure legends   877

Figure 1. µCT images of trabecular and cortical bone of the femora of representative 878

male CC mice. **(A)** Trabecular bone. Left: IL-2452, Right: IL-1513. **(B)** Cortical   879

bone. Left: IL-785, Right: IL-2689.   880

Figure 2 Trabecular and cortical traits distributions across the CC lines. X-axis is the 881

lines, y-axis is the trait means **A**. From top left, counter-clockwise: BV/TV (%), Tb.N 882

(mm^{-1}), Tb.Th (um), Conn.D (mm^{-3}), SMI, and Tb.Sp (mm). **B**. Left, vBMD   883

$(mgHA/cm^{3})$, right, Ct.Th (mm). Color codes group line(s) which significantly differ 884

from other groups. Lines are ordered inconsistently among the traits, per trait-specific   885

descending order. Refer to Table S3 for more details.   886

Figure 3 Haplotype association maps for the trabecular and cortical traits. X-axis is the   888

position on the chromosome, y-axis is the –logP value of the association. Lower   889

threshold represents the 95th percentile of 200 simulations, and top represents the 99th 890

percentile. Loci above the 99% cut-off were further investigated. From top to bottom: 891

BV/TV, Tb.N, Tb.Th, Tb.Sp, vBMD, and Ct.Th.   892

Figure 4. Ancestral effects relative to WSB. Y axis is the strain deviation relative to   895

WSB, x axis is the different strains of the eight CC founders. **(A)** to **(F)**: *Trl7* to *Trl9*,   896

Crl1, and *Crl2*, respectively.   897

Figure 5 Traits distribution at the marker UNC20471277 across bearers of   899

homozygous and heterozygous alleles, separated by sex. X-axis is the allelic variation   900

at the marker, y-axis is the trait value.   901

Figure 6 Merge analysis. Readings below logP = 4 are elided for brevity. X axis is   902

the position on the genome in Mb; y left axis is the logP score; y right axis is the   903

recombination rate scale; colored bars are genes (note that only strong putative   904

candidate genes are shown.); cyan line is the recombination rate; black continues line   905

is the haplotype test's peak; dashed line is the 99% permutation threshold. **(A)** to **(F)**:   906

Trl7 of BV/TV, Trl7 of Tb.N, Trl8 of Tb.Th, Trl9 of Tb.Sp, Crl1 of Ct.Th, and Crl2   907

of vBMD, respectively.   908

Figure 7 *Rhbdf2* knockout versus wildtype for each of the studied traits. Left is KO,   910

right is WT. PV is the confounder-adjusted P value. The unadjusted P value is in   911

brackets. PV = 0 means PV < 0.001.   912

Figure 8 μCT images of three-dimensional representative cortical and trabecular   914

bones reconstructions for *Rhbdf2* knockout and wildtype. Left: KO, right: WT. **(A)**   915

Trabecular bone. **(B)** Cortical bone. All samples were of male mice, aged 11 weeks.   916

Supporting information captions   917

Figure S1 *Trait distributions for Males and Females.* **A1, A2**: Trabecular and cortical   918

bone, respectively, males. **B1, B2**: trabecular and cortical bone, respectively, females.   919

X-axis is the lines, y-axis is the trait means **A**. From top left, counter-clockwise: 920

BV/TV (%), Tb.N (mm-1), Tb.Th (um), Conn.D (mm-3), SMI, and Tb.Sp (mm). **B**:   921

Left, vBMD (mgHA/cm3); right, Ct.Th (mm). Color codes group line(s) which   922

significantly differ from other groups. Lines are ordered inconsistently among the   923

traits, according to trait-specific descending order.   924

Figure S2 *Haplotype association maps for the trabecular and cortical traits.* X-axis   926

is the position on the chromosome, y-axis is the –logP value of the association. Lower   927

threshold represents the 95th percentile of 200 simulations, and top represents the 9th   928

percentile. Loci above the 99% cut-off were further investigated. From top to bottom: 929

Conn.D, SMI. 930

Figure S3 *Confidence interval simulations.* Loci at a neighborhood of 3-5 Mb around 931
the original locus were simulated by permuting the residual sum of squares of the 932
related phenotype. Maximum logP was obtained along with its relative position in Mb 933
to the original QTL (histograms, left panels), and with the number of markers from 934
the original QTL (boxplots, right panels). (*A*) and (*B*) show simulations results for the 935
BV/TV and Tb.N loci. These determined with high confidence that the peak QTL is 936
responsible for the effect seen in the haplotype scan, thus the narrow CI; (*C*) to (*F)* 937
show simulation results for Tb.Th, Tb.Sp, Ct.Th, and vBMD, 938
respectively; note the narrow CI for *Crl2* (vBMD), wide for *Trl8* (Tb.Th), and wider 939
still for *Crl1*(Ct.Th). 940

Figure S4 *RNA-seq of osteoclasts.* Gene names are on the right of each plot. Green 941
represents plus-stranded genes, black represents minus-stranded genes. Y-axis is the 942
raw expression count, where the negative scale refers to minus-stranded gene count. 943
Each bracket corresponds to a particular gene; left-to-right green (black) brackets fit 944
green (black) top-to-bottom gene-names. 945

Figure S5 *RNA-seq of osteocytes.* Gene names are on the right of each plot. Orange 946
represents plus-stranded genes, black represents minus-stranded genes. Y-axis is the 947
raw expression count, where the negative scale refers to minus-stranded gene count. 948
Each bracket corresponds to a particular gene; left-to-right orange (black) brackets fit 949
orange (black) top-to-bottom gene names 950

Table S1 *Pearson's correlations between measured μCT traits.* Pairwise correlations 951
for each trait are given as Pearson's *r*. In bold are the 5 most strong correlations. 952

Table S2 *Covariates effects. LogP is the negative logarithm of P value*. Effects of   953
covariates (i.e., the degree to which each covariate explains the phenotypic   954
difference). Values were determined by regressing the covariates along with the CC   955
lines, and running an ANOVA test. Note the covariates prominent effect on the   956
cortical traits. The batch effect was the strongest, affecting Tb.Th, Ct.Th, and vBMD.   957

Table S3 *Trait values per CC line included in this study (for both sexes combined).*   959
Trait means for each line, including number of members and standard error (SE). NA   960
means data were not available.   961

Table S4 *A comprehensive list of all genes under the 95% confidence-interval (CI)*   963
for the QTLs identified in this study. Light blue is the 95% CI region; blue is the 90%   964
CI region; and pink is the 50% CI region. Merge strength column gives the   965
normalized dosage of merge logP values as outlined in the methods section. Gene   966
symbols in bold type are discussed in the manuscript**.** Genes under the QTLs for the   967
cortical and trabecular traits. Light blue is the 95% CI, blue is the 90% CI, and red is   968
the 50% CI. "Merge Strength" refers to the proportion of merge logP values   969
constricted to the region of the specific gene. Note that for T*rl7* which is common to   970
BV/TV and Tb.N, the average values between the two are provided.   971

Fig.2A

Fig.2B

Fig. 4

Fig. 5

BV/TV

PV < 0.001

Tb.N

PV < 0.001

Tb.Th

PV < 0.001

Conn.D

PV = 0.006

SMI

PV < 0.001

Tb.Sp

PV = 0.2

Ct.Th

PV = 0.07

vBMD

PV = 0.8

Fig. 6

BV/TV
PV = 0.002 (0.031)

Tb.N
PV = 0 (0.027)

Tb.Th
PV = 0.424 (0.222)

Conn.D
PV = 0.008 (0.046)

SMI
PV = 0.033 (0.156)

Tb.Sp
PV = 0 (0.017)

vBMD
PV = 0.157 (0.062)

Ct.Th
PV = 0.018 (0.807)

Supporting Tables

Click here to access/download
Supporting Information
SuppTables.xlsx

Raw Meta-Data Set

Click here to access/download
Supporting Information - Compressed/ZIP File Archive
supportingInfo.rar