Title 1

Genome-wide association study in Collaborative Cross mice reveals a role for 2

Rhbdf2 **in skeletal homeostasis** 3

Short title 4

GWAS in mice reveals a new skeletal role for *Rhbdf2* 5

Authors 6

Roei Levy[1,2], Clemence Levet[3], Keren Cohen[1], Matthew Freeman[3], Richard Mott[4], 7

Fuad Iraqi[5], Yankel Gabet[1] 8

Affiliations 9

[1] Department of Anatomy and Anthropology, [2] Department of Human Molecular 10

Genetics and Biochemistry, and [5] Department of Clinical Microbiology and 11

Immunology, Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv 69978, 12

Israel. [3] Dunn School of Pathology, South Parks Road, Oxford OX1 3RE, UK. [4] UCL 13

Genetics Institute, University College London, Gower St., London, WC1E 6BT, UK. 14

Keywords Bone; Collaborative Cross; Rhbdf2; GWAS; iRhom2; cancellous bone; 15
bone mass; causal genes; femur; RIL 16

Abstract   17

Osteoporosis, the most common bone disease, is characterized by a low bone mass and   18

increased risk of fractures. Importantly, individuals with the same bone mineral density   19

(BMD) measured on two dimensional (2D) radiographs, have different risks for   20

fracture, suggesting that microstructural architecture is an important determinant of   21

skeletal strength. Here we took advantage of the rich phenotypic and genetic diversity   22

of the Collaborative Cross (CC) mice. Using microcomputed tomography, we   23

examined key structural parameters in the femoral cortical and trabecular compartments   24

of male and female mice from 34 CC lines. These traits included the trabecular bone   25

volume fraction (BV/TV), trabecular number, thickness, connectivity, and spacing, as   26

well as structural morphometric index. In the mid-diaphyseal cortex, we recorded   27

cortical thickness and volumetric BMD.   28

The broad-sense heritability of these traits ranged between 50 to 60%. Our   29

genome-wide association approach revealed 5 quantitative traits loci (QTL)   30

significantly associated with 6 of the traits. We refined each locus by combining   31

information obtained from the known ancestry of the mice and RNA-Seq data from   32

existing databases, to shortlist potential candidate genes. We found strong evidence for   33

new candidate genes, including *Rhbdf2*, which association to BV/TV and Tb.N was   34

strongly suggested by our analysis. We then examined knockout mice, and validated   35

the causal action of *Rhbdf2* on bone mass accrual and microarchitecture.   36

Our approach revealed new genome-wide QTLs and a series of genes that have   37

never been associated with bone microarchitecture. This study demonstrates for the first   38

time the skeletal role of *Rhbdf2* on the physiological remodeling of both the cortical   39

and trabecular bone. This newly assigned function for *Rhbdf2* can prove useful in   40

Author summary   43

In this study, we used the novel mouse reference population, the Collaborative Cross   44

(CC), in a study aimed at identifying new causal genes in the regulation of bone   45

microarchitecture, a critical determinant of bone strength. Our approach provides a   46

clear advantage in terms of resolution and 3D measurements of the morphometric   47

features (versus humans) and rich allelic diversity (versus classical mouse   48

populations), over current practices of genome-wide ssociation studies of bones.   49

Our genome-wide study revealed 5 loci significantly associated with microstructural   50

traits in the cortical and trabecular bone. We found strong evidence for new candidate   51

genes, in particular, *Rhbdf2*. We then validated the specific role of *Rhbdf2* on bone   52

mass accrual and microarchitecture using knockout mice. Importantly, this study is   53

the first demonstration of a physiological role for *Rhbdf2*.   54

Introduction   56

Osteoporosis is the most common bone disease in humans, affecting nearly half the US   57

and European population over the age of 50 years. With the globally increasing life   58

expectancy, osteoporosis and related bone fractures are becoming a pandemic health   59

and economic concern. By 2050, the world-wide incidence of hip fractures is expected   60

to increase by 2.5 to 6 fold [1,2]. Importantly, the mortality rate in the 12 months   61

following bone fracture is as high as 20% [3]. Risk of fracture is determined largely by   62

bone density and quality/strength, which are the end result of peak values achieved at   63

skeletal maturity and subsequent age and menopause-related bone loss. Genetic factors   64

have a major role in determining the wide range in the so-called "normal" peak bone   65

mass. Measures of bone status are inherently complex traits, as opposed to Mendelian 66

traits; i.e. they are controlled by the cumulative effect and interactions of numerous 67

genetic loci and environmental factors. 68

Genome-wide association studies (GWAS), including a large meta-analysis, 69

have identified more than 50 loci associated with bone mineral density (BMD) [4–10]. 70

However, many other genes that were experimentally associated with bone mass were 71

not confirmed by GWAS in human cohorts [11–13]. This suggests that the BMD 72

phenotype does not capture the structural complexity of the bone; there may be other 73

relevant bone phenotypes not yet studied in human GWAS [12], which hitherto have 74

generally relied on areal bone mineral density (aBMD) as the sole bone feature. aBMD 75

measured by dual energy x-ray absorptiometry (DXA) is a two dimensional projection 76

that cannot measure bone size, individual bone compartments' shape (whether 77

trabecular or cortical) or underlying microstructure, and thus likely conceals important 78

features which are assumed to be controlled by unique genetic determinants. Indeed, 79

there is a growing body of evidence that argues for distinct genetic influences of the 80

cortical and trabecular bone and thus they should be accordingly distinguished [8,9]. A 81

recent GWAS in the Collaborative Cross (CC) mice based on DXA failed to find any 82

heritability of BMD [14], whereas another report based on the same mouse panel 83

showed highly significant heritability levels in most of the cortical and trabecular 84

microstructural parameters measured by micro-computed tomography (μCT)[15] 85

Traditional peripheral quantitative CT (pQCT) has the capacity to distinguish 86

between the cortical and trabecular bone compartments, but it lacks the required 87

resolution to detect microstructural differences. A recent report based on high 88

resolution pQCT (HR-pQCT) data in humans, identified two novel bone-related loci, 89

thus far undetected by DXA and pQCT-based GWAS [9]. Another [16], found strong

genetic correlations between 1047 adult participants of the Framingham heart study,

therefore indicating that the heritability of bone microstructure constitutes a phenotypic

layer which is at least partially independent of DXA-derived BMD. Like HR-pQCT

studies in humans, understanding the genetic regulation of bone microstructural

parameters using μCT in small animals is likely to identify genetic factors distinct from

those previously identified for DXA-derived traits.

The CC mouse panel is designed to provide high resolution analysis of complex

traits, with particular emphasis on traits relevant to human health [17,18]. This unique

resource currently consists of a growing number of recombinant inbred lines (RIL)

generated from full reciprocal breeding of eight divergent strains of mice [19]. In

contrast to commonly used laboratory mouse strains, the ancestry of the CC lines

includes wild-derived inbred strains that encompass genetic variations accumulated

over ~1 million years [20]; more than 50 million single nucleotide polymorphisms

segregate in founders of the CC. The high genetic diversity means that QTLs can be

mapped using this panel that would have been invisible in a population that involved

only classical strains [21,22].

This claim is substantiated in a recent study that identified a genome-wide

significant association between *Oxt* (oxytocin) and *Avp* (vasopressin) and skeletal

microarchitecture in CC mice (Levy et al. 2015). Here, our GWAS in the CC mouse

panel identified a novel gene, *Rhbdf2,* associated with bone traits and using a specific

knockout model we validated its role in the regulation of cortical and trabecular bone

structure. This exemplifies the effectiveness and relative ease by which a GWAS with

a small CC population can associate a bone-related function to novel genes, and to 113

reveal overlooked key players in skeletal biology. 114

Results 115

CC lines widely differ in bone microarchitecture traits 116

We examined the variation in femoral cortical and trabecular microstructure between 117

34 unique CC lines totaling in 174 mice (71 females and 103 males, with an average of 118

4.25 mice per line). In the trabecular bone compartment we measured bone volume 119

fraction (BV/TV), trabecular number (Tb.N), thickness (Tb.Th), connectivity 120

(Conn.D), and spacing (or separation; Tb.Sp), as well as structural morphometric index 121

(SMI) of the trabecular framework. In the mid-diaphyseal cortex, we recorded cortical 122

thickness (Ct.Th) and volumetric bone mineral density (vBMD). These traits were 123

approximately normally distributed; BV/TV ranged from 0.017 to 0.26 (i.e. 1.7% to 124

26%; mean = 10.2%); Tb.N from 0.52 to 6.11 mm^{-1} (mean = 2.7 mm^{-1}); Tb.Th from 31 125

to 69 μm (mean = 47 μm); Conn.D from 10.9 to 268.3 mm^{-3} (mean = 104.2 mm^{-3}); SMI 126

from 0.6 to 3.3 (mean = 2.3); Tb.Sp from 0.16 to 0.7 mm (mean = 0.33 mm); Ct.Th 127

from 0.14 to 0.29 mm (mean = 0.2 mm); and vBMD from 402.5 to 809.2 $mgHA/cm^3$ 128

(mean = 581.1 $mgHA/cm^3$). μCT images taken from two mice with distinct cortical 129

(Fig. 1A) and trabecular (Fig. 1B) characteristics demonstrate the great variation in 130

bone traits due solely to the genetic background. Color-codes on the graphs in Fig. 2 131

indicate Duncan's least significance range (LSR), which dictates whether the mean 132

value of a line, or a group of lines, for a given trait differs to a degree of at least *P*-value 133

< 0.001 from any other group. LSR allows for a visual representation of the 134

heterogeneity amongst the lines. 135

With 11 distinct groups, vBMD (Fig. 2B) is the most heterogeneous trait, while   136

SMI and Conn.D are the least, with only 3 significantly distinct groups (Fig. 2A).   137

Notably, the heterogeneity of females is greater than that of males for cortical traits but   138

milder for trabecular traits (Figs. S1A1, 2 for males and S1B1, 2 for females).   139

To examine the inter-dependency between the traits, we assessed the correlation   141

between all the measured parameters, in a pairwise fashion using Pearson's correlation   142

test. The strongest correlation was between BV/TV and Tb.N (Pearson's $r = 0.94$), in   143

line with our previous findings [15], while the weakest was between Tb.N and vBMD   144

($r < 0.01$). There was also a moderately high correlation between Ct.Th and Tb.Th ($r =$ 145

0.61; and see Table S1). The correlation between sexes for each trait (Table 1) ranged   146

from $r = 0.75$ (Tb.Sp) to $r = 0.20$ (Ct.Th). Body weight (range = 17.4 - 35.0 gr) did not   147

significantly correlate with any of the traits ($r = 0.01$ for Conn.D to $r = 0.19$ for Ct.Th;   148

data not shown). After separating males from females the correlation slightly increased,   149

yet remained low. Weak correlation was found between weight and Tb.N, SMI, and   150

Ct.Th for females (Pearson's $r = -0.20$, 0.23, and 0.25 respectively), and between weight   151

and Tb.Th and Tb.Sp (Pearson's $r = 0.25$ and -0.25) for males.   152

Table 1 Heritability, sex correlations, and covariate interactions for trabecular and cortical traits

Trait	H^2	logP	H^2n	Sex Cor.	Interactions %
BV/TV	0.61	12.43	0.87	0.6695754	-
Tb.N	0.63	13.76	0.88	0.7628464	-
Tb.Th	0.54	9.08	0.83	0.7039887	34.70
Conn.D	0.56	9.90	0.84	0.5188039	-
SMI	0.55	9.43	0.84	0.2984166	26.02
Tb.Sp	0.63	13.45	0.88	0.754563	-
vBMD	0.62	12.15	0.87	0.6268818	53.92
Ct.Th	0.51	7.63	0.82	0.2055684	41.07

H^2 is the broad-sense heritability (which includes epistatic and environmental influences); logP is the negative 10-base logarithm of the P-value; H^2n is the line-mean heritability; Sex Cor. Is the sex correlation of each trait; and interactions % refers to the relative contribution of the cumulative covariate-interactions, which include sex, age, batch, month, season, year, and experimenter (see table S2).

Table 2 Positions of QTLs associated with trabecular and cortical traits												
						Simulation						
				99th % thresh	H^2_r	50% CI (Mb)		90% CI (Mb)		95% CI (Mb)	
QTL	Trait	Chr	logP			Position	Width	Position	Width	Position	Width
Trl7	BV/TV	11	7.60	4.90	0.69	116.6-116.7	0.12	113.6-118.1	4.50	112.1-118.3	6.40
Trl7	Tb.N	11	6.80	4.84	0.71	116.6-116.87	0.29	114.2-118.35	4.15	112.41-118.68	6.27
Trl8	Tb.Th	4	8.00	4.50	0.61	117.2-117.58	0.32	113.05-125.54	12.49	110.87-126.52	15.65
Trl9	Tb.Sp	5	9.40	6.01	0.78	105.78-106.14	0.35	101.6-109.11	7.54	99.8-110.39	10.62
Crl1	Ct.Th	4	8.20	4.80	0.80	9.29-9.72	0.43	4.0-11.7	7.70	3.4-11.8	8.49
Crl2	vBMD	3	9.80	4.93	0.86	97.2-97.4	0.20	94.4-103.1	8.50	93.22-104.3	10.80
Chr = chromosome; logP = negative 10-base logarithm of P value; Sig = genome-wide significance level; 99th % threshold logP = threshold used to define cut-off for QTL peaks (Fig. 4); H^2_r = regional heritability (the proportion to which the locus explains the phenotypic variability). Positions and widths of the simulation-based 50, 90, and 95% CIs are given.											

While in most lines the traits' correlations were predictive of a given line's rank,   169

in others a less expected pattern was observed; e.g., IL-1513 displayed unusually   170

extreme phenotypes for all trabecular traits and was at the higher end for BV/TV, Tb.N,   171

Tb.Th, and Conn.D and at the lower end for SMI and Tb.Sp, but IL-188 was more   172

discordant between these same traits (Fig. 2 and Table S3), illustrating unexpected co-   173

variation of the traits in the CC.   174

Heritability and confounder-control   175

We quantitated the effects of the covariates sex, age, batch, month, season, year, and   176

experimenter on each trait. Age ranged from 9 (n=6) to 13 (n=9) weeks and the mice   177

were dissected in 20 batches over a three-year course across 8 months during winter, 178

spring and summer, by two experimenters. Whereas age alone had no effect on any 179

trait, sex affected only Ct.Th; batch affected Tb.Th, vBMD, and Ct.Th; month affected 180

Tb.Sp, vBMD, and Ct.Th; season and year affected vBMD and Ct.Th; and Tb.Th and 181

Ct.Th were affected by experimenter. The cumulative effect of the covariates' pairwise 182

interactions was noted for Tb.Th, SMI, vBMD, and Ct.Th. (Table S2). 183

We then estimated the broad-sense heritability (H^2) of each trait among the CC 184

lines, which includes additive and non-additive epistatic effects and gene-environment 185

interactions. The greatest H^2 is seen for Tb.N (0.63, logP = 13.76; where logP stands 186

for the negative 10-base logarithm of the P value and tests the null hypothesis that the 187

heritability is zero), and the smallest for Ct.Th (0.51, logP = 7.63). 188

We calculated the heritability for the mean values in each line to get a better 189

representation of the percentage of genetic contribution to the phenotypic heterogeneity 190

by incorporating H^2 and the average number of lines [23]. This defines H^2n, which is 191

directly proportional to H^2 (Methods; Table 1) and ranges between 82 (Ct.Th) and 88% 192

(Tb.N and Tb.Sp). 193

Overall the cortical traits seemed more prone to covariate variation; they were 194

particularly sensitive to sex, batch, and season. This stands in contrast to our previous 195

results [15] where BV/TV, Tb.N, and Conn.D displayed a profound sex effect, although 196

there cortical traits were not measured. This means there is a deeper, complex layer of 197

sex effect dependent upon cooperative environmental and genetical factors which 198

requires further work to fully comprehend. 199

Association analysis for microarchitectural traits highlights 5 QTLs 200

We first measured statistical association between each trait and the founder haplotype 201

at each locus in the genome. Association analyses of the cortical and trabecular traits to 202

the haplotypes segregating in the CC (as defined by the ~70 K MegaMuga SNPs) 203

yielded 5 distinct QTLs. For BV/TV and Tb.N we recognized a marked peak at a locus 204

of length ~0.45 Mb between 116.5 and 116.9 Mb on chromosome 11, with peak logP 205

values of 7.6 and 6.8, which extended above the 99[th] percentile permutation-threshold 206

by 2.7 and 1.94 logP units, respectively. In Tb.Th, Tb.Sp, Ct.Th, and vBMD we 207

identified different QTLs on chromosomes 4, 5, 4 and 3, with logPs of 8.0, 9.4, 8.2, and 208

9.8, respectively, above threshold (Fig. 3 and Table 2). To account for false positive 209

results, we kept the false discovery rate (FDR) at 1% for each scan, by employing the 210

Benjamini–Hochberg multiple testing procedure on the logPs of the haplotype 211

associations to the traits. Conn.D and SMI lacked significant peaks above the stringent 212

permutated threshold and thus were not further analyzed (Fig. S2), but Conn.D 213

displayed a borderline peak in a region that matches the peak identified for BV/TV and 214

Tb.N. The 5 QTLs we describe are hereafter referred to as *Trl* (trabecular related locus) 215

7-9, and *Crl* (cortical related locus) 1-2 (respectively for BV/TV and Tb.N, Tb.Th, 216

Tb.Sp, Ct.Th, and vBMD, and in keeping with our previous report [15] that introduced 217

Trl 1-6). The 95% widths of the confidence intervals ranged from 6.4 to 15.6 Mb for 218

the *Trl*s, and were between 8.5-10.8 Mb for the *Crl*s (Table 2 and Fig. S3). 219

We measured the contribution of each CC founder to the QTLs, relative to the 220

wild-derived strain WSB/EiJ (Fig. 4). *Trl*7 is mostly affected by the classic laboratory 221

strains 129S1/SvImJ, NOD/LtJ, and NZO/HiLtJ; notably, the other traits were more 222

strongly driven by the following wild strains: *Trl*8 and *Crl*2 by PWK/PhJ; *Trl*9 by 223

WSB/EiJ; and *Crl*1 by CAST/Ei. 224

For *Trl*7, at the SNP most adjacent to the QTL peak UNC20471277, we found 225

that the majority of lines with a TT allelic variant (where T refers to the nucleic acid 226

Thymine) mostly congregate at the higher end of the BV/TV and Tb.N values (Mean 227

BV/TV = 17%); lines with a CC variant (where C refers to the nucleic acid Cytosine) 228

are at the lower end (mean BV/TV = 10%); and those with a CT variant are at the 229

intermediate range (Fig. 5). Largely, the more the trait examined is distantly correlated 230

with BV/TV, the less differentiated the CC and TT variants are, at the SNP 231

UNC20471277. This is accentuated in vBMD where there is a weak correlation with 232

BV/TV (Table S1) and leveled CC and TT groups (*P* value = 0.8 Welch's two sample 233

t-test). 234

Candidate genes identified by merge analysis and RNA-seq 236

To identify the gene most likely driving the skeletal trait, we next performed a merge 237

analysis and RNA-seq analysis and used a scoring system to rank the potential 238

candidates. 239

Merge analysis uses the catalogue of variants segregating in the eight CC 240

founders to impute the genotype dosage of each SNP in each CC line, based on the 241

haplotype reconstruction used for haplotype association [24]. Candidate causal 242

variants, if they exist, would be expected to be more significant (have higher logP 243

values) than the haplotype-based test in the flanking region. We found that *Trl*7 had the 244

highest density of polymorphisms (grey and crimson dots) with merge-logP values 245

above the haplotype logPs (continuous black line), while *Trl*8 and *Crl*2 had very few. 246

The two latter loci congregated more upstream, in accordance with the left-skewness 247

of their respective CI simulations (Fig. S3). By calculating the relative density of merge   248

logP values which are considerably higher than the haplotype merge logPs -and above   249

the 99th percentile threshold for each scan - at intervals defined by each gene within the   250

QTLs (in meaningful regions; usually between the 50th and 90th CI percentile) we   251

devised a gene ranking scale (Table S4). For example, while the proportion of merge   252

analysis SNPs for BV/TV and Tb.N with logPs greater than that of the haplotype scan   253

is 1.4% at the genome-wide scale (as well as at the region spanning the 95% CI, between   254

~112 – 118 Mb), it is 9.4% and ranked 5/36 (for BV/TV) or 14.07% and ranked 3/36   255

(for Tb.N) at the region in which the gene *Rhbdf2* is situated (~116.5 – 116.6 Mb) (See   256

Table S4 and further discussion below).   257

To strengthen the criteria that classifies potential putative candidate gene as true   258

positives, we analyzed RNA-seq datasets of osteoclasts (Fig. S4) and osteocytes (Fig.   259

S5) made publicly available by Kim *et al* [25] (Gene Expression Omnibus accession   260

number GSE72846) and St John *et al* [26] (Gene Expression Omnibus accession   261

number GSE54784). We focused on local maximas that span ~0.5 Mb in and around   262

the peaks suggested by the merge analysis, for each QTL. From the raw count reads we   263

found that *Trl7* had the strongest gene expression differential; e.g while *Mxra7* in the   264

osteoclasts was expressed to a negligible degree (Fig. S4A), it had a strong presence in   265

osteocytes (Fig. S5A), whereas the genes of the other loci had much less prominent   266

differences. This suggests that genes at Trl7 are differentially expressed between   267

osteocytes and osteoclasts more prominently than in the other loci.   268

For each of the genes with the highest merge analysis scores in *Trl7*, we   269

attributed an *osteoclast* and *osteocyte* RNA-seq ($S_{RNA\text{-}seq1}$ and $S_{RNA\text{-}seq2}$, respectively)   270

based on the following formula: $S_{RNA\text{-}seq} = 1 - \frac{r}{n}$; where r is the local gene rank, n is   271

the total number of genes at the locus where if n>10 then n=10. For example, *Ube2o* had an *osteoclast* RNA-seq score of 0.8 (ranked 2[nd] out of n>10 genes) and *osteocyte* RNA-seq score of 0.8. *Rhbdf2* had an *osteoclast* RNA-seq score of 0.9 (ranked 1[st]) and *osteocyte* RNA-seq score of 0.7.

We then summed up the cumulative *MS* (**M**erge, **S**equencing) score for each gene defined as $MS = ln(merge) + (S_{RNA-seq1} + S_{RNA-seq2})$. Our analytical approach and ranking strategy enabled us to shortlist the most plausible causal genes at the QTLs. In *Tlr7*, *Ube2o* had an $MS_{Ube2o} = 2.4 + (0.8 + 0.8) = 4.0$; while *Aanat* scored $MS_{Aanat} = 2.7 + (0 + 0) = 2.7$, and *Rhbdf2* scored $MS_{Rhbdf2} = 2.6 + (0.9 + 0.7) = 4.2$. In the other loci, the highest ranked genes were *Klf17* and *Kdm4a* for *Trl8*; *Barhl2* and *Zfp644* for *Trl9*; *Asph* and *Gdf6* for *Crl1*; and *Hfe2*, *Acp6*, *Bcl9*, and *Notch2* for *Crl2*.

Because BV/TV is a predominant parameter in bone biology, we first focused on *Trl7*. *Rhbdf2* had a merge strength of 14% (the 3[rd] strongest at the QTL and 2[nd] at the 50% CI), and a local maxima at the RNA-seq of the osteoclasts. It was located near the haplotype mapping peak, and because it received the highest cumulative *MS* score, *Rhbdf2* was retained for validation. The comprehensive list of the genes under the 50, 90, and 95% CI of the QTL, is supplied in table S4.

Validation of the skeletal role of *Rhbdf2* in knock-out mice

Femora of male mice (n=14) null at *Rhbdf2* (on a C57Bl/6J background) were collected, on which we measured the same morphometric traits as above, including BV/TV, Tb.N, Tb.Th, SMI, Tb.SP, Ct.Th, and vBMD. These were compared to their wild-type (WT) counterparts (n=13), after adjusting for batch, age and weight.

Strikingly, we found that *Rhbdf2*[-/-] mice had a significant bone phenotype. In line with our GWAS data, *Rhbdf2*[-/-] mice displayed a highly significant increase in BV/TV and Tb.N (Fig. 7, 8). As expected, *Rhbdf2* KO also affected other microstructural parameters, partly due to the high correlation between the trabecular traits. After adjusting for confounders, we observed a significant difference between KO and WT animals in Tb.Sp (P value < 0.001; uncorrected P value = 0.017), SMI (P value = 0.03; uncorrected P value = 0.156) and Conn.D (P value = 0.008; uncorrected P value = 0.046). Tb.Th and vBMD were not affected by the knockout. Although the cortical compartment did not display a haplotype peak at the vicinity of *Trl*7 in the CC animals. However, after adjusting for confounders, we observed a significant difference in Ct.Th between KO and WT bones (P value = 0. 01), suggesting that the role of *Rhbdf2* is not limited to the trabecular compartment.

Discussion

Genetic reference population (GRP) are very efficient for the study of complex traits and biological systems, because (i) genotyping is only required once ("genotype once, phenotype many times", see below), and (ii) replicate individuals with the same genotype can be generated at will allowing for optimal experimental designs [27].

This article is the second to present the results of an ongoing quest to delineate the genetic determinants that govern microstructural bone traits [15]. Here we characterize several key microstructural properties of the mouse femoral bone to assess the extent to which they are heritable; to what environmental perturbations they are prone; and to identify candidate genes by which they are controlled. Our approach narrowed down a small number of putative candidate genes for 6 of the 8 examined phenotypes. Following merge analysis and RNA-seq, we validated our leading

candidate gene, *Rhbdf2,* using a knockout model, which confirmed the critical role of 318

Rhbdf2 on bone mass accrual and homeostasis. 319

While the heritability rates assessed here - determined to be over 60% for all 320
traits - confirmed our previous findings, the degree to which sex explains the 321
phenotypic variation was very subtle, and appeared only for the cortical traits; this 322
discrepancy may be due to the specific cohort composition used in this study (Table 323
S3), which includes a sex bias due to smaller number of females than males. We found 324
a total of five QTLs in six traits; BV/TV and Tb.N shared one QTL, and Tb.Th, Tb.Sp, 325
vBMD, and Ct.Th yielded one each. Importantly, although bone microarchitecture 326
factors are complex traits, our analyses highlighted no more than two loci for each trait; 327
it is likely that analyzing a larger number of CC lines would result in the identification 328
of further loci. 329

Our analyses yielded three QTLs for the trabecular traits and 2 QTLs for the 330
cortical traits. These are referred to as *Trl*7-9, and *Crl*1-2, respectively. In addition, and 331
in close proximity, to *Rhbdf2* (Rhomboid 5 Homolog 2; elaborated below), *Trl*7 332
includes *Ube2o* (Ubiquitin Conjugating Enzyme E2 O), which encodes an enzyme that 333
is an important interactant of SMAD6. Ube2o monoubiquitinates SMAD6, and thereby 334
facilitates the latter to bind BMP 1 receptors [28]. The signal transduction of BMP 1 is 335
in turn limited [29,30], and endochondral bone formation, instead of ossification, is 336
favored. Importantly, 4 week-old SMAD6-overexpressed mice have significantly lower 337
humeral and vertebral BV/TV ratios than their controls [29]. 338

At *Trl*8, *Klf17, Kdm4a,* and *Dmap1* are likely putative candidate genes. Since 339
Klf17 (Kruppel-Like Factor 17) is part of a network that includes BMPs [31] it is more 340
likely than a nearby gene, *St3gal3* (ST3 Beta-Galactoside Alpha-2,3-Sialyltransferase 341

3), to affect bone traits, although the latter has a greater merge strength. *Kdm4a* (Lysine 342 Demethylase 4A) encodes a histone demethylase that promotes the differentiation of 343 embryonal stem cells (ESCs) to an endothelial fate [32]; endothelial cells are implied 344 in regulation of bone formation [33]. *Dmap1* (DNA Methyltransferase 1 Associated 345 Protein) which encodes a DNA-methyl transferase known to regulate obesity 346 complications, and is differentially methylated in women with polycystic ovary 347 syndrome [34,35] had the highest meaningful merge density, and it might 348 epigenetically regulate bone formation as well. *Trl*9 includes two genes of interest to 349 bone biology: *Barhl2* and *Zfp644*. By interacting with caspase3, which is essential for 350 ossification [36], *Barhl2* (BarH Like Homeobox 2) can inhibit β-catenin activation [37], 351 and regulate the expression of chordin, a BMP signaling-detrimental protein [38]. 352 *Zfp644* (Zinc Finger Protein 644), which encodes a transcription repressor zinc-finger 353 protein, is upregulated in eight week-old ovariectomized mice following treatment with 354 estradiol [39], a treatment associated with reduced bone loss [40]. Further support for 355 the candidacy of *Barhl2* and *Zfp644* is given by the role of *Barhl2* in the development 356 of amacrine cells [41,42] and the association of *Zfp644* with myopia [43], a condition 357 speculated to propagate from amacrine cell signaling [44]; interestingly myopia was 358 linked to reduced postnatal bone mineral content in humans [45] and decreased 359 expression of BMP 2 and 5 in guinea pigs [46]. 360

The first of two cortical loci, *Crl*1 contains as likely candidates the genes *Asph* 361 and *Gdf6*. *Asph* (Aspartate Beta-Hydroxylase) encodes a protein that has a role in 362 regulating calcium homeostasis, which may affect bone metabolism [47]. *Gdf6* (Growth 363 Differentiation Factor 6) is bone morphogenetic protein 13: mice with mutated *Gdf6* 364 exhibit deformed bone formation in various skeletal sites; it is among the earliest known 365 markers of limb joint formation [48], expressed in joints of ankle and knee. In *Gdf6* 366

homozygous mutant mice, bones fuse at the joints early at the segmentation stage [49]. 367

For *Crl2*, we found *Hfe2*, *Bcl9*, *Notch2,* and *Prkab2* as potential candidate genes. *Hfe2* 368
(Hemochromatosis Type 2 (Juvenile)) encodes the BMP co-receptor hemojuvelin 369
which is expressed in skeletal muscles [50] and is responsible for juvenile 370
hemochromatosis, a condition linked to sex hormones depletion and osteoporosis [51]. 371
Bcl9 (B-Cell CLL/Lymphoma 9), the mammalian ortholog of the gene *Legless*, encodes 372
a protein essential to the Wnt/beta-catenin signaling which is important for bone 373
metabolism [52], without which the nuclear localization of β-catenin and myocyte 374
differentiation are compromised [53]. Of note, there are mutual effects between bone 375
and muscle, and accumulating evidence suggest many genes show pleiotropism with 376
respect to muscle strength and bone parameters [54]. *Notch2* encodes a member of the 377
notch protein family, which influence both osteoblasts and osteoclasts [55]; 378
specifically, *Notch2* is associated with the rare Hajdu-Cheney syndrome, that includes 379
severe osteoporosis as one of its main symptoms [56,57]. For this gene, we did not find 380
any significant merge logPs included within its limits. Interestingly, *Sec22b*, an 381
adjacent gene, had the strongest merge logP marks in this locus but no documented link 382
to bone biology. The third-strong gene in terms of merge values was *Prkab2* (Protein 383
Kinase AMP-Activated Non-Catalytic Subunit Beta 2). It encodes an enzyme which is 384
the regulatory subunit of mitogen-activated protein kinase (AMPK). AMPK widely 385
affects bone metabolism [58]. 386

Rhbdf2 is not yet supported by peer-reviewed reports as bearing a relation to 387
bone. Based on its closeness to the *Tlr7* peak (within the 50% CI), its merge strength 388
and RNA-seq in bone cells, we identified *Rhbdf2* as a likely causal gene associated with 389
BV/TV and Tb.N. We therefore analyzed the bone phenotype of *Rhbdf2$^{-/-}$* mice, to 390
validate the role of this gene in the modeling of the femoral cortex and trabeculae. 391

Rhbdf2 deletion affected all the examined trabecular traits as well as Ct.Th. While the   392

effects on BV/TV and Tb.N were in line with the haplotype mapping, the *Rhbdf2* locus   393

did not appear in any of the other traits. This however is expected, because the genetic   394

architecture of the working cohort is such that the assumed contributing variant of   395

Rhbdf2 is diluted and compensated, resulting in a QTL detected only for the most   396

affected traits. Noticeably, Tb.Sp differed greatly between the $Rhbdf2^{-/-}$ and control   397

mice but did not show up at the haplotype mapping; this might be due to (i) the great   398

diversity of the wild-type mice in Tb.Sp, and/or (ii) the need for complete knockout   399

rather than a mere SNP to detect significant changes in Tb.Sp, and/or (iii) the SNPs   400

giving rise to *Trl7* are functioning variants, with differential behavior affecting only   401

BV/TV and Tb.N. A similar interpretation may be valid for the cortical phenotype of   402

the $Rhbdf2^{-/-}$ mice. Importantly, the significant QTL peak we found in our GWAS for   403

BV/TV and Tb.N ended up revealing a gene that has an important skeletal function in   404

both the trabecular and cortical bone compartments.   405

    Rhbdf2 encodes the iRhom2 protein, a polytopic membrane protein that is a   406

catalytically inactive member of the rhomboid intramembrane serine proteases   407

superfamily [59]. iRhom2 is necessary in macrophages for the maturation and release   408

of the inflammatory cytokine tumor necrosis factor α (TNFα): it acts in the trafficking   409

of TACE, the protease that releases active TNFα from its membrane-tethered precursor   410

[60,61]. iRhom2 is also implicated in EGF-family growth factor signaling [62–64].   411

With a recent report of its role in trafficking of another protein, STING, it appears that   412

iRhom2 may have a wider role in regulating membrane trafficking [65]. iRhom2 was   413

also implicated in the regulation of CSF1R (macrophage stimulating factor 1 receptor),   414

a critical regulator of osteoclasts differentiation and survival [61,66–68]. *In vivo,*   415

Rhbdf2 has been implicated in esophageal cancer, wound healing, bone marrow 416

repopulation by monocytic cells, and inflammatory arthritis [63,69–71]. 417

Further work will be needed to identify the mechanism by which iRhom2 418

controls bone homeostasis; a possible direction could involve a positive feedback loop 419

that leads to differentiation of macrophages to osteoclasts. Indeed, iRhom2 stimulates 420

the secretion of TNFα by macrophages [68,72]; hyperactivates EGFR [73,74]; and 421

regulates CSF1R [75,76]. Although *Rhbdf2* is expressed in both the osteocyte and 422

osteoclast lineages, one cannot rule out the possibility that this gene regulates bone 423

remodeling by virtue of its expression in non-skeletal cells. 424

In summary, our analyses disclose several putative genes, several of which are 425

newly linked to a role in bone biology. A confirmation of one such gene, *Rhbdf2*, 426

provides the first conclusive evidence for its effects on bone microstructure. 427

Importantly, this study is the first demonstration of a physiological role of *Rhbdf2*. This 428

finding prompts future investigations to elucidate the exact mechanism of action of 429

Rhbdf2 and its contribution to osteoporosis in humans. 430

Materials and Methods 431

<u>Mice</u> 432

Mice aged 10 to 13 weeks (male $n = 103$; female $n = 71$), from 34 different CC lines 433

(average of 5 mice per line) were used in this study. The mice were at inbreeding 434

generations of 11 to 37, which correspond to 80-99.9% genetic homozygosity, 435

respectively. The mice were bred and maintained at the small animal facility of the 436

Sackler Faculty of Medicine, Tel Aviv University (TAU), Israel. They were housed on 437

hardwood chip bedding in open-top cages, with food and distilled water available *ad* 438

libitum, in an identical controlled environment (temperature = 25 ± 2°C; 60% ≤   439

humidity ≤ 85%) and a 12-hour light/dark cycle. All experiments protocols were   440

approved by the Institutional Animal Care and Use Committee (IACUC M-13-014) at   441

TAU, which follows the NIH/USA animal care and use protocols. The *Rhbdf2* knock   442

out mice and their WT counterparts were bred and maintained at the University of   443

Oxford as approved by license PPL80/2584 of the UK Home Office.   444

Specimen collection   445

Mice were intraperitoneally euthanized with cervical dislocation performed   446

approximately one minute after breathing stops owing to 5% Isoflurane inhalation. The   447

Rhbdf2 knock out mice and their WT counterparts were euthanized by inhalation of a   448

rising concentration of carbon dioxide followed by dislocation of the neck. Left femora   449

were harvested and fixed for 24 hours in 4% paraformaldehyde solution, and then stored   450

in 70% ethanol.   451

μCT evaluation   452

Whole left femora from each mouse were examined as described previously [77] by a   453

μCT system (μCT 50, Scanco Medical AG, Switzerland). Briefly, scans were   454

performed at a 10-μm resolution in all three spatial dimensions. The mineralized tissues   455

were differentially segmented by a global thresholding procedure [78]. All   456

morphometric parameters were determined by a direct 3D approach [79]. Parameters   457

analyzed were determined in the metaphyseal trabecular bone, which included   458

trabecular bone volume fraction (BV/TV; %), trabecular thickness (Tb.Th; μm),   459

trabecular number (Tb.N; mm^{-1}), trabecular connectivity density (Conn.D; mm^{-3}),   460

trabecular structure model index (SMI), and trabecular separation (Tb.Sp; mm). Two   461

additional parameters are characteristics of the mid-shaft diaphysis section, and include   462

volumetric bone mineral density (vBMD; mgHA/cm^3 [mg Hydroxy-Apatite per cm^3]) 463

and cortical thickness (Ct.Th; mm). All parameters were generated and termed 464

according to the Guidelines for assessment of bone microstructure in rodents using 465

micro–computed tomography [80]. 466

Genotyping 467

A representative male mouse from each line was initially genotyped with a high mouse 468

diversity array (MDA), which consists of 620,000 SNPs (Durrant et al., 2011). After 469

about two intervals of 4 generations of inbreeding, all the CC lines were regenotyped 470

by mouse universal genotype array (MUGA, 7,500 markers) and finally with the 471

MegaMuga (77,800 markers) SNP array to confirm their genotype status [19]. The 472

founder-based mosaic of each CC line was reconstructed using a hidden Markov model 473

(HMM) in which the hidden states are the founder haplotypes and the observed states 474

are the CC lines, to produce a probability matrix of descent from each founder. This 475

matrix was then pruned to about 11,000 SNPs by averaging across a window of 20 476

consecutive markers for faster analyses and reduction of genotyping errors [81]. 477

Statistical analyses and data acquisition 478

All statistical analyses were performed with the statistical software R (R core 479

development team 2009), including the package happy.hbrem [82]. 480

Heritability and covariate effects. Broad-sense heritability (H^2) was obtained for each 481

trait by fitting the trait (the independent variable) to the CC line label in a linear 482

regression model that incorporates relevant covariates (sex, age, batch, month, season, 483

year, and experimenter). ANOVA test was used to compare a null model (in which all 484

dependent variables are set to 0) with linear models that fit the covariates and the CC 485

line labels to the examined trait. Practically, the difference between the residual sum of squares (RSS; $\sum_1^n (\mu_i - \hat{\mu}_i)^2$) of the covariates model and that of the CC-line labels can be seen as the net genetic contribution to the trait. Thus, this difference divided by that of the covariate model gives an estimation of the heritability. Each covariate was calculated separately, by dividing the RSS difference between the null and full model with that of the null model. Let F_0 be the model that fits the trait to the covariates; F_1 the model that fits the trait to the covariates and the CC line label; and F_{00} the null model. Then, employing ANOVA, heritability is:

$$H^2 = (RSS(F_0) - RSS(F_1)) \,/\, RSS(F_{00}).$$

Similarly, the effects for each covariate were computed separately, by fitting each in F_0. The covariate effect is thus:

$$(RSS(F_{00}) - RSS(F_1)) \,/\, RSS(F_{00}).$$

H^2n was derived from H^2 according to Atamni *et al* [23].

Haplotype mapping. Each trait was fitted in a multiple linear regression model to the probability matrix of descent from each founder, including sex and age as covariates. The expected trait value from two ancestors, termed the genetic fit, is:

$$\mu_i = \mu + \sum_{s,t} F_{Li}(s,t)(\beta_s + \beta_t) = \mu + \sum_s \sum_t F_{Li}(s,t)\beta_s$$

where μ is a normally distributed trait mean, with sex and age incorporated; $F_{Li}(s,t)$ is the probability of descent from founders s and t; and $\beta_s + \beta_t$ is the additive effect of founders s and t. Because $\sum_s \sum_t F_{Li}(s,t) = 2$ for a diploid organism, the maximum likelihood estimates $\hat{\beta}_s$ are not independent. Thus, they are expressed here as differences from the WSB/EiJ founder effect, so that $\hat{\beta}_{WSB} = 0$. Number of members

per line was weighted and integrated in the linear model. ANOVA was then used to   508

compare this model with a null model where the founder effects are all set to 0; the   509

resulting *F*-statistic yielded the significance of the genetic model *vs*. the null model and   510

the negative 10-base logarithms of the P values (logP) were recorded.   511

Permutations of the CC lines between the phenotypes were used to set   512

significance thresholds levels. Founder effects are the estimates derived from the   513

multiple linear regression fit above.   514

Regional heritability (H_r^2) was hereafter computed by ANOVA as in the broad-   515

sense heritability computation, except that here null linear regression fit was compared   516

with a genetic linear regression fit with the probability matrix of the founder descent at   517

the peak QTL as the explanatory variable.   518

False discovery rate (FDR) was calculated using the p.adjust function in R, with   519

the method "BH" (Benjamini-Hochberg [83]).   520

Confidence intervals. Confidence intervals (CIs) were obtained both by simulations and   521

by the quick method of Li, 2011 [84]. In the simulations, we resampled the residuals of   522

the original linear regression fit at the peak of each QTL and rescanned 100 intervals   523

within 7-10 Mb of the original loci to find the highest logP. Accordingly, following   524

Durrant *et al.* [22], 1000 QTLs were simulated: if $\hat{t}_i$ is a random permutation of the   525

residuals of fitted genetic model at the QTL peak, and *K* is a marker interval in a   526

neighborhood of 3.5 to 5 Mb of the QTL peak *L*, a set of values for each trait, Z_{iK} is   527

provided by: 528

$$Z_{iK} = \hat{t}_i \exp(\hat{\mu} + \sum_s X_{Kis}\hat{\beta}_s).$$ 529

In the merge analysis the eight founder strains are partitioned and merged according to 531

the strain distribution pattern (SDP) of the alleles at the quantitative trait nucleotides 532

(QTN) within a given QTL (formerly obtained by the initial mapping). If we denote the 533

polymorphism as p, then $X_p = 1$ if s has allele a at p, and $X_p = 0$ otherwise [24]. Then, 534

at p, the probability of i to inherit alleles *a and b* from *s and t, respectively,* within L is 535

$$G_{pi}(a, b) = \sum_{s,t} X_p(a, s) X_p(b, t) F_{Li}(s, t).$$ 536

This merges the founder strains by p. The expected trait value in the merged strains 537

can now be inferred by 538

$$\sum_{a,b} G_{pi}(a, b)(\beta_a + \beta_b).$$ 539

Because this is a sub-model of the QTL model, it is expected to yield higher logP values 540

due to a reduction in the degrees of freedom. Significance was obtained by comparing 541

the merge model with the QTL model. Individual genes were extracted from the Sanger 542

mouse SNP repository (http://www.sanger.ac.uk/sanger/Mouse_SnpViewer). 543

Merge strength. We ranked the list of genes under each QTL according to the density 544

of merge logPs associated with them: only genes that had merge logPs above the 545

haplotype mapping reading, and above the threshold, plus logP=1 were included. We 546

then computed the relative density according to the density of a given gene's merge 547

logPs versus the locus' merge logP density. Let g be the region encompassed by a gene; 548

l the region encompassed by a QTL; and *mp* the merge logP values above the haplotype 549

P values plus 1. Then $g_i(mp) = 1$ if at SNP i there exists a *mp* and 0 otherwise. 550

Similarly $l_i(mp) = 1$ if at SNP j there exists a mp and 0 otherwise. The merge strength   551

(MS) is therefore: 552

$$MS\ [\%] = 100 * \frac{\sum_i g_i(mp)}{\sum_j l_j(mp)}\,\%.$$ 553

RNA-seq data   554

RNA-seq data from osteoclasts and osteocytes was obtained from gene expression   555
omnibus (GEO) database (accession numbers GSE72846 and GSE54784) and mapped   556
to the *mus musculus* assembly mm10 using tophat v. 2 [85]. Read counts were then   557
casted on the loci of interest using the R (R Core Team 2015) package   558
GenomicAlignments and raw read counts were taken. For the osteocytes, the data of   559
basal level day 3 was averaged.   560

Acknowledgments   561

This study was supported by Tel Aviv University starter funds and by Israel Science   562
Foundation (ISF) grant 1822/12 to YG, by Wellcome Trust grants 085906/Z/08/Z,   563
075491/Z/04, and 090532/Z/09/Z to RM, core funding by Tel-Aviv University to FI,   564
Wellcome Trust grant 101035/Z/13/Z and the Medical Research Council (programme   565
number U105178780) to MF, and by a fellowship from the Edmond J. Safra Center for   566
Bioinformatics at Tel-Aviv University to RL.   567

References   568

1.   Burge R, Dawson-Hughes B, Solomon DH, Wong JB, King A, Tosteson A.   569
      Incidence and economic burden of osteoporosis-related fractures in the United   570
      States, 2005-2025. J Bone Miner Res. 2007;22: 465–75.   571

doi:10.1359/jbmr.061113   572

2.   Dhanwal DK, Dennison EM, Harvey NC, Cooper C. Epidemiology of hip   573
fracture: Worldwide geographic variation. Indian J Orthop. 2011;45: 15–22.   574
doi:10.4103/0019-5413.73656   575

3.   Center JR, Nguyen T V, Schneider D, Sambrook PN, Eisman JA. Mortality   576
after all major types of osteoporotic fracture in men and women: an   577
observational study. Lancet (London, England). 1999;353: 878–82.   578
doi:10.1016/S0140-6736(98)09075-8   579

4.   Mizuguchi T, Furuta I, Watanabe Y, Tsukamoto K, Tomita H, Tsujihata M, et   580
al. LRP5, low-density-lipoprotein-receptor-related protein 5, is a determinant   581
for bone mineral density. J Hum Genet. 2004;49: 80–6. doi:10.1007/s10038-   582
003-0111-6   583

5.   Richards JB, Rivadeneira F, Inouye M, Pastinen TM, Soranzo N, Wilson SG, et   584
al. Bone mineral density, osteoporosis, and osteoporotic fractures: a genome-   585
wide association study. Lancet. 2008;371: 1505–12. doi:10.1016/S0140-   586
6736(08)60599-1   587

6.   Styrkarsdottir U, Halldorsson B V, Gretarsdottir S, Gudbjartsson DF, Walters   588
GB, Ingvarsson T, et al. Multiple genetic loci for bone mineral density and   589
fractures. N Engl J Med. 2008;358: 2355–65. doi:10.1056/NEJMoa0801197   590

7.   Trikalinos TA, Salanti G, Zintzaras E, Ioannidis JPA. Meta-analysis methods.   591
Adv Genet. 2008;60: 311–34. doi:10.1016/S0065-2660(07)00413-0   592

8.   Estrada K, Styrkarsdottir U, Evangelou E, Hsu Y-H, Duncan EL, Ntzani EE, et   593
al. Genome-wide meta-analysis identifies 56 bone mineral density loci and   594
reveals 14 loci associated with risk of fracture. Nat Genet. 2012;44: 491–501.   595

doi:10.1038/ng.2249 596

9. Paternoster L, Lorentzon M, Vandenput L, Karlsson MK, Ljunggren O, 597

    Kindmark A, et al. Genome-wide association meta-analysis of cortical bone 598

    mineral density unravels allelic heterogeneity at the RANKL locus and 599

    potential pleiotropic effects on bone. PLoS Genet. 2010;6: e1001217. 600

    doi:10.1371/journal.pgen.1001217 601

10. Paternoster L, Lorentzon M, Lehtimäki T, Eriksson J, Kähönen M, Raitakari O, 602

    et al. Genetic determinants of trabecular and cortical volumetric bone mineral 603

    densities and bone microstructure. PLoS Genet. 2013;9: e1003247. 604

    doi:10.1371/journal.pgen.1003247 605

11. Jovanovich A, Bùzková P, Chonchol M, Robbins J, Fink HA, de Boer IH, et al. 606

    Fibroblast growth factor 23, bone mineral density, and risk of hip fracture 607

    among older adults: the cardiovascular health study. J Clin Endocrinol Metab. 608

    2013;98: 3323–31. doi:10.1210/jc.2013-1152 609

12. Hsu Y-H, Kiel DP. Clinical review: Genome-wide association studies of 610

    skeletal phenotypes: what we have learned and where we are headed. J Clin 611

    Endocrinol Metab. 2012;97: E1958-77. doi:10.1210/jc.2012-1890 612

13. Styrkarsdottir U, Thorleifsson G, Sulem P, Gudbjartsson DF, Sigurdsson A, 613

    Jonasdottir A, et al. Nonsense mutation in the LGR4 gene is associated with 614

    several human diseases and other traits. Nature. Nature Publishing Group, a 615

    division of Macmillan Publishers Limited. All Rights Reserved.; 2013;497: 616

    517–20. doi:10.1038/nature12124 617

14. Iraqi F a, Athamni H, Dorman A, Salymah Y, Tomlinson I, Nashif A, et al. 618

    Heritability and coefficient of genetic variation analyses of phenotypic traits 619

provide strong basis for high-resolution QTL mapping in the Collaborative 620

Cross mouse genetic reference population. Mamm Genome. 2014;25: 109–19. 621

doi:10.1007/s00335-014-9503-5 622

15. Levy R, Mott RF, Iraqi FA, Gabet Y. Collaborative cross mice in a genetic 623

association study reveal new candidate genes for bone microarchitecture. BMC 624

Genomics. 2015;16: 1013. doi:10.1186/s12864-015-2213-x 625

16. Karasik D, Demissie S, Zhou Y, Lu D, Broe KE, Bouxsein ML, et al. 626

Heritability and Genetic Correlations for Bone Microarchitecture: The 627

Framingham Study Families. J Bone Miner Res. 2016; doi:10.1002/jbmr.2915 628

17. Threadgill DW, Hunter KW, Williams RW. Genetic dissection of complex and 629

quantitative traits: from fantasy to reality via a community effort. Mamm 630

Genome. 2002;13: 175–8. doi:10.1007/s00335-001-4001-Y 631

18. Churchill G a, Airey DC, Allayee H, Angel JM, Attie AD, Beatty J, et al. The 632

Collaborative Cross, a community resource for the genetic analysis of complex 633

traits. Nat Genet. 2004;36: 1133–7. doi:10.1038/ng1104-1133 634

19. Collaborative Cross Consortium. The genome architecture of the Collaborative 635

Cross mouse genetic reference population. Genetics. 2012;190: 389–401. 636

doi:10.1534/genetics.111.132639 637

20. Keane TM, Goodstadt L, Danecek P, White M a, Wong K, Yalcin B, et al. 638

Mouse genomic variation and its effect on phenotypes and gene regulation. 639

Nature. 2011;477: 289–94. doi:10.1038/nature10413 640

21. Roberts A, Pardo-Manuel de Villena F, Wang W, McMillan L, Threadgill DW. 641

The polymorphism architecture of mouse genetic resources elucidated using 642

genome-wide resequencing data: implications for QTL discovery and systems 643

genetics. Mamm Genome. 2007;18: 473–81. doi:10.1007/s00335-007-9045-1   644

22.   Durrant C, Tayem H, Yalcin B, Cleak J, Goodstadt L, Villena FP De, et al.   645

Collaborative Cross mice and their power to map host susceptibility to   646

Aspergillus fumigatus infection Collaborative Cross mice and their power to   647

map host susceptibility to Aspergillus fumigatus infection. 2011;   648

doi:10.1101/gr.118786.110   649

23.   Atamni HJA-T, Mott R, Soller M, Iraqi FA. High-fat-diet induced development   650

of increased fasting glucose levels and impaired response to intraperitoneal   651

glucose challenge in the collaborative cross mouse genetic reference   652

population. BMC Genet. BioMed Central; 2016;17: 10. doi:10.1186/s12863-   653

015-0321-x   654

24.   Yalcin B, Flint J, Mott R. Using progenitor strain information to identify   655

quantitative trait nucleotides in outbred mice. Genetics. 2005;171: 673–81.   656

doi:10.1534/genetics.104.028902   657

25.   Kim K, Punj V, Kim J-M, Lee S, Ulmer TS, Lu W, et al. MMP-9 facilitates   658

selective proteolysis of the histone H3 tail at genes necessary for proficient   659

osteoclastogenesis. Genes Dev. 2016;30: 208–19. doi:10.1101/gad.268714.115   660

26.   St. John HC, Bishop KA, Meyer MB, Benkusky NA, Leng N, Kendziorski C,   661

et al. The Osteoblast to Osteocyte Transition: Epigenetic Changes and   662

Response to the Vitamin D $_3$ Hormone. Mol Endocrinol. 2014;28: 1150–1165.   663

doi:10.1210/me.2014-1091   664

27.   Broman KW. The genomes of recombinant inbred lines. Genetics. 2005;169:   665

1133–46. doi:10.1534/genetics.104.035212   666

28.   Zhang X, Zhang J, Bauer A, Zhang L, Selinger DW, Lu CX, et al. Fine-tuning   667

BMP7 signalling in adipogenesis by UBE2O/E2-230K-mediated 668

monoubiquitination of SMAD6. EMBO J. 2013;32: 996–1007. 669

doi:10.1038/emboj.2013.38 670

29. Horiki M, Imamura T, Okamoto M, Hayashi M, Murai J, Myoui A, et al. 671

Smad6/Smurf1 overexpression in cartilage delays chondrocyte hypertrophy and 672

causes dwarfism with osteopenia. J Cell Biol. 2004;165: 433–45. 673

doi:10.1083/jcb.200311015 674

30. Estrada KD, Retting KN, Chin AM, Lyons KM. Smad6 is essential to limit 675

BMP signaling during cartilage development. J Bone Miner Res. 2011;26: 676

2498–510. doi:10.1002/jbmr.443 677

31. Kotkamp K, Mössner R, Allen A, Onichtchouk D, Driever W. A Pou5f1/Oct4 678

dependent Klf2a, Klf2b, and Klf17 regulatory sub-network contributes to EVL 679

and ectoderm development during zebrafish embryogenesis. Dev Biol. 680

2014;385: 433–47. doi:10.1016/j.ydbio.2013.10.025 681

32. Wu L, Wary KK, Revskoy S, Gao X, Tsang K, Komarova YA, et al. Histone 682

Demethylases KDM4A and KDM4C Regulate Differentiation of Embryonic 683

Stem Cells to Endothelial Cells. Stem cell reports. 2015;5: 10–21. 684

doi:10.1016/j.stemcr.2015.05.016 685

33. Collin-Osdoby P. Role of vascular endothelial cells in bone biology. J Cell 686

Biochem. 1994;55: 304–9. doi:10.1002/jcb.240550306 687

34. Kokosar M, Benrick A, Perfilyev A, Fornes R, Nilsson E, Maliqueo M, et al. 688

Epigenetic and Transcriptional Alterations in Human Adipose Tissue of 689

Polycystic Ovary Syndrome. Sci Rep. 2016;6: 22883. doi:10.1038/srep22883 690

35. Kamei Y, Suganami T, Ehara T, Kanai S, Hayashi K, Yamamoto Y, et al. 691

Increased expression of DNA methyltransferase 3a in obese adipose tissue: 692

studies with transgenic mice. Obesity (Silver Spring). 2010;18: 314–21.   693

doi:10.1038/oby.2009.246   694

36. Miura M, Chen X-D, Allen MR, Bi Y, Gronthos S, Seo B-M, et al. A crucial   695

role of caspase-3 in osteogenic differentiation of bone marrow stromal stem   696

cells. J Clin Invest. 2004;114: 1704–13. doi:10.1172/JCI20427   697

37. Juraver-Geslin HA, Ausseil JJ, Wassef M, Durand BC. Barhl2 limits growth of   698

the diencephalic primordium through Caspase3 inhibition of beta-catenin   699

activation. Proc Natl Acad Sci U S A. 2011;108: 2288–93.   700

doi:10.1073/pnas.1014017108   701

38. Larraín J, Bachiller D, Lu B, Agius E, Piccolo S, De Robertis EM. BMP-   702

binding modules in chordin: a model for signalling regulation in the   703

extracellular space. Development. 2000;127: 821–30. Available: 704

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2280033&tool=pm   705

centrez&rendertype=abstract   706

39. Davis AM, Mao J, Naz B, Kohl JA, Rosenfeld CS. Comparative effects of   707

estradiol, methyl-piperidino-pyrazole, raloxifene, and ICI 182 780 on gene   708

expression in the murine uterus. J Mol Endocrinol. 2008;41: 205–17.   709

doi:10.1677/JME-08-0029   710

40. Kameda T, Mano H, Yuasa T, Mori Y, Miyazawa K, Shiokawa M, et al.   711

Estrogen inhibits bone resorption by directly inducing apoptosis of the bone-   712

resorbing osteoclasts. J Exp Med. 1997;186: 489–95. Available: 713

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2199029&tool=pm   714

centrez&rendertype=abstract   715

41. Ding Q, Chen H, Xie X, Libby RT, Tian N, Gan L. BARHL2 differentially   716

regulates the development of retinal amacrine and ganglion neurons. J   717

Neurosci. 2009;29: 3992–4003. doi:10.1523/JNEUROSCI.5237-08.2009   718

42. Mo Z, Li S, Yang X, Xiang M. Role of the Barhl2 homeobox gene in the   719

specification of glycinergic amacrine cells. Development. 2004;131: 1607–18.   720

doi:10.1242/dev.01071   721

43. Shi Y, Li Y, Zhang D, Zhang H, Li Y, Lu F, et al. Exome sequencing identifies   722

ZNF644 mutations in high myopia. PLoS Genet. 2011;7: e1002084.   723

doi:10.1371/journal.pgen.1002084   724

44. Chen JC, Brown B, Schmid KL. Evaluation of inner retinal function in myopia   725

using oscillatory potentials of the multifocal electroretinogram. Vision Res.   726

2006;46: 4096–103. doi:10.1016/j.visres.2006.07.033   727

45. Pohlandt F. Hypothesis: myopia of prematurity is caused by postnatal bone   728

mineral deficiency. Eur J Pediatr. 1994;153: 234–6. Available: 729

http://www.ncbi.nlm.nih.gov/pubmed/8194552   730

46. Wang Q, Xue M-L, Zhao G-Q, Liu M-G, Ma Y-N, Ma Y. Form-deprivation   731

myopia induces decreased expression of bone morphogenetic protein-2, 5 in   732

guinea pig sclera. Int J Ophthalmol. 2015;8: 39–45. doi:10.3980/j.issn.2222-   733

3959.2015.01.07   734

47. Pruitt KD, Brown GR, Hiatt SM, Thibaud-Nissen F, Astashyn A, Ermolaeva O,   735

et al. RefSeq: an update on mammalian reference sequences. Nucleic Acids   736

Res. 2014;42: D756-63. doi:10.1093/nar/gkt1114   737

48. Chang SC, Hoang B, Thomas JT, Vukicevic S, Luyten FP, Ryba NJ, et al.   738

Cartilage-derived morphogenetic proteins. New members of the transforming   739

growth factor-beta superfamily predominantly expressed in long bones during 740

human embryonic development. J Biol Chem. 1994;269: 28227–34. Available: 741

http://www.ncbi.nlm.nih.gov/pubmed/7961761 742

49. Settle SH, Rountree RB, Sinha A, Thacker A, Higgins K, Kingsley DM. 743

Multiple joint and skeletal patterning defects caused by single and double 744

mutations in the mouse Gdf6 and Gdf5 genes. Dev Biol. 2003;254: 116–30. 745

Available: http://www.ncbi.nlm.nih.gov/pubmed/12606286 746

50. Verga Falzacappa MV, Casanovas G, Hentze MW, Muckenthaler MU. A bone 747

morphogenetic protein (BMP)-responsive element in the hepcidin promoter 748

controls HFE2-mediated hepatic hepcidin expression and its response to IL-6 in 749

cultured cells. J Mol Med (Berl). 2008;86: 531–40. doi:10.1007/s00109-008- 750

0313-7 751

51. Angelopoulos NG, Goula AK, Papanikolaou G, Tolis G. Osteoporosis in HFE2 752

juvenile hemochromatosis. A case report and review of the literature. 753

Osteoporos Int. 2006;17: 150–5. doi:10.1007/s00198-005-1920-6 754

52. Baron R, Kneissel M. WNT signaling in bone homeostasis and disease: from 755

human mutations to treatments. Nat Med. Nature Publishing Group, a division 756

of Macmillan Publishers Limited. All Rights Reserved.; 2013;19: 179–92. 757

doi:10.1038/nm.3074 758

53. Brack AS, Murphy-Seiler F, Hanifi J, Deka J, Eyckerman S, Keller C, et al. 759

BCL9 is an essential component of canonical Wnt signaling that mediates the 760

differentiation of myogenic progenitors during muscle regeneration. Dev Biol. 761

2009;335: 93–105. doi:10.1016/j.ydbio.2009.08.014 762

54. Karasik D, Kiel DP. Genetics of the musculoskeletal system: a pleiotropic 763

approach. J Bone Miner Res. 2008;23: 788–802. doi:10.1359/jbmr.080218   764

55.   Bai S, Kopan R, Zou W, Hilton MJ, Ong C, Long F, et al. NOTCH1 regulates   765

        osteoclastogenesis directly in osteoclast precursors and indirectly via osteoblast   766

        lineage cells. J Biol Chem. 2008;283: 6509–18. doi:10.1074/jbc.M707000200   767

56.   Regan J, Long F. Notch signaling and bone remodeling. Curr Osteoporos Rep.   768

        2013;11: 126–9. doi:10.1007/s11914-013-0145-4   769

57.   Canalis E, Zanotti S. Hajdu-Cheney syndrome: a review. Orphanet J Rare Dis.   770

        BioMed Central; 2014;9: 200. doi:10.1186/s13023-014-0200-y   771

58.   Jeyabalan J, Shah M, Viollet B, Chenu C. AMP-activated protein kinase   772

        pathway and bone metabolism. J Endocrinol. 2012;212: 277–90.   773

        doi:10.1530/JOE-11-0306   774

59.   Lemberg MK, Freeman M. Functional and evolutionary implications of   775

        enhanced genomic analysis of rhomboid intramembrane proteases. Genome   776

        Res. 2007;17: 1634–1646. doi:10.1101/gr.6425307   777

60.   Adrain C, Zettl M, Christova Y, Taylor N, Freeman M, References A.   778

        Supporting Online Material for. 2012;225. doi:10.1126/science.1214400   779

61.   McIlwain DR, Lang PA, Maretzky T, Hamada K, Ohishi K, Maney SK, et al.   780

        iRhom2 Regulation of TACE Controls TNF-Mediated Protection Against   781

        Listeria and Responses to LPS. Science (80-). 2012;335: 229–232.   782

        doi:10.1126/science.1214448   783

62.   Siggs OM, Grieve A, Xu H, Bambrough P, Christova Y, Freeman M. Genetic   784

        interaction implicates iRhom2 in the regulation of EGF receptor signalling in   785

        mice. Biol Open. 2014;3.   786

63.   Hosur V, Johnson KR, Burzenski LM, Stearns TM, Maser RS, Shultz LD.   787

Rhbdf2 mutations increase its protein stability and drive EGFR hyperactivation   788

through enhanced secretion of amphiregulin. Proc Natl Acad Sci. 2014;111:   789

E2200–E2209. doi:10.1073/pnas.1323908111   790

64.   Li X, Maretzky T, Weskamp G, Monette S, Qing X, Issuree PDA, et al.   791

iRhoms 1 and 2 are essential upstream regulators of ADAM17-dependent   792

EGFR signaling. Proc Natl Acad Sci U S A. National Academy of Sciences;   793

2015;112: 6080–5. doi:10.1073/pnas.1505649112   794

65.   Luo W-W, Li S, Li C, Lian H, Yang Q, Zhong B, et al. iRhom2 is essential for   795

innate immunity to DNA viruses by mediating trafficking and stability of the   796

adaptor STING. Nat Immunol. 2016;17: 1057–1066. doi:10.1038/ni.3510   797

66.   Siggs OM, Xiao N, Wang Y, Shi H, Tomisato W, Li X, et al. iRhom2 is   798

required for the secretion of mouse TNFα. Blood. American Society of   799

Hematology; 2012;119: 5769–71. doi:10.1182/blood-2012-03-417949   800

67.   Adrain C, Zettl M, Christova Y, Taylor N, Freeman M. Tumor Necrosis Factor   801

Signaling Requires iRhom2 to Promote Trafficking and Activation of TACE.   802

Science (80-). 2012;335: 225–228. doi:10.1126/science.1214400   803

68.   Udagawa N, Takahashi N, Akatsu T, Tanaka H, Sasaki T, Nishihara T, et al.   804

Origin of osteoclasts: mature monocytes and macrophages are capable of   805

differentiating into osteoclasts under a suitable microenvironment prepared by   806

bone marrow-derived stromal cells. Proc Natl Acad Sci U S A. National   807

Academy of Sciences; 1990;87: 7260–4. Available: 808

http://www.ncbi.nlm.nih.gov/pubmed/2169622   809

69.   Issuree PDA, Maretzky T, McIlwain DR, Monette S, Qing X, Lang PA, et al.   810

iRHOM2 is a critical pathogenic mediator of inflammatory arthritis. J Clin   811

Invest. American Society for Clinical Investigation; 2013;123: 928–32.   812

doi:10.1172/JCI66168   813

70.   Qing X, D. Rogers L, Mortha A, Lavin Y, Redecha P, Issuree PD, et al.   814

iRhom2 regulates CSF1R cell surface expression and non-steady state   815

myelopoiesis in mice. Eur J Immunol. 2016;46: 2737–2748.   816

doi:10.1002/eji.201646482   817

71.   Blaydon DC, Etheridge SL, Risk JM, Hennies H-C, Gay LJ, Carroll R, et al.   818

RHBDF2 mutations are associated with tylosis, a familial esophageal cancer   819

syndrome. Am J Hum Genet. Elsevier; 2012;90: 340–6.   820

doi:10.1016/j.ajhg.2011.12.008   821

72.   Kobayashi K, Takahashi N, Jimi E, Udagawa N, Takami M, Kotake S, et al.   822

Tumor necrosis factor alpha stimulates osteoclast differentiation by a   823

mechanism independent of the ODF/RANKL-RANK interaction. J Exp Med.   824

2000;191: 275–86. Available: 825

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2195746&tool=pm   826

centrez&rendertype=abstract   827

73.   Yi T, Lee H-L, Cha J-H, Ko S-I, Kim H-J, Shin H-I, et al. Epidermal growth   828

factor receptor regulates osteoclast differentiation and survival through cross-   829

talking with RANK signaling. J Cell Physiol. 2008;217: 409–422.   830

doi:10.1002/jcp.21511   831

74.   Hosur V, Johnson KR, Burzenski LM, Stearns TM, Maser RS, Shultz LD.   832

Rhbdf2 mutations increase its protein stability and drive EGFR hyperactivation   833

through enhanced secretion of amphiregulin. Proc Natl Acad Sci. 2014;111:   834

E2200–E2209. doi:10.1073/pnas.1323908111   835

75. Hung JY, Horn D, Woodruff K, Prihoda T, LeSaux C, Peters J, et al. Colony-stimulating factor 1 potentiates lung cancer bone metastasis. Lab Investig. 2014;94: 371–381. doi:10.1038/labinvest.2014.1

76. Qing X, D. Rogers L, Mortha A, Lavin Y, Redecha P, Issuree PD, et al. iRhom2 regulates CSF1R cell surface expression and non-steady state myelopoiesis in mice. Eur J Immunol. 2016;46: 2737–2748. doi:10.1002/eji.201646482

77. Hiram-Bab S, Liron T, Deshet-Unger N, Mittelman M, Gassmann M, Rauner M, et al. Erythropoietin directly stimulates osteoclast precursors and induces bone loss. FASEB J. 2015;29: 1890–900. doi:10.1096/fj.14-259085

78. Rüegsegger P, Koller B, Müller R. A microtomographic system for the nondestructive evaluation of bone architecture. Calcif Tissue Int. 1996;58: 24–29. doi:10.1007/BF02509542

79. Hildebrand T, Laib A, Müller R, Dequeker J, Rüegsegger P. Direct three-dimensional morphometric analysis of human cancellous bone: microstructural data from spine, femur, iliac crest, and calcaneus. J Bone Miner Res. 1999;14: 1167–74. doi:10.1359/jbmr.1999.14.7.1167

80. Bouxsein ML, Boyd SK, Christiansen BA, Guldberg RE, Jepsen KJ, Müller R. Guidelines for assessment of bone microstructure in rodents using micro-computed tomography. J Bone Miner Res. 2010;25: 1468–86. doi:10.1002/jbmr.141

81. Hall M, Manship G, Morahan G, Pettit K, Scholten J, Tweedie K, et al. The genome architecture of the Collaborative Cross mouse genetic reference population. Genetics. 2012;190: 389–401. doi:10.1534/genetics.111.132639

82. Mott R, Talbot CJ, Turri MG, Collins a C, Flint J. A method for fine mapping
     quantitative trait loci in outbred animal stocks. Proc Natl Acad Sci U S A.
     2000;97: 12649–54. doi:10.1073/pnas.230304397

83. Hochberg Y. Controlling the False Discovery Rate : A Practical and Powerful
     Approach to Multiple Testing Author (s): Yoav Benjamini and Yosef
     Hochberg Source : Journal of the Royal Statistical Society . Series B (
     Methodological), Vol . 57 , No . 1 (1995), Publi. 2016;57: 289–300.

84. Li H. A quick method to calculate QTL confidence interval. J Genet. 2011;90:
     355–60. Available: http://www.ncbi.nlm.nih.gov/pubmed/21869489

85. Durrant C, Tayem H, Yalcin B, Cleak J, Goodstadt L, de Villena FP-M, et al.
     Collaborative Cross mice and their power to map host susceptibility to
     Aspergillus fumigatus infection. Genome Res. 2011;21: 1239–48.
     doi:10.1101/gr.118786.110

86. Trapnell C, Pachter L, Salzberg SL. TopHat: discovering splice junctions with
     RNA-Seq. Bioinformatics. Oxford University Press; 2009;25: 1105–11.
     doi:10.1093/bioinformatics/btp120

Figure legends

Figure 1. µCT images of trabecular and cortical bone of the femora of representative
male CC mice. **(A)** Trabecular bone. Left: IL-2452, Right: IL-1513. **(B)** Cortical
bone. Left: IL-785, Right: IL-2689.

Figure 2 Trabecular and cortical traits distributions across the CC lines. X-axis is the
lines, y-axis is the trait means **A**. From top left, counter-clockwise: BV/TV (%), Tb.N
(mm^{-1}), Tb.Th (um), Conn.D (mm^{-3}), SMI, and Tb.Sp (mm). **B**. Left, vBMD
(mgHA/cm^3), right, Ct.Th (mm). Color codes group line(s) which significantly differ

from other groups. Lines are ordered inconsistently among the traits, per trait-specific   884

descending order. Refer to Table S3 for more details.   885

Figure 3 Haplotype association maps for the trabecular and cortical traits. X-axis is the   887

position on the chromosome, y-axis is the −logP value of the association. Lower   888

threshold represents the 95th percentile of 200 simulations, and top represents the 99th 889

percentile. Loci above the 99% cut-off were further investigated. From top to bottom: 890

BV/TV, Tb.N, Tb.Th, Tb.Sp, vBMD, and Ct.Th.   891

Figure 4. Ancestral effects relative to WSB. Y axis is the strain deviation relative to   894

WSB, x axis is the different strains of the eight CC founders. **(A)** to **(F)**: *Trl7* to *Trl9*,   895

Crl1, and *Crl2*, respectively.   896

Figure 5 Traits distribution at the marker UNC20471277 across bearers of   898

homozygous and heterozygous alleles, separated by sex. X-axis is the allelic variation   899

at the marker, y-axis is the trait value.   900

Figure 6 Merge analysis. Readings below logP = 4 are elided for brevity. X axis is   901

the position on the genome in Mb; y left axis is the logP score; y right axis is the   902

recombination rate scale; colored bars are genes (note that only strong putative   903

candidate genes are shown.); cyan line is the recombination rate; black continues line   904

is the haplotype test's peak; dashed line is the 99% permutation threshold. **(A)** to **(F)**:   905

Trl7 of BV/TV, Trl7 of Tb.N, Trl8 of Tb.Th, Trl9 of Tb.Sp, Crl1 of Ct.Th, and Crl2   906

of vBMD, respectively.   907

Figure 7 *Rhbdf2* knockout versus wildtype for each of the studied traits. Left is KO,   909

right is WT. PV is the confounder-adjusted P value. The unadjusted P value is in   910

brackets. PV = 0 means PV < 0.001.   911

Figure 8 μCT images of three-dimensional representative cortical and trabecular   913

bones reconstructions for *Rhbdf2* knockout and wildtype. Left: KO, right: WT. **(A)**   914

Trabecular bone. **(B)** Cortical bone. All samples were of male mice, aged 11 weeks.   915

Supporting information captions   916

Figure S1 *Trait distributions for Males and Females.* **A1, A2**: Trabecular and cortical   917

bone, respectively, males. **B1, B2**: trabecular and cortical bone, respectively, females.   918

X-axis is the lines, y-axis is the trait means **A**. From top left, counter-clockwise: 919

BV/TV (%), Tb.N (mm-1), Tb.Th (um), Conn.D (mm-3), SMI, and Tb.Sp (mm). **B**:   920

Left, vBMD (mgHA/cm3); right, Ct.Th (mm). Color codes group line(s) which   921

significantly differ from other groups. Lines are ordered inconsistently among the   922

traits, according to trait-specific descending order.   923

Figure S2 *Haplotype association maps for the trabecular and cortical traits.* X-axis   925

is the position on the chromosome, y-axis is the –logP value of the association. Lower   926

threshold represents the 95th percentile of 200 simulations, and top represents the 9th   927

percentile. Loci above the 99% cut-off were further investigated. From top to bottom:

Conn.D, SMI.

Figure S3 *Confidence interval simulations.* Loci at a neighborhood of 3-5 Mb around

the original locus were simulated by permuting the residual sum of squares of the

related phenotype. Maximum logP was obtained along with its relative position in Mb

to the original QTL (histograms, left panels), and with the number of markers from

the original QTL (boxplots, right panels). (*A*) and (*B*) show simulations results for the

BV/TV and Tb.N loci. These determined with high confidence that the peak QTL is

responsible for the effect seen in the haplotype scan, thus the narrow CI; (*C*) to (*F)*

show simulation results for Tb.Th, Tb.Sp, Ct.Th, and vBMD,

respectively; note the narrow CI for *Crl2* (vBMD), wide for *Trl8* (Tb.Th), and wider

still for *Crl1*(Ct.Th).

Figure S4 *RNA-seq of osteoclasts.* Gene names are on the right of each plot. Green

represents plus-stranded genes, black represents minus-stranded genes. Y-axis is the

raw expression count, where the negative scale refers to minus-stranded gene count.

Each bracket corresponds to a particular gene; left-to-right green (black) brackets fit

green (black) top-to-bottom gene-names.

Figure S5 *RNA-seq of osteocytes.* Gene names are on the right of each plot. Orange

represents plus-stranded genes, black represents minus-stranded genes. Y-axis is the

raw expression count, where the negative scale refers to minus-stranded gene count.

Each bracket corresponds to a particular gene; left-to-right orange (black) brackets fit

orange (black) top-to-bottom gene names

Table S1 *Pearson's correlations between measured μCT traits.* Pairwise correlations

for each trait are given as Pearson's *r*. In bold are the 5 most strong correlations.

Table S2 *Covariates effects. LogP is the negative logarithm of P value.* Effects of   952

covariates (i.e., the degree to which each covariate explains the phenotypic   953

difference). Values were determined by regressing the covariates along with the CC   954

lines, and running an ANOVA test. Note the covariates prominent effect on the   955

cortical traits. The batch effect was the strongest, affecting Tb.Th, Ct.Th, and vBMD.   956

Table S3 *Trait values per CC line included in this study (for both sexes combined).*   958
Trait means for each line, including number of members and standard error (SE). NA   959
means data were not available.   960

Table S4 *A comprehensive list of all genes under the 95% confidence-interval (CI)*   962

for the QTLs identified in this study. Light blue is the 95% CI region; blue is the 90%   963

CI region; and pink is the 50% CI region. Merge strength column gives the   964

normalized dosage of merge logP values as outlined in the methods section. Gene   965

symbols in bold type are discussed in the manuscript. Genes under the QTLs for the   966

cortical and trabecular traits. Light blue is the 95% CI, blue is the 90% CI, and red is   967

the 50% CI. "Merge Strength" refers to the proportion of merge logP values   968

constricted to the region of the specific gene. Note that for T*rl7* which is common to   969

BV/TV and Tb.N, the average values between the two are provided.   970

BV/TV
PV < 0.001

Tb.N
PV < 0.001

Tb.Th
PV < 0.001

Conn.D
PV = 0.006

SMI
PV < 0.001

Tb.Sp
PV = 0.2

Ct.Th
PV = 0.07

vBMD
PV = 0.8

BV/TV

PV = 0.002 (0.031)

Tb.N

PV = 0 (0.027)

Tb.Th

PV = 0.424 (0.222)

Conn.D

PV = 0.008 (0.046)

SMI

PV = 0.033 (0.156)

Tb.Sp

PV = 0 (0.017)

vBMD

PV = 0.157 (0.062)

Ct.Th

PV = 0.018 (0.807)