Title: Genomes of all known members of a *Plasmodium* subgenus reveal paths to
virulent human malaria

Authors: Thomas D. Otto[1,†,#,*], Aude Gilabert[2,†], Thomas Crellen[1,3], Ulrike Böhme[1], Céline
Arnathau[2], Mandy Sanders[1], Samuel Oyola[1,4], Alain Prince Okouga[5], Larson Boundenga[5], Eric
Willaume[6], Barthélémy Ngoubangoye[5], Nancy Diamella Moukodoum[5], Christophe Paupy[2], Patrick
Durand[2], Virginie Rougeron[2,5], Benjamin Ollomo[5], François Renaud[2], Chris Newbold[1,7], Matthew
Berriman[1,*] & Franck Prugnolle[2,5,*]

Affiliations:

[1] Wellcome Trust Sanger Institute, Wellcome Genome Campus, Hinxton CB10 1SA, United
Kingdom

[2] Laboratoire MIVEGEC, UMR 5290-224 CNRS 5290-IRD 224-UM, Montpellier, France

[3] Department of Infectious Disease Epidemiology, Imperial College London, St Mary's Campus,
Norfolk Place, London W2 1PG, United Kingdom

[4] International Livestock Research Institute, Box 30709, Nairobi, Kenya (current address)

[5] Centre International de Recherches Médicales de Franceville, Franceville, Gabon

[6] Sodepal, Parc of la Lékédi, Bakoumba, Gabon

[7] Weatherall Institute of Molecular Medicine, University of Oxford, John Radcliffe Hospital,
Oxford OX3 9DS, United Kingdom

Current Address: Centre of Immunobiology, Institute of Infection, Immunity & Inflammation,
College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, United
Kingdom

[*]Correspondence to: Thomas D. Otto (thomasdan.otto@glasgow.ac.uk), Matthew Berriman
(mb4@sanger.ac.uk) or Franck Prugnolle (franck.prugnolle@ird.fr)

[†]These authors contributed equally.

Abstract: *Plasmodium falciparum*, the most virulent agent of human malaria, shares a recent
common ancestor with the gorilla parasite *P. praefalciparum*. Little is known about the other gorilla
and chimpanzee-infecting species in the same (*Laverania*) subgenus as *P. falciparum* but none of
them are capable of establishing repeated infection and transmission in humans. To elucidate
underlying mechanisms and the evolutionary history of this subgenus, we have generated multiple
genomes from all known *Laverania* species. The completeness of our dataset allows us to conclude
that interspecific gene transfers as well as convergent evolution were important in the evolution of
these species. Striking copy number and structural variations were observed within gene families
and one, *stevor* shows a host specific sequence pattern. The complete genome sequence of the
closest ancestor of *P. falciparum* enables us to estimate confidently for the first time the timing of
the beginning of speciation to be 40,000-60,000 years ago followed by a population bottleneck
around 4,000-6,000 years ago. Our data allow us also to search in detail for the features of *P.
falciparum* that made it the only member of the *Laverania* able to infect and spread in humans.

38
39

Main Text:

The evolutionary history of *Plasmodium falciparum*, the most common and deadliest human malaria parasite, has been the subject of uncertainty and debate[1,2]. Recently it has become clear that *P. falciparum* is derived from a group of parasites infecting African Great Apes and known as the *Laverania* subgenus[2]. Until 2009, the only other species known in this subgenus was a parasite of chimpanzees known as *P. reichenowi*, for which only one isolate was available[3]. It is now clear that there are a total of at least seven species in Great Apes that naturally infect chimpanzees (*P. gaboni*, *P. billcollinsi* and *P. reichenowi*), gorillas (*P. praefalciparum*, *P. blacklocki* and *P. adleri*)[4,5], or humans (*P. falciparum*) (Fig. 1a). Within this group, *P. falciparum* is the only parasite that has successfully adapted to humans after a transfer from gorillas and subsequently spread all over the world[2].

Over time there have been various estimates concerning the evolutionary history of *P. falciparum* with the speciation event having been estimated to be anywhere between 10,000 to 5.5 million years ago, the latter falsely based on the date of the chimpanzee–human split[6,7]. Others report a bottleneck less than 10,000 years ago[8], but suggest a drop to a single progenitor parasite. The latter seems unlikely due to the presence of allelic dimorphisms that predate speciation events and therefore could not have both been transmitted if a new species were founded by a single individual infection. Also, the dating of the speciation cannot be accurately estimated without the genome sequence of *P. praefalciparum,* the closest living sister species to *P. falciparum*.

The absence of *in vitro* culture or a usable animal model has precluded obtaining sufficient DNA for full genome sequencing and has hindered investigation of the *Laverania.* So far just the full draft genome of *P. reichenowi* [9] and a nearly complete draft sequence of *P. gaboni*[6] are available. These data together with additional PCR based approaches[10] have provided important insights into the evolution of this subgenus, including the lateral gene transfer of the *rh5* locus, the early expansion of the FIKK gene family and the observation that the common ancestor also had *var* genes. Our data confirm and significantly extend these findings. However, the lack of whole genome information for the whole subgenus (particularly *P. praefalciparum*) has severely constrained the scope of subsequent analyses.

To investigate the evolutionary history of all known members of the *Laverania* subgenus and to address the question of why *P. falciparum* is the only extant species to have adapted successfully to humans, we have sequenced multiple genotypes of all known *Laverania* species.

 Genome sequencing from six *Laverania* species

     Blood samples were taken during successive routine sanitary controls, from four gorillas and

 seven chimpanzees living in a sanctuary or quarantine facility prior to release (see Methods). A

 total of 15 blood samples were positive for ape malaria parasites by PCR. Despite low parasitemia

 in most animals, a combination of host DNA depletion, parasite cell sorting and amplification

 methods enabled sufficient parasite DNA templates to be obtained for short-read (Illumina) and

 long read (Pacific Bioscience) sequencing (Table 1). Mixed-species infections were frequent but

 resolved by utilising sequence data from single infections, resulting in 19 genotypes

 (Supplementary Table 1). The dominant genotype in each sample was assembled *de novo* (see

 Methods) using long read technology into a reference genome for six malaria parasite species: *P.*

 praefalciparum, *P. blacklocki*, *P. adleri, P. billcollinsi*, *P. gaboni* and *P. reichenowi*. The

 assemblies comprised 44-97 scaffolds (Table 1), with large contigs containing the subtelomeric

 regions and internal gene clusters that house multigene families known in *P. falciparum* and *P.*

 reichenowi to be involved in virulence and host-pathogen interactions. The high quality of the

 assemblies compared to those obtained previously is illustrated by the good representation of multi-

 gene family members (Supplementary Table 2) and the large number of one-to-one orthologues

 obtained between the different reference genomes (4,350 among the seven species and 4,826

 between *P. falciparum*, *P. praefalciparum* and *P. reichenowi*). Two to four additional genomes

 were obtained for each species except for *P. blacklocki* (Table 1, Supplementary Table 1).

 Speciation history in the *Laverania* sub-genus

     Conservation of gene content and synteny is striking between these complete genomes and

 enabled us to reconstruct with confidence the relationships between different *Laverania* species, to

 compare their relative genetic diversity (Fig 2a, Supplementary Fig. 1) and to estimate the age of

 the different speciation events that led to the extant species. The latter has been problematic in the

 past due to the lack of both complete genome data and accurate estimates of mutation rate and

 generation time. Using the most divergent estimates of generation time and measured mutation

 rates from the *P. falciparum* literature, we found the data converge to 0.9-1.5 mutations per year per

 genome (Supplementary Note 1). We observed a similar substitution rate *in vivo* by examining

 existing sequence data for five geographically diverse isolates, covering a 200-kb region

 surrounding the PfCRT gene that is relatively conserved due to a selective sweep resulting from

 chloroquine use (Supplementary Note 1; Supplementary Figure 2). The fact that these two figures

 are similar suggests that the *in vitro* mutation rate may have been underestimated since many

mutations will be lost by genetic drift. Since no data is available for the other species we have

assumed hereon that these values generalise across the subgenus. From our Bayesian whole-genome

estimates, the ancestor of all current day parasites of this subgenus existed 0.7–1.2 million years

ago, a time at which the subgenus divided into two main clades, A (*P. adleri* and *P. gaboni*) and B

that includes the remaining species (Fig. 1a). Our range of values is far more recent than previous

estimates[3,11]. Following the Clade A/B subdivision, several speciation events occurred leading

either to new chimpanzee or gorilla parasites. Interestingly, the divergence between *P. adleri* and *P.*

gaboni in one lineage and *P. reichenowi* and the ancestor of *P. praefalciparum/P. falciparum* in the

other lineage occurred at approximately the same time (140–230 thousand years ago; Fig. 1a, Table

2). Based on our coalescence estimates, *P. falciparum* begun to emerge in humans from *P.*

praefalciparum around 40–60 thousand years ago (Fig. 1a), significantly later than the evolution of

the first modern humans and their spread throughout Africa[12]. Our analysis also indicates

significant gene flow between these two parasite species after their initial divergence (Table 2).

P. falciparum has strikingly low diversity (π =0.0004), compared with the other *Laverania*

species (0.002–0.0049) (Supplementary Fig. 1). It has been proposed that *P. falciparum* arose from

a single transfer of *P. praefalciparum* into humans[6] and based in part on the paucity of neutral SNPs

within the genome, that *P. falciparum* emerged from a bottleneck of a single parasite around 10,000

years ago, after agriculture was established[6,8]. In light of our results, we estimate that the

P. falciparum population declined around 11,000 years ago and reached a minimum about 5,000

years ago (Fig. 1b) with an effective population size (*Ne*) of around 3,000 (Supplementary Note 1;

generally the census number of parasites is higher than *Ne*[13]). It is important to note that the

methods we used to estimate *Ne* do not rely on the Fisher-Wright model, that could give problems

as previously reported[14]. The hypothesis of a single progenitor is also inconsistent with the

observation of several ancient gene dimorphisms that have been observed in *P. falciparum*. A

previous analysis using *P. reichenowi* and *P. gaboni* sequence data, provided some evidence that

different dimorphic loci diverged at different points in the tree[15]. Looking at each of these

P. falciparum loci across the *Laverania*, we found different patterns of evolution at the *msp1*,

var1csa, and *msp3* loci (Supplementary Fig. 3a). Most strikingly, a mutually exclusive dimorphism

(described as MAD20/K1[16]) in the central 70% of the *msp1* sequence, clearly pre-dates the

P. falciparum–*P. praefalciparum* common ancestor and dimorphism in *var1csa* (an unusual *var*

gene of unknown function that is transcribed late in the asexual cycle) occurred before the split with

P. reichenowi.

In contrast, the gene *eba-175* that encodes a parasite surface ligand involved in red blood cell

invasion contains a dimorphism that arose after the emergence of *P. falciparum* (Supplementary

Fig. 3b). The time to the most recent common ancestor of *eba-175* has been estimated as 130–140

thousand years in an analysis[17] that assumed *P. falciparum* and *P. reichenowi* diverged 6 million

years ago. However, based on our new estimate for *P. falciparum*–*P. reichenowi* divergence, we

recalibrated their estimate of the most recent common ancestor of the *eba-175* alleles to be around

4,000 years ago, which is in good agreement with our divergence time for *P. falciparum*

(Supplementary Note 1). The recent dimorphism cannot however explain the observation of an

ancient dimorphism near the human and ape loci for glycophorin[18] – an EBA-175 binding protein.

The formation and maintenance of all of these dimorphic loci has therefore been shaped by different

balancing selection pressures over time.

***P. falciparum*-specific evolution**
151
During its move away from gorillas, *P. falciparum* had to adapt to new environmental

conditions, namely a new vertebrate host (human) and new vector species (e.g. *Anopheles*

gambiae)[19]. To infer *P. falciparum* specific adaptive changes, we considered the *P. falciparum* /

P. praefalciparum /*P. reichenowi* genome trio and then applied two lineage based tests to find

positive selection that occurred in the *P. falciparum* branch (see methods). The two tests identified

172 genes (out of 4,826) with signatures of positive selection in the human parasite species only

(Supplementary Table 3). Two genes (*rop14* and PF3D7_0609900) were significant in both tests.

Among the 172 genes, almost half (n=82) encoded proteins of unknown function. Analysis of those

with functional annotation indicated that genes involved in pathogenesis, entry into host, actin

movement and organization and in drug response were significantly over-represented. Other genes,

expressed in different stages of the *P. falciparum* life cycle (e.g. *sera4* and *emp3*, involved during

the erythrocytic stages; *trsp* and *lisp*1, involved in the hepatic stages; and *plp4*, *CelTOS* and

Cap380, involved in the mosquito stages) also showed a significant signal of adaptive evolution

(Supplementary Table 3).

Evolution through introgression, gene transfer and convergence

Frequent mixed species infections in apes and mosquitoes[19] provide clear opportunities for

interspecific gene flow between these parasites. A recent study[6] reported a gene transfer event

between *P. adleri* and the ancestor of *P. falciparum* and *P. praefalciparum* of a region on

chromosome 4 including key genes involved in erythrocyte invasion (*rhr5* and *cyrpA*). Because

such events preserve the phylogenetic history of the genes involved, we systematically examined

the evidence for introgression or gene transfer events across the complete subgenus by testing the

congruence of each gene tree to the species tree for genes with one-to-one orthologues. Beyond the

region that includes *rh5* (Fig 2b, Supplementary Fig. 4a), few signals of gene flow between

parasites infecting the same host species were obtained (n=11) suggesting that these events were

rare or usually strongly deleterious (Supplementary Fig. 5).

The *Laverania* subgenus evolved to infect chimpanzees and gorillas but, on a genome-wide

scale, the convergent evolution of host-specific traits has not left a signature (Supplementary Note

2). We therefore examined each CDS independently and were able to identify genes with

differences fixed within specific hosts, falling into three categories: 53 in chimpanzee-infective

parasites, 49 in gorilla-infective and 12 with fixed traits in both host species (Fig. 2; Supplementary

Table 4a). For at least 67 genes, these differences were unlikely to have arisen by chance (p <0.05)

and GO term enrichment analysis revealed that several of these genes are involved in host invasion

and pathogenesis (Supplementary Table 4b) including *rh5* (which has a signal for convergent

evolution even when the introgressed tree topology is taken into consideration; Supplementary Fig.

4b). *Rh5* is the only gene identified in *P. falciparum* that is essential for erythrocyte recognition

during invasion, via binding to Basigin. *P. falciparum rh5* cannot bind to gorilla Basigin and binds

poorly to the chimpanzee protein[20]. We notice that one of the convergent sites is known to be a

binding site for the host receptor Basigin[21] (Supplementary Fig. 4b). The gene *eba-165* encodes a

member of the erythrocyte binding like (EBL) super family of proteins that are involved in

erythrocyte invasion. Although *eba-165* is a pseudogene in *P. falciparum*[22], it is not a pseudogene

in the other *Laverania* species and may therefore be involved in erythrocyte invasion, like other

EBL members. The protein has three convergent sites in gorillas. One falls inside the F2 region, a

domain involved in the interactions with erythrocyte receptors. The role of this protein and of these

convergent sites in the invasion of gorilla red cells remains to be determined. Finally, genes

involved in gamete fertility (the 6-cysteine protein P230) or implicated in *Plasmodium* invasion of

erythrocytes (doc2[23]) also displayed signals of convergent evolution. Twelve parasite coding

sequences had fixed differences at the same amino-acid position in chimpanzees and gorillas. Of

these P230 was the only one found with a position that was different and fixed across all three host

species. P230 is involved in gamete development and trans-specific reproductive barriers[24], possibly

through enabling male gametes to bind to erythrocytes prior to exflagellation[25]. Host-specific

residues observed in P230 might affect the efficiency of the binding to the erythrocyte receptors and

result from co-evolution between the parasite molecule and the host receptor.

Subtelomeric gene families

To date, the only in-depth data on the subtelomeric gene families of the *Laverania* have come from *P. reichenowi* and *P. falciparum*. As a result of using long read sequencing, these families are well represented in our assemblies (Supplementary tables 2 and 5A). We provide for the first time a comprehensive picture of the evolution of these important families.

Most gene families were likely present in the ancestor of all Laverania. The same general pattern of one-to-one orthology throughout the subgenus indicates that many underwent gene duplication early (e.g. FIKK) or prior (e.g. ETRAMP, PHIST and SURFIN) to the development of a distinct Laverania lineage. Only a subset displayed contractions or expansions between specific Laverania species (Fig. 3 and Supplementary Table 5a, Supplementary Figure 7). For these latter families, Clade A and most species of Clade B clearly differ in their composition. *P. blacklocki* (Clade B) is intermediate in its composition. Some gene families, like the group of exported proteins *hyp4, hyp5*, *mc-2tm* and *EPF1*, have expanded only in *P. praefalciparum* and *P. falciparum* (and even more in *P. falciparum* for *hyp4* and *hyp5*). Since all four are components of Maurer's clefts, an organelle involved in protein export[26], some evolution of function in this organelle may have been an important precursor to human infection. The family of acyl-CoA synthetase genes, reported to be expanded and diversified in *P. falciparum*[27] is in fact expanded across the *Laverania* and has four fewer copies in *P. falciparum* (Supplementary Fig. 6). Other genes that show clade or group specific expansion include DBLmsp, glycophorin binding protein and CLAG (Supplementary Fig. 7).

One striking inter-clade difference concerns the largest gene family that is likely common to all other malaria species: the *Plasmodium* interspersed repeat family (*pir,* which includes the *rif* and *stevor* families in *P. falciparum*) (Fig. 3, 4). This family has been proposed to be involved in important functions such as antigenic variation, immune evasion, signalling, trafficking, red cell rigidity and adhesion[28] and yet has expanded only in Clade B, after the *P. blacklocki* split (Fig 3). The *rif* genes comprise a small conserved group and a much larger group of more diverse members that contains just 13 genes from Clade A species and at least 180 members per Clade B species (Fig. 4). There is however no evidence for host-specific adaptation in these sequences.

In contrast, a subset of *stevor* genes showed strong host-specific sequence diversification (Fig 4 and Supplementary Fig. 8). Based on full-length alignments, there is a deep phylogenetic split between *stevor* genes but when only short conserved protein motifs are considered, a group of Stevor proteins (*stevor II,* Fig 4a) forms a cluster comprising almost entirely of members from

238 gorilla-infecting species. Since *stevor* genes are known to be involved in host–parasite interactions
239 (such as binding to host glycophorin C in *P. falciparum*[29]), this host specific sequence may reflect
240 sequence differences in host-specific factors in gorillas.
241
Evolution of *var* genes
The *var* genes, crucial mediators of pathogenesis and the establishment of chronic infection through
cytoadherence and immune evasion, are the best studied *P. falciparum* multi-gene family and
unique to the *Laverania*[30]. They are two-exon genes and their products have three types of major
domain; exon 1 encodes Duffy Binding like (DBL) and Cysteine Rich Interdomain Regions (CIDR)
and exon 2 encodes Acidic Terminal Sequence (ATS)[31]. Similar to *P. falciparum*, our data are
consistent with all *Laverania* species having *var* genes (Fig. 3) that retain a two-exon structure and
are organized into subtelomeric or internal *var* gene clusters. There are however three notable
features of the evolution of this family within the sub-genus.
First, there is a deep division in how the repertoire is organised between the major clades. The
var genes of Clade B parasites, with the exception of *P. blacklocki*, resemble those of *P. falciparum*
in terms of genomic organisation, domain types and numbers (Fig 5, Supplementary Table 6). In
contrast, the repertoires of Clade A parasites and *P. blacklocki* (treated as one group hereafter in
this section) differ in their domain composition, contain a novel CIDR-like domain (CIDRn, Fig 5a,
Supplementary Fig. 9) and have lower sequence diversity per domain but cluster into more sub-
groups than Clade B domains (Fig 5b, Supplementary Fig. 10). The paucity of domains similar to
those in *P. falciparum* (such as CIDRα) that are involved in cytoadherence to some specific and
common host receptors, means that if endothelial cytoadherence was important in Clade A, some
alternative receptors must have been utilised.
Second, in total there are 10 internal *var* gene clusters (confirmed by contiguous sequence data)
but 8 are oppositely oriented between the two clades (Supplementary Fig. 11, Supplementary Table
7). Clade B parasites also show a much greater number of associated GC-rich RNAs of unknown
Function (RUF) elements than Clade A (Supplementary Table 7).
Third, the ATS domains cluster tightly within Clade A. Within Clade B there is clear evidence
of species-specific diversification, except in *P. praefalciparum* and *P. falciparum* reflecting their
recent speciation. There is one intact ATS from *P. falciparum* as well as several pseudogenes that
cluster with Clade A (Fig 5c). Moreover, of seven internal *var* arrays (Supplementary Fig. 11) in
P. falciparum, containing a functional *var* gene, five terminate with one of these pseudogenes (on
the opposite DNA strand) suggesting that they may be remnants from ancient rearrangements. The

intact *P. falciparum* gene is *var2csa*, a *var*-like gene that is highly conserved between *P. falciparum*

isolates[32], involved both in cytoadherence in the placenta in primigravidae, and proposed to be a

central intermediate in *var* gene switching during antigenic variation[33]. We therefore propose

var2csa is a remnant of an ancient multigene family that has been maintained as a single complete

gene in *P. falciparum*, for the dual purposes of *var*-switching and placental cytoadherence.

There is other evidence of retention of ancient *var* gene sequence across the subgenus. First,

in Clade B we find a nearly full length *var* pseudogene that has highest similarity to *P. adleri* and *P.*

gaboni var genes, within an internal *var* cluster on chromosome 4 in *P. falciparum* and *P.*

praefalciparum but on the opposite strand to the other *var* genes. It is found in all *P. falciparum*

isolates, but not in *P. reichenowi*. Second, in *P. gaboni* and *P. adleri,* three genes have the N-

terminal DBLα/CIDRα architecture typical of Clade B genes and their domains cluster within Clade

B based on similarity (Fig. 5b, larger nodes). Directly adjacent to two of these *var* genes are two *rif*

pseudogenes that also show greatest similarity to those from Clade B. Last, we find a further nine

rif pseudogenes of Clade A parasites that cluster with Clade B *rif* genes (Fig. 4). If these

observations reflect retention of ancient copies, their high sequence conservation suggests that they

are under extremely unusual selection pressure. Alternatively, they may represent relics of gene

transfer between species that occurred after the Clade A/B split.

Conclusion

We have produced high quality genomes and used mutation rates and generation times, covering the

full range of most recent estimates, to calculate the date of speciation for all known members of the

Laverania, with only a small margin of error. In our analysis, we have shown that the successful

infection of humans by *P. falciparum* occurred quite recently and involved numerous parasites

rather than a single one as previously proposed. After the establishment in its new host, the parasite

population went through a bottleneck around 5,000 years ago during the period of rapid human

population expansion due to farming (Fig. 1b). We summarise the major genomic events during the

evolution of the *Laverania* in Fig. 6.

As a result of our analyses we propose the following series of events for the emergence of *P.*

falciparum as a major human pathogen. First, the crucial lateral transfer event of the *rh5* locus

between Clade A and B parasites may also have involved *var* and *rifin* genes in other parts of the

genome that, because of their orientation on the opposite strand, were not lost during later

recombination. Next, facilitatory mutations are likely to have occurred in *rh5* that in the first

instance allowed invasion of both gorilla and human red cells. Modern humans emerged more than

300,000 years ago[34] and existed as small isolated populations[12]. Our evidence suggests that *P.*

falciparum and *P. praefalciparum* started to diverge around 40,000-60,000 years ago. In the

following 40,000 years with low population densities in humans and gorillas there would have not

been high selection pressure to optimise infectivity in either the hosts or vectors, enabling at least

some movement of parasites between hosts. We find evidence for gene flow between lineages

throughout this period. The expansion of the human population with the advent of farming likely

led to strong evolutionary pressure for mosquito species (specifically *An. gambiae*) to feed

primarily on humans[35]. Therefore, the existing human infective (*P. falciparum)* genotypes would be

selected for human and appropriate vector success and the fittest would rapidly expand. Subsequent

rapid accumulation of mutations that favoured growth in humans, and in the anthropophilic vectors

such as *An. gambiae*, are likely to have occurred to increase human-specific reproductive success.

The resulting specific parasite genotypes that expanded (and appeared as an emergence from a

bottleneck), would have had a much lower probability of a direct transfer back to apes. With

experiments on gorillas and chimpanzees not possible it will be difficult directly to prove the

precise combination of different alleles that allowed the emergence of *P. falciparum*. However,

existing data suggest that the genes that we have implicated in this process are expressed throughout

the life cycle but that half remain uncharacterised (www.genedb.org, www.plasmodb.org), opening

up new opportunities for future studies on host specificity and host adaptation in *Plasmodium*.

Online Methods

Sample collection

All but two infected blood samples from chimpanzees (*Pan troglodytes troglodytes*) and

gorillas (*Gorilla gorilla gorilla*) were obtained from the sanctuary "Parc de La Lékédi", Bakoumba

(Haut-Ogooué, Gabon), during routine sanitary controls of the animals. This park holds various

primate species, including gorillas, chimpanzees and monkeys (*Cercopithecinae*), that have been

orphaned due to bushmeat-poaching activities and have been confiscated by the Gabonese

Government, quarantined at the Centre International de Recherches Médicales de Franceville

(CIRMF, Gabon) and finally released into semi-free ranging enclosures in the sanctuary. Every six

months, chimpanzees (12 individuals) and gorillas (2 individuals) are anesthetized for medical

check up. Blood samples were collected from the animals during sanitary controls (July 2011,

September 2012, May 2013 and December 2013). Two additional infected blood samples were

obtained from gorilla orphans (GG05, GG06) seized by the Gabonese government in 2011 and 2013

and sent to the CIRMF for a quarantine before being released in a sanctuary. All animal work was

conducted according to relevant national and international guidelines. From each animal, 15 ml of

whole blood were collected in EDTA tubes. For all samples but three, white blood cell depletion

was performed on 10 ml of the freshly collected samples using cellulose columns as described in [36].

Remaining blood was subsequently used for DNA extraction and detection of *Plasmodium*

infections as described in Ollomo et al[3]. Overall, 15 blood samples from 7 chimpanzees and 4

gorillas were found to contain the *Laverania* samples used in the present study (Table 1).

Sample preparation

Three methods were used for DNA amplification prior to sequencing (Supplementary Table 1).

For all but one sample, whole genome amplification (WGA) was performed with a REPLI-g Mini

Kit (Qiagen) following a modified protocol[37] to enrich genomic DNA. The genome of *P. blacklocki*

was generated using selective WGA (sWGA) as indicated in[38] using 20 primers, followed by a

WGA. Finally, for the PprfG03 (a *P. praefalciparum* isolate) and PadlG02 (a *P. adleri* isolate)

samples, we used a cell sorting approach[39].

Sample sequencing

All samples were first sequenced with Illumina. Amplification-free Illumina libraries of 400-

600 bp were prepared from the enriched genomic DNA[40] and run on MiSeq and HiSeq 2000 (v3

chemistry) Illumina machines.

After the Illumina sequencing, six samples with a combination of the least number of multiple

infections (see below) and the lowest level of host contamination were chosen for long read

sequencing, using Pacific Biosciences (PacBio). The DNA of the samples (after WGA) was size-

selected to 8 kb and sequenced with the C3/P5 chemistry. The number of SMRT cells (Pacific

Bioscience sequencing runs) used varied between samples (Supplementary Table 1).

Genome assembly, genome QC, split of infection & annotation

Determination of multiple infections

To initially quantify multiple infections and so allow samples to be selected for PacBio

sequencing from those comprising a low number of species, Illumina reads from each sample were

mapped against a concatenation of all available *Cox 3* and *CytB* genes of the *Laverania* from NCBI,

using SNP-o-Matic[41] (parameter chop=5) to position reads only where they aligned perfectly. SNP-

o-Matic returns all the positions of repetitive mapping reads. This output allowed us to count the

read depth of these two genes across all species and therefore determine the number and relative

amount of different malaria species per sample.

Whole genome amplification (WGA) bias

The uneven coverage that resulted from WGA bias, host contamination and multiple infections

presented a challenge for sequence assembly. To overcome the bias and the host contamination,

each DNA sample was sequenced deeper than normally necessary. Lower coverage of the

subtelomeres was obtained for the sWGA sample (*P. blacklocki*) meaning that the subtelomeres in

that assembly were not as complete as those in the assemblies for other species.

Long reads (Pacific Bioscience) assemblies

Six reference genomes were assembled using HGAP[42], with different settings for the genome

size parameter, ranging from 23 Mb (*P. reichenowi*) to 72 Mb (*P. billcollinsi*). This parameter

encodes how many long reads are corrected for use in the assembly and depends on the host

contamination and the amount of different isolates in the samples. The obtained contigs from

HGAP were ordered with ABACAS[43] against a *P. falciparum* 3D7 reference that has no

subtelomeric regions. Assembly errors and WGA artefacts were manually corrected using ACT[44].

After this step, three iterations of ICORN2[45] were run, followed by another ABACAS step,

allowing overlapping contigs to be merged (parameter: ABA_CHECK_OVERLAP=1). For the

PrG01, PgabG01 and PadlG01 assembly, we also ran PBjelly to close some of the sequencing

gaps[46].

Host decontamination

To detect and remove sequence data derived from host DNA, contigs were compared with the

chimpanzee or gorilla genomes using BLAST. Contigs were considered as host contamination if

more than 50% of their BLAST hits had higher than 95% identity to any of the great ape genomes.

Unordered contigs with a GC content >32% were searched against the non-redundant nucleotide

database, to detect and remove further contaminants.

Resolving multiple infections

The first assembled genome was a single *P. reichenowi* infection, PrG01. We detected low

levels of *P. vivax-like* and virus contamination (TT virus, AB038624.1), which were excluded. For

quality control, the assembly was compared against the existing PrCDC[9] reference genome. The

number of *Plasmodium* interspersed repeats (PIRs) was similar, and there were no breaks in

synteny. There were however significantly fewer sequencing gaps and 17 Rep20 regions could be

found (a known repeat close to the telomeres in *P. falciparum)*. Thus, the assembly of PacBio data

(PrG01; Supplementary Table 2) appears to be of higher quality than the existing *P. reichenowi*

PrCDC reference.

The *P. adleri* sample comprised a single infection. Because a large number of cycles of

amplification were used, a greater number of SMRT cells were sequenced (Supplementary Table 1)

to overcome the problem of uneven coverage resulting in under-represented regions. An estimated

genome size of 60 Mb was chosen for the HGAP analysis to ensure that all regions were covered.

PgabG01 was a *P. gaboni* isolate with a *P. vivax-like* co-infection. To detect contigs of *P.*

vivax, unordered contigs (those that could not be placed against Pf3D7 using ABACAS) were

searched against the protein sequences of *P. falciparum* 3D7 and the *P. vivax* PvP01 reference

genome using TBLASTx. For each contig, the relative number of genes hitting against the two

genomes was used to assign it to *P. gaboni* or *P. vivax*. In most cases, all genes for a given contig

consistently hit only one genome so that the attribution to either species was clear. Overall, 14 Mb

of *P. vivax-like* sequences were obtained that will be described elsewhere.

The *P. billcollinsi* genome (PbilcG01) was obtained from a co-infection with a *P. gaboni*

genome (PgabG02). Rather than ordering the contigs just against Pf3D7 with ABACAS, contigs

were ordered against a combined reference comprising *P. gaboni* (PgabG01) and the Pf3D7

(parameters: overlap 500 bp, identity 90%). The species designation of contigs was confirmed with

a TBLASTx searches of annotated genes against a combination of the proteomes of PgabG01 and

PrCDC. For subtelomeric gene families, contigs were attributed to species if the hit was significant

for one species, not the other. Some of the contigs could not be attributed unambiguously and were

discarded. Due to sequencing gaps, some of the core genes are missing from the final assembly.

The sample used to produce the *P. praefalciparum* genome (PprfG01) had a high level of host

contamination, a low level of co-infection with *P. adleri* and contained two distinct

P. praefalciparum genotypes. For the core genome, we used iCORN to identify and utilise the

dominant genotype at each position. Where it was not possible to phase the genotypes, due to a lack

of variation, we assumed that they were identical. In the subtelomeres however, it was possible to

distinguish but not phase the two *P. praefalciparum* genotypes resulting in approximately twice the

number of *var* genes as seen in *P. falciparum*. Due to contamination of construction vectors (*E.*

coli) and host, 29 SMRT cells were sequenced and the HAGP parameter for the assembly size was

set to 60 Mb. Contigs were screened against *P. adleri* and *P. falciparum* to exclude a *P. adleri* co-

infection. All of the contigs that had a *P. falciparum* BLAST hit or had no clear hit (such as those

containing species-specific gene families) were attributed to the *P. praefalciparum* assembly. Last,

all samples (Supplementary Table 1) including five *P. falciparum* genomes were mapped against

the Pf3D7, *P. praefalciparum and P. adleri* assemblies. Contigs were excluded where more

normalized hits to the three *P. adleri* samples were found than to one of the two other *P.*

praefalciparum samples. Similarly, this method was used to eliminate the remote possibility that

any of the contigs in the *P. praefalciparum* assembly were in fact derived from *P. falciparum* co-

infection.

The *P. blacklocki* sample was from a single infection. Due to sWGA, the PacBio sequence data

covered regions not covered by Illumina but due to the bias of the primers, the subtelomeres were

not covered fully. However, the internal *var* gene clusters are all assembled. Some of the core genes

from this species are also missing.

Annotation

The genomes were annotated as described in[47]. In short, the annotation of *P. falciparum*

(version July 2015) was transferred with RATT[48] and new gene models were called with

Augustus[49]. Obvious structural errors in core genes were manually corrected in Artemis[50].

Mapping - generation of further samples

To generate the gene sequence for different samples, Illumina reads were mapped against a set

of reference genomes using BWA[51] and default parameters. For the gorilla samples, we mapped

against the combined PacBio reference genomes of *P. adleri*, *P. blacklocki* and *P. praefalciparum*

and for the chimpanzee samples, the combined references of *P. gaboni* (PgabG01), *P. billcollinsi*

and *P. reichenowi* (PrG01). SNPs with Phred score ≥ 100 were called using GATK

UnifiedGenotyper[52] v2.0.35 (parameters: -pnrm POOL -ploidy 2 -glm POOLBOTH). From these

SNP calls we constructed the new gene set, masking regions in genes with less than 10x coverage

of 'properly' (correct distance and orientation) mapped paired reads. To generate the sequences of

the other 13 isolates, homozygous SNP calls were obtained (consensus program from bcftools-

1.2[53]). We quality controlled the SNP calling by regenerating PrCDC and PgabG02 gene set from

PrG01 and PgabG01, respectively and confirmed that they were placed with nearly no differences

in a phylogenetic tree.

Orthologous group determination and alignment

Orthologous groups were identified using OrthoMCL v1.4[54] across: (i) the seven core

Laverania genomes; (ii) the seven core genomes, the *Laverania* isolates PgabG02, PrCDC and *P.*

falciparum IT, as well as two outgroup genomes *Plasmodium vivax* Sal1 and *Plasmodium knowlesi*

strain H; and (iii) just Pf3D7, PprfG01 and PrG01. *P. praefalciparum* II was excluded due to its

partial genome. From these groups, different complete sets of 1:1 orthologues were extracted:

        (1) "Lav12sp" set of 3,369 orthologues across the seven core *Laverania* species, the PrCDC

and *P. falciparum* IT isolates, *P. vivax* and *P. knowlesi*

(2) "Lav25st" set of 424 1:1 orthologues from across the 25 *Laverania* isolates, including the

previously published *P. reichenowi* CDC and five *P. falciparum* isolates (3D7, IT, DD2,

HB3 and 7G8[9]).

(3) "Lav7sp" set of 4,350 orthologues from across the seven *Laverania* reference genomes

(4) "Lav15st" set of 3,808 orthologues, with two representative sequences per species,

excluding *P. blacklocki* and *P. praefalciparum*.

(5) "Lav3sp" set of 4,826 1:1 orthologues across all the *P. reichenowi*, *P. praefalciparum* and

P. falciparum isolates

The first two sets were used to reconstruct the species tree, the third one for the comparative

genomic analyses (introgression, convergence and gene family evolution), the fourth one for the

analyses of within species polymorphism and the fifth one for the analysis of *P. falciparum*

adaptive evolution.

To reduce the rate of false positives in the evolutionary analyses due to misalignments (e.g.[55]),

codon-based multiple alignments were performed using PRANK[56,57] with the -codon and +F

options, as it was shown to outperform other programs in the context of the detection of positive

selection[58,59]. Prior to aligning codons, low complexity regions were excluded in the nucleotide

sequences using dustmasker[60] and in amino acid sequences using segmasker[61] from NCBI-BLAST.

Poorly aligned regions were excluded using Gblocks[62], with default settings.

Analysis of interspecific gene flow, introgression or gene transfer

Species-tree inference

Two ML trees were performed using RAxMLv8.1.20[63] to illustrate the phylogenetic

relationships between the *Laverania* species and genotypes studied here using the "Lav12sp" and

the "Lav25st" set of orthologues. For each tree, multiple nucleotide alignments of each orthologous

group were conducted as described above. Trees were then constructed from the concatenated

alignments of the "Lav12sp" set of orthologues for the species tree and the "Lav25st" set for the

strain tree using RAxML and the following options "-m GTRGAMMA –f a -# 100". Trees were

rooted afterwards using *P. vivax* and *P. knowlesi* for the species tree and the *P. adleri/P. gaboni*

clade for the genotype tree.

Tree topology test

Interspecific gene flow was investigated by testing congruence between each gene tree

topology and the species tree topology. We performed the Shimodaira-Hasegawa test (SH test[64])

using RAxMLv8.1.20 to test whether the phylogenetic tree for each gene significantly differed from

the *Laverania* species tree. Topology tests were based on multiple nucleotide alignments of the

4,350 "Lav7sp" set of orthologues. For each coding sequence, RAxML was called with the options

"-m GTRGAMMA –f h".

Convergent evolution analyses

Genome-wide test of convergent evolution

Convergent substitutions can occur by chance and the number of random convergent

substitutions between two lineages is correlated with the number of divergent substitutions

observed in these two lineages[65,66]. Excess of convergent substitutions in specific branch pairs can

thus be identified by analyzing the correlation between the number of convergent and divergent

substitutions between all the branch pairs in a phylogeny using orthogonal regression, and looking

for outlier branch pairs: branch pairs with a high positive residual show an excess of convergent

substitutions relatively to the number of divergent substitutions[65]. We used the software Grand-

Convergence (available at https://github.com/dekoning-lab/grand-conv) to estimate for each

chromosome the numbers of divergent and convergent substitutions between all branch pairs in the

Laverania tree and investigate whether branch pairs including *Laverania* species infecting the same

host species (gorilla or chimpanzee) presented an excess of convergence. Analyses were performed

under different models of amino-acid evolution: LG, WAG, JONES and DAYHOFF.

Gene-based detection of convergent evolution throughout the Laverania

For each orthologue of the "Lav7sp" set, the number and percentage of fixed amino acid

differences between parasites infecting the same host were calculated, *i.e.* the number of positions

showing the same amino acid within a host species but different amino acid between host species.

Alignments of all the available sequences ("Lav15st") from all the sequenced isolates were then

used to determine what number of host-specific differences were fixed within each host and each

species. To evaluate whether the observed number of host-specific fixed differences in an alignment

can be attributed to neutral evolution/purifying selection alone (with no positive selection), we used

a simulation-based approach. For each coding sequence, 1,000 sequences of the same size were

simulated, evolving along the same tree with the same specified branch lengths, substitution model,

codon frequencies and omega (d_N/d_S), using the program Evolver from PAML v4.8a[67]. The

program Codeml from PAML v4.8a[67] was first used to estimate the tree, the codon frequencies and

the average omega values for each of the coding sequences with fixed amino acid differences. For

each simulated dataset, the number of fixed amino-acid differences between the parasites infecting

a same host was estimated. The probability of observing *n* fixed differences was then computed as

the proportion of the simulated dataset of 1000 sequences that showed at least the same number of

fixed differences as observed in the real data.

Tests for positive selection

Branch site tests

To search for genes that have been subjected to positive selection in the *P. falciparum* lineage

alone after the divergence from *P. praefalciparum*, we used an updated branch site test [68]

implemented in PAML v4.4c [67]. This test detects sites that have undergone positive selection in a

specific branch of the phylogenetic tree (foreground branch). The "Lav3sp" set of 4,826

orthologous groups between *P. reichenowi*, *P. praefalciparum I* and *P. falciparum* was used for the

test. d_N/d_S ratio estimates per branch and gene were obtained using Codeml (PAML v4.4c) with a

free-ratio model of evolution. This identified 139 genes with a significant signal of positive

selection in *P. falciparum* only.

Using the "Lav7sp" dataset, a branch site test was also applied for each gene (Figure 2),

comparing the sequence from an individual species to the inferred sequence of its closest branch

point on the tree (e.g. *P. falciparum* compared with the sequence at the *P. falciparum*–

P. praefalciparum split).

McDonald–Kreitman (MK) tests

Selection in *P. falciparum* was also tested using McDonald–Kreitman (MK) tests [69] to compare

the polymorphism within species to the divergence between species, using *P. praefalciparum* as the

outgroup. Analyses were performed using the 4,826 "Lav3sp" set of orthologues. MK tests were

performed as described before[9]. Thirty-five genes had an MK ratio considerably higher than 1.

Gene Ontology enrichment analyses

Analysis of Gene Ontology (GO) term-enrichment was performed in R, using TopGO[70] with

default parameters. GO annotations from GeneDB were used but with unreviewed automated

annotations excluded.

Gene family analyses

To estimate the differential abundance of gene families across species, the Gene products and

the Pfam domains were counted and analysed by the variance of the occurrence. Unless otherwise

stated, trees were constructed using PhyML[71] (default parameters) or RAxML[63] (model estimated)

from alignments generated with Muscle[72] and trimmed with Gblocks[62] in Seaview[73] with default

values. Many of the findings were confirmed manually through ACT and bamview[50]. The analysis

of the *var* genes was performed on *var* genes larger than 2.5kb. Domains were called with the

HMMer models from varDom[74]. Distance matrices were generated based on BLASTp scores,

without filtering low complexity regions. Representation was done in R through the heatmap.2

program from gplot (see also Supplementary Note 3).

Allelic dimorphisms

For the analysis of dimorphism in *msp*, all sequences available for the *Laverania* were

downloaded from Uniprot[75]. Data were subsampled to obtain a similar number of sequences for

each group. Phylogenetic trees were constructed with PhyML[71], using default parameters and drawn

in Figtree. The *eba-175* alignment was visualized with Jalview[76].

Divergence Dating

Alignments of the *Laverania* included intergenic regions where possible. Assuming 402–681

mitotic events per year (Supplementary Note 1) and a mutation rate of 3.78E-10 for 4 mitotic

events[77,78](mutation rate from latter paper was taken from Pf3D7 line without drugs), equivalent to

around 0.9–1.5 mutations per genome per year. Although we observed similar mutation rates in

clinical samples (Supplementary Note 1), these estimates have potential errors and therefore we

report ratios of divergence times in the figures that are robust to errors in these parameters. For

coalescence based estimates of speciation times, G-Phocs[79] was used and multiple sequentially

Markovian coalescent (MSMC) on segregating sites[80] was used to estimate the *P. falciparum*

bottleneck.

Author Contributions: TDO, BO, FR, CN, MB, FP designed the study. CA, APO, LB, EW, BN,

ND, CP, PD, VR, FP collected and assessed samples. CA performed the WGA and cell sorting on

one sample. SO performed the WGA on the samples; MS organised the sequencing. TDO did

assembly and annotation. UB did manual gene curation; AG, FP performed the evolutionary

analyses on core genomes. TDO, CN, MB performed the analyses of gene families

and dimorphisms. TC performed the dating analyses. TDO, AG, CN, MB, FP wrote the manuscript.

All authors read and approved the paper.

Acknowledgments:

This work was funded by ANR ORIGIN JCJC 2012, LMI ZOFAC, CNRS, CIRMF, IRD and the

Wellcome Trust (grants WT 098051 and WT 206194 to the Sanger Institute, 104792/Z/14/Z to

CN). TC holds a MRC DTP Studentship. We thank Gavin Rutledge for performing the sWGA and

Julian Rayner and Francisco J. Ayala for helpful discussion. We thank the PlasmoDB team for

promptly making these data available.

Data availability

All sequences have been submitted to the European Nucleotide Archive. The accession numbers of

the raw reads, and assembly data can be found in Supplementary Table 8. The genomes are being

submitted to EBI, project ID PRJEB13584. The genomes are available from plasmodb.org and

from ftp://ftp.sanger.ac.uk/pub/project/pathogens/Plasmodium/Laverania/.

Competing financial interests.

None
628

629 **Computer code**

630 Custom computer code is available on request.

632
633

 Figures

 Figure 1. Overview of the dating of the evolution of the *Laverania*. (**a**) Maximum likelihood tree
 of the *Laverania* based on the "Lav12sp" set of orthologues. All bootstrap values are 100.
 Coalescence based estimates of the timing of speciation events are displayed on nodes (MYA -
 million years ago), based on intergenic and genic alignments. (**b**) Multiple sequentially Markovian
 coalescent estimates of the effective population size (*Ne*) in the *P. falciparum* and *P.*
 praefalciparum population. Assuming our estimate of the number of mitotic events per year, a
 bottleneck occurred in *P. falciparum* 4,000-6,000 years ago. The y-axis shows the natural logarithm
 (Ln) of *Ne*. Bootstrapping (pale lines) was performed by randomly resampling segregating sites
 from the input 50 times.

 Figure 2. Overview of the analyses of core genes over all *Laverania* genomes. (a) Summary of
 evolution of core genes. From outer to inner track: scatterplot of branch site test for each genome
 (see Supplementary Table 3 for *P. falciparum* data); per-species d_N/d_S values ($0.5 < d_N/d_S < 2$);
 orthologues represented by vertical black lines under the chromosome track represent, with dots
 representing *P. falciparum* 3D7 *var* genes on the forward (blue) or reverse strands (red), or *var*
 pseudogenes (black); average of the relative polymorphism (π) across species, with the underlying π
 for each species calculated from multiple strains ("Lav25st" dataset) and normalized by the average
 for that species; signatures of convergent evolution based on host-specific fixed differences analysis
 with the chromosome 4 region that includes the *Rh5* locus highlighted (black box). (b) Magnified
 view of the *Rh5* region that is enriched with host specific fixed differences.   Convergent evolution
 analysis was performed using orthologues conserved across the *Laverania*. Filled circles represent
 the subset of differences that were fixed within all the isolates available ("Lav15st" set) and for
 which we could reject neutral evolution (for the gene list see Supplementary Table 4).

 Figure 3. Gene families in the *Laverania*. Distribution of major multigene families including *var*
 and those that show significant copy number variation among lineages. Data from
 P. praefalciparum include the subtelomeric gene families from the two infecting genotypes.
 Assembly of *P. billcollinsi* is incomplete in the subtelomeres.

Figure 4: Clustering of Pir (Rifin and Stevor) proteins families. Graphical representation of similarity between all *pir* proteins > 250aa, coloured by species. A BLAST cut-off of 45% global identity was used (see methods). More connected genes are more similar. Black circles highlight Clade A rifin proteins that cluster with Clade B rifin proteins.

Figure 5. Evolution of *var* gene domains in the *Laverania* (a) Heatmap of numbers of *var* gene domains in each *Laverania* species. Duffy represents regions closest to the Pfam Duffy binding domain. CIDRn is a new domain discovered in this study in Clade A. Only domains from *var* genes longer than 2.5 kb were considered. Heat map colours blue-yellow-white indicate decreasing copy numbers. **(b)** Graphical representation of similarity between domains, using domains from *var* genes longer than 2.5kb. Domains are coloured by species and clustered by a minimum BLAST cut-off of 45% global identity. Larger circles denote *var* genes in the opposite orientation. **(c)** Maximum likelihood trees of the Acidic Terminal Sequence (ATS). Apparent ATS sequences from clade A that cluster with clade B are indicated (**).

Figure 6. Overview of the genomic evolution of the *Laverania* subgenus. The values of polymorphism (π) within the species are indicated by triangles of different size at the end of the tree branches, as well the bottleneck in *P. falciparum* (constricted branch width), ~ 5,000 years ago. Also shown are the gene transfers that occurred between certain Clade A and B species and the huge genomic differences that accumulated in Clade B after the divergence with *P. blacklocki*.

Species	Sample ID	Primate species	Assembly size Mb	Contigs	Scaffolds	Genes	Primary co-infection [b]
P. praefalciparum	**PprfG01**	**Chimp**	**26**	**142**	**73**	**6,476**	***P. adleri***
	PprfG02	Gorilla	-	-	-	-	none
	PprfG04	Co-infection, see PadlG03					*P. adleri*
	PprfG03	Gorilla	-	-	-	-	none
P. reichenowi	**PrG01**	**Chimp**	**24.5**	**66**	**48**	**5,941**	***none***
	PrCDC [a]	Chimp	24.1	374	2,465	5,895	none
	PrG02	Chimp	-	-	-	-	none
	PrG03	Chimp	-	-	-	-	none
P. billcollinsi	**PbilcG01**	**Co-infection PgabG02**	**23.1**	**306**	**89**	**5,637**	***P. gaboni***
	PbilcG03	Co-infection, see PgabG04					*P. gaboni*
	PbilcG04	Co-infection, see PgabG05					*P. gaboni*
P. blacklocki	**PblacG01**	**Gorilla**	**22**	**311**	**97**	**5,346**	**none**
P. gaboni	**PGAB01**	**Chimp**	**21.8**	**121**	**96**	**5,421**	***P. vivax***
	PgabG02	**Chimp**	**20.9**	**76**	**44**	**5,249**	***P. billcollinsi***
	PgabG03	Chimp	-	-	-	-	*P. vivax*
	PgabG04	Chimp	-	-	-	-	*P. billcollinsi*
	PgabG05	Chimp	-	-	-	-	*P. reichenowi*
P. adleri	**PadlG01**	**Gorilla**	**22.2**	**102**	**82**	**5,515**	**none**
	PadlG02	Gorilla	-	-	-	-	none
	PadlG03	Gorilla	-	-	-	-	*P. praefalciparum*

[a] Data from Otto *et al.* [9]

[b] Based on percentage of reads mapping to the reference for each *Laverania* species (see
Supplementary Table 1)

Table 1: Overview of all *Laverania* samples used in study. All samples generated for this study,
except PrCDC, were sequenced using Illumina technology with a range of read lengths from 100–
250 bp. For isolates in bold, assemblies were produced using long-reads from Single-Molecule
Real-Time sequencing (Pacific Biosciences) and the total assembled size, number of contigs,
scaffolds and predicted genes are shown. Fold coverage is reported based on mapping reads to
the *P. falciparum* 3D7 reference genome (v3.1). *P. billcollinsi* sequences were obtained from *P.*
gaboni co-infections, of which some also harboured *P. reichenowi* species.

Population	Status	Coalescence, years (x 10^5)			Effective Population size, Ne (x 10^5)			Migration target	Probability of Parasite Migration		
		CDS	Intergenic	Intergenic no UTR	CDS	Intergenic	Intergenic no UTR		CDS	Intergenic	Intergenic no UTR
P. falciparum	Present	NA	NA	NA	1.9 - 3.2 (1.8 - 3.4)	0.4 - 0.7 (0 - 2.1)	0.4 - 0.7 (0 - 2.1)	NA	0	0	0
P. praefalciparum	Present	NA	NA	NA	8.8 - 14.8 (8.1 - 16)	12.2 - 20.7 (11.6 - 21.8)	10 - 17 (9.2 - 18.4)	Pf	0.02 (0.0 - 0.03)	0.14 (0.09- 0.19)	0.16 (0.08- 0.27)
P. reichenowi	Present	NA	NA	NA	11.3 - 19.2 (3.7 - 19.9)	15.9 - 26.9 (15.4 - 27.7)	12.3 - 20.7 (11.7 - 21.8)	NA	0	0	0
P. billcollinsi	Present	NA	NA	NA	9.3 - 15.8 (8.9 - 16.5)	-	-	NA	0	-	-
P. gaboni	Present	NA	NA	NA	18.1 - 30.6 (17.6 - 31.6)	-	-	NA	0	-	-
P. adleri	Present	NA	NA	NA	17.7 - 29.9 (16.9 - 31.2)	-	-	NA	0	-	-
Pf - Pprf	Ancestral	0.2 - 0.3 (0.1 -0.3)	0.5 - 0.8 (0.4 - 0.8)	0.4 - 0.6 (0.3 - 0.7)	6.5 - 11.1 (6 - 11.9)	16.5 - 27.9 (15.4 - 29.7)	12.4 - 21 (11.1 - 23.2)	NA	0	0	0
Pf - Pprf - Pr	Ancestral	0.9 - 1.5 (0.8 -1.5)	1.6 - 2.7 (1.5 - 2.7)	1.4 - 2.3 (1.3 - 2.4)	29.7 - 50.3 (27.7 - 53.7)	68.6 - 116.2 (65.4 - 121.6)	42.3 - 71.5 (39.1 - 77.1)	NA	0	0	0
Pf -Pprf - Pr-Pbilc	Ancestral	2.3 - 4 (2.3 - 4)	4.2 - 7.1*	3.7 - 6.2*	65.8 - 111.3 (62 - 118.5)	-	-	Pgab -Padl	0.605 (0.45 - 0.77)	-	-
Pgab - Padl	Ancestral	0.9 - 1.4 (0.8 -1.5)	1.5 - 2.6*	1.3 - 2.2*	28.3 - 47.9 (26.7 - 50.7)	-	-	NA	0	-	-
All species	Ancestral	4.5 - 7.6 (4.2 - 8)	8.1 -13.6*	7 - 11.8*	98.9 - 167.4 (93.2 - 178.4)	-	-	NA	0	-	-

Table 2: Estimated dates of speciation (time to coalescence), effective population size (Ne) and migration rates for present and ancestral *Laverania* populations. Estimated dates of speciation (time to coalescence), effective population size (Ne) and migration rates for Laverania populations, including inferred ancestors. Values inferred using GPhoCS coalescence model. The results are presented from sequence alignments of protein coding (CDS) (genic) regions as well as apparent intergenic regions inferred in the presence or absence of 0.5 kb flanking (UTR) sequences. The coalescence model parameters have been scaled based on an assumption of 401–681 mitotic events per year (Supplementary Note 1). The number in parentheses is the range based on the prediction (95% confidence interval). Values with (*) are linear estimates, as good alignments could not be obtained between those species due to the low GC content. The migration rate refers to the probability of a parasite from the source lineage migrating into the target over the time period. The data from these estimates are presented in Fig. 1.

References

Prugnolle, F. *et al.* African great apes are natural hosts of multiple related malaria species, including Plasmodium falciparum. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 1458-1463, doi:10.1073/pnas.0914440107 (2010).

Liu, W. *et al.* Origin of the human malaria parasite *Plasmodium falciparum* in gorillas. *Nature* **467**, 420-U467, doi:10.1038/nature09442 (2010).

Ollomo, B. *et al.* A New Malaria Agent in African Hominids. *PLoS Pathog* **5**, e1000446, doi:10.1371/journal.ppat.1000446 (2009).

Liu, W. *et al.* Multigenomic Delineation of Plasmodium Species of the Laverania Subgenus Infecting Wild-Living Chimpanzees and Gorillas. *Genome biology and evolution* **8**, 1929-1939, doi:10.1093/gbe/evw128 (2016).

Boundenga, L. *et al.* Diversity of malaria parasites in great apes in Gabon. *Malar J* **14**, 111, doi:10.1186/s12936-015-0622-6 (2015).

Sundararaman, S. A. *et al.* Genomes of cryptic chimpanzee Plasmodium species reveal key evolutionary events leading to human malaria. *Nature communications* **7**, 11078, doi:10.1038/ncomms11078 (2016).

Silva, J. C., Egan, A., Arze, C., Spouge, J. L. & Harris, D. G. A New Method for Estimating Species Age Supports the Coexistence of Malaria Parasites and Their Mammalian Hosts. *Molecular biology and evolution* **32**, 1354-1364, doi:10.1093/molbev/msv005 (2015).

Volkman, S. K. *et al.* Recent origin of Plasmodium falciparum from a single progenitor. *Science* **293**, 482-484, doi:10.1126/science.1059878 (2001).

Otto, T. D. *et al.* Genome sequencing of chimpanzee malaria parasites reveals possible pathways of adaptation to human hosts. *Nature communications* **5**, 4754, doi:10.1038/ncomms5754 (2014).

Larremore, D. B. *et al.* Ape parasite origins of human malaria virulence genes. *Nature communications* **6**, 8368, doi:10.1038/ncomms9368 (2015).

Pacheco, M. A. *et al.* Timing the origin of human malarias: the lemur puzzle. *BMC evolutionary biology* **11**, 299, doi:10.1186/1471-2148-11-299 (2011).

Behar, D. M. *et al.* The dawn of human matrilineal diversity. *American journal of human genetics* **82**, 1130-1140, doi:10.1016/j.ajhg.2008.04.002 (2008).

Palstra, F. P. & Fraser, D. J. Effective/census population size ratio estimation: a compendium and appraisal. *Ecology and evolution* **2**, 2357-2365, doi:10.1002/ece3.329 (2012).

Chang, H. H. & Hartl, D. L. Recurrent bottlenecks in the malaria life cycle obscure signals of positive selection. *Parasitology* **142 Suppl 1**, S98-S107, doi:10.1017/S0031182014000067 (2015).

Roy, S. W. The Plasmodium gaboni genome illuminates allelic dimorphism of immunologically important surface antigens in P. falciparum. *Infection, genetics and evolution : journal of molecular epidemiology and evolutionary genetics in infectious diseases* **36**, 441-449, doi:10.1016/j.meegid.2015.08.014 (2015).

Tanabe, K., Mackay, M., Goman, M. & Scaife, J. G. Allelic dimorphism in a surface antigen gene of the malaria parasite Plasmodium falciparum. *Journal of molecular biology* **195**, 273-287, doi:10.1016/0022-2836(87)90649-8 (1987).

Yasukochi, Y., Naka, I., Patarapotikul, J., Hananantachai, H. & Ohashi, J. Genetic evidence for contribution of human dispersal to the genetic diversity of EBA-175 in Plasmodium falciparum. *Malar J* **14**, 293, doi:10.1186/s12936-015-0820-2 (2015).

Malaria Genomic Epidemiology, N., Band, G., Rockett, K. A., Spencer, C. C. & Kwiatkowski, D. P. A novel locus of resistance to severe malaria in a region of ancient balancing selection. *Nature* **526**, 253-257, doi:10.1038/nature15390 (2015).

Makanga, B. *et al.* Ape malaria transmission and potential for ape-to-human transfers in Africa. *Proceedings of the National Academy of Sciences of the United States of America*, 1603008113-, doi:10.1073/pnas.1603008113 (2016).

Wanaguru, M., Liu, W., Hahn, B. H., Rayner, J. C. & Wright, G. J. RH5–Basigin interaction plays a major role in the host tropism of Plasmodium falciparum. *Proceedings of the National Academy of Sciences* **110**, 20735-20740, doi:10.1073/pnas.1320771110 (2013).

Wright, K. E. *et al.* Structure of malaria invasion protein RH5 with erythrocyte basigin and blocking antibodies. *Nature* **515**, 427-+, doi:10.1038/nature13715 (2014).

Triglia, T., Thompson, J. K. & Cowman, A. F. An EBA175 homologue which is transcribed but not translated in erythrocytic stages of Plasmodium falciparum. *Mol Biochem Parasitol* **116**, 55-63, doi:10.1016/S0166-6851(01)00303-6 (2001).

Farrell, A. *et al.* A DOC2 protein identified by mutational profiling is essential for apicomplexan parasite exocytosis. *Science* **335**, 218-221, doi:10.1126/science.1210829 (2012).

Ramiro, R. S. *et al.* Hybridization and pre-zygotic reproductive barriers in Plasmodium. *Proceedings of the Royal Society B-Biological Sciences* **282**, doi:10.1098/rspb.2014.3027 (2015).

Eksi, S. *et al.* Malaria transmission-blocking antigen, Pfs230, mediates human red blood cell binding to exflagellating male parasites and oocyst production. *Molecular Microbiology* **61**, 991-998, doi:10.1111/j.1365-2958.2006.0528.x (2006).

Mundwiler-Pachlatko, E. & Beck, H. P. Maurer's clefts, the enigma of Plasmodium falciparum. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 19987-19994, doi:10.1073/pnas.1309247110 (2013).

Bethke, L. L. *et al.* Duplication, gene conversion, and genetic diversity in the species-specific acyl-CoA synthetase gene family of Plasmodium falciparum. *Mol Biochem Parasitol* **150**, 10-24, doi:10.1016/j.molbiopara.2006.06.004 (2006).

Cunningham, D., Lawton, J., Jarra, W., Preiser, P. & Langhorne, J. The pir multigene family of Plasmodium: antigenic variation and beyond. *Mol Biochem Parasitol* **170**, 65-73, doi:10.1016/j.molbiopara.2009.12.010 (2010).

Niang, M. *et al.* STEVOR is a Plasmodium falciparum erythrocyte binding protein that mediates merozoite invasion and rosetting. *Cell Host Microbe* **16**, 81-93, doi:10.1016/j.chom.2014.06.004 (2014).

Kraemer, S. M. & Smith, J. D. A family affair: var genes, PfEMP1 binding, and malaria disease. *Curr Opin Microbiol* **9**, 374-380, doi:10.1016/j.mib.2006.06.006 (2006).

Gardner, M. J. *et al.* Genome sequence of the human malaria parasite *Plasmodium falciparum. Nature* **419**, 498-511, doi:10.1038/nature01097 (2002).

Bordbar, B. *et al.* Genetic diversity of VAR2CSA ID1-DBL2Xb in worldwide Plasmodium falciparum populations: impact on vaccine design for placental malaria. *Infection, genetics and evolution : journal of molecular epidemiology and evolutionary genetics in infectious diseases* **25**, 81-92, doi:10.1016/j.meegid.2014.04.010 (2014).

Frank, M., Dzikowski, R., Amulic, B. & Deitsch, K. Variable switching rates of malaria virulence genes are associated with chromosomal position. *Mol Microbiol* **64**, 1486-1498, doi:10.1111/j.1365-2958.2007.05736.x (2007).

Scally, A. & Durbin, R. Revising the human mutation rate: implications for understanding human evolution. *Nature reviews. Genetics* **13**, 745-753, doi:10.1038/nrg3295 (2012).

Carter, R. & Mendis, K. N. Evolutionary and historical aspects of the burden of malaria. *Clinical microbiology reviews* **15**, 564-594, doi:10.1128/CMR.15.4.564-594.2002 (2002).

Auburn, S. *et al.* An effective method to purify plasmodium falciparum dna directly from clinical blood samples for whole genome high-throughput sequencing. *PLoS ONE* **6**, doi:10.1371/journal.pone.0022213 (2011).

Oyola, S. O. *et al.* Optimized whole-genome amplification strategy for extremely AT-biased template. *DNA research : an international journal for rapid publication of reports on genes and genomes* **21**, 661-671, doi:10.1093/dnares/dsu028 (2014).

Oyola, S. O. *et al.* Whole genome sequencing of Plasmodium falciparum from dried blood spots using selective whole genome amplification. *bioRxiv*, doi:10.1101/067546 (2016).

Boissiere, A. *et al.* Isolation of Plasmodium falciparum by flow-cytometry: implications for single-trophozoite genotyping and parasite DNA purification for whole-genome high-throughput sequencing of archival samples. *Malaria Journal* **11**, 163, doi:10.1186/1475-2875-11-163 (2012).

Quail, M. A. *et al.* Optimal enzymes for amplifying sequencing libraries. *Nat Methods* **9**, 10-11, doi:10.1038/nmeth.1814 (2012).

Manske, H. & Kwiatkowski, D. SNP-o-matic. *Bioinformatics* **25**, 2434-2435, doi:10.1093/bioinformatics/btp403 (2009).

Chin, C. S. *et al.* Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* **10**, 563-569, doi:10.1038/nmeth.2474 (2013).

Assefa, S., Keane, T. M., Otto, T. D., Newbold, C. & Berriman, M. ABACAS: algorithm-based automatic contiguation of assembled sequences. *Bioinformatics Vol. 25 No. 15*, 1968-1969 (2009).

Carver, T. *et al.* Artemis and ACT: viewing, annotation and comparing sequences stored in relational database. *Bioinformatics* **24**, 2672-2676, doi:10.1093/bioinformatics/btn529 (2008).

Otto, T. D., Sanders, M., Berriman, M. & Newbold, C. Iterative Correction of Reference Nucleotides (iCORN) using second generation sequencing technology. *Bioinformatics* **26**, 1704-1707, doi:10.1093/bioinformatics/btq269 (2010).

English, A. C. *et al.* Mind the gap: upgrading genomes with Pacific Biosciences RS long-read sequencing technology. *PLoS One* **7**, e47768, doi:10.1371/journal.pone.0047768 (2012).

Otto, T. D. From sequence mapping to genome assemblies. *Methods Mol Biol* **1201**, 19-50, doi:10.1007/978-1-4939-1438-8_2 (2015).

Otto, T. D., Dillon, G. P., Degrave, W. S. & Berriman, M. RATT: Rapid Annotation Transfer Tool. *Nucleic Acids Research* **39**, e57, doi:10.1093/nar/gkq1268 (2011).

Stanke, M. *et al.* AUGUSTUS: ab initio prediction of alternative transcripts. *Nucleic Acids Research* **34**, W435-439, doi:10.1093/nar/gkl200 (2006).

Carver, T. *et al.* BamView: visualizing and interpretation of next-generation sequencing read. *Briefings in bioinformatics*, doi:10.1093/bib/bbr073 (2013).

Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754-1760, doi:10.1093/bioinformatics/btp324 (2009).

McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res* **20**, 1297-1303, doi:10.1101/gr.107524.110 (2010).

Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156-2158, doi:10.1093/bioinformatics/btr330 (2011).

Li, L., Stoeckert, C. J., Jr. & Roos, D. S. OrthoMCL: identification of ortholog groups for eukaryotic genomes. *Genome Res* **13**, 2178-2189, doi:10.1101/gr.1224503 (2003).

Jordan, G. & Goldman, N. The Effects of Alignment Error and Alignment Filtering on the Sitewise Detection of Positive Selection. *Molecular biology and evolution* **29**, 1125-1139, doi:10.1093/molbev/msr272 (2012).

Loytynoja, A. & Goldman, N. An algorithm for progressive multiple alignment of sequences with insertions. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 10557-10562, doi:10.1073/pnas.0409137102 (2005).

Loytynoja, A. & Goldman, N. Phylogeny-aware gap placement prevents errors in sequence alignment and evolutionary analysis. *Science* **320**, 1632-1635, doi:10.1126/science.1158395 (2008).

Fletcher, W. & Yang, Z. The Effect of Insertions, Deletions, and Alignment Errors on the Branch-Site Test of Positive Selection. *Molecular biology and evolution* **27**, 2257-2267, doi:10.1093/molbev/msq115 (2010).

Markova-Raina, P. & Petrov, D. High sensitivity to aligner and high rate of false positives in the estimates of positive selection in the 12 Drosophila genomes. *Genome Research* **21**, 863-874, doi:10.1101/gr.115949.110 (2011).

Morgulis, A., Gertz, E. M., Schäffer, A. A. & Agarwala, R. A Fast and Symmetric DUST Implementation to Mask Low-Complexity DNA Sequences. *Journal of Computational Biology* **13**, 1028-1040, doi:10.1089/cmb.2006.13.1028 (2006).

Wootton, J. C. & Federhen, S. Statistics of local complexity in amino acid sequences and sequence databases. *Computers & Chemistry* **17**, 149-163, doi:10.1016/0097-8485(93)85006-X (1993).

Castresana, J. Selection of Conserved Blocks from Multiple Alignments for Their Use in Phylogenetic Analysis. *Molecular biology and evolution* **17**, 540-552, doi:10.1093/oxfordjournals.molbev.a026334 (2000).

Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312-1313, doi:10.1093/bioinformatics/btu033 (2014).

Shimodaira, H. & Hasegawa, M. Multiple Comparisons of Log-Likelihoods with Applications to Phylogenetic Inference. *Molecular biology and evolution* **16**, 1114, doi:10.1093/oxfordjournals.molbev.a026201 (1999).

Castoe, T. A. *et al.* Evidence for an ancient adaptive episode of convergent molecular evolution. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 8986-8991, doi:10.1073/pnas.0900233106 (2009).

Thomas, G. W. C. & Hahn, M. W. Determining the Null Model for Detecting Adaptive Convergence from Genomic Data: A Case Study using Echolocating Mammals. *Molecular biology and evolution* **32**, 1232-1236, doi:10.1093/molbev/msv013 (2015).

Yang, Z. PAML 4: Phylogenetic Analysis by Maximum Likelihood. *Molecular biology and evolution* **24**, 1586-1591, doi:10.1093/molbev/msm088 (2007).

Zhang, J., Nielsen, R. & Yang, Z. Evaluation of an Improved Branch-Site Likelihood Method for Detecting Positive Selection at the Molecular Level. *Molecular biology and evolution* **22**, 2472-2479, doi:10.1093/molbev/msi237 (2005).

McDonald, J. H. & Kreitman, M. Adaptive protein evolution at the *Adh* locus in *Drosophila*. *Nature* **351**, 652-654, doi:10.1038/351652a0 (1991).

Rahnenfuhrer, A. A. a. J. topGO: Enrichment analysis for Gene Ontology. *R package*, doi:10.18129/B9.bioc.topGO (2010).

Guindon, S. *et al.* New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol* **59**, 307-321, doi:10.1093/sysbio/syq010 (2010).

Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* **32**, 1792-1797, doi:10.1093/nar/gkh340 (2004).

Gouy, M., Guindon, S. & Gascuel, O. SeaView version 4: A multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Molecular biology and evolution* **27**, 221-224, doi:10.1093/molbev/msp259 (2010).

Rask, T. S., Hansen, D. A., Theander, T. G., Gorm Pedersen, A. & Lavstsen, T. Plasmodium falciparum erythrocyte membrane protein 1 diversity in seven genomes--divide and conquer. *PLoS Comput Biol* **6**, doi:10.1371/journal.pcbi.1000933 (2010).

UniProt, C. UniProt: a hub for protein information. *Nucleic Acids Res* **43**, D204-212, doi:10.1093/nar/gku989 (2015).

Waterhouse, A. M., Procter, J. B., Martin, D. M., Clamp, M. & Barton, G. J. Jalview Version 2--a multiple sequence alignment editor and analysis workbench. *Bioinformatics* **25**, 1189-1191, doi:10.1093/bioinformatics/btp033 (2009).

Claessens, A. *et al.* Generation of antigenic diversity in Plasmodium falciparum by structured rearrangement of Var genes during mitosis. *PLoS Genet* **10**, e1004812, doi:10.1371/journal.pgen.1004812 (2014).

Bopp, S. E. *et al.* Mitotic evolution of Plasmodium falciparum shows a stable core genome but recombination in antigen families. *PLoS Genet* **9**, e1003293, doi:10.1371/journal.pgen.1003293 (2013).

Gronau, I., Hubisz, M. J., Gulko, B., Danko, C. G. & Siepel, A. Bayesian inference of ancient human demography from individual genome sequences. *Nat Genet* **43**, 1031-1034, doi:10.1038/ng.937 (2011).

Schiffels, S. & Durbin, R. Inferring human population size and separation history from multiple genome sequences. *Nat Genet* **46**, 919-925, doi:10.1038/ng.3015 (2014).

a

Branch site test

d_N/d_S

Chr

Orthologs/ *var* genes

π

Fixed diff.

P. adleri
P. gaboni
P. blacklocki
P. billcollinsi
P. reichenowi
P. praefalciparum
P. falciparum

b

fixed in great ape parasites
fixed in gorilla parasites
fixed in chimpanzee parasites

fixed differences

PF3D7_0424000

EBA-165

RH5

RH4

CyRPA

PF3D7_0423900

position on chromosome 4

1,060,000   1,080,000

P. adleri   L F I P S Y N T Y D E N I T N D F
P. gaboni   L F M P S Y N S Y D E N I A N N F
P. blacklocki   L F I P S N N T Y D E N V S N D F
P. billcolllinsi L F M P S N N I Y D E N M A N D F
P. reichenowi   L F M P S N N I Y D E N V A N D F
P. praefalciparum L F I P S N K T Y D E N V D N D F
P. falciparum

	P. falciparum	*P. praefalciparum*	*P. reichenowi*	*P. billcollinsi*	*P. blacklocki*	*P. gaboni*	*P. adleri*
var ≥ 2.5kb	67	112	92	35	43	61	58
rif > 250aa	183	582	440	280	13	12	19
stevor > 250aa	41	78	54	45	21	1	13
hyp4	9	2	1	1	0	0	0
hyp5	9	2	2	1	5	3	0
Maurer	13	12	5	4	7	0	0
EPF1	8	6	4	3	5	1	1
RESA-like	6	5	2	2	1	3	3
CLAG	5	7	6	7	35	16	24
DBLmsp	1	1	1	1	1	4	7
glycophorin binding	3	4	5	6	1	3	5
MSP7-like	8	8	7	6	4	7	14
Acyl-CoA	13	17	17	11	18	16	26

P. adleri   *P. blacklocki*   *P. reichenowi*   *P. falciparum*
P. gaboni   *P. billcollinsi*   *P. praefalciparum*

	CIDRa	CIDRb	CIDRd	CIDRg	CIDRn	CIDRpam	DBLpam1	DBLpam2	DBLpam3	DBLa	DBLb	DBLd	DBLe	DBLg	DBLz	Duffy	ATS
P. adleri	1	1	0	0	14	4	1	15	9	2	63	0	106	67	20	77	39
P. gaboni	1	1	0	0	8	16	3	30	20	2	47	0	84	43	16	48	41
P. blacklocki	0	1	0	0	0	0	0	0	0	1	17	0	16	55	7	0	34
P. billcollinsi	31	28	0	5	0	0	1	0	0	30	9	34	4	2	1	0	28
P. reichenowi	86	61	1	27	0	1	2	1	1	90	50	85	18	43	8	0	85
P. praefalciparum	85	48	5	30	0	5	5	5	5	94	86	72	97	84	34	0	105
P. falciparum	56	37	2	17	0	1	3	1	1	59	18	53	15	16	7	0	65

* Pfvar2csa /
** Clade A

P. adleri   *P. gaboni*   *P. blacklocki*   *P. billcollinsi*   *P. reichenowi*   *P. praefalciparum*   *P. falciparum*