Control of Lipid Domain Organization by a Biomimetic Contractile Actomyosin Cortex

Sven K. Vogel[1†], Ferdinand Greiss[1,2,3†], Alena Khmelinskaia[1,3] and Petra Schwille[1]

[1]Max-Planck Institute of Biochemistry, Am Klopferspitz 18, D-82152 Martinsried, Germany
[2]Systems Biophysics, Physics Department, Ludwig-Maximilans-University, Amalienstr. 54, D-80799 Munich, Germany
[3]Graduate School of Quantitative Biosciences, Ludwig-Maximilans-University, Feodor-Lynen-Str. 25, D-81377 Munich, Germany

† Contributed equally

Abstract

The cell membrane is a heterogeneously organized composite with lipid-protein micro-domains. The contractile actin cortex may govern the lateral organization of these domains in the cell membrane, yet the underlying mechanisms are not known. We recently reconstituted minimal actin cortices (MACs) (Vogel et al, 2013b) and here advanced our assay to investigate effects of rearranging actin filaments on the lateral membrane organization by introducing various phase-separated lipid mono- and bilayers to the MACs. The addition of actin filaments reorganized membrane domains. We found that the process reached a steady state where line tension and lateral crowding balanced. Moreover, the phase boundary allowed myosin driven actin filament rearrangements to actively move individual lipid domains, often accompanied by their shape change, fusion or splitting. Our findings illustrate how actin cortex remodeling in cells may control dynamic rearrangements of lipids and other molecules inside domains without directly binding to actin filaments.

Introduction

The spatiotemporal organization of lipids, proteins and other molecules at and within the cell membrane is pivotal for many fundamental cellular processes, such as signal transduction from the extracellular to the intracellular space (Groves & Kuriyan, 2010). Recent findings suggest that the cell membrane should be considered as a heterogeneous lipid protein layer with coexisting small micro domains and clusters of lipids and proteins that are assumed to dynamically form and reorganize in response to external and internal cues (Engelman, 2005; Simons & Gerl, 2010). Whether and how their spatiotemporal organization is actively regulated and maintained by the cell remains to be revealed. *In vivo* and *in vitro* studies suggest an important role of the eukaryotic actin cytoskeleton that interacts with the cell membrane via membrane-associated proteins (Heinemann et al, 2013; Koster et al, 2016; Murase et al, 2004; Sheetz et al, 1980). Actin structures were found to mediate the lateral organization of membrane proteins (Gudheti et al, 2013) and to modulate their diffusive behavior (Heinemann et al, 2013; Honigmann et al, 2014; Murase et al, 2004). Theoretical considerations have proposed a key role of the actin motor myosin for organizing and forming distinct protein and lipid micro-domains in cell membranes (Gowrishankar et al, 2012). However, direct experimental evidence for a control of lipid micro-domains by actomyosin rearrangements is still lacking.

In eukaryotic cells, the actin cortex is constantly rearranged by the motor protein myosin II and dozens of actin binding proteins. Therefore, reducing the complexity of experimental conditions, e.g. reducing dimensionality or exploiting a minimal biomimetic system, and utilizing microscopic techniques with a high temporal resolution is beneficial for studying these processes. Phase-separated lipid bilayers with controlled lipid compositions are well-established test beds to mimic lipid micro-domains in cell membranes and biological processes, e.g. the lateral organization of proteins that are otherwise difficult to observe *in vivo*. Ternary mixtures of lipids below their characteristic melting temperature (Tm) can phase separate into liquid disordered (L_d) and liquid ordered (L_o) domains. Liu and Fletcher used phase-separated giant unilamellar vesicles (GUVs) as a model system to study the influence of branched actin networks on the miscibility transition temperature (Liu & Fletcher, 2006). They reported that the formation of localized actin networks on PIP_2-containing phase-separated GUVs could act as switch for the orientation of lipid

domain growth sites.

However, the effects of an actin meshwork on individual lipid domains are technically

difficult to study, due to the 3-dimensional architecture of the GUVs. We therefore

made use of a minimal biomimetic system of planar geometry that we recently

developed (Vogel et al, 2013a; Vogel et al, 2013b) and combined the minimal actin

cortex (MAC) with supported phase-separated membranes and free-standing lipid

monolayers. We then visualized and studied the effects of actin filament adhesion and

myosin driven rearrangements on the lateral organization of membrane domains with

total internal reflection microscopy (TIRFM) and confocal spinning disk microscopy.

Results

Actin filament partitioning on phase-separated membranes

The effects of adhesion and myosin induced contraction of an actin meshwork on the

lateral organization of lipid domains was studied by combining phase-separated lipid

membranes with our established assay featuring contractile MACs (Vogel, 2016;

Vogel et al, 2013a; Vogel et al, 2013b) (Fig. 1A). For the adhesion assay a ternary

lipid mixture with DOPC, DPPC and Cholesterol in a 1:2:1 molar ratio was prepared

(see Material and Methods). Similar lipid compositions were described to form L_o and

L_d phases in free-standing membranes up to 30 °C (Veatch & Keller, 2003; Veatch &

Keller, 2005). The low miscibility transition temperature of the mixture avoids

thermal degradation of proteins and allowed us to study the phase transition behavior

in the presence of actin filaments. The fluorescent probe DiD (0.03 mol%) was used

to label the L_d phases (Garcia-Saez et al, 2007). The density of biotinylated actin

filaments was controlled by the concentration of biotinylated lipid DSPE-PEG(2000)-

Biotin (see also (Vogel, 2016; Vogel et al, 2013b)). As observed with TIRF

microscopy, the membrane separated into micrometer-sized L_o and L_d domains (Fig.

1B-E) and homogenized at ~37 ºC. The observed shift of 7 °C in Tm compared with

studies from (Veatch & Keller, 2003) agreed with our expectations considering the

interaction of lipid molecules with the mica support (Garcia-Saez et al, 2007).

The adhesion of actin filaments via neutravidin to biotinylated DSPE provided a

stable link between the MAC and the lipid bilayer over a wide range of temperatures.

To validate the partitioning preference of all molecular components, the fluorescence

signals of labeled neutravidin and actin filaments were acquired after domain

formation when cooled below Tm (Fig. 1B-E). We found that both neutravidin and

actin filaments partitioned into the L_o phase (Fig. 1B and C). We therefore concluded

that the biotinylated lipid DSPE partitioned strongly to the L_o phase.

Actin filament crowding effects on supported phase-separated membranes

We prepared low (0.01 mol % DSPE-PEG(2000)-Biotin) and medium (0.1 mol %

DSPE-PEG(2000)-Biotin, Fig. 2B) dense MACs to investigate the effect of actin

filament density on phase-separated membranes. To this end, the ternary lipid mixture

was incubated with non-labeled neutravidin and placed on a temperature-controlled

microscope objective. A fluorescence image was acquired as a reference before actin

filaments were added (Fig. 2A and B (left column)). A homogeneous membrane was

produced by heating the microscope objective to 42 °C (above Tm). Actin filaments

were added to the membrane above Tm (Fig. 2A and B (middle column)) and the

sample was subsequently incubated for ~45 min at 42 °C (Fig. 2A). The membrane

was completely covered by actin filaments after approximately 30 minutes (Fig. 2A

and B (middle column)). As final step, the membrane was slowly cooled down to 30

°C without active cooling and the fluorescence signal was recorded after complete

phase separation (Fig. 2A and B (right column)). While the low density MAC did not

show any influence on membrane domain properties (data not shown), the medium

dense MAC caused the formation of smaller domains (Fig. 2B, right column). Both,

actin filaments and DSPE-PEG(2000)-Biotin partitioned into the L_o phase (Fig. 2B

and C), confirming our initial observations (Fig. 1B - E). We conclude that actin

filaments act as nucleation sites for domain formation and drive their lateral spatial

distribution reproducing the observations reported by Honigmann et al (Honigmann et

al, 2014).

To further investigate the relation between lateral membrane organization and the

binding of actin filaments, we exposed high density MACs containing a lipid mixture

with 1.0 mol% biotinylated DSPE to different temperatures below Tm (Fig. 2C - F).

Fluorescently labeled actin filaments were added to the sample chamber at specified

temperatures below Tm and imaged for ~30 min using TIRF microscopy (Fig. 2C).

Because of the strong binding affinity of neutravidin and biotin, we found that the

change in fluorescence intensity of labeled actin filaments (which we used as a

measure for actin filament binding) was independent of the applied temperature (Fig. 2F). Hence, the adhesion process of actin filaments to the membrane was constant over the employed temperature range. The contour length (L) (Fig. 2G) of membrane domains was extracted and tracked over time (Fig. 2D). After ~1 min of actin filament addition, first domain deformations could be observed and the contour length was found to grow until a steady state at ~20 min was reached (Fig. 2D). The final contour length increased with temperature (Fig. 2E, final change in contour length: L_{Final} = 188±4 µm and τ = 9.5±0.5 min at 24 °C, L_{Final} = 314.0±1.5 µm and τ = 5.6±0.1 min at 29 °C, L_{Final} = 706.3±2.5 µm and τ = 4.4±0.1 min at 32 °C). The line tension γ between L_o and L_d domains is known to be a linear function of temperature with $\gamma \approx \gamma_0 (T_C - T)/T_C$ where T_C is the critical temperature (Baumgart et al, 2003; Honerkamp-Smith et al, 2008; Veatch et al, 2008). With the temperature-independent actin filament binding and the boundary energy given by $E = \gamma L$ (Yang et al, 2016) where E is the boundary energy, the final contour length L should increase with

$$L = \frac{E}{\gamma_0} \frac{T_C}{T_C - T} \, . \qquad (1)$$

The model describes the experimental data well (Fig. 2E) and gives a length change (E/γ_0) of 86±9 µm at T = 0 °C with T_C = 37 °C.

Actomyosin contraction governs lateral membrane domain organization

Aster shaped actomyosin clusters form *in vivo* (Luo et al, 2013; Munro et al, 2004; Verkhovsky et al, 1997) and *in vitro* (Backouche et al, 2006; Koster et al, 2016; Murrell & Gardel, 2012; Soares e Silva et al, 2011; Vogel et al, 2013b) upon myosin's contractile activity. Synthetic myofilaments contract the MAC in the presence of ATP and, hence rearrange actin filaments to form stable actomyosin clusters (Fig. 3A, C and G; Video 1 and 5). For studying the effects of myofilament induced actin rearrangements we prepared two lipid mixtures for lipid mono- and bilayer experiments, namely DOPC, DPPC and Cholesterol in a 1:2:1 molar ratio and DOPC, PSM and Cholesterol in a 3:3:1 molar ratio containing 0.1 or 1 mol% DSPE-PEG(2000)-Biotin DSPE (see Material and Methods).

We found that upon contraction, membrane domains deformed at the phase boundaries within minutes and resulted in various splitting and fusion events (Fig. 3A,

D, F and G; Videos 2-4 and 6). Domain shape changes, such as inward ingression,

were observed and correlated with the movement of actomyosin clusters (Fig. 3D;

Videos 3 and 4). Actomyosin contraction against smaller L_d domains resulted in their

net movement and occasionally led to splitting events of the domain (Fig. 3D; Video

4) while the contraction for L_o domains led to deformation and fusion with

neighboring domains (Fig. 3F; Video 6). For both mixtures the area of the L_o phase

remained constant over the course of actomyosin contraction (Fig. 3B and E, left

panels). The number of L_d domains increased with time due to their splitting for the

DOPC, DPPC, Cholesterol mixture (Fig. 3B, right panel; Videos 2-4). The number of

L_o domains decreased in the DOPC, PSM, Cholesterol mixture due to their fusion and

occasional disappearance during actomyosin contraction (Fig. 3E (right panel) and G;

Videos 6 and 7). A similar behavior in L_o domain disappearance has been observed by

Koster et al (Koster et al, 2016). In contrast to their work we did not observe a

decrease of the total L_o phase domain area in neither of the lipid mixtures (Fig. 3B

and E, left panels). In summary, we showed active lateral reorganization of lipid

domains for two different lipid mixtures, indicating the general validity of our

experimental observations.

In order to mimic free-standing membranes without support-induced friction as in the

SLBs but keeping the technical advantages of a planar geometry, we made use of a

recently developed lipid monolayer system (Chwastek & Schwille, 2013) with an air-

liquid interface (Fig. 4A - D). Actin filaments were coupled to ternary phase-

separated lipid monolayers similar to the situation in the supported lipid bilayer (SLB)

system via biotinylated lipids (1 mol%) and the use of neutravidin. Contrary to the

situation in the SLB system, actin filaments preferentially bind to the liquid extended

(disordered) (L_e) phase where also the neutravidin anchor protein mainly partitioned

to (Fig. 4A - D). We observed this behavior for both lipid mixtures that were also

used for the SLBs (see Material and Methods). In low and medium dense monolayer

MACs the liquid condensed (ordered) L_c domains acquire circular shapes and the

actin filaments close to the phase boundaries align to their circular shape (Fig. 4A). In

the case of low and medium densities we expect that the line tension energy

dominates over the actin filament wetting energy at the phase boundary and hence the

L_c can assume an unrestricted shape with aligned actin filaments. Similar effects have

been observed using bacterial cytoskeleton proteins (Arumugam et al, 2015). The

addition of myosin filaments in the presence of ATP led to the contraction of the actin

layer deforming and rearranging the L_c domains in both lipid mixtures similar to what

we observed in the SLB systems (Fig. 4B – D; Video 8). The shape changes included

fusion and stretching of the L_c domains (Fig. 4B – D; Video 8). Note that these

"active" deformation forces were only exerted in the presence of myosin filaments

and ATP. Interestingly, the obtained lipid domain shapes seemed to be stabilized after

the active contraction period, probably by the remaining actin filaments. Using the

monolayer system also tells us that the frictional force caused by a solid support in

SLBs does not play a significant role in the observed phenomena.

Discussion

As cells may need to quickly adapt the macro-and microscopic organization of lipid

and protein aggregates within the cell membrane due to external or internal cues, we

propose that actomyosin driven reorganization of actin filaments may aid to quickly

govern their lateral distribution. Recent evidence exists that the presence of an actin

meshwork influences the lateral diffusion behavior of lipids and proteins in

membranes *in vivo* (Murase et al, 2004) and *in vitro* (Heinemann et al, 2013) and that

it impacts the behavior of phase-separated membranes (Honigmann et al, 2014; Liu &

Fletcher, 2006). The important role of myosin is further supported by recent

theoretical and *in vitro* studies (Gowrishankar et al, 2012; Koster et al, 2016). It is

therefore tempting to speculate that cortical actomyosin contractility may be a generic

model for eukaryotic organisms not only to control their mechanical stability and

shape but also to quickly and actively control the lateral lipid and protein organization

at the cell membrane.

In our MACs that were combined with ternary lipid mixtures we found that binding of

actin filaments to a homogeneous bilayer at temperatures above Tm induced spatial

alignment of L_o domains to actin bound locations upon cooling below Tm (Fig. 2).

Here, actin filaments serve as nucleation sites for domain formation and drive their

lateral spatial distribution. Already phase separated, we could show that membrane

domains reorganized with the adhesion of actin filaments depending on line tension

and actin filament density. We further give direct evidence that the dynamic

reorganization of actin filaments by myosin motors actively changes the macroscopic

organization of membrane domains in our reconstituted phase-separated lipid bilayers

and monolayers. We propose that the transition energy between the L_d and L_o phase

(Baumgart et al, 2003; Garcia-Saez et al, 2007; Honerkamp-Smith et al, 2008)

enabled the lipid anchored actin filaments to exert lateral forces on the phase

boundaries leading to a macroscopic motion of lipid domains that eventually resulted

in splitting, fusion or deformation of the lipid domains and in their overall change in

number during and after the actomyosin contraction (Fig. 3 B and E, right panels;

Videos 2-4, 6 and 8). In a model we would first consider a biotinylated lipid that is

dragged by an actomyosin filament (Fig. 4E). The actomyosin filaments are unequally

associated with the lipid phases (L_o and L_d) with different viscosities. The drag is

counteracted by friction and, through the different viscosities, leads to domain

deformation and rearrangement. Here, the force propagation would be independent of

phase boundaries. As a second consideration, the biotinylated lipid would need to

overcome the transition barrier between phases while being dragged over a boundary

(Fig. 4E). Since our contraction experiments primarily showed that the boundaries

deformed locally at actin filament sites and that actomyosin foci formed at the vicinity

of phase boundaries, the high transition energy is favored as the dictating driving

force for phase deformation. Furthermore, domain deformation was commonly found

to happen when actin filaments were dragged over the L_d phase that impedes the free

contraction process through the strong partitioning preference of DSPE-PEG(2000)-

Biotin to the L_o phase.

With an estimated line tension of 1 pN for a 1:2:1 DOPC:DPPC:Cholesterol

membrane (Baumgart et al, 2003; Garcia-Saez et al, 2007; Kuzmin et al, 2005; Veatch

& Keller, 2005) and a reported value of ~1.5 $k_B T$ for the transfer of DPPC between

phases (Veatch et al, 2004), the energy for one DPPC molecule to transition is

roughly 6-fold higher than the energy that is needed to elongate the phase boundary

by 1 nm. Further experiments should elaborate on the physical model since

actomyosin contraction is clearly a non-equilibrium process and line tension itself was

reported to depend on the applied lateral tension (Akimov et al, 2007). Together with

our experimental findings, we can conclude that actively rearranging actin filaments

will lead to an extensive deformation of membrane domains through the physical link

with a small subset of membrane constituents. A plausible mechanism is therefore

apparent that explains active lateral rearrangements of membrane components by

actomyosin contractions without the need of binding directly to actin filaments.

Fig. 1. Minimal actin cortex on supported phase-separated lipid membranes. (A) Illustration depicting the preparation of the SLB and the coupling of actin filaments to the membrane (Adapted from (Vogel et al, 2013a)). (B) Since DiD is known to partition specifically to the L_d phase (1:2:1 DOPC:DPPC:Cholesterol), the fluorescence signal of Alexa-488-phalloidin labeled actin filaments showed that DSPE-PEG(2000)-Biotin partitioned into the L_o phase. (C) Likewise, Oregon Green labeled neutravidin colocalized with the L_o phase. (D and E) Line profiles of the normalized fluorescence signal of the actin cortex, DiD and neutravidin as was measured along the superimposed arrows in (B) and (C). Scale bar, 10 μm.

Figure 2

280
Fig. 2. Impact of actin filaments on the lateral organization of phase-separated membranes. (A) Scheme
of a phase separation experiment with and without MAC is shown with circles indicating time points
when fluorescent images were acquired. (B) Size distribution of lipid domains changed in the presence
of a MAC while undergoing phase separation. TIRF images of a medium dense MAC with DiD labeled
L_d phase (1:2:1 DOPC:DPPC:Cholesterol) are shown below Tm (30 °C) (left column), above Tm (42
°C) in the presence of bound Alexa-488-phalloidin labeled actin filaments (middle column) and below
Tm (30 °C) in the presence of bound actin (right column). The final size of L_d domains decreased
through the presence of actin filaments. (C) Change in contour length through the binding of actin

filaments to the already phase-separated membrane at various temperatures below Tm. A high density
MAC with a DiD labeled L_d phase before (0 min) and after the addition (32 min) of actin filaments at
24 °C is shown. (D) Contour length of L_d domains upon F-actin adhesion with time and mono-
exponential decay fits at 24 °C (blue dots), 29 °C (green dots) and 32 °C (red dots). (Note that the
observed local dip in contour length between 2 and 5 min at 24 °C is because of a focal misalignment
and blurring of the images after the addition of actin filaments). (E) Final change in contour length for
the different temperatures. With a constant lateral pressure, the linearly decreasing line tension with
temperature leads to an increase in final contour length according to equation (1). Dashed vertical line
indicates Tm of 37 °C. (F) The binding kinetics of actin filaments was tracked with the normalized
fluorescence signal over time at 24 °C (blue dots), 29 °C (green dots) and 32 °C (red dots). (G)
Illustrative sketch for the change in contour length before and after F-actin adhesion. Scale bars, 10
μm.

Figure 3

Fig. 3. Actomyosin contraction governs lateral membrane organization for various lipid compositions. (A) TIRFM time-lapse images of a contracting MAC with Alexa-488-phalloidin labeled actin filaments and DiD labeled L_d domains of a 1:2:1 DOPC:DPPC:Cholesterol mixture. Scale bar, 10 µm. (B) For the 1:2:1 DOPC:DPPC:Cholesterol mixture the area of the L_o phase was found to remain constant while the number of L_d domains increased with time through splitting. (C) Distance to neighboring actomyosin clusters for the lipid composition of 1:2:1 DOPC:DPPC:Cholesterol. (D) Representative example (1:2:1 DOPC:DPPC:Cholesterol mixture) showing movement (white asterisks) and deformation (inward ingression, yellow arrows) of a phase boundary upon actomyosin contraction and its splitting into two separated domains (yellow arrowheads). Scale bar, 10 µm. (E) For a lipid

composition of 3:3:1 DOPC:PSM:Cholesterol, the area of the L_o phase remained again constant and the
number of L_o domains decreased. (F) Pulling of a L_o domain (white asterisk) for a lipid composition of
3:3:1 DOPC:PSM:Cholesterol by actomyosin led to fusion (yellow arrowheads) and relaxation into a
larger neighboring domain. Scale bar, 2.5 μm. (G) Time-lapse montage of the actomyosin contraction
on a 3:3:1 DOPC:PSM:Cholesterol phase-separated membrane. Small L_o domains are moved and fused
to produce larger domains. Scale bar, 5 μm.

Figure 4

317
Fig. 4. Lipid monolayer domain shape changes and fusion events induced by actomyosin contractions.
(A) Confocal Spinning Disk microscope images of Alexa-488-phalloidin labeled actin filaments
coupled to a ATTO655-DOPE labeled lipid monolayer (3:3:1 DOPC:PSM:Cholesterol) in the absence

of myofilaments. Actin filaments close to the phase boundaries of lipid monolayer domains align to
their circular shape. (B-D) Spinning Disk microscope time-lapse images of Alexa-488-phalloidin
labeled actin filaments coupled to ATTO655-DOPE labeled lipid monolayers during myofilament
induced contractions of actin filaments. Actomyosin contractions lead to shape changes and fusion
events (yellow arrows and arrowheads) of the lipid monolayer L_c domains. (E) Scheme of a
microscopic model. Scale bars, 10 μm.

Video legends

Video 1: Myosin induced actin rearrangements in a minimal actin cortex (MAC)
combined with a supported phase-separated lipid bilayer

MAC with a supported phase-separated membrane (1:2:1 DOPC:DPPC:Cholesterol)

containing Alexa-488-phalloidin labeled actin filaments (green) exhibits dynamic

rearrangements of actin filaments after the addition of myofilaments in the presence

of ATP and eventually forms actomyosin clusters. The phase-separated membrane

containing DiD labeled L_d domains (red) is shown in the upper channel. The middle

channel shows Alexa-488-phalloidin labeled actin filaments (green) that bind to the

L_o domains. The lower channel shows the merge of both channels. TIRFM image

sequence was acquired at 5 sec. time intervals and contains 200 frames. The video is

displayed at 15 frames per second (fps). Total time: 16.6 min. Scale bar, 10 μm.

(Compressed JPG avi; 10.2 MB).

Video 2: Shape changes, rearrangements and fusion events of L_d domains during
actomyosin contraction

The phase-separated membrane (1:2:1 DOPC:DPPC:Cholesterol) containing DiD

labeled L_d domains (red) is shown in the left channel. The middle channel shows

Alexa-488-phalloidin labeled actin filaments that bind to the L_o domains. The right

channel shows the merge of both channels. TIRFM image sequence was acquired at 5

sec. time intervals and contains 124 frames. The video is displayed at 15 frames per

second (fps). Total time: 10.3 min. Scale bar, 10 μm. (Compressed JPG avi; 0.8 MB).

Video 3: Splitting, shape changes and deformations of L_d domains during
actomyosin contraction

The phase-separated membrane (1:2:1 DOPC:DPPC:Cholesterol) containing DiD

labeled L_d domains (red) is shown in the left channel. The middle channel shows

Alexa-488-phalloidin labeled actin filaments that bind to the L_o domains. The right

channel shows the merge of both channels. TIRFM image sequence was acquired at 5
sec. time intervals and contains 200 frames. The video is displayed at 15 frames per
second (fps). Total time: 16.6 min. Scale bar, 10 μm. (Compressed JPG avi; 0.8 MB).
359

Video 4: L_d domain movement, splitting and ingression during actomyosin
contraction
The phase-separated membrane (1:2:1 DOPC:DPPC:Cholesterol) containing DiD
labeled L_d domains (red) is shown in the left channel. The middle channel shows
Alexa-488-phalloidin labeled actin filaments that bind to the L_o domains. The right
channel shows the merge of both channels. TIRFM image sequence was acquired at 5
sec. time intervals and contains 200 frames. The video is displayed at 15 frames per
second (fps). Total time: 16.6 min. Scale bar, 5 μm. Corresponds to Figure 3D.
(Compressed JPG avi; 0.4 MB).
369

Video 5: Shape changes, rearrangements and fusion events of L_o domains during
actomyosin contraction
The phase-separated and DiD labeled (red) membrane (3:3:1
DOPC:PSM:Cholesterol) containing L_o domains (dark) is shown in the upper channel.
The middle channel shows Alexa-488-phalloidin labeled actin filaments (green) that
bind to the L_o domains. The lower channel shows the merge of both channels. TIRFM
image sequence was acquired at 5 sec. time intervals and contains 500 frames. The
video is displayed at 15 frames per second (fps). Total time: 41.6 min. Scale bar, 10
μm. (Compressed JPG avi; 28 MB).
379

Video 6: Fusion event of a L_o domain during actomyosin contraction
A small L_o domain is stretched by actin filaments thereby pulling the domain towards
a larger domain leading eventually to fusion of both L_o domains. The phase-separated
and DiD labeled (red) membrane (3:3:1 DOPC:PSM:Cholesterol) containing L_o
domains (dark) is shown in the left channel. The middle channel shows Alexa-488-
phalloidin labeled actin filaments (green) that bind to the L_o domains. The right
channel shows the merge of both channels. TIRFM image sequence was acquired at 5
sec. time intervals and contains 337 frames. The video is displayed at 120 frames per
second (fps). Total time: 28 min. Scale bar, 2.5 μm. Corresponds to Figure 3F.
(Compressed JPG avi; 0.4 MB).

Video 7: Disappearance of small L_o domains during actomyosin contraction

Small L_o domains often vanish during actomyosin contractions. The phase-separated

and DiD labeled (red) membrane (3:3:1 DOPC:PSM:Cholesterol) containing L_o

domains (dark) is shown in the left channel. The middle channel shows Alexa-488-

phalloidin labeled actin filaments (green) that bind to the L_o domains. The right

channel shows the merge of both channels. TIRFM image sequence was acquired at 5

sec. time intervals and contains 493 frames. The video is displayed at 120 frames per

second (fps). Total time: 41 min. Scale bar, 5 μm. Corresponds to Figure 3G.

(Compressed JPG avi; 4.1 MB).

Video 8: Shape changes and fusion events during actomyosin contraction of L_c

domains in a MAC combined with a phase-separated lipid monolayer

Myofilaments in the presence of ATP led to the contraction of the actin layers and to

shape changes and fusion events of the L_c domains. The phase-separated lipid

monolayer (3:3:1 DOPC:PSM:Cholesterol) containing the ATTO655-DOPE labeled

L_e phase (red) is shown in the left channel. The middle channel shows Alexa-488-

phalloidin labeled actin filaments that bind to the L_e phase. The right channel shows

the merge of both channels. Confocal Spinning Disk image sequence was acquired at

20 sec. time intervals and contains 64 frames. The video is displayed at 15 frames per

second (fps). Total time: 21 min. Scale bar, 10 μm. Corresponds to Figure 4C.

(Compressed JPG avi; 1.7 MB).

Material and Methods

Actin labeling and polymerization

F-actin preparation was performed as described in (Vogel et al, 2013b). Briefly, a

39.6 μM actin solution (Actin/Actin-Biotin ratio of 5:1) was prepared by mixing

rabbit skeletal actin monomers (32 μl, 2 mg/ml, Molecular Probes) with biotinylated

rabbit actin monomers (1.6 μl, 10 mg/ml, tebu-bio/Cytoskeleton Inc.). F-buffer

(1 mM DTT, 1 mM ATP, 10 mM Tris-HCl (pH 7.4), 2 mM MgCl$_2$ and 50 mM KCl)

was added to the mixture in order to start polymerization. Actin polymers were

labeled and stabilized with Alexa Fluor 488 Phalloidin according to the

manufacturer's protocol (Molecular Probes). Finally, the 2 μM Alexa-488-Phalloidin

labeled biotinylated actin filament solution was stored at 4 °C.

MAC (minimal actin cortex) preparation

1,2-dioleoyl-*sn*-glycero-3-phosphorcholine (DOPC), 1,2-dipalmitoyl-*sn*-glycero-3-

phosphocholine (DPPC), and cholesterol were added in a molar ratio of 1:2:1 or

DOPC, N-palmitoyl-D-sphingomyelin (PSM), and Cholesterol in a molar ratio of

3:3:1 to a final lipid concentration of 5 mg/ml (Avanti Polar Lipids, Inc). The lipid

bilayer was further supplemented with 0.03 mol% DiD (Molecular Probes, Eugene,

OR) and 0.01, 0.1 or 1 mol% DSPE-PEG(2000)-Biotin (Avanti Polar Lipids, Inc.).

The solution was dried under continuous nitrogen flux and then placed under vacuum

for 1 h to remove chloroform residuals. The pellet was rehydrated in SLB buffer (150

mM KCl, 25 mM Tris-HCl, pH 7.5) by vigorous vortexing and sonication.

20 μl of clear lipid suspension was then diluted in 130 μl A-buffer (50 mM KCl, 2

mM $MgCl_2$, 1 mM DTT and 10 mM Tris-HCl, pH 7.5) and heated to 55°C.

Meanwhile, freshly cleaved mica was fixated with immersion oil (Carl Zeiss, Jena,

Germany) on a cover slip (22x22 mm, #1.5, Menzel Gläser, Thermo Fisher) and

covered with the center part of a cut 1.5 mL Eppendorf tube. The Eppendorf tube was

glued with UV-sensitive glue (Norland Optical Adhesive 63, Cranbury, USA). The

chamber was filled with 75 μl of small unilamellar vesicles and incubated at 55 °C for

45 minutes with 1 mM $CaCl_2$. Non-fused vesicles were removed by washing the

suspension with 2 ml warmed A-buffer and gentle pipetting. The chamber's

temperature was slowly cooled. Next, 2 μl of unlabeled or Oregon-Green labeled

neutravidin (1mg/ml, Molecular Probes) diluted in 200 μl A-buffer were added to the

sample and incubated for 10 min. Finally, unbound proteins were removed by gently

washing the solution with 2 ml A-buffer.

Phase separation with MAC

The phase-separated membrane was heated with an objective heater (Carl Zeiss, Jena,

Germany) to the setpoint of 42 °C and cooled down to 30 °C, both with and without

Alexa-488-Phalloidin labeled biotinylated actin filaments. The area distribution of

domains at equilibrium (45 min after 30 °C was reached) was then compared between

the different concentrations of DSPE-PEG(2000)-Biotin (0.1, 0.01 mol%).

Crowding-Effect

The sample was placed on the TIRF microscope objective with attached objective heater and slowly warmed to the setpoint of 42 °C. The lipid bilayer with 1.0 % DSPE-PEG(2000)-Biotin was equilibrated at different temperatures below the melting point (~37 °C for 1:2:1 DOPC:DPPC:Cholesterol). 20 µl of Alexa-488-Phalloidin labeled biotinylated actin filaments were then carefully added to the chamber. Binding of actin filaments to the membrane was recorded by acquiring images in interleaved mode (488 nm and 640 nm) every 2.5 seconds, generating a time-lapse movie with a 5 sec delay between subsequent images.

F-actin network contraction by myofilaments

After the addition of 20 µl Alexa-488-Phallodin labeled biotinylated actin filaments to the supported lipid bilayer at room temperature (~24 °C), the mixture was incubated for approximately 45 min in order to ensure full binding of actin filaments to the membrane. Subsequently, residual actin was removed by gently exchanging the solution with 2 ml A-buffer. Once the properly assembled MAC was verified with TIRF microscopy, a solution of 20 µl myofilaments and 1 µl ATP (0.1 M) were added to start the compaction of actin filaments.

Images were acquired every 2.5 sec in interleaved mode, which eliminated the cross talk between color channels of actin filaments and the phase-separated membrane.

TIRF microscopy

Fluorescent imaging of labeled proteins and membrane was carried out on a custom-built TIRF microscope. The setup was integrated into an Axiovert 200 microscope (Zeiss). The probe was illuminated and imaged through a Plan-Apochromat 100x/NA 1.46 oil immersion objective with a 488-nm and 630-nm laser. Images were acquired with an Andor Solis EMCCD camera (electron gain = 300, exposure time = 50 ms, frame interval = 2.5 or 5 sec) in interleaved mode.

Data analysis

Image processing, analysis and data visualization was performed with Fiji and the scientific packages for Python. Multichannel beads were used to align double-color image stacks with the Fiji plugin Descriptor-based series registration. The contour

length between the lipid domains was extracted by detecting edges (Canny edge

detection; (van der Walt et al, 2014)) in single fluorescent images acquired from

phase-separated membranes. The contour length over time was fitted to $L =$

$L_{Final} (1 - \exp(-1/\tau (t - t_0)))$ with $t_0 = 1$ min. The actomyosin clusters were

detected using the Laplacian of Gaussian method (scikit-image (van der Walt et al,

2014)) and assigned to neighbors using the Delaunay triangulation and its method

vertex_neighbor_vertices.

To detect the lipid domains, the fluorescence images were firstly corrected for an

uneven background signal. Then, the DiD signal was classified with a local adaptive

thresholding algorithm and the corresponding phase boundaries with the Canny edge

detection algorithm. The binary image stacks were combined with the XOR operator

and further refined using the morphological closing operation. Small holes were

removed from the binary images. Small objects were detected with the Laplacian of

Gaussian method and as a final step, combined with the binary images. The script was

implemented in Python with the image processing package scikit-image.

MAC assembly on lipid monolayers

For the lipid monolayer assays the same lipid mixtures and molar ratios were used as

for the supported bilayers with 1 mol% of DSPE-PEG(2000)-Biotin. The lipids were

mixed, dried under nitrogen flux for 15 min, subsequently put into vacuum for 30 min

and dissolved in chloroform (total lipid concentration of 1 mg/ml). The mixtures were

further diluted to a final lipid concentration of 0.1 mg/ml and labeled by addition of

0.1 mol% of ATTO655-DOPE (ATTO-TEC GmbH, Siegen, Germany). The total

lipid concentration was confirmed by gravimetry.

To form a chamber (see Scheme 1), chamber spacers were cut from a 5mm thick sheet

of PTFE by a laser cutter. The spacers were sonicated step by step in acetone,

chloroform, isopropanol and ethanol (15 min each). Glass cover slips of 15 mm

(Gerhard Menzel GmbH, Braunschweig, Germany) were fixed to the spacer by

picodent twinsil® 22 two component glue (picodent®, Wipperfuerth, Germany). The

chambers were washed alternately with ethanol and water, air dried and air plasma-

cleaned for 10 min in order to make the glass hydrophilic. The surface was then

passivated by covering the glass surface with PLL-PEG(2000) (SuSos AG,

Dübendorf, Switzerland) 0.5 mg/mL solution in PBS buffer and incubated for

minimum half an hour. After throughout wash with water (5 times 200 μl) and

reaction buffer (3 times 200 μl), the chambers were ready to use.

Lipid monolayers were formed by drop-wise deposition of the lipid mixture on the

buffer-air interface (for further details, see also (Chwastek & Schwille, 2013)). A lipid

mixture volume corresponding to a lipid surface density of 70 $Å^2$ / molecule was

deposited drop-wise on the surface of the buffer solution.

The samples were imaged using a Yokogawa scan head CSU10-X1 spinning disk

system connected to a Nikon Eclipse Ti inverted microscope (Nikon, Japan) with an

Andor Ixon Ultra 512x512 EMCCD camera and a 3i solid state diode laser stack with

488 nm, 561 nm and 640 nm laser lines (3il33, Denver, Colorado USA). For

simultaneous Alexa-488-phalloidin and ATTO655-DOPE excitation, the 488 nm and

the 640 nm laser lines and an UPLanSApo 60x/1.20 Water UIS2 objective (Olympus,

Japan) were used. The time interval between the recorded images was 20 s.

After confirming the formation of a phase-separated lipid monolayer by imaging,

100 μl of neutravidin solution (0.01 μg/μl) was added to the sample twice and was

incubated for 5 min. Note that all protein solutions or other solutions are applied

directly to the liquid subphase by dipping the pipette tip through the monolayer. Next,

the subphase was washed 5 times with buffer (100 μl steps) to remove unbound

neutravidin. Subsequently, 20 μl of Alexa-488-phalloidin labeled actin filaments

(2 μM) was added to the subphase and incubated for at least 60 min, since binding of

actin filaments to the interface was assumed to occur relatively slowly. When the

binding of the actin filaments was confirmed by imaging, the monolayer was

thoroughly washed (7 to 10 steps of 100 μl) with reaction buffer containing 1 μM

ATP and 100 μl of myofilaments (0.3 μM) containing 1 μM ATP (enzymatically

regenerated see above) was added to the subphase twice. The sample was sealed by a

glass cover slide with grease to avoid subphase evaporation, allowing for long sample

observation. The lipid monolayer MAC system started to contract after a few minutes

of incubation, resulting in the formation of actomyosin clusters and deformation of

the lipid domains.

Scheme 1: Top (A) and lateral (B) schematic view of the PTFE chamber (adapted from (Chwastek & Schwille, 2013).

Acknowledgements

We are grateful for the financial support by the Daimler und Benz foundation (Project Grant PSBioc8216), the Gottfried Wilhelm Leibniz-Program of the DFG (SCHW716/8-1), the support of the Graduate School of Quantitative Biosciences Munich and the MaxSynBio consortium, which is jointly funded by the Federal Ministry of Education and Research of Germany and the Max Planck Society.

References

Akimov SA, Kuzmin PI, Zimmerberg J, Cohen FS (2007) Lateral tension increases the line tension between two domains in a lipid bilayer membrane. *Phys Rev E* **75**

Arumugam S, Petrov EP, Schwille P (2015) Cytoskeletal pinning controls phase separation in multicomponent lipid membranes. *Biophys J* **108:** 1104-1113

Backouche F, Haviv L, Groswasser D, Bernheim-Groswasser A (2006) Active gels: dynamics of patterning and self-organization. *Phys Biol* **3:** 264-273

Baumgart T, Hess ST, Webb WW (2003) Imaging coexisting fluid domains in biomembrane models coupling curvature and line tension. *Nature* **425:** 821-824

Chwastek G, Schwille P (2013) A monolayer assay tailored to investigate lipid-protein systems. *Chemphyschem* **14:** 1877-1881

583   Engelman DM (2005) Membranes are more mosaic than fluid. *Nature* **438:** 578-580
584

585   Garcia-Saez AJ, Chiantia S, Schwille P (2007) Effect of line tension on the lateral
586   organization of lipid membranes. *The Journal of biological chemistry* **282:** 33537-
587   33544
588

589   Gowrishankar K, Ghosh S, Saha S, C R, Mayor S, Rao M (2012) Active remodeling
590   of cortical actin regulates spatiotemporal organization of cell surface molecules. *Cell*
591   **149:** 1353-1367
592

593   Groves JT, Kuriyan J (2010) Molecular mechanisms in signal transduction at the
594   membrane. *Nat Struct Mol Biol* **17:** 659-665
595

596   Gudheti MV, Curthoys NM, Gould TJ, Kim D, Gunewardene MS, Gabor KA, Gosse
597   JA, Kim CH, Zimmerberg J, Hess ST (2013) Actin mediates the nanoscale membrane
598   organization of the clustered membrane protein influenza hemagglutinin. *Biophys J*
599   **104:** 2182-2192
600

601   Heinemann F, Vogel SK, Schwille P (2013) Lateral membrane diffusion modulated
602   by a minimal actin cortex. *Biophys J* **104:** 1465-1475
603

604   Honerkamp-Smith AR, Cicuta P, Collins MD, Veatch SL, den Nijs M, Schick M,
605   Keller SL (2008) Line tensions, correlation lengths, and critical exponents in lipid
606   membranes near critical points. *Biophys J* **95:** 236-246
607

608   Honigmann A, Sadeghi S, Keller J, Hell SW, Eggeling C, Vink R (2014) A lipid
609   bound actin meshwork organizes liquid phase separation in model membranes. *Elife*
610   **3:** e01671
611

612   Koster DV, Husain K, Iljazi E, Bhat A, Bieling P, Mullins RD, Rao M, Mayor S
613   (2016) Actomyosin dynamics drive local membrane component organization in an in
614   vitro active composite layer. *Proc Natl Acad Sci U S A*
615

616   Kuzmin PI, Akimov SA, Chizmadzhev YA, Zimmerberg J, Cohen FS (2005) Line
617   tension and interaction energies of membrane rafts calculated from lipid splay and tilt.
618   *Biophysical Journal* **88:** 1120-1133
619

620   Liu AP, Fletcher DA (2006) Actin polymerization serves as a membrane domain
621   switch in model lipid bilayers. *Biophys J* **91:** 4064-4070
622

623   Luo W, Yu CH, Lieu ZZ, Allard J, Mogilner A, Sheetz MP, Bershadsky AD (2013)
624   Analysis of the local organization and dynamics of cellular actin networks. *J Cell Biol*
625   **202:** 1057-1073
626

627   Munro E, Nance J, Priess JR (2004) Cortical flows powered by asymmetrical
628   contraction transport PAR proteins to establish and maintain anterior-posterior
629   polarity in the early C. elegans embryo. *Developmental cell* **7:** 413-424
630

631   Murase K, Fujiwara T, Umemura Y, Suzuki K, Iino R, Yamashita H, Saito M,
632   Murakoshi H, Ritchie K, Kusumi A (2004) Ultrafine membrane compartments for

molecular diffusion as revealed by single molecule techniques. *Biophysical journal* **86:** 4075-4093

Murrell MP, Gardel ML (2012) F-actin buckling coordinates contractility and severing in a biomimetic actomyosin cortex. *Proc Natl Acad Sci U S A* **109:** 20820-20825

Sheetz MP, Schindler M, Koppel DE (1980) Lateral mobility of integral membrane proteins is increased in spherocytic erythrocytes.

Simons K, Gerl MJ (2010) Revitalizing membrane rafts: new tools and insights. *Nature Reviews Molecular Cell Biology* **11:** 688-699

Soares e Silva M, Depken M, Stuhrmann B, Korsten M, MacKintosh FC, Koenderink GH (2011) Active multistage coarsening of actin networks driven by myosin motors. *Proc Natl Acad Sci U S A* **108:** 9408-9413

van der Walt S, Schonberger JL, Nunez-Iglesias J, Boulogne F, Warner JD, Yager N, Gouillart E, Yu T, scikit-image c (2014) scikit-image: image processing in Python. *PeerJ* **2:** e453

Veatch SL, Cicuta P, Sengupta P, Honerkamp-Smith A, Holowka D, Baird B (2008) Critical fluctuations in plasma membrane vesicles. *Acs Chem Biol* **3:** 287-293

Veatch SL, Keller SL (2003) Separation of liquid phases in giant vesicles of ternary mixtures of phospholipids and cholesterol. *Biophys J* **85:** 3074-3083

Veatch SL, Keller SL (2005) Miscibility phase diagrams of giant vesicles containing sphingomyelin. *Phys Rev Lett* **94:** 148101

Veatch SL, Polozov IV, Gawrisch K, Keller SL (2004) Liquid domains in vesicles investigated by NMR and fluorescence microscopy. *Biophys J* **86:** 2910-2922

Verkhovsky AB, Svitkina TM, Borisy GG (1997) Polarity sorting of actin filaments in cytochalasin-treated fibroblasts. *Journal of cell science* **110 (Pt 15):** 1693-1704

Vogel SK (2016) Reconstitution of a Minimal Actin Cortex by Coupling Actin Filaments to Reconstituted Membranes. *Methods Mol Biol* **1365:** 213-223

Vogel SK, Heinemann F, Chwastek G, Schwille P (2013a) The design of MACs (minimal actin cortices). *Cytoskeleton (Hoboken)* **70:** 706-717

Vogel SK, Petrasek Z, Heinemann F, Schwille P (2013b) Myosin Motors Fragment and Compact Membrane-Bound Actin Filaments. *eLife* **2013;2:e00116**

Yang ST, Kiessling V, Tamm LK (2016) Line tension at lipid phase boundaries as driving force for HIV fusion peptide-mediated fusion. *Nature communications* **7**