Creating a National Vector Surveillance System: Integrated mosquito trap data and digital epidemiology

Samuel S.C. Rund[1,2,3] (Samuel.Rund@ed.ac.uk) and
Micaela Elvira Martinez[1,2,4] (micaelam@princeton.edu)

[1]These authors contributed equally to this article.
[2]These authors were co–principal investigators.

[3]Centre for Immunology, Infection and Evolution, University of Edinburgh,
Edinburgh, Scotland, UK

[4]Princeton University, New Jersey, USA

Major Subject Areas

- Ecology
- Epidemiology and Global Health

Abstract

Fluctuating population dynamics and shifting ranges, behavior, and phenology make monitoring of mosquito populations essential in controlling emerging pathogens. Despite ~1000 US mosquito control agencies, there is no centralized collation of their data. We provide a roadmap for the creation of a National Vector Surveillance System, for mosquito control agencies to routinely report standardized data. We characterized the extent of current monitoring, and collated mosquito abundance between 2009-2016. Despite a minority of agencies publically reporting data, our data set consists of records on >12 million mosquitoes. We demonstrate the

utility of a National Vector Surveillance System by providing novel insight into

nationwide mosquito ecology and show that Digital Epidemiology could provide

indirect mosquito monitoring, with Google queries for "mosquito" covarying with

mosquito abundance. We demonstrate a National Vector Surveillance System would

provide a cost effective means to combat vector borne disease emergence and can be

integrated with wide-scale Digital Epidemiology.

Introduction

The emergence of arboviruses ("arthropod-borne viruses") —dengue virus

(DENV), West Nile Virus (WNV), chikungunya virus (CHIKV), and Zika virus

(ZIKV)—is a global problem[1]. There are multiple hypotheses regarding the epidemic

emergence of arboviruses. For example, increasing globalization and climate change

could be driving geographic expansion of vectors and increased transmission from

vectors to humans[2,3]. The occurrence of an arbovirus epidemic requires the presence

of vectors in sufficient abundance to sustain transmission. Mosquito monitoring has,

therefore, long been recognized as an important component of disease surveillance

and control efforts[4].

Large scale surveillance efforts have provided insight into the distribution of

disease vectors worldwide, including Aedes, Anopheles, and Culex mosquitoes which

are primary vectors for DENV, malaria, and WNV, respectively[5–7]. Maps of vector

geographic distribution (*i.e.*, presence/absence) are often used to assess disease

risk[2,5,8]. Most recently, vector mapping has been used in the US to infer risk of ZIKV

transmission[9]. It was predicted that the risk of ZIKV transmission in the US is

relatively localized to southern states. As predicted[9,10], ZIKV transmission established

in Florida, specifically, Miami-Dade County. As of October 12, 2016 there have been

128 reports of locally-acquired ZIKV cases[11]. Given the public health threat of

emerging (*e.g.*, ZIKV, CHIKV) and endemic (*e.g.*, WNV) arboviruses in the US,

there is great incentive to invest in studies of vector distribution and abundance.

In the US, mosquito control and surveillance efforts are ongoing at state- and

local-levels[12]. Public health and environmental agencies are tasked with (1)

systematic monitoring of vector abundance and mosquito infection status (*i.e.,* using

adult mosquito trapping and mosquito larval pool testing for viral presence), and (2)

reducing disease transmission risk by controlling vector population size using

larvicide, larval-site management, and/or insecticidal fogging[13]. Many agencies have

active surveillance programs that entail daily, weekly, and/or monthly trapping and

counting of mosquitoes, with taxonomic identification of the genus or species level in

many instances. Some vector surveillance programs were started in the 1940s at US

military installations to combat malaria transmission, and have since continued[4,14,15],

resulting in the generation of long-term ecological data. Time series data of vector

abundance that result from such long-term monitoring are invaluable for the study of

vector population dynamics and quantifying the risk vectors pose to human health.

There are ~1000 mosquito control agencies in the US, with a mosquito control

agency broadly defined as the local government authority responsible for the

surveillance and control of mosquitoes. Surprisingly, the exact number of agencies is

unknown, (*personal communication*, Joseph M Conlon, Technical Advisor, American

Mosquito Control Association). Mosquito control responsibilities may lie with the

local health department, or separate entities such as mosquito abatement districts. The

trapping and identification of mosquitoes requires considerable economic investment

in the form of personnel, equipment, and infrastructure. Despite these efforts, there is

no centralized reporting or repository in the US to facilitate the integration of such

publicly funded data. At present, in the US, the majority of mosquito surveillance data

remain isolated at local and/or state agencies and are inaccessible to the wider

mosquito surveillance and research communities[3,16].

In the US, mosquito-borne diseases are nationally notifiable with state-level

reporting to the CDC National Notifiable Disease Surveillance System[17]. We propose

a parallel system be created for mosquito population surveillance that is: standardized,

centralized, and regular. More specifically, data of a pre-specified type (standardized)

should be reported to a centralized system (*e.g.*, such as one hosted by the CDC,

PAHO, or WHO), and reporting should be done at regular intervals (weekly or

monthly). Having such a surveillance system in place would not only facilitate the

creation of Big Data on mosquito populations, but would also serve as a repository for

the wealth of surveillance data that exist to date. The creation and deployment of a

National Vector Surveillance System would be a relatively simple and low cost means

of establishing the risk of current and future arbovirus threats, and inform future

ecological and epidemiological studies. As proof of principle, here we summarize the

breath of mosquito surveillance efforts in the US, curate the limited data currently

accessible from mosquito control boards, and analyze these data to provide a glimpse

into vector population ecology, highlighting the potential power of a National Vector

Surveillance System data repository.

Results

Mosquito Abundance Data

We compiled a list and identified a web presence for 997 US mosquito control

agencies in 48 US states (Figure 1a, Supplemental File 6). We found 91 agencies

(9%) in 22 states provided historical data for at least one year in 2009-2015. Further,

54 agencies (5%) in 18 states provided live data (*i.e.*, 2016 data *reported* at least

monthly), which are indications of present and ongoing efforts to generate and make

abundance data publicly available online. 50 agencies (5%) provide both live and

historical data. While long, multi-year observations in the data were scarce, a total of

95 agencies provide live and/or historical data (Supplemental File 6).

We collated data from a subset of these agencies (76 of 91 agencies) where

data was presented in a mineable format such as online tables, graphs, and GIS maps

where population abundance could be followed over time. All data were limited to

what was publicly available online. The full data set can be found in Supplemental

File 1. Our final data set contains 144,234 records covering mosquito trapping data

from 2009-2015 (as well as a further 5235 records from 2016). We standardized dates

and taxonomic identifiers. The data covers 551 location identifiers, some trap level,

some a geographic feature (*e.g.,* a particular park name), and some a whole

jurisdiction or multi-agency region (*e.g.,* 'Northwestern Rural'). Our final data set

represented over 12 million individual mosquitoes counted (some data were provided

as averages, this is a minimum estimate of total mosquitoes trapped). From 2009-

2015, approximately 80% of the mosquitoes have genus level identification, and

~11% have species level identification.

Temporal resolution of the data was variable, with some data reported as

weekly, some monthly, and some a combination of the two. We consistently found a

lack of continuous sampling throughout the year, as surveillance efforts were targeted

to seasonal windows that varied between agencies. This frequently resulted in the

onset and/or offsets of the mosquito season being unobserved.

Taxonomic identification and reporting varied between agencies. While we

were able to standardize subtle differences in taxonomic reporting (*e.g., Aedes* vs.

Ochlerotatus or *Aedes* vs. *Ae.* there are some differences in reporting, however, that

could not be standardized. For example, in some districts, taxonomic descriptors such

as 'summer floodwater', 'other', 'non-culex', and 'nuisance species', were used to

identify at least some mosquitoes. However, we note that 56 agencies report species

level identification of at least some of the mosquitoes they collected. This reflects

local knowledge of the important vector and nuisance mosquito species in a given

control agency's jurisdiction.

Surveillance System

Reviewing the data across the many agencies provided insights into how a

nation-wide surveillance program could be designed. First, states could coordinate

with control agencies to place sentinel sites for trapping with particular emphasis on

ecological conditions in terms of urban/rural or climate zones within a state. New

Jersey, for example, divides their state into ten ecologically relevant zones[18]. The

traps in a given location should be of fixed type and attractant, and these traps should

ideally be checked at least weekly and reported weekly during the mosquito season.

We propose continuous trapping throughout the year be performed. This could be as

simple as adding monthly trapping in the winter to characterize the onset of mosquito

season more precisely. Given climate projections of earlier springs in the northern

hemisphere[2,3], there is a need to test for phenology shifts in disease vectors.

The data, units, traps, and metadata associated with the records varied widely

between agencies in our study. We do not suggest significant changes in operational

protocols, but it is important that agencies report full metadata, including trapping

strategy so their data can be interpreted. The most useable data will include trap

location name and/or GPS location, attractants used, the date the trap was counted,

and the duration of time the trap was set/reset. Finally, it is critically important that

the unit used for reporting mosquito count is standardized. Control boards should

submit trap-level data, and not aggregations over areas, and it must be made clear if

the data reported is male and female mosquitoes, or only female mosquitoes.

We found that some agencies attempted to identify each mosquito collected to

the species level. This effort is certain to yield valuable insights, but it is not feasible

for nation-wide standardized surveillance. Most districts, however, did identify

mosquitoes to the genus level. Due to their status as viral vectors of ZIKV, CHIKV,

and DENV, counts of *Ae. aegypti* and *Ae. albopictus* should always be reported.

Importantly, their absence should also be recorded. A similar system would be

desirable, but likely not feasible on a national scale, for members of the

morphologically similar invasive *Culex pipiens/quinquefasciatus* group which vector

West Nile and St. Louis Encephalitis viruses

Reporting of collected data would be submitted electronically, in a

standardized spreadsheet or online web form. We provide a mock-up reporting form

(Figure 1b) to illustrate the format of clean, unambiguous data that is ideal for

analyzing across agencies and states. Whenever possible and feasible, corresponding

historical data should also be represented in the database/national surveillance system.

Mosquito Seasonality

Due to the large amount of variation in the unstandardized data reported from

mosquito control agencies, we did not analyze the full data set. Instead, we searched

the compiled data for time series that would allow us to highlight the types of

analyses that could be done if there were a standardized national surveillance system.

Specifically, a national surveillance system would allow researchers to study (1)

geographic variation in mosquito populations, (2) interspecific/inter-genus variation

in phenology, and (3) interannual variation in mosquito abundance.

For any given taxonomic group, seasonal phenology may vary geographically

due to variation in environmental conditions. We searched the compiled data for a

taxonomic group for which we had 3+ years of data from multiple states. Data from

Culex pipiens in Minnesota, Iowa, and California fit these criteria and were of

particular interest because *C. pipiens* is the vector of West Nile Virus. We found that

the *C. pipiens* season was restricted to a narrow time frame (late summer) in

Minnesota (Figure 2a, d), the most northern of the three states. In Iowa and California,

the *C. pipiens* season extended later into the fall (Figure 2b-c, e-f), which might be

due to more mild autumnal weather in Iowa and California, relative to Minnesota.

Interestingly, in California the *C. pipiens* season began in the early spring. This could

be due to springtime environmental conditions being favorable throughout the state.

California, however, spans a large latitudinal range, and the springtime presence of *C.*

pipiens may be localized within the state, but by aggregating data from two districts in

California, we may have masked geographic variation within the state.

The benefit of genus or species level identification within a state is that it will

allow mosquito control agencies to identify differences in phenology among different

genus/species. The Culex genus was of interest because of the public health risk posed

by *C. pipiens*' transmission of WNV. We therefore searched the compiled data for a

state with species identification of Culex mosquitoes. Minnesota reported data for *C.*

pipiens, *C. restuans*, and *C. tarsalis*. We found that the seasonal timing of peak

abundance varied for these species, with *C. tarsalis* having the earliest peak, *C.*

restuans   intermediate,   and   *C.   pipiens*   the   latest   (

204
Figure 3). The seasonal staggering between taxonomic groups is important from a

public health perspective, because, if this pattern is consistent year-upon-year and

applies across states, then *C. tarselis* could be used as an early warning of the *C.*

pipiens season.

Continuous (throughout the year), long term times series of mosquito

abundance, such as would be generated by a National Vector Surveillance System,

will allow the research community to not only characterize mosquito seasonality, but

also study variation in mosquito abundance at larger temporal scales. California

provided continuous data for *C. pipiens* for 2011-2013. This time series, although it

only represents three years, revealed that *C. pipiens* abundance can vary greatly

between years. This demonstrates that mosquito abundance can be characterized on

two temporal scales (i) seasonal, characterized by the mosquito season vs. the off-

season, and (ii) interannual, characterized by high abundance years vs. low years

(Figure 4a). High abundance years might be tied to climate conditions or other

environmental variables. When continuous long-term time series are available, they

can be combined with data on environmental or other ecological variables, and

population dynamic models, to test for mechanistic drivers of variation in abundance.

A mechanistic understanding of vector population dynamics could allow for the

forecasting of high abundance years and allow for interventions to mitigate vector

borne diseases. Importantly, the data from California highlight that vector abundance

does not necessarily translate to higher disease risk. Although the abundance of

female *C. pipiens* was higher in 2011 than 2012 or 2013, there were fewer human

WNV cases in California in 2011 than in 2012 or 2013. There were 158, 479, 379

human cases in 2011, 2012, and 2013, respectively[19]. To study disease risk, mosquito

population dynamics would need to be factored into disease transmission models[20].

Google Queries as a Proxy for Mosquito Abundance

For the states for which we had both mosquito trap data and Google Trends

data during the same time period, we found that the abundance of Google queries

covaried with direct measures of mosquito abundance. Google queries for "mosquito"

in New York showed the same seasonal cycle (*i.e.*, high mosquito abundance in the

summer) as that observed in trap data (Figure 4b). Not only did the Google query data

capture seasonal variation in mosquito abundance, but in Illinois, a state with low

Culex abundance in 2011 and 2012, followed by high abundance in 2013, 2014, and

2015, Google queries captured both the seasonal variation in abundance and the

variation in abundance in high years vs. low years (Figure 4c). These data suggest

digital epidemiology can be used to supplement on-the-ground mosquito surveillance,

help inform early warning systems, and afford more surveillance coverage of the

population. We submit, however, that more mosquito trap data are needed in order to

perform a formal statistical evaluation of the digital epidemiology for mosquito

surveillance.

As the Google query data covaried with mosquito trap data in the US, we

therefore used Google query data of the language-specific search term "mosquito" as

an indirect measure of mosquito abundance in 17 countries, representing 39 locations

with sufficient data for testing seasonality (Figure **5**). There were multiple Google

trends time series for some countries with high search volume. Of the 39 locations,

wavelet analysis detected significant seasonality in 38 of them, with 2 countries

(Thailand and Taiwan) having a biannual signature. For those locations with

significant seasonality, GAMM model predictions demonstrated that the seasonal

variation in mosquito abundance was geographically structured.

Discussion

CDC guidelines call for standardized and repeated collection of mosquito

vectors[13] – a point recently reaffirmed by others[21,22]. We argue that nation-wide

aggregation and sharing of standardized data would allow the research community to

better understand the ecology of vectors, and this in turn would inform disease control

efforts. Clearly many states and agencies recognize the need to generate and

disseminate mosquito abundance data, as some states already operate

repositories[18,23,24]. We identified 49 agencies that provided abundance data on their

websites, independent of a state repository, indicating these agencies recognize the

importance of open data.

By collating publicly available data from 76 mosquito control agencies in the

US, we provide an extensive set of high temporal resolution mosquito abundance

data, comprised of >144,000 records. Importantly, these data represent only a small

fraction of the data that exist. The majority of mosquito abundance data are not

readily available to the scientific community. We demonstrate that mosquito

abundance data, collated from multiple data-silos, can give insight into the

spatiotemporal ecology of disease vectors. As we discovered, there is currently a great

deal of variation in how mosquito trap data are collected and reported. The lack of

data standardization presents hurdles to data integration. The mosquito control

community, however, could come together to decide upon the most efficient and

effective way to collect and report data in a standardized manner that would facilitate

real-time data release and data integration. Given the need to respond to emerging

mosquito-borne diseases in the Americas, the integration of mosquito abundance data

should be made a priority.

Enormous human effort and public expense has gone into the collection and

identification of the more than 12 million mosquitoes in our data set. Since the

generation of such data is publicly funded, future and past data should be made

publicly available in the spirit of The White House's Open Government Initiative[25,26].

A central repository is the most efficient way to promote data reuse and accessibility

of publicly funded data[16] across agencies/states - or even international and language

boundaries. Examples of similar interjurisdictional databases for the reporting and

dissemination of disease incidence include ArboNET[27] (US arbovirus incidence), the

CDC National Notifiable Disease Surveillance System[17], and the PAHO database[28]

(dengue surveillance in the Americas).

A significant hurdle to data integration is that the units of reporting varied

among agencies. As to be expected, agencies deploy a variety of traps, with a variety

of attractants. Comparing absolute mosquito trap numbers between localities is

unlikely to be useful to researchers or control professionals – it is the relative change

in abundance that is of interest, regardless of units.   However, successful

interpretation of species composition is dependent on knowing trapping methodology

– as traps have species bias[29].

A significant shortfall we identified is the lack of historical reporting of the

important invasive disease vectors, *Ae. aegypti* and *Ae. albopictus*. Only Iowa and

New Jersey report live *Ae. albopictus* abundance data*,* and no state reported live *Ae.*

aegypti abundance data. Further, it is often unclear if "0" or no data represents true

zeros (surveillance was performed, with zero members of the genus/species trapped)

versus a lack of specific surveillance for these critical species. Reliable

presence/absence data are necessary for distribution maps and modeling efforts[3,21,22].

We only used data generated by US mosquito control agencies. Substantial

data, primarily historical species surveillance from US military installations going

back to 1947, are held by VectorMap[15]. Further, NSF funded NEON projects sites are

coming online[30]. These twenty nationwide sites will record highly standardized, year-

round mosquito surveillance, identifications, and virus-status[30]. Our proposed

surveillance system, along with other research based data sets and international

governmental and non-governmental data sources are promising data sources for

integration.

Considering the caveats of our collated data, and the fact that it represents only

a small fraction of the mosquito abundance data that exists, we were nevertheless able

to uncover spatiotemporal patterns of mosquito populations that are relevant to public

health. This is enticing evidence that the data generated by centralized, regular,

standardized surveillance would be a powerful method for anticipating and

responding to arboviral threats. High temporal resolution time series data

(weekly/monthly) are powerful tools. As we have highlighted with our analyses, such

time series would allow for the characterization of spatial and temporal mosquito

population ecology. Characterizing mosquito population cycles and understanding the

mechanisms generating seasonal and interannual variation would enable us to:

(1) Infer when to anticipate epidemics/outbreaks;

(2) Quantify geographic variation in risk (how bad can epidemics become), especially

at geographic margins of vectors;

(3) Monitor virus persistence in source and sink populations and determine if vectors

overwinter in sufficient numbers to sustain transmission;

(4) Identify windows of opportunity for spraying, larval site management, ramping up

surveillance for pathogen (testing for virus), public outreach campaigns;

(5) Study vector-borne disease transmission dynamics; and

(6) Coordinate responses across states/counties, with the goal of controlling epidemics

and creating early warning systems.

Having rich data sets of mosquito abundance over time will allow for analyses

that have not yet been feasible, and the combination of data from multiple sources

would allow for analyses that can only be done in aggregate[31]. Modelers, for example,

require high quality data for model training and validation[3,22]. Data for aggregation,

need not only be mosquito abundance data, but truly new and powerful analyses

would be possible if abundance time series are combined with other sources of Big

Data. These include arbovirus disease incidence data, mosquito pool testing for virus

(ArboNET[27]); land use data (National Land Cover Database[32]); meteorological data

(Land Data Assimilation Systems, LDAS[33]), and climate change predictions (NCAR-

CCSM 4.0[34]). The data could be used to motivate action when neighboring

communities have vectors of concern[22], and tools could be developed to assist local

control agencies, such as the automated generation of up-to-date disease risk maps,

and other publicly disseminated reports. Finally, the data could be integrated into

biodiversity databases such as the Global Biodiversity Information Facility

(GBIF)[31,35].

The problems and opportunities we addressed here are not unique to the

United States. Emerging arboviruses are a global problem. Here we used the United

States as an operational example, but the data curation and application applies

worldwide. Furthermore, this is the appropriate time for the US and other countries to

initiate country-wide mosquito surveillance efforts due to the recent emergence of

ZIKV and CHIKV, and the continued threat that DENV poses in >100 countries

across six continents, causing 96 million cases per year[36]. The United States is

currently allocating resources for emerging vector-borne diseases and the Americas

are battling the ongoing Zika outbreak. We propose standardized reporting by existing

mosquito surveillance programs would be a low cost but powerful means of

empowering control efforts for vector borne diseases.

Methods

Mosquito Abundance Data

We systematically searched online to identify mosquito control agencies

within the US. Specifically, we used Google searches, the American Mosquito

Control Association website[37], state government websites (*e.g.*, Florida Department of

Agriculture & Consumer Services[38]), and state mosquito control association web

pages (*e.g.,* Texas, Michigan, and California[39–41]) to find the names/locations of

mosquito control agencies. We then supplemented our list of mosquito control

agencies by cross-referencing it with lists of mosquito control agencies generated by

other researchers. For each agency identified (Supplemental File 6), we searched for a

web presence, either an agency website or agency representation in a state-wide

mosquito abundance repository (*e.g.,* such as those that exist for Iowa, New Jersey,

and South Dakota[18,23,24]). The mosquito control agency websites/repositories were

searched for the presence of mosquito abundance surveillance data at a temporal

resolution of daily, weekly, and/or monthly from 2009 to mid-2016. We used data

from fixed traps. Data from temporary traps were excluded because they cannot be

adequately used to reconstruct time series of abundance. We collated data from 76 of

the 91 agencies presented in a mineable format such as online tables, graphs, and GIS

maps where population abundance could be followed over time. All data were limited

to what was publically available online.

Mosquito Abundance Data Analysis

In order to obtain state-level time series of mosquito abundance, the mosquito

trap data were aggregated within each state. We worked with states with data that

spanned multiple years: California, Colorado, Florida, Illinois, Iowa, Massachusetts,

Michigan, Minnesota, New Jersey, New York, and Ohio. The data were subset by

state, taxonomic group (*i.e.*, genus, with the exception of Culex which was analyzed

to the species-level), temporal resolution (*i.e.*, weekly or monthly data), and unit of

measure. The data were then aggregated within each subset by either summing or

averaging—depending on which was appropriate for each data type—across traps

collected at the same time. Data aggregation resulted in time series of mosquito

abundance for each combination of state and taxonomic group represented in our data.

The taxonomic groups represented in the aggregated data included *Culex, Culiseta,*

Aedes, Anopheles, Mansonia, Psorophora, Coquillettidia, Uranotaenia, and

Orthopodomyia. All aggregated data can be found in the Supplemental File 2. We

analyzed the time series for seasonal fluctuations in mosquito abundance and

characterized differences in seasonal phenology across states and taxonomic groups.

The unit of measure for reporting mosquito abundance differed among

mosquito control boards. Some mosquito control boards reported total mosquitoes

trapped, while others reported females per trap night, or mean females per trap. When

comparing seasonality across time series, we standardized the data so each time series

took values from 0-1. For time series observation, $y_{i,j}$, observation i in year j, the

following was used to standardized the data $s_{i,j} = (y_{i,j} - \min(y_j))/(\max(y_j)-\min(y_j))$.

Digital Epidemiology

Google Trends[42] were downloaded for queries of "mosquito" in the US, as

well as language-specific queries of "mosquito" in other countries (Supplemental

Files 3-5 contain the data and search term used for each country). Google Trends data

were downloaded for countries in each hemisphere, for all years for which data were

available, typically 2004-2015/2016, depending on the date of download. Google

Trends data represent the relative abundance of Google queries reported as values

from 0-100. In order to test for seasonality in Google queries of mosquito, we ran a

wavelet analysis on each time series, using the R package biwavelet[43]. Google Trends

time series with a significant 1-year period (significant when tested against time series

with the same lag-1 autocorrelation) were $\log_{10}$ transformed and detrended to make

them more sinusoidal and remove long term trends (which we assumed were due to

changes in internet query habits or Google algorithms, rather than changes in

mosquito ecology). Time series were standardized to values between 0-1 and a

GAMM model was fit to each standardized time series with the explanatory variable

being the time of the year. Predictions from the fitted GAMM models were used to

characterize mosquito seasonality on a global scale.

Acknowledgements

This work would not be possible without the efforts of the 95 local mosquito

control agencies highlighted in Supplemental File 6 who collected mosquito

population abundance data and made their data available online, as well as Iowa State

University Entomology Department, the Rutgers Center for Vector Biology, and

South Dakota State University for their state-level repositories[18,23,24].

We thank Frank Collins, Cynthia Lord, Paul Hickner, Lauren Cator, Luke

McNally, and Ryan Smith for their helpful discussions during the development of this

project.

We are incredibly grateful for Nathan Nixon, Ben Weise, Ben Chan, Jason

Cleland, Ollie Lee, and Emma Stotter for their work in locating and digitizing the

mosquito population abundance data.

Micaela Martinez was funded by a US National Science Foundation

Postdoctoral Fellowship in Biology Award Number 1523757, and is currently funded

by the NIH Director's Early Independence Award. Research reported in this

publication was supported by the Office of the Director, National Institutes of Health,

under Award Number DP5OD023 l00. The content is solely the responsibility of the

authors and does not necessarily represent the official views of the National Institutes

of Health.

Samuel S.C. Rund is funded by the Royal Society (NF140517) and a strategic

award from the Wellcome Trust (No. 095831).

Figures

Figure 2: *Culex pipiens* **seasonality. (a-c) Black points show the raw data from each trap within the state and correspond to the left y-axis. Blue time series are aggregated data across traps within the state, corresponding to the right y-axis. (d-f) Standardized data corresponding to the time series in a-c. The standardized time series were plotted individually for each year to show how mosquito abundance varied by time-of-year and among states.**

Figure 3: Variation in seasonal phenology among species. (a) *C. restuans* and *C. pipiens* abundance as reported in Minnesota. (b) *C. tarsalis* and *C. pipiens* abundance as reported in Minnesota. (c-d) Standardized data corresponding to the time series in a-b. The standardized time series are plotted individually for each year to show how mosquito abundance varied by month and among species.

Figure 4: (a) Time series of female *C. pipiens* abundance in California. Top line above the time series indicates the high abundance year vs. the low abundance years. Bottom line above the time series indicates the mosquito season vs. the off season. **(b-c)** Mosquitoes trapped in New York state and Illinois (black time series corresponding to the left y-axis), along with Google queries for the search term "mosquito" in each state during the same time period (yellow time series corresponding to the right y-axis). **(d)** Google queries for "mosquito" from 2010-2015 used for a wavelet analysis with power spectrum shown in **(e)** where the red area indicates significant 1-year periodicity of the time series. **(f)** Yellow points indicate the Google queries from New York, detrended and $\log_{10}$ transformed. Red curve indicates the predicted seasonal abundance as predicted by a GAMM model with time-of-year as the independent variable.

Figure 5: Mosquito seasonality around the world. The seasonal structure of mosquito abundance as measured indirectly via Google Trends using the language-specific search term "mosquito". Each row of the heatmap shows mosquito seasonality as predicted by the GAMM model for the location listed on the y-axis. The x-axis is the time-of-year. Dark blue represents the mosquito off-season and the bright green indicates the mosquito season.

Supplementary Material

Supplemental File 1: CSV file containing compiled mosquito population data from 2009-2016. See Mosquito Abundance Data in the Methods section for more details.

Supplemental File 2: CSV file containing time series of mosquito abundance created by aggregating data across traps within each state. The data are from California, Colorado, Florida, Illinois, Massachusetts, Minnesota, New Jersey, Iowa, Michigan, New York, and Ohio.

Supplemental File 3: CSV file containing monthly Google Trends data from multiple countries.

Supplemental File 4: CSV file containing weekly Google Trends data from multiple countries.

Supplemental File 5: CSV file containing Google Trends data from US with multiple years of mosquito trap data in Supplemental File 1.

Supplemental File 6: CSV file containing identified mosquito control agencies. The name and state of each mosquito control agency is provided. We provide a representative postcode and GPS coordinate in the geographic vicinity of each agency. Each agency is assigned a unique ID (AgencyID) that corresponds with population data recorded in Supplemental File 1.

References

1.   Fauci A, Morens D. Serving two masters. *N Engl J Med*. 2016;374(7):601-604. doi:10.1056/NEJMp1002530.
2.   Proestos Y, Christophides GK, Ergüler K, Tanarhte M, Waldock J, Lelieveld J. Present and future projections of habitat suitability of the Asian tiger mosquito, a vector of viral pathogens, from global climate simulation. *Philos Trans R Soc Lond B Biol Sci*. 2015;370(1665):20130554-. doi:10.1098/rstb.2013.0554.
3.   Campbell LP, Luther C, Moo-Llanes D, Ramsey JM, Danis-Lozano R, Peterson AT. Climate change influences on global distributions of dengue and chikungunya virus vectors. *Philos Trans R Soc B*. 2015;370:20140135-. doi:10.1098/rstb.2014.0135.
4.   Carpenter SJ. Insect control at army training camps. *J Econ Entomol*. 1944;37(4):470-474. http://jee.oxfordjournals.org/content/37/4/470. Accessed October 19, 2016.
5.   Hay SI, Sinka ME, Okara RM, et al. Developing global maps of the dominant Anopheles vectors of human malaria. *PLoS Med*. 2010;7(2):e1000209. doi:10.1371/journal.pmed.1000209.

6. Vazeille M, Mousson L, Rakatoarivony I, et al. Population genetic structure
and competence as a vector for dengue type 2 virus of Aedes aegypti and
Aedes albopictus from Madagascar. *Am J Trop Med Hyg*. 2001;65(5):491-
497. http://www.ncbi.nlm.nih.gov/pubmed/11716103. Accessed October
19, 2016.
7. Turell MJ, Dohm DJ, Sardelis MR, O'guinn ML, Andreadis TG, Blow JA. An
update on the potential of North American mosquitoes (Diptera: Culicidae)
to transmit west Nile virus. *J Med Entomol*. 2005;42(1):57-62.
8. Kraemer MUG, Sinka ME, Duda KA, et al. The global distribution of the
arbovirus vectors Aedes aegypti and Ae. Albopictus. *Elife*. 2015;4:e08347.
doi:10.7554/eLife.08347.
9. Bogoch II, Brady OJ, Kraemer MUG, et al. Anticipating the international
spread of Zika virus from Brazil. *Lancet*. 2016;387:335-336.
doi:10.1016/S0140-6736(16)00080-5.
10. Messina JP, Kraemer MU, Brady OJ, et al. Mapping global environmental
suitability for Zika virus. *Elife*. 2016;5:42-47. doi:10.7554/eLife.15272.
11. Centers for Disease Control and Prevention. Maps of Zika in the United
States | Zika virus. http://www.cdc.gov/zika/intheus/maps-zika-us.html.
Published 2016. Accessed October 19, 2016.
12. Council of state and territorial epidemiologists. *Assessment of Capacity in*
2012 for the Surveillance, Prevention and Control of West Nile Virus and
Other Mosquito-Borne Virus Infections in State and Large City/county
Health Departments and How It Compares to 2004.; 2014.
13. Centers for Disease Control and Prevention. *West Nile Virus in the United*
States: Guidelines for Surveillance, Prevention, and Control.; 2013.
14. Foley DH, Maloney FAJ, Harrison FJ, Wilkerson RC, Rueda LM. Online
spatial database of US Army Public Health Command Region-West 29
mosquito surveillance records: 1947-2009. *United States Army Med Dep J*.
2011;July-Sep:29-36. http://vectormap.si.edu/downloads/AMEDDJ.pdf.
Accessed October 19, 2016.
15. The Walter Reed Biosystematics Unit. VectorMap.
http://vectormap.si.edu/. Published 2016. Accessed October 20, 2016.
16. Sikes DS, Copas K, Hirsch T, Longino JT, Schigel D. On natural history
collections, digitized and not: A response to Ferro and Flick. *Zookeys*.
2016;618:145-158. doi:10.3897/zookeys.618.9986.
17. Centers for Disease Control and Prevention. National Notifiable Diseases
Surveillance System (NNDSS). https://wwwn.cdc.gov/nndss/. Published
2016. Accessed October 19, 2016.
18. Rutgers Center for Vector Biology. New Jersey Statewide Adult Mosquito
Surveillance Reports. http://vectorbio.rutgers.edu/reports/mosquito/.
Published 2016. Accessed October 20, 2016.
19. The California Department of Public Health. West Nile Virus Website.
http://westnile.ca.gov/. Published 2016. Accessed December 6, 2016.
20. Martinez ME. Preventing zika virus infection during pregnancy using a
seasonal window of opportunity for conception. Read AF, ed. *PLOS Biol*.
2016;14(7):e1002520. doi:10.1371/journal.pbio.1002520.
21. Hahn MB, Eisen RJ, Eisen L, et al. Reported Distribution of Aedes
(Stegomyia) aegypti and Aedes (Stegomyia) albopictus in the United
States, 1995-2016 (Diptera: Culicidae). *J Med Entomol*. 2016;ePub ahead:1-

7. doi:10.1093/jme/tjw072.

22.   Monaghan AJ, Morin CW, Steinhoff DF, et al. On the seasonal occurrence
and abundance of the Zika virus vector mosquito Aedes Aegypti in the
contiguous United States. *PLoS Curr Outbreaks*. 2016;1.
doi:10.1371/currents.outbreaks.50dfc7f46798675fc63e7d7da563da76.Au
thors.

23.   Iowa State University Entomology Department. Iowa Mosquito
Surveillance. http://mosquito.ent.iastate.edu/. Published 2016.

24.   South Dakota State University. South Dakota Mosquito Information
Systems. http://mosquito.sdstate.edu/. Accessed October 20, 2016.

25.   Costello MJ, Michener WK, Gahegan M, Zhang ZQ, Bourne PE. Biodiversity
data should be published, cited, and peer reviewed. *Trends Ecol Evol*.
2013;28(8):454-461. doi:10.1016/j.tree.2013.05.002.

26.   The White House. Open Government Initiative.
https://www.whitehouse.gov/open.

27.   CDC ArboNET. Diseasemaps. http://diseasemaps.usgs.gov/mapviewer/.
Published 2016. Accessed October 20, 2016.

28.   PAHO, WHO. Dengue | PAHO/WHO Data, Maps and Statistics.
http://www.paho.org/hq/index.php?option=com_topics&view=readall&ci
d=3273&Itemid=40734&lang=en. Published 2016. Accessed November 1,
2016.

29.   Acuff VR. Trap biases influencing mosquito collecting. *Mosq News*.
1976;36(2):173-176.

30.   National Science Foundation. NEON: National Ecological Observatory
Network. http://www.neonscience.org/.

31.   Costello MJ, Wieczorek J. Best practice for biodiversity data management
and publication. *Biol Conserv*. 2014;173:68-73.
doi:10.1016/j.biocon.2013.10.018.

32.   Homer C, Dewitz J, Yang L, et al. Completion of the 2011 National Land
Cover Database for the conterminous United States – Representing a
decade of land cover change information. *Photogramm Eng Remote Sens*.
2015:345-354.

33.   NASA Goddard. Land Data Assimilation Systems (LDAS).

34.   National Center for Atmospheric Research. Community Climate System
Model (CCSM). http://www.cesm.ucar.edu/models/ccsm4.0/. Accessed
October 20, 2016.

35.   Global Biodiversity Information Facility. GBIF. http://www.gbif.org/.
Accessed October 20, 2016.

36.   Guzman MG, Harris E. Dengue. *Lancet*. 2015;385:453-465.
doi:10.1016/S0140-6736(14)60572-9.

37.   American Mosquito Control Association. Links.
http://www.mosquito.org/links. Published 2016. Accessed October 20,
2016.

38.   Florida Department of Agriculture & Consumer Services. Mosquito
Directory. http://www.freshfromflorida.com/Divisions-
Offices/Agricultural-Environmental-Services/Consumer-
Resources/Mosquito-Control/Mosquito-Directory. Published 2016.
Accessed October 20, 2016.

39.   Texas Mosquito Control Association. Organized Mosquito Control Districts

in Texas. http://www.texasmosquito.org/Districts.html. Published 2016.
Accessed October 20, 2016.
40.   Mosquito and vector control association of California. Member Agencies -
MVCAC. http://www.mvcac.org/resources/member-agencies/. Published
2016. Accessed October 20, 2016.
41.   Michigan Mosquito Control Association. Information Links.
http://www.mimosq.org/links/links.htm. Accessed October 20, 2016.
42.   Google. Google Trends. https://www.google.com/trends/. Published 2016.
Accessed October 20, 2016.
43.   Gouhier TC, Grinsted A, Simko V. Package "biwavelet." 2016.
https://github.com/tgouhier/biwavelet. Accessed October 27, 2016.
617
618