TITLE PAGE

Title

Neural precursors of decisions that matter—an ERP study of deliberate and arbitrary choice

Authors

U. Maoz[1,2,3]*, G. Yaffe[4], C. Koch[5], L. Mudrik[6]

Affiliations

[1] Department of Psychology and Brain Institute, Chapman University, Orange, CA, USA.

[2] Department of Psychology, University of California, Los Angeles, CA, USA.

[3] Division of Humanities and Social Sciences, California Institute of Technology, Pasadena, CA, USA.

[4] Yale Law School, Yale University, New Haven, CT, USA.

[5] Allen Institute for Brain Science, Seattle, WA, USA.

[6] School of Psychological Science and Sagol School of Neuroscience, Tel Aviv University, Tel Aviv, Israel.

* urim@caltech.edu

Abstract

The onset of the readiness potential (RP)—a key neural correlate of upcoming action—was
repeatedly found to precede subjects' reports of having made an internal decision. This has
been taken by some as evidence against a causal role for consciousness in human decision-
making and thus as a denial of free-will. Yet those studies focused on purposeless, unreasoned,
arbitrary decisions, bereft of consequences. It remains unknown to what degree these specific
neural precursors of action generalize to deliberate decisions, which are more ecological and
relevant to real life. We therefore directly compared the neural correlates of deliberate and
arbitrary decision-making during a \$1000-donation task to non-profit organizations among
subjects prescreened for social involvement. While we found the expected RPs for arbitrary
decisions, they were strikingly absent for deliberate ones. Our results and a drift-diffusion
model we constructed are congruent with the RP representing the accumulation of noisy,
random fluctuations, which drive arbitrary—but not deliberate—decisions. The absence of RPs
in deliberate decisions further points to different neural mechanisms underlying deliberate and
arbitrary decisions and thus challenges the generalizability of studies that argue for no causal
role for consciousness in decision making from arbitrary to deliberate, real-life decisions.

Significance Statement

The extent of human free will has been debated for millennia. Previous studies demonstrated
that neural precursors of action—especially the readiness potential—precede subjects' reports
of deciding to move. Some viewed this as evidence against free-will. However, these
experiments focused on arbitrary decisions—e.g., randomly raising the left or right hand. We
directly compared deliberate (actual $1000 donations to NPOs) and arbitrary decisions, and
found readiness potentials before arbitrary decisions, but—critically—not before deliberate
decisions. This supports the interpretation of readiness potentials as byproducts of
accumulation of random fluctuations in arbitrary but not deliberate decisions and points to
different neural mechanisms underlying deliberate and arbitrary choice. Hence, it challenges
the generalizability of previous results to deliberate decisions.

MAIN TEXT

Introduction

Humans typically experience freely selecting between alternative courses of action, say, when
ordering a particular item off a restaurant menu. Yet a series of human studies using EEG
(Haggard & Eimer, 1999; Libet, Gleason, Wright, & Pearl, 1983; Salvaris & Haggard, 2014),
fMRI (S. Bode & Haynes, 2009; S Bode et al., 2011; Soon, Brass, Heinze, & Haynes, 2008;
Soon, He, Bode, & Haynes, 2013), intracranial (Perez et al., 2015), and single-cell recordings
(Fried, Mukamel, & Kreiman, 2011) challenged the validity of this common experience,
finding neural correlates of decision processes hundreds of milliseconds and even seconds
prior to the moment that subjects reported having consciously decided. Some have claimed,
following these and other findings, that the subjective human experience of freely deciding is
but an illusion, because human actions are unconsciously initiated before the conscious
decision to act (Harris, 2012; Libet et al., 1983; Wegner, 2002). This debate has been
captivating scholars from many disciplines in and outside of academia (Frith, Blakemore, &
Wolpert, 2000; Haggard, 2008; Jeannerod, 2006; Lau, Rogers, Haggard, & Passingham, 2004;
Mele, 2006; Wegner, 2002).

Critically, in the above studies, subjects were either told to arbitrarily move their right hand or
flex their right wrist; or they were instructed to arbitrarily move either the right or left hand
(Haggard, 2008; Hallett, 2016; A. Roskies, 2010). Thus, their decisions were always
unreasoned, purposeless, and bereft of any real consequence. This stands in sharp contrast to
real-life decisions that are reasoned, purposeful, and bear consequences (Ullmann-Margalit &
Morgenbesser, 1977)—which clothes to wear, what route to take to work, as well as more
formative decisions about life partners, career choices, and so on. Such deliberate decisions are
also at the center of the philosophical debate on free will (Breitmeyer, 1985; A. Roskies,
2010). They typically involve more conscious and lengthy deliberation and might thus be more
tightly bound to conscious processes.

Deliberate decisions have been widely studied in the field of neuroeconomics (Kable &
Glimcher, 2009; A. Sanfey, Loewenstein, McClure, & Cohen, 2006) and in perceptual tasks
(Gold & Shadlen, 2007). Yet, interestingly, little has been done in that field to assess the
relation between decision-related activity, subjects' conscious experience of deciding, and the
neural activity instantaneously contributing to this experience. Though some studies compared,
for example, internally driven and externally cued decisions (Thut et al., 2000; Wisniewski,
Goschke, & Haynes, 2016), or stimulus-based and intention-based actions (Waszak et al.,
2005), these were typically arbitrary decisions and actions that had no real implications.
Therefore, the results of these studies provide evidence for comparing arbitrary and deliberate
internal decisions.

Here, we compared neural precursors of deliberate and arbitrary decisions—and in particular
the readiness potentials (RP) on the same subjects, in an EEG experiment. We focused on the
RP because this component was the focus of so many previous studies of voluntary action. Our
experiment utilized a donation-preference paradigm, in which a pair of non-profit
organizations (NPOs) were presented in each trial. In deliberate-decision trials, subjects chose
to which NPO they would like to donate $1000. In arbitrary-decision trials, both NPOs
received an equal donation of $500, irrespective of subjects' key presses (Fig. 1). In both
conditions, subjects were instructed to report their decisions as soon as they made them, and
their hands were placed on the response keys, to make sure they could do so as quickly as

possible. Notably, while the visual inputs and motor outputs were identical between deliberate and arbitrary decisions, the decisions' meaning for the subjects was radically different: in deliberate blocks, the decisions were meaningful and consequential—reminiscent of important, real-life decisions—while in arbitrary blocks, the decisions were meaningless and bereft of consequences—mimicking previous studies of volition. Demonstrating differences in RP between arbitrary and deliberate decisions would first challenge the generalizability of the RP (from arbitrary to deliberate decisions) as an index for internal decision-making. Second, it would more generally suggest that different neural mechanisms might be at play between deliberate and arbitrary decisions. This, in turn, would question the generalizability of studies focused on arbitrary decisions to everyday, ecological, deliberate decisions—regardless of whether these studies relied on the RP or not.

Figure 1: Experimental paradigm. The experiment included deliberate (red, left panel) and arbitrary (blue, right panel) blocks, each containing nine trials. In each trial, two NPO names were presented, and subjects were asked to either choose to which NPO they would like to donate (deliberate), or to simply press either right or left, as both NPOs would receive an equal donation (arbitrary). They were specifically instructed to respond as soon as they reached a decision, in both conditions. Within each block, some of the trials were easy (lighter colors) decisions, where the subject's preferences for the two NPOs substantially differed (based on a previous rating session), and some were hard decisions (darker colors), where the preferences were more similar; easy and hard trials were intermixed within each block. To make sure subjects were paying attention to the NPO names, even in arbitrary trials, and to better equate the cognitive load between deliberate and arbitrary trials, memory tests (in light grey) were randomly introduced, where subjects were asked to determine which of four NPO names appeared in the immediately previous trial. For a full list of NPOs and causes see Supplementary Table 1.

Results

Behavioral Results

Subjects' reaction times (RTs) were analyzed using a 2-way ANOVA along decision type (arbitrary/deliberate) and difficulty (easy/hard). This was carried out on log-transformed data (raw RTs violated the normality assumption; W=0.94, p=0.001). As expected, subjects were substantially slower for deliberate than for arbitrary decisions (Fig. 2, left; $F_{(1,17)}=126.11$, p<0.0001 for the main effect of decision type). A main effect of decision difficulty was also found $F_{(1,17)}=18.76$, p=0.0004. Importantly, subjects were slower for hard vs. easy decisions in the deliberate case (hard vs. easy deliberate decisions: t(17)=4.78, p=0.0002), yet not for the arbitrary case (t(17)=1.01, p=0.33; $F_{(1,17)}=20.12$, p=0.0003 for the interaction between decision type and decision difficulty). This validates our experimental manipulation, and further demonstrates that, in deliberate decisions, subjects were making meaningful decisions, affected by the difference in the values of the two NPOs, while in arbitrary decisions they were not. What is more, the roughly equal RTs between easy and hard arbitrary decisions provide evidence against concerns that subjects were deliberating during arbitrary decisions.

Figure 2: Behavioral results. Response Times (RTs; left) and Consistency Grades (CG; right) in arbitrary (blue) and deliberate (red) decisions. Each dot represents the average RT/CG for easy and hard decisions for an individual subject (hard decisions: x-coordinate; easy decisions: y-coordinate). Group means and SEs are represented in dark red and dark blue crosses. The histograms at the bottom-left corner of each plot sum the number of dots with respect to the solid diagonal line. The dashed diagonal line represents equal RT/CG for easy and hard decisions; data points below that diagonal indicate longer RTs or higher CGs for hard decisions. In both measures, arbitrary decisions are more centered around the diagonal than deliberate decisions, showing no or substantially reduced differences between easy and hard decisions.

The consistency between subjects' choices throughout the main experiment and the NPO
ratings they gave prior to the main experimental session was also analyzed using a 2-way
ANOVA (see Methods). As expected, subjects were highly consistent with their own, previous
ratings when making deliberate decisions, but not when making arbitrary ones (Fig. 2, right;
$F(1,17)=946.55$, $p<0.0001$) for the main effect of decision type. A main effect of decision
difficulty was also found ($F(1,17)=57.39$, $p<0.0001$). Again, decision type and decision
difficulty interacted ($F(1,17)=25.96$, $p<0.0001$): subjects were much more consistent with their
choices in easy vs. hard deliberate decisions ($t(17)=11.15$, $p<0.0001$), than they were in easy
vs. hard arbitrary decisions ($t(17)=2.50$, $p=0.028$). Nevertheless, though subjects were around
chance (i.e., 0.5) in their consistency in arbitrary decisions (ranging between 0.39 and 0.64), it
seems that some subjects were slightly influenced by their preferences in easy-arbitrary
decisions trials, resulting in the significant difference between hard-arbitrary and easy-arbitrary
decisions above. Finally, no differences were found between subjects' tendency to press the
right vs. left key in the different conditions (both main effects and interaction: $F<1$).

EEG Results: Readiness Potential (RP)

The RP is generally held to index unconscious readiness for upcoming movement (Haggard,
2008; Kornhuber & Deecke, 1990; Libet et al., 1983; Shibasaki & Hallett, 2006); although
more recently alternative interpretations of the RP have been suggested (Miller, Shepherdson,
& Trevena, 2011; Schmidt, Jo, Wittmann, & Hinterberger, 2016; Schurger, Sitt, & Dehaene,
2012; Trevena & Miller, 2010; Verleger, Haake, Baur, & Śmigasiewicz, 2016). It has
nevertheless been the standard component studied in EEG versions of the Libet paradigm
(Haggard, 2008; Haggard & Eimer, 1999; Hallett, 2007; Libet, 1985; Libet et al., 1983; Libet,
Wright, & Gleason, 1982; Miller et al., 2011; Schurger et al., 2012; Shibasaki & Hallett, 2006;
Trevena & Miller, 2010). Here, the RP was measured over electrode Cz in the different
conditions by averaging the activity across trials in the 2 s prior to subjects' movement.
Focusing on the last 500 ms before movement onset for our statistical tests, we found a clear
RP in arbitrary decisions, yet RP amplitude was not significantly different from 0 in deliberate
decisions (Fig. 3A; $F(1,17)=11.86$, $p=0.003$ for the main effect of decision type; in t-tests
against 0, corrected for multiple comparisons, an effect was only found for arbitrary decisions
(hard: $t(17)=5.75$, $p<0.0001$; easy: $t(17)=5.09$, $p=0.0004$) and not for deliberate ones (hard:
$t(17)=1.24$, $p>0.5$; easy: $t(17)=1.84$, $p=0.34$). In a similar manner, regressing voltage against
time for the last 1000 ms before response onset, the downward trend was significant for
arbitrary decisions (Fig. 3B; $p<0.0001$, for both easy and hard conditions) but not for deliberate
decisions (hard: $p>0.5$, easy: $p=0.35$; all Bonferroni corrected for multiple comparisons).
Notably, this pattern of results was also manifested for single-subject analysis (Fig. 4; 14 of the
18 subjects had significant downward slopes for arbitrary decisions—i.e., $p<0.05$, Bonferroni
corrected for multiple comparisons—when regressing voltage against time for every trial over
the last 1000 ms before response onset; but only 5 of the 18 subjects had significant downward
slopes for the same regression analysis for deliberate decisions; see Methods. In addition, the
average slopes for deliberate and arbitrary decisions were -0.43±0.31 and -2.30±0.44
(mean±SE), respectively, a significant difference: $t(17)=3.51$, $p=0.001$).

To test whether the null result for RP amplitude in deliberate decisions stems from a genuine
absence of effect or from insufficient or underpowered data, we used Bayesian statistics.
Specifically, the Bayes factor allowed us to compare the probability of observing the data
given H_0 (i.e., no RP in deliberate decisions) against the probability of observing the data given
H_1 (i.e., RP exists in deliberate decisions). We followed the convention that a BF < 0.33
implies substantial evidence for lack of an effect (that is, the data is three times more likely to

be observed given H_0 than given H_1), $0.33 < BF < 3$ suggests insensitivity of the data, and BF
> 3 denotes substantial evidence for the presence of an effect (H_1) (Jeffreys, 1998). We found
strong evidence that arbitrary trials (pooled across difficulty for more statistical power) are
different from 0 (BF=$2.82 \cdot 10^6$), but inconclusive evidence for deliberate trials (BF=1.46, again
for pooled trials). Notably, however, this result might reflect the general, slow negative trend
of the deliberate trials data (Fig. 3A) rather than a typical RP, which is locked to movement-
onset and shows a sharp decline towards that. To remove the effect of this slow negative drift,
we used a baseline period of -1000 ms to -500 ms relative to *movement* onset (i.e., a baseline
that immediately preceded our time of interest window) instead of our usual baseline that
preceded stimulus onset (see Methods). Under this analysis, we found evidence that deliberate
decisions are not different from 0 (BF=0.332), supporting the claim that the RP during the last
500 ms before response onset was completely absent (BF for arbitrary decisions was $5.07 \cdot 10^4$).

Figure 3: The readiness potentials for deliberate and arbitrary decisions. (A) Mean and SE
of the Readiness Potential (RP) in deliberate (red shades) and arbitrary (blue shades) easy and
hard decisions in electrode Cz, as well as scalp distributions. Zero refers to time of right/left
movement, or response, made by the subject. Notably, the RP significantly differs from zero
and displays a typical scalp distribution for arbitrary decisions only. The scalp distribution was
calculated over the averaged activity during the last 500 ms before response, across subjects.
The inset shows the mean amplitude of the RP, with 95% confidence intervals, over the same
time window. Response-locked potentials with an expanded timecourse, and stimulus-locked
potentials are given in Fig. 6B and 6A, respectively. The same (response-locked) potentials as
here, but with a *movement-locked baseline* of -1 to -0.5 s (same as in our Bayesian analysis
below), are given in Fig. 6C. **(B)** Individual subjects' Cz activity in the four conditions (n=18).
The linear-regression line for voltage against time over the last 1000 ms before response onset
is designated by a dashed, black line. Note that the waveforms converge to an RP only in
arbitrary decisions.

We further tested whether differences in reaction time between the conditions, eye movements,
the hand with which the subjects executed the movement, and subjects' consistency scores
might explain our effect. We also tested whether the RPs might reflect some stimulus-locked
potentials or be due to baseline considerations.

Figure 4: Individual-subjects RPs. Six examples of for individual subjects' RPs for deliberate decisions (in red) and arbitrary ones (in blue).

Differences in reaction times (RT) between conditions, including stimulus-locked potentials and baselines, do not drive the effect

RTs in deliberate decisions were typically more than twice as long as RTs in arbitrary decisions. We therefore wanted to rule out the possibility that the absence of RP in deliberate decisions stemmed from the difference in RTs between the conditions. We carried out four analyses for this purpose. First, we ran a median split analysis—dividing the subjects into two groups based on their RTs: lower and higher than the median, for deliberate and arbitrary trials, respectively. We then ran the same analysis using only the faster subjects in the deliberate condition (M=1.91 s, SD=0.25) and the slower subjects in the arbitrary condition (M=1.25 s, SD=0.23). If RT length affects RP amplitudes, we would expect the RP amplitudes to be more similar between these two groups. However, though there were only half the data points, a similar pattern of results to those over the whole dataset was observed (Fig. 5A). Deliberate

and arbitrary decisions were still reliably different (F(1,17)=5.22, p=0.03), with significant RPs
found in arbitrary (easy: t(8)=4.57, p=0.0018; hard: t(8)=4.09, p=0.0035), but not deliberate
(easy: t(8)=1.92, p=0.09; hard: t(8)=0.63, p=0.54) decisions. In addition, the RPs for arbitrary
decisions were not significantly different between the subjects with above-median RTs and the
entire population for the easy or hard conditions (easy: t(25)=0.14, p>0.5; hard: t(25)=0.56,
p>0.5). Similarly, the RPs for deliberate decisions were not significantly different between the
subjects with below-median RTs and the entire population for the easy or hard conditions
(easy: t(25)=-0.34, p>0.5; hard: t(25)=0.17, p>0.5). This suggest that RTs do not reliably affect
Cz activation for deliberate or arbitrary decisions in our results.

Second, we regressed the difference between RPs in deliberate and arbitrary decisions
(averaged over the last 500 ms before response onset) against the difference between the RTs
in these two conditions for each subject (Fig. 5B). Again, if RT length affects RP amplitudes,
we would expect differences between RTs in deliberate and arbitrary conditions to correlate
with differences between RPs in the two conditions. But no correlation was found between the
two measures (r=0.22, t(17)=0.86, p=0.4). We further tried regressing the RP differences on
RT differences. The regression did not produce any reliable relation between RT and RP
differences (regression line: y = 0.54 [CI -0.8, 1.89] x - 0.95 [CI -2.75, 0.85]; the R^2 was very
low, at 0.05 (as expected from the r value above), and, as the confidence intervals suggest, the
slope was not significantly different from 0, F(1,17)=0.74, p=0.4).

A third concern that could relate to the RT differences among the conditions is that the RP in
arbitrary blocks might actually be some potential evoked by the stimuli (i.e., the presentations
of the two causes), specifically in arbitrary blocks, where the RTs are shorter (and thus stimuli-
evoked effects could still affect the decision). In particular, a stimulus-evoked potential might
just happen to bear some similarity to the RP when plotted locked to response onset. To test
this explanation, we plotted the potentials in all conditions, locked to the onset of the stimulus
(Fig. 6A). We also plotted the response-locked potentials across an expanded timecourse for
comparison (Fig. 6B). If the RP-like shape we see in Figs. 3A and 6B is due to a stimulus-
locked potential, we would expect to see the following before the 4 mean response onset times
(indicated by vertical lines at 0.98 and 1.00, 2.13, and 2.52 s for arbitrary easy, arbitrary hard,
deliberate easy, and deliberate hard, respectively) in the stimulus-locked plot (Fig. 6A):
Consistent potentials, which precede the mean response times, that would further be of a
similar shape and magnitude to the RPs found in the decision-locked analysis in the arbitrary
condition (though potentially more smeared for stimulus locking). We thus calculated a
stimulus-locked version of our ERPs, using the same baseline (Fig. 6A). As the comparison
between Fig. 6A and 6B clearly shows, no such consistent potentials were found before the 4
response times, nor were these potentials similar to the RP in either shape or magnitude (their
magnitudes are at the most around 1μV, while the RP magnitudes we found are around 2.5 μV;
Figs. 3A, 6B). This analysis thus suggests that it is unlikely that a stimulus-locked potential
drives the RP we found.

Figure 5: Relations between RTs and RPs. (A) The subjects with above-median RTs for arbitrary decisions (in blue) and below-median RTs for deliberate decisions (in red), show the same pattern that was found in the main analysis (compare Fig. 3A in main text). **(B)** A regression of the difference between the RPs versus the difference between the RTs for deliberate and arbitrary decisions for each subject. The equation of the regression line is y = 0.54 [CI -0.8, 1.89] x - 0.95 [CI -2.75, 0.85]. The R^2 is 0.05. One subject, #7, had an RT difference between deliberate and arbitrary decisions that was more than 6 interquartile ranges (IQRs) away from the median difference across all subjects. That same subject's RT difference was also more than 5 IQRs higher than the 75th percentile across all subjects. That subject was therefore designated an outlier and removed only from this regression analysis.

Figure 6: Stimulus- and response-locked Cz-electrode ERPs with different baselines and overlaid events. (A) Stimulus-locked waveforms including the trial onset range, baseline period, and mean reaction times for all four experimental conditions. **(B)** Response-locked waveforms with mean stimulus onsets for all four conditions as well as the offset of the highlighting of the selected cause and the start of the next trial. **(C)** Same potentials and timeline as Fig. 3A, but with a *response-locked* baseline of -1 to -0.5 s—the same baseline used for our Bayesian analysis.

A fourth concern is that the differences in RTs may affect the results in the following manner:
Because the main baseline period we used thus far was 1 to 0.5 s before stimulus onset, the
duration from the baseline to the decision also varied widely between the conditions. To make
sure this difference in temporal distance between the baseline period and the response to which
the ERPs were locked does not drive our results, we recalculated the potentials for all
conditions with a *response-locked* baseline of -1 to -0.5 s (Fig. 6C)—the same baseline we
used for the Bayesian analysis above. The rationale behind this choice of baseline was to have
the time that elapsed from baseline to response onset be the same across all conditions. As is
evident in Fig. 6C, the results for this new baseline were very similar to those for the stimulus-
locked baseline we used before. Focusing again on the -0.5 to 0 s range before response onset
for our statistical tests, we found a clear RP in arbitrary decisions, yet RP amplitude was not
significantly different from 0 in deliberate decisions (Fig. 6C; ANOVA $F_{(1,17)}=12.09$,
p=0.003 for the main effect of decision type; in t-tests against 0, corrected for multiple
comparisons, an effect was only found for arbitrary decisions (hard: $t_{(17)}=4.13$, p=0.0007;
easy: $t_{(17)}=4.72$, p=0.0002) and not for deliberate ones (hard: $t_{(17)}=0.38$, p>0.5; easy:
$t_{(17)}=1.13$, p=0.27). This supports the notion that the choice of baseline does not strongly
affect our main results. Taken together, all the results above provide strong evidence against
the claim that the differences in RPs stem from or are affected by the differences in RTs
between the conditions.

Eye movements do not affect the results

Though ICA was used to remove blink artifacts and saccades (see Methods), we wanted to
make sure our results do not stem from differential eye movement patterns between the
conditions. We therefore computed a saccade-count metric (SC; see Methods) for each trial for
all subjects. Focusing again on the last 500 ms before response onset, we computed mean (±
s.e.m.) SC values of 1.65±0.07 and 1.67±0.06 saccades for easy and hard deliberate decisions,
respectively, and 1.69±0.07 and 1.73±0.07 saccades for easy and hard arbitrary decisions,
respectively. We found no reliable differences between the number of saccades during
deliberate and arbitrary trials ($F_{(1,17)}=2.56$, p=0.13 for the main effect of decision type).

We further investigated potential effects of saccades by running a median-split analysis—
dividing the trials for each subject into two groups based on their SC score: lower and higher
than the median, for deliberate and arbitrary trials, respectively. We then ran the same analysis
using only the trials with more saccades in the deliberate condition (SC was 2.02±0.07 and
2.04±0.07 for easy and hard, respectively) and those with less saccades for the arbitrary
condition (SC was 1.33±0.07 and 1.31±0.08 for easy and hard, respectively). If the number of
saccades affects RP amplitudes, we would expect that the differences in RPs between arbitrary
and deliberate trials will diminish, or even reverse (as now we had more saccades in the
deliberate condition). However, though there were only half the data points for each subject in
each condition, a similar pattern of results to those over the whole dataset was observed:
Deliberate and arbitrary decisions were still reliably different within the median-split RPs
($F_{(1,17)}=16.70$, p<0.001), with significant RPs found in arbitrary (easy: $t_{(17)}=4.79$, p=0.002;
hard: $t_{(17)}=5.77$, p<0.001), but not deliberate (easy: $t_{(17)}=0.90$, p=0.38; hard: $t_{(17)}=0.30$,
p>0.5) decisions. In addition, we compared the RP data across all the trials with the median-
split RP data above. No significant differences were found for arbitrary decisions (easy:
$t_{(17)}=1.02$, p=0.32; hard: $t_{(17)}=0.75$, p=0.46) or for deliberate decisions (easy: $t_{(17)}=1.63$,
p=0.12; hard: $t_{(17)}=1.47$, p=0.16). Taken together, the analyses above provide strong evidence
against the involvement of eye movements in our results.

Testing alternative explanations

We took a closer look at subjects' behavior in the easy arbitrary condition, where some
subjects had a consistency score that was further above 0.5 (chance) than others. It seems like
those subjects had a greater difficulty ignoring their preferences, despite the instructions to do
so. We therefore wanted to test to what extent the RP of those subjects was similar to the RPs
of the other subjects. Focusing on the 8 subjects that had a consistency score above 0.55
(M=0.59, SD=0.03) and comparing their RPs to those of the 10 other subjects (consistency
M=0.50, SD=0.06) in easy arbitrary trials, we found no reliable differences (t(16)=0.94,
p=0.36). This is not surprising, as the mean consistency score of these subjects—though higher
than chance—was still far below their consistency score for easy deliberate decisions (M=0.99,
SD=0.02).

EEG Results: Lateralized Readiness Potential (LRP)

The LRP, which reflects activation processes within the motor cortex for action preparation
after action selection (Eimer, 1998; Masaki, Wild-wall, Sangals, & Sommer, 2004), was
measured by subtracting the difference potentials (C3-C4) in right-hand response trials from
this difference in left-hand responses trials and averaging the activity over the same time
window (Eimer, 1998; Haggard & Eimer, 1999). In this purely motor component, no
difference was found between the two decision types (Fig. 7; all Fs<1). It should be noted that
the LRP we calculated was relatively large, compared to Haggard and Eimer (1999),
potentially due to referencing differences (see Methods). Our analysis of EOG channels
suggests that some of that LRP might be driven by eye movements (we repeated the LRP
computation on the EOG channels instead of C3 and C4). However, the shape of the eye-
movement-induced LRP is very different from the LRP we calculated from C3 and C4. Also,
the differences that we found between conditions in the EOG LRP are not reflected in the
C3/C4 LRP. So, while our LRP might be boosted by eye movements, it is not strictly driven by
these eye movements.

371
372 **Figure 7: Lateralized readiness potential.** The lateralized readiness potential (LRP) for
373 deliberate and arbitrary, easy and hard decisions. No difference was found between the
374 conditions (ANOVA all Fs<1).

Drift Diffusion Model (DDM)

The main finding of this study—the absence of RP in deliberate decisions, suggesting different
neural underpinnings of arbitrary and deliberate decisions—is in line with recent work that
used a drift-diffusion model (DDM) to claim that the RP might arise from the nonlinearities
associated with threshold crossing together with time-locking neural activity to response onset,
rather than from specific preparation for or ballistic-like initiation of movement (Schurger et
al., 2012). DDMs of decision-making generally feature a process that rises toward a threshold.
The crossing of that threshold reflects the onset of the decision in the model, typically leading
to action. Schurger and colleagues modelled arbitrary decisions, and suggested that there the
nonlinear threshold crossing leading to response onset is largely determined by spontaneous
subthreshold fluctuations of the neural activity (Schurger et al., 2012). This challenged the
common view of the RP as a neural correlate of unconscious preparation for upcoming action
(Shibasaki & Hallett, 2006). Instead, Schurger and colleagues claimed, time-locking to
response onset ensures that these spontaneous fluctuations appear, when averaged over many
trials, as a gradual increase in neural activity.

To further assess this interpretation of the RP, we expanded the model developed by Schurger
et al. (2012) to a DDM that was composed of a *value-assessment* component and a *noise-
generation* component (see Methods). Under this assumption, Cz-electrode activity mainly
reflects the noise-generation component. (Note that we suggest that noise generation might be
a key function of the (pre-)SMA and other brain regions underneath the Cz electrode during
this specific task. When subjects make arbitrary decisions, these might be based on some
symmetry-breaking mechanism, which is driven by random fluctuations that are here simulated
as noise, following Schurger et al. Thus, we neither claim nor think that noise generation is the
main purpose or function of these brain regions in general.) In addition to the value-assessment
and noise-generation components, each trial was modeled as a race to threshold between the
NPO pair that served as the stimulus. One option was to select the NPO that was rated higher
than the other in the earlier rating session (the *congruent* option; see Methods); the other was to
select the lower-rated NPO (the *incongruent* option). Each decision was thus modeled as a race
between two leaky stochastic accumulators (each composed of a value-assessment and noise-
generation component; see Methods for more details and model parameters; Fig. 8D shows
some runs of the noise-generation component of the model). Note that the longer runs of the
model during deliberate decisions show a more pronounced tendency for what could be
perceived as decision reversals (or "changes of mind") than arbitrary decisions, as might be
expected a priory (Fig. 8D).

We fit our DDM to our average empirical reaction-times, which were 2.13, 2.52, 0.98 and 1.00
s for the different conditions (henceforth, magnitudes are given for deliberate easy, deliberate
hard, arbitrary easy, and arbitrary hard, respectively, in this order). The model's corresponding
mean RTs were 2.04, 2.46, 0.94, and 0.96 s for these conditions (Fig. 8A, B). The model was
further fit to the empirical consistency ratios (the proportions of congruent decisions), which
were 0.99, 0.83, 0.54 and 0.49. The model's corresponding consistency ratios were 1.00, 0.84,
0.53 and 0.53. The model then predicted the shape of the ERP in its noise component (assumed
to be reflected by Cz-electrode activity) for each decision type: a continuing, RP-like increase
in activity (with a negative sign) for arbitrary decisions, but only a very slight increase in
activity for deliberate decisions (Fig. 8C). This was in line with our empirical results (compare
Fig. 3A). Note that that the Schurger model aims to account for neural activity leading up to
the decision to move, but no further (Schurger et al., 2012). Similarly, we expect our DDM to

fit Cz neural data only up to around -0.1 s (100 ms before response onset). We also make no claims that ours is the only, or even optimal, model that explains our results. Rather, by extending the Schurger model, our goal was to show how that interpretation of the RP could also be applied to our more-complex paradigm. (We refer the reader to work by Schurger and colleagues (Schurger, 2018; Schurger et al., 2012) for further discussions about the model, its comparison to other models, and the relation to conscious-decision onset).

Figure 8: Empirical and model RTs and model prediction for Cz activity. (A) The empirical distributions of subjects' RTs across the four decision types. **(B)** The equivalent distributions of RTs for the model. **(C)** The model's prediction for the ERP activity in electrode Cz across all four decision types. **(D)** The first 20 model runs for the noise-generation component for all four decision conditions. The integration threshold, at 0.15, is designated by a dashed line in all decision conditions. Here t = 0 s designates the beginning of the model's run.

Discussion

Since the publication of Libet's seminal work claiming that neural precursors of action, in the form of the RP, precede subjects' reports of having consciously decided to act (Libet et al., 1983), a vigorous discussion has been raging among neuroscientists, philosophers, and other

scholars about the meaning of these findings for the debate on free will (recent collections
include (Mele, 2015; Pockett, Banks, & Gallagher, 2009; Sinnott-Armstrong & Nadel, 2011)).
Some claim that these results have removed conscious will from the causal chain leading to
action (Haggard, 2005, 2008; Wegner, 2002). Others are unconvinced that these results are
decisive for, or even applicable to, the free-will debate (Breitmeyer, 1985; Mele, 2009;
Nahmias, Shepard, & Reuter, 2014; A. Roskies, 2010). At the heart of much of this debate lies
the RP, thought to represent unconscious decision/planning mechanisms that initiate subjects'
decisions prior to their conscious experience of deciding (Kornhuber & Deecke, 1990; Libet et
al., 1983).

Notably, the RP and similar findings showing neural activations preceding the conscious
decision to act have typically been based on arbitrary decisions of different types (Haggard &
Eimer, 1999; Lau et al., 2004; Libet, 1985; Libet et al., 1983; Sirigu et al., 2004; Soon et al.,
2008; Soon et al., 2013). This, among other reasons, rested on the notion that for an action to
be completely free, it should not be determined in any way by external factors (Libet, 1985)—
which is the case for arbitrary, but not deliberate, decisions (where each decision alternative is
associated with a value, and the values of alternatives typically guide one's decision). But this
notion of freedom faces several obstacles. First, most discussions of free will focus on
deliberate decisions, asking when and whether these are free (Frankfurt, 1971; Hobbes, 1994;
Wolf, 1990). This might be because everyday decisions to which we associate freedom of
will—like choosing a more expensive but more environmentally friendly car, helping a friend
instead of studying more for a test, donating to charity, and so on—are generally deliberate, in
the sense of being reasoned, purposeful, and bearing consequences (although see
Deutschländer, Pauen, and Haynes (2017)). In particular, the free will debate is often
considered in the context of moral responsibility (e.g., was the decision to harm another person
free or not) (Fischer, 1999; Haggard, 2008; Maoz & Yaffe, 2016; A. L. Roskies, 2012; Sinnott-
Armstrong, 2014; Strawson, 1994), and free will is even sometimes defined as the capacity that
allows one to be morally responsible (Mele, 2006, 2009). In contrast, it seems meaningless to
assign blame or praise to arbitrary decisions. Thus, though the scientific operationalization of
free will has typically focused on arbitrary decisions, the common interpretations of these
studies—in neuroscience and across the free will debate—have often alluded to deliberate
ones. Here, we show that inference from arbitrary to deliberate decisions may not be justified,
as the neural precursors of arbitrary decisions, and in particular the RP, do not generalize to
meaningful ones (Breitmeyer, 1985; A. Roskies, 2010). These potentially different neural
mechanisms therefore challenge previous studies relying on arbitrary decisions, regardless of
whether they were based on the RP or not.

Interestingly, while the RP was present in arbitrary decisions but absent in deliberate ones, the
LRP—a long-standing, more-motor ERP component—was indistinguishable between the
different decision types. This provides evidence that, at the motor level, the neural
representation of the deliberate and arbitrary decisions that our subjects made may have been
indistinguishable, as was our intention when designing the task.

Our finding and the model thus suggest that two different mechanisms may be involved in
arbitrary and deliberate decisions. Earlier literature demonstrated that deliberate, reasoned
decision-making—which was mostly studied in the field of neuroeconomics (Kable &
Glimcher, 2009) or using perceptual decisions (Gold & Shadlen, 2007)—elicited activity in the
prefrontal cortex (PFC; mainly the dorsolateral (DLPFC) part (A. G. Sanfey, Rilling, Aronson,
Nystrom, & Cohen, 2003; J. D. Wallis & Miller, 2003) and ventromedial (VMPFC)
part/orbitofrontal cortex (OFC) (Ramnani & Owen, 2004; Jonathan D Wallis, 2007) and the

anterior cingulate cortex (ACC) (Bush, Luu, & Posner, 2000; Carter et al., 1998). Arbitrary, meaningless decisions, in contrast, were mainly probed using variants of the Libet paradigm, showing activations in the Supplementary Motor Area (SMA), alongside other frontal areas like the medial frontal cortex (Brass & Haggard, 2008; Krieghoff, Waszak, Prinz, & Brass, 2011) or the frontopolar cortex, as well as the posterior cingulate cortex (Fried et al., 2011; Soon et al., 2008) (though see Hughes, Schütz-Bosbach, and Waszak (2011), which suggests that a common mechanism may underlie both decision types). Possibly then, arbitrary and deliberate decisions may differ not only with respect to the RP, but be subserved by different underlying neural circuits, which makes generalization from one class of decisions to the other more difficult. Deliberate decisions are associated with more lateralized and central neural activity while arbitrary ones are associated with more medial and frontal ones. This appears to align with the different brain regions associated with the two decision types above, as also evidenced by the differences we found between the scalp distributions of arbitrary and deliberate decisions (Fig. 3A). Further studies are needed to explore this potential divergence in the neural regions between the two decision types.

To be clear, and following the above, we do not claim that the RP captures all unconscious processes that precede conscious awareness. However, some have suggested that the RP represents unconscious motor-preparatory activity before any kind of decision (e.g., Libet, 1985). But our results provide evidence against that claim, as we do not find an RP before deliberate decisions, which also entail motor preparation. What is more, in deliberate decisions in particular, it is likely that there are neural precursors of upcoming actions—possibly involving the above neural circuits as well as circuits that represents values—which are unrelated to the RP. Note also that we did not attempt to separately measure the timing of subjects' conscious decision to move. Rather, we instructed them to hold their hands above the relevant keyboard keys and press their selected key as soon as they made up their minds. This was both to keep the decisions in this task more ecological and because we think that the key method of measuring decision onset (using some type of clock to measure Libet's W-time) is highly problematic (see Methods). Some might also claim that unconscious decision-making could explain our results, suggesting that in arbitrary decisions subjects engage in unconscious deliberation or in actively inhibiting their urge to follow their preference as well as in free choice, while in deliberate decisions only deliberation is required. But this interpretation is unlikely because the longer RTs in deliberate decisions suggest, if anything, that more complex mental processes (conscious or unconscious) took place before deliberate and not arbitrary decisions. What is more, these interpretations should impede our chances of finding the RP in arbitrary trials (as the design diverges from the original Libet task), yet the RP was present, rendering them less plausible.

Aside from highlighting the neural differences between arbitrary and deliberate decisions, this study also challenges a common interpretation of the function of the RP. If the RP is not present before deliberate action, it does not seem to be a necessary link in the general causal chain leading to action. Schurger et al. (2012) suggested that the RP reflects the accumulation of stochastic fluctuations in neural activity that lead to action, following a threshold crossing, when humans arbitrarily decide to move. The shape of the RP therefore results from the manner in which it is computed: averaged over trials that are locked to response onset (that directly follows the threshold crossing). Our results and our model are in line with that interpretation and expand upon it, suggesting that the RP represents the accumulation of noisy, random fluctuations that drive arbitrary decisions, while deliberate decisions are mainly driven by the values associated with the decision alternatives (Maoz et al., 2013).

Our drift-diffusion model was based on the assumption that every decision is driven by a
component based on the values of the decision alternatives (the subject's support for the two
NPOs in our case) and by another component representing noise—random fluctuations in
neural activity. The value component plays little to no role in arbitrary decisions, so action
selection and timing depend on when the accumulation of noise crosses the decision threshold
for the congruent and incongruent decision alternatives. In deliberate decisions, in contrast, the
value component drives the decisions, while the noise has a smaller effect. Thus, in arbitrary
decisions, action onset closely tracks threshold crossings of the noise accumulation. But, in
deliberate decisions, the noise component is at more random levels at the moment of response
onset. Hence, locking the ERP to response onset and averaging over trials to obtain the RP,
leads to a relatively flat signal for deliberate decisions but to the expected RP shape in arbitrary
decisions. This provides strong evidence that the RP does not reflect subconscious movement
preparation. Rather, it is induced by threshold crossing of random fluctuations in arbitrary
decisions, which do not drive deliberate decisions; accordingly, the RP is not found there.
Further studies of the causal role of consciousness in deliberate versus arbitrary decisions are
required to test this claim.

Nevertheless, two possible, alternative explanations of our results can be raised. First, one
could claim that—in the deliberate condition only—the NPO names act as a cue, thereby
turning what we term internal, deliberate decisions into no more than simple responses to
external stimuli. Under this account, if the preferred NPO is on the right, it is immediately
interpreted as "Press right". It would therefore follow that subjects are actually not making
decisions in deliberate trials, which in turn is reflected by the absence of the RP in those trials.
However, the reaction time and consistency results that we obtained provide evidence against
this interpretation. We found longer reaction times for hard-deliberate decisions than for easy-
deliberate ones (2.52 versus 2.13 s, on average, respectively; Fig. 2 left) and higher
consistencies with the initial ratings for easy-deliberate decisions than for hard-deliberate
decisions (0.99 versus 0.83, on average, respectively; Fig. 2 right). If the NPO names acted as
mere cues, we would have expected no differences between reaction times or consistencies for
easy- and hard-deliberate decisions. In addition, there were 50 different causes in the first part
of the experiment. So, it is highly unlikely that subjects could memorize all 1225 pairwise
preferences among these causes and simply transform any decision between a pair of causes
into a stimulus instructing to press left or right.

Another alternative interpretation of our results is that subjects engage in (unconscious)
deliberation also during arbitrary decisions (Tusche, Bode, & Haynes, 2010), as they are trying
to find a way to break the symmetry between the two possible actions. If so, the RP in the
arbitrary decisions might actually reflect the extra effort in those types of decisions, which is
not found in deliberate decisions. However, this interpretation entails a longer reaction time for
arbitrary than for deliberate decisions, because of the heavier cognitive load, which is the
opposite of what we found (Fig. 2A). We would also expect the simpler deliberation in
arbitrary-easy trials to result in a shorter reaction-time than that of arbitrary-hard. But this is
not what we find (Fig. 2A).

In conclusion, our study suggests that RPs do not precede deliberate decisions. In addition, it
suggests that RPs represent an artificial accumulation of random fluctuations rather than
serving a genuine marker of an unconscious decision to initiate voluntary movement. This
further motivates future investigations into other precursors of action besides the RP using
EEG, fMRI, or other techniques. It would be of particular interest to find the neural activity

that precedes deliberate decisions. And it would also be of interest to find neural activity,
which is not motor activity, that is common to both deliberate and arbitrary decisions.

Materials and Methods

Subjects

Twenty healthy subjects participated in the study. They were California Institute of
Technology (Caltech) students as well as members of the Pasadena community. All subjects
had reported normal or corrected-to-normal sight and no psychiatric or neurological history.
They volunteered to participate in the study for payment ($20 per hour). Subjects were
prescreened to include only participants who were socially involved and active in the
community (based on the strength of their support of social causes, past volunteer work, past
donations to social causes, and tendency to vote). The data from 18 subjects was analyzed; two
subjects were excluded from our analysis (see *Sample size and exclusion criteria* below). The
experiment was approved by Caltech's Institutional Review Board, and informed consent was
obtained from all participants after the experimental procedures were explained to them.

Sample size and exclusion criteria

We ran a power analysis based on the findings of Haggard and Eimer (1999). Their RP in a
free left/right-choice task had a mean of 5.293 µV and standard deviation of 2.267 µV. Data
from a pilot study we ran before this experiment suggested that we might obtain smaller RP
values in our task (they referenced to the tip of the nose and we to the average of all channels,
which typically results in a smaller RP). Therefore, we conservatively estimated the magnitude
of our RP as half of that of Haggard & Eimer, 2.647 µV, while keeping the standard deviation
the same at 2.267 µV. Our power analysis therefore suggested that we would need at least 16
subjects to reliably find a difference between an RP and a null RP (0 µV) at a p-value of 0.05
and power of 0.99. This number agreed with our pilot study, where we found that a sample size
of at least 16 subjects resulted in a clear, averaged RP. Following the above reasoning, we
decided beforehand to collect 20 subjects for this study, taking into account that some could be
excluded as they would not meet the following predefined inclusion criteria: at least 30 trials
per experimental condition remaining after artifact rejection; and averaged RTs (across
conditions) that deviated by less than 3 standard deviations from the group mean.

Subjects were informed about the overall number of subjects that would participate in the
experiment when the NPO lottery was explained to them (see below). So, we had to finalize
the overall number of subjects who would participate in the study—but not necessarily the
overall number of subjects whose data would be part of the analysis—before the experiment
began. After completing data collection, we ran only the EEG preprocessing and behavioral-
data analysis to test each subject against the exclusion criteria. This was done before we looked
at the data with respect to our hypothesis or research question. Two subjects did not meet the
inclusion criteria: the data of one subject (#18) suffered from poor signal quality, resulting in
less than 30 trials remaining after artifact rejection; another subject (#12) had RTs longer than
3 standard deviations from the mean. All analyses were thus run on the 18 remaining subjects.

Stimuli and apparatus

620 Subjects sat in a dimly lit room. The stimuli were presented on a 21" Viewsonic G225f (20"
621 viewable) CRT monitor with a 60-Hz refresh rate and a 1024×768 resolution using

622 Psychtoolbox version 3 and Mathworks Matlab 2014b (Brainard, 1997; Pelli, 1997). They
623 appeared with a gray background (RGB values: [128, 128,128]). The screen was located 60 cm
624 away from subjects' eyes. Stimuli included names of 50 real, non-profit organizations (NPOs).
625 Twenty organizations were consensual (e.g., the Cancer Research Institute, or the Hunger
626 project), and thirty were more controversial: we chose 15 causes that were widely debated
627 (e.g., pro/anti guns, pro/anti abortions), and selected one NPO that supported each of the two
628 sides of the debate. This was done to achieve variability in subjects' willingness to donate to
629 the different NPOs. In the main part of the experiment, succinct descriptions of the causes
630 (e.g., pro-marijuana legalization, pro-child protection; for a full list of NPOs and causes see
631 Supplementary Table 1) were presented in black Comic Sans MS.

632 *Study Design*

633 The objective of this study was to compare ERPs elicited by arbitrary and deliberate decision-
634 making, and in particular the RP. We further manipulated decision difficulty to validate our
635 manipulation of decisions type: we introduced hard and easy decisions which corresponded to
636 small and large differences between subjects' preferences for the pairs of presented NPOs,
637 respectively. We reasoned that if the manipulation of decision type (arbitrary vs. deliberate)
638 was effective, there would be behavioral differences between easy and hard decisions for
639 deliberate choices but not for arbitrary choices (because differences in preferences should not
640 influence subjects' arbitrary decisions). Our 2 x 2 design was therefore decision type (arbitrary
641 vs. deliberate) by decision difficulty (easy vs. hard). Each condition included 90 trials,
642 separated into 10 blocks of 9 trials each, resulting in a total of 360 trials and 40 blocks. Blocks
643 of different decision types were randomly intermixed. Decision difficulty was randomly
644 counterbalanced across trials within each block.

645 *Experimental Procedure*

646 In the first part of the experiment, subjects were presented with each of the 50 NPOs and the
647 causes with which the NPOs were associated separately (see Supplementary Table 1). They
648 were instructed to rate how much they would like to support that NPO with a $1000 donation
649 on a scale of 1 ("I would not like to support this NPO at all) to 7 ("I would very much like to
650 support this NPO"). No time pressure was put on the subjects, and they were given access to
651 the website of each NPO to give them the opportunity to learn more about the NPO and the
652 cause it supports.

653 After the subjects finished rating all NPOs, the main experiment began. In each block of the
654 experiment, subjects made either deliberate or arbitrary decisions. Two succinct cause
655 descriptions, representing two actual NPOs, were presented in each trial (Fig. 1). In deliberate
656 blocks, subjects were instructed to choose the NPO to which they would like to donate $1000
657 by pressing the <Q> or <P> key on the keyboard, for the NPO on the left or right, respectively,
658 as soon as they decided. Subjects were informed that at the end of each block one of the NPOs
659 they chose would be randomly selected to advance to a lottery. Then, at the end of the
660 experiment, the lottery will take place and the winning NPO will receive a $20 donation. In
661 addition, that NPO will advance to the final, inter-subject lottery, where one subject's NPO
662 will be picked randomly for a $1000 donation. It was stressed that the donations were real and
663 that no deception was used in the experiment. To persuade the subjects that the donations were
664 real, we presented a signed commitment to donate the money, and promised to send them the
665 donation receipts after the experiment. Thus, subjects knew that in deliberate trials, every

choice they made was not hypothetical, and could potentially lead to an actual $1020 donation
to their chosen NPO.

Arbitrary trials were identical to deliberate trials except for the following crucial differences.
Subjects were told that, at the end of each block, the pair of NPOs in one randomly selected
trial would advance to the lottery together. And, if that pair wins the lottery, both NPOs would
receive $10 (each). Further, the NPO pair that would win the inter-subject lottery would
receive a $500 donation each. Hence it was stressed to the subjects that there was no reason for
them to prefer one NPO over the other in arbitrary blocks, as both NPOs would receive the
same donation regardless of their button press. Subjects were told to therefore simply press
either <Q> or <P> as soon as they decided to do so.

Thus, while subjects' decisions in the deliberate blocks were meaningful and consequential,
their decisions in the arbitrary blocks had no impact on the final donations that were made. In
these trials, subjects were further urged not to let their preferred NPO dictate their response.
Importantly, despite the difference in decision type between deliberate and arbitrary blocks, the
instructions for carrying out the decisions were identical: Subjects were instructed to report
their decisions as soon as they made them in both conditions. They were further asked to place
their fingers on the response keys, so they could respond as quickly as possible. Note that we
did not ask subjects to report their "W-time" (time of consciously reaching a decision), because
this measure was shown to rely on neural processes occurring after response onset (Lau,
Rogers, & Passingham, 2007) and to potentially be backward inferred from movement time
(Banks & Isham, 2009). Even more importantly, clock monitoring was demonstrated to have
an effect on RP size (Miller et al., 2011), so it could potentially confound our results (Maoz et
al., 2015).

Decision difficulty (Easy/Hard) was manipulated throughout the experiment, randomly
intermixed within each block. Decision difficulty was determined based on the rating
difference between the two presented NPOs. NPO pairs with 1 or at least 4 rating-point
difference were designated hard or easy, respectively. Based on each subject's ratings, we
created a list of NPO pairs, half of each were easy choices and the other half hard choices.

Each block started with an instruction written either in dark orange (Deliberate: "In this block
choose the cause to which you want to donate $1000") or in blue (Arbitrary: "In this block
both causes may each get a $500 donation regardless of the choice") on a gray background that
was used throughout the experiment. Short-hand instructions appeared at the top of the screen
throughout the block in the same colors as that block's initial instructions; Deliberate: "Choose
for $1000" or Arbitrary: "Press for $500 each" (Fig. 1).

Each trial started with the gray screen that was blank except for a centered, black fixation
cross. The fixation screen was on for a duration drawn from a uniform distribution between 1
and 1.5 s. Then, the two cause-descriptions appeared on the left and right side of the fixation
cross (left/right assignments were randomly counterbalanced) and remained on the screen until
the subjects reported their decisions with a key press—<Q> or <P> on the keyboard for the
cause on the left or right, respectively. The cause corresponding to the pressed button then
turned white for 1 s, and a new trial started immediately. If subjects did not respond within 20
s, they received an error message and were informed that, if this trial would be selected for the
lottery, no NPO would receive a donation. However, this did not happen for any subject on any
trial.

To assess the consistency of subjects' decisions during the main experiment with their ratings
in the first part of the experiment, subjects' choices were coded in the following way: each
binary choice in the main experiment was given a consistency grade of 1, if subjects chose the
NPO that was rated higher in the rating session, and 0 if not. Then an averaged consistency
grade for each subject was calculated as the mean consistency grade over all the choices. Thus,
a consistency grade of 1 indicates perfect consistency with one's ratings across all trials, 0 is
perfect inconsistency, and 0.5 is chance performance.

We wanted to make sure subjects were carefully reading and remembering the causes also
during the arbitrary trials to better equate memory load, attention, and other cognitive aspects
between deliberate and arbitrary decisions—except those aspects directly associated with the
decision type, which was the focus of our investigation. We therefore randomly interspersed 36
memory catch-trials throughout the experiment (thus more than one catch trial could occur per
block). On such trials, four succinct descriptions of causes were presented, and subjects had to
select the one that appeared in the previous trial. A correct or incorrect response added or
subtracted 50 cents from their total, respectively. (Subjects were informed that if they reached
a negative balance, no money will be deducted off their payment for participation in the
experiment.) Thus, subjects could earn $18 more for the experiment, if they answered all
memory test questions correctly. Subjects typically did well on these memory questions, on
average erring in 2.5 out of 36 memory catch trials (7% error) and gaining additional $16.75
(SD=3.19). Subjects' error rates in the memory task did not differ significantly between the
experimental conditions (2-way ANOVA; decision type: $F(1,17)=2.51$, $p=0.13$; decision
difficulty: $F(1,17)=2.62$, $p=0.12$; interaction: $F(1,17)=0.84$, $p=0.37$).

ERP recording methods

The EEG was recorded using an Active 2 system (BioSemi, the Netherlands) from 64
electrodes distributed based on the extended 10–20 system and connected to a cap, and seven
external electrodes. Four of the external electrodes recorded the EOG: two located at the outer
canthi of the right and left eyes and two above and below the center of the right eye. Two
external electrodes were located on the mastoids, and one electrode was placed on the tip of the
nose. All electrodes were referenced during recording to a common-mode signal (CMS)
electrode between POz and PO3. The EEG was continuously sampled at 512 Hz and stored for
offline analysis.

ERP analysis

ERP analysis was conducted using the "Brain Vision Analyzer" software (Brain Products,
Germany) and in-house Mathworks Matlab scripts. Data from all channels were referenced
offline to the average of all channels, which is known to result in a reduced-amplitude RP
(because the RP is such a spatially diffuse signal). The data were then digitally high-pass
filtered at 0.1 Hz using a Finite Impulse Response (FIR) filter to remove slow drifts. A notch
filter at 59-61 Hz was applied to the data to remove 60-Hz electrical noise. The signal was then
cleaned of blink and saccade artifacts using Independent Component Analysis (ICA)
(Junghofer, Elbert, Tucker, & Rockstroh, 2000). Signal artifacts were detected as amplitudes
exceeding ±100 µV, differences beyond 100 µV within a 200 ms interval, or activity below
0.5 µV for over 100 ms (the last condition was never found). Sections of EEG data that
included such artifacts in any channel were removed (150 ms before and after the artifact). We
further excluded single trials in which subjects pressed the wrong button as well as trials where
subjects' RTs were less than 200 ms, more than 10s, or more than 3 standard deviations away

from that subject's mean in that condition (mean number of excluded trials =7.17, SD=2.46,
which are 1.99% of the trials). Overall, the average number of included trials in each
experimental cell was 70.38 trials with a range of 36-86 out of 90 trials per condition. Channels
that consistently had artifacts were replaced using interpolation (4.2 channels per subject, on
average). No significant differences were found in the number of excluded trials across
conditions (2-way ANOVA; decision type: $F(1,17)=3.31$, $p=0.09$; decision difficulty:
$F(1,17)=1.83$, $p=0.19$; interaction: $F(1,17)=0.42$, $p=0.53$).

The EEG was segmented by locking the waveforms to subjects' movement onset, starting 2s
prior to the movement and ending 0.2s afterwards, with the segments averaged separately for
each decision type (Deliberate/Arbitrary x Easy/Hard) and decision content (right/left). The
baseline period was defined as the time window between -1000 ms and -500 ms prior to
stimulus onset, that is, the onset of the causes screen, rather than prior to movement onset. In
addition to the main baseline, we tested another baseline—from -1000 ms to -500 ms relative
to *movement* onset—to investigate whether the baseline period influenced our main results (see
Results). Furthermore, we segmented the EEG based on *stimulus* onset, using the same
baseline, for stimulus-locked analysis (again, see Results).

To assess potential effects of eye movements during the experiment, we defined the radial eye
signal as the average over all 4 EOG channels, when band-pass filtered to between 30 and 100
Hz. We then defined a saccade as any signal that was more than 2.5 standardized IQRs away
from the median of the radial signal for more than 2 ms. Two consecutive saccades had to be at
least 50 ms apart. The saccade count (SC) was the number of saccades during the last 500 ms
before response onset (Keren, Yuval-Greenberg, & Deouell, 2010) (see also (Croft & Barry,
2000; Elbert, Lutzenberger, Rockstroh, & Birbaumer, 1985; Shan, Moster, & Roemer, 1995)).

Statistical Analysis

EEG differences greater than expected by chance were assessed using two-way ANOVAs with
decision type (deliberate, arbitrary) and decision difficulty (easy, hard), using IBM SPSS
statistics, version 24. For both RP and LRP signals, the mean amplitude from 500 ms before to
button-press onset were used for the ANOVAs. Greenhouse–Geisser correction was never
required as sphericity was never violated (Picton et al., 2000).

Trend analysis on all subjects' data was carried out by regressing the voltage for every subject
against time for the last 1000 ms before response onset using first-order polynomial linear
regression (see Results). We used every 10th time sample for the regression (i.e., the 1st, 11th,
21st, 31st samples, and so on) to conform with the individual-subject analysis (see below). For
the individual-subject analysis, the voltage on all trials was regressed against time in the same
manner (i.e., for the last 1000 ms before response onset and using first-order polynomial linear
regression). As individual-trial data is much noisier than the mean over all trials in each
subject, we opted for standard robust-regression using iteratively reweighted least squares
(implemented using the *robustfit()* function in Mathworks Matlab). The iterative robust-
regression procedure is time consuming. So, we used every 10th time sample instead of every
sample to make the procedure's run time manageable. Also, as EEG signals have a 1/f power
spectrum, taking every 10th sample further better conforms with the assumption of i.i.d. noise
in linear regression.

Model and Simulations

All simulations were performed using Mathworks Matlab 2014b. The model was devised off
the one proposed by Schurger et al. (2012). Like them, we built a drift-diffusion model
(Ratcliff, 1978; Usher & McClelland, 2001), which included a leaky stochastic accumulator
(with a threshold on its output) and a time-locking/epoching procedure. The original model
amounted to iterative numerical integration of the differential equation

$$\delta x_i = (I - k x_i)\Delta t + c\xi_i\sqrt{\Delta t} \tag{1}$$

where I is the drift rate, k is the leak (exponential decay in x), ξ is Gaussian noise, and c is a
noise-scaling factor (we used $c = 0.05$). Δt is the discrete time step used in the simulation (we
used $\Delta t = 0.001$, similar to our EEG sampling rate). The model integrates x_i until it crosses a
threshold, which represents a decision having been made.

In such drift-diffusion models, for a given k and c, I and the threshold together determine how
quickly a decision will be reached, on average. If we further fix the threshold, a higher drift
rate, I, represents a faster decision, on average. (The drift rate alone can thus be viewed as a
constant "urgency to respond" (using the original Schurger term) that is inherent in the demand
characteristics of the task, evidenced by the fact that no subject took more than 20 s to make a
decision on any trial. The leak term, k, ensures that the model would not be too linear; i.e., it
prevented the drift rate from setting up a linear trajectory for the accumulator toward the
threshold. Hence, due to the leak term, doubling the magnitude of the threshold would make
the accumulator rarely reach the threshold, instead of reaching it in roughly twice the amount
of time (up to the noise term).

Our model differed from Schurger's in two main ways. First, it accounted for both arbitrary
and deliberate decisions and was built to fit our empirical results. It was thus composed of two
components, each generally described by Eq. (1), but with different parameter values. The first
accumulated activity that drove arbitrary decisions (i.e., random fluctuations (Schurger et al.,
2012)). We term it the *Noise* component. The second component drove deliberate decisions
based on subjects' values associated with the decision alternatives. We term it the *Value*
components. Our model mainly relied on its noise component for arbitrary decisions and on its
value one for deliberate decisions.

Second, Schurger and colleagues modeled only the decision *when* to move (during arbitrary
decisions). But our subjects decided both *when* and *which hand* to move. So, we had to extend
the Schurger model in that respect as well. We did this using a race-to-threshold mechanism
between the two decision alternatives. In our empirical paradigm, the difference in rating of the
two causes was either 1 (for hard decisions) or 4-6 (for easy decisions; see "Experimental
Procedure" in Methods), so there was always an alternative that was ranked higher than the
other. Choosing the higher- or lower-ranked alternative was termed a congruent or incongruent
choice with respect to the initial ratings, respectively. Hence, we modeled each decision the
subjects made as a race to threshold between the congruent and incongruent alternatives in the
noise component (for arbitrary decisions) or value component (for deliberate ones).

Using a parameter sweep, we found the values of the thresholds, drift rate, and leak that best fit
our average empirical reaction times for (easy, hard) x (deliberate, arbitrary) decisions as well
as our empirical consistency ratios for those 4 decision types. The model's reaction time was
defined as the overall time that it took until the first threshold crossing in the race-to-threshold

pair (again, each step took $\Delta t = 0.001$ s). We used the same threshold value of 0.15 and leak
value of $k=0.5$ for all model types. The only parameter that was modulated across (deliberate,
arbitrary) x (easy, hard) decisions x (congruent, incongruent) decision alternatives was the drift
rate, I (Table 1). All of these parameters were then fixed when we used the model to derive the
simulated Cz activity across all conditions.

Table 1: Values of the drift-rate parameter across decision types. Values of the
drift-rate parameter, I, in our model across (deliberate, arbitrary) x (easy, hard)
decisions x (congruent, incongruent) decision alternatives.

Drift rate (I) values	Congruent		Incongruent	
	Easy	Hard	Easy	Hard
Deliberate	0.0400	0.1010	0.0228	0.0000
Arbitrary	0.1650	0.1648	0.1650	0.1566

Each simulation consisted of either 120 runs of the model, the same as the number of empirical
trials per condition, or 10000 runs of the model for a smoother reaction-time distribution for
the model (see Results). For each run of the model, we identified the first threshold crossing
point and extracted the last second (1000 steps) before the crossing in each run. If the first
crossing was earlier than sample no. 1,000 by $n > 0$ samples, we padded the beginning of the
epoch with n null values (NaN or "not-a-number" in Matlab). These values did not contribute
to the average across simulated trials, so the simulated average RP became noisier at earlier
time points in the epoch. Hence, our model was similarly limited to the Schurger model in its
inability to account for activity earlier than the beginning of the trial (see Results).

References

Banks, W. P., & Isham, E. A. (2009). We infer rather than perceive the moment we decided to
act. *Psychological Science, 20*(1), 17.
Bode, S., & Haynes, J. D. (2009). Decoding sequential stages of task preparation in the human
brain. *Neuroimage, 45*(2), 606-613.
Bode, S., He, A., Soon, C., Trampel, R., Turner, R., & Haynes, J. (2011). Tracking the
unconscious generation of free decisions using uitra-high field fMRI. *PLoS One, 6*(6),
e21612.
Brainard, D. H. (1997). The psychophysics toolbox. *Spatial vision, 10*, 433-436.
Brass, M., & Haggard, P. (2008). The what, when, whether model of intentional action. *The
Neuroscientist, 14*(4), 319.
Breitmeyer, B. G. (1985). Problems with the psychophysics of intention. *Behavioral and brain
sciences, 8*(04), 539-540.
Bush, G., Luu, P., & Posner, M. I. (2000). Cognitive and emotional influences in anterior
cingulate cortex. *Trends in Cognitive Sciences, 4*(6), 215-222.
Carter, C. S., Braver, T. S., Barch, D. M., Botvinick, M. M., Noll, D., & Cohen, J. D. (1998).
Anterior cingulate cortex, error detection, and the online monitoring of performance.
Science, 280(5364), 747-749.
Croft, R. J., & Barry, R. J. (2000). Removal of ocular artifact from the EEG: a review.
Neurophysiologie Clinique/Clinical Neurophysiology, 30(1), 5-19.
Deutschländer, R., Pauen, M., & Haynes, J.-D. (2017). Probing folk-psychology: Do Libet-
style experiments reflect folk intuitions about free action? *Consciousness and
cognition, 48*, 232-245.

Eimer, M. (1998). The lateralized readiness potential as an on-line measure of central response
activation processes. *Behavior Research Methods, Instruments, & Computers, 30*(1),
146-156.
Elbert, T., Lutzenberger, W., Rockstroh, B., & Birbaumer, N. (1985). Removal of ocular
artifacts from the EEG—a biophysical approach to the EOG. *Electroencephalography*
and Clinical Neurophysiology, 60(5), 455-463.
Fischer, J. M. (1999). Recent work on moral responsibility. *Ethics, 110*(1), 93-139.
Frankfurt, H. G. (1971). Freedom of the Will and the Concept of a Person. *The Journal of*
Philosophy, 68(1), 5-20.
Fried, I., Mukamel, R., & Kreiman, G. (2011). Internally generated preactivation of single
neurons in human medial frontal cortex predicts volition. *Neuron, 69*, 548-562.
Frith, C., Blakemore, S., & Wolpert, D. (2000). Abnormalities in the awareness and control of
action. *Philosophical Transactions of the Royal Society B: Biological Sciences,*
355(1404), 1771.
Gold, J., & Shadlen, M. (2007). The neural basis of decision making.
Haggard, P. (2005). Conscious intention and motor cognition. *Trends in Cognitive Sciences,*
9(6), 290-295.
Haggard, P. (2008). Human volition: towards a neuroscience of will. *Nature Reviews*
Neuroscience, 9(12), 934-946.
Haggard, P., & Eimer, M. (1999). On the relation between brain potentials and the awareness
of voluntary movements. *Experimental Brain Research, 126*(1), 128-133.
Hallett, M. (2007). Volitional control of movement: The physiology of free will. *Clinical*
Neurophysiology, 118(6), 1179-1192.
Hallett, M. (2016). Physiology of free will. *Annals of neurology, 80*(1), 5-12.
Harris, S. (2012). *Free will.* New York, NY: Simon & Schuster, Inc.
Hobbes, T. (1994). *Leviathan: with selected variants from the Latin edition of 1668*: Hackett
Publishing Company.
Hughes, G., Schütz-Bosbach, S., & Waszak, F. (2011). One action system or two? Evidence
for common central preparatory mechanisms in voluntary and stimulus-driven actions.
Journal of Neuroscience, 31(46), 16692-16699.
Jeannerod, M. (2006). *Motor cognition: What actions tell the self*: Oxford University Press.
Jeffreys, H. (1998). *The theory of probability*: OUP Oxford.
Junghofer, M., Elbert, T., Tucker, D., & Rockstroh, B. (2000). Statistical control of artifacts in
dense array EEG/MEG studies. *Psychophysiology, 37*(04), 523-532.
Kable, J. W., & Glimcher, P. W. (2009). The neurobiology of decision: consensus and
controversy. *Neuron, 63*(6), 733-745.
Keren, A. S., Yuval-Greenberg, S., & Deouell, L. Y. (2010). Saccadic spike potentials in
gamma-band EEG: characterization, detection and suppression. *Neuroimage, 49*(3),
2248-2263.
Kornhuber, H., & Deecke, L. (1990). Readiness for movement—the Bereitschafts potential-
story. *Current Contents Life Sciences, 33*(4), 14.
Krieghoff, V., Waszak, F., Prinz, W., & Brass, M. (2011). Neural and behavioral correlates of
intentional actions. *Neuropsychologia, 49*(5), 767-776.
Lau, H., Rogers, R., Haggard, P., & Passingham, R. (2004). Attention to intention. *Science,*
303(5661), 1208.
Lau, H., Rogers, R., & Passingham, R. (2007). Manipulating the experienced onset of intention
after action execution. *Journal of cognitive neuroscience, 19*(1), 81-90.
Libet, B. (1985). Unconscious cerebral initiative and the role of conscious will in voluntary
action. *Behavioral and brain sciences, 8*, 529-539.

Libet, B., Gleason, C., Wright, E., & Pearl, D. (1983). Time of conscious intention to act in
relation to onset of cerebral activity (readiness-potential): The unconscious initiation of
a freely voluntary act. *Brain, 106*(3), 623.
Libet, B., Wright, E., & Gleason, C. (1982). Readiness-potentials preceding unrestricted
'spontaneous' vs. pre-planned voluntary acts. *Electroencephalography and Clinical*
Neurophysiology, 54(3), 322-335.
Maoz, U., Mudrik, L., Rivlin, R., Ross, I., Mamelak, A., & Yaffe, G. (2015). On reporting the
onset of the intention to move. In A. R. Mele (Ed.), *Surrounding Free Will:*
Philosophy: Psychology, Neuroscience: Oxford University Press.
Maoz, U., Rutishauser, U., Kim, S., Cai, X., Lee, D., & Koch, C. (2013). Predeliberation
activity in prefrontal cortex and striatum and the prediction of subsequent value
judgment. *Frontiers in neuroscience, 7*, 225.
Maoz, U., & Yaffe, G. (2016). What does recent neuroscience tell us about criminal
responsibility? *Journal of Law and the Biosciences, 3*(1), 120-139.
Masaki, H., Wild-wall, N., Sangals, J., & Sommer, W. (2004). The functional locus of the
lateralized readiness potential. *Psychophysiology, 41*(2), 220-230.
Mele, A. (2006). *Free will and luck*: Oxford University Press.
Mele, A. (2009). *Effective intentions: the power of conscious will*: Oxford University Press,
USA.
Mele, A. (2015). *Surrounding free will: philosophy, psychology, neuroscience*: Oxford
University Press.
Miller, J., Shepherdson, P., & Trevena, J. (2011). Effects of Clock Monitoring on
Electroencephalographic Activity. *Psychological Science, 22*(1), 103-109.
Nahmias, E., Shepard, J., & Reuter, S. (2014). It's OK if 'my brain made me do it': People's
intuitions about free will and neuroscientific prediction. *Cognition, 133*(2), 502-516.
Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics: Transforming
numbers into movies. *Spatial vision, 10*(4), 437-442.
Perez, O., Mukamel, R., Tankus, A., Rosenblatt, J. D., Yeshurun, Y., & Fried, I. (2015).
Preconscious prediction of a driver's decision using intracranial recordings. *Journal of*
cognitive neuroscience.
Picton, T., Bentin, S., Berg, P., Donchin, E., Hillyard, S., Johnson, R., . . . Rugg, M. (2000).
Guidelines for using human event-related potentials to study cognition: recording
standards and publication criteria. *Psychophysiology, 37*(02), 127-152.
Pockett, S., Banks, W. P., & Gallagher, S. (2009). *Does consciousness cause behavior?* : Mit
Press.
Ramnani, N., & Owen, A. M. (2004). Anterior prefrontal cortex: insights into function from
anatomy and neuroimaging. *Nature Reviews Neuroscience, 5*(3), 184-194.
Ratcliff, R. (1978). A theory of memory retrieval. *Psychological Review, 85*(2), 59-108.
Roskies, A. (2010). How Does Neuroscience Affect Our Conception of Volition? *Annual*
Review of Neuroscience, 33, 109-130.
Roskies, A. L. (2012). How does the neuroscience of decision making bear on our
understanding of moral responsibility and free will? *Current Opinion in Neurobiology,*
22(6), 1022-1026.
Salvaris, M., & Haggard, P. (2014). Decoding Intention at Sensorimotor Timescales. *PLoS*
One, 9(2), e85100.
Sanfey, A., Loewenstein, G., McClure, S., & Cohen, J. (2006). Neuroeconomics: cross-
currents in research on decision-making. *Trends in Cognitive Sciences, 10*(3), 108-116.
Sanfey, A. G., Rilling, J. K., Aronson, J. A., Nystrom, L. E., & Cohen, J. D. (2003). The neural
basis of economic decision-making in the ultimatum game. *Science, 300*(5626), 1755-
1758.

Schmidt, S., Jo, H.-G., Wittmann, M., & Hinterberger, T. (2016). 'Catching the waves'–slow
cortical potentials as moderator of voluntary action. *Neuroscience & Biobehavioral*
Reviews, 68, 639-650.
Schurger, A. (2018). Specific Relationship between the Shape of the Readiness Potential,
Subjective Decision Time, and Waiting Time Predicted by an Accumulator Model with
Temporally Autocorrelated Input Noise. *eNeuro, 5*(1), ENEURO. 0302-0317.2018.
Schurger, A., Sitt, J. D., & Dehaene, S. (2012). An accumulator model for spontaneous neural
activity prior to self-initiated movement. *Proceedings of the National Academy of*
Sciences, 109(42), E2904-E2913.
Shan, Y., Moster, M. L., & Roemer, R. A. (1995). The effects of time point alignment on the
amplitude of averaged orbital presaccadic spike potential (SP).
Electroencephalography and Clinical Neurophysiology, 95(6), 475-477.
Shibasaki, H., & Hallett, M. (2006). What is the Bereitschaftspotential? *Clinical*
Neurophysiology, 117(11), 2341-2356.
Sinnott-Armstrong, W. (2014). *Moral Psychology: Free Will and Moral Responsibility* (Vol.
4): Mit Press.
Sinnott-Armstrong, W., & Nadel, L. (Eds.). (2011). *Conscious will and responsibility: A*
tribute to Benjamin Libet: Oxford University Press.
Sirigu, A., Daprati, E., Ciancia, S., Giraux, P., Nighoghossian, N., Posada, A., & Haggard, P.
(2004). Altered awareness of voluntary action after damage to the parietal cortex.
Nature Neuroscience, 7(1), 80-84.
Soon, C., Brass, M., Heinze, H., & Haynes, J. (2008). Unconscious determinants of free
decisions in the human brain. *Nature Neuroscience, 11*(5), 543-545.
Soon, C., He, A., Bode, S., & Haynes, J. (2013). Predicting free choices for abstract intentions.
Proceedings of the National Academy of Sciences, 110(15), 6217-6222.
Strawson, G. (1994). The impossibility of moral responsibility. *Philosophical Studies, 75*(1), 5-
24.
Thut, G., Hauert, C.-A., Viviani, P., Morand, S., Spinelli, L., Blanke, O., . . . Michel, C.
(2000). Internally driven vs. externally cued movement selection: a study on the timing
of brain activity. *Cognitive Brain Research, 9*(3), 261-269.
Trevena, J., & Miller, J. (2010). Brain preparation before a voluntary action: Evidence against
unconscious movement initiation. *Consciousness and cognition, 19*(1), 447-456.
Tusche, A., Bode, S., & Haynes, J.-D. (2010). Neural responses to unattended products predict
later consumer choices. *Journal of Neuroscience, 30*(23), 8024-8031.
Ullmann-Margalit, E., & Morgenbesser, S. (1977). Picking and choosing. *Social Research,*
44(4), 757-783.
Usher, M., & McClelland, J. L. (2001). The time course of perceptual choice: the leaky,
competing accumulator model. *Psychological Review, 108*(3), 550.
Verleger, R., Haake, M., Baur, A., & Śmigasiewicz, K. (2016). Time to Move Again: Does the
Bereitschaftspotential Covary with Demands on Internal Timing? *Frontiers in human*
neuroscience, 10.
Wallis, J. D. (2007). Orbitofrontal cortex and its contribution to decision-making. *Annu. Rev.*
Neurosci., 30, 31-56.
Wallis, J. D., & Miller, E. K. (2003). Neuronal activity in primate dorsolateral and orbital
prefrontal cortex during performance of a reward preference task. *Eur J Neurosci,*
18(7), 2069-2081. doi:2922 [pii]
Waszak, F., Wascher, E., Keller, P., Koch, I., Aschersleben, G., Rosenbaum, D. A., & Prinz,
W. (2005). Intention-based and stimulus-based mechanisms in action selection.
Experimental Brain Research, 162(3), 346-356.
Wegner, D. (2002). *The illusion of conscious will*: MIT Press.

Wisniewski, D., Goschke, T., & Haynes, J.-D. (2016). Similar coding of freely chosen and
externally cued intentions in a fronto-parietal network. *Neuroimage, 134*, 450-458.
Wolf, S. (1990). *Freedom within reason*: Oxford University Press.

Acknowledgments: We thank Ralph Adolph for his invaluable guidance and support in
designing and running the experiment as well as for very useful discussions of the results. We
thank Ram Rivlin for various conceptual discussions about deliberate versus arbitrary decision-
making and about the initial experimental paradigm design. We thank Caitlin Duncan for her
help in patiently and meticulously gathering the EEG data. We thank Daw-An Wu for
discussions about EEG data collection and preprocessing and for his help with actual data
collection. We thank Daniel Grossman for his help in carefully preprocessing the data and
suggesting potential interpretations of it. We thank Ueli Rutishauser for discussions about the
model and its simulations. We thank Shlomit Yuval-Greenberg and Leon Deouell for
important discussions about EEG processing and analysis. Last, we thank the anonymous
reviewers for their invaluable comments, which greatly improved this manuscript. **Funding:**
This research was supported by Florida State University's Big Questions in Free Will Initiative,
funded by the John Templeton Foundation, to U.M., G.Y., and C.K.; by the Ralph Schlaeger
Charitable Foundation to U.M.; by the Bial Foundation to U.M. and to U.M. and L.M.; and by
the German-Israeli Foundation for Scientific Research and Development to L.M.. C.K. thanks
the Allen Institute founders, Paul G. Allen and Jody Allen, for their vision, encouragement,
and support. **Author contributions:** U.M, L.M, G.Y., and C.K. conceived the project and
designed the experiments. L.M. and U.M. analyzed the results. U.M. designed and simulated
the model. L.M. and U.M. wrote the manuscript. G.Y. and C.K. suggested revisions to the
manuscript. **Competing interests:** The authors declare that they have no competing interests.
Data and materials availability: All data needed to evaluate the conclusions in the paper are
present in the paper. Additional data related to this paper may be requested from the authors.

1053

1054   Neural precursors of decisions that matter—an ERP study of deliberate and arbitrary choice

1055   Maoz U, Yaffe G, Koch C, and Mudrik L

1056   Supplementary Data

1057   **Supplementary Table 1: NPO names and causes acronyms**

NPO	Cause	NPO website
Consensual NPOs		
American Society on Aging	Pro Quality of Life for the Elderly	http://asaging.org/
Conservation Fund	Pro Environment protection	http://www.conservationfund.org/
Bill & Melinda Gates Foundation	Pro Education	http://www.gatesfoundation.org/
Global Fund for Women	Pro Women's Rights	https://www.globalfundforwomen.org/
The Hunger Project	Pro Hunger Relief	https://www.thp.org/
Oxfam International	Pro Poverty & Disaster Relief	http://www.oxfam.org/
World Wild Life Fund (WWF)	Pro Species Conservation	http://worldwildlife.org/
Cancer Research Institute	Pro Cancer Research	http://www.cancerresearch.org/
Habitat for Humanity	Pro Housing for All	http://www.habitat.org/
Reading is Fundamental	Pro Advancement of Literacy	http://www.rif.org/
International Institute for Conservation of Historic and Artistic Works	Pro Culture & Arts Preservation	https://www.iiconservation.org/
Big Brothers and Big Sisters of America	Pro Youth Development	http://www.bbbs.org/site/c.9iILI3NGKhK6F/ b.5962335/k.BE16/Home.htm
United Nations Children's Fund (UNICEF)	Pro Child Protection	http://www.unicef.org/
Doctors without Borders (Medecins sans frontieres)	Pro Disaster Medical Care	http://www.msf.org/
Soldiers' Angels	Pro Veterans & Military	http://www.soldiersangels.org/heroes/index.php
Disability Rights International	Pro Disabilities Rights	http://www.disabilityrightsintl.org/

National Crime Prevention Council (NCPC)	Pro Crime Prevention	http://www.ncpc.org/
Amnesty International	Pro Human Rights	https://www.amnesty.org/
Peace Corps	Pro Peace & Development	http://www.peacecorps.gov/
World Health Organization	Pro World Health	http://www.who.int/en/
Controversial NPOs		
Planned Parenthood	Pro Abortion & Family Planning	http://www.plannedparenthood.org/
Pro-Life Alliance	Anti Abortion & Family Planning	http://www.prolifealliance.com/
Human Rights Campaign	Pro LBGTQ Rights	http://www.hrc.org/
National Organization for Marriage	Anti LBGTQ Rights	https://www.nationformarriage.org/
Stem for Life Foundation	Pro Stem Cell Research	http://www.stemforlife.org/
Christian Dental & Medical Association	Anti Stem Cell Research	http://www.cmda.org/
Greenpeace	Pro Action Against Climate Change	http://www.greenpeace.org/international/en/
Global Climate Scam	Anti Action Against Climate Change	http://www.globalclimatescam.com/
National Association for Gun Rights	Pro Gun Rights	http://www.nationalgunrights.org/
Coalition to Stop Gun Violence	Pro Gun Control	http://csgv.org/
American Gas Association	Pro Fracking for Natural Gas	http://www.aga.org/Pages/default.aspx
Americans Against Fracking	Anti Fracking for Natural Gas	http://www.americansagainstfracking.org/
StandWithUs (Israel)	Pro Israel	http://www.standwithus.com/
Palestinian Centre for Human Rights	Pro Palestine	http://www.pchrgaza.org/portal/en/
National Organization for the Reform of Marijuana Laws	Pro Marijuana Legalization	http://norml.org/
Citizens Against Legalizing Marijuana	Anti Marijuana Legalization	http://www.calmca.org/
Understanding Animal Research	Pro Scientific Experiments on Animals	http://www.understandinganimalresearch.org.uk/

International Association Against Painful Experiments on Animals	Anti Scientific Experiments on Animals	http://www.iaapea.com/
Federation for American Immigration Reform	Pro Immigration Reform	http://www.fairus.org/
American Immigration Control	Anti Immigration Reform	http://www.immigrationcontrol.com/
Human Cloning Foundation	Pro Human Cloning	http://www.humancloning.org/
Americans to Ban Cloning	Anti Human Cloning	http://www.cloninginformation.org/
Americans United for Separation of Church and State	Pro Separation of Church & State	https://www.au.org/
Christian Coalition of America	Anti Separation of Church & State	http://www.cc.org/
Death with Dignity National Center	Pro Euthanasia (Assisted Suicide)	http://www.deathwithdignity.org/
Euthanasia Prevention Coalition	Anti Euthanasia (Assisted Suicide)	http://www.epcc.ca/
The Alliance for Better Foods	Pro Genetically Modified Foods	http://www.betterfoods.org/
Non-GMO Project	Anti Genetically Modified Foods	http://www.nongmoproject.org/
Answers in Genesis	Pro Creationism Teaching	https://answersingenesis.org
National Center for Science Education	Pro Evolution Teaching	http://ncse.com/

1058

1059

1060

1061