Title

- Neural precursors of decisions that matter—an ERP study of deliberate and arbitrary choice

Authors

U. Maoz[1,2,3]*, G. Yaffe[4], C. Koch[5], L. Mudrik[6]

Affiliations

[1] Department of Psychology and Brain Institute, Chapman University, Orange, CA, USA.

[2] Department of Psychology, University of California, Los Angeles, CA, USA.

[3] Division of Humanities and Social Sciences, California Institute of Technology, Pasadena, CA, USA.

[4] Yale Law School, Yale University, New Haven, CT, USA.

[5] Allen Institute for Brain Science, Seattle, WA, USA.

[6] School of Psychological Science and Sagol School of Neuroscience, Tel Aviv University, Tel Aviv, Israel.

* urim@caltech.edu

Abstract

The onset of the readiness potential (RP)—a key neural correlate of upcoming action—was repeatedly found to precede subjects' reports of having made an internal decision. This has been taken by some as evidence against a causal role for consciousness in human decision-making and thus as a denial of free-will. Yet those studies focused on purposeless, unreasoned, arbitrary decisions, bereft of consequences. It remains unknown to what degree these specific neural precursors of action generalize to deliberate decisions, which are more ecological and relevant to real life. We therefore directly compared the neural correlates of deliberate and arbitrary decision-making during a $1000-donation task to non-profit organizations. While we found the expected RPs for arbitrary decisions, they were strikingly absent for deliberate ones. Our results are congruent with the RP representing the accumulation of noisy, random fluctuations, which drive arbitrary—but not deliberate—decisions. The absence of RPs in deliberate decisions challenges the generalizability of studies that argue for no causal role for consciousness in decision making from arbitrary to deliberate, real-life decisions.

MAIN TEXT

Introduction

Humans typically experience freely selecting between alternative courses of action, say, when ordering a particular item off a restaurant menu. Yet a series of human studies using EEG (*1-3*), fMRI (*4-7*), intracranial (*8*), and single-cell recordings (*9*) challenged the validity of this common experience, finding neural correlates of decision processes hundreds of milliseconds and even

seconds prior to the moment that subjects reported having consciously decided. Some have claimed, following these and other findings, that the subjective human experience of freely deciding is but an illusion, because human actions are unconsciously initiated before the conscious decision to act (*1, 10*). This debate has been captivating scholars from many disciplines in and outside of academia (*10-15*).

Critically, in the above studies, subjects were either told to move their right hand or flex their right wrist; or they were instructed to move either the right or left hand. (*13, 16, 17*). Thus, their decisions were always unreasoned, purposeless, and bereft of any real consequence. This stands in sharp contrast to real-life decisions that are reasoned, purposeful, and bear consequences (*18*)—which clothes to wear, what route to take to work, as well as more formative decisions about life partners, career choices, and so on. Such deliberate decisions are also at the center of the philosophical debate on free will (*16, 19*). They typically involve more conscious and lengthy deliberation, and might thus be more tightly bound to conscious processes.

Deliberate decisions have been widely studied in the field of neuroeconomics (*20, 21*) and in perceptual tasks (*22*). Yet, interestingly, little has been done to assess the relation between decision-related activity, subjects' conscious experience of deciding, and the neural activity instantaneously contributing to this experience. Though some studies compared, for example, internally driven and externally cued decisions (e.g., *23, 24*), these were typically arbitrary decisions that had no real implications. Here, we compared neural precursors of deliberate and arbitrary decisions, on the same subjects, in an EEG experiment. Our experiment utilized a donation-preference paradigm, in which a pair of non-profit organizations (NPOs) were presented in each trial. In deliberate-decision trials, subjects chose to which NPO they would like to donate $1000. In arbitrary-decision trials, both NPOs received an equal donation of $500, irrespective of subjects' key presses (Fig. 1). Notably, while the visual inputs and motor outputs were identical between deliberate and arbitrary decisions, the decisions' meaning for the subjects was radically different: in deliberate blocks, the decisions were meaningful and consequential—reminiscent of important, real-life decisions—while in arbitrary blocks, the decisions were meaningless and bereft of consequences—mimicking previous studies of volition.

Results

Behavioral Results

To validate the experimental paradigm, we manipulated decision difficulty as well as decision type (Fig. 1). We reasoned that this should affect only deliberate decisions, and not arbitrary ones. This was born out in our subjects' behavioral patterns. Subjects' reaction times (RTs) were substantially slower for deliberate than for arbitrary decisions (Fig. 2, left; $F_{(1,17)}=126.11$, $p<0.0001$ for the main effect of decision type, as revealed by a 2-way ANOVA with decision type (arbitrary/deliberate) and difficulty (easy/hard). The ANOVA also showed a main effect of decision difficulty, $F_{(1,17)}=18.76$, $p=0.0004$; all analyses were performed on log-transformed data, because the raw RTs violated the normality assumption ($W=0.94$, $p=0.001$)). Moreover, subjects were slower for hard vs. easy deliberate decisions ($F_{(1,17)}=20.12$, $p=0.0003$ for the interaction between decision type and decision difficulty; hard vs. easy deliberate decisions: $t_{(17)}=4.78$, $p=0.0002$, yet not for arbitrary decisions: $t_{(17)}=1.01$, $p=0.325$). This further demonstrates that, in deliberate decisions, subjects were making meaningful decisions, affected by the difference in the values of the two NPOs, while in arbitrary decisions they were not.

Figure 1: Experimental paradigm. The experiment included deliberate (red, left) and arbitrary (blue, right) blocks, each containing nine trials. In each trial, two NPO names were presented, and subjects were asked to either choose to which NPO they would like to donate (deliberate), or to simply press either right or left, as both NPOs would receive an equal donation (arbitrary). Within each block, some of the trials were easy (lighter colors) decisions, where the subject's preferences for the two NPOs substantially differed (based on a previous rating session), and some were hard decisions (darker colors), where the preferences were more similar (easy and hard trials were intermixed within each block). To make sure subjects were paying attention to the NPO names, even in arbitrary trials, and to better equate the cognitive load between deliberate and arbitrary trials, memory tests (in light grey) were randomly introduced, where subjects were asked to determine which of four NPO names appeared in the immediately previous trial.

The consistency between subjects' choices throughout the main experiment and the NPO ratings they gave prior to the main experimental session was also analyzed (see Methods). As expected, subjects were highly consistent with their own, previous ratings when making deliberate decisions, but not when making arbitrary ones (Fig. 2, right; 2-way ANOVA: $F_{(1,17)}=946.55$, $p<0.0001$) for the main effect of decision type. A main effect of decision difficulty was also found ($F_{(1,17)}=57.39$, $p<0.0001$). Again, decision type and decision difficulty interacted ($F_{(1,17)}=25.96$, $p<0.0001$: subjects were much more consistent with their choices in easy vs. hard deliberate decisions ($t_{(17)}=11.15$, $p<0.0001$), than they were in easy vs. hard arbitrary decisions ($t_{(17)}=2.50$, $p=0.028$). Nevertheless, though subjects were around chance (i.e., 0.5) in their consistency in arbitrary decisions (ranging between 0.39 to 0.64), it seems that some subjects were slightly influenced by their preferences in easy-arbitrary decisions trials, resulting in the significant difference between hard-arbitrary and easy-arbitrary decisions above. Finally, no differences were found between subjects' tendency to press the right vs. left button in the different conditions (both main effects and interaction: $F<1$).

Figure 2: Behavioral results. Response Times (RTs; left) and Consistency Grades (CG; right) in arbitrary (blue) and deliberate (red) decisions. Each dot represents the average RT/CG for easy and hard decisions for an individual subject (hard decisions: x-coordinate; easy decisions: y-coordinate). Group means and SEs are represented in dark red and dark blue crosses. The histograms at the bottom-left corner of each plot sum the number of dots with respect to the solid diagonal line. The dashed diagonal line represents equal RT/CG for easy and hard decisions; data points below that diagonal indicate longer RTs or higher CGs for hard decisions. In both measures, arbitrary decisions are more centered around the diagonal than deliberate decisions, showing no or substantially reduced differences between easy and hard decisions.

EEG Results: Readiness Potential (RP)

The RP, generally held to index unconscious readiness for upcoming movement (*1, 13, 25, 26*) (though see (*27-29*)), was measured over electrode Cz in the different conditions by averaging the activity across trials in the 2 s prior to subjects' movement. Focusing on the last 500 ms before movement onset for our statistical tests, we found a clear RP in arbitrary decisions, yet RP amplitude was not significantly different from 0 in deliberate decisions (Fig. 3A; ANOVA F(1,17)=11.86, p=0.003 for the main effect of decision type; in t-tests against zero, corrected for multiple comparisons, an effect was only found for arbitrary decisions (hard: t(17)=5.75, p<0.0001; easy: t(17)=5.09, p=0.0004) and not for deliberate ones (hard: t(17)=1.24, p>0.5; easy: t(17)=1.84, p=0.34). In a similar manner, regressing voltage against time for the last 1000 ms before movement onset, the downward trend was significant for arbitrary decisions (p<0.0001, for both easy and hard conditions) but not for deliberate decisions (hard: p>0.5, easy: p=0.35; all Bonferroni corrected for multiple comparisons). Notably, this pattern of results was also manifested for single-subject analysis (Fig. S1; 14 of the 18 subjects had significant downward slopes for arbitrary decisions—i.e., p<0.05, Bonferroni corrected for multiple comparisons— when regressing voltage against time for every trial over the last 1000 ms before movement onset; but only 5 of the 18 subjects had significant downward slopes for the same regression analysis for deliberate decisions; see Methods. In addition, the average slopes for deliberate and arbitrary decisions were -0.43±0.31 and -2.30±0.44 (mean±SE), respectively, a significant difference: t(17)=3.51, p=0.001).

Readiness Potentials

Figure 3: The readiness potentials for deliberate and arbitrary decisions. (A) Mean and SE of the Readiness Potential (RP) in deliberate (red shades) and arbitrary (blue shades) easy and hard decisions in electrode Cz, as well as scalp distributions. Zero refers to time of right/left response made by the subject. Notably, the RP significantly differs from zero and displays a typical scalp distribution for arbitrary decisions only. The scalp distribution was calculated over the averaged activity during the last 500 ms before response, across subjects. The inset shows the mean amplitude of the RP, with 95% confidence intervals over the same time window. **(B)** Individual subjects' Cz activity in the four conditions (n=18). The linear-regression line for voltage against time over the last 1000 ms before movement onset is designated by a dashed, black line. Note that the waveforms converge to an RP only in arbitrary decisions.

To test whether the null result for RP amplitude in deliberate decisions stems from a genuine absence of effect or from insufficient or underpowered data, we used Bayesian statistics. Specifically, the Bayes factor allowed us to compare the probability of observing the data given H_0 (i.e., no RP in deliberate decisions) against the probability of observing the data given H_1 (i.e., RP exists in deliberate decisions). We followed the convention that a BF < 0.33 implies substantial evidence for lack of an effect (that is, the data is three times more likely to be observed given H_0 than given H_1), 0.33 < BF < 3 suggests insensitivity of the data, and BF > 3 denotes substantial evidence for the presence of an effect (H_1) (*30*). We found strong evidence that arbitrary trials (pooled across difficultly for more statistical power) are different from 0 (BF=2.815·10^6), but inconclusive evidence for deliberate trials (BF=1.457, again for pooled trials). Notably, however, this result might reflect the general, slow negative trend of the deliberate trials data (Fig. 3A) rather than a typical RP, which is locked to decision-onset and shows a sharp decline towards that. This difference might therefore be due to differences in RTs between the conditions. To examine whether such a decline occurred in deliberate decisions and to remove the effect of this slow negative drift, we used a baseline period of -1000ms to -500ms relative to *movement onset* (i.e., a baseline that immediately preceded our time of interest window) instead of our usual baseline that preceded *stimulus* onset (see Methods). Under this analysis, we found evidence that deliberate decisions are not different from 0 (BF=0.332), supporting the claim that the RP during the last 500 ms before movement onset was completely absent (BF for arbitrary decisions was 5.07·10^4).

Differences in reaction times (RT) between conditions do not drive the effect

RTs in deliberate decisions were typically more than twice as long as RTs in arbitrary decisions. We therefore wanted to rule out the possibility that the absence of RP in deliberate decisions stemmed from the difference in RTs between the conditions. We carried out two analyses for this purpose. First, we ran a median split analysis—dividing the subjects into two groups based on their RTs: lower and higher than the median, for deliberate and arbitrary trials, respectively. We then ran the same analysis using only the faster subjects in the deliberate condition (M=1.91s, SD=0.25) and the slower subjects in the arbitrary condition (M=1.25s, SD=0.23). If RT length affects RP amplitudes, we would expect the RP amplitudes to be more similar between these two groups. However, though there were only half the data points, a similar pattern of results to those over the whole dataset was observed (Fig. S2A). Deliberate and arbitrary decisions were still reliably different ($F_{(1,17)}$=5.22, p=0.03), with significant RPs found in arbitrary (easy: t(8)=4.57, p=0.0018; hard: t(8)=4.09, p=0.0035), but not deliberate (easy: t(8)=1.92, p=0.09; hard: t(8)=0.63, p=0.54) decisions. In addition, the RPs for arbitrary decisions were not significantly different between the subjects with above-median RTs and the entire population for the easy or hard conditions (easy: t(25)=0.14, p>0.5; hard: t(25)=0.56, p>0.5). Similarly, the RPs for deliberate decisions were not significantly different between the subjects with below-median RTs and the entire population for the easy or hard conditions (easy: t(25)=-0.34, p>0.5; hard: t(25)=0.17, p>0.5). This suggest that RTs do not reliably affect Cz activation for deliberate or arbitrary decisions in our results.

Second, we regressed the difference between RPs in deliberate and arbitrary decisions (averaged over the last 500 ms before movement onset) against the difference between the RTs in these two conditions for each subject (Fig. S2B). Again, if RT length affects RP amplitudes, we would expect differences between RTs in deliberate and arbitrary conditions to correlate with differences between RPs in the two conditions. But no correlation was found between the two measures (r=0.22, T(17)=0.86, p=0.4). We further tried regressing the RP differences on RT differences. The regression did not produce any reliable relation between RT and RP differences (regression line: y = 0.54 [CI -0.8, 1.89] x - 0.95 [CI -2.75, 0.85]; the R^2 was very low, at 0.05 (as expected from the r value above), and, as the confidence intervals suggest, the slope was not significantly different from 0, $F_{(1,17)}$=0.74, p=0.4). Taken together, these results provide strong evidence against the claim that the difference in RPs stems from or is affected by the difference in RTs between the conditions.

Eye movements do not affect the results

Though ICA was used to remove blink artifacts and saccades (see Methods), we wanted to make sure our results do not stem from differential eye movement patterns between the conditions. We therefore computed a saccade-count metric (SC; see Methods) for each trial for all subjects. Focusing again on the last 500 ms before movement onset, we computed mean (± s.e.m.) SC values of 1.65±0.07 and 1.67±0.06 saccades for easy and hard deliberate decisions, respectively, and 1.69±0.07 and 1.73±0.07 saccades for easy and hard arbitrary decisions, respectively. We found no reliable differences between the number of saccades during deliberate and arbitrary trials (ANOVA $F_{(1,17)}$=2.56, p=0.13 for the main effect of decision type).

We further investigated potential effects of saccades by running a median-split analysis—dividing the trials for each subject into two groups based on their SC score: lower and higher than the median, for deliberate and arbitrary trials, respectively. We then ran the same analysis using only the trials with more saccades in the deliberate condition (SC was 2.02±0.07 and 2.04±0.07 for easy and hard, respectively) and those with less saccades for the arbitrary condition (SC was 1.33±0.07 and 1.31±0.08 for easy and hard, respectively). If the number of saccades affects RP

amplitudes, we would expect that the differences in RPs between arbitrary and deliberate trials will diminish, or even reverse (as now we had more saccades in the deliberate condition). However, though there were only half the data points for each subject in each condition, a similar pattern of results to those over the whole dataset was observed (Fig. S3): Deliberate and arbitrary decisions were still reliably different within the median-split RPs ($F(1,17)=16.70$, $p<0.001$), with significant RPs found in arbitrary (easy: $t(17)=4.79$, $p=0.002$; hard: $t(17)=5.77$, $p<0.001$), but not deliberate (easy: $t(17)=0.90$, $p=0.38$; hard: $t(17)=0.30$, $p>0.5$) decisions. In addition, we compared the RP data across all the trials with the median-split RP data above. No significant differences were found for arbitrary decisions (easy: $t(17)=1.02$, $p=0.32$; hard: $t(17)=0.75$, $p=0.46$) or for deliberate decisions (easy: $t(17)=1.63$, $p=0.12$; hard: $t(17)=1.47$, $p=0.16$). Taken together, the analyses above provide strong evidence against the involvement of eye movements in our results.

Testing other factors that could potentially account for the results

We further tested whether our results depend on the hand with which subjects executed the movement. Again, the same pattern of results persisted when separating left-hand button presses from right-hand ones (Fig. S4). Second, we took a closer look at subjects' behavior in the easy arbitrary condition, where some subjects had a consistency score that was further above 0.5 (chance) than others. It seems like those subjects had a greater difficulty ignoring their preferences, despite the instructions to do so. We therefore wanted to test to what extent the RP of those subjects was similar to the RPs of the other subjects. Focusing on the 8 subjects that had a consistency score above 0.55 (M=0.59, SD=0.03) and comparing their RPs to those of the 10 other subjects (consistency M=0.50, SD=0.06) in easy arbitrary trials, we found no reliable differences ($t(16)=0.94$, $p=0.36$). This is not surprising, as the mean consistency score of these subjects – though indeed higher than chance – was still far below their consistency score for easy deliberate decisions (M=0.99, SD=0.02).

EEG Results: Lateralized Readiness Potential (LRP)

The LRP, which reflects activation processes within the motor cortex for action preparation after response selection (*31, 32*), was measured by subtracting the difference potentials (C3-C4) in right-hand response trials from this difference in left-hand responses trials and averaging the activity over the same time window (*2, 31*). In this purely motor component, no difference was found between the two decision types (Fig. S5; ANOVA all Fs<1).

Drift Diffusion Model (DDM)

The main finding of this study—the absence of RP in deliberate decisions – is in line with recent work that used a Drift-diffusion model (DDM) to claim that the RP might arise from time-locking neural activity to movement onset rather than from specific preparation for or ballistic-like initiation of movement (*27*). DDMs of decision-making typically feature a process that rises toward a threshold. The crossing of that threshold reflects the onset of the decision in the model, possibly leading to action. Schurger and colleagues (*27*) modelled arbitrary decisions, and suggested that there the threshold crossing leading to movement onset is largely determined by spontaneous subthreshold fluctuations of the neural activity. This challenged the common view of the RP as a neural correlate of unconscious preparation for upcoming action (*26*). Instead, Schurger and colleagues claimed, time-locking to movement onset ensures that these spontaneous fluctuations appear, when averaged over many trials, as a gradual increase in neural activity.

To further assess this interpretation of the RP, we expanded the model developed by Schurger and colleagues (*27*) to a DDM that was composed of a *value-assessment* component and a *noise-generation* component. Under this assumption, Cz-electrode activity mainly reflects the noise-generation component. Each trial was further modeled as a race to threshold between two alternatives: one where the subject chose the NPO that was rated higher in the earlier rating session (the *congruent* option) and the other where the subject preferred the lower-rated NPO (the *incongruent* option). Each race was represented as a leaky stochastic accumulator (see Methods for details and model parameters).

Figure 4: Empirical and model RTs and model prediction for Cz activity. (A) The empirical distributions of subjects' RTs across the four decision types. **(B)** The equivalent distributions of RTs for the model. **(C)** The model's prediction for the ERP activity in electrode Cz across all four decision types.

We fit our DDM to our average empirical reaction-times, which were 2.13, 2.52, 0.98 and 1.00 s for the different conditions (henceforth, magnitudes are given for deliberate easy, deliberate hard, arbitrary easy, and arbitrary hard, respectively). The model's corresponding mean RTs were 2.04, 2.46, 0.94, and 0.96 s for these conditions (Fig. 4A, B). The model was further fit to the empirical consistency ratios (the proportions of congruent decisions), which were 0.99, 0.83, 0.54 and 0.49. The model's corresponding consistency ratios were 1.00, 0.84, 0.53 and 0.53. The model then predicted the shape of the ERP in its noise component (assumed to reflected by Cz-electrode activity) for each decision type: a continuing, RP-like increase in activity (with a negative sign) for arbitrary decisions, but only a very slight increase in activity for deliberate decisions (Fig. 4C; see also Fig. S6). This was in line with our empirical results (Fig. 3A). Note that that the Schurger

model aims to account for neural activity leading up to the decision to move, but no further (*27*). Similarly, we expect our DDM to fit Cz neural data only up to around -0.1 s (100 ms before movement onset). We also make no claims that ours is the only, or even optimal, model that explains our results. Rather, by extending the Schurger model, we show how that interpretation of the RP could also be applied to our more-complex paradigm.

Discussion

Since the publication of Libet's seminal work claiming that neural precursors of action, in the form of the RP, precede subjects' reports of having consciously decided to act (*1*), a vigorous discussion has been raging between neuroscientists, philosophers, and other scholars about the meaning of these findings for the debate on free will (recent collections include (*33-35*)). Some claim that these results have removed conscious will from the causal chain leading to action (*13, 36, 37*). Others are unconvinced that these results are decisive for, or even applicable to, the free-will debate (*16, 19, 38*). At the heart of much of this debate lies the RP, thought to represent unconscious decision/planning mechanisms that initiate subjects' decisions prior to their conscious experience of deciding (*1, 25*).

Notably, the RP and similar findings showing neural activations preceding the conscious decision to act have typically been based on arbitrary decisions of different types (*1, 2, 4, 5, 14, 39, 40*). This, among other reasons, rested on the notion that for an action to be completely free, it should not be determined in any way by external factors (*41*)—which is the case for arbitrary, but not deliberate, decisions (where each decision alternative is associated with a value, and the values of alternatives typically guide one's decision). But this notion of freedom faces several obstacles. First, most discussions of free will focus on deliberate decisions, asking when and whether these are free (*42-44*). This might be because everyday decisions to which we associate freedom of will—like choosing a more expensive but more environmentally friendly car, helping a friend instead of studying more for a test, donating to charity, and so on—are generally deliberate, in the sense of being reasoned, purposeful, and bearing consequences (although see (*45*)). In particular, the free will debate is often considered in the context of moral responsibility (e.g., was the decision to harm another person free or not) (*13, 46-50*), and free will is even sometimes defined as the capacity that allows one to be morally responsible (*38, 51*). In contrast, it seems meaningless to assign blame or praise to arbitrary decisions. Thus, though the scientific operationalization of free will has typically focused on arbitrary decisions, the common interpretations of these studies—in neuroscience and across the free will debate—have often alluded to deliberate ones. Here, we show that inference from arbitrary to deliberate decisions may not be justified, as the neural precursors of arbitrary decisions do not generalize to meaningful ones (*16, 19*). Interestingly, while the RP was present in arbitrary decisions but absent in deliberate ones, the LRP—a long-standing, more-motor ERP component—was indistinguishable between the different decision types. This provides evidence that, at the motor level, the neural representation of the deliberate and arbitrary decisions that our subjects made may have been indistinguishable, as was our intention when designing the task.

Our finding and the model thus suggest that two different mechanisms may be involved in arbitrary and deliberate decisions. Earlier literature demonstrated that deliberate, reasoned decision-making—which was mostly studied in the field of neuroeconomics (*20*) or using perceptual decisions (*22*)—elicited activity in the prefrontal cortex (PFC; mainly the dorsolateral (DLPFC) part (*52, 53*) and ventromedial (VMPFC) part/orbitofrontal cortex (OFC) (*54, 55*) and the anterior cingulate cortex (ACC) (*56, 57*). Arbitrary, meaningless decisions, in contrast, were mainly probed using variants of the Libet paradigm, showing activations in the Supplementary Motor Area (SMA), alongside other frontal areas like the medial frontal cortex (*58, 59*) or the

frontopolar cortex, as well as the posterior cingulate cortex (*4, 9*) (though see (*60*), which suggests that a common mechanism may underlie both decision types). Possibly then, arbitrary and deliberate decisions may differ not only in respect to the RP, but may be subserved by different underlying neural circuits, which makes generalization from one class of decisions to the other more difficult. Deliberate decisions are associated with more lateralized and central neural activity while arbitrary ones are associated with more medial and frontal ones. This appears to align with the different brain regions associated with the two decision types above, as also evidenced by the differences we found between the scalp distributions of arbitrary and deliberate decisions (Fig. 3A).. Future studies are needed to further explore this potential divergence in the neural regions between the two decision types.

To be clear, and following the above, we do not claim that the RP captures all unconscious processes that precede conscious awareness. In deliberate decisions in particular, there may be neural precursors of upcoming actions—possibly involving the above neural circuits as well as circuits that represents values—which are unrelated to the RP. In addition, we did not attempt to separately measure the timing of subjects' conscious decision to move, but rather instructed subjects to press the buttons as soon as they consciously made up their minds. This is both to keep the decisions in this task more ecological and because we think that the key method of measuring decision onset (using some type of clock to measure Libet's W-time) is highly problematic (see Methods). Alternatively, some have claimed that the RP represents unconscious motor-preparatory activity before any kind of decision. But our results provide evidence against that claim, as we do not find an RP before deliberate decisions.

Aside from highlighting the differences between arbitrary and deliberate decisions, this study also challenges a common interpretation of the function of the RP. If the RP is not present before deliberate action, it does not seem to be a necessary link in the general causal chain leading to action. Schurger and colleagues (*27*) suggested that the RP reflects the accumulation of stochastic fluctuations in neural activity that lead to action following a threshold crossing when humans arbitrarily decide to move. The shape of the RP therefore results of the manner in which it is computed: averaged over trials that are locked to movement onset (that directly follows the threshold crossing). Our results and our model are in line with that interpretation and expand upon it, suggesting that the RP represents the accumulation of noisy, random fluctuations that drive arbitrary decisions, while deliberate decisions are mainly driven by the values associated with the decision alternatives (*61*). Our DDM was based on the assumption that every decision is driven by a component based on the values of the decision alternatives (the subject's support for the two NPOs in our case) and by another component representing noise—random fluctuations in neural activity. The value component plays little to no role in arbitrary decisions, so action selection and timing depend on when the accumulation of noise crosses the decision threshold for the congruent and incongruent decision alternatives. In deliberate decisions, in contrast, the value component drives the decisions, while the noise has a smaller effect. Thus, in arbitrary decisions, action onset closely tracks threshold crossings of the noise accumulation. But, in deliberate decisions, the noise component is at more random levels at movement onset. Hence, locking the ERP to movement onset and averaging over trials to obtain the RP, leads to a relatively flat signal for deliberate decisions but to the expected RP shape in arbitrary decisions. This provides strong evidence that the RP does not reflect subconscious movement preparation. Rather, it is induced by threshold crossing of random fluctuations in arbitrary decisions, that do not drive deliberate decisions; accordingly, the RP is not found there. Further studies of the causal role of consciousness in deliberate versus arbitrary decisions are required to test this claim.

Another, alternative explanation of our results is that—in the deliberate condition only—the NPO names act as a cue, thereby turning what we term internal, deliberate decisions into no more than

simple responses to external stimuli. Under this account, if the preferred NPO is on the right, it is immediately interpreted as "Press right". It would therefore follow that the absence of decisions in deliberate trials results in the absence of the RP in those trials. However, the reaction time and consistency values that we observed provide evidence against this interpretation. We found longer reaction times for hard-deliberate decisions than for easy-deliberate ones (2.52 versus 2.13 s, on average, respectively; see also Fig. 2) and higher consistencies with the initial ratings for easy-deliberate decisions than for hard-deliberate decisions (0.99 versus 0.83, on average, respectively; see also Fig. 2). If the NPO names acted as mere cues, we would have expected no differences between reaction times or consistencies for easy- and hard-deliberate decisions. In addition, there were 50 different causes in the first part of the experiment. So, it is highly unlikely that subjects could memorize all 1225 pairwise preferences among these causes and simply transform any decision between a pair of causes into a stimulus instructing to press left or right.

Yet another interpretation of our results might be that subjects engage in (unconscious) deliberation also during arbitrary decisions (*62*). If so, the RP in the arbitrary decisions might reflect the extra effort required to break the symmetry under this condition. However, this interpretation entails a longer reaction time for arbitrary than for deliberate decisions, because of the heavier cognitive load, which is the opposite of what we found (Fig. 2A). We would also expect the simpler deliberate in arbitrary-easy trials to result in a shorter reaction-time than that of arbitrary-hard. But this is not what we find (Fig. 2A).

Our study, which suggests that RPs do not precede deliberate decisions, further motivates future investigations into other precursors of action besides the RP using EEG, fMRI or other techniques. It would be of interest to find the neural activity that precedes deliberate decisions in particular. And it would also be of interest to find neural activity, which is not motor activity, that is common to both deliberate and arbitrary decisions.

Materials and Methods

Study Design

The objective of this study was to compare ERPs elicited by arbitrary and deliberate decision-making. We further manipulated decision difficulty to validate our manipulation of decisions type: we introduced hard and easy decisions which corresponded to small and large differences between subjects' preferences for the pairs of presented NPOs, respectively. We reasoned that if the manipulation of decision type (arbitrary vs. deliberate) was effective, there would be behavioral differences between easy and hard decisions for deliberate choices but not for arbitrary choices (because differences in preferences should not influence subjects' arbitrary decisions). Our 2 x 2 design was therefore decision type (arbitrary/deliberate) by decision difficulty (easy/hard). Each condition included 90 trials divided into 10 blocks of 9 trials each, leading to a total of 360 trials and 40 blocks. Blocks of different decision types were randomly intermixed. Decision difficulty was randomly counterbalanced across trials within each block.

In a preliminary study that we ran, we found that a sample size of at least 16 subjects results in a clear, averaged RP. So, we decided beforehand to collect 20 subjects for this study, taking into account that some could be excluded as they would not meet the following predefined inclusion criteria: at least 30 trials per experimental condition left after artifact rejection; and averaged RTs (across conditions) which deviated by 3 or less standard deviations from the group mean. Subjects were informed about the overall number of subjects that would participate in the experiment when the NPO lottery was explained to them (see below). So, we had to finalize the overall number of subjects who would participate in the study—but not necessarily the overall number of subjects

whose data would be part of the analysis—before the experiment began. After completing data collection, we ran only the EEG preprocessing and behavioral-data analysis to test each subject against the exclusion criteria. This was done before we looked at the data with respect to our hypothesis or research question. Two subjects did not meet the inclusion criteria: the data of one subject (#18) suffered from poor signal quality which resulted in less than 30 trials after artifact rejection; another subject (#12) had RTs longer than 3 standard deviations from the mean. All analyses were thus run on the 18 remaining subjects.

We further excluded single trials in which subjects pressed the wrong button as well as trials where subjects' RTs were less than 200ms, more than 10s, or more than 3 standard deviations away from that subject's mean in that condition (mean number of excluded trials =7.17, SD=2.46, which are 1.99% of the trials).

Subjects

Twenty healthy subjects participated in the study. They were California Institute of Technology (Caltech) students as well as members of the Pasadena community. All subjects had reported normal or corrected-to-normal sight and no psychiatric or neurological history. They volunteered to participate in the study for payment ($20 per hour). Subjects were prescreened to include only participants who were socially involved and active in the community (based on strength of their support of social causes, past volunteer work, past donations to social causes, and tendency to vote). The data from 18 subjects was analyzed; two subjects were excluded from our analysis (see Study Design above). The experiment was approved by Caltech's Institutional Review Board, and informed consent was obtained from all participants after the experimental procedures were explained to the subjects.

Stimuli and apparatus

Subjects sat in a dimly lit room. The stimuli were presented on a 21" Viewsonic G225f (20" viewable) CRT monitor with a 60-Hz refresh rate and a 1024×768 resolution using Psychtoolbox version 3 and Mathworks Matlab 2014b (*63, 64*). They appeared with a gray background (RGB values: [128, 128,128]). The screen was located 60 cm away from subjects' eyes. Stimuli included names of 50 real non-profit organizations (NPOs). Twenty organizations were consensual (e.g., the Cancer Research Institute, or the Hunger project), and thirty were more controversial: we chose 15 causes that are widely debated (e.g., pro/anti guns, pro/anti abortions), and selected one NPO that supports each of the two sides of the debate. This was done to achieve variability in subjects' willingness to donate to the different NPOs. In the main part of the experiment, succinct descriptions of the causes (e.g., pro-marijuana legalization, pro-child protection; for a full list of NPOs and causes see Table S1) were presented in black Comic Sans MS.

Experimental Procedure

In the first part of the experiment, subjects were presented with each NPO separately. They were instructed to rate how much they would like to support that NPO with a $1000 donation on a scale of 1 ("I would not like to support this NPO at all) to 7 ("I would very much like to support this NPO"). No time pressure was put on the subjects, and they were given access to the website of each NPO to give them the opportunity to learn more about the NPO and the cause it supports.

After the subjects finished rating all NPOs, the main experiment began. In each block of the experiment, subjects made either deliberate or arbitrary decisions. Two succinct cause descriptions, representing two actual NPOs, were presented in each trial (Fig. 1). In deliberate

blocks, subjects were instructed to choose the NPO to which they would like to donate $1000 by pressing the <Q> or <P> key on the keyboard, for the NPO on the left or right, respectively, as soon as they decided. Subjects were informed that at the end of each block one of the NPOs they chose would be randomly selected to advance to a lottery. Then, at the end of the experiment, the lottery will take place and the winning NPO will receive a $20 donation. In addition, that NPO will advance to the final, inter-subject lottery, where one subject's NPO will be picked randomly and will be given a $1000 donation. It was stressed that the donations were real and that no deception was used in the experiment. To persuade the subjects that the donations are real, we presented a signed commitment to donate the money, and promised to send them the donation receipts after the experiment. Thus, subjects knew that in deliberate trials, every choice they made was not hypothetical, and could potentially lead to an actual $1020 donation to their chosen NPO.

Arbitrary trials were identical to deliberate trials except for the following crucial differences. Subjects were told that, at the end of each block, the pair of NPOs in one randomly selected trial would advance to the lottery together. And, if that pair wins the lottery, both NPOs would receive $10 each. Further, the NPO pair that would win the inter-subject lottery would receive a $500 donation each. Hence it was stressed to the subjects that there was no reason for them to prefer one NPO over the other in arbitrary blocks, as both NPOs would receive the same donation regardless of their button press. Subjects were told to therefore simply press either <Q> or <P> when they felt the urge to do so.

Thus, while subjects' decisions in the deliberate blocks were meaningful and consequential, their decisions in the arbitrary blocks had no impact on the final donations that were made. In these trials, subjects were further urged not to let their preferred NPO dictate their response. Note that we did not ask subjects to report their "W-time" (time of consciously reaching a decision), because this measure was shown to rely on neural processes occurring after movement onset (65) and to potentially be backward inferred from movement time (66). Even more importantly, clock monitoring was demonstrated to have an effect on RP size (67), so it could potentially confound our results (68).

In addition, we manipulated decision difficulty (Easy/Hard) throughout the experiment, randomly intermixed within each block. Decision difficulty was determined based on the rating difference between the two presented NPOs. NPO pairs with 1 or 4 or more rating point difference were designated hard or easy, respectively. Based on each subject's ratings, we created a list of NPO pairs, half of each were easy choices and the other half hard choices.

Each block started with an instruction written either in dark orange (Deliberate: "In this block choose the cause to which you want to donate $1000") or in blue (Arbitrary: "In this block both causes may each get a $500 donation regardless of the choice"). Short-hand instructions appeared at the top of the screen throughout the block in the same colors as that block's initial instructions; Deliberate: "Choose for $1000" or Arbitrary: "Press for $500 each" (Fig. 1). Each trial started with a fixation cross, with a duration drawn from a uniform distribution between 1 and 1.5s. Then, the two cause-descriptions appeared on the left and right side of the fixation cross (left/right assignments were randomly counterbalanced), and remained on the screen until the subjects responded with a key press. The cause corresponding to the pressed button then turned white, and a new trial started. If subjects did not respond within 20s, they received an error message and were informed that if this trial would be selected for the lottery, no NPO would receive a donation. However, this did not happen for any subject on any trial.

To assess the consistency of subjects' decisions during the main experiment with their ratings in the first part of the experiment, subjects' choices were coded in the following way: each binary

choice in the main experiment was given a consistency grade of 1, if subjects chose the NPO that was rated higher in the rating session, and 0 if not. Then a consistency grade was calculated as the mean over all the choices. A consistency grade of 1 indicates perfect consistency with one's ratings, 0 – perfect inconsistency, and 0.5 – chance performance.

To better equate memory load, attention, and other cognitive aspects between deliberate and arbitrary decisions—except those aspects directly associated with the decision type, which was the focus of our investigation—we wanted to make sure subjects were carefully reading and remembering the causes also during the arbitrary trials. We therefore randomly interspersed 36 memory catch-trials throughout the experiment (thus more than one catch trial could occur per block). On such trials, four succinct descriptions of causes were presented and subjects had to select the one that appeared in the previous trial. A correct or incorrect response added or subtracted 50 cents from their total, respectively. (Subjects were informed that if they reached a negative balance, no money will be deducted off their payment for participation in the experiment.) Thus, subjects could earn $18 more for the experiment, if they answered all memory test questions correctly. Subjects typically did well on these memory questions, on average erring in 2.5 out of 36 memory catch trials (7% error) and gaining an additional $16.75 (SD=3.19).

ERP recording methods

The EEG was recorded using an Active 2 system (BioSemi, the Netherlands) from 64 electrodes distributed based on the extended 10–20 system and connected to a cap, and seven external electrodes. Four of the external electrodes recorded the EOG: two located at the outer canthi of the right and left eyes and two above and below the center of the right eye. Two external electrodes were located on the mastoids, and one electrode was placed on the tip of the nose. All electrodes were referenced during recording to a common-mode signal (CMS) electrode between POz and PO3. The EEG was continuously sampled at 512 Hz and stored for offline analysis.

ERP analysis

ERP analysis was conducted using the "Brain Vision Analyzer" software (Brain Products, Germany) and in-house Mathworks Matlab scripts. Data from all channels were referenced offline to the average of all channels. The data were then digitally high-pass filtered at 0.1 Hz using a Finite Impulse Response (FIR) filter to remove slow drifts. A notch filter at 59-61 Hz was applied to the data to remove 60-Hz electrical noise. The signal was then cleaned of blink and saccade artifacts using Independent Component Analysis (ICA) (*69*). Signal artifacts were detected as amplitudes exceeding ±100 µV, differences beyond 100 µV within a 200 ms interval, or activity below 0.5 mV for over 100 ms (the last condition was never found). Sections of EEG data that included such artifacts in any channel were removed (150ms before and after the artifact), leaving an average number of 70.38 trials with a range of 36-86 out of 90 trials per condition. Channels that consistently had artifacts were replaced using interpolation (4.2 channels per subject, on average).

The EEG was segmented by locking the waveforms to subjects' decision onset, starting 2s prior to the decision and ending 0.2s afterwards, with the segments averaged separately for each decision type (Deliberate/Arbitrary x Easy/Hard) and decision content (right/left). The baseline period was defined as the time window between -1000ms and -500ms prior to the beginning of the trial (i.e., prior to *stimulus* onset), rather than prior to decision onset. It is in particular suitable here because the differences in reaction times between deliberate and arbitrary decisions make baselines locked to decision times inappropriate.

To assess eye movements during the experiment, we defined the radial eye signal as the average over all 4 EOG channels, when band-pass filtered to between 30 and 100 Hz. We then defined a saccade as any signal that was more than 2.5 standardized IQRs away from the median of the radial signal for more than 2 ms. Two consecutive saccades had to be at least 50 ms apart. The saccade count (SC) was the number of saccades during the last 500 ms before movement onset (*70*).

Statistical Analysis

Differences greater than expected by chance were assessed using two-way ANOVAs with decision type (deliberate, arbitrary) and decision difficulty (easy, hard), using IBM SPSS statistics, version 24. For both RP and LRP signals, the mean amplitude from 500 ms before to button-press onset were used for the ANOVAs. Greenhouse–Geisser correction was never required as sphericity was never violated (*71*).

Trend analysis on all subjects' data (Fig. 3B) was carried out by regressing the voltage for every subject against time for the last 1000 ms before movement onset using first-order polynomial linear regression. We used every 10^{th} time sample for the regression (i.e., the 1^{st}, 11^{th}, 21^{st}, 31^{st} samples, and so on) to conform with the individual-subject analysis (see below). For the individual-subject analysis, the voltage on all trials was regressed against time in the same manner (i.e., for the last 1000 ms before movement onset and using first-order polynomial linear regression). As individual-trial data is much noisier than the mean over all trials in each subjects, we opted for standard robust-regression using iteratively reweighted least squares (implemented using the *robustfit()* function in Mathworks Matlab). The iterative robust-regression procedure is time consuming. So, we used every 10^{th} time sample instead of every sample to make the procedure's run time manageable.

Model and Simulations

All simulations were performed using Mathworks Matlab 2014b. The model was devised off the one proposed by Schurger and colleagues (*27*). They built a drift-diffusion model (*72, 73*) for arbitrary decisions only, which included a leaky stochastic accumulator (with a threshold on its output) and a time-locking/epoching procedure. Their model amounted to iterative numerical integration of the differential equation

$$\delta x_i = (I - kx_i)\Delta t + c\xi_i\sqrt{\Delta t}$$

where *I* is the drift rate, *k* is the leak (exponential decay in *x*), ξ is Gaussian noise, and *c* is a noise-scaling factor (we used *c* = 0.05). *Δt* is the discrete time step used in the simulation (we used *Δt* = 0.001, similar to our EEG sampling rate). Here *I* represents a constant urgency to respond that is inherent in the demand characteristics of the task, evidenced by the fact that no subject took more than 20 s to decide on any trial. The leak term, *k*, ensures that the model would not be too linear; i.e., it prevents the urgency from setting up a linear trajectory for the accumulator toward the threshold. Hence, due to the leak term, doubling the magnitude of the threshold would make the accumulator rarely reach the threshold, instead of reaching it in roughly twice the amount of time (up to the noise term) without a leak term.

Our model accounted for both arbitrary and deliberate decisions and was built to fit our empirical results, based on with two Schurger-like components. The first one accumulated activity that drove arbitrary decisions (i.e., random fluctuations (*27*)). The second component drove deliberate

decisions based on subjects' values associated with the decision alternatives. Henceforth we term these *Noise* and *Value* components for ease of description. Our model used its Noise component for arbitrary decisions and its Value one for deliberate decisions.

Schurger and colleagues modeled only the decision when to move. But our subjects decided both when and which hand to move. So, we had to extend the Schurger model in that respect as well. We did this using a race-to-threshold mechanism between the two decision alternatives. In our paradigm, the difference in rating was either 1 (hard decisions) or 4-6 (easy decisions; see "Procedure & Experimental Design" in Methods), so there was always an alternative that was ranked higher than the other. Choosing the higher or lower alternative was termed a congruent or incongruent choice with the initial ratings, respectively.

Using a parameter sweep, we found the values of the thresholds, urgency, and leak that best fit our average empirical reaction times for (easy, hard) x (deliberate, arbitrary) decisions as well as our empirical consistency ratios for those 4 decision types. The model's reaction time was defined as the overall time (where each step took $\Delta t = 0.001$ s) that it took until the first threshold crossing in the race-to-threshold pair. We used the same threshold value of 0.15 and leak value of $k=0.5$ for all model types. The only parameter that was modulated across (deliberate, arbitrary) x (easy, hard) decisions x (congruent, incongruent) decision alternatives was the urgency, I (Table 1). These parameters were then fixed when fitting the simulated Cz activity across all conditions.

Each simulation consisted of either 120 runs of the model, the same as the number of empirical trials per condition (Fig. 4C), or 10000 runs of the model for a smoother reaction-time distribution for the model (Fig. 4B). For each run of the model, we identified the first threshold crossing point and extracted the last second (1000 steps) before the crossing in each run. If the first crossing was earlier than sample no. 1,000 by $n > 0$ samples, we padded the beginning of the epoch with n null values (NaN or "not-a-number" in Matlab). These values did not contribute to the average across simulated trials, so the simulated average RP became noisier at earlier time points in the epoch. Our model was therefore similarly limited to the Schurger model in its inability to account for activity earlier than the beginning of the trial. (Fig. 3C).

Urgency (*I*) values	Congruent		Incongruent	
	Easy	Hard	Easy	Hard
Deliberate	0.0400	0.1010	0.0228	0.0000
Arbitrary	0.1650	0.1648	0.1650	0.1566

Table 1: Values of the urgency parameter across decision types. Values of the urgency parameter, *I*, in our model across (deliberate, arbitrary) x (easy, hard) decisions x (congruent, incongruent) decision alternatives.

Supplementary Materials

Figure S1: Individual-subjects RPs.
Figure S2: Relations between RTs and RPs.
Figure S3: Relations between eye movements and RPs.
Figure S4: RPs for left-hand and right-hand trials.
Figure S5: Lateralized readiness potential.
Figure S6: Model runs.
Table S1: NPO names and causes

References and Notes

1. B. Libet, C. A. Gleason, E. W. Wright, D. K. Pearl, Time of conscious intention to act in relation to onset of cerebral activity (readiness-potential). The unconscious initiation of a freely voluntary act. *Brain* **106**, 623-642 (1983).
2. P. Haggard, M. Eimer, On the relation between brain potentials and the awareness of voluntary movements. *Experimental Brain Research* **126**, 128-133 (1999).
3. M. Salvaris, P. Haggard, Decoding Intention at Sensorimotor Timescales. *PloS one* **9**, e85100 (2014).
4. C. S. Soon, M. Brass, H. J. Heinze, J. D. Haynes, Unconscious determinants of free decisions in the human brain. *International Journal of Psychology* **43**, 238-238 (2008).
5. C. S. Soon, A. H. He, S. Bode, J. D. Haynes, Predicting free choices for abstract intentions. *Proceedings of the National Academy of Sciences* **110**, 6217-6222 (2013).
6. S. Bode, C. Bogler, J. D. Haynes, Similar neural mechanisms for perceptual guesses and free decisions. *Neuroimage* **65**, 456–465 (2012).
7. S. Bode *et al.*, Predicting perceptual decision biases from early brain activity. *The Journal of Neuroscience* **32**, 12488-12498 (2012).
8. O. Perez *et al.*, Preconscious prediction of a driver's decision using intracranial recordings. *Journal of cognitive neuroscience*, (2015).
9. I. Fried, R. Mukamel, G. Kreiman, Internally generated preactivation of single neurons in human medial frontal cortex predicts volition. *Neuron* **69**, 548-562 (2011).
10. D. M. Wegner, *The Illusion of Conscious Will*. (MIT Press, Cambridge, Mass, 2002).
11. C. D. Frith, D. M. Wolpert, Abnormalities in the awareness and control of action. *Philosophical Transactions of the Royal Society B: Biological Sciences* **355**, 1771-1788 (2000).
12. M. Jeannerod, *Motor cognition: What actions tell the self.* (Oxford University Press, Oxford, 2006), vol. 42.
13. P. Haggard, Human volition: towards a neuroscience of will. *Nat. Rev. Neurosci.* **9**, 934-946 (2008).
14. H. Lau, R. Rogers, P. Haggard, R. E. Passingham, Attention to intention. *Science* **303**, 1208-1210 (2004).
15. A. R. Mele, *Free will and luck*. (Oxford University Press, 2006).
16. A. Roskies, How Does Neuroscience Affect Our Conception of Volition? *Annual Review of Neuroscience* **33**, 109-130 (2010).
17. M. Hallett, Physiology of free will. *Annals of neurology* **80**, 5-12 (2016).
18. E. Ullman-Margalit, S. Morgenbesser, Picking and choosing. *Social Research* **44**, 757-785 (1977).
19. B. G. Breitmeyer, Problems with the psychophysics of intention. *Behavioral and brain sciences* **8**, 539-540 (1985).
20. P. W. Glimcher, C. Camerer, E. Fehr, R. A. Poldrack, Eds., *Neuroeconomics: Decision making and the brain*, (Academic Press, London, 2009).
21. A. G. Sanfey, G. Loewenstein, S. M. McClure, J. D. Cohen, Neuroeconomics: cross-currents in research on decision-making. *Trends Cogn. Sci.* **10**, 108-116 (2006).
22. J. I. Gold, M. N. Shadlen, The neural basis of decision making. *Annu. Rev. Neurosci.* **30**, 535-574 (2007).
23. W. D., T. Goschke, J. D. Haynes, Similar coding of freely chosen and externally cued intentions in a fronto-parietal network. *Neuroimage* **134**, 450-458 (2016).
24. G. Thut *et al.*, Internally driven vs. externally cued movement selection: a study on the timing of brain activity. *Cognitive Brain Research* **9**, 261-269 (2000).

25.   H. H. Kornhuber, L. Deecke, Readiness for movement—the Bereitschafts potential-story. *Current Contents Life Sciences* **33**, 14 (1990).

26.   H. Shibasaki, M. Hallett, What is the Bereitschaftspotential? *Clinical Neurophysiology* **117**, 2341-2356 (2006).

27.   A. Schurger, J. D. Sitt, S. Dehaene, An accumulator model for spontaneous neural activity prior to self-initiated movement. *Proceedings of the National Academy of Sciences* **109**, E2904-E2913 (2012).

28.   J. Trevena, J. Miller, Brain preparation before a voluntary action: Evidence against unconscious movement initiation. *Consciousness and cognition* **19**, 447-456 (2010).

29.   J. Miller, P. Shepherdson, J. Trevena, Effects of clock monitoring on electroencephalographic activity is unconscious movement initiation an artifact of the clock? *Psychological Science* **22**, 103-109 (2011).

30.   H. Jeffreys, *The theory of probability*. (OUP Oxford, 1998).

31.   M. Eimer, The lateralized readiness potential as an on-line measure of central response activation processes. *Behavior Research Methods, Instruments, & Computers* **30**, 146-156 (1998).

32.   H. Masaki, N. Wild-wall, J. Sangals, W. Sommer, The functional locus of the lateralized readiness potential. *Psychophysiology* **41**, 220-230 (2004).

33.   A. R. Mele, Ed., *Surrounding Free Will: Philosophy, Psychology, Neuroscience*, (Oxford University Press, Oxford, 2015).

34.   W. Sinnott-Armstrong, L. Nadel, Eds., *Conscious will and responsibility: A tribute to Benjamin Libet*, (Oxford University Press, 2011).

35.   S. Pockett, W. P. Banks, S. Gallagher, *Does consciousness cause behavior?* , (Mit Press, 2009).

36.   P. Haggard, Conscious intention and motor cognition. *Trends Cogn. Sci.* **9**, 290-295 (2005).

37.   D. Wegner, *The illusion of conscious will*. Bradford Books (MIT Press, 2002).

38.   A. Mele, *Effective intentions: the power of conscious will*. (Oxford University Press, USA, 2009).

39.   B. Libet, Subjective antedating of a sensory experience and mind-brain theories: reply to Honderich (1984). *J Theor Biol* **114**, 563-570 (1985).

40.   A. Sirigu *et al.*, Altered awareness of voluntary action after damage to the parietal cortex. *Nature Neuroscience* **7**, 80-84 (2003).

41.   B. Libet, Unconscious cerebral initiative and the role of conscious will in voluntary action. *Behavioral and brain sciences* **8**, 529-539 (1985).

42.   H. G. Frankfurt, Freedom of the Will and the Concept of a Person. *The Journal of Philosophy* **68**, 5-20 (1971).

43.   T. Hobbes, in *Leviathan, with selected variants from the Latin edition of 1668,* E. Curley, Ed. (Hackett, Indianapolis, 1994).

44.   S. Wolf, *Freedom Within Reason*. (Oxford University Press, New York, 1990).

45.   R. Deutschländer, M. Pauen, J.-D. Haynes, Probing folk-psychology: Do Libet-style experiments reflect folk intuitions about free action? *Consciousness and Cognition* **48**, 232-245 (2017).

46.   G. Strawson, The impossibility of moral responsibility. *Philosophical Studies* **75**, 5-24 (1994).

47.   J. M. Fischer, Recent Work on Moral Responsibility*. *Ethics* **110**, 93-139 (1999).

48.   W. Sinnott-Armstrong, *Moral psychology: Free will and moral responsibility*. (MIT Press, 2014).

49.   A. L. Roskies, How does the neuroscience of decision making bear on our understanding of moral responsibility and free will? *Current opinion in neurobiology* **22**, 1022-1026 (2012).

50.   U. Maoz, G. Yaffe, What does recent neuroscience tell us about criminal responsibility? *Journal of Law and the Biosciences* **3**, 120-139 (2016).

51.   A. Mele, *Free will and luck.* (Oxford University Press, 2006).

52.   A. G. Sanfey, J. K. Rilling, J. A. Aronson, L. E. Nystrom, J. D. Cohen, The neural basis of economic decision-making in the ultimatum game. *Science* **300**, 1755-1758 (2003).

53.   J. D. Wallis, E. K. Miller, Neuronal activity in primate dorsolateral and orbital prefrontal cortex during performance of a reward preference task. *European Journal of Neuroscience* **18**, 2069-2081 (2003).

54.   N. Ramnani, A. M. Owen, Anterior prefrontal cortex: insights into function from anatomy and neuroimaging. *Nature Reviews Neuroscience* **5**, 184-194 (2004).

55.   J. D. Wallis, Orbitofrontal cortex and its contribution to decision-making. *Annu. Rev. Neurosci.* **30**, 31-56 (2007).

56.   G. Bush, P. Luu, M. I. Posner, Cognitive and emotional influences in anterior cingulate cortex. *Trends in Cognitive Sciences* **4**, 215-222 (2000).

57.   C. S. Carter *et al.*, Anterior cingulate cortex, error detection, and the online monitoring of performance. *Science* **280**, 747-749 (1998).

58.   V. Krieghoff, F. Waszak, W. Prinz, M. Brass, Neural and behavioral correlates of intentional actions. *Neuropsychologia* **49**, 767-776 (2011).

59.   M. Brass, P. Haggard, The what, when, whether model of intentional action. *The Neuroscientist* **14**, 319-325 (2008).

60.   G. Hughes, S. Schütz-Bosbach, F. Waszak, One action system or two? Evidence for common central preparatory mechanisms in voluntary and stimulus-driven actions. *The Journal of Neuroscience* **31**, 16692-16699 (2011).

61.   U. Maoz *et al.*, Predeliberation activity in prefrontal cortex and striatum and the prediction of subsequent value judgment. *Front Neurosci* **7**, 225 (2013).

62.   A. Tusche, S. Bode, J. D. Haynes, Neural responses to unattended products predict later consumer choices. *J Neurosci* **30**, 8024-8031 (2010).

63.   D. Brainard, The psychophysics toolbox. *Spatial Vision* **10**, 433-436 (1997).

64.   D. G. Pelli, The VideoToolbox software for visual psychophysics: Transforming numbers into movies. *Spatial Vision* **10**, 437-442 (1997).

65.   H. Lau, R. Rogers, R. Passingham, Manipulating the experienced onset of intention after action execution. *Journal of cognitive neuroscience* **19**, 81-90 (2007).

66.   W. P. Banks, E. A. Isham, We infer rather than perceive the moment we decided to act. *Psychological Science* **20**, 17 (2009).

67.   J. Miller, P. Shepherdson, J. Trevena, Effects of Clock Monitoring on Electroencephalographic Activity. *Psychological Science* **22**, 103-109 (2011).

68.   U. Maoz *et al.*, in *Surrounding Free Will: Philosophy: Psychology, Neuroscience,* A. R. Mele, Ed. (Oxford University Press, 2015).

69.   T. P. Jung *et al.*, Removing electroencephalographic artifacts by blind source separation. *Psychophysiology* **37**, 163-178 (2000).

70.   A. S. Keren, S. Yuval-Greenberg, L. Y. Deouell, Saccadic spike potentials in gamma-band EEG: characterization, detection and suppression. *Neuroimage* **49**, 2248-2263 (2010).

71.   T. W. Picton *et al.*, Guidelines for using human event-related potentials to study cognition: Recording standards and publication criteria. *Psychophysiology* **37**, 127-152 (2000).

72.   R. Ratcliff, A theory of memory retrieval. *Psychological Review* **85**, 59-108 (1978).

73.   M. Usher, J. L. McClelland, The time course of perceptual choice: the leaky, competing accumulator model. *Psychological review* **108**, 550 (2001).

Acknowledgments: We thank Ralph Adolph for his invaluable guidance and support in designing and running the experiment as well as for very useful discussions of the results. We thank Ram Rivlin for various conceptual discussions about deliberate versus arbitrary decision-making and about the initial experimental paradigm design. We thank Caitlin Duncan for her help in patiently and meticulously gathering the EEG data. We thank Daw-An Wu for discussions about EEG data collection and preprocessing and for his help with actual data collection. We thank Daniel Grossman for his help in carefully preprocessing the data and suggesting potential interpretations of it. We thank Ueli Rutishauser for discussions about the model and its simulations. **Funding:** This research was supported by Florida State University's Big Questions in Free Will Initiative, funded by the John Templeton Foundation, to U.M. and C.K.; by the Ralph Schlaeger Charitable Foundation to U.M.; by the Bial Foundation to U.M. and to U.M. and L.M.; and by the German-Israeli Foundation for Scientific Research and Development to L.M.. C.K. thanks the Allen Institute founders, Paul G. Allen and Jody Allen, for their vision, encouragement, and support. **Author contributions:** U.M, L.M, G.Y., and C.K. conceived the project and designed the experiments. L.M. and U.M. analyzed the results. U.M. designed and simulated the model. L.M. and U.M. wrote the manuscript. G.Y. and C.K. suggested revisions to the manuscript. **Competing interests:** The authors declare that they have no competing interests. **Data and materials availability:** All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. Additional data related to this paper may be requested from the authors.