**Integration of visual information in auditory cortex promotes auditory scene analysis through**

**multisensory binding**

Huriye Atilgan[1], Stephen M. Town[1], Katherine C. Wood[1], Gareth P. Jones[1], Ross K. Maddox[2,3], Adrian

K.C. Lee[3] and Jennifer K. Bizley[1]

[1]The Ear Institute, University College London, UK

[2]Department of Biomedical Engineering, University of Rochester, Department of Neuroscience,

University of Rochester, Rochester, USA

[3]Institute for Learning and Brain Sciences and Department of Speech and Hearing Sciences,

University of Washington, Seattle, USA

7 Figures

59600 characters

Corresponding Author and Lead Contact: Jennifer Bizley j.bizley@ucl.ac.uk

UCL Ear Institute, 332 Gray's Inn Road, London, WC1X 8EE.

## Summary

How and where in the brain audio-visual signals are bound to create multimodal objects remains

unknown. One hypothesis is that temporal coherence between dynamic multisensory signals

provides a mechanism for binding stimulus features across sensory modalities. Here we report that

when the luminance of a visual stimulus is temporally coherent with the amplitude fluctuations of

one sound in a mixture, the representation of that sound is enhanced in auditory cortex. Critically,

this enhancement extends to include both binding and non-binding features of the sound. We

demonstrate that visual information conveyed from visual cortex, via the phase of the local field

potential, and is combined with auditory information within auditory cortex. These data provide

evidence that early cross-sensory binding provides a bottom-up mechanism for the formation of

cross-sensory objects and that one role for multisensory binding in auditory cortex is to support

auditory scene analysis.

When listening to a sound of interest, we frequently look at the source. However, how auditory and

visual information are integrated to form a coherent perceptual object is unknown. The temporal

properties of a visual stimulus can be exploited to detect correspondence between auditory and

visual streams(Crosse et al., 2015; Denison et al., 2013; Rahne et al., 2008), can bias the perceptual

organisation of a sound scene (Brosch et al., 2015), and can enhance or impair listening performance

depending on whether the visual stimulus is temporally coherent with a target or distractor sound

stream (Maddox et al., 2015). Together, these behavioural results suggest that temporal coherence

between auditory and visual stimuli can promote binding of cross-modal features to enable the

formation of an auditory-visual (AV) object (Bizley et al., 2016).

Visual stimuli can both drive and modulate neural activity in primary and non-primary auditory

cortex (Bizley et al., 2007a; Chandrasekaran et al., 2013; Ghazanfar et al., 2005; Kayser et al., 2008;

Perrodin et al., 2015) but the contribution that visual activity in auditory cortex makes to auditory

function remains unknown. One possibility is that the integration of cross-sensory information into

early sensory cortex provides a bottom-up substrate for the binding of multisensory stimulus

features into a single perceptual object (Bizley et al., 2016). We have recently argued that binding is

a distinct form of multisensory integration that underpins perceptual object formation. We

hypothesise that binding is associated with a modification of the sensory representation and can be

identified by demonstrating a benefit in the behavioural or neural discrimination of a stimulus

feature orthogonal to the features that link crossmodal stimuli (Fig. 1a). Therefore, in order to

demonstrate binding an appropriate crossmodal stimulus should elicit not only enhanced neural

encoding of the stimulus features that bind auditory and visual streams (the "binding features"), but

that there should be enhancement in the representation of *other* stimulus features ("non-binding

features" associated with the source (Fig. 1c).

Here we test the hypothesis that the incorporation of visual information into auditory cortex can

determine the neuronal representation of an auditory scene through multisensory binding (Fig.1).

We demonstrate that when visual luminance changes coherently with the amplitude changes of one

sound in a mixture, auditory cortex is biased towards representing the temporally coherent sound.

Consistent with these effects reflecting cross-modal binding, the encoding of auditory timbre, a non-

binding stimulus feature, is subsequently enhanced in the temporally coherent auditory stream.

Finally, we demonstrate that the site of multisensory convergence is in auditory cortex and that

visual information is conveyed via the local field potential directly from visual cortex.

**Figure 1: Hypothesis and experimental design**

**a** Conceptual model illustrating how binding can be identified as a distinct form of multisensory
integration. Multisensory binding is defined as a subset of multisensory integration that results in
the formation of a crossmodal object. During binding, all features of the audio-visual object are
linked and enhanced including both those features that bind the stimuli across modalities (here

temporal coherence between auditory (A) intensity and visual (V) luminance) and orthogonal features such as auditory pitch and timbre, and visual colour and size. Other forms of multisensory integration would result in only the features that promote binding being enhanced - here auditory intensity and visual luminance. To identify binding therefore requires a demonstration that non-binding features (e.g. here pitch, timbre, colour or size) are enhanced. **b** When two competing sounds (red and blue waveforms) are presented they can be separated on the basis of their features, but may elicit overlapping neuronal representations in auditory cortex. **c** Hypothesised enhancement in auditory stream segregation when a temporally coherent visual stimulus enables multisensory binding. When the visual stimulus changes coherently with the red sound (A1, top) this sound is enhanced and the two sources are better segregated. Perceptually this would result in more effective auditory scene analysis and an enhancement of the non-binding features. **d** Stimulus design: Auditory stimuli were two artificial vowels (denoted A1 and A2), each with distinct pitch and timbre and independently amplitude modulated with a noisy low pass envelope. **e** Visual stimulus: a luminance modulated white light was presented with one of two temporal envelopes derived from the amplitude modulations of A1 and A2. **f** illustrates the stimulus combinations that were tested experimentally in *single stream* (a single auditory visual pair) and *dual stream* (two sounds and one visual stimulus) conditions.

# Results

We recorded neuronal responses in the auditory cortex of awake passively listening ferrets (n=9

ferrets, 221 single units, 311 multi-units) in response to naturalistic time-varying auditory and visual

stimuli adapted from Maddox et al (2015). The stimuli are designed to share properties with natural

speech; they are modulated at approximately syllable rate and, like competing voices, can be

separated on the basis of their fundamental frequency (F0, the physical determinant of pitch). These

sounds are devoid of any linguistic content rendering them suitable for animal models and

permitting the separation of general sensory processing mechanisms from language-specific ones.

We additionally recorded neural responses in medetomidine-ketamine anesthetised ferrets (n=5

ferrets, 426 single units, 772 multi units) which allowed us to eliminate attentional effects and thus

isolate bottom-up processing. These experiments also permitted longer recording durations for

additional control stimuli and enabled simultaneous characterization of neural activity across cortical

laminae. In a subset of these animals we were able to reversibly silence visual cortex during

recording in order to determine the origin of visual-stimulus elicited neural changes.  Recordings

were made in awake animals while they held their head at a drinking spout but were not engaged in

a behavioral task. and allowed us to measure neural activity free from any confounds associated

with pharmacological manipulation and in the absence of task-directed attention.

The stimuli were two auditory streams comprised of two vowels, each with a distinct pitch and

timbre (denoted A1: [u], fundamental frequency (F0) = 175 Hz and A2: [a], F0 = 195 Hz, Fig. 1) and

independently amplitude modulated with a low-pass (<7 Hz) envelope (Fig. 1d). A full-field visual

stimulus accompanied the auditory stimuli, the luminance of which was temporally modulated with

the modulation envelope from one of the two auditory streams (Fig. 1e). We tested stimulus

conditions in which both auditory streams were presented ("dual stream") and the visual stimulus

was temporally coherent with one or other of the auditory streams (A12V1 or A12V2, Fig. 1e). We

also tested conditions in which a single AV stimulus pair was presented ('single stream' stimuli),

where the auditory and visual streams could be temporally coherent (A1V1, A2V2) or independently

modulated (A1V2, A2V1).

**Auditory-visual temporal coherence shapes the representation of a sound scene in**

**auditory cortex**

We first asked whether the temporal dynamics of a visual stimulus could enhance the

representation of one sound in a mixture.  We therefore recorded responses to auditory scenes

composed of two sounds (A1 and A2), presented simultaneously, with a visual stimulus that was

temporally coherent with one or other auditory stream (A12V1 or A12V2). A visual stimulus is known

to enhance the representation of the envelope of an attended speech stream in auditory cortex

(Zion Golumbic et al., 2013; Park et al., 2016). To test whether we could observe a similar

phenomena in single neurons in the absence of selective attention we used neural responses to

temporally coherent single stream stimuli (i.e. A1V1 and A2V2) to determine to what extent the

neural response to the sound mixture was specific to one or other sound stream.

**Figure 2: Visual stimuli can determine which sound stream auditory cortical neurons follow in a mixture.**

Spiking responses from an example unit in response to **a,** single stream AV stimuli used as decoding templates and **b**, dual stream stimuli. In each case rasters and PSTHs are illustrated. When the visual component of the dual stream was V1, the majority of trials were classified as A1V1 (82% (19/23) of trials), and A2V2 when the visual stimulus was V2 (26% (6/23) of responses classified as A1V1 (see also green data point in c). **c-f** population data for awake (**c,d,** 541 units) and anesthetised (**e,f** 661 units) datasets. In each case the left panel (c,e) shows the distribution of decoding values according to the visual condition and the right panel (**d,f**) shows the population mean (± SEM) Pairwise comparisons revealed significant effect of visual condition on decoding in both datasets (p<0.001).

Figure 2 illustrates this approach for a single unit: responses to the single stream AV stimuli (Fig. 2a)

formed templates which were used to decode the responses to the dual stream stimuli (Fig. 2b)

using a Euclidean distance based spike pattern classifier (additional example: Fig S1).  Responses to

the dual stream stimuli were more commonly decoded as A1V1 when the visual stimulus was V1,

and A2V2 when the visual stimulus was V2. For any unit in which we were able to decode the single

stream template responses at a level better than chance (p<0.05, permutation test; 541 units, awake

dataset; 661 units anesthetised dataset) we proceeded to use these single stream responses to

decode dual stream responses. Enhancement of the coherent auditory stimulus representation was

visible at the population level (Fig. 2c-f): Auditory cortical responses to dual-stream stimuli most

closely resembled responses to the single stream stimulus with the same visual component. This

finding was robust in both awake (Fig. 2d, pairwise t-test: $t_{540}$ = 6.073, p<0.001) and anesthetised

animals (Fig. 2f, $t_{660}$ = 9.5137, p<0.001) suggesting that these effects were not mediated by attention.

Modulation of dual stream responses by visual stimulus identity was not simply a consequence of

the shared visual component of single stream and dual stream stimuli and was observed in neurons

in which visual or auditory identity could be decoded (Fig S1). If this effect was only apparent in

visual neurons in auditory cortex then eliminating the visual element of the single stream stimuli

should impair decoding performance for the dual stream stimuli. Additional control experiments

(n=89 units, awake animals) demonstrated that this was not the case: the enhancement of the

temporally coherent sound in the sound mixture was evident whether dual stream stimuli (A12V1

and A12V2) were decoded using responses to auditory-only stimuli (A1 or A2) or auditory-visual

stimuli (Fig. 3a). We also determined that removing the visual stimulus from the dual-stream

eliminated any decoding difference in responses observed. A two-way repeated measures ANOVA

on decoded responses with factors of visual stream (V1, V2, no visual), and template type (AV or A)

demonstrated a significant effect of visual stream identity on dual stream decoding (Fig. 3d, $F_{(2, 528)}$ = 19.320, p <0.001), but there was no effect of template type ( $F_{(1, 528)}$ = 0.073, p = 0.787) or

interaction between factors ($F_{(2, 528)}$ = 0.599, p = 0.550). Post-hoc comparisons revealed that

without visual stimulation there was no tendency to respond preferentially to either stream but that

visual stream identity significantly influenced the classification of dual stream responses.

168
**Figure 3: Visual stimuli shape the neural representation of an auditory scene.**

In an additional control experiment (n=89 units recorded in awake animals), the responses to
coherent AV and auditory-only (A Only) single stream stimuli were used as templates to decode dual
stream stimuli either accompanied by visual stimuli (V1/V2) or in the absence of visual stimulation
(no visual). Spiking responses from an example unit in response to **a,** single stream auditory stimuli
which were used as decoding templates to decode the responses to dual stream stimuli in **b**, in each
case the auditory waveform, rasters and PSTHs are shown. **c**, population data showing the
proportion of responses classified as A1 when the visual stimulus was V1 or V2 when decoded with
auditory-only templates. **d,** Mean (± SEM) values for these units when decoded with A-only
templates (i.e. data from **c**), AV templates (as in Fig.2) or in the absence of a visual stimulus. The
green data point in **c** depicts the example in **a, b**. Pairwise comparisons revealed significant effect
across visual conditions in both datasets (* indicates p<0.001).

The ability of auditory-visual temporal coherence to enhance one sound in a mixture was observed

across all cortical layers (defined by current source density analysis, see methods, Fig.S2, a,b) and

across three tonotopic fields (Fig.S2 c,d) of anaesthetized subjects. While present in all layers, the

influence of the visual stimulus was strongest in the supra-granular layers (Fig.S2). Separating units

into auditory, visual, and auditory-visual units according to responsiveness to classic

neurophysiological stimuli (noise bursts and light flashes) revealed that the impact of a visual

stimulus on the representation of a sound mixture was present across functional sub-populations

but strongest in visual and auditory-visual units (Fig. S3, anesthetised dataset). Finally, we observed

these effects in both single and multi-units (Fig.S4).

## Auditory-visual temporal coherence enhances non-binding sound features

A hall-mark of an object-based rather than feature-based representation is that all stimulus features

are bound into a unitary perceptual construct, including those features which do not directly

mediate binding (Desimone and Duncan, 1995). We predicted that binding across modalities would

be promoted via synchronous changes in auditory intensity and visual luminance (Fig. 1b) and

observed that the temporal dynamics of the visual stimulus enhanced the representation of

temporally coherent auditory streams (Fig. 2e-f and 3d-f). To determine whether temporal

synchrony of visual and auditory stimulus components also enhanced the representation of

orthogonal stimulus features (Fig. 1c) and thus fulfil a key prediction of binding, we introduced brief

timbre perturbations into our acoustic stimuli (two in each of the A1 and A2 streams). Each deviant

lasted for 200 ms during which the spectral timbre smoothly transitioned to the identity of another

vowel and back to the original vowel. Such deviants could be detected by human listeners and were

better detected when the auditory stream in which they were embedded was temporally coherent

with an accompanying visual stimulus (Maddox et al., 2015). We hypothesised that, despite

containing no information about the timing or occurrence of deviants, a temporally coherent visual

stimulus would enhance the representation of timbre deviants in the responses of auditory cortical

neurons.

**Figure 4: Temporally coherent changes in visual luminance and auditory intensity enhance the representation of auditory timbre**

**a** Example unit (from the awake dataset) showing the influence of visual temporal coherence on spiking responses to dual stream stimuli with (red PSTH) or without (black PSTH) timbre deviants. Shaded rectangles indicate the 200 ms window over which the timbre deviant occurred and over which analysis was conducted. **b-d** timbre deviant discrimination in the awake dataset. Two deviants were included in each auditory stream giving a possible maximum of 4per unit **b,** Histogram showing the number of deviants (out of 4) that could be discriminated from spiking responses **c,** Box plots showing the average timbre deviant discrimination scores in the single stream condition across different visual conditions (Coh: coherent, ind: independent). The boxes show the upper and lower quartile values, and the horizontal lines indicates the median. **d,** Discrimination scores for timbre deviant detection in dual stream stimuli. Discrimination scores are plotted according to the auditory

stream in which the deviant occurred and the visual stream that accompanied the sound mixture. **e-**
**g** show the same as **b-d** but for the anesthetised dataset.

To isolate neural responses to the timbre change from those elicited by the on-going amplitude

modulation, we extracted the 200ms epochs of the neuronal response during which the timbre

deviant occurred and compared these to epochs from responses to otherwise identical stimuli

without deviants. We observed that the spiking activity of many units differed between deviant and

no-deviant trials (e.g. Fig. 4a) and we were able to discriminate deviant from no-deviant trials with a

spike pattern classifier. We first considered the influence of temporal coherence between auditory

and visual stimuli on the representation of timbre deviants in the single stream condition (A1V1,

A1V2 etc.). We found that a greater proportion of units detected at least one deviant when the

auditory stream in which deviants occurred was temporally coherent with the visual stimulus

relative to the temporally independent condition. This was true both for awake (Fig. 4b; Pearson chi-

square statistic, $\chi^2 = 322.617$, $p < 0.001$) and anesthetised animals (Fig. 4e; $\chi^2 = 288.731$, $p < 0.001$).

For units that discriminated at least one deviant, discrimination scores were significantly higher

when accompanied by a temporally coherent visual stimulus (Fig.4c, awake dataset, pairwise t-test

$t_{300} = 3.599$ $p<0.001$; Fig. 4f, anesthetised data $t_{262} = 4.444$ $p<0.001$).

We also observed an enhancement in the representation of timbre changes in the context of a

sound scene (Fig 4d,g): timbre changes were more reliably encoded when the sound stream in which

they were embedded was accompanied by a temporally coherent visual stimuli. We performed a

two-way repeated measures ANOVA on deviant discrimination performance with visual condition

(V1/V2) and the auditory stream in which the deviants occurred (A1/A2) as factors. We predicted

that enhancement of the representation of timbre deviants in the temporally coherent auditory

stream would be revealed as a significant interaction term in the dual stream data. Significant

interactions were seen in both the awake (Fig. 4d, $F(1, 600) = 29.138$, $p<0.001$) and anesthetised

datasets (Fig. 4g, $F(1, 524) = 16.652$, $p<0.001$). We also observed significant main effects of auditory

and visual conditions in awake (main effect of auditory stream, $F_{(1, 600)} = 4.565$, $p = 0.033$; main

effect of visual condition, $F_{(1, 600)} = 2.650$, $p = 0.010$) but not anesthetised animals (main effect of

auditory stream, $F_{(1,524)} = 0.004$, $p = 0.948$; main effect of visual condition, $F_{(1, 524)} = 1.355$, $p =$

0.245). Thus these data demonstrate the hypothesised enhancement in the neural representation of

both binding (i.e. auditory amplitude) and non-binding features (here auditory timbre) that are

orthogonal to those that promote binding between auditory and visual streams and so the effects

we observe in auditory cortex fulfil our definition of multisensory binding. Next we turn to the

question of how these effects are mediated and whether they emerge within or outside of auditory

cortex.

**Auditory cortical spike patterns differentiate dynamic auditory-visual stimuli more**

**effectively when stimuli are temporally coherent**

We used the responses to single stream stimuli to classify neurons according to whether they were

modulated primarily by auditory or visual temporal dynamics. To determine whether the auditory

amplitude envelope reliably modulated spiking we used a spike-pattern classifier to decode the

auditory stream identity, collapsed across visual stimuli (i.e. we decoded auditory stream identity

from the combined responses to A1V1 and A1V2 stimuli and the combination of A2V1 and A2V2

responses). An identical approach was taken to determine if neuronal responses reliably

distinguished visual modulation (i.e. we decoded visual identity from the combined responses to

A1V1 and A2V1 stimuli and the combined responses elicited by A1V2 and A2V2). Neuronal

responses which were informative about auditory or visual stimulus identity at a level better than

chance (estimated with a bootstrap resampling) were classified as auditory-discriminating (Fig. 5a-b)

and / or visual-discriminating (Fig. 5c-d) respectively.

**Figure 5: Auditory-visual temporal coherence enhances neural coding in auditory cortex**.

A pattern classifier was used to determine whether neuronal responses were informative about auditory or visual stimuli. The responses to single stream stimuli are shown for two example units, with responses grouped according to the identity of the auditory (**a, b**, for an auditory discriminating unit) or visual stream (**c, d**, for a visual discriminating unit). In each case the stimulus amplitude (a,b) / luminance (c,d) waveform is shown in the top panel with the resulting raster plots and PSTHs below. **e, f**: Decoder performance (mean ± SEM) for discriminating stimulus identity (coherent: A1V1 vs. A2V2, purple; independent: A1V2 vs. A2V1, blue) in auditory and visual classified units recorded in awake (e) and anaesthetised (f) ferrets. Pairwise comparisons for decoding of coherent versus independent stimuli (*** indicates p<0.001 ).

In awake animals, 39.5% (210/532) of driven units were auditory-discriminating, 11.1% (59/532)

were visual-discriminating, and only 0.4% (2/532) discriminated both auditory and visual stimuli.

Overall a smaller proportion of units represented the identity of auditory or visual streams in the

anesthetised dataset: 20.2% (242/1198) were auditory-discriminating, 6.8% (82/1198) were visual

discriminating, and 0.6% (7/1198) discriminated both. During recordings made under anaesthesia,

we also recorded responses to noise bursts and light flashes (both 100 ms duration) presented

separately and together to compare how the proportion of auditory / visual discriminating units

measured to naturalistic dynamic stimuli compared to more traditional artificial stimuli. Specifically,

responsiveness was defined using a two-way ANOVA (factors: auditory stimulus [on/off] and visual

stimulus [on/off]) on spike counts measured during stimulus presentation. We defined units as being

sound-driven (main effect of auditory stimulus, no effect of visual stimulus or interaction), light-

driven (main effect of visual stimulus, no effect of auditory stimulus or interaction) or both (main

effect of both auditory and visual stimuli or significant interaction; $p < 0.05$) as in (Bizley et al.,

2007b). Using such stimuli revealed that the classification of units by visual / auditory discrimination

of single stream stimuli selected a subset of light or sound driven units and that the proportions of

auditory, visual and AV units recorded in our sample were in line with previous studies from ferret

auditory cortex (65.1% (328/504) of units were driven by noise bursts, 16.1% (81/504) by light

flashes and 14.1% (71/504) by both. For units which were classified as auditory or visual

discriminating, and for which we recorded responses to noise bursts and light flashes, 53% (160/307)

were classified as auditory, 17% (53/307) as visual and 31% (94/307) as auditory-visual when

classified with simple stimuli.

We hypothesised that the effects we observed in the dual-stream condition might be a consequence

of temporal coherence between auditory and visual stimuli enhancing the discriminability of neural

responses. We confirmed this prediction by using the same spike pattern decoder to compare our

ability to discriminate temporally coherent (A1V1 vs. A2V2) and temporally independent (A1V2 vs.

A2V1) stimuli (Fig. 5e, f): Temporally coherent AV stimuli produced more discriminable spike

patterns than those elicited by temporally independent ones in both awake (Fig. 5e, pairwise t-test,

auditory-discriminating $t_{418}$ = 11.872, p<0.001; visual-discriminating $t_{116}$ = 6.338, p<0.001; All $t_{540}$ =

13.610, p<0.001) and anesthetised recordings (Fig.5f, auditory-discriminating $t_{482}$ = 17.754, p<0.001;

visual-discriminating $t_{162}$ = 8.186, p<0.001; All $t_{664}$ = 19.461, p<0.001).

## Dynamic visual stimuli elicit reliable changes in LFP phase

Temporal coherence between auditory and visual stimulus streams results in more discriminable

spike trains in the single stream condition, and an enhancement of the representation of the

temporally coherent sound when that sound forms part of an auditory scene. What might underlie

the enhanced discriminability observed for temporally coherent cross-modal stimuli? The phase of

on-going oscillations determines the excitability of the surrounding cortical tissue (Azouz and Gray,

1999; Okun et al., 2010; Szymanski et al., 2011). LFP phase is reliably modulated by naturalistic

stimulation (Chandrasekaran et al., 2010; Kayser et al., 2009; Luo and Poeppel, 2007b; Ng et al.,

2012; Schyns et al., 2011) and has been implicated in multisensory processing (Golumbic et al., 2013;

Lakatos et al., 2007). We hypothesised that sub-threshold visual inputs could modulate spiking

activity by modifying the phase of the local field potential such that when auditory and visual stimuli

were temporally coherent visual stimuli 'reset' the LFP phase to maximise the likelihood that feed

forward sound-evoked activity elicited robust spiking responses. This would provide a mechanism by

which neuronal spiking was enhanced when auditory and visual streams are temporally coherent.

Stimulus evoked changes in the local field potential (LFP) were evident from the recorded voltage

traces and analysis of inter-trial phase coherence demonstrated that there were reliable changes in

phase across repetitions of identical AV stimuli (Fig. 6 a, b). To isolate the influence of visual activity

on the LFP at each recording site, and address the hypothesis that visual stimuli elicited reliable

changes in the LFP, we calculated phase and power dissimilarity functions for stimuli with identical

auditory signals but differing visual stimuli (Luo and Poeppel, 2007b). Briefly, this analysis assumes

that if the phase (or power) within a particular frequency band differs systematically between

responses to two different stimuli, then inter-trial phase coherence (ITPC) across repetitions of the

same stimulus will be greater than across randomly selected stimuli. For each frequency band in the

LFP, we therefore compared "within-stimulus" ITPC for responses to each stimulus (A1 stream Fig.

6c; A2 stream Fig. 3d) with "across-stimulus" ITPC calculated from stimuli with *identical auditory*

*components* but randomly selected visual stimuli (e.g. randomly drawn from A1V1 and A1V2; Fig.

6c). The difference between within-stimulus and across-stimulus ITPC was then calculated across

frequency and described as the phase dissimilarity index (PDI) (Fig. 6e, f) with positive PDI values

indicating reliable changes in phase coherence elicited by the visual component of the stimulus.

Importantly, because both test distributions and the null distribution contain identical sounds any

significant PDI value can be attributed directly to the visual component of the stimulus.

**Figure 6: Visual stimuli elicit reliable changes in the phase of the local field potential**

**a, b** Example LFP responses to single stream stimuli across visual conditions. Data obtained from the recording site at which multiunit spiking activity discriminated auditory stream identity in Fig. 5 a and b. The amplitude waveforms of the stimuli are shown in the top panel, with the evoked LFP underneath (mean across 21 trials). The resulting inter-trial phase coherence (ITPC) values are shown in the bottom two panels. **c, d** ITPC was calculated for coherent and independent AV stimuli separately and compared to a null distribution (ITPC across). Single stream phase dissimilarity values (PDI) were calculated by comparing ITPC within values to the ITPC across null distribution (**e, f**). **g,l** Population mean inter-trial phase coherence (ITPC) values across frequency for coherent (**g**) and independent (**i**) conditions. Dots indicate frequencies at which the ITPC-within values were

significantly greater than the ITPC-across values (Pairwise t-test, $\alpha$ = 0.0012, Bonferroni corrected for
43 frequencies). **k:** Mean (±SEM) single stream phase dissimilarity index (PDI) values for coherent
and independent stimuli in animals. Black dots indicate frequencies at which the temporally
coherent single stream PDI is significantly greater than in the independent conditions (p<0.001). **h,j,l**
as **g,l,k** for anesthetised dataset. **m, n,** mean ±SEM dual stream PDI values for awake (m) and
anesthetised (n) datasets.

We calculated PDI values for each of the four single stream stimuli and grouped conditions by

coherency (coherent: A1V1 / A2V2, or independent: A1V2 / A2V1). To determine at what

frequencies the across-trial phase reliability was significantly positive, we compared the within-

stimulus values with the across-stimulus (null distribution) values for each frequency band (paired t-

test with Bonferroni correction for 43 frequencies, $\alpha$ = 0.0012). In awake subjects we identified a

restricted range of frequencies between 10 and 20 Hz where visual stimuli enhanced the phase

reliability (Fig. 3g, h). In anesthetised animals, average PDI values were larger than in awake animals

and all frequencies tested had single stream PDI values that were significantly non-zero (Fig. 3j,k).

We therefore conclude that visual stimulation elicited reliable changes in the LFP phase in auditory

cortex. In contrast to LFP phase, a parallel analysis of across trial power reliability showed no

significant effect of visual stimuli on LFP power in any frequency band (Fig.S5).

If visual information was only conveyed in the case of temporally coherent stimuli this might indicate

that the locus of binding was outside of auditory cortex. While there were significant PDI values for

both temporally independent and coherent stimuli we next asked whether there were any

frequencies at which phase coherence was significantly greater in temporally coherent AV stimuli

compared to temporally independent AV stimuli. To address this we performed a pairwise

comparison of single stream PDI values yielded from temporally coherent and independent stimuli,

for all frequency points. In awake animals, PDI values were similar for temporally coherent and

temporally independent stimuli, except in the 11-14 Hz band where coherent stimuli elicited

significantly greater phase coherence (Fig. 6i). In anaesthetised animals, the single stream PDI did

not differ between coherent and independent stimuli at any frequency (Fig. 6l). Together these data

suggests that visual inputs modulate the phase of the low frequency field potential in auditory

cortex largely independently of any temporal coherence between auditory and visual stimuli. This

finding supports the conjecture that multisensory binding occurs within auditory cortex.

To understand whether similar mechanisms could underlie the visual-stimulus induced

enhancement of a temporally coherent sound in a mixture we performed similar analysis on the

data collected in response to the dual stream stimuli. We generated within-stimulus ITPC values for

each dual-stream stimulus (i.e. A12V1 and A12V2, Fig. 6a) and across-stimulus ITPC by randomly

selecting responses across visual conditions (Fig. 6b). We then expressed the difference as the dual

stream phase dissimilarity index (dual stream PDI, Fig. 6c). Since the auditory components were

identical in each dual stream stimulus, the influence of the visual component on LFP phase could be

isolated as non-zero dual stream PDI values (paired t-test, bonferoni corrected, $\alpha = 0.0012$). In

awake animals, the dual stream PDI was significantly non- zero at 11-14Hz and 16-19 Hz (Fig. 6d, e)

whereas in anesthetised animals, we found positive dual stream PDI values across all frequencies

tested (Fig. 6f, g). In anesthetised animals where we were able to use the responses of units to noise

and light flashes to categorise units as auditory, visual or auditory-visual, and were able to confirm

significant PDI values in the LFP recorded on the same electrode as units in each of these

subpopulations (Fig. S3c). In awake animals, we tested auditory visual stimuli presented at three

different modulation rates (7, 12 and 17Hz) and confirmed that significant PDI values were obtained

at very similar frequencies across these modulation rates consistent with these being evoked phase

resets rather than stimulus-induced oscillations (Fig. S6).

**Visual cortex mediates visual-stimulus induced LFP changes in auditory cortex**

Visual inputs to auditory cortex potentially originate from many sources: In the ferret visual multiple

visual cortical fields are known to innervate auditory cortex (Bizley et al., 2007b), but frontal and

thalamic areas are additional candidates for sources of top-down and bottom-up multisensory

innervation. Therefore, in order to determine the origin of the visual effects that we observe in

auditory cortex, we performed an additional experiment in which we cooled the gyral surface of the

posterior suprasylvian sulcus where visual cortical fields SSY (Cantone et al., 2006) and Area 21

(Innocenti et al., 2002) are located (Fig. 7a). Neural tracer studies have demonstrated that these

areas directly project to auditory cortex in the ferret (Bizley et al., 2007b). We used a cooling loop

cooled to 9-10°C to reversible silence neural activity within <500 um of the loop (see Fig. 3 (Wood et

al., 2017). Using simple noise bursts and light flashes we verified that cooling visual cortex did not

alter the response to noise bursts in auditory cortex (repeated measures ANOVA $F_{(2,164)}$ = 0.42

p=0.88) but did reversibly attenuate the spiking response to light flashes in visual cortical sites

>500um from the cooling loop (repeated measures ANOVA $F_{(2,92)}$ = 6.83 p=0.001, post-hoc

comparisons shows pre-cool and cool-post were significantly different, pre-post were not

significantly different indicating the effects were reversible). We measured responses to the single

stream stimuli in auditory and visual cortex before and during cooling, while the cooling loop was

held at 9-10°C. From the LFP, we calculated the across-trial-phase coherence and phase dissimilarity

indexes (as in Fig. 6). Cooling significantly decreased the magnitude of the single stream PDI values in

auditory cortex (Fig 7e,h). A 3-way repeated measures ANOVA with factors of visual condition

(coherent/independent), frequency, and cortical temperature (warm/cooled) on the SS PDI values

obtained in auditory cortex showed a main effect of frequency ($F_{(88,22605)}$ = 47.91, $p<0.1*10^9$)

and temperature ($F_{(1,22605)}$ = 1072, $p<0.1*10^9$,) but not visual condition (p=0.49). In contrast, those

in visual cortex (away from the loop) were unaffected by cooling (3-way ANOVA demonstrated that

the magnitude of the SS PDI value was influenced by frequency ($F_{(88,17265)}$ = 24.73, $p<0.1*10^9$), but

not temperature (p=0.75) or visual condition (p=0.29)). From these data we conclude that the

influence of visual stimuli on the auditory cortical field potential phase is mediated, at least in part,

by inputs from visual cortical areas SSY and 21. While cooling does not allow us to confirm that these

are direct mono-synaptic inputs the observation that the phase effects in other areas of visual cortex

are uninfluenced suggests that we are influencing these areas selectively rather than supressing

visual processing generally.

436
**Figure 7: Visual stimulus induced LFP phase changes in auditory cortex are mediated in visual**

**cortex**

**a** Schematic showing the location of auditory cortical recording sites and the location of a cooling
loop (black, grey line marks the 500 um radius over which cooling is effective (Wood et al., 2017)
which was used to inactivate visual cortex. **b** Spike rate responses in Auditory Cortex (top row) and
visual cortex (bottom row, sites >500 um from the loop) in response to noise bursts or light flashes
before, during and after cooling. **c,d**, inter-trial phase coherence values for the coherent (**c**) and
independent stimuli (**d**) AV stimuli recorded in auditory cortex (AC) prior to cooling visual cortex (VC)
compared to the shuffled null distribution (ITPC across). Asterisks indicate the frequencies at which
the ITPC values are significantly different from the shuffled ITPC-across distribution **e** single stream
Phase Dissimilarity Index values calculated from the ITPC values in **c** and **d**. **f-h** - as c-e but while
visual cortex was cooled to 9 degrees. **i-n** as c-h but for sites in visual cortex >500um from the
cooling loop. c-h includes data from 83 sites from 6 electrode penetrations, i-n includes data from 47
sites from 5 penetrations.

# Discussion

Here we provide insight into how and where auditory-visual binding occurs, and provide evidence

that this effect is mediated by cortical-cortical interactions between visual and auditory cortex.

Visual stimuli elicit reliable changes in the phase of the local field potential in auditory cortex,

irrespective of auditory-visual temporal coherence, indicating that the inputs to auditory cortex

reflect the unisensory stimulus properties. When the visual and auditory stimuli are temporally

phase resets elicited by the visual stimulus via the LFP interact with the feed-forward sound evoked

activity and result in an enhanced spiking output from auditory cortex. These results are consistent

with the binding of cross-modal information to form a multisensory object because they result in a

modification of the representation of the sound in auditory cortex which is not restricted to the

features that link auditory and visual signals but extends to other non-binding features. These data

provide a physiological underpinning for the pattern of performance observed in human listeners

performing an auditory selective attention task, in which the detection of a pertubation in a stimulus

stream is enhanced or impaired when a visual stimulus is temporally coherent with the target or

masker auditory stream respectively (Maddox et al., 2015). The effects of the visual stimulus on the

representation of an auditory scene can be observed in anesthetised animals ruling out any top-

down effect of attentional modulation.

Previous investigations of the impact of visual stimuli on auditory scene analysis have frequently

used speech stimuli. Being able to see a talker's mouth provides listeners with rhythm information

and information about the amplitude of the speech waveform which may help listeners by cueing

them to pay attention to the auditory envelope (Peelle and Sommers, 2015) as well as provide cues

to the place of articulation that can disambiguate different consonants(Sumby and Pollack, 1954).

However, the use of speech stimuli makes it difficult to dissociate general multisensory mechanisms

from speech-specific ones. Therefore, in order to probe more general principles we chose to employ

continuous naturalistic non-speech stimuli, that utilized modulation rates that fell within the range

of syllable rates in human speech but lacked any linguistic content(Chandrasekaran et al., 2009).

Previous work has demonstrated that a visual stimulus can enhance the neural representation of the

speech amplitude envelope both in quiet and in noise(Crosse et al., 2015; Crosse et al., 2016; Luo et

al., 2010)'(Park et al., 2016), but functional imaging methods make if difficult to demonstrate

enhanced neural encoding of features beyond the amplitude envelope. The implication of our

findings is that the representation of the spectro-temporal features that allow speech recognition

would be enhanced in auditory cortex when a listener views a talker's face, including features such

as pitch that help pull apart voices in an auditory stream.

Visual speech information is hypothesised to be relayed to auditory cortex through multiple routes

in parallel to influence the processing of auditory speech: Our data support the idea that early

integration of visual information occurs (Möttönen et al., 2004; Okada et al., 2013; Peelle and

Sommers, 2015; Schroeder et al., 2008) and is likely to reflect a general phenomenon whereby visual

stimuli can cause phase-entrainment in the local field potential. Within this framework, cross-modal

binding results from an interaction between the bottom-up auditory spiking information and the

visual stimulus induced phase changes of the LFP. Indeed low-frequency entrainment to modulations

in an on-going stimulus are observed in the human brain including to lip movements(Park et al.,

2016) and have been shown to optimize listening performance for non-speech stimuli (Henry and

Obleser, 2012). In contrast, later integration is likely to underlie information about speech gestures

that might be used to constrain lexical identity (Peelle and Sommers, 2015).

Consistent with previous studies, our analysis of local field potential activity revealed that visual

information reliably modulated the phase of oscillatory activity in auditory cortex (Chandrasekaran

et al., 2013; Ghazanfar et al., 2005; Kayser et al., 2008; Perrodin et al., 2015). This occurred

independently of the modulation frequency of the stimulus suggesting that rather than entraining

oscillations at the stimulus modulation frequency particular features within the stream (presumably

points at which the luminance changed rapidly from low-high amplitude) triggered relatively broad-

band phase resets. Neuronal excitability varies with LFP phase (Jacobs et al., 2007; Klimesch et al.,

2007; Lakatos et al., 2013; Lőrincz et al., 2009) and such phase resets may be the physiological

mechanism through which cross-sensory information is integrated. Our analysis allowed us to isolate

changes in LFP phase that were directly attributable to the visual stimulus and identified reliable

changes in the LFP phase irrespective of whether the visual stimulus was temporally coherent with

the auditory stimulus. Such a finding is consistent with the idea that the LFP phase synchronization

arises from fluctuating inputs to cortical networks(Lakatos et al., 2007; Mazzoni et al., 2008;

Szymanski et al., 2011), and the loss of these effects with cooling suggests that visual cortex is a key

component of this network. Our finding that visual stimulation elicited reliable phase modulation in

both awake and anesthetised animals suggests that bottom-up cross-modal integration interacts

with selective attention, which also modulates phase information in auditory cortex(Golumbic et al.,

2013),(Park et al., 2016). While our data suggest that cross-modal binding can occur in the absence

of attention, it is likely that the effects we observe in auditory cortex are the substrates on which

selective attention acts to further boost the representation of cross-modal objects.

In the awake animal the impact of visual stimulation on LFP phase reliability was smaller than in the

anesthetised animal and was restricted to a narrower range of frequencies, consistent with a

dependence of oscillatory activity on behavioural state (Tukker et al., 2007; Voloh and Womelsdorf,

2016; Wang, 2010). Since the neural correlates of multisensory binding are evident in the

anesthetised animal, the specific increase in alpha phase reliability that occurred in awake animals in

response to temporally coherent auditory-visual stimulus pairs (Fig. 3c, 6e) may indicate an

attention-related signal triggered by temporal coherence between auditory and visual signals, or an

additional top-down signal conveying cross-modal information. Phase resetting or synchronisation of

alpha phase has been associated both with enhanced functional connectivity (Voloh and

Womelsdorf, 2016) and as a top-down predictive signal for upcoming visual information (Samaha et

al., 2015). Disambiguating these possibilities would require simultaneous recordings in auditory and

visual cortex and/or recording during the performance of a task designed to explicitly manipulate

attention.

Temporal coherence between sound elements has been proposed as a fundamental organising

principle for auditory cortex (Elhilali et al., 2009; O'Sullivan et al., 2015b) and here we extend this

principle to the formation of cross-modal constructs. Our data provide evidence that one role for the

early integration of visual information into auditory cortex is to resolve competition between

multiple sound sources within an auditory scene and that these neural computations occur pre-

attentively. While some proponents of a temporal coherence based model for auditory streaming

have stressed the importance of attention in auditory stream formation (Lu et al., 2017), neural

signatures of temporal-coherence based streaming are present in passively listening subjects

(O'Sullivan et al., 2015a; Teki et al., 2016). Previous studies have demonstrated a role for visual

information in conveying lip movement information to auditory cortex (Chandrasekaran et al., 2013;

Crosse et al., 2015; Ghazanfar et al., 2005; Golumbic et al., 2013), but such stimuli make it difficult to

separate sensory information from linguistic cues. Our data generated using non-speech stimuli

provide evidence that at least part of the boost provided by visualising a speaker's mouth arises

from a more general (language-independent) phenomenon whereby visual temporal cues facilitate

auditory scene analysis through the formation of cross-sensory objects. Our data are supportive of

visual cortical providing at least one source of information. Other visual cortical fields and sub-

cortical structures innervate tonotopic auditory cortex (Bizley et al., 2007a; Budinger et al., 2006)

and may potentially provide additional visual inputs to auditory cortex. Further dissecting the origin

of visual innervation requires experiments that allow pathway specific manipulation of neuronal

activity (for example by silencing the terminal fields of neurons that project from a candidate area

into auditory cortex)..

In summary, activity in auditory cortex was reliably affected by visual stimulation in a manner that

enhanced the representation of temporally coherent auditory information. Enhancement of auditory

information was observed for sounds presented alone or in a mixture and for sound features that

were related to (amplitude) and orthogonal to (timbre) variation in visual input. Such processes

provide mechanistic support for a coherence-based model of cross-modal binding in object

formation and indicate that one role for the early integration of visual information in auditory cortex

is to support auditory scene analysis.

## Figure Legends

**Figure 1: Hypothesis and experimental design**

**a** Conceptual model illustrating how binding can be identified as a distinct form of multisensory integration. Multisensory binding is defined as a subset of multisensory integration that results in the formation of a crossmodal object. During binding, all features of the audio-visual object are linked and enhanced including both those features that bind the stimuli across modalities (here temporal coherence between auditory (A) intensity and visual (V) luminance) and orthogonal features such as auditory pitch and timbre, and visual colour and size. Other forms of multisensory integration would result in only the features that promote binding being enhanced - here auditory intensity and visual luminance. To identify binding therefore requires a demonstration that non-binding features (e.g. here pitch, timbre, colour or size) are enhanced. **b** When two competing sounds (red and blue waveforms) are presented they can be separated on the basis of their features, but may elicit overlapping neuronal representations in auditory cortex. **c** Hypothesised enhancement in auditory stream segregation when a temporally coherent visual stimulus enables multisensory binding. When the visual stimulus changes coherently with the red sound (A1, top) this sound is enhanced and the two sources are better segregated. Perceptually this would result in more effective auditory scene analysis and an enhancement of the non-binding features. **d** Stimulus design: Auditory stimuli were two artificial vowels (denoted A1 and A2), each with distinct pitch and timbre and independently amplitude modulated with a noisy low pass envelope. **e** Visual stimulus: a luminance modulated white light was presented with one of two temporal envelopes derived from the amplitude modulations of A1 and A2. **f** illustrates the stimulus combinations that were tested experimentally in *single stream* (a single auditory visual pair) and *dual stream* (two sounds and one visual stimulus) conditions.

**Figure 2: Visual stimuli can determine which sound stream auditory cortical neurons follow in a mixture.**

Spiking responses from an example unit in response to **a,** single stream AV stimuli used as decoding templates and **b**, dual stream stimuli. In each case rasters and PSTHs are illustrated. When the visual component of the dual stream was V1, the majority of trials were classified as A1V1 (82% (19/23) of trials), and A2V2 when the visual stimulus was V2 (26% (6/23) of responses classified as A1V1 (see also green data point in c). **c-f** population data for awake (**c,d,** 541 units) and anesthetised (**e,f** 661 units) datasets. In each case the left panel (c,e) shows the distribution of decoding values according to the visual condition and the right panel (**d,f**) shows the population mean (± SEM) Pairwise comparisons revealed significant effect of visual condition on decoding in both datasets (p<0.001).

**Figure 3: Visual stimuli shape the neural representation of an auditory scene.**

In an additional control experiment (n=89 units recorded in awake animals), the responses to coherent AV and auditory-only (A Only) single stream stimuli were used as templates to decode dual stream stimuli either accompanied by visual stimuli (V1/V2) or in the absence of visual stimulation (no visual). Spiking responses from an example unit in response to **a,** single stream auditory stimuli which were used as decoding templates to decode the responses to dual stream stimuli in **b**, in each

case the auditory waveform, rasters and PSTHs are shown. **c**, population data showing the proportion of responses classified as A1 when the visual stimulus was V1 or V2 when decoded with auditory-only templates. **d,** Mean (± SEM) values for these units when decoded with A-only templates (i.e. data from **c**), AV templates (as in Fig.2) or in the absence of a visual stimulus. The green data point in **c** depicts the example in **a, b**. Pairwise comparisons revealed significant effect across visual conditions in both datasets (* indicates p<0.001).

**Figure 4: Temporally coherent changes in visual luminance and auditory intensity enhance the representation of auditory timbre**

**a** Example unit (from the awake dataset) showing the influence of visual temporal coherence on spiking responses to dual stream stimuli with (red PSTH) or without (black PSTH) timbre deviants. Shaded rectangles indicate the 200 ms window over which the timbre deviant occurred and over which analysis was conducted. **b-d** timbre deviant discrimination in the awake dataset. Two deviants were included in each auditory stream giving a possible maximum of 4per unit **b,** Histogram showing the number of deviants (out of 4) that could be discriminated from spiking responses **c,** Box plots showing the average timbre deviant discrimination scores in the single stream condition across different visual conditions (Coh: coherent, ind: independent). The boxes show the upper and lower quartile values, and the horizontal lines indicates the median. **d,** Discrimination scores for timbre deviant detection in dual stream stimuli. Discrimination scores are plotted according to the auditory stream in which the deviant occurred and the visual stream that accompanied the sound mixture. **e-g** show the same as **b-d** but for the anesthetised dataset.

**Figure 5: Auditory-visual temporal coherence enhances neural coding in auditory cortex**.

A pattern classifier was used to determine whether neuronal responses were informative about auditory or visual stimuli. The responses to single stream stimuli are shown for two example units, with responses grouped according to the identity of the auditory (**a, b**, for an auditory discriminating unit) or visual stream (**c, d**, for a visual discriminating unit). In each case the stimulus amplitude (a,b) / luminance (c,d) waveform is shown in the top panel with the resulting raster plots and PSTHs below. **e, f**: Decoder performance (mean ± SEM) for discriminating stimulus identity (coherent: A1V1 vs. A2V2, purple; independent: A1V2 vs. A2V1, blue) in auditory and visual classified units recorded in awake (e) and anaesthetised (f) ferrets. Pairwise comparisons for decoding of coherent versus independent stimuli (*** indicates p<0.001 ).

**Figure 6: Visual stimuli elicit reliable changes in the phase of the local field potential**

**a, b** Example LFP responses to single stream stimuli across visual conditions. Data obtained from the recording site at which multiunit spiking activity discriminated auditory stream identity in Fig. 5 a and b. The amplitude waveforms of the stimuli are shown in the top panel, with the evoked LFP underneath (mean across 21 trials). The resulting inter-trial phase coherence (ITPC) values are shown in the bottom two panels. **c, d** ITPC was calculated for coherent and independent AV stimuli separately and compared to a null distribution (ITPC across). Single stream phase dissimilarity values (PDI) were calculated by comparing ITPC within values to the ITPC across null distribution (**e, f**). **g,I** Population mean inter-trial phase coherence (ITPC) values across frequency for coherent (**g**) and independent (**i**) conditions. Dots indicate frequencies at which the ITPC-within values were significantly greater than the ITPC-across values (Pairwise t-test, $\alpha = 0.0012$, Bonferroni corrected for 43 frequencies). **k:** Mean (±SEM) single stream phase dissimilarity index (PDI) values for coherent and independent stimuli in animals. Black dots indicate frequencies at which the temporally coherent single stream PDI is significantly greater than in the independent conditions (p<0.001). **h,j,l**

as **g,l,k** for anesthetised dataset. **m, n,** mean ±SEM dual stream PDI values for awake (m) and
anesthetised (n) datasets.

**Figure 7: Visual stimulus induced LFP phase changes in auditory cortex are mediated in visual**
**cortex**

**a** Schematic showing the location of auditory cortical recording sites and the location of a cooling
loop (black, grey line marks the 500 um radius over which cooling is effective (Wood et al., 2017)
which was used to inactivate visual cortex. **b** Spike rate responses in Auditory Cortex (top row) and
visual cortex (bottom row, sites >500 um from the loop) in response to noise bursts or light flashes
before, during and after cooling. **c,d**, inter-trial phase coherence values for the coherent (**c**) and
independent stimuli (**d**) AV stimuli recorded in auditory cortex (AC) prior to cooling visual cortex (VC)
compared to the shuffled null distribution (ITPC across). Asterisks indicate the frequencies at which
the ITPC values are significantly different from the shuffled ITPC-across distribution **e** single stream
Phase Dissimilarity Index values calculated from the ITPC values in **c** and **d**. **f-h** - as c-e but while
visual cortex was cooled to 9 degrees. **i-n** as c-h but for sites in visual cortex >500um from the
cooling loop. c-h includes data from 83 sites from 6 electrode penetrations, i-n includes data from 47
sites from 5 penetrations.

## Acknowledgments

This work was funded by grants to each author: JKB: Wellcome Trust / Royal Society WT098418MA;

Biotechnology and Biological Sciences Research Council (BB/H016813/1), and an Action on Hearing

Loss Studentship (596: UEI: JB); RKM: NIH K99DC014288 and Hearing Health Foundation Emerging

Research Grant; AKCL: NIH R01DC013260; and an International Exchanges Scheme award from the

Royal Society to JKB and AKCL.

## Author contributions

HA, RKM, AKCL, JKB Conception and design, HA, SMT, KCW, GPJ, JKB Acquisition of data, HA, JKB

Analysis and interpretation of data, HA, SMT, RKM, AKCL, JKB Drafting or revising the article.

## References

Azouz, R., and Gray, C.M. (1999). Cellular mechanisms contributing to response variability of cortical
neurons in vivo. The Journal of neuroscience 19, 2209-2223.
Bizley, J.K., Maddox, R.K., and Lee, A.K. (2016). Defining Auditory-Visual Objects: Behavioral Tests
and Physiological Mechanisms. Trends in Neurosciences.

Bizley, J.K., Nodal, F.R., Bajo, V.M., Nelken, I., and King, A.J. (2007a). Physiological and anatomical
evidence for multisensory interactions in auditory cortex. Cereb Cortex 17, 2172-2189.

Bizley, J.K., Nodal, F.R., Bajo, V.M., Nelken, I., and King, A.J. (2007b). Physiological and anatomical
evidence for multisensory interactions in auditory cortex. Cereb Cortex 17, 2172-2189.

Bizley, J.K., Walker, K.M.M., King, A.J., and Schnupp, J.W.H. (2013). Spectral timbre perception in
ferrets: Discrimination of artificial vowels under different listening conditions. J Acoust Soc Am 133,
365-376.

Bizley, J.K., Walker, K.M.M., Silverman, B.W., King, A.J., and Schnupp, J.W.H. (2009). Interdependent
Encoding of Pitch, Timbre, and Spatial Location in Auditory Cortex. Journal of Neuroscience 29, 2064-
2075.

Brosch, M., Selezneva, E., and Scheich, H. (2015). Neuronal activity in primate auditory cortex during
the performance of audiovisual tasks. European Journal of Neuroscience 41, 603-614.

Budinger, E., Heil, P., Hess, A., and Scheich, H. (2006). Multisensory processing via early cortical
stages: connections of the primary auditory cortical field with other sensory systems. Neuroscience
143, 1065-1083.

Cantone, G., Xiao, J., and Levitt, J.B. (2006). Retinotopic organization of ferret suprasylvian cortex.
Visual neuroscience 23, 61-77.

Chandrasekaran, C., Lemus, L., and Ghazanfar, A.A. (2013). Dynamic faces speed up the onset of
auditory cortical spiking responses during vocal detection. Proceedings of the National Academy of
Sciences 110, E4668-E4677.

Chandrasekaran, C., Trubanova, A., Stillittano, S., Caplier, A., and Ghazanfar, A.A. (2009). The natural
statistics of audiovisual speech. Plos Comput Biol 5, e1000436.

Chandrasekaran, C., Turesson, H.K., Brown, C.H., and Ghazanfar, A.A. (2010). The influence of natural
scene dynamics on auditory cortical activity. The Journal of Neuroscience 30, 13919-13931.

Crosse, M.J., Butler, J.S., and Lalor, E.C. (2015). Congruent visual speech enhances cortical
entrainment to continuous auditory speech in noise-free conditions. The Journal of Neuroscience 35,
14195-14204.

Crosse, M.J., Di Liberto, G.M., and Lalor, E.C. (2016). Eye Can Hear Clearly Now: Inverse Effectiveness
in Natural Audiovisual Speech Processing Relies on Long-Term Crossmodal Temporal Integration. The
Journal of Neuroscience 36, 9888-9895.

Denison, R.N., Driver, J., and Ruff, C.C. (2013). Temporal structure and complexity affect audio-visual
correspondence detection. Front Psychol 3.

Desimone, R., and Duncan, J. (1995). Neural mechanisms of selective visual attention. Annual review
of neuroscience 18, 193-222.

Elhilali, M., Ma, L., Micheyl, C., Oxenham, A.J., and Shamma, S.A. (2009). Temporal Coherence in the
Perceptual Organization and Cortical Representation of Auditory Scenes. Neuron 61, 317-329.

Foffani, G., and Moxon, K.A. (2004). PSTH-based classification of sensory stimuli using ensembles of
single neurons. J Neurosci Methods 135, 107-120.

Ghazanfar, A.A., Maier, J.X., Hoffman, K.L., and Logothetis, N.K. (2005). Multisensory integration of
dynamic faces and voices in rhesus monkey auditory cortex. The Journal of Neuroscience 25, 5004-
5012.

Golumbic, E.Z., Cogan, G.B., Schroeder, C.E., and Poeppel, D. (2013). Visual input enhances selective
speech envelope tracking in auditory cortex at a "cocktail party". The Journal of Neuroscience 33,
1417-1426.

Henry, M.J., and Obleser, J. (2012). Frequency modulation entrains slow neural oscillations and
optimizes human listening behavior. Proceedings of the National Academy of Sciences 109, 20095-
20100.

Innocenti, G.M., Manger, P.R., Masiello, I., Colin, I., and Tettoni, L. (2002). Architecture and callosal
connections of visual areas 17, 18, 19 and 21 in the ferret (Mustela putorius). Cereb Cortex 12, 411-
422.

Jacobs, J., Kahana, M.J., Ekstrom, A.D., and Fried, I. (2007). Brain oscillations control timing of single-
neuron activity in humans. The Journal of neuroscience 27, 3839-3844.
Kaur, S., Rose, H., Lazar, R., Liang, K., and Metherate, R. (2005). Spectral integration in primary
auditory cortex: laminar processing of afferent input, in vivo and in vitro. Neuroscience 134, 1033-
1045.
Kayser, C., Petkov, C.I., and Logothetis, N.K. (2008). Visual modulation of neurons in auditory cortex.
Cerebral Cortex 18, 1560-1574.
Kayser, C., Petkov, C.I., and Logothetis, N.K. (2009). Multisensory interactions in primate auditory
cortex: fMRI and electrophysiology. Hearing Res 258, 80-88.
Klimesch, W., Sauseng, P., and Hanslmayr, S. (2007). EEG alpha oscillations: The inhibition–timing
hypothesis. Brain Research Reviews 53, 63-88.
Lakatos, P., Chen, C.-M., O'Connell, M.N., Mills, A., and Schroeder, C.E. (2007). Neuronal oscillations
and multisensory interaction in primary auditory cortex. Neuron 53, 279-292.
Lakatos, P., Musacchia, G., O'Connel, M.N., Falchier, A.Y., Javitt, D.C., and Schroeder, C.E. (2013). The
spectrotemporal filter mechanism of auditory selective attention. Neuron 77, 750-761.
Lőrincz, M.L., Kékesi, K.A., Juhász, G., Crunelli, V., and Hughes, S.W. (2009). Temporal framing of
thalamic relay-mode firing by phasic inhibition during the alpha rhythm. Neuron 63, 683-696.
Lu, K., Xu, Y., Yin, P., Oxenham, A.J., Fritz, J.B., and Shamma, S.A. (2017). Temporal coherence
structure rapidly shapes neuronal interactions. Nature communications 8, 13900.
Luo, H., Liu, Z., and Poeppel, D. (2010). Auditory cortex tracks both auditory and visual stimulus
dynamics using low-frequency neuronal phase modulation. PLoS Biol 8, e1000445.
Luo, H., and Poeppel, D. (2007a). Phase patterns of neuronal responses reliably discriminate speech
in human auditory cortex. Neuron 54, 1001-1010.
Luo, H., and Poeppel, D. (2007b). Phase patterns of neuronal responses reliably discriminate speech
in human auditory cortex. Neuron 54, 1001-1010.
Maddox, R.K., Atilgan, H., Bizley, J.K., and Lee, A.K. (2015). Auditory selective attention is enhanced
by a task-irrelevant temporally coherent visual stimulus in human listeners. Elife 4, e04995.
Mazzoni, A., Panzeri, S., Logothetis, N.K., and Brunel, N. (2008). Encoding of naturalistic stimuli by
local field potential spectra in networks of excitatory and inhibitory neurons. PLoS Comput Biol 4,
e1000239.
Möttönen, R., Schürmann, M., and Sams, M. (2004). Time course of multisensory interactions during
audiovisual speech perception in humans: a magnetoencephalographic study. Neuroscience letters
363, 112-115.
Ng, B.S.W., Schroeder, T., and Kayser, C. (2012). A precluding but not ensuring role of entrained low-
frequency oscillations for auditory perception. The Journal of Neuroscience 32, 12268-12276.
O'Sullivan, J.A., Shamma, S.A., and Lalor, E.C. (2015a). Evidence for Neural Computations of
Temporal Coherence in an Auditory Scene and Their Enhancement during Active Listening. J
Neurosci 35, 7256-7263.
O'Sullivan, J.A., Shamma, S.A., and Lalor, E.C. (2015b). Evidence for neural computations of temporal
coherence in an auditory scene and their enhancement during active listening. The Journal of
Neuroscience 35, 7256-7263.
Okada, K., Venezia, J.H., Matchin, W., Saberi, K., and Hickok, G. (2013). An fMRI study of audiovisual
speech perception reveals multisensory interactions in auditory cortex. Plos One 8, e68959.
Okun, M., Naim, A., and Lampl, I. (2010). The subthreshold relation between cortical local field
potential and neuronal firing unveiled by intracellular recordings in awake rats. The Journal of
neuroscience 30, 4440-4448.
Park, H., Kayser, C., Thut, G., and Gross, J. (2016). Lip movements entrain the observers' low-
frequency brain oscillations to facilitate speech intelligibility. eLife 5.
Peelle, J.E., and Sommers, M.S. (2015). Prediction and constraint in audiovisual speech perception.
Cortex 68, 169-181.

Perrodin, C., Kayser, C., Logothetis, N.K., and Petkov, C.I. (2015). Natural asynchronies in audiovisual
communication signals regulate neuronal multisensory interactions in voice-sensitive cortex.
Proceedings of the National Academy of Sciences 112, 273-278.
Rahne, T., Deike, S., Selezneva, E., Brosch, M., Konig, R., Scheich, H., Bockmann, M., and Brechmann,
A. (2008). A multilevel and cross-modal approach towards neuronal mechanisms of auditory
streaming. Brain Research 1220, 118-131.
Samaha, J., Bauer, P., Cimaroli, S., and Postle, B.R. (2015). Top-down control of the phase of alpha-
band oscillations as a mechanism for temporal prediction. Proceedings of the National Academy of
Sciences 112, 8439-8444.
Schroeder, C.E., Lakatos, P., Kajikawa, Y., Partan, S., and Puce, A. (2008). Neuronal oscillations and
visual amplification of speech. Trends in cognitive sciences 12, 106-113.
Schyns, P.G., Thut, G., and Gross, J. (2011). Cracking the code of oscillatory activity. PLoS Biol 9,
e1001064.
Sumby, W.H., and Pollack, I. (1954). Visual contribution to speech intelligibility in noise. The journal
of the acoustical society of america 26, 212-215.
Szymanski, F.D., Rabinowitz, N.C., Magri, C., Panzeri, S., and Schnupp, J.W. (2011). The laminar and
temporal structure of stimulus information in the phase of field potentials of auditory cortex. The
Journal of Neuroscience 31, 15787-15801.
Teki, S., Barascud, N., Picard, S., Payne, C., Griffiths, T.D., and Chait, M. (2016). Neural Correlates of
Auditory Figure-Ground Segregation Based on Temporal Coherence. Cereb Cortex 26, 3669-3680.
Tukker, J.J., Fuentealba, P., Hartwich, K., Somogyi, P., and Klausberger, T. (2007). Cell type-specific
tuning of hippocampal interneuron firing during gamma oscillations in vivo. The journal of
neuroscience 27, 8184-8189.
Voloh, B., and Womelsdorf, T. (2016). A Role of Phase-Resetting in Coordinating Large Scale Neural
Networks During Attention and Goal-Directed Behavior. Frontiers in systems neuroscience 10.
Wang, X.-J. (2010). Neurophysiological and computational principles of cortical rhythms in cognition.
Physiological reviews 90, 1195-1268.
Wood, K.C., Town, S.M., Atilgan, H., Jones, A.K., and Bizley, J.K. (2017). Acute Inactivation of Primary
Auditory Cortex Causes a Sound Localisation Deficit in Ferrets. PLoS One DOI:
10.1371/journal.pone.0170264.
Zion Golumbic, E., Cogan, G.B., Schroeder, C.E., and Poeppel, D. (2013). Visual input enhances
selective speech envelope tracking in auditory cortex at a "cocktail party". J Neurosci 33, 1417-1426.

## Methods

**Animal preparation:** The experiments were approved by the Animal Welfare and Ethical Review Board of University College London and The Royal Veterinary College, and performed under license from the UK Home Office (PPL 70/7267) and in accordance with the Animals Scientific Procedures Act 1986. Neural responses were recorded in a total of 11 awake pigmented adult female ferrets (Mustela putorius furo; 1-5 years old). Data from 9 of these animals was used for the main experiment (532 units), data from 8 other animals (6/9 plus two other ferrets, 89 units) was collected for additional control analysis (Fig. 3e, Fig S3, S6). Animals were chronically implanted with recording electrodes and passively listening/watching stimuli while holding their head at a water spout and receiving continuous water reward. These animals were trained in various listening tasks for other studies. An additional 5 adult females were used to record responses under anaesthesia. Recording electrodes in awake animals targeted tonotopic auditory cortex (fields A1 and AAF and their low frequency boarders with tonotopic belt areas PPF and PSF).

Full methods for recording under anaesthesia can be found in Bizley et al., (2009). Briefly, ferrets were anesthetized with medetomidine (Domitor; 0.022mg/kg/h; Pfizer, Sandwich, UK) and ketamine (Ketaset; 5mg/kg/h; Fort Dodge Animal Health, Southampton, UK). The animal was intubated and the left radial vein was cannulated in order to provide a continuous infusion (5 ml/h) of a mixture of medetomidine and ketamine in lactated ringers solution augmented with 5% glucose, atropine sulfate (0.06 mg/kg/h; C-Vet Veterinary Products) and dexamethasone (0.5 mg/kg/h, Dexadreson; Intervet, UK). The ferret was placed in a stereotaxic frame in order to implant a bar on the skull, enabling the subsequent removal of the stereotaxic frame. The left temporal muscle was largely removed, and the suprasylvian and pseudosylvian sulci were exposed by a craniotomy, revealing auditory cortex (Kelly et al., 1986). The dura was removed over auditory cortex and the brain protected with 3% agar solution. The eyes were protected with zero-refractive power contact lenses.

The animal was then transferred to a small table in a sound-attenuating chamber. Body

temperature, end-tidal CO2, and the electrocardiogram were monitored throughout the experiment.

Experiments typically lasted between 36 and 56 h. Neural activity was recorded with multisite silicon

electrodes (Neuronexus Technologies) in a 1x 16, 2x 16 or 4x 8 (shank x number of sites)

configuration. For experiments in which visual cortex was cooled we extended the craniotomy

caudally to expose visual cortex and placed a cooling loop over the posterior suprasylvian gyrus.

Details of the manufacture of the cooling loop and validation of its efficacy in the ferret animal

model are provided in full in (Wood et al., 2017).

Full surgical methods for recording implanting electrode arrays to facilitate recording from awake

animals are available in Bizley et al. (2013). Briefly, animals were bilaterally implanted with WARP-16

drives (Neuralynx, Montana, USA) loaded with high impedance tungsten electrodes (FHC, Bowdoin,

USA) under general anaesthesia (medetomidine and ketamine induction, as above, isoflurane

maintenance 1-3%). Craniotomies were made over left and right auditory cortex, a small number of

screws were inserted into the skull for anchoring and grounding the arrays, and the WARP-16 drive

was anchored with dental acrylic and protected with a capped well. Animals were allowed to recover

for a week before the electrodes were advanced into auditory cortex. Pre-operative, peri-operative

and post-operative analgesia were provided to animals under veterinary advice. Recordings were

made over the next 1-2.5 years, with electrodes individually advanced every few weeks until the

thickness of auditory cortex was traversed.

Stimulus presentation: All stimuli were created using TDT System 3 hardware (Tucker-Davis

Technologies, Alachua, FL) and controlled via MATLAB (Mathworks, USA). For recordings in awake

animals, sounds were presented over two loud speakers (Visaton FRS 8). Water deprived ferrets

were placed in a dimly lit testing box (69 x 42 x 52 cm length x width x height) and received water

from a central reward spout located between the two speakers. Sound levels were calibrated using a

Brüel and Kjær (Norcross, GA) sound level meter and free-field ½-inch microphone (4191). Auditory

streams were presented at 65 dB SPL (Fig. 1a). Visual stimuli were delivered by illuminating the

spout with a white LED which provided full field illumination (Precision Gold N76CC Luxmeter, 0 to

36.9 lux). The animals were not required to do anything other than maintain their heads in position

at the spout where they were freely rewarded. Recording was terminated when animals were sated.

For anesthetised recordings, acoustic stimuli were presented using Panasonic headphones

(Panasonic RP-HV297, Bracknell, UK) at 65 dB SPL. Visual stimuli were presented with a white Light

Emitting Diode (LED) which was placed in a diffuser at a distance of roughly 10 cm from the

contralateral eye so that it illuminated virtually the whole contralateral visual field.

Stimuli and data acquisition: *Auditory stimuli* were artificial vowel sounds that were created in

Matlab (MathWorks, USA). In the behavioural experiment that motivated this study(Maddox et al.,

2015), stimuli were 14 seconds in duration. However, we adapted the stimulus duration in awake

recordings to 3 seconds in order to collect sufficient repetitions of all stimuli, and to ensure animals

maintained their head position facing forwards for the whole trial duration. The animals were

observed constantly via a webcam and recording was terminated / paused if the animal's head

moved from the centre spout. In the anesthetised recording stimulus streams were 14 seconds long,

as in the human psychophysics but we only analysed the first 3 seconds to ensure datasets were

directly comparable (see also Fig. S7 which replicates analysis for 3 second and 14 second stimuli).

Stimulus A1 was the vowel [u] (formant frequencies F1-4: 460, 1105, 2857, 4205 Hz, F0= 195Hz), A2

was [a] (F1-4: 936, 1551, 2975, 4263 Hz, F0= 175Hz). Streams were amplitude modulated with a

noisy lowpass (7 Hz cut-off) envelope. Unless specifically noted, the timbre of the auditory stream

remained fixed throughout the trial. However, we also recorded responses to auditory streams that

included brief timbre deviants. As in our previous behavioural study, deviants were 200ms epochs in

which the identity of the vowel was varied by smoothly changing the first and second formant

frequencies to and from those identifying another vowel. Stream A1 was morphed to/from [ɛ] (730,

2058, 2857, 4205 Hz) and A2 to/from [i] (437, 2761, 2975, 4263 Hz).

*Visual stimuli* were generated using an LED whose luminance was modulated with dynamics that

matched the amplitude modulation applied to A1 or A2. In single stream conditions a single auditory

and single visual stream were presented (e.g. A1V1, A1V2, A2V1, or A2V2) whereas in dual stream

conditions both auditory streams were presented simultaneously, accompanied by a single visual

stimulus (A12V1, A12V2, A12V1 A12V2) (Fig. 1e). Auditory streams were always presented from both

speakers so that spatial cues could not facilitate segregation, and stimulus order was varied pseudo-

randomly. In the anesthetised recordings each stimulus was presented 20 times. In the awake

dataset, where recording duration was determined by how long the ferret remained at the central

location (mean repetitions: 20, minimum: 14, maximum: 34).

During anaesthetised recordings, pure tone stimuli (150 Hz to 19 kHz in 1/3-octave steps, from 10 to

80 dB SPL in 10 dB, 100 ms in duration, 5 ms cosine ramped) were also presented. These allowed us

to characterize individual units and determine tonotopic gradients, so as to confirm the cortical field

in which any given recording was made. Additionally broadband noise bursts and diffuse light flashes

(100 ms duration, 70 dB SPL) were presented and used to classify a stimulus as auditory, visual or

auditory visual. LFPs were subjected to current source density analysis to identify sources and sinks

as described by Kaur et al. (2005)

Data Analysis: Electrophysiological data were analysed offline. Spiking activity and local field

potential signals were extracted from the broadband voltage waveform by filtering at 0.3-5kHz and

1-150 Hz respectively. Spikes were detected, extracted and then sorted with a spike-sorting

algorithm (WaveClus, Quiroga et al., 2004).

We used a Euclidean distance based pattern classifier (Foffani and Moxon, 2004) with leave-one-out

cross validation to determine whether the neuronal responses to different stimuli could be

discriminated. Spiking responses to a given stimulus were binned into a series of spike counts from

stimulus onset (0 s) to offset (3s) in 20 ms bins. The average across-repetition response to each

stimulus (excluding the to-be-classified response) were used as templates and the response to a

single stimulus presentation was classified by calculating the Euclidean distance between itself and

the template sweeps and assigning it to the closest template. To determine whether the classifier

performed significantly better than expected by chance, a 1000 iteration permutation test was

performed where trials were drawn (with replacement) from the observed data and randomly

assigned to a stimulus that was then used for template formation / decoding. A neural response was

considered to be significantly informative about stimulus identity if the observed value exceeded the

95th percentile of the randomly-drawn distribution.

This approach allowed us to classify units according to their functional properties: auditory units

discriminated two auditory stimuli based on the amplitude modulation of sound (A1 versus A2)

regardless of visual dynamics, (Fig. 2a, b), visual units discriminated visual presentations based on

temporal envelope of visual stimuli (V1 versus V2) regardless of auditory presentation (Fig.  2c, d)

and AV units could do both. This approach was extended to classify dual stream responses by using

the average response to each of the temporally coherent single stream stimuli (A1V1 or A2V2) as

templates. Performance was (arbitrarily) expressed as the proportion of responses classified as being

from the A1, and compared for the two dual stream stimuli with different visual conditions (Figure

5). To be considered in this analysis the response of a unit had to be informative about the single

stream stimuli (i.e. classified as either auditory and/or visual discriminating).

Phase/power dissimilarity analysis:  Local field potential recordings were considered for all sites at

which there was a significant driven spiking response, irrespective of whether that response could

discriminate auditory or visual stream identity. For the single stream trials, we computed a single

stream Phase Dissimilarity Index (PDI), which characterizes the consistency and uniqueness of the

temporal phase/power pattern of neural responses to continuous auditory stimuli (Luo and Poeppel,

2007a). This analysis compares the phase (or power) consistency across repetitions of the same

stimulus with a baseline of phase-consistency across trials in which different stimuli were presented.

In the first stage of PDI analysis, we obtained a time-frequency representation of each response

using wavelet decomposition with complex 7-cycle Morlet wavelets in 0.5 steps between 2.5–45 Hz,

resulting in 86 frequency points. Next, we calculated the inter-trial phase-coherence value (ITPC;

Equ.1) at each time-frequency point, across all trials in which the same stimulus was presented. For

each frequency band, the ITPC time-course was averaged over the duration of the analysis window

and across all repetitions to obtain the average *within-stimulus ITPC*.

958     
$$ITPC_{t,f} = \left| \frac{\sum_{k=1}^{N} e^{i\theta k,t,f}}{N} \right| \qquad \text{Equ.1}$$

In which N is equal to the number of trials, and $\theta$ is the phase of trial *k* at a given frequency (*f*) and

time (*t*). The *across-stimuli ITPC* was estimated using the same approach but using shuffled data,

such that the ITPC was computed across trials with the same auditory stimulus but randomly drawn

visual stimuli. The single stream phase dissimilarity index (Single stream PDI) was computed as the

difference between the ITPC value calculated for *within* trials and the ITPC values calculated *across*

visual trials (Equ.2). y. The dissimilarity function for each frequency bin *i* was defined as;

966     
$$Single\ Stream\ PDI_i = \frac{\sum_{j=1}^{N} ITPC_{ij}\ within_{vis}}{N} - \frac{\sum_{j=1}^{N} ITPC_{ij}\ ,across_{vis}}{N} \qquad \text{Equ.2}$$

Large positive PDI indicate that responses to individual stimuli have a highly consistent response on

single trials. Single stream PDI values were calculated for each stimulus type and then averaged

across stimuli to calculate values for temporally coherent and temporally independent auditory

visual stimuli. Single stream PDI was positive if within stimulus ITPC was larger than across-stimulus

ITPC (pairwise t-test, p<0.05 Bonferroni correction for 86 frequencies points) and was considered

significant if a minimum of 2 adjacent bins exceeded the corrected threshold. PDI magnitude values

were calculated by summing the PDI values across all significant frequencies.

Dual stream phase dissimilarity index (dual stream PDI) values were calculated by extending this

approach for dual stream stimuli with the goal of determining how the temporal envelope of the

visual stimulus influences the neural response to a sound mixture. To this end, we calculated the

*within-dual ITPC* from the A12V1 trials and A12V2 trials separately and *across-dual ITPC* by randomly

selecting trials from both stimuli (Equ.3). The within-dual and across-dual ITPCs were then averaged

over time and subtracted to yield the dual stream PDI (Equ.3).

$$Dual\ Stream\ PDI_i = \frac{\sum_{j=1}^{N} ITPC_{ij}\ within_{dual}}{N} - \frac{\sum_{j=1}^{N} ITPC_{ij}\ across_{dual}}{N}$$    Equ.3

Positive dual stream PDI values indicate that the time course of the neural responses was influenced

by visual input, despite the identical acoustic input. We determined whether the dual stream PDI

was greater if the *within_dual ITPC* was significantly larger than *across_dual ITPC* (pairwise t-test,

p<0.05 Bonferroni correction, as above). PDI magnitude values were calculated by summing the PDI

values across all significant frequencies.

Timbre deviant analysis: In order to determine how a visual stimulus influenced the ability to

decode timbre deviants embedded within the auditory streams we used the cross-validated pattern

classifier described above for analysing single stream stimuli to discriminate deviant from no-deviant

trials. Responses were considered over the 200 ms time window that the deviant occurred (or the

equivalent time point in the no-deviant stimulus) binned with a 10 ms resolution. Significance was

assessed by a 1000 iteration permutation test in which trials were randomly drawn with

replacement from deviant and no-deviant responses. The discrimination score was calculated as the

proportion of correctly classified trials.

**Code and Data Availability**

Data and MATLAB code is available from the authors upon request.

1000

1001 Supplementary Figures

1002

1003 **Supplementary figure 1: related to Figure 2**

1004 **Visual stimuli can determine which sound stream auditory cortical neurons follow in a mixture**
1005 **in auditory-discriminating units**

1006 The spiking responses of an example unit are shown to coherent single stream (**a**) and dual stream
1007 stimuli (**b**). This example unit was an auditory discriminating unit recorded from an awake animal.

When the visual stimulus was V1, 67% of trials were classified as A1V1, when the visual stimulus was V2 only 29% of trials were classified as A1V1.

**Supplementary figure 2: Related to Figure 2**

**Preference to the auditory stream with which the visual stimulus was coherent observed across all cortical layers and in three tonotopic fields.**

In the anesthetised animal, we determined tonotopic gradients from responses to pure tone stimuli to accurately confirm the cortical field in which any given recording was made. Recordings were made with linear shank electrodes, facilitating current source density analysis to identify the cortical layers. **a** Proportion of responses classified as A1 when the visual stimulus was V1 or V2 are shown for supragranular layers (SG, $t_{700}=5.686$, p<0.001), granular layer (G, $t_{878}=3.481$ p<0.001) and infragranular layers (IG, $t_{690}=4.418$, p<0.001). Data shown as mean (± SEM) across units. A two-way ANOVA across visual condition and cortical layers showed only a significant effect of visual condition (F (1, 2273) =64.288, p<0.001) but not of layers (F (1, 2273) =0.91, p =0.404) and no interaction was observed (F (1, 2273) =2.679, p=0.068). Significant post-hoc comparisons (p<0.05) are shown with an asterisk. **b,** In order to compare the magnitude of the effect of the visual stimulus across cortical layers we calculated visual modulation as the decoding score when the visual stimulus was V1 – the decoder score when the visual stimulus was V2. Plotted are the mean (± SEM) visual modulation across different layers. A one-way ANOVA across cortical layers showed a significant effect of cortical layer (F (2, 1134) =3.15, p =0.043). Post hoc comparison revealed that visual modulation in SG is greater than G. **c,** as for a but for different cortical fields (A1, $t_{798}$ =5.435, p<0.001; AAF, $t_{636}$ =4.302, p<0.001; PPF, $t_{510}$ =3.609, p<0.001; PSF, $t_{317}$ =0.932, p=0.352). A two way ANOVA across visual condition and cortical field showed only a significant effect of visual condition (Visual condition: F (1, 2267) =40.301, p<0.001; fields: F (1, 2267) =1.937, p=0.121; visual condition x fields: F (1, 2267) =1.804, p=0.144). Significant post-hoc comparisons (p<0.05) are shown with an asterisk. In order to compare the magnitude of the effects across cortical fields we again calculated a visual modulation value. **b,** Mean (± SEM) visual modulation across different

fields. A one-way ANOVA across cortical layers showed a no significant effect of cortical field (F (3, 1130) =2.19, p =0.0877).

1040

1041

1042

1043

1044

**Supplementary figure 3: Related to Figure 3**

**Auditory, visual and auditory-visual units are influenced by visual stimuli.**

In anesthetised animals we also recorded responses to 100 ms noise bursts and/or LED flashes in order to apply conventional classification of units as either auditory, visual or auditory-visual. Unit classification was based on a two way ANOVA of spike counts calculated over a 200ms window following stimulus onset with auditory and visual stimuli as factors. Units in which both auditory and visual stimuli significantly modulated spiking or in which there was a significant interaction between stimuli were classified as auditory-visual (AV). This analysis yielded recording sites at which units were classified as auditory only (A; n=160 units), visual only (V, n=53) and AV (grey, n=94). This allowed us to explore the visual modulation effects according to independently determined functional definitions of units as auditory, visual or auditory visual.

**a**, The proportion of responses classified as A1 when visual stimuli as V1 or V2 is plotted with each unit color coded according classification as A, V or AV. **b**, Mean (± SEM) values across the population. A two-way ANOVA revealed a significant effect of visual condition (F (1, 608) = 18.142, p<0.001), but no effect of unit type (F (2,608) = 0.765, p=0.4657), or interaction between factors (F (1,608) =2.474, p=0.0851) on the proportion of responses classified as A1.   Pairwise post-hoc comparisons confirmed a significant effect of visual stimulus in all subgroups (p<0.05). **c**, Mean (± SEM) dual stream phase dissimilarity index (PDI) values for recording sites categorised according to the spiking responses recorded there. Symbols indicate the frequencies at which the dual stream PDI index was significant (pairwise t-test, p<0.001 with correction). While the phase effects are greatest at the sites where visual activity was recorded, significant dual stream PDI values were observed in all three unit types. In all three cases significant phase coherence was seen at 12Hz, 13.5Hz-14.5Hz and 42.5-44.5Hz. Modulation at 10-12 Hz was only observed at sites in which AV and V responses were recorded.

1069

**Supplementary figure 4: Related to Figure 3**

**The effects of temporal coherence on single stream decoding and of visual identity on dual stream decoding are evident in both single and multi-units.**

**a,b** Discrimination accuracy for single stream decoding is indistinguishable for single units and multi-unit activity. Two way ANOVA revealed a significant effect of visual condition and no effect of unit type in discrimination accuracy values in awake recordings (a, Visual condition: $F(1,851) = 14.6607$ $p<0.01$, Unit Type: $F(1,851)= 1.02$, $p=0.312$; interaction: $F(1,851)=0.005$, $p=0.449$) and anaesthetised recordings (b, Visual condition: $F(1,1847) = 24.514$, $p<0.01$, Unit Type: $F(1,1847)= 0$, $p=0.986$; interaction: $F(1,1847)=0.66$, $p=0.418$) **c, d**. Impact of visual stimulus on dual stream decoding in single units and multi units. A two way ANOVA revealed a significant effect of visual condition but no effect of unit type in discrimination accuracy values in awake recordings (c, Visual condition: $F(1,449) = 19.41$ $p<0.01$, Unit Type: $F(1,851)= 1.07$, $p=0.414$; interaction: $F(1,851)=0.33$, $p=0.568$ ) and anaesthetised recordings (d, Visual condition: $F(1,1847) = 15.41$, $p<0.01$, Unit Type: $F(1,1847)= 1.1$, $p=0.506$;interaction: $F(1,1847)=0.66$, $p=0.418$).

1091

**Supplementary figure 5, related to Fig.6**

**Dynamic visual stimuli do not elicit reliable changes in LFP power.**

**a, b:** Mean (±SEM) single stream phase dissimilarity index values for coherent (A1V1 and A2V2) and independent (A1V2 and A2V1) stimuli at all recording sites at which there was a driven spiking response for awake (a) and anaesthetised (b) animals.

1097

1098

1099

1100

1101

**Supplementary figure 6: related to figure 6**

**Dual stream PDI values in the alpha range are independent of amplitude modulation rate.**

Stimuli were generated with three different amplitude modulation rates (<7Hz, as in the main experiment, <12Hz and <17Hz, values picked to avoid harmonics of 7 Hz) and responses to these were recorded in 92 units. Symbols indicate where the dual stream phase selectivity index was significant (pairwise t-test, p<0.05 with correction). In all three cases significant phase coherence is seen between 10Hz-11.5Hz, 19Hz-20Hz and 24-26 Hz.

1109

**Supplementary figure 7: related to experimental methods**

**Stimulus duration does not influence the frequency range across which significant single stream PDI values and dual stream PDI values are observed.**

We observed that phase coherence values were higher for longer duration stimuli and therefore in order to directly compare anesthetised and awake datasets, all analysis in the anesthetised dataset was restricted to the first three seconds of the stimulus. At longer stimulus durations the ITPC profile and resulting PDI varied more smoothly with frequency (a, c); However at both durations phase values were significantly different from zero at all frequencies. The pattern of significant phase selectivity values was also preserved across stimulus durations. (**b, d**). Frequency points at which the single stream PDI value and dual stream PDI values were similar in 3 second length (**a, b**) and 14 second length (**c, d**) Blue, red and black symbols indicate where the PDI was significant (pairwise t-test, a = 0.0012 with correction).