Title: Comparing three types of dietary samples for prey DNA decay in an insect

generalist predator

STEFANIYA KAMENOVA[1,2,3], REBECCA MAYER[4], OSKAR R. RUBBMARK[4], ERIC

COISSAC[5,6]*,* MANUEL PLANTEGENEST[2] and MICHAEL TRAUGOTT[4]

[1]Centre d'Etudes Biologiques de Chizé, Villiers-en-Bois, France, [2]Agrocampus Ouest

UMR1349 IGEPP, 65 rue de Saint-Brieuc, CS 84215, F-35042 Rennes, France,

[3]Integrative Biology, University of Guelph, Guelph, ON, Canada, [4]Mountain Agriculture

Research Unit, Institute of Ecology, University of Innsbruck, Technikerstraße 25, 6020

Innsbruck, Austria, [5]Université Grenoble-Alpes, Laboratoire d'Ecologie Alpine (LECA),

F-38000 Grenoble, France, [6]Centre National de la Recherche Scientifique, Laboratoire

d'Ecologie Alpine (LECA), F-38000 Grenoble, France

Corresponding author: Stefaniya Kamenova

Integrative Biology, University of Guelph, 50 Stone Road East, N1G 2W1, Guelph, ON,

Canada

stefaniya.kamenova@gmail.com

Keywords: diagnostic PCR, molecular diet analysis, prey detection, Carabidae, trophic

interactions

Running title: DNA decay in three types of dietary samples

Abstract

The rapidly growing field of molecular diet analysis is becoming increasingly popular

among ecologists, especially when investigating methodologically challenging groups

such as invertebrate generalist predators. Prey DNA detection success is known to be

affected by multiple factors, however the type of dietary sample has rarely been

considered. Here, we address this knowledge gap by comparing prey DNA detection

success from three types of dietary samples. In a controlled feeding experiment, using

the carabid beetle *Pterostichus melanarius* as a model predator, we collected

regurgitates, feces and whole consumers (including their gut contents) at different time

points post-feeding. All dietary samples were analyzed using multiplex PCR targeting

three different length DNA fragments (128 bp, 332 bp and 612 bp). Our results show

that both the type of dietary sample and the size of the DNA fragment contribute to a

significant part of the variation found in the detectability of prey DNA. Specifically, we

observed that in both regurgitates and whole consumers prey DNA was detectable

significantly longer for all fragment sizes than for feces. Based on these observations,

we conclude that prey DNA detected from regurgitates and whole consumers DNA

extracts are comparable, whereas prey DNA detected from feces, though still

sufficiently reliable for ecological studies, will not be directly comparable to the former.

Therefore, regurgitates and feces constitute an interesting, non-lethal source for dietary

information that could be applied to field studies in situations when invertebrate

predators should not be killed.

Introduction

Introduction

DNA-based diet analysis is rapidly being employed as a widespread tool for

empirically characterizing diet and trophic interactions in a broad range of vertebrates

and invertebrates (Traugott *et al.* 2013; Clare 2015). DNA-based methods for diet

analysis typically rely on the detection of short fragments of prey DNA, recovered from

predator's gut contents (*e.g.* Leray *et al.* 2015; Mollot *et al.* 2014) or other types of

dietary samples such as feces, regurgitates, or whole consumers (*e.g.* Ibanez *et al.*

2013; Kartzinel *et al.* 2015; Thalinger *et al.* 2016; Wallinger *et al.* 2015). The success of

DNA-based approaches to analyze trophic interactions is mainly due to the fact that

they allow the direct and accurate identification of trophic links from minute amount of

starting material, even for very small-sized organisms such as mites (Pérez-Sayas *et al.*

2015) or zooplankton (Durbin *et al.* 2012). Furthermore, the rapid growth of public

sequence databases and methodological improvements in detection sensitivity and

high-throughput technology, offer time- and cost-effective procedures applicable to a

great variety of ecological systems and to large sample sizes (*e.g.* Valentini *et al.* 2009;

Pompanon *et al.* 2012; Sint *et al.* 2011).

DNA-based diet analysis is particularly valuable for studying invertebrate

generalist predators (Symondson 2012). Indeed, DNA methods offer a sensitive and

flexible alternative to traditional behavioural or dissecting techniques that often fail to

detect prey that does not leave hard remains in these cryptic liquid feeders (Traugott *et*

al. 2013). DNA techniques are, however, also subject to bias and prey DNA detection

success could be hampered by a variety of factors among which the type of dietary

sample could play an important role (King *et al.* 2008; Pompanon *et al.* 2012; Traugott

et al. 2013). In the case of arthropods, whole body extracts are usually the most

convenient source of dietary DNA that avoid laborious dissections. Besides the

drawbacks of a lethal approach (e.g. sacrificing rare or endangered species; affecting

population dynamics, etc.), whole body extracts may pose additional challenges

especially in the case of DNA metabarcoding diet analysis. As DNA metabarcoding

combines general primers and high-throughput sequencing, the concomitant

amplification of consumer DNA usually compromises the detection success of scarcer

and degraded prey DNA (*e.g.* Shehzad *et al.* 2012; Piñol *et al.* 2014).

Waldner & Traugott (2012) demonstrated that regurgitates, a fluid mixture

containing semi-digested prey remains and digestive enzymes, obtained from predatory

carabid beetles provided superior prey DNA detection rates compared to whole body

DNA extracts. Another prospective source of food DNA are feces, although their use as

a dietary source in invertebrates is still uncommon (*e.g.* Ibanez *et al.* 2013; Redd *et al.*

2014; Sint *et al.* 2015). Usually, both regurgitates and feces seem to provide similar or

better detection rates compared to whole body extracts (Durbin *et al.* 2012; Egeter *et al.*

2015; Unruh *et al.* 2016), and contain comparatively much less consumer DNA, making

them putatively an ideal source for metabarcoding diet analysis. Nevertheless, to date

we lack a comparative and quantitative assessment of the respective efficiency in

detection success between whole bodies, regurgitates and feces as well as prospective

interactions with other sources of non-dietary variation such as the target DNA fragment

size.

In this study, we address this knowledge gap by comparing the prey DNA

detection rates for three types of dietary samples: whole consumers including their gut

content, regurgitates and feces. Using a controlled feeding experiment involving a

widespread carabid predator, *Pterostichus melanarius* (Coleoptera: Carabidae) we test

the following hypotheses: (i) post feeding prey DNA detection success should be similar

or better in regurgitates compared to whole beetles due to lesser degradation of prey

DNA in the former; (ii) prey DNA detection success should be lower in feces compared

to regurgitates and whole bodies as faecal material represents the final stage of the

digestion process; and (iii) prey DNA detection should decrease with increasing DNA

fragment size and the time post-feeding for all types of samples.

Material & Methods

Sampling and maintenance of predators

Pterostichus melanarius individuals were collected by dry pitfall traps in two

adjacent maize fields situated at the experimental site of INRA Le Rheu (Ille-et-Vilaine,

France; GPS coordinates: 48.10744282N; 1.78830482W). Regular 24-hour trapping

sessions were conducted in July – August 2013 until a sufficient number of individuals

had been collected. All living beetles were brought to the laboratory where they were

identified to species level and individually placed in plastic containers filled with loam.

Beetles were stored at room temperature and continuously provided with water and

food (field-collected earthworms and small pieces of apple).

Feeding experiment

Prior to the feeding experiment, beetles were starved for 96 h in fresh individual

plastic Petri dishes (5 cm diameter) containing only a droplet of water. After the

starvation period, all beetles were transferred to a new Petri dish and provided with one

freshly freeze-killed mealworm (*Tenebrio molitor*, Coleoptera: Tenebrionidae) cut in half.

Carabids were allowed to feed for one hour in a dark climatic chamber at 20°. After

feeding all beetles which had fully consumed the mealworm were transferred into fresh

Petri dishes with no food. Beetles were stored at room temperature and continuously

provided with water during the experiment.

For the "whole beetle" treatment, batches of 10 randomly chosen carabids were

frozen in 2-mL reaction tubes by immersion in liquid nitrogen at 0, 12, 24, 36, 48, 60, 72

and 96 hours post-feeding. Immersion in liquid nitrogen was necessary as previous

tests showed that living beetles do not die immediately after placement at -20°C,

leading them to regurgitate into the reaction tube. After immersion, all whole beetles

were stored at -20°C. Thirteen starved beetles were never allowed to feed and they

were freeze-killed at 0 h to be used as negative controls. For the "regurgitate" treatment,

batches of 10 randomly chosen individuals were allowed to regurgitate on a cotton wool

tip according to the protocol described in Waldner & Traugott (2012) at 0, 12, 24, 36, 48,

60, 72 and 96 h post-feeding. After regurgitation, all beetles per given time-point were

freeze-killed and stored at -20°C. Exactly the same procedure at each time point was

applied on a control tip without touching a beetle for checking potential DNA carry-over

contaminations. All samples were stored at -20°C prior to DNA extraction and PCR. For

the "feces" treatment, 20 carabid beetles were placed in new Petri dishes after feeding

with a droplet of clean water. Carabids were continuously checked for feces production

at every 6 hours. Detected feces were immediately frozen within the Petri dish at -20°C

whereupon the corresponding carabid individual was transferred into a new Petri dish.

Feces production was monitored until all beetles died.

Molecular diet analysis

Regurgitate and fecal samples were directly lysed in 200 µl TES Lysis Buffer

(Macherey-Nagel, Germany) and 5 µl Proteinase K (10 mg/mL) overnight at 56°C. The

whole beetles were previously ground using three 4 mm stainless steel beads (Lemoine

S.A.S, Rennes, France) within a volume of 620 µl TES Lysis Buffer and 10 µl

Proteinase K (10 mg/mL) per beetle. Tissues were disrupted by a 1-minute bead-

beating step using a professional paint mixer (Fluid Management Inc., Wheeling, IL,

USA). All samples were incubated overnight at 56°C. Respectively 2, 6, and 2 lysate

blanks (i.e. no DNA material) were carried out for the whole beetles, fecal and

regurgitate treatments. DNA was extracted in batches of 92 samples using the Biosprint

96 DNA Blood Kit (Qiagen, Hilden, Germany) on a Biosprint 96 extraction robotic

platform (Qiagen) following the manufacturer's instructions. DNA was finally diluted in

200 µl TE buffer (0.1 M TRIS, pH 8, 10 mM EDTA) and the extracts were stored at -28

°C. To avoid contamination, DNA extractions were done in a separate pre-PCR

laboratory using a UVC-equipped laminar flow hood. To check for sample-to-sample

cross-contamination, four extraction negative controls (PCR-grade RNase-free water

instead of lysate) were included within each batch of 92 samples. All of these controls

tested negative using the diagnostic PCR assay described below.

The DNA extracts were screened with a multiplex PCR assay targeting three

overlapping COI mtDNA fragments of *T. molitor,* i.e. 128 bp, 332 bp and 612 bp. The

primer mix contained 6 µM of primers Ten-mol-S210 (5'-

TACCGTTATTCGTATGAGCAGTAT-3') and Ten-mol-A212 (5'-

CGCTGGGTCAAAGAAGGAT-3') as well as 2 µM of primers Ten-mol-S232 (5'-

TAATAAGAAGAATTGTAGAAAACGGG-3') and Ten-mol-S231 (5'-

TCATTTTTGGAGCGTGATCC-3') (Oehm *et al*. 2011; Sint *et al*. 2011). Each 10 µl PCR

consisted of 1.5 µl template DNA, 5.0 µl of 2x Multiplex PCR Kit reaction mix (Qiagen),

1.0 µl of primer mix, 0.5 µl of bovine serum albumin (BSA, 10 mg ml^{-1}), and 2.0 µl of

PCR-grade RNase-free water (Qiagen) to adjust the volume. Thermocycling was

conducted in Eppendorf Mastercyclers (Eppendorf, Hamburg Germany) and cycling

conditions were 15 min at 95 °C, 35 cycles of 30 sec at 94 °C, 90 sec at 63 °C, 1 min at

72 °C, and final elongation 10 min at 72 °C. To check for amplification success and

DNA carry-over contamination, two positive (mealworm DNA) and two negative controls

(PCR water instead of DNA) were included within each PCR, respectively.

The PCR products obtained were visualized using QIAxcel, an automated capillary

electrophoresis system (Qiagen), with method AL320. The results were scored with

Biocalculator Fast Analysis Software version 3.0 (Qiagen) and the threshold was set to

0.07 relative fluorescent units. Samples above this threshold and showing the expected

fragment length were counted as positives. All DNA extracts that were tested negative

in the first run were re-tested with general primers (Folmer *et al*. 1994) in a second PCR

to check for any amplifiable DNA (all of these samples tested positive). To ensure

contamination-free conditions, PCR preparation and visualization of PCR products were

done in two separate laboratories (workflow: from pre- to post-PCR areas).

Statistical analyses

A generalized linear mixed model was built to fit a logistic regression on the DNA

detection data. We integrated three fixed effects into the model: two qualitative factors,

the marker size (128 bp, 332 bp, 612 bp) and the sample type (regurgitates, feces or

whole body DNA extracts), and one continuous variable, the time post-feeding. To

compensate for non-independence in collection of feces individuals were included as a

random effect. The model was fitted using the *glmm* function from the R package

"*glmm"* (https://cran.r-project.org/web/packages/glmm). Models were fit using a Monte

Carlo sample size of 1024 with 10,000 iterations. The distribution of each of the model

parameters was approximated to a normal distribution using the maximum goodness-of-

fit estimation with the "*fitdist"* function available in the R package "*fitdistrplus"*

(Delignette-Muller & Dutang 2015). The variance in detectability rates explained by the

model was estimated using the coefficient of determination method (Tjur 2009). Tests of

the differences between mean detectability rates for each of the qualitative factors

(marker length and sample type) were conducted using a Z-test. The time point for a

prey detection probability of 50% (i.e. the time point at which on average half of the

individuals show positive for the target prey) was determined for each dietary sample

and DNA fragment size. To compensate for false discovery rate in multiple testing

comparisons between fragments were based on 95% confidence limits (CI) as

suggested by Greenstone et al (2013). All statistical analyses were conducted using the

R software (R Core Team 2013).

Results

Detectability of mealworm DNA in *P. melanarius* decreased with increasing post-

feeding time and prey DNA fragment length for the three dietary samples (Fig. 1, small

vs medium and small vs large fragments: $p<0.001$; medium *vs* large fragment: $p=0.08$),

with post-feeding detection time intervals being longest for the shortest DNA fragment

(Fig. 1 A, B, C). We also observed a significant effect of the dietary sample type, with

prey DNA detection success being significantly lower in feces compared to regurgitates

and whole beetles for all the three fragment sizes (Fig. 1, in all cases $p<0.001$). There

was also a tendency for longer post-feeding detection periods in regurgitates compared

to whole beetles (Fig. 1A, B) but differences were not significant ($p=0.6$). Our model

fitted the data well for all of the three dietary samples: regurgitates (Fig. 1A), whole

beetles (Fig. 1B) and feces (Fig. 1C), and explained 50% of the variance in DNA

detectability. Raw data are presented in Table 1. For the small prey DNA fragment, 50%

detection time was the highest for regurgitates (94 hours) but the value significantly

dropped by more than half for the medium fragment (42 hours) and was significantly

shortest for the largest prey DNA fragment (30.6 hours; Table 2). In feces 50%

detection probabilities were the lowest for all the three DNA fragment sizes, with only 19

hours for the largest DNA fragment (612 pb) and a significantly shorter detection

probability for the medium prey DNA fragment when compared to the regurgitate

samples (Table 2).

Discussion

How long prey DNA can be detected in a sample is determined by a range of

interacting factors related to the environment, the predator-prey system and the

molecular techniques used. These might affect results, but how is difficult to disentangle

without conducting comprehensive experiments that explicitly account for factor

multiplicity. Here, we compared multiple dietary samples from one species of

invertebrate consumer, in a controlled feeding experiment, and assessed how the

combined effects of the type of dietary sample and DNA fragment size will affect the

prey DNA detection probability over time since feeding. Our results show that each of

these factors significantly affects the rate at which the probability of detecting prey DNA

decreases over time. Consistent with our hypothesis, the time during which DNA could

be detected was the longest for regurgitates, for each of the three tested prey DNA

fragment sizes. While this was not significantly different from DNA detected from whole

beetles, prey DNA contained in feces was detectable for a significantly shorter time for

all three fragment sizes.

Our results support the general assumption that regurgitates constitute a good

alternative source of prey DNA (Waldner & Traugott 2012; Wallinger *et al*. 2015). Such

an alternative could be particularly useful in manipulative food web experiments, where

the removal and killing of the targeted predators during sampling could disturb the

system under study. As 79% of predaceous land-dwelling arthropods use extra-oral

digestion (Cohen 1995), this approach is potentially applicable to a large array of taxa

and ecological situations. Furthermore, by containing comparatively less predator DNA,

regurgitates could also be a valuable source of dietary data in DNA metabarcoding

studies involving the use of general primers (Waldner & Traugott, 2012).

Nevertheless, the use of regurgitates could entail some additional limitations

such as the detection of only the most recent diet items, and probably represents

merely a narrow fraction of individual's diet, especially in generalist feeders with

frequent switching behavior such as carabids (Lövei & Sunderland, 1996). Thus, the

choice of the most appropriate dietary sample will most likely consist in a trade-off

between DNA detection rates and representativeness in terms of diet according to the

focus of interest.

Fecal samples could provide a more integrated picture of individual's diet. Our

results show that overall prey DNA detection was lower compared to regurgitates and

whole bodies. Note, however, that this was true only for the medium sized fragment

when considering the 50% prey DNA detection probability, with significantly lower post-

feeding interval found in feces compared to regurgitates. This indicates that feces

overall are a good source of dietary DNA at least in *P. melanarius* beetles. Similarly,

earlier study in wolf spiders showed that prey DNA was detectable in spider feces albeit

in lower rates compared to whole body DNA extracts (Sint *et al.* 2015). As spiders

represent an important group of generalist feeders that typically do not regurgitate, the

sole non-lethal dietary sample that could be collected are feces. Also, we cannot rule

out the possibility that in our case DNA prey detection success in feces was lower

simply due to the constraints of the experiment. As carabids were checked for feces

every 6 hours, feces deposited earlier within that timeframe could have experienced

higher DNA degradation due to longer exposure to ambient temperature, thus resulting

in increased variability in DNA prey detection success.

In a recent paper Unruh et al. (2016) even show that there is no difference in

DNA detection between whole bodies and feces in the insect predator *Forficula*

auricularia. While the authors do not discuss the possible mechanisms behind this

observation, results tend to suggest that feces could be at least as good dietary source

as whole body extracts for predatory insects such as *F. auricularia*. Hence, feces

remain a viable non-lethal dietary source in certain situations, as detection rates are

generally high.

Yet, the question of the time it takes for prey DNA to travel through the digestive

tract of insects and how this varies across different taxa remains. Having a better

understanding about the temporal aspects of digestion in insects in general, and

particularly in carabid beetles is important. Carabids are generalist, mobile feeders with

frequent switching behavior (Lövei & Sunderland, 1996) meaning that frequent diet

shifts but long prey DNA retention periods may result for instance in an overestimation

of consumption rates or in a mismatch between diet composition and estimations of

prey availability at the place where dietary samples were collected. We also usually do

not consider whether this problem could be exacerbated in herbivorous species as the

digestion process of plant DNA in insects can last much longer as compared to animal

DNA (Staudacher *et al.* 2011; Wallinger *et al.* 2013, 2015). For instance, results have

shown that [14]C-inulin labelled prey in carabid beetles could still be detected in feces up

to five days post-feeding (Cheeseman and Gillott 1987). It will be interesting to confront

these findings with observations about prey DNA transit.

Here, we also show that prey DNA detection continuously decreases over time

for all the three types of dietary samples, with longer fragments (332-612 bp) decaying

more rapidly compared to the shorter one (128 bp). These results meet our

expectations and corroborate the general idea that digested DNA molecules break

down relatively quickly and that the size of the targeted prey DNA fragment affects post-

feeding prey DNA detection (Agustí *et al.* 2003; von Berg *et al.* 2008). In line with

previous studies, our results support the idea that targeting short to medium size DNA

fragments in DNA diet analysis is essential in order to maximize the prey detection

(Deagle *et al.* 2006; Valentini *et al.* 2009). Nonetheless, if a recent feeding event is the

focus, then targeting longer fragments might actually be a better strategy to ensure that

only the most recent prey items are detected. Additionally, as in DNA metabarcoding

diet analysis there is generally a trade-off between DNA fragment length and taxonomic

resolution, targeting longer DNA fragments – within a certain range - could indeed

improve the taxonomic discrimination of prey species (Pompanon *et al.* 2012). In this

study, the most important observed source of variation in terms of prey DNA detection,

besides time post feeding, is DNA fragment size. This could have profound implications

in metabarcoding studies where the DNA fragment size usually needs to be optimized in

order to meet criteria for both optimal detectability and taxonomic resolution (Taberlet *et*

al. 2012). It would be interesting to simultaneously explore the decay rate of a larger

array of DNA fragments of different lengths in order to assess whether a general

relationship between DNA length and detectability can be drawn despite the many other

sources of variability detected in previous studies. One might speculate that a

consistent relationship between DNA detection success and DNA fragment size could

be further used as a raw predictor of post feeding prey DNA detection intervals based

solely on prey DNA fragment length.

In general, our findings show that quantitative analyses of diet based on different

DNA fragment sizes or on different dietary samples are not directly comparable. Our

study suggests that for estimating and comparing consumption rates for the same

species between studies using different DNA fragment sizes or different dietary samples

(whole beetles/regurgitates vs feces), values should be corrected after taking into

account differences in detection probabilities (e.g. Greenstone *et al.* 2010).

Nevertheless, prey DNA detection depends on numerous additional factors including

species identity of the prey or the predator (Hosseini *et al.* 2008; Wallinger *et al.* 2013),

the feeding mode (Greenstone *et al.* 2007, 2013), the time since the last meal, the

number/size or the quality of prey consumed (Hoogendoorn & Heimpel 2001; Harper *et*

al. 2005; Eitzinger *et al.* 2014), which we did not investigate here. The next step

therefore would be the integration of multiple sources of variation in a complex

multispecies, multifactorial experimental design where the different sources of variation

could be quantified at once, and hierarchized (Welch *et al.* 2014).

Acknowledgements

We thank Paul Abrams for his invaluable help with early analyses of data presented

here. This study was founded by the French National Research Agency (Peerless

project ANR-12-AGRO-0006). S.K. was funded by the Atlantic Canada Opportunities

Agency. R.M. was supported by a grant of the regional government of Tyrol ("Land

Tirol") provided by the Mountain Agriculture Research Unit of the University of

Innsbruck. We also thank the three anonymous referees for their pertinent suggestions

which helped improve the present manuscript.

Data Accessibility

All the data used in this manuscript are included in the figure and the table presented

within the paper.

Author Contributions

MT, SK and MP designed the experiment. SK realized the field work and the feeding

experiment. RM carried out molecular analyses. ORR and EC realized data analyses.

SK wrote the manuscript with input from all the authors.

Conflicts of Interest: The authors have declared that no competing interests exist.

Bibliography

Agustí N, Shayler SP, Harwood JD, Vaughan IP, Sunderland KD, Symondson WOC (2003) Collembola as alternative prey sustaining spiders in arable ecosystems: prey detection within predators using molecular markers. *Molecular Ecology*, **12**, 3467e3475.

Cheeseman MT, Gillott C (1987) Organization of protein digestion in adult *Calosoma calidum* (Coleoptera: Carabidae). *Journal of Insect Physiology*, **33**, 1-8.

Clare EL (2015) Molecular detection of trophic interactions: Emerging trends, distinct advantages, significant considerations and conservation applications. *Evolutionary Applications*, **7**, 1144-1157.

Cohen AC (1995) Extra-oral digestion in predaceous terrestrial Arthropoda. *Annual review of entomology*, **40**, 85–103.

Deagle BE, Eveson JP, Jarman SN (2006) Quantification of damage in DNA recovered from highly degraded samples–a case study on DNA in faeces. *Frontiers in Zoology*, **3**, 1.

Delignette-Muller ML, Dutang C (2015) fitdistrplus: An R package for fitting distributions. *Journal of statistical software*, **4**, 1-34.

Durbin EG, Casas MC, Rynearson TA (2012) Copepod feeding and digestion rates using prey DNA and qPCR. *Journal of Plankton Research*, **34**, 72-82.

Egeter B, Bishop PJ, Robertson BC (2015) Detecting frogs as prey in the diets of introduced mammals: a comparison between morphological and DNA-based diet analyses. *Molecular Ecology Resources*, **15**, 306–316.

Eitzinger B, Unger EM, Traugott M, Scheu S (2014) Effects of prey quality and predator body size on prey DNA detection success in a centipede predator. *Molecular Ecology*, **23**, 3767-3776.

Folmer O, Black M, Hoeh W et al. (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology*, **3**, 294-299.

Greenstone MH, Payton ME, Weber DC, Simmons AM (2013) The detectability half-life in arthropod predator-prey research: what it is, why we need it, how to measure it, and how to use it. *Molecular Ecology*, **23**, 3799-3813.

Greenstone MH, Rowley DL, Weber DC, Payton ME, Hawthorne DJ (2007) Feeding mode and prey detectability half-lives in molecular gut-content analysis: an example

with two predators of the Colorado potato beetle. *Bulletin of Entomological Research*, **97**, 201–209.

Greenstone MH, Szendrei Z, Payton ME, Rowley DL, Coudron TC, Weber DC (2010) Choosing natural enemies for conservation biological control: use of the prey detectability half-life to rank key predators of Colorado potato beetles. *Entomologia Experimentalis et Applicata*, **136**, 97-107.

Harper GL, King RA, Dodd CS, Harwood JD, Glen DM et al. (2005) Rapid screening of invertebrate predators for multiple prey DNA targets. *Molecular Ecology*, **14**, 819-827.

Hoogendoorn M, Heimpel GE (2001) PCR-based gut content analysis of insect predators: using ribosomal ITS-1 fragments from prey to estimate predation frequency. *Molecular Ecology*, **10**, 2059-2067.

Hosseini R, Schmidt O, Keller MA (2008) Factors affecting detectability of prey DNA in the gut contents of invertebrate predators: a polymerase chain reaction-based method. *Entomologia Experimentalis et Applicata*, **126**, 194–202.

Ibanez S, Manneville O, Miquel C *et al.* (2013) Plant functional traits reveal the relative contribution of habitat and food preferences to the diet of grasshoppers. *Oecologia*, **173**, 1459–1470.

Kartzinel TR, Chen PA, Coverdale TC *et al.* (2015) DNA metabarcoding illuminates dietary niche partitioning by African large herbivores. *Proceedings of the National Academy of Sciences*, **112**, 8019–8024 (2015).

King RA, Read DS, Traugott M, Symondson WOC (2008) Molecular analysis of predation: a review of best practice for DNA-based approaches. *Molecular Ecology*, 17, 947-963.

Leray M, Meyer CP, Mills SC (2015) Metabarcoding dietary analysis of coral dwelling predatory fish demonstrates the minor contribution of coral mutualists to their highly partitioned, generalist diet. *PeerJ*, **3,** e1047.

Lövei GL, Sunderland KD (1996) Ecology and behavior of ground beetles (Coleoptera: Carabidae). *Annual review of entomology*, **41**, 231–256.

Mollot G, Duyck P-F, Lefeuvre P *et al.* (2014) Cover Cropping Alters the Diet of Arthropods in a Banana Plantation: A Metabarcoding Approach. *PLoS ONE*, **9**, e93740.

Oehm J, Juen A, Nagiller K et al. (2011) Molecular scatology: how to improve prey DNA detection success in avian faeces? Improving prey DNA detection in avian faeces. *Molecular Ecology Resources*, **11**, 620–628 (2011).

446
447 Pérez-Sayas, C, Pina T, Gómez-Martínez MA *et al.* (2015) Disentangling mite
448 predator-prey relationships by multiplex PCR. *Molecular Ecology Resources*, **15**,
449 1330–1345.
450
451 Piñol J, San Andrés V, Clare EL et al. (2014) A pragmatic approach to the analysis
452 of diets of generalist predators: the use of next-generation sequencing with no
453 blocking probes. *Molecular Ecology Resources*, **14**, 18–26.
454
455 Pompanon F, Deagle BE, Symondson WOC *et al.* (2012) Who is eating what: diet
456 assessment using next generation sequencing. *Molecular Ecology*, **21**, 1931–1950.
457
458 Redd KS, Ling SD, Frusher SD, Jarman S, Johnson CR (2014) Using molecular
459 prey detection to quantify rock lobster predation on barrens-forming sea urchins.
460 *Molecular Ecology*, **23**, 3849–3869.
461
462 Shehzad W, Riaz T, Nawaz MA *et al.* (2012) Carnivore diet analysis based on next-
463 generation sequencing: application to the leopard cat (*Prionailurus bengalensis*) in
464 Pakistan. *Molecular Ecology*, **21**, 1951–1965.
465
466 Sint D, Raso L, Kaufmann R et al. (2011) Optimizing methods for PCR-based
467 analysis of predation. *Molecular Ecology Resources*, **11**, 795–801.
468
469 Sint D, Thurner I, Kaufmann R, Traugott M (2015) Sparing spiders: Faeces as a
470 non-invasive source of DNA. *Frontiers in Zoology*, *12*, doi:10.1186/s12983-015-
471 0096-y
472
473 Staudacher K, Wallinger C, Schallhart N, Traugott M (2011) Detecting ingested plant
474 DNA in soil-living insect larvae. *Soil Biology & Biochemistry*, **43**, 346-350.
475
476 Symondson, W.O.C., 2012. The molecular revolution: using polymerase chain
477 reaction based methods to explore the role of predators in terrestrial food webs. In:
478 Gurr, G.M., Wratten, S.D., Snyder, W.E., Read, D.M.Y. (Eds.), Biodiversity and
479 Insect Pests: Key Issues for Sustainable Management. Wiley and Sons, pp. 166–
480 184.
481
482 Thalinger B, Oehm J, Mayr H *et al.* (2016) Molecular prey identification in Central
483 European piscivores. *Molecular Ecology Resources*, **16**, 123–137.
484
485 Tjur T (2009) Coefficients of Determination in Logistic Regression Models—A New
486 Proposal: The Coefficient of Discrimination. *The American statistician*, **63**, 366–372.
487
488 Traugott M, Kamenova S, Ruess L et al. (2013) Empirically characterising trophic
489 networks: what emerging DNA-based methods, stable isotope and fatty acid
490 analyses can offer. *Advances in Ecological Research*, **49**, 177.
491

492   Unruh TR, Miliczky ER, Horton DR *et al.* (2016) Gut content analysis of arthropod
493   predators of codling moth in Washington apple orchards. *Biological Control,*
494   doi:10.1016/j.biocontrol.2016.05.014
495

496   Valentini A, Miquel C, Muhammad AN et al. (2009) New perspectives in diet analysis
497   based on DNA barcoding and parallel pyrosequencing: the trnL approach. *Molecular*
498   *Ecology Resources*, **9**, 51-60.
499

500   Von Berg K, Traugott M, Symondson WOC (2008) The effects of temperature on
501   detection of prey DNA in two species of carabid beetle. *Bulletin of Entomological*
502   *Research* **98,** 263-269.
503

504   Waldner T, Sint D, Juen A et al. (2013) The effect of predator identity on post-
505   feeding prey DNA detection success in soil-dwelling macro-invertebrates. *Soil*
506   *Biology and Biochemistry*, **63**, 116–123.
507

508   Waldner T, Traugott M (2012) DNA-based analysis of regurgitates: a noninvasive
509   approach to examine the diet of invertebrate consumers. *Molecular Ecology*
510   *Resources*, **12**, 669–675.
511

512   Wallinger C, Sint D, Baier F et al. (2015) Detection of seed DNA in regurgitates of
513   granivorous carabid beetles. *Bulletin of Entomological Research*, **1**, 1-8.
514

515   Wallinger C, Staudacher K, Schallhart N, Peter E, Dresch P, Juen A, Traugott M
516   (2013). The effect of plant identity and the level of plant decay on molecular gut
517   content analysis in a herbivorous soil insect. *Molecular Ecology Resources*, **13**, 75–
518   83.
519

520   Welch KD, Schofield MR, Chapman EG, Harwood JD (2014) Comparing rates of
521   springtail predation by web-building spiders using Bayesian inference. *Molecular*
522   *Ecology*, **23**, 3814-3825.
523

Figure 1 Prey DNA detection success in the predatory carabid beetle *Pterostichus*
melanarius for regurgitates (A), whole bodies (B) and feces (C). Detection rates are
provided for the different time points examined within each dietary sample and for the
three target DNA fragment sizes. Circles and dashed lines indicate actual measures.
Bold solid lines indicate the logistic regressions estimated from the glmm model and the
shaded area the 95% confidence interval envelopes of the fit. The horizontal line
represents the 50% prey DNA detection probability. Corresponding lower and upper
95% confidence limits are presented in Table 2.
533
534 **Figure 2** Estimated time points post-feeding for a 50% prey DNA detection probability
535 for the different types of dietary samples and DNA fragment sizes. Provided are the
536 50% prey detection probabilities in hours post-feeding.
537

538 **Table 1** Detection rates of small (128 bp), medium (332 bp) and large (612 bp) prey
539 DNA fragments of the mealworm *Tenebrio molitor* fed to the carabid *Pterostichus*
540 *melanarius* in whole beetles, regurgitates, and feces. *N* is the number of samples
541 analyzed per digestion time.
542

Dietary sample	Digestion time (h)	n	Detection rate per fragment size (%)		
			Small (128 bp)	Medium (332 bp)	Large (612 bp)
Whole bodies	0	20	95	90	95
	12	10	100	100	60
	24	10	100	80	40
	36	9	78	44	22
	48	10	100	40	40
	60	10	80	20	0
	72	10	70	10	0
	96	9	56	0	0
Regurgitates	0	11	100	100	100
	12	10	100	100	80
	24	10	100	80	50
	36	10	100	30	30
	48	9	100	33	33
	60	10	70	10	0
	72	9	78	0	0
	96	9	78	22	11
Feces	3	8	100	87,5	100
	6	13	92	69	69
	9	5	100	100	60
	12	4	100	100	50
	15	7	100	43	57
	21	2	100	50	50
	24	4	100	25	0
	27	12	100	75	50
	30	2	100	50	50
	33	5	100	60	40
	39	9	100	22	11
	42	3	67	0	0
	45	2	100	0	0
	48	3	100	100	100
	51	4	100	0	25
	54	2	50	0	0
	57	1	100	100	0
	58	1	100	100	0
	60	1	0	0	0
	64	1	100	0	0
	70	3	67	0	0
	76	1	0	0	0