Potassium channels contribute to activity-dependent scaling of dendritic inhibition.

Jeremy T. Chang and Michael J. Higley

Department of Neuroscience

Program in Cellular Neuroscience, Neurodegeneration and Repair

Kavli Institute

Yale School of Medicine

New Haven, CT 06510

14   Correspondence: m.higley@yale.edu

Abstract

GABAergic inhibition plays a critical role in the regulation of neuronal activity. In the neocortex, inhibitory interneurons that target the dendrites of pyramidal cells influence both electrical and biochemical postsynaptic signaling. Voltage-gated ion channels strongly shape dendritic excitability and the integration of excitatory inputs, but their contribution to GABAergic signaling is less well understood. Here, we examine the actions of potassium channels in regulating dendritic inhibition in mouse visual cortex. By combining 2-photon calcium imaging and focal GABA uncaging, we show that A-type channels normally suppress the GABAergic inhibition of calcium signals evoked by back-propagating action potentials in dendritic spines and shafts. Moreover, the voltage-dependent inactivation of these channels leads to enhancement of dendritic inhibition following somatic spiking. Overall, our findings highlight the interaction between intrinsic and synaptic properties and reveal a novel mechanism for the activity-dependent scaling of GABAergic inhibition.

Introduction

Inhibition in the neocortex is primarily mediated by the neurotransmitter gamma-aminobutyric acid (GABA) through synaptic contacts made by interneurons. These synapses are distributed across the entire somatodendritic arbor and work to counteract excitatory glutamatergic input. GABAergic synapses that target the axon initial segment and soma exert a strong influence on somatic voltage, and consequently play important roles in regulating the generation and timing of action potentials (Higley and Contreras, 2006; Pouille and Scanziani, 2001; Wehr and Zador, 2003; Zhu et al., 2004). However, the vast majority of inhibitory inputs are formed onto pyramidal cell dendrites (Beaulieu and Somogyi, 1990), and the role of dendrite-targeting inhibition has been an area of growing interest (Bloss et al., 2016; Lovett-Barron et al., 2012; Miles et al., 1996). One important function of dendritic inhibition, in addition to action potential regulation, is the regulation of dendritic calcium signals which are thought to play an instructive role in synaptic plasticity (Chiu et al., 2013; Palmer et al., 2012; Tsubokawa and Ross, 1996). Recent reports in the neocortex and hippocampus have described varying efficacy of dendritic calcium inhibition, ranging from spatial compartmentalization within individual spines to complete abolition of actively propagating action potentials (Chiu et al., 2013; Kanemoto et al., 2011; Marlin and Carter, 2014; Mullner et al., 2015; Stokes et al., 2014). The mechanisms underlying this heterogeneity are unclear, but one contributing factor may be variations in intrinsic dendritic properties, like voltage-dependent channels, whose impact on GABAergic inhibition is not well understood.

The expression of voltage-gated ion channels within neuronal dendrites regulates cellular excitability and strongly influences synaptic integration (Cook and Johnston, 1999; Johnston and Narayanan, 2008; Miller et al., 1985; Poirazi and Mel, 2001; Shepherd et al., 1985). One example is the A-type potassium (KA) conductance, characterized by fast inactivation and sensitivity to 4-aminopyridine (4-AP)(Bekkers, 2000; Clark et al., 2008; Korngreen and Sakmann, 2000; Serodio and Rudy, 1998). Within cortical and hippocampal pyramidal neurons, KA channels are expressed throughout the dendritic arbor and have been implicated in the regulation of back-propagating action potentials (bAPs) and excitatory synaptic integration (Cai et al., 2004; Carrasquillo et al., 2012; Gasparini, 2011; Harnett et al., 2013; Hoffman et al., 1997; Ramakers and Storm, 2002). Despite clearly influencing dendritic excitation, KA channels have been shown to preferentially co-localize with GABAergic synapses, suggesting they may also play a role in the control of inhibition (Burkhalter et al., 2006; Jinno et al., 2005).

Here, we examine how voltage-gated potassium channels alter GABAergic inhibition of bAP-evoked Ca2+ signals (ΔCa2+) in dendrites of L5 pyramidal neurons in mouse visual cortex. We show that the blockade of these channels specifically in the apical dendrite enhances both the amplitude and inhibition of bAP-evoked ΔCa2+. Moreover, our data suggest the involvement of dendritic Kv4 channels, previously shown to play a role in dendritic excitability. We also show that the voltage-dependent inactivation of these channels gives rise to a scaling of dendritic GABAergic inhibition, such that inhibitory efficacy is enhanced following strong somatic activity. Thus, our findings demonstrate that intrinsic excitability interacts with GABAergic synaptic input to dynamically regulate dendritic Ca2+ signaling.

Results

In order to investigate the impact of potassium channels on dendritic inhibition, we performed

two-photon calcium imaging of bAP-evoked dendritic Ca2+ transients in layer 5 pyramidal neurons

(L5PNs) of mouse visual cortex (Figure 1A). Ca2+ signals were measured in dendritic spines and

neighboring shafts along the primary apical dendrite, 100-150 μm from the soma (Figure 1B). To probe

the effects of GABAergic inhibition on ΔCa2+, we compared uninhibited ΔCa2+ from bAPs induced by

somatic current injection with ΔCa2+ from bAPs preceded (15 ms) by local (at the imaging site) uncaging

of RuBi-GABA (Rial Verde et al., 2008). To compare observations across different recordings, GABAergic

inhibition of ΔCa2+ was quantified as in previous studies as (ΔCa2+$_{Ctl}$-ΔCa2+$_{Inh}$)/ΔCa2+$_{Ctl}$ (Chiu et al.,

2013). The magnitude of this Ca2+ inhibition was measured before and after bath-application of the

potassium channel blocker 4-aminopyridine (4-AP)(Figure 1A). Treatment with 4-AP broadened the

somatic action potential (p=0.0020)(Figure 1C) and increased the average peak ΔCa2+ evoked by a

single bAP for both spines (p=0.0059) and neighboring shafts (p=0.0137) (Figure 1D-G). Moreover, 4-AP

significantly enhanced the average GABAergic inhibition of ΔCa2+ for both spines (p=0.0195) and

neighboring shafts (p=0.0098) (Figure 1D-G). Importantly, this result was not seen when slices were pre-

treated with the GABA$_A$R antagonist picrotoxin (Figure 1 - figure supplement 1) nor with application of

control saline, which demonstrated the presence of a run-down in peak ΔCa2+ over time without change

in Ca2+ inhibition (Figure 1 - figure supplement 2), consistent with previous findings (Canepari et al.,

2007; Chalifoux and Carter, 2011; Fukumoto et al., 2012; Kameyama et al., 1989).

Given the enhanced GABAergic inhibition of dendritic ΔCa2+ following 4-AP treatment, we next

asked whether 4-AP also altered the somatic inhibitory postsynaptic potential (IPSP) evoked by

GABAergic dendritic input. We analyzed IPSPs evoked by either single uncaging events or bursts of three

uncaging events (10 Hz), directed to the proximal apical dendrite as above. In the presence of 4-AP,

IPSP amplitude was enhanced for both single (p=0.0391) and triple (p=0.0039) uncaging protocols

(Figure 1— figure supplement 3A-C). Concurrent with the increase in IPSP amplitude, the decay of the

IPSP was also significantly slowed after treatment with 4-AP (p=0.0039), leading to enhanced temporal

summation for bursts of IPSPs (Figure 1—figure supplement 3C-D). The IPSP alterations could be

explained by a significant increase in somatic input resistance measured using a series of negative

current steps (p=0.0391)(Figure 1—figure supplement 3E). In sum, these data suggest voltage-

dependent potassium channels normally suppress the somatic and dendritic effects of GABAergic

inhibition in the apical dendrite.

Next, we asked whether the actions of 4-AP-sensitive channels on inhibition of dendritic ΔCa2+

generalized to other subcellular regions, as previous reports have suggested differential contributions to

excitability in basal and apical dendrites (Sandler et al., 2016). We performed similar imaging experiments

in basal dendrites (100-150 µm from the soma) (Figure 1—figure supplement 4A-D). Again, the bAP-

evoked ΔCa2+ was significantly increased by 4-AP for both spines (p= 0.0026) and neighboring shafts

(p= 0.0020). However, GABAergic inhibition of $\Delta Ca2+$ was much stronger in the basal dendrites and was

not further enhanced by application of 4-AP for either spines (p= 0.1602) or neighboring shafts (p=

0.2754) (Figure 1—figure supplement 3B-D). Note that while the fractional inhibition of $\Delta Ca2+$ was not

altered, the absolute inhibition of $\Delta Ca2+$ influx was enhanced by 4-AP. Thus, our results suggest that

potassium channels differentially shape GABAergic efficacy in apical versus basal dendrites.

As bath application of 4-AP altered both somatic and apical dendritic effects of GABAergic

inhibition, we investigated whether these actions arise from similar or distinct pools of potassium

channels. We used a puffer pipette to locally apply 4-AP at different locations along the somatodendritic

axis. When applied to the proximal apical dendrite (at the site of GABA uncaging), 4-AP replicated the

effects of bath-application on the magnitude of bAP-evoked $\Delta Ca2+$ and its inhibition by GABA.

Specifically, 4-AP increased $\Delta Ca2+$ in spines (p=0.0156) and neighboring shafts (p=0.0313) (Figure 2A-

D). GABAergic inhibition of $\Delta Ca2+$ was also enhanced in spines (p=0.0469) and neighboring shafts

(p=0.0313) (Figure 2A-D). In contrast, application of 4-AP to either the cell body or apical tuft had no

impact on peak $\Delta Ca2+$ or its inhibition by GABA within the proximal apical dendrite (Figure 2— figure

supplement 1A-G). Conversely, the effects of bath-applied 4-AP on somatic IPSPs evoked by dendritic

GABA uncaging were mimicked by focal application to the soma but not the proximal apical dendrite

(Figure 2— figure supplement 2). Thus, our data suggest that the somatic and dendritic impacts of

dendritic GABAergic inhibition are modulated by two spatially separated pools of potassium channels.

While 4-AP-sensitive potassium currents can be mediated by a variety of molecularly-defined

channels, A-type Kv4.2 channels have been specifically linked to the control of dendritic excitability

(Carrasquillo et al., 2012; Harnett et al., 2013; Losonczy et al., 2008; Sandler et al., 2016). We therefore

investigated the contribution of these channels to the regulation of GABAergic inhibition using the

scorpion toxin AmmTx3, which specifically blocks Kv4.2 and Kv4.3 channels in the presence of the

auxiliary subunit dipeptidyl-peptidaselike-protein 6 (DPP6) (Maffie et al., 2013). Similar to our results with

4-AP, GABAergic inhibition of bAP-evoked $\Delta Ca2+$ was significantly enhanced following bath application

of 200 nM AmmTx3 for both spines (p=0.0195) and neighboring shafts (p=0.0488) (Figure 3A-D).

Following AmmTx3 application, there was no change in the amplitude of the bAP-evoked $\Delta Ca2+$ in spines

(p=0.3223) and $\Delta Ca2+$ was significantly reduced in dendritic shafts (p=0.0137) relative to baseline (Figure

3B-D). However, the observation that $\Delta Ca2+$ exhibits considerable run-down over time (see Figure 1 -

figure supplement 2) suggests that AmmTx3 likely produces a small enhancement of bAP-evoked $\Delta Ca2+$

relative to control conditions that is masked by the run-down. Consistent with previous studies (Pathak et

al., 2016), AmmTx3 had no effect on either somatic input resistance or IPSP amplitude (Figure 3— figure

supplement 1).

As the effects of AmmTx3 application did not completely recapture those of 4-AP, we quantified

the contribution of Kv4 channels to L5PN membrane currents by recording from somatic outside-out

macropatches. Application of 4-AP blocked a fast outward current by -73.5 ± 6.8%. In contrast,

application of AmmTx3 produced a significantly smaller -43.1 ± 11.2% change in current (p=0.0417,

unpaired t-test), demonstrating that Kv4 channels sensitive to AmmTx3 constitute a fraction of the

voltage-gated potassium channels blocked by 4-AP in L5PNs (Figure 3 - figure supplement 2). Altogether,

these results suggest that the activity of A-type Kv4 channels is sufficient to influence dendritic Ca2+

signaling and GABAergic inhibition.

One notable feature of KA channels is their fast, voltage-dependent inactivation, which limits their

conductance during periods of high neuronal activity (Bekkers, 2000; Kim et al., 2005). We therefore

asked whether KA channels might enable dendritic GABAergic inhibition to dynamically scale with

somatic firing. To test this hypothesis, we compared GABAergic inhibition of ΔCa2+ evoked by a bAP

alone or preceded 20 ms by a train of 5 bAPs at 100 Hz (Figure 4A). Similar to 4-AP, the preceding train

significantly enhanced the peak ΔCa2+ for both spines (p= 0.002) and neighboring shafts (p= 0.002) and

also enhanced GABAergic inhibition of ΔCa2+ for spines (p=0.0059) and neighboring shafts (p= 0.002)

(Figure 4B-D). In contrast, preceding spike trains at 50 Hz had minimal effect on either ΔCa2+ amplitude

or GABAergic inhibition (Figure 4— figure supplement1A-D). Importantly, the ability of the 100 Hz train to

enhance dendritic inhibition was occluded by prior bath application of 4-AP, suggesting that KA channels

are required for activity-dependent scaling (Figure 4— figure supplement 1E-H).

We next asked whether activity-dependent scaling of inhibition could be seen with synaptic GABA

release. To test this, we expressed channelrhodopsin-2 (ChR2) in a subset of dendrite-targeting cortical

interneurons expressing somatostatin (SOM-INs)(Figure 5A). Brief pulses of blue light were used to

activate SOM-INs and produce postsynaptic IPSPs. We repeated experiments comparing the GABAergic

inhibition of ΔCa2+ evoked by a bAP alone or preceded by a 100 Hz train. As with GABA uncaging, trains

of somatic action potentials significantly enhanced the magnitude of ΔCa2+ in spines (p=0.0098) and

neighboring shafts p=0.0059) and led to stronger GABAergic inhibition of ΔCa2+ for both spines

(p=0.0273) and neighboring shafts (p=0.0273) (Figure 5B-D). Taken together, these results demonstrate

that KA channels in the apical dendrite play a key role in reducing the impact of synaptic GABAergic

inhibition on bAP-evoked ΔCa2+.

Finally, to examine the biophysical mechanisms underlying the interaction of dendritic potassium

channels with GABAergic signaling, we simulated an active dendritic compartment and tested the impact

of varying voltage-gated conductances on the magnitude of Ca2+ inhibition (Figure 6). We first explored

whether increased GABAergic inhibition was mediated by 4-AP-induced enhancement of dendritic

depolarization and the resulting increase in chloride driving force. The model revealed the surprising

finding that increasing bAP-dependent depolarization by increasing the sodium reversal potential actually

reduced GABAergic inhibition (Figure 6A-B). In contrast, reducing KA channel conductance recapitulated

our data, increasing both bAP amplitude and the amount of Ca2+ current inhibited by GABAergic

signaling (Figure 6A-C). Thus, alterations in peak depolarization attained during the bAP alone do not

account for the enhanced Ca2+ inhibition (Figure 6D). Instead, we found that manipulations that

increased the bAP duration (e.g., decreasing either E_{Na} or g_A) led to increased Ca2+ inhibition (Figure

6E). This phenomenon is illustrated in Figure 6A, where similar inhibition of the bAP peak leads to greater

reduction in the integrated Ca2+ current (yellow shaded areas) for broader bAP waveforms. In summary,

these observations strongly suggest that the contribution of potassium conductance to bAP duration is an

important factor in the magnitude of Ca2+ inhibition.

Discussion

Rapidly activating and inactivating A-type potassium channels are widely recognized as key

modulators of neuronal excitability as well as synaptic integration and plasticity (Carrasquillo et al., 2012;

Foeger et al., 2012; Harnett et al., 2013; Hoffman et al., 1997; Losonczy et al., 2008; Magee and Carruth,

1999). In the present work, we have described a role for KA channels in the regulation of dendritic

GABAergic inhibition and its control over dendritic Ca2+ signaling. Using a combination of

electrophysiology, 2-photon Ca2+ imaging, and focal GABA uncaging, we show that blocking KA

channels either pharmacologically or via activity-dependent inactivation enhances both bAP-evoked Ca2+

influx and GABAergic inhibition of these transients in the apical dendrites of L5PNs. Moreover, this

modulation of dendritic signaling can occur in the absence of changes to the somatically recorded IPSP.

These results provide evidence that KA channels compartmentalize their regulation of GABAergic

signaling.

Dendritic potassium channels comprise a diverse molecular group, including both Kv1-, Kv3, and

Kv4-type channels (Carrasquillo et al., 2012; Serodio et al., 1994). Our pharmacological analysis

indicates a contribution from Kv4 channels, previously shown to be expressed in both cortical and

hippocampal pyramidal dendrites. Indeed the ability of AmmTx3 to replicate the actions of 4-AP on

GABAergic Ca2+ inhibition suggests that Kv4 channels in L5PNs include the auxiliary subunit dipeptidyl-

peptidase-like-protein 6 (DPP6), in agreement with previous studies (Foeger et al., 2012; Pathak et al.,

2016). Nevertheless, AmmTx3 produces a smaller change in inhibition compared with 4-AP, suggesting

that other voltage-gated potassium channels also play a role in regulating dendritic GABAergic signaling.

This conclusion is supported by our data from outside-out somatic patches. Notably, the expression of

Kv4.2 channels is regulated by the activity of NMDA-type glutamate receptors, indicating the possibility

for multiple feedback loops within pyramidal neuron dendrites (Kim et al., 2007; Sandler et al., 2016).

Several previous studies have implicated KA channels in the regulation of both dendritic

excitability and glutamatergic synaptic integration. In both CA1 and cortical pyramidal neurons, the

presence of KA channels limits the spread of voltage between distinct compartments, such as the distal

and proximal apical dendrite, regulating both the back propagation of action potentials and the spread of

synaptically evoked dendritic spikes (Cai et al., 2004; Frick et al., 2003; Harnett et al., 2013; Kim et al.,

2007; Kim et al., 2005; Losonczy et al., 2008). Our study demonstrates that KA channels similarly restrict

the efficacy of GABAergic inhibition, suggesting that KA channels generally serve as dendritic "shock

absorbers", limiting the impact of synaptic inputs from all sources (Yuste, 1997). An intriguing possibility is

that voltage-gated potassium channels asymmetrically regulate excitation and inhibition, potentially

leading to moment-to-moment alterations of the balance between these opposing drives. Disruption of the

normal balance of excitatory and inhibitory signaling is thought to occur in a number of neuropsychiatric

disorders, including autism and schizophrenia. Indeed, in Fragile X syndrome, loss of FMRP produces a

dysregulation of KA channels that could lead to an imbalance of excitation and inhibition (Brager and

Johnston, 2014; Gross et al., 2011; Kalmbach et al., 2015). Future studies are necessary to elucidate the

precise role of KA channels in shaping glutamatergic and GABAergic integration.

In addition to regulating dendritic excitability, KA channels have been implicated in shaping long-

term plasticity of glutamatergic synapses. In particular, studies have focused on spike-timing dependent

plasticity (STDP), where bAPs can potentiate or depress synaptic inputs depending on their relative

timing to synaptic activity (Magee and Johnston, 1997; Markram et al., 1997). For example, EPSPs in

CA1 pyramidal neuron dendrites can inactivate KA channels, enhancing the dendritic invasion of somatic

action potentials and subsequent plasticity (Hoffman et al., 1997). Recent experimental and

computational studies have also suggested a key role for GABAergic inhibition in spike-timing dependent

plasticity (Cichon and Gan, 2015; Hayama et al., 2013; Paille et al., 2013; Wilmes et al., 2016). For

example, focal activation of GABAergic synapses in CA1 dendrites was shown to convert long-term

potentiation to depression due to negative regulation of dendritic Ca2+ influx (Hayama et al., 2013).

Together, these various findings suggest that the interaction of NMDARs, KA channels, and GABAergic

inhibition may strongly contribute to the development and maintenance of cortical circuits.

It is intriguing to speculate that expression patterns of KA channels may also explain some of the

recent diversity in studies examining GABAergic control of dendritic ΔCa2+. Previous work from our lab

showed that inhibition could be highly compartmentalized in layer 2/3 pyramidal neurons, with

neighboring spines exhibiting markedly different amounts of inhibition (Chiu et al., 2013). In contrast, work

from other groups has shown that more broad dendritic inhibition can occur in L5PNs and hippocampal

CA1 pyramidal neurons (Marlin and Carter, 2014; Mullner et al., 2015; Stokes et al., 2014). Differential

expression and recruitment of voltage-gated conductances, such as KA channels, would be expected to

contribute to the heterogeneity of inhibitory function across cell types.

Our computational modeling provides additional insight into the biophysical mechanism

underlying the interaction of KA channels and GABAergic inhibition. We found that decreasing the

potassium conductance, rather than increasing the bAP amplitude per se, was necessary to enhance

dendritic Ca2+ inhibition. This phenomenon is directly due to the broadening of the bAP duration and

subsequent reduction in the integral Ca2+ current (see Figure 6A). This finding suggests that a major

contributor to the efficacy of GABAergic regulation of action potentials is the relationship between bAP

duration and Ca2+ current. In addition, our computational modeling suggests that the enhancement of

GABAergic inhibition is a general biophysical feature of channels that broaden action potentials. AP width

has been linked to the coupling of AP into dendrites and dendritic location (Golding et al., 2001; Stuart et

al., 1997). Furthermore KA channels have been linked to both activity dependent as well as cell type

dependent alteration of action potential repolarization (Kim et al., 2005; Pathak et al., 2016). Our model

and experimental findings together thus predict that dendritic channels that regulate bAP shape are likely

to be important factors in altering the dendritic influence of GABAergic synapses, driving cell type- and subcellular location-dependent differences in the efficacy of GABAergic inhibition.

Finally, we found that the voltage-dependent inactivation of KA channels allows for the enhancement of GABAergic inhibition in the presence of high frequency somatic spike generation. This suggests that dendritic inhibition may exert greater control over Ca2+ signaling during periods of high network activity or somatic depolarization, essentially acting as a source of homeostatic control. These findings are consistent with previous experimental and computational studies demonstrating the activity-dependent amplification of trains of bAPs and bAP width in L5PN apical dendrites (Grewe et al., 2010; Kim et al., 2005; Larkum et al., 1999). Our results suggest that the dynamic properties of active dendritic conductances such as KA channels enable the alteration of GABAergic inhibition over short millisecond time frames, providing the basis for a context-dependent, flexible role of GABAergic signaling in shaping biochemical and electrical signaling in dendrites.

Materials and Methods
Slice Preparation

All animal handling was performed in accordance with guidelines approved by the Yale Institutional Animal Care and Use Committee and federal guidelines. For GABA uncaging experiments, subjects were male wild-type C57-BL6 mice, ages P30-40 (Harlan). For optogenetic experiments, subjects were male and female SOM-Cre mice, ages P30-40 (IMSR Cat# JAX:013044, RRID:IMSR_JAX:013044). Under isofluorane anesthesia, mice were decapitated and coronal slices (300 μm thick) containing primary visual cortex were cut in ice cold external solution containing (in mM): 110 choline, 25 NaHCO3, 1.25 NaH2PO4, 2.5 KCl, 7 MgCl2, 0.5 CaCl2, 20 glucose, 11.6 sodium ascorbate, and 3.1 sodium pyruvate, bubbled with 95% O2 and 5% CO2. After an incubation period of 20 minutes at 34°C, slices were transferred to artificial cerebrospinal fluid (ACSF) containing in (mM): 127 NaCl, 25 NaHCO3, 1.25 NaH2PO4, 2.5 KCl, 1 MgCl2, 2 CaCl2, and 20 glucose bubbled with 95% O2 and 5% CO2 and maintained at room temperature (20-22°C) for at least 20 min until use.

Electrophysiology and imaging

Experiments were conducted at room temperature in a submersion type recording chamber. Whole-cell patch clamp recordings were obtained from layer 5 pyramidal neurons (500 µm to 600 µm from the pial surface) identified with video infrared-differential interference contrast. For current-clamp recordings, glass electrodes (2-4 MΩ tip resistance) were filled with internal solution containing (in mM): 135 KMeSO3, 10 HEPES, 4 MGCl2, 4 Na2ATP, 0.5 NaGTP, and 10 sodium creatine phosphate, adjusted to pH 7.3 with KOH. For Ca2+ imaging experiments, red fluorescent Alexa Fluor-568 (40 μM) and green fluorescent Ca2+-sensitive Fluo-5F (300 μM) were included in the pipette solution to visualize cell morphology and changes of intracellular Ca2+ concentration, respectively. In experiments where only

somatic IPSPs or outside-out patches were recorded, Alexa Fluor-568 (40 μM) and EGTA (100 μM) were

included in the internal solution. Electrophysiological recordings were made using a Multiclamp 700B

amplifier (Molecular Devices), filtered at 4 kHz, and digitized at 10 kHz. For all recordings, membrane

potential was adjusted to -64 mV using current injection through the pipette. Steady state somatic chord

input resistance was determined by linearly fitting the amplitude of response to 600 ms current injections

of -300, -200, and -100 pA.

Outside-out macropatch data was obtained in voltage clamp configuration. Patches were formed

after whole-cell configuration was obtained by withdrawing the pipet without additional suction. Holding

potential was maintained at -75 mV. After a 1 s voltage step to -90 mV, voltage dependent membrane

currents were recorded in response to a 500 ms voltage step to +35 mV. Passive leak was removed

using a P/N subtraction method of the response from -110 mV to -140 mV. A-type currents were

quantified over a 50 ms window beginning 5 ms after the depolarizing voltage step to avoid stimulus

artifact.

Two-photon imaging was performed with a custom-modified Olympus BX51-WI microscope,

including components manufactured by Mike's Machine Company. Fluorophores were excited using 840

nm light from a pulsed titanium-sapphire laser. Emissions were separated using appropriate optical filters

(Chroma, Semrock) and collected by photomultiplier tubes (Hamamatsu). A mechanical shutter was

placed in front of the collectors to prevent damage during blue light stimulation. For Ca2+ imaging, signals

were collected during a 500 Hz line scan across a spine and neighboring dendritic shaft on the main

apical trunk 100 μm to 150 μm from the cell body. Back-propagating action potentials (bAPs) were

evoked using a brief depolarizing current pulse (0.5 ms, 1.5-2.5 nA) through the recording pipette. Trials

including bAP alone, IPSP-bAP, and IPSP alone were interleaved with a 45 second inter-trial interval. In a

subset of experiments, trains of action potentials at 50 Hz and 100 Hz were elicited by current pulse

injections through the recording pipette, ending 20 ms prior to a single current pulse. In this case, trials

including single bAP alone, train-bAP alone, IPSP-bAP, train-IPSP-bAP, IPSP alone, train-IPSP, and train

alone were interleaved with a 45 ms inter-trial interval. Fluorescent traces were computed for individual

cells as the average of 10 trials.

Reference frame scans were taken between each acquisition to correct for small spatial drift over

time. Ca2+ signals were first quantified as changes in green fluorescence from baseline normalized to the

average red fluorescence (ΔG/R). To permit comparison of the imaging data across various microscope

configurations, we expressed fluorescence changes as the fraction of the G/R ratio measured in

saturating Ca2+ (ΔG/Gsat).

Data acquisition and analysis

Imaging and physiology data were acquired using custom software written in MATLAB. Off-line

analysis was performed using custom routines written in MATLAB (MATLAB, RRID:SCR_001622) and

IgorPro (Wavemetrics Software, RRID:SCR_000325). Ca2+ responses were calculated as the integral of

the fluorescence transient over the first 100 ms after bAP initiation. In order to enable comparisons across cells, Ca2+ inhibition was expressed as in previous studies (Chiu et al., 2013) as ($\Delta Ca2+_{Ctl}$ - $\Delta Ca2+_{Inh}$)/$\Delta Ca2+_{Ctl}$. IPSPs were computed as the average of 10 trials, and amplitudes were calculated by finding the peak of the voltage trace and averaging values across a surrounding 3 ms window. IPSP decay kinetics were calculated by fitting an exponential to the falling phase of the IPSP. All statistical comparisons were made using the non-parametric Wilcoxon matched pairs signed rank test in GraphPad Prism version 7.01 (GraphPad Prism, RRID:SCR_002798) unless otherwise noted.

Pharmacology

For all GABA uncaging experiments, ACSF included 3 μM CGP-55845 hydrochloride (Tocris Cat. No. 1248) to block GABAB receptors, 10 μM (R)-CPP (Tocris Cat. No. 0247) to block NMDA receptors, and 10 μM NBQX disodium salt (Tocris Cat. No. 1044) to block AMPA receptors. For a subset of experiments, the ACSF included 5 mM 4-aminopyradine (Tocris Cat. No. 0940), 200 nM AmmTx3 (Smartox Biotechnology), or 100 μM picrotoxin (Tocris Cat. No 1128). Outside-out potassium currents were recorded in the presence of 1 μM tetrodotoxin (Tocris Ca. No 1078). Local application of 25 mM 4-AP was achieved using a glass puffer pipette (< 2 μm tip) coupled to a Picospritzer. Drugs were ejected continuously with 10-17 psi, and pipettes were position 30-70 μm from the targeted structure at the surface of the slice. In experiments where one-photon uncaging was performed with local drug application, 10.8 μM RuBi-GABA was included in the puffer pipette. In a subset of cells, somatic current injections elicited bursts of action potentials in the presence of 4-AP and were excluded from subsequent analysis.

Visible light-evoked GABA uncaging was accomplished using RuBi-GABA (10.8 μM) bath-applied in the ACSF(Rial Verde et al., 2008). We overfilled the back aperture of the microscope objective (60x, 1.0 NA) with collimated blue light from a fiber-coupled 473 nm laser. Spherical aberrations due to fiber-coupling resulted in a 15-20 μm diameter disc of light at the focal plane centered on the field of view. A brief (0.5 ms) pulse of light (1-2 mW at the sample) reliably evoked uncaging-evoked IPSPs. For Ca2+ imaging experiments, a blue light photo-artifact was corrected by subtracting fluorescence traces on uncaging-alone trials from those with Ca2+ imaging. For all experiments, GABA uncaging occurred 15 ms prior to bAP initiation.

ChR2 Expression and Activation

To stimulate SOM-INs, SOM-Cre mice were injected 13-23 days prior to slice preparation into the primary visual cortex with recombinant adeno-associated virus (AAV) driving conditional expression of a ChR2-eYFP fusion protein under the Ef1a-promoter (AAV-DIO-Ef1a-ChR2-EYFP)(UNC Vector Core). Optogenetic stimulation was accomplished using the same light source and path as one-photon GABA uncaging (see above). Brief (2-3 mW, 0.5 ms) pulses were used to stimulate SOM-INs 15 ms prior to bAP initiation.

NEURON Modeling

Multi-compartment time-dependent simulations were run using NEURON v7.4 (NEURON,
RRID:SCR_005393, available free at http://neuron.med.yale.edu) and analyzed using custom scripts
written in Jupyter Notebooks 4.1.0 using Python 3.5.2. We modified a previously published ball and stick
model (Chiu et al., 2013) adding compartments for a basal dendrite and two apical tuft dendrites. A single
dendritic spine (1 μm diameter) was attached to the apical dendrite 122.5 μm from the cell body by a neck
(1 μm length, 0.07 μm diameter). A medium voltage-gated calcium channel was inserted into the dendritic
spine and neighboring dendrite such that currents through these channels would minimally impact
membrane potential (Almog and Korngreen, 2014). GABAergic synapses were modeled as an
exponential synapse contacting the dendritic shaft located 123.5 μm from the cell body. We modeled KA
currents using a previously published channel definition that fits observed A-type potassium currents in
distal dendrites and altered the expression pattern of this channel and sodium currents in the apical
dendrite (Acker and Antic, 2009; Migliore et al., 1999). In order to reproduce our experimental conditions,
an iterative search was conducted to find a somatic current injection that maintained the somatic resting
potential at 64.00 +/- 0.0001 mV at the cell body for each condition tested. Similar to our experiments we
quantified calcium flux over a 100 ms window in order to calculate percent calcium change due to
inhibition. To speed up simulation time, simulations were run in parallel using the built-in message
passing interface of NEURON.

Acknowledgements
387   The authors thank Drs. Jess Cardin, Susumu Tomita, and Michael Crair as well as members of the Higley
388   lab for constructive comments during the preparation of this manuscript. Experiments were supported by
389   funding from the March of Dimes (Basil O'Connor Award) and the NIH (R01 MH099045, T32 EY22312).

References

Acker, C.D., and Antic, S.D. (2009). Quantitative assessment of the distributions of membrane conductances involved in action potential backpropagation along basal dendrites. J Neurophysiol *101*, 1524-1541. doi:10.1152/jn.00651.2007

Almog, M., and Korngreen, A. (2014). A quantitative description of dendritic conductances and its application to dendritic excitation in layer 5 pyramidal neurons. J Neurosci *34*, 182-196. doi:10.1523/JNEUROSCI.2896-13.2014

Beaulieu, C., and Somogyi, P. (1990). Targets and Quantitative Distribution of GABAergic Synapses in the Visual Cortex of the Cat. Eur J Neurosci *2*, 896. doi:10.1111/j.1460-9568.1990.tb00421.x

Bekkers, J.M. (2000). Distribution and activation of voltage-gated potassium channels in cell-attached and outside-out patches from large layer 5 cortical pyramidal neurons of the rat. J Physiol *525 Pt 3*, 611-620. doi:10.1111/j.1469-7793.2000.t01-2-00611.x

Bloss, E.B., Cembrowski, M.S., Karsh, B., Colonell, J., Fetter, R.D., and Spruston, N. (2016). Structured Dendritic Inhibition Supports Branch-Selective Integration in CA1 Pyramidal Cells. Neuron *89*, 1016-1030. doi:10.1016/j.neuron.2016.01.029

Brager, D.H., and Johnston, D. (2014). Channelopathies and dendritic dysfunction in fragile X syndrome. Brain Res Bull *103*, 11-17. doi:10.1016/j.brainresbull.2014.01.002

Burkhalter, A., Gonchar, Y., Mellor, R.L., and Nerbonne, J.M. (2006). Differential expression of I(A) channel subunits Kv4.2 and Kv4.3 in mouse visual cortical neurons and synapses. J Neurosci *26*, 12274-12282. doi:10.1523/JNEUROSCI.2599-06.2006

Cai, X., Liang, C.W., Muralidharan, S., Kao, J.P., Tang, C.M., and Thompson, S.M. (2004). Unique roles of SK and Kv4.2 potassium channels in dendritic integration. Neuron *44*, 351-364. doi:10.1016/j.neuron.2004.09.026

Canepari, M., Djurisic, M., and Zecevic, D. (2007). Dendritic signals from rat hippocampal CA1 pyramidal neurons during coincident pre- and post-synaptic activity: a combined voltage- and calcium-imaging study. J Physiol *580*, 463-484. doi:10.1113/jphysiol.2006.125005

Carrasquillo, Y., Burkhalter, A., and Nerbonne, J.M. (2012). A-type K+ channels encoded by Kv4.2, Kv4.3 and Kv1.4 differentially regulate intrinsic excitability of cortical pyramidal neurons. J Physiol *590*, 3877-3890. doi:10.1113/jphysiol.2012.229013

Chalifoux, J.R., and Carter, A.G. (2011). GABAB receptor modulation of voltage-sensitive calcium channels in spines and dendrites. J Neurosci *31*, 4221-4232. doi:10.1523/JNEUROSCI.4561-10.2011

Chiu, C.Q., Lur, G., Morse, T.M., Carnevale, N.T., Ellis-Davies, G.C., and Higley, M.J. (2013). Compartmentalization of GABAergic inhibition by dendritic spines. Science *340*, 759-762. doi:10.1126/science.1234274

Cichon, J., and Gan, W.B. (2015). Branch-specific dendritic Ca(2+) spikes cause persistent synaptic plasticity. Nature *520*, 180-185. doi:10.1038/nature14251

Clark, B.D., Kwon, E., Maffie, J., Jeong, H.Y., Nadal, M., Strop, P., and Rudy, B. (2008). DPP6 Localization in Brain Supports Function as a Kv4 Channel Associated Protein. Front Mol Neurosci *1*, 8. doi:10.3389/neuro.02.008.2008

Cook, E.P., and Johnston, D. (1999). Voltage-dependent properties of dendrites that eliminate location-dependent variability of synaptic input. J Neurophysiol *81*, 535-543.

Foeger, N.C., Norris, A.J., Wren, L.M., and Nerbonne, J.M. (2012). Augmentation of Kv4.2-encoded currents by accessory dipeptidyl peptidase 6 and 10 subunits reflects selective cell surface Kv4.2 protein stabilization. J Biol Chem *287*, 9640-9650. doi:10.1074/jbc.M111.324574

Frick, A., Magee, J., Koester, H.J., Migliore, M., and Johnston, D. (2003). Normalization of Ca2+ signals by small oblique dendrites of CA1 pyramidal neurons. J Neurosci *23*, 3243-3250.

Fukumoto, N., Kitamura, N., Niimi, K., Takahashi, E., Itakura, C., and Shibuya, I. (2012). Ca2+ channel currents in dorsal root ganglion neurons of P/Q-type voltage-gated Ca2+ channel mutant mouse, rolling mouse Nagoya. Neurosci Res *73*, 199-206. doi:10.1016/j.neures.2012.04.006

Gasparini, S. (2011). Distance- and activity-dependent modulation of spike back-propagation in layer V pyramidal neurons of the medial entorhinal cortex. J Neurophysiol *105*, 1372-1379. doi:10.1152/jn.00014.2010

Golding, N.L., Kath, W.L., and Spruston, N. (2001). Dichotomy of action-potential backpropagation in CA1 pyramidal neuron dendrites. J Neurophysiol *86*, 2998-3010.

Grewe, B.F., Bonnan, A., and Frick, A. (2010). Back-Propagation of Physiological Action Potential Output
in Dendrites of Slender-Tufted L5A Pyramidal Neurons. Front Cell Neurosci *4*, 13.
doi:10.3389/fncel.2010.00013
Gross, C., Yao, X., Pong, D.L., Jeromin, A., and Bassell, G.J. (2011). Fragile X mental retardation protein
regulates protein expression and mRNA translation of the potassium channel Kv4.2. J Neurosci *31*,
5693-5698. doi:10.1523/JNEUROSCI.6661-10.2011
Harnett, M.T., Xu, N.L., Magee, J.C., and Williams, S.R. (2013). Potassium channels control the
interaction between active dendritic integration compartments in layer 5 cortical pyramidal neurons.
Neuron *79*, 516-529. doi:10.1016/j.neuron.2013.06.005
Hayama, T., Noguchi, J., Watanabe, S., Takahashi, N., Hayashi-Takagi, A., Ellis-Davies, G.C., Matsuzaki,
M., and Kasai, H. (2013). GABA promotes the competitive selection of dendritic spines by controlling
local Ca2+ signaling. Nat Neurosci *16*, 1409-1416. doi:10.1038/nn.3496
Higley, M.J., and Contreras, D. (2006). Balanced excitation and inhibition determine spike timing during
frequency adaptation. J Neurosci *26*, 448-457. doi:10.1523/JNEUROSCI.3506-05.2006
Hoffman, D.A., Magee, J.C., Colbert, C.M., and Johnston, D. (1997). K+ channel regulation of signal
propagation in dendrites of hippocampal pyramidal neurons. Nature *387*, 869-875. doi:10.1038/43119
Jinno, S., Jeromin, A., and Kosaka, T. (2005). Postsynaptic and extrasynaptic localization of Kv4.2
channels in the mouse hippocampal region, with special reference to targeted clustering at gabaergic
synapses. Neuroscience *134*, 483-494. doi:10.1016/j.neuroscience.2005.04.065
Johnston, D., and Narayanan, R. (2008). Active dendrites: colorful wings of the mysterious butterflies.
Trends Neurosci *31*, 309-316. doi:10.1016/j.tins.2008.03.004
Kalmbach, B.E., Johnston, D., and Brager, D.H. (2015). Cell-Type Specific Channelopathies in the
Prefrontal Cortex of the fmr1-/y Mouse Model of Fragile X Syndrome. eNeuro *2*.
doi:10.1523/ENEURO.0114-15.2015
Kameyama, M., Kameyama, A., Kaibara, M., Nakayama, T., and Irisawa, H. (1989). Intracellular
mechanisms involved in "run-down" of calcium channels. Adv Exp Med Biol *255*, 111-117.
Kanemoto, Y., Matsuzaki, M., Morita, S., Hayama, T., Noguchi, J., Senda, N., Momotake, A., Arai, T., and
Kasai, H. (2011). Spatial distributions of GABA receptors and local inhibition of Ca2+ transients
studied with GABA uncaging in the dendrites of CA1 pyramidal neurons. PLoS One *6*, e22652.
doi:10.1371/journal.pone.0022652
Kim, J., Jung, S.C., Clemens, A.M., Petralia, R.S., and Hoffman, D.A. (2007). Regulation of dendritic
excitability by activity-dependent trafficking of the A-type K+ channel subunit Kv4.2 in hippocampal
neurons. Neuron *54*, 933-947. doi:10.1016/j.neuron.2007.05.026
Kim, J., Wei, D.S., and Hoffman, D.A. (2005). Kv4 potassium channel subunits control action potential
repolarization and frequency-dependent broadening in rat hippocampal CA1 pyramidal neurones. J
Physiol *569*, 41-57. doi:10.1113/jphysiol.2005.095042
Korngreen, A., and Sakmann, B. (2000). Voltage-gated K+ channels in layer 5 neocortical pyramidal
neurones from young rats: subtypes and gradients. J Physiol *525 Pt 3*, 621-639. doi:10.1111/j.1469-
7793.2000.00621.x
Larkum, M.E., Zhu, J.J., and Sakmann, B. (1999). A new cellular mechanism for coupling inputs arriving
at different cortical layers. Nature *398*, 338-341. doi:10.1038/18686
Losonczy, A., Makara, J.K., and Magee, J.C. (2008). Compartmentalized dendritic plasticity and input
feature storage in neurons. Nature *452*, 436-441. doi:10.1038/nature06725
Lovett-Barron, M., Turi, G.F., Kaifosh, P., Lee, P.H., Bolze, F., Sun, X.H., Nicoud, J.F., Zemelman, B.V.,
Sternson, S.M., and Losonczy, A. (2012). Regulation of neuronal input transformations by tunable
dendritic inhibition. Nat Neurosci *15*, 423-430, S421-423. doi:10.1038/nn.3024
Maffie, J.K., Dvoretskova, E., Bougis, P.E., Martin-Eauclaire, M.F., and Rudy, B. (2013). Dipeptidyl-
peptidase-like-proteins confer high sensitivity to the scorpion toxin AmmTX3 to Kv4-mediated A-type
K+ channels. J Physiol *591*, 2419-2427. doi:10.1113/jphysiol.2012.248831
Magee, J.C., and Carruth, M. (1999). Dendritic voltage-gated ion channels regulate the action potential
firing mode of hippocampal CA1 pyramidal neurons. J Neurophysiol *82*, 1895-1901.
Magee, J.C., and Johnston, D. (1997). A synaptically controlled, associative signal for Hebbian plasticity
in hippocampal neurons. Science *275*, 209-213. doi:10.1126/science.275.5297.209
Markram, H., Lubke, J., Frotscher, M., and Sakmann, B. (1997). Regulation of synaptic efficacy by
coincidence of postsynaptic APs and EPSPs. Science *275*, 213-215.
doi:10.1126/science.275.5297.213

Marlin, J.J., and Carter, A.G. (2014). GABA-A receptor inhibition of local calcium signaling in spines and
dendrites. J Neurosci *34*, 15898-15911. doi:10.1523/JNEUROSCI.0869-13.2014
Migliore, M., Hoffman, D.A., Magee, J.C., and Johnston, D. (1999). Role of an A-type K+ conductance in
the back-propagation of action potentials in the dendrites of hippocampal pyramidal neurons. J
Comput Neurosci *7*, 5-15.
Miles, R., Tóth, K., Gulyás, A.I., Hájos, N., and Freund, T.F. (1996). Differences between Somatic and
Dendritic Inhibition in the Hippocampus. Neuron *16*, 815-823. doi:10.1016/S0896-6273(00)80101-4
Miller, J.P., Rall, W., and Rinzel, J. (1985). Synaptic amplification by active membrane in dendritic spines.
Brain Res *325*, 325-330. doi:10.1016/0006-8993(85)90333-6
Mullner, F.E., Wierenga, C.J., and Bonhoeffer, T. (2015). Precision of Inhibition: Dendritic Inhibition by
Individual GABAergic Synapses on Hippocampal Pyramidal Cells Is Confined in Space and Time.
Neuron *87*, 576-589. doi:10.1016/j.neuron.2015.07.003
Paille, V., Fino, E., Du, K., Morera-Herreras, T., Perez, S., Kotaleski, J.H., and Venance, L. (2013).
GABAergic circuits control spike-timing-dependent plasticity. J Neurosci *33*, 9353-9363.
doi:10.1523/JNEUROSCI.5796-12.2013
Palmer, L., Murayama, M., and Larkum, M. (2012). Inhibitory Regulation of Dendritic Activity in vivo. Front
Neural Circuits *6*, 26. doi:10.3389/fncir.2012.00026
Pathak, D., Guan, D., and Foehring, R.C. (2016). Roles of specific Kv channel types in repolarization of
the action potential in genetically identified subclasses of pyramidal neurons in mouse neocortex. J
Neurophysiol *115*, 2317-2329. doi:10.1152/jn.01028.2015
Poirazi, P., and Mel, B.W. (2001). Impact of active dendrites and structural plasticity on the memory
capacity of neural tissue. Neuron *29*, 779-796. doi:10.1016/S0896-6273(01)00252-5
Pouille, F., and Scanziani, M. (2001). Enforcement of temporal fidelity in pyramidal cells by somatic feed-
forward inhibition. Science *293*, 1159-1163. doi:10.1126/science.1060342
Ramakers, G.M., and Storm, J.F. (2002). A postsynaptic transient K(+) current modulated by arachidonic
acid regulates synaptic integration and threshold for LTP induction in hippocampal pyramidal cells.
Proc Natl Acad Sci U S A *99*, 10144-10149. doi:10.1073/pnas.152620399
Rial Verde, E.M., Zayat, L., Etchenique, R., and Yuste, R. (2008). Photorelease of GABA with Visible
Light Using an Inorganic Caging Group. Front Neural Circuits *2*, 2. doi:10.3389/neuro.04.002.2008
Sandler, M., Shulman, Y., and Schiller, J. (2016). A Novel Form of Local Plasticity in Tuft Dendrites of
Neocortical Somatosensory Layer 5 Pyramidal Neurons. Neuron *90*, 1028-1042.
doi:10.1016/j.neuron.2016.04.032
Serodio, P., Kentros, C., and Rudy, B. (1994). Identification of molecular components of A-type channels
activating at subthreshold potentials. J Neurophysiol *72*, 1516-1529.
Serodio, P., and Rudy, B. (1998). Differential expression of Kv4 K+ channel subunits mediating
subthreshold transient K+ (A-type) currents in rat brain. J Neurophysiol *79*, 1081-1091.
Shepherd, G.M., Brayton, R.K., Miller, J.P., Segev, I., Rinzel, J., and Rall, W. (1985). Signal
enhancement in distal cortical dendrites by means of interactions between active dendritic spines.
Proc Natl Acad Sci U S A *82*, 2192-2195.
Stokes, C.C., Teeter, C.M., and Isaacson, J.S. (2014). Single dendrite-targeting interneurons generate
branch-specific inhibition. Front Neural Circuits *8*, 139. doi:10.3389/fncir.2014.00139
Stuart, G., Schiller, J., and Sakmann, B. (1997). Action potential initiation and propagation in rat
neocortical pyramidal neurons. J Physiol *505 (Pt 3)*, 617-632.
Tsubokawa, H., and Ross, W.N. (1996). IPSPs modulate spike backpropagation and associated [Ca2+]i
changes in the dendrites of hippocampal CA1 pyramidal neurons. J Neurophysiol *76*, 2896-2906.
Wehr, M., and Zador, A.M. (2003). Balanced inhibition underlies tuning and sharpens spike timing in
auditory cortex. Nature *426*, 442-446. doi:10.1038/nature02116
Wilmes, K.A., Sprekeler, H., and Schreiber, S. (2016). Inhibition as a Binary Switch for Excitatory
Plasticity in Pyramidal Neurons. PLoS Comput Biol *12*, e1004768. doi:10.1371/journal.pcbi.1004768
Yuste, R. (1997). Potassium channels. Dendritic shock absorbers. Nature *387*, 851, 853.
doi:10.1038/43063
Zhu, Y., Stornetta, R.L., and Zhu, J.J. (2004). Chandelier cells control excessive cortical excitation:
characteristics of whisker-evoked synaptic responses of layer 2/3 nonpyramidal and pyramidal
neurons. J Neurosci *24*, 5101-5108. doi:10.1523/JNEUROSCI.0544-04.2004

Figure Legends

Figure 1. GABAergic inhibition of ∆Ca2+ is enhanced by blockade of KA channels.

(**A**) Whole Cell patch recording where performed in L5PNs of visual cortex. Ca2+ imaging and GABA uncaging were performed along the proximal apical dendrite, as shown in the example cell (A1) and schematic (A2). Scale bar 25 µm. (**B**) Example spine-dendrite pair and the associated line-scanned response to a bAP. (**C**) Average ± SEM somatic voltage recorded before (black) and after (green) treatment with 4-AP for action potentials (upper traces) and uncaging-evoked IPSPs. (**D**) Example bAP-evoked ∆Ca2+ for the apical dendritic region shown in (B) for bAP alone (black, blue) or paired with GABA uncaging (red, orange) for the spine and neighboring dendrite, before (left) and after (right) treatment with 5 mM 4-AP. (**E**) Average (n=10) ∆Ca2+ for the population of imaged spines and dendrites, colors as in (D). (**F-G**) Population data (n=10) showing the magnitude of ∆Ca2+ inhibition and peak bAP-evoked ∆Ca2+ for spines (F) and neighboring dendrites (G) before (black) and after (green) treatment with 5 mM 4-AP (Wilcoxon matched-pairs signed rank test, *p<0.05).

Figure 1 — figure supplement 1. Picrotoxin blocks the effects of GABA uncaging.

Figure 1 — figure supplement 2. Saline does not alter GABAergic inhibition of ∆Ca2+.

Figure 1 — figure supplement 3. Somatic IPSPs are enhanced by blockade of KA channels.

Figure 1 — figure supplement 4. GABAergic inhibition of ∆Ca2+ is not enhanced by 4-AP in basal dendrites

Figure 1— figure supplement 1. Picrotoxin blocks the effects of GABA uncaging.

(**A**) Schematic of recording and imaging configuration. Experiments are performed in the presence of picrotoxin (100 µM). (**B**) Average ± SEM (n=6) Ca2+ transients in proximal apical dendrites for bAP alone or paired with GABA uncaging before (left) and after (right) treatment with 4-AP (colors as in Fig. 1). (**C-D**) Population data (n=6) showing the magnitude of Ca2+ inhibition and peak bAP-evoked Ca2+ for spines (C) and neighboring dendritic shafts (D) in control (black) and 4-AP (green) conditions (Wilcoxon matched-pairs signed rank test, *p<0.05).

Figure 1 — figure supplement 2. Saline does not alter GABAergic inhibition of ∆Ca2+.

(**A**) Schematic of recording and imaging configuration. (**B**) Average ± SEM (n=7) Ca2+ transients in proximal apical dendrites for bAP alone or paired with GABA uncaging before (left) and after (right) treatment with control saline (colors as in Fig. 1). (**C-D**) Population data (n=7) showing the magnitude of Ca2+ inhibition and peak bAP-evoked Ca2+ for spines (C) and neighboring dendritic shafts (D) in control (black) and saline (green) conditions (Wilcoxon matched-pairs signed rank test, *p<0.05).

Figure 1 — figure supplement 3. Somatic IPSPs are enhanced by blockade of KA channels.

(**A**) Schematic of recording configuration. (**B**) Average ± SEM (n=9) IPSPs before (black) and after
(green) treatment with 5 mM 4-AP for single (left) and triple (right) GABA uncaging events. (**C**) Population
data (n=9) showing peak IPSP amplitude (left) and ratio of third:first amplitude (right) in control (black)
and 4-AP (green) conditions (Wilcoxon matched-pairs signed rank test, *p<0.05). (**D**) Population data
(n=9) showing decay time constant for single IPSPs (Wilcoxon matched-pairs signed rank test, *p<0.05).
(**E**) Population data (n=9) showing somatic input resistance (Wilcoxon matched-pairs signed rank test,
p>0.05)
605

Figure 1 — figure supplement 4. GABAergic inhibition of ΔCa2+ is not enhanced by 4-AP in basal
dendrites
(**A**) Schematic of recording and imaging configuration. (**B**) Average ± SEM (n=10) Ca2+ transients in
basal dendrites for bAP alone or paired with GABA uncaging before (left) and after (right) treatment with
4-AP (colors as in Fig. 1). (**C-D**) Population data (n=10) showing the magnitude of Ca2+ inhibition and
peak bAP-evoked Ca2+ for spines (C) and neighboring dendritic shafts (D) in control (black) and 4-AP
(green) conditions (Wilcoxon matched-pairs signed rank test, *p<0.05).
613

Figure 2. GABAergic inhibition of ΔCa2+ is enhanced by local blockade of KA channels.
(**A**) Schematic of recording and imaging configuration, illustrating location of puffed 4-AP. (**B**) Average ±
SEM (n=7) Ca2+ transients for bAP alone or paired with GABA uncaging before (left) and after (right)
application of 4-AP to the proximal apical dendrite (colors as in Fig. 1). (**C-D**) Population data (n=7)
showing the magnitude of Ca2+ inhibition and peak bAP-evoked Ca2+ for spines (C) and neighboring
dendritic shafts (D) in control (black) and 4-AP (green) conditions (Wilcoxon matched-pairs signed rank
test, *p<0.05).
Figure 2 — figure supplement 1. GABAergic inhibition of ΔCa2+ in the proximal apical dendrite is
not affected by somatic or distal dendritic KA channel blockade.
Figure 2 — figure supplement 2. Effect of blockade of potassium channels at proximal apical
dendrite and soma on somatic IPSP.
625

Figure 2 — figure supplement 1. GABAergic inhibition of ΔCa2+ in the proximal apical dendrite is
not affected by somatic or distal dendritic KA channel blockade.
(**A**) Schematic of recording and imaging configuration, illustrating somatic location of puffed 4-AP. (**B**)
Average ± SEM (n=6) Ca2+ transients for bAP alone or paired with GABA uncaging before (left) and after
(right) application of 4-AP to the soma (colors as in Fig. 1). (**C-D**) Population data (n=6) showing the
magnitude of Ca2+ inhibition and peak bAP-evoked Ca2+ for spines (C) and neighboring dendritic shafts
(D) in control (black) and 4-AP (green) conditions (Wilcoxon matched-pairs signed rank test, p>0.05). (**E**)
Schematic of experimental configuration, illustrating distal dendritic location of puffed 4-AP. (**F**) Average ±
SEM (n=6) Ca2+ transients for bAP alone of paired with GABA uncaging before (left) and after (right)

application of 4-AP to the distal dendrite (colors as in Fig. 1). (**G-H**) Population data (n=6) showing the magnitude of Ca2+ inhibition and peak bAP-evoked Ca2+ for spines (G) and neighboring dendritic shafts (H) in control (black) and 4-AP (green) conditions (Wilcoxon matched-pairs signed rank test, p>0.05).

Figure 2 — figure supplement 2. Effect of blockade of potassium channels at proximal apical dendrite and soma on somatic IPSP

(**A**) Schematic of recording configuration, illustrating dendritic location of puffed 4-AP. (**B**) Average ± SEM (n=9) IPSPs before (black) and after (green) application of 4-AP to the apical dendrite for single (upper) and triple (lower) GABA uncaging events. (**C**) Population data (n=9) showing peak IPSP amplitude (left) and ratio of third:first amplitude (right)(Wilcoxon matched-pairs signed rank test, *p<0.05). (**D**) Population data (n=9) showing decay time constant for single IPSPs (Wilcoxon matched-pairs signed rank test, p<0.05). (**E**) Population data (n=9) showing somatic input resistance (Wilcoxon matched-pairs signed rank test, p<0.05). (**F-J**) As in (A-E) for local application of 4-AP to the soma (n=9, Wilcoxon matched-pairs signed rank test, *p<0.05).

Figure 3. GABAergic inhibition of ∆Ca2+ is enhanced by blockade of Kv4 channels using AmmTx3.

(**A**) Schematic showing recording and imaging configuration. (**B**) Average ± SEM (n=10) Ca2+ transients for bAP alone or paired with GABA uncaging, before (left) and after (right) treatment with the selective Kv4 blocker AmmTx3 (200 nM) (colors as in Fig. 1). (**C-D**) Population data (n=10) showing the magnitude of Ca2+ inhibition and peak bAP-evoked Ca2+ for spines (C) and neighboring dendritic shafts (D) in control (black) and AmmTx3 (green) conditions (Wilcoxon matched-pairs signed rank test, *p<0.05).
Figure 3 — figure supplement 1. Blockade of Kv4 by AmmTx3 has no effect on somatic IPSP
Figure 3 — figure supplement 2. Outside-out recordings reveal expression of Kv4 channels.

Figure 3 — figure supplement 1. Blockade of Kv4 by AmmTx3 has no effect on somatic IPSP

(**A**) Schematic of recording configuration. (**B**) Average ± SEM (n=6) IPSPs before (black) and after (green) treatment with 200 nM AmmTx3 for single (left) and triple (right) GABA uncaging events. (**C**) Population data (n=6) showing peak IPSP amplitude (left) and ratio of third:first amplitude (right) in control (black) and AmmTx3 (green) conditions (Wilcoxon matched-pairs signed rank test, *p<0.05). (**D**) Population data (n=6) showing decay time constant for single IPSPs (Wilcoxon matched-pairs signed rank test, p>0.05). (**E**) Population data (n=6) showing somatic input resistance (Wilcoxon matched-pairs signed rank test, *p>0.05).

Figure 3 — figure supplement 2. Outside-out recordings reveal expression of Kv4 channels.

(**A**) Peak-normalized average ± SEM (n=6) currents evoked in somatic outside-out patches before (black) and after (green) treatment with 4-AP. (**B**) Peak-normalized average ± SEM (n=6) currents evoked in

 outside-out patches before (black) and after (green) treatment with AmmTx3. (**C**) Population data (n=6)

showing reduction in current produced by 4-AP or AmmTx3 for the measured intervals marked in

(A)(Wilcoxon matched-pairs signed rank test for single columns, Unpaired t-test for comparison between

treatments, *p<0.05).

Figure 4. Somatic activity enhances GABAergic inhibition of dendritic ∆Ca2+.

(**A**) Average ± SEM (n=10) of somatic recordings for a single action potential (black) or when preceded 20

ms by a 100 Hz train of action potentials (green). Inset shows change in spike waveform. (**B**) Average ±

SEM (n=10) Ca2+ transients for bAP alone or paired with GABA uncaging, presented either singly (left) or

following a 100 Hz train of action potentials (right)(colors as in Fig. 1). (**C-D**) Population data (n=10)

showing the magnitude of Ca2+ inhibition and peak bAP-evoked Ca2+ for spines (C) and neighboring

dendritic shafts (D), presented either singly (black) or following a 100 Hz train of action potentials

(green)(Wilcoxon matched-pairs signed rank test, *p<0.05).

Figure 4— figure supplement1. Activity-dependent enhancement of GABAergic inhibition is

frequency-dependent and occluded by blockade of KA channels.

Figure 4— figure supplement1. Activity-dependent enhancement of GABAergic inhibition is

frequency-dependent and occluded by blockade of KA channels.

(**A**) Average ± SEM (n=10) of somatic recordings for a single action potential (black) or when preceded

20 ms by a 50 Hz train of action potentials (green). Inset shows change in spike waveform. (**B**) Average ±

SEM (n=10) Ca2+ transients for bAP alone or paired with GABA uncaging, presented either singly (left) or

following a 50 Hz train of action potentials (right)(colors as in Fig. 1). (**C-D**) Population data (n=10)

showing the magnitude of Ca2+ inhibition and peak bAP-evoked Ca2+ for spines (C) and neighboring

dendritic shafts (D), presented either single (black) or following a 50 Hz train of action potentials

(green)(Wilcoxon matched-pairs signed rank test, *p<0.05). (**E**) Average ± SEM (n=10) of somatic

recordings for a single action potential (black) or when preceded 20 ms by a 100 Hz train of action

potentials (green), in the presence of 5 mM 4-AP. (**F-H**) As in (B-D) for a 100 Hz preceding train and in

the presence of 5 mM 4-AP (n=10, Wilcoxon matched-pairs signed rank test, *p<0.05).

Figure 5. Somatic activity enhances synaptic GABAergic inhibition of ∆Ca2+.

(**A**) Schematic showing recording and imaging configuration. ChR2 was virally expressed in somatostatin-

containing interneurons (SOM-INs) and activated with blue light pulses. (**B**) Average ± SEM (n=10) Ca2+

transients for bAP alone or paired with optical stimulation of SOM-INs, presented either singly (left) or

following a 100 Hz train of action potentials (right)(colors as in Fig. 1). (**C-D**) Population data (n=10)

showing the magnitude of Ca2+ inhibition and peak bAP-evoked Ca2+ for spines (C) and neighboring

dendritic shafts (D), presented either singly (black) or following a 100 Hz train of action potentials

(green)(Wilcoxon matched-pairs signed rank test, *p<0.05).

Figure 6. Computational simulations reveal mechanisms underlying KA-dependent regulation of
inhibition.
(**A**) Simulated Ca2+ currents under three different conditions (indicated in figure) for bAPs evoked alone
(black) or preceded by GABAergic inhibition (red). Yellow shading indicates integral of inhibited Ca2+. (**B**)
Relationship between sodium reversal potential and either peak membrane potential (left) or Ca2+
inhibition (right). For peak V_m data, values for bAP alone (black), bAP paired with inhibition (red), and
resting V_m (green) are shown. (**C**) Relationship between KA conductance and either peak membrane
potential (left) or Ca2+ inhibition (right). Colors are as in (B). (**D**) Relationship between peak V_m and Ca2+
inhibition under conditions where either sodium reversal potential (closed circles) or KA conductance
(open circles) was varied. (**E**) Relationship between action potential duration and Ca2+ inhibition under
conditions where either sodium reversal potential (closed circles) or KA conductance (open circles) was
varied.
721
722

Figure 1

Figure 1 S1

Figure 1 S2

Figure 1 S3

Figure 1 S4

Figure 2

Figure 2 S1

Figure 2 S2

Figure 3

Figure 3 S1

Figure 3 S2

Figure 4

Figure 4 S1

Figure 5

A_i

E_{Na} = 60 mV
g_A = 12.1 mS/cm²
ΔCa_{Inh} = -22%

2 ms

A_{ii}

E_{Na} = 75 mV
g_A = 12.1 mS/cm²
ΔCa_{Inh} = -20%

A_{iii}

E_{Na} = 60 mV
g_A = 2.4 mS/cm²
ΔCa_{Inh} = -26%

B

Peak V_m (mV)

E_{Na} (mV)

ΔCa^{2+} Inhibition (% change)

E_{Na} (mV)

C

Peak V_m (mV)

g_A (mS/cm²)

ΔCa^{2+} Inhibition (% change)

g_A (mS/cm²)

D

ΔCa^{2+} Inhibition (% change)

Peak V_m (mV)

E

ΔCa^{2+} Inhibition (% change)

AP FWHM (ms)

○ Vary g_A
● Vary E_{Na}

Figure 6