Watching the clock for 25 years in

FlyClockbase:

Variability in circadian clocks of *Drosophila melanogaster*

as uncovered by biological model curation

Katherine S. **Scheuer**[*,†], Bret **Hanlon**[‡], Jerdon W. **Dresel**[*,†], Erik D. **Nolan**[*,†],

John C. **Davis**[‡], Laurence **Loewe**[*,†]

[*]Systems Biology Theme, Wisconsin Institute for Discovery,

University of Wisconsin-Madison, Madison, WI, 53715

[†]Laboratory of Genetics, University of Wisconsin-Madison, Madison, WI 53706

[‡]Department of Statistics, University of Wisconsin-Madison, Madison, WI, 53706

General Article Summary

Circadian clocks impact health and fitness by controlling daily rhythms of gene-expression through complex gene-regulatory networks. Deciphering how they work requires experimentally tracking changes in amounts of clock components. We designed FlyClockbase to simplify data-access for biologists and modelers and curated over 400 time series observed in wildtype fruit flies from 25 years of research on clocks. We found differences in peak time variance of the clock-proteins 'PERIOD' and 'TIMELESS', which probably stem from differences in phosphorylation-network complexity. Combining in-depth circadian-biology, model-curation, and compiler logic, our trans-disciplinary research shows how biology-friendly compilers could simplify model curation enough to democratize it.

Running title: FlyClockbase time series variance curation

Keywords / Key phrases:

Drosophila melanogaster circadian clock model biological data input review,

experimental time series peak-valley variance and outlier clock observations,

differential variance hypothesis on PERIOD - TIMELESS amount peak times,

data repository for estimating parameters of mechanistic simulation models,

compiler logic enabling human error analysis simplifying biological model curation.

Corresponding author:

Laurence Loewe

Wisconsin Institute for Discovery,   University of Wisconsin-Madison,

330 N Orchard St,   Madison, WI, 53715

Email and Phone: loewe@wisc.edu   (608) 316-4324

Statement of data availability and stabilizing versioning number: **QQv1r1**

For review purposes: QQv1r1 zip-archive in Supplemental Material or upon request.

Before final publication: FlyClockbase will be at ***https://github.com/FlyClockbase***

Abbreviations: Table 1: Core clock components Table 2: Concepts in FlyClockbase

Supporting Material:

Supporting Text and Tables,   Supplemental Statistical Analysis (87 pages),

R-Script zip file (>12K lines, QQv1), FlyClockbase zip file (QQv1).

Abstract

High-quality model curation provides insights by organizing biological knowledge-fragments. We aim to integrate published results about circadian clocks in *Drosophila melanogaster* while exploring economies of scale in model curation. Clocks govern rhythms of gene-expression that impact fitness, health, cancer, memory, mental functions, and more. Human clock insights have been pioneered in flies. Flies simplify investigating complex gene regulatory networks, which express proteins cyclically using environmentally entrained interlocking feedback loops. Simulations could simplify research further, but currently few models test their quality directly against experimentally observed time series scattered across publications. We designed FlyClockbase for robust efficient access to such scattered data for biologists and modelers, prioritizing simplicity and openness to encourage experimentalists to preserve more annotations and raw-data. Such details could multiply long-term value for modelers interested in meta-analyses, parameter estimates, and hypothesis testing. Currently FlyClockbase contains over 400 wildtype time series of core circadian components systematically curated from 86 studies published between 1990 and 2015. Using FlyClockbase, we show that PERIOD protein amount peak time variance unexpectedly exceeds that of TIMELESS. We hypothesize, PERIOD's exceedingly more complex phosphorylation rules are responsible. Human error analysis improved data quality and revealed significance-degrading outliers, possibly violating presumed absence of wildtype heterogeneity or lab evolution. We found PCR-measured peak time variances exceed those from other methods, pointing to initial count stochasticity. Our trans-disciplinary analyses demonstrate how compilers with more biology-friendly logic could simplify, guide, and naturally distribute biological model curation. Resulting quality increases and cost reductions benefit curation-dependent grand challenges like personalizing medicine.

Table of Contents

TABLES OVEREIW – MAIN TEXT

FIGURES OVEREIW – MAIN TEXT

RELATED ANALYSES

Hypotheses comparing timing in PER and TIM: data and analyses in Table 4, Table S1, and Table 6, Table 7, Figure 6, Figure 7, as well as parts of the Supplemental Statistical Analysis are related.

Comparing PCR vs non-PCR methods for measuring time series in per mRNA: data and analyses in Table 8, Figure 8, Figure 9, Figure 10, and parts of the Supplemental Statistical Analysis are related.

Watching the clock for 25 years in FlyClockbase:
Variability in circadian clocks of *Drosophila melanogaster* as uncovered by biological model curation

Katherine S. **Scheuer**[*,†], Bret **Hanlon**[‡], Jerdon W. **Dresel**[*,†],
Erik D. **Nolan**[*,†], John C. **Davis**[‡], Laurence **Loewe**[*,†]

[*]Systems Biology Theme, Wisconsin Institute for Discovery, [†]Laboratory of Genetics,
[‡]Department of Statistics, University of Wisconsin-Madison

Appendix: Supplemental Material

Table of Contents

Intentionally free

2017-01-17

INTRODUCTION

Several grand challenges of our time, such as personalizing medicine or mapping

genotypes to phenotypes, critically depend on the careful curation of biological

knowledge-fragments into integrated resources of intermediate size that are easier to

handle. Such resources can provide comprehensive overviews of integrated models or

experimental results on a given topic. If these resources are organized well enough and

are machine readable, then biological models and datasets can be explored in more

automated ways and thus greatly accelerate biological discovery.

We aim to make it easier to create such intermediate resources by improving the

efficiency of high-quality biological model curation. The goal of such curation work is to

integrate 'all current knowledge' that is relevant for a research topic of reasonable

complexity *while* keeping all deposited information well-organized and machine

readable. This requires computational solutions that are best developed *while*

simultaneously engaging deeply with the complexities of real-world biological research

where results can be less than clear-cut and relevant data may be scattered across

diverse sources. Curating such diffuse and scattered data can be prohibitively

complicated without appropriate strategies for handling recurrent problems.

We chose the study of circadian clocks in fruit flies as our area of in-depth

biological research in order to provide a real-world context for developing strategies that

improve curation efficiency. While climbing onto the shoulders of giants in fly clock

research, we integrated as much fly clock expertise as we could. In our opinion, a full

integration is currently far beyond the scope of any single study if it is to efficiently point

readers to the detailed, evidence-based evaluations of the strengths and weaknesses of

a state-of-the-art fly clock model. Thus, we focused on integrating all *Drosophila*

melanogaster wildtype time series observations from 25 years of research (often

reported as wildtype control experiments for evaluating effects of mutants). Despite this

substantial reduction of scope, our integration task is far from trivial if we aim to ensure

the reproducibility, stability, and rigor of integration.

Reproducibility is pivotal for science. It also does not come easy. We aimed to

increase the reproducibility of results from our research in fly clock biology while

exploring strategies for simplifying reproducibility in research. Our main biological

findings are differences in variances of certain time series traits that we observed

between different clock components. We hypothesize that these differences hold

important clues for improving our mechanistic understanding of circadian clocks.

Variances are easily affected by errors that also affect reproducibility and are

independent from underlying biological mechanisms. Therefore, we deem it essential to

include our progress towards reproducibility in the scope of this study. We mitigate the

inevitable increase of length with headings that simplify navigating its various aspects.

Our study contributes to the foundations of a system for integrating all expertise

on the fly clock. We provide detailed experimental observations of time series ready for

linking to statements in 'big-picture' fly clock models. We simultaneously explore how to

integrate more efficiently the underpinning diffuse and scattered data. We find that such

integration work is best accomplished by biological model curators with a deep

biological interest in the research results that are being integrated. We also find that the

efficiency of integrating and curating results can be greatly increased by access to

strategies and tools designed to handle complex biological observations. We note that

current computational tools repeatedly restrict representation possibilities to options that

almost fit observed data – but not *entirely*. Such cases force model curators to

inappropriately ignore data or subtly bias results by defining the closest representation

as 'good enough'. We ask if well-known data structures and logic formalisms originally

developed outside of biology can appropriately capture real-world observations in

biology. If not, is there merit in breaking the mold? Can economies of scale be

leveraged in model curation? Abstraction is critical. If solutions are too abstract, they

work everywhere – albeit poorly; if not abstract enough, they work perfectly for *one*

problem – but force reinventing the wheel next time. To guide our development of

computational abstractions, we find it essential to constantly face the challenging

complexities of real-world experimental data as we work to advance biological research

in circadian clocks of flies. This is where useful abstractions emerge naturally.

Accordingly, this study has three strands: (i) introduce FlyClockbase, the new

resource we produced, and measure its reliability; (ii) present new biological insights on

clocks in flies from analyzing data in FlyClockbase; (iii) evaluate emergent opportunities

for abstraction as seen by a programming language compiler architect aiming to

improve the efficiency of navigating the tension between the clear-cut logic formalisms

in computers and the uncertain, incomplete and noisy biological data. We found that all

three strands significantly strengthened each other. Each presents a distinct view on the

integrated body of trans-disciplinary research presented here.

Circadian clocks are biochemical pathways characterized by cyclical protein expression. They play a critical role in a wide variety of behavioral and physiological processes, and a better understanding of their genetic and biochemical bases could advance research in many areas (PREUSSNER AND HEYD 2016; SHARMA *et al.* 2016), including consciousness and sleep (CIRELLI 2009), feeding and metabolism (XU *et al.* 2008; HURLEY *et al.* 2016), learning and memory (XU *et al.* 2008; CHOUHAN *et al.* 2015), stress and immunity (DUMBELL *et al.* 2016), inflammation (CARTER *et al.* 2016), cancer (SEPHTON AND SPIEGEL 2003; MASRI *et al.* 2015; SALAVATY 2015; MOLINA-RODRÍGUEZ AND ÁLVAREZ 2016), and psychological functioning (MCCLUNG 2013; PAREKH *et al.* 2015; COOGAN *et al.* 2016).

Model organisms. Many model organisms have been used to study circadian rhythms, including *Synechococcus elongatus*, *Neurospora crassa*, *Arabidopsis thaliana, Mus musculus*, and *D. melanogaster* (BELL-PEDERSEN *et al.* 2005). Each model organism presents benefits and challenges. Here we focus on *D. melanogaster*, which is known for its ease of genetic manipulation (STANEWSKY 2003; ÖZKAYA AND ROSATO 2012) and its well-characterized genome (DOS SANTOS *et al.* 2015). Human circadian rhythms are certainly more complex than those of *D. melanogaster*. For example, it is not uncommon for one fly circadian clock component (e.g., *period* or *cryptochrome*) to correspond to multiple mammalian circadian clock components (e.g., *period1* and *period2*, or *cryptochrome1* and *cryptochrome2*) as reviewed elsewhere (YOUNG AND KAY 2001). Despite these differences in complexity, the *D. melanogaster* clock is similar in

many aspects to the mammalian clock (YOUNG AND KAY 2001; ROSATO *et al.* 2006).

Insights from the fly clock have substantially contributed to understanding aspects of the

mammalian clock in general and in particular related disorders such as familial

advanced sleep phase syndrome (FASPS) (ROSATO *et al.* 2006), pancreatic cancer

(POGUE-GEILE *et al.* 2006), and bipolar disorder (KO *et al.* 2010; MCCLUNG 2013).

Increased knowledge of the *D. melanogaster* circadian clock could continue to provide

important information for future work in a variety of areas pertaining to the mammalian

clock, including sleep disorders (WAGER-SMITH AND KAY 2000), Alzheimer's disease

(LONG *et al.* 2014), and psychiatric disorders (MCCLUNG 2013; ZORDAN AND SANDRELLI

2015).

Math models. Computer simulations of mathematical models are powerful tools

for studying the dynamics of complex non-linear systems such as circadian clocks. They

have been used for decades in many disciplines (KURTZ 1972; CROSBY 1973;

TARANTOLA AND VALETTE 1982; ASCHER AND PETZOLD 1998; LAW AND KELTON 2000;

ZEIGLER *et al.* 2000; TARANTOLA 2005; ANDERSON 2007; GILLESPIE 2007; GILLESPIE 2008;

ANDERSON *et al.* 2011; KARR *et al.* 2012; MAVELLI 2012; WILKINSON 2012; ZEIGLER 2012;

DISTEFANO 2013; GILLESPIE *et al.* 2013; SANGHVI *et al.* 2013; KARR *et al.* 2014; CHYLEK

et al. 2015; KARR *et al.* 2015a). To be useful for the study of circadian clocks,

mathematical models need to mirror relevant aspects of real-world clocks, which may

include key mechanisms, reaction rates, and/or other parameters or traits. Models

integrate the specified details to enable simulations of time series of amounts of

circadian clock components that are based on the assumptions of the *in silico* model.

The simulated distributions of amounts of different clock components at specified times is expected to match observable real-world time series if a model's assumptions are correct. Such simulations are facilitated by rigorous simulation algorithms that have a rich history in modeling biochemical reaction networks (KURTZ 1972; GILLESPIE 1977; ANDERSON 2007; GILLESPIE 2007; GILLESPIE 2008; ANDERSON *et al.* 2011; KARR *et al.* 2012; MAVELLI 2012; WILKINSON 2012; DISTEFANO 2013; GILLESPIE *et al.* 2013; SANGHVI *et al.* 2013; KARR *et al.* 2014; CHYLEK *et al.* 2015; KARR *et al.* 2015a).

Estimating unknown rates from observed time series. If the structure of a model is essentially correct but its parameter values are not, then time series observed in the real world can, in principle, be used to narrow the margins of uncertainty around poorly-known rate parameters (TARANTOLA AND VALETTE 1982; TARANTOLA 2005). Thus, access to a high-quality collection of observed time series could substantially contribute to improving the quality of biological insights gained from computational models. Mechanistic models with firm mathematical underpinnings can be used to explore hypotheses that are impractical to investigate in the laboratory for reasons that may include the effort required to produce mutants or the difficulty of measuring particular clock features experimentally (LOEWE AND HILLSTON 2008; LOEWE 2009; LOEWE 2016). By exploring potential hypotheses of interest *in silico*, models ideally inform future wet-lab experiments.

Biological example. The clock component *clockwork orange* (*cwo*) was thought to indirectly inhibit transcription of a number of key clock genes, including *period* (*per*), *PER-aryl-domain protein 1* (*pdp1*), and *vrille* (*vri*) (KADENER *et al.* 2007; LIM *et al.* 2007),

so flies with decreased *cwo* expression were expected to show increased levels of *per*, *pdp1*, and *vri*. Experimental results, however, indicated that *cwo* mutants exhibited decreased expression of these clock components (MATSUMOTO *et al.* 2007; RICHIER *et al.* 2008). In an attempt to explain these results, FATHALLAH-SHAYK et al. (2009) created a clock model that included *cwo*. This model was able to predict the experimental results previously shown and was used to develop a novel hypothesis which described a more complex interaction between *cwo* and the rest of the clock. Rather than simply repressing a transcriptional activator, the authors of this model postulated that the interaction between weak repression by *cwo* and strong activation by the transcriptional activator led to indirect activation of a different part of the clock. This explained the experimental results from *cwo* mutants and suggested that *cwo* plays a role in reducing minor variations in the clock known as "jitters" ((FATHALLAH-SHAYKH 2010); more details (SCRIBNER AND FATHALLAH-SHAYKH 2011)). This is just one example of the powerful ways in which modeling can act as a thinking tool, helping us to understand biology better.

  Models and reality. Computational models simulations models are fundamentally attempts to represent a simplified version of reality, and their utility hinges on their ability to faithfully capture the most important aspects of reality (TARANTOLA AND VALETTE 1982; TARANTOLA 2005). Complex processes with well-defined inputs and outputs are easily simplified by assuming that the timing of these processes remains essentially unchanged; then complicated sub-models of such processes can be substituted by simple transformations that merely reproduce the correct timing. If a model is to help understand, describe, or predict a biological system, such simplifications must be grounded in observations and a biological understanding of the phenomena to be

modeled (WOOLEY AND LIN 2005; BRODLAND 2015). For example, a model for predicting

credible functions for *cwo* in the clock (FATHALLAH-SHAYKH *et al.* 2009) needed to be

able to replicate previous experimental results first, before it could make useful

predictions. Such work is facilitated by the simplifying assumption that transcription,

translation, and degradation in clock models can be replaced by simple reactions with

rates appropriately chosen to match experimental data. Thus, access to a broad array

of curated, high-quality experimental observations is critical for efficiently constructing

and refining computational models.

Reproducibility of research. The advanced mechanistic simulations and complex

statistical inference methods above are necessary for arriving at a rigorous

understanding of circadian clocks. They require a complex software stack and

substantial efforts to implement dedicated code, workflows, and data organization

schemes. It is not easy to develop scientific computing solutions of such complexity

without loss of usability or sacrificing the reproducibility of earlier results. Yet the

importance of reproducibility for science is undisputed and has recently received some

attention (IOANNIDIS 2005b; JASNY *et al.* 2011; HUANG AND GOTTARDO 2013; MCNUTT

2014; AARTS *et al.* 2015; FREEDMAN *et al.* 2015a; ALLISON *et al.* 2016; BAKER 2016;

BARBA 2016; LEWIS *et al.* 2016; STODDEN *et al.* 2016). Reproducibility is an extremely

broad topic that frequently requires input from many experimental, statistical,

computational, theoretical and applied disciplines to arrive at rigorous solutions (HUANG

AND GOTTARDO 2013). Conducting research reproducibly requires more effort than

commonly realized (DONOHO 2009; STODDEN *et al.* 2014; JAMES *et al.* 2015; LOEWE *et al.*

2016b; MESNARD AND BARBA 2016). Yet, the steep upfront costs of entry seem to pay off: research teams with a reproducible research workflow report substantial benefits (DONOHO 2009; MESNARD AND BARBA 2016). Evaluations of computational tools Such reports inspired us to work towards improving reproducibility in our efforts. While useful recommendations and tools exist e.g. see https://www.xsede.org/web/reproducibility and (INCE *et al.* 2012; STODDEN *et al.* 2014; POLDRACK AND POLINE 2015; LEWIS *et al.* 2016; STODDEN *et al.* 2016), there is no silver bullet and standards are still evolving. We aimed to keep computational requirements to a minimum.

Firm foundations. Here we cannot investigate the reproducibility of complex mechanistic circadian clock simulations or their underpinning parameter estimates. However, we can prepare a firm foundation for later studies. Rigorous reproducible reports of new parameter estimates need to provide many of the details reported here. This includes details on (i) literature database search strategies, (ii) initial screening processes and criteria, (iii) filtering of candidate studies and other special selection methods, (iv) reasons for combining some datasets but not others, (v) justifications for approaches that handle noise, outliers, or other complications, (vi) bibliographic references, (vii) a version of all raw data before preprocessing, and (viii) a well-integrated final version of the modified data used for inference after preprocessing. Ideally, such a study would include additional analyses like (ix) a human error analysis providing estimates of some low-level error rates quantifying the quality of internal raw data handling, (x) an error analysis and justifiable correction strategies for errors inherent to the raw data as received by this study, (xi) some high-level summary statistics description of the observed data, (xii) justifications and results from conducting

various reasonable consistency checks, (xiii) reviews of reasonable biological

interpretations of typical observations, outliers, or other patterns of interest, (xiv) critical

assessments summarizing sufficient biological and other context to help readers

evaluate thoroughly, skeptically, and efficiently how much trust is justified by the quality

of the best and most complete dataset in this study, and (xv) any other potential

limitations. Since circadian clocks in flies have been an active area of research for some

time, a substantial number of studies report time series of potential interest. Time series

of clock components are foundational for understanding circadian clocks. Appropriately

integrating them raises various subtle issues that require decisions in order to build a

strong foundation for further studies. We initially underestimated the complexity of

dealing with their combined impact on the reproducibility of a steep data processing

pyramid that aims to eventually integrate parameter estimation and biologically

reasonable fly clock model ensembles. Since later steps such as parameter estimation

cannot correct quality problems at earlier steps, we decided to dedicate this study to

ensuring the availability of a durable high-quality set of time series observations ready

to serve beyond this study. Such efforts rival the complexity of wet-lab experiments,

except that they occur in a dry lab. The substantial investments in manual curation of

high quality datasets are justified by the well-known GIGO principle that applies to

simulations and experiments alike (Garbage In ➜ Garbage Out). Thus, we next present

some background on questions of basic reproducibility and data quality that arise for

integration efforts at the scale of our study.

Problems with label reproducibility. Irreproducibility can be caused by seemingly

trivial errors while executing deceptively simple work, such as pipetting errors (BROMAN

et al. 2015) or (mis)labeling a line of descent in the lab (LORSCH *et al.* 2014; FREEDMAN

et al. 2015b). Assigning sequence annotations in GeneOntology databases is neither

trivial nor always correct (JONES *et al.* 2007); incorrect assignments can replicate via

uninformed users and can also be generated easily by using spreadsheets with

inappropriate auto-conversion (ZEEBERG *et al.* 2004; ZIEMANN *et al.* 2016). Activities like

labeling or pipetting in array shaped micro-titer plates appear simple, but their simplicity

is deceptive because they involve naming – an often-underestimated problem of

extremely varying complexity (LOEWE *et al.* 2016b). The stakes are high and have led to

calls for systematically improving research at the bench and beyond (COLLINS AND TABAK

2014; LORSCH *et al.* 2014; ALLISON *et al.* 2016). The impact of these problems on our

study is immediate. We have little choice but to start with the assumption that *all errors*

had been corrected by the time of publication, implying correctness of all name-related

operations of all researchers involved in the production of the over 400 time series we

report below. Time of publication matters, as authors and readers might struggle to get

errors corrected if found later (ALLISON *et al.* 2016). The list of implications is long: no

accidental swaps anywhere, neither in any fly strain used throughout its relevant history

of descent, nor in the vials of final experiments, nor in the raw data, nor in averaging

repeated observations, nor in labeling the final plots for publication. We appreciate that

every single team of authors did their best to ensure that all errors were corrected in the

final publication. Also, label-errors in published time series of clock components are

probably less frequent than extrapolations of single-person initial error rates might

suggest (assuming scrutiny from co-authors and peer review). However, human error

analyses performed over decades in very diverse disciplines and for tasks of varying

complexity have quantified in numerous experiments that "*to err is human*" (PANKO

2016). Measured error rates observed in one type of experiment do not easily transfer

to other contexts, but the existence of labeling errors in labs is well documented

(LORSCH *et al.* 2014; BROMAN *et al.* 2015). Thus, it would be surprising if not a single

error existed in the published time series data we integrated. Equally, it would be

surprising if such a complex set of diffuse and scattered data could be integrated

without adding a single error from data handling. These observations highlight the

importance of assessing error rates and providing a defined protocol for reducing data

handling errors. Thus, high-quality data curation requires (in reverse order) mature

strategies for efficiently

- monitoring and handling all relevant error types,

- defining data structures that enable true data integration (and efficient querying)

- collecting all relevant scattered data in one place (and pre-sort for integration).

All this requires substantial efforts and biological model curators could probably learn

from the substantial methodologies for human error analysis that have been developed

elsewhere (NASA *et al.* 2006-07; NASA *et al.* 2011). Some of these approaches are too

complex for the application to individual studies in biology. However, meta-analyses

aiming to draw conclusions from noisy biological data need to find a way of handling the

errors that occur during data handling. They also have to address reproducibility in the

domain of statistics.

　　　Statistical reproducibility. A substantial fraction of recent problems with

reproducibility is caused by a lack of statistical reproducibility (AARTS *et al.* 2015;

HALSEY *et al.* 2015; STODDEN 2015). These problems easily arise while designing

experiments or analyzing data without the necessary statistical background (SALSBURG

1985; VAUX 2012). Here, interpretations of 'necessary' are the subject of much

discussion (STERNE 2003; CUMMING 2013; SHARPE 2013; LEEK AND PENG 2015) as

guidelines on statistical best practices are being updated (ALTMAN *et al.* 1983; MILLIS

2003; PLOWMAN 2008; MAZUMDAR *et al.* 2010; CUMMING 2013; DRUMMOND AND VOWLER

2013; JOHNSON 2013; CUMMING 2014; HUANG *et al.* 2015; SAVALEI AND DUNN 2015;

TRAFIMOW AND MARKS 2015; WOOLSTON 2015; TARONI *et al.* 2016; WASSERSTEIN AND

LAZAR 2016), but not necessarily followed everywhere (LEW 2012; TRESSOLDI *et al.*

2013). For example, a short-sighted over-reliance on *P*-values easily generates

irreproducible or misleading results (LEW 2012; CUMMING 2013; NUZZO 2014; HALSEY *et*

al. 2015; LEEK AND PENG 2015), a criticism with history (LOFTUS 1993; STERNE 2003).

Briefly, *P*-values are the probability that an observation can be explained by a given

null-hypothesis, which usually represents the 'most boring explanation'. Thus, *P*-values

are often seen as 'null-hypothesis significance tests', but they do not make any

statements about alternative hypotheses, of which there could be many. Yet

researchers often use *P*-values to draw unsafe conclusions of deceptive simplicity about

their respective favorite alternative hypotheses (LEW 2012; TRESSOLDI *et al.* 2013). They

do so with such regularity that this error's pervasiveness might one day motivate a

fascinating human error analysis. *P*-values may offer substantial attractions as they

combine the apparent reassurance of a precise number, the obvious simplicity of a

single dimension, and the clear choice between a boring and a seemingly interesting

option. In comparison, careful time-consuming analyses might be less appealing as they

often reveal complex ensembles of less-than-clear-cut alternatives in a world of multi-

dimensional trade-offs, requiring qualitative reasoning to decide which quantitative

methods to use for producing precise numbers. Such analyses offer more nuance,

albeit at greater cost and require more expertise in advanced statistics (WILCOX 2012),

and aspects of type systems (PIERCE 2002), semantics, and naming (LOEWE *et al.*

2016b). These complex analyses underscore a conclusion that is intuitively well

understood: biology does not present itself in a black and white picture of only

interesting or boring parts. Instead it offers not only shades (allowing for gradients in

addition to cutoffs at significance thresholds), but also colors (additional dimensions that

otherwise might be inappropriately collapsed into a single dimension). The recent

interest in statistical reproducibility has produced guidelines that recommend a closer

look at some of these additional dimensions by estimating confidence intervals and

other measures instead of testing arbitrary significance thresholds (KILLEEN 2005;

NAKAGAWA AND CUTHILL 2007; CURRAN-EVERETT 2009; CUMMING 2013; CUMMING 2014;

DEMIDENKO 2016). This does not mean that *P*-values have no merit (MURTAUGH 2014;

STANTON-GEDDES *et al.* 2014) and hence a pragmatic approach might be most

appropriate (BOOS AND STEFANSKI 2011), if the high variability of *P*-values is accounted

for (HALSEY *et al.* 2015). In either case, close attention to the robustness of statistical

methods is warranted (WILCOX 2012), and any statistical conclusions should be

supported by some analysis of their statistical reproducibility (HALSEY *et al.* 2015).

Finally, showing more raw data is preferable (LOFTUS 1993; DRUMMOND AND VOWLER

2011).

     Statistical error iceberg. Recent interest in statistical reproducibility has drawn

attention to other aspects of statistical analysis workflows. In this context, *P*-values have

been described as the tip of the iceberg (LEEK AND PENG 2015). To arrive at fully

rigorous conclusions requires investigating numerous detailed decisions about which

data to include, which outliers to remove, which tests to use, and which simplifying

assumptions to employ. Such analyses are more complex to produce and read, but they

are currently essential for exploring the most efficient approaches for arriving at reliable

statistical results. The fundamental nature of time series data for understanding clocks

in flies motivated us to invest in corresponding statistical reliability. Therefore, we

explore below several alternative ways to construct the statistical analysis pipeline for

this study.

Reproducibility in genetics. The recent surge of interest in reproducibility has

resulted in a number of studies of additional relevance to FlyClockbase. For example,

the reproducibility of genotype-phenotype associations has been investigated (NCI-

NHGRI WORKING GROUP ON REPLICATION IN ASSOCIATION STUDIES *et al.* 2007; IOANNIDIS

et al. 2009b; JANSSENS *et al.* 2009; KRAFT *et al.* 2009). Analysis of gene expression are

an important tool of genetic analysis and have therefore seen substantial standardizing

efforts (BAMMLER *et al.* 2005). Analysis of standardized and non-standardized

measurements have found improved reproducibility when standardized experimental

protocols were used (BAMMLER *et al.* 2005). For independent studies collected from the

literature, repeatability of microarray gene expression analyses has met limited success

(IOANNIDIS *et al.* 2009a). Reasons for failure included the unavailability of data,

incomplete annotations, and missing documentation on data processing (IOANNIDIS *et al.*

2009a). Other relevant observations that can hamper reproducibility include pipetting

errors (Bʀᴏᴍᴀɴ *et al.* 2015) and pedigree errors (Bʀᴏᴍᴀɴ 1999).

Versioned Biological Information Resources (VBIRs). The importance of

biological information resources is undisputed and has motivated the construction of

hundreds of heterogeneous resources as reviewed elsewhere (Bʀᴏᴏᴋsʙᴀɴᴋ *et al.* 2005;

Nɢ *et al.* 2006; Lᴀɪʙᴇ ᴀɴᴅ Lᴇ Nᴏᴠᴇʀᴇ 2007; Wɪᴇʀʟɪɴɢ *et al.* 2007; ᴠᴀɴ Gᴇɴᴅ ᴀɴᴅ Sɴᴏᴇᴘ

2008; Sᴜʟʟɪᴠᴀɴ *et al.* 2010; Dʀᴀɢᴇʀ ᴀɴᴅ Pᴀʟssᴏɴ 2014; Nᴀᴊᴀꜰɪ *et al.* 2014). Sizes vary,

as do scope and topics, ranging from general (e.g.

https://datascience.nih.gov/commons ; https://kbase.us), to organism specific (e.g.

http://flybase.org organized around *Drosophila* genomes (Gʀᴀᴍᴀᴛᴇs *et al.* 2016;

Mᴀʀʏɢᴏʟᴅ *et al.* 2016)), modeling specific (Lᴇ Nᴏᴠᴇʀᴇ *et al.* 2006; Cʜᴇʟʟɪᴀʜ *et al.*

2015), approach specific (Cᴜsɪᴄᴋ *et al.* 2009), and down to pathway or molecule specific

resources (e.g. ClotBase (Sᴏɴᴀᴡᴀɴɪ *et al.* 2010) or SwissLipids (Aɪᴍᴏ *et al.* 2015)).

Their heterogeneity remains a challenge and motivated development of the FAIR

Principles (Wɪʟᴋɪɴsᴏɴ *et al.* 2016). The FAIR Principles were designed for evaluating

credible solutions for the problem of exchanging data in biology and emphasize

important principles that make data sharing FAIR and data Findable, Accessible,

Interoperable, and Reusable (Wɪʟᴋɪɴsᴏɴ *et al.* 2016). The FAIR Principles do not aim to

provide any standards or implementations, but leave the actual development of

solutions to others, such as the proposed standard for Minimal Information Requested

In the Annotation of biochemical Models (MIRIAM) (Lᴀɪʙᴇ ᴀɴᴅ Lᴇ Nᴏᴠᴇʀᴇ 2007), Brief,

Explicit, Summarizing, Technical (BEST) Names (Lᴏᴇᴡᴇ *et al.* 2016b), and the many

proposals reviewed elsewhere (Dʀᴀɢᴇʀ ᴀɴᴅ Pᴀʟssᴏɴ 2014). Overall solutions will need

to solve the extraordinarily difficult challenge of data integration (DOAN *et al.* 2012).

Accordingly, efforts to exchange detailed data more efficiently in these complex

contexts have become top priorities in biological research contexts (NIH *et al.* 2012;

DRAGER AND PALSSON 2014; NIH 2015; NIH 2016; WILKINSON *et al.* 2016).

Importance of versioned data integrators. Versioned interoperable information

resources of intermediate size are likely to play a permanent role as hubs of integration

for the biological expertise in an area. The versioning is important to enable users to

access a stable state of information, without the unrealistic demand that these

resources have reached their final stage of development. Any *VBIR* worth developing

will likely be updated and improved for an extended period of time. While rates of such

change are likely to vary substantially, none of these changes should imperil the

reproducibility of some result that is based on the earlier state of the resourced as

accessed by the authors of that result. How to achieve interoperable long-term stable

versioning that is flexible enough to accommodate the broad range of needs of

resources as heterogeneous as *VBIR*s is an open question for research in the

semantics of naming. Currently, innumerable, incompatible, and inconsistent versioning

systems are actively used by numerous projects. Integrating them without reflection

would create a system that is almost incomprehensible and inflict on users intolerable

amounts of inessential complexity. Experience has shown that such a complex system

would be very brittle and would jeopardize reproducibility by its complexity. However,

the value of consistent easily reproducible integration is in the quality to which usable

resources offer expert curated relevant data that is continuously updated. Updates

could be triggered by detecting errors or integrating future experiments, possibly

expanding scope or precision through improved data models (but always increasing

some versioning number). Managing such updating processes works best for *VBIR*s of

some intermediate size. Thus, *VBIR*s size is defined at the lower end by exceeding the

limited scope of single publications, reviews, or meta-analyses that are all frozen in time

once completed. At their typical size, *VBIR*s enable the functional, ongoing integration

of information evaluating multiple studies and reviews from the perspective of a well-

defined scope. At the upper end, *VBIR*s generally remain at a much lower complexity

than grand challenges, and thereby avoid many additional complications caused by

their excessive complexity. This intermediate size and their stability enables *VBIR* to act

as reliable building blocks for accumulating biological expertise and address existing

grand challenges more efficiently.

Flexibility of VBIRs. These minimal constraints allow *VBIR*s to take on a great

diversity of organizational forms and the size of their scope may vary widely. However,

not all biological information resources currently perceived as useful do satisfy these

requirements. For example, not all repositories and databases in biology that aim to

continually integrate information provide reliable access to clearly defined states in the

past that are easy to access and to cite. Traditionally, the biological information

resources that are easiest to cite are journal articles, but these do not usually provide

information in a form that is structured enough for further processing and they do not

usually update information (ALLISON *et al.* 2016). Some authors complement their

articles with online databases or more static resources that can contain valuable

material. However, the lack of standards and tools that are easy to use means that such

efforts usually require substantial programming and data science expertise when they

                     QQv1r1

are set up and when they are to be maintained. Such barriers of entry make it very difficult for non-programming biologists with interesting datasets to set up and publish a VBIR in a form that facilitates further data integration.

Database integration is a special case of data integration. Both are enormous general challenges, whenever non-trivial datasets are to be used together (DOAN *et al.* 2012). For example, a substantial research collaboration worked towards integrating data scattered across 81 geospatial temporal ecology datasets from 7 provider types in an effort to build LAGOS, the LAke multi-scaled GeOSpatial & temporal database (SORANNO *et al.* 2015). The substantial supporting online material of the initial LAGOS description (SORANNO *et al.* 2015) provides an impression of the numerous data-handling and type system synchronization challenges the LAGOS team had to face in order to obtain some state of data integration (see http://csilimno.cse.msu.edu/lagos_status.php for updates).

Cochrane reviews provide a completely different approach to data integration. To reduce arbitrary bias that easily arises in more limited non-systematic reviews, http://www.cochranelibrary.com aims to stimulate the use of systematic methods for finding and integrating all peer-reviewed information about a given topic. The resulting retrospective and prospective meta-analyses have substantially advanced the integration of biomedical observations. Ongoing development of methodologies for systematic bias reduction has greatly increased awareness and approaches available for reducing the influence of important biasing factors (THARYAN 1998; ADES *et al.* 2008; MCKENZIE *et al.* 2013; DOI 2014; ONITILO 2014; DEBRAY *et al.* 2015; EFTHIMIOU *et al.* 2016). This success does not imply that further improvements are impossible; there are

biases that are notoriously difficult to address, such as certain types of ascertainment

bias (AMOS *et al.* 2003; CLARK *et al.* 2005; LACHANCE AND TISHKOFF 2013; MINIKEL *et al.*

2014) and biases against the publication of negative results (JOHNSON AND DICKERSIN

2007). In fact, it can be argued that biases are almost everywhere (e.g.: IOANNIDIS

2005a; PATIL *et al.* 2015). Thus, quantifying biases and analyzing their impact

appropriately may be more important than demanding their absence. Cochrane reviews

are not VBIRs, because traditional publications cannot be updated regularly. If VBIRs

could be published easily, without the currently required database programming

overheads, then Cochrane reviews with a reasonably well defined data model could in

principle become VBIRs.

Genome projects as a model for VBIRs development on a broader scale.

*VBIR*s increase the speed of hypothesis testing and greatly add to the long-term value

of properly annotated wet-lab data. They offer the raw material for diverse meta-

analyses, opening up entirely new research perspectives. In this respect, *VBIR*s mirror

similar efficiencies known from genome projects. The similarities do not end here.

*VBIR*s development also shows similar strong dependencies on software and data

organization. Genome projects demonstrated efficiencies of scale by separating data

collection and various stages of data interpretation (LANDER *et al.* 2001; VENTER *et al.*

2001). The efficiency of post-genomic sequencing workflows critically depends on the

development of appropriate data structures, processing tools and exchange protocols

(WILKINSON *et al.* 2016). We expect similar boosts to efficiency from developing *VBIR*-

tools. For example, they could support more biology-friendly data structures to increase

the efficiency and precision of integrating inherently imprecise biological observations.

They could also greatly accelerate the adoption of sophisticated statistical analysis by

the biological community, simply by implementing the appropriate statistical methods

into the corresponding automated workflows. This would reduce problems of confusing

SD and SEM (SALSBURG 1985), and opens up a new avenue for communicating

recommended best practice for statistical analyses, that could be provided right next to

a user-friendly implementation (MAZUMDAR *et al.* 2010; SHARPE 2013).

Genome projects have revolutionized biology. Here we want to explore whether

efficiencies of scale in biological model curation organized in a VBIRs project might hold

a similar potential for accelerating the pace of biological discovery.

The success of genome projects and other targeted efforts has been built on

efficiently organizing and exchanging new data and interpretations (WILKINSON *et al.*

2016). For VBIRs exchanging data is more complicated, because their data is more

structured and more diverse than typical genome data. Accordingly, efforts to exchange

details data more efficiently in these complex contexts have become top priorities (NIH

et al. 2012; DRAGER AND PALSSON 2014; NIH 2015; NIH 2016; WILKINSON *et al.* 2016).

Such work is essential for progress towards meeting increasingly complex grand

challenges like personalizing medicine, constructing genotype-phenotype-maps, or

predicting how cancer cell populations evolve using mechanistic models in evolutionary

systems biology (LOEWE 2016). By definition, these grand challenges all exceed the

problem-solving skills of any single research unit, and therefore critically depend on the

efficient communication of the latest progress. This progress could be captured in high-

quality *VBIR*s that probably will bring the same efficiency benefits as genome projects,

once corresponding tools become available for dealing with their more diverse data

types.

Importance of biological model curation. None of the benefits above can be

realized without substantial human input in the form of biological model curation.

Despite many advances in machine learning, the gold standard for data curation is still

the eye of a domain expert. While machines are extraordinary in exploiting regular

patterns, it has been difficult to teach machines how to correctly handle the many

exceptions that are readily recognized by human experts as deviations from 'common

sense' (BURKHARDT *et al.* 2006; SALIMI AND VITA 2006). Biological model curation by itself

is not new. In fact, one could argue that very essence of research is the construction

and curation of biological expertise that could also be described as a model. Thus, the

curation of biological information is at least as old as Linnaean taxonomy (LINNÉ 1758).

None of the recent resources that systems biology depends on could have been put

together without substantial model curation efforts (see e.g. DRAGER AND PALSSON

2014). The work of biological model curators, or biocurators, has only recently come

into focus as an increasingly important avenue of biological research (BOURNE AND

MCENTYRE 2006; BURKHARDT *et al.* 2006; SALIMI AND VITA 2006; HOWE *et al.* 2008; ST

PIERRE AND MCQUILTON 2009; BATEMAN 2010; BURGE *et al.* 2012; HIRSCHMAN *et al.* 2012;

ZHANG *et al.* 2014b; MITCHELL *et al.* 2015; ORCHARD AND HERMJAKOB 2015; RODRIGUEZ-

ESTEBAN 2015; GIBSON *et al.* 2016; KIM *et al.* 2016; REISER *et al.* 2016; SINGHAL *et al.*

2016). By now biocurators have an international society (http://biocuration.org) and an

official journal (http://database.oxfordjournals.org). Funding for digital depositories in

biology has historically been complicated because few have realized the essential

contributions of biological model curators to the overall scientific enterprise. This

problem has been recognized and efforts are underway to address this discrepancy

(EMBER *et al.* 2013). One potential contribution to these efforts could be to find a way

that substantially reduces the cost of initiating, growing, and maintaining *VBIR*s. We

explore a potential approach for simplifying model curation by exploiting advances in

computer science that have greatly simplified the design and construction of compliers

for new programming languages.

How a compiler could help in biological model curation. Programing

language compliers are extraordinary efficient tools for guaranteeing that a given

collection of texts (source code) conform to a well-defined standard (the complier's

language) and are transformed into output that is guaranteed to conform to strict rules.

The construction of compliers requires an advanced understanding of computer

science, but decades of research have produced a substantial body of data structures,

algorithms, and tools that greatly simplify the construction of compilers today (e.g.

COOPER 2012; GRUNE 2012). Thus, the question today is not if a compiler can be

constructed for a given language, but rather what should a language look like for which

it is worth constructing a compiler. After the construction of uncounted programing

languages each with their own strengths and weaknesses, rather compelling reasons

are necessary for creating a new one and any such efforts should learn from the diverse

shortcomings of their many predecessors (MANDRIOLI AND PRADELLA 2015). Traditional

approaches to designing new programming languages have not included the very large

amounts of feedback from research biologists that are necessary for creating a

language design that could efficiently support biologists in their work (LOEWE *et al.*

2016b).

   Enforce best practices. For example, such a compiler could greatly reduce the

confusion between SD and SEM that has plagued the reporting of biological results for

some time (SALSBURG 1985). A compiler that implements the latest statistical testing

methods could greatly improve the adoption of statistical best practices in the biological

research community (MAZUMDAR *et al.* 2010; SHARPE 2013). Such a compiler could also

advance standards for facilitating interoperability in systems biology (DRAGER AND

PALSSON 2014) and thereby contribute towards solving the extraordinarily difficult

challenge of data integration (DOAN *et al.* 2012), improve the semantic reproducibility of

biological data (LOEWE *et al.* 2016b), facilitate the sharing of meaningful data based on

the FAIR Principles (WILKINSON *et al.* 2016), and encourage biologists to provide the

Minimal Information Requested In the Annotation of biochemical Models (MIRIAM)

(LAIBE AND LE NOVERE 2007). VBIRs construction does not require the existence of such

a compiler, as every task can also be performed manually. Manual work is slower, but

also more flexible, and can therefore better attend to the needs of high-quality biological

model curation of a given data set. If such curation work is combined with the

perspective of a compiler architect, then it provides extraordinary opportunities for

designing efficient abstractions for data structures and tasks that can later be supported

by a fully automated compiler.

Efficiencies of scale. Many VBIRs are likely to have similar needs that can be served by the same compiler if they share a standard for storing data. Thus, the costs of compiler development can then benefit several *VBIR*s where they reduce the cost of *VBIR* development and maintenance, which have been difficult to fund (EMBER *et al.* 2013). Compiler development is also an excellent opportunity for detecting problems in logic formalisms; such errors have the potential for causing exorbitant costs (e.g. HOARE 2009; KAMP 2011; LOEWE *et al.* 2016b). Therefore, constructing such a compiler could already be a cost-effective decision for the longer-term development and maintenance of a single VBIR alone. We aim to observe during our biological research where commonly used logic formalisms made it more complicated to accurately represent biological observations with their usual uncertainty. Representing uncertain biological data in computational structures is a sufficiently frequent problem to cause substantial frustration in efforts towards curating biological models at a reasonably high quality.

Opportunity. Our systematic study of circadian clock gene expression patterns offers intriguing opportunities for engaging with the timely questions of reliable data handling, control experiment repeatability, human error analysis, reproducible computing, statistical reproducibility, and the semantic reproducibility of source code in research computing. These questions might be easy to dismiss at first sight, but as discussed above, in the broader context of growing data sets in VBIRs, low rates of diverse and individually rare errors can combine into a pervasive fog of confusion that can render a valuable collection of scientific results unusable. Usually these problems cannot be investigated at the level of a single experimental study, but this does not

imply that errors in *VBIR*s are rare, or without consequence (e.g. see (ZEEBERG *et al.* 2004; JONES *et al.* 2007; SCHNOES *et al.* 2009)). The resulting irreproducibility is not cheap. For example, non-clinical biomedical studies with an estimated cost of about $7Bn/yr throughout the US come with difficulties in data analysis and reporting that hamper their reproducibility (FREEDMAN *et al.* 2015a). Pervasive biases in biological datasets (IOANNIDIS 2005a) and statistical difficulties that can lead to substantially wrong conclusions (IOANNIDIS 2005b) can interfere with scientific discovery. To address these problems, it is important to invest in efforts towards opening science (BARTLING AND FRIESIKE 2014), sharing data (PACKER 2016; WILKINSON *et al.* 2016), and improving reproducibility in various areas (IOANNIDIS 2005b; DONOHO 2009; HUANG AND GOTTARDO 2013; LOEWE AND KEEL 2014; STODDEN *et al.* 2014; FREEDMAN *et al.* 2015a; JAMES *et al.* 2015; STODDEN 2015; BARBA 2016; LOEWE *et al.* 2016a; LOEWE *et al.* 2016b). Reproducibility frameworks greatly facilitate individual scientific research studies, but take much more effort to put into place than could possibly be expected of the investigators in any individual study. Therefore, it is important to find efficient ways to achieve these goals at institutional and national scales (NIH *et al.* 2012; NIH 2015; NIH 2016). Our study is different from many typical studies in that we have attempted to simultaneously conduct high quality research while working towards a framework for improving reproducibility.

Purpose of this study. Here we interweave several perspectives integral to one body of trans-disciplinary research. We aim to improve amount and quality of experimental time series available for parameter estimation in mechanistic simulations

                    QQv1r1

along with our overall understanding of *D. melanogaster* circadian clock models. To this

end we present FlyClockbase, a new carefully curated biological information resource

designed to maximize accessibility and ease of use for experimental biologists and

modelers. We show how to use it for testing hypotheses and report our own new

findings about the variability of peak times in the clock. Finally, we make trans-

disciplinary observations at the interface of experimental biology, data curation,

reproducibility, and the applicability of logic formalisms in biology. We present our

process for working towards constructing a compiler that would substantially reduce the

effort required for developing and maintaining data resources like FlyClockbase. These

perspectives are next explained in a more detailed overview.

Formal organization of Versioned Biological Information Resources (VBIRs).

FlyClockbase is a *VBIR*. *VBIR*s store biological information using controlled immutable

versioning numbers for marking each publicly released variant of the resource to ensure

that previously released data remains accessible under that number. As we developed

FlyClockbase, we aimed to separate the special from the general *VBIR* aspects to help

make our design more applicable to the development of future *VBIR*s. Below we

discuss why more *VBIR*s are needed. For ease of use, quality control, and future

maintainability, we designed FlyClockbase as a file-based data repository that follows a

well-defined scheme for collecting tables in text files grouped in folders. This

organization as a set of tables is conceptually similar to the structure of a relational

database, albeit without the speed, rule enforcement, and other amenities provided by

modern database systems. As a result, our approach maximizes flexibility and

openness, while minimizing certain types of administrational and long-term maintenance

costs (to increase chances of long-term survival; see Discussion). The resulting system

is even more general than *VBIR*s and we nicknamed it 'TabFS'. The name TabFS

highlights the central role of tabs (for delimiting), table-files (for storing) and the file-

system (for organizing data). To simplify the implementation of other *VBIR*s, we have

been separating the specific details of implementing FlyClockbase from general abstract

features. In TabFS we aim to capture the abstractions and rules required for

implementing *VBIR*s with a long-term view to developing a reliable *VBIR* standard. Our

goal is to provide the simplest and most efficient organization possible without

sacrificing the flexibility curators need for defining new data types that represent diverse

types of complex and uncertain biological observations. Efficiently integrating new

datasets in FlyClockbase requires this flexibility. It facilitates focusing on clock biology

and minimizes distractions from defining or decoding data types. This approach enables

FlyClockbase to integrate a diverse array of wildtype and wildtype-like time series along

with the attributes necessary for documenting a broad range of experimental details.

Each time series records the relative amount of a clock component as observed at

various points in time.

Data integrated by curation. In FlyClockbase we provide a curated overview of 25

years of published observations of *D. melanogaster* clock components, which we use

for retrospective meta-analyses. The types of molecules reported in FlyClockbase are

based on the biological *D. melanogaster* clock model we abstracted from the relevant

literature (see Table 1 for brief descriptions of core clock components). FlyClockbase

contains more than 400 time series curated from the wildtype control experiments of 86

circadian clock studies. They can be compared in many ways within or between clock

components for testing diverse hypotheses of potential interest for fly clock research.

Hypothesis tested: biological variability. We use FlyClockbase for comparing the

variance of times at which the circadian mRNAs and proteins *period* (*per*) and *timeless*

(*tim*) reach their relative daily peak and valley. We find significant differences in

variance that are not easily explained as a statistical fluke and survived several rounds

of in-depth error checking (which led to interesting conclusions in their own right). Thus,

we hypothesize that the larger variance of peak times for the protein PER in comparison

to TIM might have mechanistic reasons that could help illuminate interesting aspects of

the clock if recovered in mechanistic models.

Hypothesis tested: observation method. The confluence of many diverse

independently observed time series in FlyClockbase provides a unique resource for

understanding such variability of fly clocks in a broad range of settings, as documented

in the attributes of FlyClockbase time series. This variability can also be used for

comparing the reproducibility of different approaches to measuring time series. We

compared time series measured by PCR based methods (qPCR, RT-PCR) with

methods that do not include self-replication (Northern Blot, RNAse Protection Assay).

The variability of PCR-based time series in FlyClockbase exceeds that of non-PCR

based methods; though originally surprising, this is consistent with both the exponential

nature of amplification in PCR and previous reports on the reproducibility of quantitative

measurements from PCR-based methods.

Human error analysis. We measured human error rates for a given set of tasks in

FlyClockbase. Our results are broadly comparable to previous observations. The

findings suggest that VBIRs would benefit from developing methods for ensuring that

scientific conclusions are not affected by human errors that inevitably occur when

handling or analyzing data and corrupt content or type. Designing a formal type system

capturing relevant expert insights for FlyClockbase could facilitate and ultimately

automate searches for logical inconsistencies.

Compiler logic design. We developed FlyClockbase while simultaneously

exploring design options for programming language compilers that could help construct

and maintain *VBIR*s. We have identified numerous pivotal features for supporting the

long-term stability of FlyClockbase that are most efficiently implemented by a

correspondingly designed compiler. We discuss how these and other practical aspects

of working with *VBIR*s can improve the usefulness and chances of longer-term survival

for *VBIR*s. We use an analogy to well-known results from population genetics to

illustrate what the future might hold for a newly-born *VBIR*, such as FlyClockbase.

These considerations show that cumulative practical impacts from many small

complications or innovations can be unexpectedly large. We illustrate using

FlyClockbase how it can be difficult to represent uncertain biological data in the

certainty-demanding logic formalisms of the data types commonly used in

computational tools. Most of the data types we need in FlyClockbase fall into two

categories. Some are very general data types that are very common and thus ideally

designed for interoperability (e.g. bibliographic references, tables, etc., see TabFS).

Other data types are specific to FlyClockbase and therefore are not reusable. These

types need to be defined by those expert biologists who curate FlyClockbase, and best

understand the relevant biology. To facilitate these discussions, we developed a trans-

disciplinary collaboration model to help with the necessary communication of biological

curators with a compiler architect, who needs to be capable of bridging biology and

aspects of designing compiler logic. If such a communication approach is used by a

compiler architect for informing important decisions about the relevant logic formalisms

to be implemented, then our observations suggest that the efficiency of biological model

curation could greatly increase once a user-friendly *VBIR* compiler becomes available.

Such a compiler will empower biological model curators to define their own data types

that can be shaped to more appropriately representing the uncertainty of the biological

observations they curate – without violating biological or computer science logic. Such

compilers will also allow biologists to define their own consistency checks, which can

then be automatically maintained by a correspondingly designed compiler. If this

compiler also implements sufficiently reviewed standards for interoperability and data

exchange, biological research will benefit from an unprecedented ability to combine

models and analyses from different VBIRs.

Importance of efficient biological model curation. Our work highlights why topic-

specific *VBIR*s like FlyClockbase have an essential, irreplaceable role to play in

biological research – once curated to high quality by expert biologists. As we illustrate in

FlyClockbase, *VBIR*s increase the speed of hypothesis testing and greatly add to the

long-term value of properly annotated wet-lab data. They offer the raw material for

diverse meta-analyses, opening up entirely new research perspectives. In this respect,

*VBIR*s mirror similar efficiencies previously observed in genome projects. Our trans-

disciplinary analysis suggests that the most efficient route for integrating biological

information requires more work on type systems and logic formalisms in order to better

capture the many uncertainties regularly found in biological data. To sample the

problem space well enough, more studies like ours are needed that report how in-depth

biological research challenges the expressivity of logic formalisms that have become

candidates for implementation in the discussed compiler. Combining such observations

with a rigorous in-depth usability and expert review process as defined elsewhere

(LOEWE *et al.* 2016b) will greatly accelerate the definition of more appropriate logic

formalisms, VBIRs compiler design, biological model curation, and thus progress

towards meeting the grand challenges of our time. This new efficiency would indeed

allow us to stand on the shoulders of giants and no longer have to start crawling

upwards from the elbow whenever a new question arises.

Overview of Sections. In the next Section, we review biological and computation

clock models, as well as the data model of FlyClockbase from a biological perspective.

We then describe how we selected the data in FlyClockbase, how we processed time

series and which statistical methods we used. Our Results Section first quantifies the

historic use of direct experimental time series observations in modeling studies. It then

reviews the number of time series observed for each core clock component in

Drosophila melanogaster. It provides an overview of the variability in all components

before presenting a human error analysis investigating potential impacts of data

handling errors in FlyClockbase on the variances of peak and valley timings of the

period and *timeless* gene products, which are compared after defining a basic null-

hypothesis for data in FlyClockbase. Our last result compares methods for observing

mRNA. We start our Discussion Section by explaining, how FlyClockbase facilitates

hypothesis-driven research. We then discuss the two hypotheses tested in this study

and suggest mechanistic models for further testing. Next we broaden our view to

discuss the importance of model curation for molecular systems biology data. We then

highlight observations that illustrate, how a tool with the capabilities of a specially

crafted programming language compiler could advance work in FlyClockbase and

beyond. We will pay attention to aspects like efficiency, error detection, and formal logic.

Given the high likelihood of long-term loss of biological information resources, we finally

discuss population genetics modeling results with some applicability to the fate of

FlyClockbase. We do so in order to prioritize and motivate the next steps. We conclude

with a list of the various disciplinary areas engaged by this study and how biological

model curation will facilitate critical progress towards various grand challenges of our

time. Our online material includes additional text with a more computational perspective

and a supplementary statistical analysis to which we frequently refer in our results

(including R source and data).

MODELS

Biological model of fly circadian clocks

The *D. melanogaster* clock is a gene regulatory network that receives environmental inputs (such as light and temperature) and produces its hallmark cyclical behavior through various interlocking positive and negative feedback loops (KUCZENSKI *et al.* 2007; WANG AND ZHOU 2010). Table 1 lists the most important clock components, and Figure 1 represents their key interactions in the Systems Biology Graphical Notation (MOODIE *et al.* 2011). The timing of various clock sub-processes is essential for any clock. Circadian clocks critically depend on generic cellular processes of importance in the information processing associated with proteins, such as transcription, translation, and degradation. Thus, mutations disrupting critical functions in these generic processes are also likely to affect the clock. However, they are also likely to have many other harmful consequences; hence, we do not consider them as *core* clock components (which are the exclusive focus of our study here).

Place   TABLE 1   about here.

Place   FIGURE 1   about here.

QQv1r1

Main loop. Briefly, in the core (negative) feedback loop, the proteins CLOCK

(CLK) and CYCLE (CYC) form a heterodimer and promote the transcription of *period*

and *timeless* (DARLINGTON *et al.* 1998; RUTILA *et al.* 1998). PER protein is increasingly

phosphorylated by DOUBLETIME (DBT) and several other kinases. Fully

phosphorylated PER interacts with the F-box protein SLIMB (SLMB) to be marked for

degradation unless TIM protein is present to form a PER/TIM complex. This complex

represses the effects of the transcriptional activator CLK to form a negative feedback

loop (GEKAKIS *et al.* 1995; KLOSS *et al.* 1998; LEE *et al.* 1998; PRICE *et al.* 1998; LEE *et

al.* 1999; KLOSS *et al.* 2001; CHIU *et al.* 2008). When light is present, the protein

CRYPTOCHROME (CRY) undergoes a conformational change that renders it active. As

a final step before TIM degradation, activated CRY and the kinase SHAGGY (SGG)

cause TIM in its phosphorylated form to interact with the F-box protein JETLAG (JET)

(MARTINEK *et al.* 2001; KOH *et al.* 2006). If TIM is degraded thought its phosphorylated

form, this will limit the formation of the PER/TIM complex. If this complex cannot form,

then PER will be left in an isolated form in which it can be further phosphorylated (and

thus be moved closer to its degradation). If PER pairs with TIM to form this complex,

then PER cannot be phosphorylated, and it will temporarily stop its progress towards

degradation. Thus degrading TIM facilitates the degradation of PER by allowing PER to

become fully phosphorylated (EMERY *et al.* 1998; NAIDOO *et al.* 1999; BUSZA *et al.* 2004;

OZTURK *et al.* 2011). Consequently, PER/TIM complexes no longer repress CLK

transcriptional activity, and CLK proceeds to start a new cycle by again promoting again

the transcription of *per* and *tim*.

Other loops. A second feedback loop primarily concerns *clk* transcription. CLK promotes transcription of both *PAR-domain protein 1 (pdp1)* and *vrille (vri)* (BLAU 1999; MCDONALD AND ROSBASH 2001; CYRAN *et al.* 2003). PDP1 protein then promotes the transcription of *clk*, while VRI represses the activity of PDP1 and inhibits the transcription of *clk*, creating positive and negative feedbacks, respectively (CYRAN *et al.* 2003; GLOSSOP *et al.* 2003). The relatively recently discovered *clockwork orange* (*cwo*) modulates both feedback loops by weakly repressing CLK-mediated transcription of *per*, *tim*, *pdp1*, *vri*, and *cwo* itself (KADENER *et al.* 2007; LIM *et al.* 2007; MATSUMOTO *et al.* 2007; RICHIER *et al.* 2008). The interplay between strong transcriptional activation promoted by CLK and weak repression from CWO protein counteracts "jitters," or small variations in period (FATHALLAH-SHAYKH *et al.* 2009; FATHALLAH-SHAYKH 2010; SCRIBNER AND FATHALLAH-SHAYKH 2011). Other notable circadian products involved with post-translational modification, synchronization of clock neurons, and other processes include *casein kinase 2 alpha (ck2a), protein phosphatase 2a (pp2a)*, *pigment-dispersing factor (pdf)*, *nemo (nmo),* and others (GRIMA *et al.* 2002; KO *et al.* 2002; LIN *et al.* 2002a; SATHYANARAYANAN *et al.* 2004). A more detailed review of the clock can be found elsewhere (HARDIN 2011; ÖZKAYA AND ROSATO 2012).

              QQv1r1

In silico models integrating fly clock observations

Mathematical models of circadian clocks have been contributing to our understanding of clock biology for decades.

Biological results overview. The origins of many fly clock models can be traced back over 50 years to work by PITTENDRIGH & VICTOR (PITTENDRIGH AND VICTOR 1957; GOODWIN 1964; GOODWIN 1965) and GOODWIN (PITTENDRIGH AND VICTOR 1957; GOODWIN 1964; GOODWIN 1965). Then GOLDBETER (1995) developed a model with five ordinary differential equations that use *per* mRNA and PER protein (in various phosphorylation states) to describe a negative feedback loop created when PER represses *per* mRNA transcription. LELOUP (1998a) expanded this model to include *tim* mRNA and TIM protein. Later models such as those published by UEDA (2001) and SMOLEN (2001) again expanded the feedback loops by adding CLK, and more recent models added a feedback loop based on *vri, pdp1* (SMOLEN *et al.* 2004; XIE AND KULASIRI 2007; KULASIRI AND XIE 2008) and CWO (FATHALLAH-SHAYKH *et al.* 2009). Two other common points of interest for clock models were the importance of a positive feedback loop (TYSON *et al.* 1999; SMOLEN *et al.* 2001; KUCZENSKI *et al.* 2007; WANG AND ZHOU 2010) and the influence of post-translational modifications such as phosphorylation (LEISE AND MOIN 2007; RISAU-GUSMAN AND GLEISER 2012). Because the clock is able to adjust to temperature variations, a number of models further investigated the role of temperature on the clock (KIDD *et al.* 2015). Some temperature models, such as those by HONG

(HONG AND TYSON 1997) and LELOUP (LELOUP AND GOLDBETER 1997), are based on

GOLDBETER'S 1995 model, while others (RUOFF AND RENSING 1996; RUOFF *et al.* 1997;

RUOFF *et al.* 1999) use the more general GOODWIN (1965) oscillator as a foundation.

While models have become sophisticated enough to explain observable biological

phenomena, methods have improved for choosing more realistic values for model

parameters (KURATA *et al.* 2007; XIE *et al.* 2010; LEBIEDZ *et al.* 2012). A number of

review articles compare the various clock models and present more details on their

history (SMOLEN *et al.* 2000b; SMOLEN *et al.* 2000a; GOLDBETER 2002; KUROSAWA *et al.*

2002; OGAWA *et al.* 2008; GERARD *et al.* 2009; LELOUP 2009; GONZE 2011; SCRIBNER AND

FATHALLAH-SHAYKH 2011).

Role of stochasticity. Researchers have also been using increasingly

sophisticated computational approaches for simulating the clock. Many early models

were constructed using deterministic ordinary differential equations, but some of their

underlying assumptions are not always applicable to the clock. In particular,

deterministic models assume large enough numbers of molecules so that any random

variations caused by stochastic changes in the state of individual clock components are

compensated at the level of the whole clock. Such large numbers of molecules may not

be realistic for clocks at the cellular level (RUOFF *et al.* 1999). More recently, stochastic

models have been constructed to overcome these limitations; these models are often

derived from previously published deterministic equivalents (BARKAI AND LEIBLER 2000;

ZAK *et al.* 2001; GONZE *et al.* 2002a; GONZE *et al.* 2002b; UEDA *et al.* 2002; VILAR *et al.*

2002; GONZE *et al.* 2003; GONZE *et al.* 2004; MIURA *et al.* 2008). This allowed for a better

understanding of how intracellular stochasticity generates noisy clock observations (BARKAI AND LEIBLER 2000; GONZE *et al.* 2002b; YI *et al.* 2006; LI AND LANG 2008; LERNER *et al.* 2015) and how this noise can be reduced by synchronizing clocks across groups of neurons (KATAKURA AND OHMORI 2006; BAGHERI *et al.* 2007; BAGHERI *et al.* 2008b; DIAMBRA AND MALTA 2012; RISAU-GUSMAN AND GLEISER 2014).

Shared problems. The models above examine different aspects of the clock, but they all face two common modeling challenges: estimating parameters and testing the quality of models. Both issues require the ability to access and use high-quality experimental data, yet there is a painful lack of experimentally measured rate parameters for important circadian clock processes. Researchers have used a wide range of methods to find potentially realistic parameters (LELOUP AND GOLDBETER 1998a; LELOUP AND GOLDBETER 2000; SMOLEN *et al.* 2001; SMOLEN *et al.* 2002; SMOLEN *et al.* 2004; RUOFF *et al.* 2005; XIE AND KULASIRI 2007; BAGHERI *et al.* 2008a; KULASIRI AND XIE 2008; WANG AND ZHOU 2010). These include trial-and-error approaches, but even the best systematic methods cannot guarantee finding rates that reflect nature's values. In principle, such searches aim to find combinations of input parameter values that cause models to produce time series mirroring those observed experimentally. This ideally provides ensembles of realistic parameter combinations that cannot be ruled out by experimental evidence. It is immaterial, whether such ensembles were generated by testing deterministically or stochastically proposed parameter combinations. However, in no case is it possible to "validate" any parameter combination on principal grounds due to the open nature of the models as discussed elsewhere (ORESKES *et al.* 1994;

TARANTOLA 2006). The best parameter estimates are thus "realistic" (up to a given

stringency), and conclusions based on simulations using them are reasonable (up to the

usually unknown degree to which these parameter combinations represent reality). This

indirect approach has been successful in a wide range of disciplines for estimating

parameters in complex models (e.g. (STAINFORTH *et al.* 2005)). The remaining

"unknown" degree can be narrowed for a given model with unknown parameters by

using statistically rigorous approaches (TARANTOLA AND VALETTE 1982; JAYNES AND

BRETTHORST 2003; TARANTOLA 2005; TARANTOLA 2006; MOURA NETO AND SILVA NETO

2013). The problem of unknown model parameters is widespread in many disciplines

that use modeling approaches and has also become known as the "inverse problem"; it

can be can be solved in principle by probability theory (TARANTOLA AND VALETTE 1982;

JAYNES AND BRETTHORST 2003; TARANTOLA 2005; TARANTOLA 2006). Concisely stated,

the inverse problem is the challenge to use all known data about a system for restricting

the ranges of unknown causal input factors for a model that produces simulation output

that is equivalent to data observed in the real system itself (even though the latter

ultimately remains unknown). Solving the inverse problem for increasingly realistic

biological models using growing datasets of varying quality quickly exceeds current

mathematical and computational capabilities and thus remains a research challenge.

Additional limits for such reverse-engineering of systems biology models may come

from the large variability of their kinetic rates (ERGULER AND STUMPF 2011).

Parameter estimation in complex models. Numerous algorithms can propose

sequences of input parameter combinations that repeatedly reduce computed distances

between simulated and observed data. However, few frameworks can rigorously

estimate the statistical uncertainty associated with their point estimates. Maximum

likelihood and Bayesian statistics are currently the frameworks that are most advanced

(EDWARDS 1992; HEYDE 1997; JAYNES AND BRETTHORST 2003; BISHOP 2006). They

usually require a function that directly computes the likelihood that a given system will

produce a given set of observations for a given set of input parameters. However, this

likelihood function is increasingly difficult to specify for non-linear stochastic models of

growing complexity such as the circadian clock models of interest. Help may come from

formalized frameworks for Approximate Bayesian Computation (ABC) that have been

developed in various disciplines and do not require explicit likelihood functions (TONI *et*

al. 2009a; CSILLERY *et al.* 2010a; ROBERT *et al.* 2011; SUNNAKER *et al.* 2013; WILKINSON

2013; LEE *et al.* 2014b; STUMPF 2014; BUZBAS AND ROSENBERG 2015). In theory, ABC

can solve inverse problems for all simulation models capable of producing output that is

comparable to real observations. Briefly, ABC approximates likelihoods by (i) proposing

new potentially realistic input parameters, (ii) simulating the model to predict

corresponding results, (iii) calculating the distance of these results to experimentally

observed data, and (iv) deciding which input parameters are actually supported by

experimental evidence, based on comparing these distances to predetermined

acceptance criteria. Thus, ABC generates ensembles of model variants, which describe

sets of biologically realistic parameter combinations that quantify the uncertainty

associated with a given model in the light of available data. Recent progress on

uncertainty quantification via ensemble analysis has been reviewed by BAUER *et al.*

(BAUER *et al.* 2015). The accuracy of such ensembles depends on the quality of

distance measures, acceptance criteria, and sampling density in relevant regions of

parameter space. The statistical, numerical and computational challenges associated

with ABC increase with model complexity and data diversity. In practice, sampling

speed is often limiting, and distances to observed data might have to rely on summary

statistics that can create complicated biases. These biases will matter when models

with different structures are compared and these summary statistics do not capture all

information that is relevant for fully evaluating the models (ROBERT *et al.* 2011).

Fortunately, these problems can be solved for simulations of biochemical systems (TONI

et al. 2009a; ROBERT *et al.* 2011). Generally, ABC benefits from access to raw

experimental observations such as time series to maximize the information used to

estimate parameters and minimize bias from incompletely processed data. Estimating

parameters using ABC in fly clock models of realistic complexity is very challenging and

has not yet been attempted (to the best of our knowledge; recent work in Neurospora

clocks (DENG *et al.* 2016) demonstrates some of the challenges). While the full potential

of ABC still waits to be realized, the clock modeling community has estimated

parameters either by fitting model output to abstract time series traits or by using time

series data more directly. These approaches are discussed next.

Using abstract time series traits. The use of higher-level abstractions of time

series traits can greatly simplify assessing the realism of a given circadian clock model,

at least when compared to data-intensive work with raw time series. Despite the very

general nature of abstract traits (e.g. "oscillates" or "has feedback loops"), they can

provide powerful filters for removing biologically uninteresting parameter combinations

when analyzing circadian clocks. More specific examples of such abstract traits are:

1. period close to 24 hours or modified as observed in mutants (GOLDBETER 1995;

    LELOUP AND GOLDBETER 1998a; LELOUP AND GOLDBETER 1998b; ROENNEBERG AND

    MERROW 1998; LELOUP AND GOLDBETER 2000; UEDA *et al.* 2001; FATHALLAH-

    SHAYKH *et al.* 2009; WANG AND ZHOU 2010; RISAU-GUSMAN AND GLEISER 2012),

2. phase changes based on light exposure (ROENNEBERG AND MERROW 1998;

    SMOLEN *et al.* 2004),

3. ability to account for responses to light, including light pulses (ROENNEBERG AND

    MERROW 1998; TYSON *et al.* 1999; LELOUP AND GOLDBETER 2000; LELOUP AND

    GOLDBETER 2001; PETRI AND STENGL 2001; SMOLEN *et al.* 2001; SMOLEN *et al.*

    2002; SMOLEN *et al.* 2004; RUOFF *et al.* 2005; BAGHERI *et al.* 2008a; FATHALLAH-

    SHAYKH *et al.* 2009)

4. ability to be properly entrained (SMOLEN *et al.* 2001; SMOLEN *et al.* 2002),

5. robustness to small parameter changes (SMOLEN *et al.* 2001; SMOLEN 2002;

    SMOLEN *et al.* 2004),

6. ability to replicate the behavior of mutants (ROENNEBERG AND MERROW 1998;

    TYSON *et al.* 1999; SMOLEN *et al.* 2004; RUOFF *et al.* 2005; BAGHERI *et al.* 2008a;

    FATHALLAH-SHAYKH *et al.* 2009; RISAU-GUSMAN AND GLEISER 2012),

7. delay between the peaks of a given mRNA and its protein (LELOUP AND

    GOLDBETER 1998b; SCHEPER *et al.* 1999a; SCHEPER *et al.* 1999b; SMOLEN *et al.*

    2001; SMOLEN *et al.* 2002; XIE AND KULASIRI 2007; WANG AND ZHOU 2010; RISAU-

    GUSMAN AND GLEISER 2012),

8. dynamics of the combined amounts of all forms of PER protein (PETRI AND STENGL 2001; SMOLEN *et al.* 2004), and

9. time at peak expression of a given clock component (PETRI AND STENGL 2001; FATHALLAH-SHAYKH *et al.* 2009).

Comparing simulation results and observed values for such abstract clock traits is generally easier than comparing simulations with complex experimental data. However, it unclear how much information about circadian clocks is preserved and how much is biased or lost when reducing all relevant circadian clock output to the abstract measures given here.

Using complete observed time series. To reduce these uncontrollable biases from using abstract traits, researchers might seek to incorporate all available time series data in more direct tests to compare the distance between observed and simulated time series. The core idea is to increase statistical power by including as much information as possible when estimating parameters and to use repeated observations to obtain better estimates of the underlying distributions. It is therefore desirable to integrate all observed time series of clocks in flies in a single biological information resource. More time series also facilitate recognizing genuine clock signals among the experimental noise inevitably associated with all biological observations and thus help avoid overfitting. In practice, creating models based on experimental time series is complicated by diverse challenges:

(i) All challenges of inverse problems with many dimensions discussed above are (or seem) exacerbated because time series usually provide more degrees of

freedom than lower-dimensional summaries of their features. Parameter

estimation is particularly difficult due to the complex, non-linear relationships

between clock components (TARANTOLA AND VALETTE 1982; FORGER *et al.* 2005).

While recent progress in developing statistical frameworks like ABC is

encouraging, these are not straightforward to use for models with more than a

dozen unknown parameters (TONI *et al.* 2009b; CSILLERY *et al.* 2010b;

SOUBEYRAND *et al.* 2013; SUNNAKER *et al.* 2013; WU *et al.* 2014). Exploring such

techniques is beyond the scope of this paper; here we aim to present a real-

world, research-grade dataset that provides a non-trivial versatility and

complexity test for candidate methods.

(ii) It can be difficult to choose optimal measures for comparing time series,

regardless of whether applied to simulations or experimental data. There are

many "standard measures" for comparing time series in general (e.g. the

Euclidian distance, equivalent to the assumption that amounts follow a Normal

distribution), and circadian clock time series in particular (e.g. period length).

Selecting one or more appropriate measures is not trivial (GLYNN *et al.* 2006;

REFINETTI *et al.* 2007; DING *et al.* 2008; BATISTA *et al.* 2011; JIN 2011; SUN *et al.*

2014; YIN *et al.* 2014; BANKO AND ABONYI 2015; KOTSIFAKOS *et al.* 2016; MORI *et*

al. 2016). This is particularly true when comparing time series with differently

calibrated, non-linear scales that may be associated with substantial

measurement errors, as is often the case for experimental observations. Thus,

many diverse quantitative methods can be used to calculate diverse measures of

distance between time series. However, this does not solve the substantial

qualitative need for arguing which quantitative approaches are appropriate, if any.

(iii) It is challenging to compile all relevant time series observations into one place and organize them in a uniformly accessible manner, as pertinent time series were observed using different methods in diverse contexts and span a rich body of literature across many years. Furthermore, data processing is complicated by many obstacles associated with scattered big data, which characterizes many types of biological information. Practical challenges include the degrees to which

a. data is rarely compiled in a uniform, directly usable data format,

b. different datasets require diverse manual corrections for special cases that are individually rare but aggregately comprise a substantial part of the data and are hence not ignorable,

c. time series data is always incomplete and gaps between observed time points are irregular,

d. data is often insufficiently documented such that it becomes impossible to determine essential information about the precise types and attributes that we collect (and which document the precise meaning of the data),

e. data has ascertainment and other biases as well as error rates that are poorly documented and difficult to control (see reports (CLARK *et al.* 2005; LACHANCE AND TISHKOFF 2013) of biases in initial samples of human genomes),

f. there are other practical problems that are often associated with scattered big data (see, e.g. (GITELMAN 2013; MCCALLUM 2013)), or that require too

much data wrangling before a given information resource becomes useful

(GOLDSTON 2008).

The aggregated difficulties of navigating all these challenges make it much easier to

understand why only a minority of circadian clock modelers chose to estimate

parameters directly from such time-series data, and why many others preferred to

match abstract time series traits (see Results below).

Using both, complete observed time series and abstract traits. It is obvious that

both types of observations presented above have advantages and disadvantages.

Complete experimentally observed time series increase the information available

for parameter inference, but bring the costs of handling more complex, yet incomplete

datasets associated with the inevitable problems of real-world measurements. Due to

experimental challenges, almost all such time series report relative amounts that are

comparable within a specific observed time series. It is rarely possible to obtain

reasonably precise calibrations those absolute units that matter most for modeling: the

counts of different types of molecules within their respective cellular compartments.

Furthermore, experiments may only report aggregated amounts, averaging over cells or

other biological units. Such practical details can substantially complicate the

computation of the likelihood that a given model will produce a certain observation.

Solving these problems does not determine the weights of different points of

observation in time series. Ideally, such weights maximize the impact of key information

while minimizing noise to avoid overfitting from algorithms that focus on unimportant

details, especially if no absolute calibration is available (as usual). Thus, many

researchers have historically avoided raw time series and used higher-level abstractions

of time series traits to assess the realism of circadian clock models. The relative

difficulties of implementation may have contributed to this trend reported in the Results.

Working with abstract time series traits provides the ease of using higher-level

traits, but comes at a price of its own. Abstract traits usually require fewer dimensions to

be managed and could minimize overfitting if they provide a focused view of important

clock features. However, abstract clock traits are difficult to choose and can easily omit

potentially pivotal information. Higher levels of abstraction can make it easier to find

parameter combinations that mimic observed ones 'reasonably well'. Since the quality

of such fits to observations can be judged in many ways, abstract traits might make it

too easy to produce a 'working clock'. Such model can easily omit details that are

essential for understanding a *particular* biological circadian clock. In the worst case they

degenerate into descriptions of artificial circuits that oscillate, but are unlikely to help us

understand the carbon-based circadian clocks studied in biology. Abstract time series

traits can reject many parameter combinations as biologically irrelevant and thus

pivotally contribute to the construction of useful clock models. Any given abstract trait is

not likely to extract all statistical information from the data. Thus, combining many such

traits is essential for successful modeling, yet there is no guarantee that any

combination will be statistically sufficient such that it can extract all relevant statistical

information.

Distances. Furthermore, the use of multiple abstract traits raises the question of

how to compute distances among and between simulated models and independent wet

lab observations. Many summary statistics provide distance measures that can be

adequate for some questions, yet cannot extract all information from the data and are

therefore not adequate for other questions. The pervasive non-linearity of circadian

clock systems complicates combining multiple traits into one reliable overall summary

distance statistic. This results in unknown, unpredictable and hence uncontrollable

biases when estimating parameters for clock models. To reduce these uncontrollable

biases caused by abstract traits with imperfect statistical properties, researchers have

started to incorporate more time series data in more direct comparisons of distances

between observed and simulated time series (see below).

Both sides offer advantages. It might eventually be possible to combine their

insights for improving the accuracy and robustness of parameter estimates in circadian

clocks. It often appears easier to compare abstract clock traits than the data-rich

simulated time series and corresponding experimental observations that necessarily

come with many gaps and complex nuances. However, potentially important aspects of

circadian clock mechanisms might be impossible to uncover, except by using a more

data-rich time series based approach. This extra data often adds many more

dimensions, uncertainties and complexities. It can easily overburden modeling studies

with irrelevant details and noise that may lead to overfitting. To counter such difficulties,

abstract traits may complement full time series data by acting as powerful filters that

remove unrealistic models, which might be difficult to identify in other ways. Combining

both appropaches could provide a powerful set of tests for detecting realistic oscillation

patterns in new circadian clock models. Increasing the statistical power of such tests will

make it increasingly difficult for them to be passed by random parameter combinations.

To find values that pass all filters and move beyond a given local optimum, researchers

can now use a broad array of optimization techniques combined with raw computational

power (Bussieck and Meeraus 2004; Bagheri *et al.* 2008a; Leugering 2012; 2016).

However, finding parameter combinations that have extremely rare desired properties

does not guarantee the correctness of a model (see discussion in (Loewe 2016)). The

vastness of parameter spaces requires caution when claiming that useful parameter

combinations for circadian clock models describe biological reality (see Oreskes *et al.*

(1994)), even if it was difficult to find working parameter combination. Other input might

pass the same set of tests and thus have the same claim to be in an ensemble that

might be used to represent biological reality. The purpose of FlyClockbase is to improve

the availability of data for testing clock models that might be part of such ensembles.

Studies integrating time series data. The challenges above present significant

barriers to the incorporation of experimental data into models of the *D. melanogaster*

circadian clock. Thus it is not surprising that only three of the many modeling studies we

surveyed (see Results below) used experimentally observed fly time series to estimate

clock parameters in a more direct way. Fathallah-Shaykh (2009) used published

microarray data from Kadener (2007) to fit parameters related to cry mRNA oscillation.

Kuczenski (2007) used a Monte Carlo random walk method to find a set of parameters

most similar to time series of circadian mRNA and proteins from twelve different

experimental studies (Hardin *et al.* 1992; Zeng *et al.* 1994; Sehgal *et al.* 1995; Marrus

et al. 1996; So and Rosbash 1997; Bae *et al.* 1998; Lee *et al.* 1998; Blau 1999; Bae *et*

al. 2000; Kim *et al.* 2002; Cyran *et al.* 2003; Glossop *et al.* 2003). Leise (2007)

employed a coordinate search method to estimate parameters based on time series

from three papers (Lee *et al.* 1998; Bae *et al.* 2000; Shafer *et al.* 2002). Both

Kuczenski (2007) and Leise (2007) point to the fit between the experimental and

simulation data as evidence of the quality of their models. While further discussion of

the many statistical challenges of parameter estimation in real-world datasets is beyond

the scope of this study, we note here that such discussion is rather hypothetical without

an actual real-world compilation of "all known" time series observations that can test

how many of the real-world complications can be handled by any given approach.

One purpose of our study is to provide such an integrated dataset that paves the

way for more thorough analyses of statistical approaches to assessing how good a

given simulation result might fit to "all known experimental observations" of wildtype and

wildtype-like *D. melanogaster* clocks. We created FlyClockbase to lower the barriers

that currently limit the use of real-world data for improving simulation models.

                         QQv1r1

FlyClockbase data model overview

FlyClockbase is a file-based database for collecting and organizing experimentally observed time series of *D. melanogaster*, reporting the core circadian clock components, such as mRNAs and proteins in various states. FlyClockbase is dedicated to circadian clock research in flies and can after sufficient stabilization accept observations of circadian clocks in other organisms. It is publicly available at:

> *Repositories at   **https://github.com/FlyClockbase**   will become active some time before publication. For reviewing see zip-archive; if urgent, request a copy from Laurence Loewe, who maintains FlyClockbase for the foreseeable future.*

Despite starting with a shared interest in the same model organism, FlyClockbase is completely independent from FlyBase (DOS SANTOS *et al.* 2015), a portal for genomic and other information about Drosophila as a model organism.

Place   FIGURE 2   about here.

Overview. Constructing and maintaining a highly specialized biological information resource like FlyClockbase is only feasible for skilled biologists with a passion for flies and clocks. To improve the probability of finding capable biologist curators, we deemed it important to minimize the computational expertise required for making substantial

contributions to FlyClockbase. We were aiming to minimize IT overheads of initial

construction and longer-term maintenance. As discussed below and in the

Supplemental Material, this goal informed important requirements for improving the

efficiency of biologists curating FlyClockbase and the accuracy with which its formal

type system can capture biologically relevant information. As a result, we have made a

number of unconventional database design decisions. Below we summarize key

differences between the FlyClockbase design presented here and other typical

database designs currently used. We provide more details in the Supplemental Material,

but a full technical description of FlyClockbase is beyond the scope of this study, nor

can we appropriately present the many considerations that informed the current design.

Instead, we focus here on how our choices help biologists, who are (i) interested in the

biological question of reliably observable variability in circadian clocks of flies, or are

(ii) aiming to navigate FlyClockbase for using, building on, or contributing to the quality

of data in this new resource. Figure 2 provides an overview of high-level organization

and Table 2 lists the Brief and Explicit Names of various FlyClockbase data structures

(we use these *Italicized Proper Names* to distinguish well-specified FlyClockbase data

structures from the meaning of their generic English counterparts in usual orthography).

FlyClockbase is a Versioned Biological Information Resource (*VBIR*) with two *Sections*:

- *SumS*, the *Summary Section* stores statistical summaries of time series, such as

  arithmetic averages. We extracted these as they were presented in relevant

  publications (see Materials and Methods).

- *DetS*, the *Details Section* stores all individual observations available at a stage

  where they have not been aggregated into summary statistics. This enables

independent researchers to compute the best summary statistics for investigating

specific questions. Describing *DetS* is beyond the scope of this study (raw data is

almost never reported among the publications in FlyClockbase; see below).

Each *Section* stores variants of time series in the form of *Raw* and *Modified*

Observations (each in an *ObsRaw* or *ObsMod TimeSeries ContentTable*, respectively).

The big workflow steps of importing, fully integrating data, and extracting data for

analysis are marked in Figure 2 as steps D1-D10 and S1-S8. To facilitate comparisons

across different datasets, we refine *ObsRaw* into *ObsMod* data in our current workflow

Step S5 (for more details, see Materials and Methods Section below). Each *TimeSeries*

is further characterized by some *Attributes* and may exhibit certain *Traits* (see below for

details). *Attributes* denote inherent features that need to be stored and cannot be

computed, such an observed genotype. *Traits* capture emergent features that need to

be computed from the *Content* and *Attributes* of a time series, such as 'peaks'.

Separating *Attributes* and *Traits* helps to keep FlyClockbase organized and simplifies

selecting relevant time series (see Steps S6-S8, D6-D8 in Figure 2).

Place   TABLE 2   about here.

Logic in biology. Both, *Attributes* and *Traits*, may be absent in ways that may

be of biological interest and differ fundamentally between time series. These challenges

inspired us to investigate fundamental aspects of type systems and logic in

programming languages in a search for appropriate ways of quantifying various aspects

of uncertainty, unavailability, and inability to be tested shared among many scattered

and diverse datasets of biological interest. In the Supplemental Material we discuss a

new data type termed 'BioBinary' which stores one of the four alternative states termed

OK, OKO, KO, MIS, which are defined by an enumeration termed 'OKScale'. The

BioBinary type is designed for handling statements in biology, where "*completely true"*

or "*entirely false"*, are less appropriate than "*any transient intermediate"* or "*mistake"*

(see also Discussion below, Supplemental Material, and p.16 of the online supporting

material in LOEWE et al. (2016b); a full analysis is beyond our scope here).

Other design decisions of interest to biologists discussed in the Supplemental

Material include:

- the intertwining and mutual stimulation of the development of FlyClockbase

  and the unconventional way in which the Evolvix modeling language

  (http://evolvix.org), is being developed (LOEWE *et al.* 2016b) from a first

  prototype for recording time series in pure mass action models (EHLERT AND

  LOEWE 2014) towards adding general-purpose programming capabilities with

  the triple goal of maximizing expressivity, usability, and long-term backwards

  compatibility;

- the use of a stabilizing versioning number system for facilitating review

  processes in ways that improve the possibilities of working towards long-term

  stability without frustrating innovators by turning them away, based on the

  StabilizingZone of the Project Organization Stabilizing Tool (POST) system

  (see p.74 of online supporting material in (LOEWE *et al.* 2016b));

- reasons beyond ease of implementation and installation for not choosing a conventional database system, but rather design-dedicated file-folder structures in a file system that can be copied easily across system boundaries;

While these points are important for questions of reproducibility and programming language development in biology and beyond, they do not directly apply to the biology of circadian clocks and are hence discussed in the Supplemental Material. We next highlight aspects of FlyClockbase that impact the ability of users to represent very diverse data in a surprisingly direct way: our particular choice of basic storage technology.

Simple file system storage. To increase flexibility, FlyClockbase stores data in a simple, well-defined, stable layout of files and folders in standard file systems. This design is intended to:

- maximize accessibility to biologists with very diverse levels of computational literacy and who use many different computing platforms. On all of these platforms it should be easy for any researcher to start FlyClockbase: experimental biologists, who strongly prefer to work with standard spreadsheet software as well as computational biologists, who strongly prefer direct programmatic access to raw data files to implement their own analyses,
- minimize long-term maintenance costs by delegating storage to standard file systems that maximize ease of distribution across diverse platforms,

- reduce the need for mandatory database updates that may require costly

     developer time or endanger the accessibility of valuable data.

These advantages come at the cost of requiring the discipline necessary to maintaining

consistency, and expecting users to not irresponsibly alter data that is freely accessible

to them. We expect FlyClockbase mostly to be maintained by researchers with sufficient

experience and for submissions to be appropriately reviewed so it is always easy for

beginners to get the last authoritative version in case they need a fresh start. Our

choices of technologies and formats have numerous strategic reasons further detailed

in the Supplemental Material.

Flexibility. In light of the enormous logic and error-reporting challenges faced by

any application (see Supplemental Material), our decision to design FlyClockbase

around the simpler and less restrictive technology of a filesystem has provided us with

an open field of efficient experimentation to improve our way of handling the challenges

of data curation. Our key insight here is the importance to empower experimental

biologists with little or no computing background to efficiently *launch* important decisions

about the type system and controlled lists used in FlyClockbase. Launching is not

landing; developing a stable type-system requires more experience than launching the

decision to consider adding a new biological special case. The importance of efficiently

communicating and collaborating across very different disciplines cannot be over-

emphasized: most people with enough formal experience to understand formal type

systems (PIERCE 2002; PIERCE 2005) cannot imagine the many special biological cases

that a corresponding logic would have to be able to handle. This is different for

experimental biologists: if they cannot recall these from the top of their head, a few days

or weeks in the lab will quickly help them to remember. However, acquiring the

necessary biological expertise, usually comes at the cost of less training in the abstract

art of designing consistent and stable type systems.

Practically, we developed FlyClockbase's flexible file system folder-structure to

enable the storage of content in standardized spreadsheet files easily modified by

common spreadsheet programs. We found this easy to use by experimental biologists

who regularly experience (and thus are best positioned to help reduce) the tension

between the abstract type system (aiming to restrict chaos by setting some rules) and

reality (with its own rules). Their contributions are best recorded on the spot in the most

flexible form possible to ensure they are captured at all. This requires maximal flexibility

and permissions and is simplest to implement by providing every local user of a local

FlyClockbase installation the equivalent of full (FlyClockbase-)system administrator

rights (including the ability to add, change, delete, or wreck anything and everything in

their local copy of FlyClockbase). Please consult the Supplemental Material for a

discussion of permissions, backups and the reliability of data storage.

These and other reasons beyond the scope of this study have motivated us to

forgo the obvious speed advantages of well-known standard databases. We do this to

gain the potential for a reduction in the cost of maintenance and an increase in stability,

combined with the flexibility to experiment with more nuanced type systems. These type

systems can better represent the complexity, diversity, uncertainty and occasional

contradictions that are so pervasively found in biological data. While working on

FlyClockbase, we encountered such problems regularly, as we tried to integrate all

available information about the circadian clock of *D. melanogaster.* The Discussion

reviews some of these experiences, but a full analysis is beyond the scope of this

paper.

Data types for organizing content. FlyClockbase is organized around a few

data types that help to structure its data (see Figure 2 for overview of components,

highlighted as *ItalicizedProperNouns*). These are presented below after briefly

discussing the fundamental recurring concepts of *Content*, *Attribute,* and *Trait*, which

simplify navigating FlyClockbase data structures. Without loss of generality, we illustrate

our definitions using observed time series as an example:

- *Content (Cnt)*: a container for directly storing 'the data' describing items of

  primary interest. For example, the *Content* of a time series observation in

  FlyClockbase is given by a series of pairs, each storing a time and a number -

  ordered so times keep increasing.

- *Attribute (Att)*: data about data. Each *Attribute* stores a type of value describing a

  fragment of information 'inherent' to a given data item, such as one of its owners,

  methods of observation, contexts, or types. The inherent nature of *Attributes*

  implies that they always have to be stored in addition to, and can never be

  derived from the *Content* they describe. Sometimes also called 'metadata',

  Attributes provide informal descriptions of the type or history of an item that can

  be essential for the correct interpretation of *Content*. For example, using the

  Content of time series requires *Attributes* describing which type of clock

component was observed and in what context – neither of these can be derived from the *Content* itself.

- *Trait (Tra)*: data derived from data. *Traits* capture emergent features, which are externally defined properties, patterns, or conclusions derived from a given set of *Content* and its *Attributes.* For example, a time series that only records how some amount changes over time may allow the observation of one or more peaks, but neither the steps for recognizing such *Traits*, nor the annotations of peak presence or absence are part of the *Content* to which they refer.

Irrespective of how data is packaged, *Attributes* and *Traits* both characterize *Content*, but do so in different ways. In FlyClockbase the trio *Content*, *Attributes* and *Traits* form a causality chain. For example, the real-world circadian clocks of one or more flies **X** *in natura* causally affect the observed time series **Y** *in vivo*, which causally affect results of interest **Z** inferred *in silico*. Note that the flies **X** are described incompletely by *Attributes*, the time series **Y** is observed incompletely as Content and the results of interest **Z** are derived from *Content* and *Attributes* by using a set of steps that define this *Trait*. Researchers often search for some **Z** useful for investigating a **Y**, only to find their efforts undermined by loss of pivotal information on **X** (equivalent to missing type information causing many computer bugs).

FlyClockbase has been built to reduce this very problem for circadian clock research in *D. melanogaster* by providing scientists with all information about X, Y, and Z that has been made available, ideally without increasing, reducing, or biasing any existing uncertainties about X or Y. It is possible to package the same information in a myriad of different ways by nesting and re-packaging various combinations of these

three. However, their diverse nature would make the use and maintenance of

FlyClockbase unnecessarily complicated, as each type offers a distinct value: *Content*

stores each time series that meets the required specification (a big but doable task,

aiming for completion). Associated *Attributes* store as much biological and historic

context information as possible (often impossibly difficult, making available Attributes

very valuable). *Traits* are defined at will by active researchers investigating a given

biological question or in search of new interesting *Traits* (it is always possible to define

new ones, but few are interesting on the long run). Based on these, we define:

- *ContentTable (CntTbl)*: a table of frequently used data of type 'content' *s*uch as a specialized *TimeSeries ContentTable*; *Attributes* and *Traits* are stored separately (see below);

- *AttributeTable (AttTbl)*: a table of *Attributes* for a given *ContentTable.* Currently, the most important *AttributeTables* are those for *References*, and *Summary TimeSeries*;

- *TraitTable (TraTbl)*: a table of *Traits* as determined from the *Trait* definition and a given *ContentTable.* Currently, the most important *TraitTables* are those storing the *Peak* and *Valley* timing for the first day of each *TimeSeries* of each clock component after *ObsMod6* refinement (as given in *SearchResult*);

- *Reference (Ref)*: a specific set of *Attributes*, which combine to storing the bibliographic information about a published study that reports *Summaries* of experimentally observed *TimeSeries* or other data of interest (if not prohibited by copyright, FlyClockbase includes the corresponding files as *Content* of a *Reference*);

- *Reference AttributeTable (Ref AttTbl):* stores each *Reference* (but not the files of its study), determining once and for all its unique *Reference_IDX*, an index used throughout FlyClockbase (the next largest integer available);

- *Submission*: here a submitted set of experimental observations reporting enough *Details* to enable the independent computation of diverse *Summary* statistics of individual observations;

- *SearchResult*: a set of tables compiled automatically or manually from each *TimeSeries* in the *Details* and *Summary Section*, (i) by testing whether all *Attributes* meet the search criteria and (ii) for those that do, by testing whether the *Traits* of appropriately grouped individual or aggregated *TimeSeries* meets the *Trait* search criteria. Our results below derive from a single *SearchResult* extracted from *ObsMod6*, and analyzed in various ways as described.

The basic layout of the folder structure in FlyClockbase follows the layout specified in the Project Organization Stabilizing Tool (POST) system described elsewhere (LOEWE *et al.* 2016b). We next discuss additional data types of biological interest before returning to the current scope and data collection strategy of FlyClockbase.

Identification of *TimeSeries*. This and future studies will need to refer to time series in FlyClockbase unambiguously. This requires a user-friendly system of precise and stable identification, in order to facilitate giving, using, and maintaining labels for time series with minimal effort. Defining such a system is a challenge facing various naming problems (see tables 1-2 in LOEWE *et al.* 2016b). We aimed to avoid two extremes: (i) Using the next running number for the next time series creates efficient

labels, but complicates some frequent tasks, such as determining if a pair of time series

belong to the same study. (ii) Descriptive labels including author, year, figure-panel,

plot-symbol, etc. can be informative, but are often too tedious, hard to automate or

difficult to maintain (e.g. avoid synonyms). We therefore developed a system that

combines localized integers that stand for local *Items* in different frames of reference,

each of which defines a *Context* that is itself an *Item*, nested into a bigger *Context*. The

resulting nestable index integers gives the outermost local item identifier (*IDLocal*, *IDL*)

as the first, top, left-most integer. This top *ID* is separated by a dot ('.') from the next *ID*

and provides the *Context* necessary for interpreting this second, next-to-top, next-to-left-

most integer. This *ID* in turn is separated by a dot from the third, etc., creating as many

nested *Contexts* as needed (more details are beyond the scope of this study).

In practice, naming *TimeSeries* unambiguously in FlyClockbase requires the

following three types of local identifiers for these three levels of nesting in the *Context*

provided by FlyClockbase:

1. *Reference_IDL*, points to a bibliographic reference.

    The *Context* for interpreting the *IDLocal* of a *Ref* is FlyClockbase itself;

    Ref_IDL is identical to *Reference_IDX* introduced above.

2. *Figure_IDL,* points to a *Figure* in the *Context* of a study, given by its *Ref_IDL.*

3. *TimeSeries_IDL* points to a *TimeSeries* in the *Context* of a figure panel*,*

    given by its *Figure_IDL.*

Any contiguous sequence of the elements above forms an *IDFragment* (*IDF*), which

identifies its corresponding Items. To distinguish potentially ambiguous *IDFs* from full

identifiers in a memory area, we denote the latter as 'memory identifiers', or *IDM*s for

*IDMemory. IDM*s are unambiguous *IDFs* guaranteed to refer to a unique *Item* within a

defined memory area, like unique time series IDs in FlyClockbase. Thus, we can

unambiguously identify each *TimeSeries* by its *TimeSeries_IDM* in the FlyClockbase

SummarySection by using the following form:

```
    SumS.Reference_IDL.Figure_IDL.TimeSeries_IDL        (TS_IDM),
```

where each IDL is replaced by its respective integer. Like all other *ID*s, these *IDL*s are

stored as *TimeSeriesAttribute*s in the corresponding *AttributeTable* (*SumS TS AttTbl)*,

along with *Attributes* for both identifying the figure in terms used in its publication, and

the time series in its figure (e.g. capturing line-type, color, plot-symbol, etc.). Thus, a

TimeSeries_IDM such as '*SumS*.1.2.3' refers to time series 3 in figure 2 of reference 1

in the *SummarySection* of FlyClockbase. Since the numbering of time series in a figure

panel (etc.), and the numbering of the latter in a study is far from clearly determined, the

additional *Attributes* help identify the actual figure panel and time series denoted and

would also facilitate the generation of automated reports in the future. For simplicity, we

drop the leading "`SumS.`" from *TimeSeries_IDM*s elsewhere in this text (all TS are

Summarized). This is appropriate until a *DetailSection* (*DetS*) is introduced for capturing

non-summarized time series measurements in FlyClockbase.

　　For example, four figures from one study (*Reference_IDM* "2") may have the

Figure_IDLs 1, 2, 5, or 8, resulting in FlyClockbase-wide *Figure_IDMs* 2.1, 2.2, 2.5, or

2.8; currently, gaps in *Figure_IDLs* (such as 3, 6, or 7) are allowed if unavoidable and

may indicate that data has been excluded when we later found that it did not fit criteria

for inclusion. If the above FlyClockbase-wide *Figure_IDM* 2.5 includes the pertinent

local *TimeSeries_IDL*s 3, 7, and 8, then their FlyClockbase-wide *TimeSeries_IDM*s will

be 2.5.3, 2.5.7, and 2.5.8. Each such *TimeSeries_IDM* points to a unique experimental

observation, unless marked in FlyClockbase as one of the rare cases where a review

re-publishes an older time series along with new data. In principle, a new *IDL* can be

any integer that has not yet been used in its local context. In practice, FlyClockbase will

critically depend on a single naming authority for assigning integers to their

corresponding items and ensuring that these assignments are never changed. Initially

this naming authority will be the maintainer of FlyClockbase, until this functionality can

be automated. Submissions of new studies to FlyClockbase can assign final *IDL*s for

figures and time series, but only a temporary *IDL* for a reference. The final

Reference_IDM can only be assigned once the new entry has arrived in memory area of

FlyClockbase. Following proper procedures for these naming issues is essential for the

integrity of FlyClockbase. Naming is complicated and the source of much concern for

managing biological data in *VBIR*s (NIH *et al.* 2012; LOEWE *et al.* 2016b). Naming time

series IDs provides a microcosm of the many problems that complicate naming. Still,

fully specifying a concrete dataset for further analysis requires more than particular

TimeSeries_IDMs: it also requires specifying the type of observation, as discussed next.

Raw, Mod, and Odd Observations. In FlyClockbase, each *Observation* (*Obs*),

is a *TimeSeries ContentTable* with time values measured in in *DZT* or *CZT* (as defined

below), and an associated measure of the amount of mRNA or protein at that time.

Each observation also includes measures quantifying imprecision and variability as

shown in the published figure (if any). *ObsRaw* ('raw observations') specify time as *DZT*

and contain the amount values as collected (including negative numbers or imprecision

resulting from undetected human error during data collection). Each *ObsMod* ('modified

observation') is a *TimeSeries ContentTable* measuring time in *CZT* and transforming

observations to contain amounts of mRNA or protein that are easier to compare across

time series (see steps S5,D5 in Figure 2). Recognizable problems with *ObsRaw* data

are appropriately corrected in *ObsMod*. *ObsRaw* time series over less than 6 hours or

with more than half of their *Content* marked as 'unreadable' were not simplified into

ObsMod. To identify and correct errors in FlyClockbase, we found it useful to pay close

attention to extreme values that might appear unusual or odd. We denote as *ObsOdd*.

This does not exclude them from analyses, but motivated us to revisit the whole

deduction chain that led to a given *ObsOdd*. Since any data collection will contain

human errors, we thought that users of FlyClockbase might find it useful to have

estimates of expected human error rates. We constructed repeated rounds of *ObsMod*

(see below), which were then used for our final analyses.

Data types of time measurements. We follow disciplinary conventions for

defining ZT (ZeitgeberTime) as *hours* since the last light period started (dawn). We

initially also defined this point in time as exactly ZT = 0. However, this resulted in

occasional unintended confusion of two very different meanings, simply because both

are conveniently denoted by zero: (i) a valid time measurement indicating an event

exactly at dawn as denoted by '0' and (ii) the inappropriate use of '0' for indicating that a

time was *NotGiven.* While the absence of a particular expected measurement is to be

indicated by the label '*NotGiven'* in FlyClockbase, it proved difficult to guarantee that no

unintended '0' could slip in. Elsewhere, such as for elementary addition, the use of '0'

for indicating absence as in '0 apples' is justified; it is also common enough and deeply

engrained, so that every new curator would have to spend significant learning effort to

avoid this ambiguity. Moreover, such errors are difficult to find, because a careful

analysis is required to determine, whether a particular '0' indicates '0*h*' or 'not given'. To

improve the long-term quality of FlyClockbase *and* reduce curation costs, we decided to

use a more robust Code2Brain interface (LOEWE *et al.* 2016b) instead. Hence, *DZT*=0*h*

has been declared a risky ambiguity to be removed from FlyClockbase as soon as

possible, whenever found (process is ongoing). The old *DZT*=0*h* is replaced by the new

DZT=24*h*, such that 0*h* < *DZT* ≤ 24*h*, while absence continues to be denoted as

NotGiven. We think that this new approach has a robust Code2Brain interface (LOEWE

et al. 2016b) and provides a high-quality representation of *Null* for DZT values (see

Table 2 and Discussion of Errors in Compilers below and elsewhere (WHITE *et al.*

2013)). An important difference exists between measuring fractions of *hours* in

FlyClockbase and outside. Usually, 1 hour comprises 60 minutes, but hours in

FlyClockbase are decimalized; thus, fractions of *hours* are measured in decimal

fractions and not minutes and the next hour is imminent at 0.99 *h*, not 59 min.

   CZT. To simplify analyzing several days of data in sequence, we define *Continuous*

ZT (*CZT*) to be an extension of ZT such that time increases without interruption at the

same rate over multiple days – instead of switching back to ZT = 0 at the start of each

new light period (dawn). Some studies use Circadian Time (CT), which can carry the

connotation of time series recorded in unusual light schemes such as 24-hour darkness

(DD) or 24-hour light (LL). *CZT* helps us to avoid any ambiguity. We use the term *Daily*

ZT (or *DZT*) when we mean ZT in this study and in FlyClockbase to reduce the potential

for confusion with *CZT* (also reduces ambiguity about types of time in code). Storing the

respective day together with *DZT* or ZT creates a 1:1 relation to *CZT*. For example, the

times of "lights on" (dawn) and "lights off" (dusk) over three days in the LD 12:12

scheme (12 hours of light (L) followed by 12 hours of darkness (D) each day) used in

the experiments of the initial FlyClockbase release can be given as *DZT* (24, 12, 24, 12,

24, 12), where days are implicitly assumed to form a sequence. These times are

equivalent to *CZT* (0, 12, 24, 36, 48, 60), using a different way of encoding days

implicitly. *CZT* time series simplify selecting observations only from the first day in any

given time series, as done in this study. We use italics for *DZT*, *CZT*, *h*, and *hours* to

indicate that these types are used as defined in FlyClockbase (see Table 2). We do not

italicize ZT, because we do not recommend its use in FlyClockbase.

Data types of amounts. None of the time series data collected reflected

absolute amounts or concentrations in a cell; rather, they show the amount of mRNA or

protein relative to a reference. References are different for many time series, which

presents a significant challenge when attempting to compare time series. The relative

amount of mRNA or protein at a given time cannot be compared across studies, so we

instead turned to a trait-based comparison method. We used modified observation

tables to extract two *Traits* from each day of a given time series: time of maximum

expression ("peak") and time of minimum expression ("valley"). We then combined

these two *Trait* values with time series *Attributes* to produce *PeakValleyTables*, which

represent *SearchResults* for further analysis. For more details, please refer to the

Materials and Methods.

Current definition of scope. Aligned with our interest to construct the best

possible circadian clock model for wildtype *D. melanogaster*, FlyClockbase currently

only includes time series from wild-type or wild-type-like flies (e.g., Canton-s, yw,

"control") observed in a LD 12:12 environment from studies published between 1990

and 2015 (see search criteria below). Thus, we currently exclude on purpose any

mutants that are meant to carry changes in clock genes, diversity in light-dark regimes

and other species for reducing the complexity of data curation. We include as "wildtype-

like" any mutants that were constructed without the intention of altering the dynamics of

clock components, including reporter genes (e.g. luciferase) and modifications to body

and eye color (e.g. yw). We thereby take the reported results at face value, implying that

such genetic engineering actually does not affect clock dynamics. Testing this

assumption is beyond the scope of this study and might become possible with the help

of large numbers of replicates collected in FlyClockbase. Measurement errors

associated with many observed time series are substantial and so is their variability

between time series. Thus we assume in this study that "wildtype" and "wildtype-like"

flies observed in the control experiments of many clock studies can be pooled. As a

result, we are including time series from the 86 studies cited below in the first public

release of FlyClockbase (QQv1 in the *StablizingZone* notation of the POST system

(LOEWE *et al.* 2016b), see http://evolvix.org/post). Beyond historic accident, there is no

  QQv1r1

particular reason to limit FlyClockbase to this scope, as long as expansions of scope

are coordinated carefully with corresponding data structures that enable the selection of

desired datasets.

Collection of data. Unfortunately, only one study provided individual raw time

series observations in addition to summaries (SHI *et al.* 2014). For the 85 other studies,

we extracted observed amounts and times from the time series figures published in

these papers (by plot digitizing, see details below). Future releases of the database will

allow the inclusion of individual time series observations in the *Details Section.* This will

enable meta-analyses to customize the statistics they report in order to choose

measures of variation that may be more appropriate than the arithmetic mean and

standard deviation. In our experience not all studies specify the variation measures they

report with the appropriate care (e.g. failing to specify whether a figure reports standard

devitations or standard errors of the mean; see SALSBURG (1985) for similar

experiences).

MATERIALS AND METHODS

Literature search

We searched the literature databases PubMed and Web of Science to collect references with time series data of the core components of the *D. melanogaster* circadian clock. Time series were broadly defined as any timed measurements of amounts of a relevant type of mRNA or protein that showed a daily peak or valley time for clock components, irrespective of the absence of scaling, calibration, and linearity. Search terms focused on variations of the terms "drosophila melanogaster" and "circadian clock" (plural, singular, or MeSH terms, or requiring any of the words in "circadian clock"). Marking phrases as specifically being MeSH terms did not influence the number of results. Using plural search terms (i.e., "circadian clocks," "clocks") reduced the number of results, sometimes by hundreds of articles. Requiring both words in "circadian clock" (as opposed to allowing either "circadian" or "clock") also decreased results by up to half. To reduce the likelihood that relevant data would be excluded in the initial literature search, we chose terms that produced as many results as possible. The final literature search occurred on March 26, 2015. After we removed duplicate studies, this initial search produced 1249 results.

  QQv1r1

Initial eligibility assessment. We assessed the title and abstract of each study

identified in the literature search based on the following three factors:

1. *Apparent content.* We excluded articles focusing on organisms other than *D.*

   melanogaster or centered on processes other than the core clock. We define the

   "core clock" to be comprised of genes integral to the functioning of the circadian

   clock in pacemaker cells, with a particular focus on the small ventral-lateral neurons.

   These genes include those shown in Figure 1 as well as *ck2a*, *sgg*, and *pdf* (see

   Table 1). Genes related to upstream or downstream clock processes were not

   included as part of the core clock; neither were genes that affect transcription,

   translation, or degradation rates in general. We also excluded papers if they were

   deemed unlikely to contain relevant time series based on the title. The articles we

   excluded based on this criterion focused on functional areas such as sleep, rest,

   arousal, locomotor rhythms, the visual system, metabolism and feeding.

2. *Type of data.* We only included papers with experimental data. As simulation data is

   beyond the scope of FlyClockbase, we excluded articles focusing solely on

   mathematical models since simulation data is beyond the scope of FlyClockbase.

3. *Format and availability.* We excluded the following reference formats because they

   were unlikely to contain specific experimental data: book chapters and prefaces,

   comments, dispatches, features, meeting reports, monitors, news, outlook articles,

   prediction reports, perspective articles, and reports from workshops. We also

   excluded one paper that was not available in English.

We then examined the full texts of the remaining 603 studies to determine which papers contained time series data. We were able to find only one article with raw time series data (SHI *et al.* 2014), so we used time series figures as a summarized proxy for raw measurement data. We found 149 studies with at least one time series figure.

Biological eligibility assessment. We further filtered these 149 studies with time series based on the following biological factors:

1. *External conditions.* We excluded time series with light schemes other than 12 hours each of light and darkness (12:12 LD) or with temperatures that varied over the time of collection.

2. *Observed cell specimens.* We required time series data to be based on measurements taken from biological material including at least some of the neurons closely related to the central clock in Drosophila (e.g., the small and large ventral-lateral, dorsal, dorsal-lateral, and posterior neurons, and S2 cells). For example, we included time series data taken from whole fly heads, whole flies, or the specified cell groups but excluded data from fly eyes or wings.

3. *Genotypes.* We only collected time series of wild-type or wild-type-like fly strains. We considered strains described as "wild-type", "control", "+/+", or "Canton-s" to be wild-type strains. We also included other fly strains if they were natural variants such as CRY-H and CRY-s (time series IDs 9.1.1 and 9.1.2). We excluded genotypes with mutations intentionally inserted to affect levels of protein expression, phosphorylation, binding, or light response of core clock proteins. We characterized wildtype-like flies as any animals with mutations not believed to interfere with the

operation of the clock. Examples include "yw" flies (have yellow bodies and white

eyes) and insertions of reporter genes such as luciferase, which are co-expressed

with clock genes.

4. *Amount and type of data.* We excluded time series if they were generated by

mathematical extrapolation, contained fewer than three data points, or covered less

than 12 hours.

The 86 remaining studies were defined as biologically eligible for inclusion in the initial

release of FlyClockbase and each study, relevant figure and relevant time series were

given a corresponding *Reference_IDM, Figure_IDL*, and *TimeSeries_IDL* as described

above (see Figure 2, Steps S1-S4, and Section Identification of *TimeSeries*).

TimeSeries data extraction

Since only a single study provided raw time series data in its online material (SHI *et al.*
2014), we extracted numbers for amounts and times from plotted time series in figures
as follows. We first extracted a screenshot of each figure from an appropriately
magnified downloaded PDF-formatted copy of the relevant study (using the Mac OSX
program "Grab," a simple application for taking screenshots). We then extracted the
data from each individual time series from its corresponding image using the open
source program "Plot Digitizer" (version 2.6.3 available at
http://plotdigitizer.sourceforge.net/) and recorded the result in a *SummarySection*
ObsRaw TimeSeries ContentTable (Step S2 in Figure 2). Plot Digitizer requires users to
specify the plotted values and the physical locations of minima and maxima for each of
the x- and y-axes in the figure. We did not detect significant curvature in the planes of
plots (as might be added by careless digitizing of printed copies) and therefore assume
that Plot Digitizer's linear interpolation provides reasonably accurate numerical x and y
coordinates of each point manually selected by the user. All y-axis values were directly
recorded using Plot Digitizer, unless a study specifically stated the value of mRNA or
protein at a given time. For x-axis, the value from Plot Digitizer was disregarded if the
value was noted in the text or clearly marked on the graph.

Accuracy estimates of digitized *TimeSeries* data. To assess the human
operator component of digitizing accuracy, we measured the variability of plot-digitizing
by three authors. Each operator plot-digitized the sample time series (63.1, WTLD,

          QQv1r1

green line with green squares) independently three times to produce a set of values

ready for inclusion as if it was true raw data. For each operator, time point digitized, and

axis, we calculated the following values: the mean of the absolute value of the relative

difference between each pair of the three independently produced values. Averaging

over all time points allowed us to calculate the average operator difference percentage

for the time axis, t_{aod}, representing an estimate of the relative error that one might

expect for a new value added to a time series in FlyClockbase by plot-digitizing. We

observed these intra-operator averages for time (with a maximal value of t_{max} = 48):

t_{aod} = 0.72%, 1.79%, 1.80%, including the first point, and

t_{aod} = 0.76%, 1.07%,   0.75%, excluding the first point

of the time series for operator 1, 2, and 3, respectively. We also observed the following

equivalent measures for the amount values v_{aod} on the y-axis (with the maximal value

of v_{max} = 2), resulting in averages of

v_{aod} = 0.95%, 1.15%, 1.11%, including the first point, and

v_{aod} = 1.00%, 1.23%, 1.16%, excluding the first point.

Operator 3 plot-digitized all values in FlyClockbase. We averaged all values digitized by

all operators and we found:

t_{aod} = 1.40% and v_{aod} = 1.06%, including the first point, and

t_{aod} = 0.84% and v_{aod} = 1.12%, excluding the first point.

Averaging coefficients of variation calculated separately for each time point by

combining data from all operators gives

t_{acv} = 1.72% and v_{acv} = 2.61%, including the first point, and

t_{acv} = 0.90% and v_{acv} = 2.79%, excluding the first point.

The drop in t_{aod} and t_{acv} when excluding the first point of the time series stems from the uniformity of absolute errors ('hit a point on screen'), which results in a proportionally larger impact on small values; this can help estimating precise values near zero.

Overall, these measurements indicate that relative errors introduced by plot-digitizing are small (1% or less) compared to the errors associated with the wet-lab measurements. Thus, we conclude that errors from plot-digitizing can usually be ignored. However, these are not the only potential errors in FlyClockbase. We refined each *ObsRaw TimeSeries ContentTable* (direct from plot-digitizing) into a corresponding *ObsMod TimeSeries ContentTable* (see Step S5 in Figure 2) by correcting human errors associated with data extraction and annotation, as discussed below.

Extracting *TimeSeries Attributes*. After extracting *ObsRaw TimeSeries Content* from published figures, we extracted associated *TimeSeries Attributes* and *Reference Attributes* from the corresponding published experimental studies (see Steps S3-S4, Figure 2). *Attributes* relevant to a study as a whole is recorded in the *Reference AttributesTable*, while *Attributes* specific to a given time series is in the *TimeSeries AttributesTable*. We collected *Attributes* for each *TimeSeries* to serve two purposes:

(i)   to help us to ensure *TimeSeries* fit the biological eligibility criteria previously described;

(ii)   *Attributes* enable later comparisons of biological, methodological, and other factors that could result in variability between time series.

We collected *Attributes* related to these three categories of questions:

1. Information about the time series

   A. Which *MethodRealm* does the time series belong to (*in vitro, in vivo, ex vivo, post mortem*)?

   B. What is the molecular type of the time series? Which protein or mRNA does it represent? Which isoform or splicing variant, or phosphorylation state of a given mRNA or protein was measured (if relevant)?

   C. Does the time series reflect data based on Zeitgeber Time (*DZT* or *CZT*)?

   D. Do the amounts reported in the time series come provide any information for limiting associated measurement errors? If yes, which types of errors are reported: the standard error of the mean (*SEM*), the standard deviation (*SD*) or an *UnknownErrorMeasure (UKEM)*?

2. Information about the method used to collect the time series

   A. Which method was used to observe the time series?

   B. Which machines, reagents, and software were used?

   C. Which probes or antibodies and dilutions were used (if relevant)?

   D. Which calibrations were applied, both mathematically (e.g., raw values were scaled by the maximum value) and biologically (e.g., values measured are relative to a specific standard mRNA or protein)?

   E. How many repeats were observed, and how are those repeats defined?

   F. When was the first and last data point recorded (*CZT* or *DZT* with days)?

   G. How long did the overall experiment last (in hours)?

   H. How long were the intervals between observed data points (if regular)?

   I. At which specific times were observations recorded (if specified in the text of the study or clearly marked on the time series figure)?

3. Biological and environmental information about flies used to collect time series
data

    A. Were flies exposed to light:dark schemes other than 12:12 L:D?

    B. To which temperature were the flies exposed?

    C. How long were the flies entrained?

    D. How old were the flies?

    E. What were the genotypes of the flies?

    F. Which sex(es) of flies were used?

In some cases, longer *Comments* directly copied from the text of a study were the most appropriate way of describing a given *Attribute* without introducing the potential for errors from paraphrasing. To avoid visual clutter and unnecessary bloating of the *TimeSeries AttributeTable* we introduced the notion of column locality, which is defined by a locality index column that is allowed to have identical values in consecutive rows and thereby define one *LocalColumn* for each such run of identical values. The purpose of this construct is to provide a formal check for the use of the equivalent of '*ibid*.' in FlyClockbase. For example, adding "See Method_Comments, 2.5.1" in the column "Method_Comments" at time series row 2.5.2 indicates that the longer comment stored in the row above (2.5.1) also applies to the next time series. To avoid loss of context, the FlyClockbase look-up keyword "See" must be followed by an indication of the column and row that are being referenced. To maintain readability for humans and simplify the implementation of code that understands *LocalColumns*, such pointers to previous cells of a column must not be interrupted by cells with unrelated content.

TimeSeries Traits analysis of Peaks and Valleys refined by ObsOdd checks

We manually compiled a set of *SearchResultTables* that integrated a simplified set of *Attributes* describing the nature of a given time series. In addition these included *Traits* that computed the respective times where amounts show a peak or a valley on the first observed day of an *ObsMod* time series in FlyClockbase. These *SearchResultTables* were termed '*PeakValleyTables*' and constructed for each given mRNA and protein. Each row in a peak-valley table corresponds to one time series and records several *Attributes* (from the *SummarySection TimeSeries AttributeTable*, see Steps S6 in Figure 2) and two *Traits* (peak and valley, from the *ObsMod TimeSeries ContentTables*, see Steps S7-S8 in Figure 2 and Figure 3). Each row also stores a day index, indicating the day during which the reported peak and valley were observed, counting from the start of the experiment (*CZT* = 0h). For example, a day index of two would indicate that peak and valley of the time series described by the *Attributes* in the row were observed between 24 and 48h *CZT* after initiating the observation of this time series.

<center>Place   FIGURE 3   about here.</center>

Measuring a limit for maximal peak time variance. In an effort to limit mistakes rising from human error, we refined the initially constructed *PeakValleyTables* ('*Raw*') into a series of successively modified *PeakValleyTables* shown in Figure 3 ('*Mod1*' – '*Mod6*'). To create the first set of modified *PeakValleyTables* (*Mod1*), we identified *PeakValleyTables* where the standard deviation of the peak value, valley

values, or both was greater than six hours. We based this threshold on two control

distributions. First, we created a uniform distribution with 25 artefactual regularly placed

observations covering every hour of the day effectively starting at 0*h* up to the very end

at 24*h* in 1*h* steps. This distribution had an averages and median of *DZT*=12*h* ± 7.36*h*

SD; omitting the first or last hour reduced the standard deviation to 7.07*h*. We also

randomly sampled 1000 values from a uniform distribution with a range from 0*h* to 24*h*.

Repeating this exercise three times produced *DZT* medians of 11.66, 12.41, or 12.08 *h*,

averages of 11.85, 12.47, 12.01 *h* and standard deviations of 6.81, 6.92, 7.02 *h*,

respectively. We therefore concluded that observations of *SD* ≥ 6h effectively indicate

signals that are indistinguishable from randomly distributed impulses that are not

oscillating in any discernably coordinated manner. Assuming that clock researchers

were probably correct when they reported oscillations, we explored the hypothesis that

such high variation *SD*s might have been caused by errors in acquiring or interpreting

some aspect of the data.

Factors contributing to increased trait variance. An important early insight

was the necessity to exclude peaks or valleys that coincide with the first or last point of

a time series. Although these points might appear to report the maximum and minimum

expression of a clock component, the amounts following or preceding this value (not

shown in the figure) could easily continue the local trend. For example, a time series

figure could appear to show maximum expression at the final data point (e.g. 23 *CZT*)

but reflect a system where the next theoretical data point (e.g. 25 *CZT*) corresponds to

the "true" maximum expression. We excluded these time series and made additional

corrections of observed human errors to construct the refined dataset *ObsMod1* from

ObsRaw. This early success in using odd observations for detecting potential lower-

level problems in datasets encouraged us to continue to investigate unusually extreme

values, which we then defined as *ObsOdd* peak or valley times outside of the range

defined by a given clock component's observed *Avg* ± 1 *SD*. Time series with *ObsOdd*

in *Mod1* were recorded in the peak-valley table '*Odd1*'. After correcting mistakes in

Odd1, we combined the corrected values from *Odd1* with the remainder of the data from

Mod1 to create *Mod2*. We repeated this cycle of checking for mistakes, recording

unusual values in *Odd PeakValleyTables*, and fixing errors until we created the final set

of modified and odd *PeakValleyTables*, *Mod6* and *Odd6*. In addition to correcting more

unique human errors, we also made adjustments for these potential method-based

sources of errors:

1. *Local minima in peaks*: Some time series (e.g. 85.7.2, 65.1.3, 81.4.1) report a

    local minimum, where it is easy to intuitively suspect a peak. We only adjusted

    our peak estimate if (i) a peak was also expected based on other time series of

    the same clock component, and (ii) the data points on either side of the local

    minimum are the highest two values for the respective day in the time series.

    These local dips can result from measurements outside of linear reporting

    ranges for time series observation methods such as RNase protection assays

    (RPAs) and Northern Blots. Increases beyond their linear range no longer

    produce linear increases in the intensity of signals and might even decrease

    the signal if product inhibition phenomena occur. We therefore suggest these

    local minima reflect measurement inaccuracies rather than actual decreases in

    amounts. To correct for this error, we recorded peak time as the average time

    of the two surrounding near-peak amounts.

2. *Luciferase initial spike*: Time series observed using luciferase (e.g. 62.2.1) may appear to report a peak shortly after starting to record data. The timing of this first peak can be inconsistent with peaks on other days in the same time series and other time series. Often the second peak has a *DZT* timing similar to peaks on subsequent days and in other time series. Such odd initial maxima are likely to be artifacts of the bioluminescence technique used to measure such a time series. They can be caused by an initial adjustment period that is required for accurate measurements of luciferase levels. Reported extra peaks occurs shortly after the arbitrary end of such initial periods (PLAUTZ *et al.* 1997; STANEWSKY *et al.* 1997). In such cases, we ignore the initial peak and record the values associated with the second peak on that day (as opposed to the technical maximum).

3. *Minimal Duration*: Some time series (e.g. see Figure_IDs 43.1 - 43.6, 61.1 - 61.4, 75.3) had a duration of twelve or fewer hours. Although minima and maxima can be read from such figures, their use is questionable, in particular, when the actual peak or valley times are not expected in the recorded time. Thus, we mark these peak or valley times as not given.

Linearizing *TimeSeries* data. The cyclic nature of circadian rhythms must be kept in mind when statistically describing the times of peaks or valleys in circadian time series. To illustrate this point, we will use data from the first day of *clk* mRNA time series. With no alterations, mean peak time estimates suggest *DZT* = 8.79h ± 8.59h (*SD*), which is indistinguishable from randomly distributed peaks (see above). Closer

                    QQv1r1

inspection of the data, however, shows that these values may be misleading.

Calculating the mean and standard deviation depends on finding differences between

values representing time, an operation that is substantially complicated by the circular

nature of hours in a day, where the value of 24h + 3 min results in 0.05h. This is in

sharp contrast to any linear expectation. For example, the central peak times for the *clk*

time series 35.2.1 and 35.2.2 (see Fig. S3B in KADENER *et al.* 2009) are observed at

about *CZT* 23 and about *CZT* 27, respectively. Translating them into circular circadian

time results in *DZT* 23 and *DZT* 3, respectively. If we then disregard the circular nature

of these values, we might infer a time difference of 20h between *DZT* 23 and *DZT* 3.

However, visual inspection quickly clarifies that the peak occurring at *CZT* 23 on the first

day is close to the early peak of the following day, which occurs at *CZT* 27h given by

the sum of (Day 1 *DZT* 24h) + (Day 2 *DZT* 3h). The difference between these peaks in

time series 35.2.1 and 35.2.2 is thus more accurately calculated as *CZT* 27 - *CZT* 23 =

4h. The corresponding change is equivalent to a local linearization of time when some

part of the original day crosses over into the next or last day. Peak or valley times that

have been linearized in this way are hereafter called "linearized."

We manually linearized all values in the *Mod PeakValleyTables* by moving the

minority of values to an earlier or later day (i.e. adding or subtracting 24 hours). Raw

PeakValleyTables are not linearized. Our current way of linearizing is geared towards

analyses of a single 24h period. Close direct visual inspection of all relevant time series

and claimed peaks makes it reasonably easy to linearize other periods; here it is

beyond our scope to conduct more general analyses. We found that linearization

greatly increases the overall reliability of representations of groups of time series. This

is of particular importance when estimating the variability of peak and valley times,

which is easily inflated artificially by omitting the linearization step.

Table S1 demonstrates the differences that can result from calculating summary statistics for *Raw* (circular time, no value linearized) or *Mod* (linearized time) peak and valley times of selected types of mRNA and protein. Results show that median, mean, and *SD* of linearized times may (but are not required to) differ dramatically from those calculated for circular raw times. For example, the linearized mean peak time for *clk* mRNA occurs 6 hours before its circular raw equivalent. Similarly, the *SD* of linearized peaks is about one-third of the *SD* of peaks measured in raw circular time.

Comparisons to summary statistics of uniform random distributions (see above) are instructive; as a rule of thumb, peak and valley times (or any daily event times) with a standard deviation greater than six hours should be treated with suspicion; they might be difficult to distinguish from randomly distributed times or could be the result of a lack of linearization. The latter affects *Raw SD* values for *clk* mRNA peak time, PER protein peak time, and *per* mRNA valley time (Table S1), which are close to *SD* values for uniform randomly drawn samples. In contrast, after linearization, *SD* for these *Traits* are much more similar to the *SD* of *Traits* of other linearized time series. Overall, this and other experiences suggest that linearization is an important step in obtaining trustworthy summary statistics from circular values such as *DZT*, even though linearization makes no difference in cases where times are already linear (e.g. TIM protein peak with peak *CZT* 18.41 ± 2.54 in Table S1).

                                            QQv1r1

Statistical analyses

Our initial screening for variability of the Traits we call *Peak* time and *Valley* time
(observed on day 1) surprised us by suggesting that variability might differ significantly
among clock components. Given the various sources of spurious variability described
above, we aimed to remove all artifacts that might randomly inflate variability estimates,
including potential biases that might be introduced from analyzing more than the first
day of a time series (after entrainment). Our interest in reproducible results and
disappointing experiences with large untested data collections has inspired various
rounds of error checking and increasingly rigorous statistical analysis.

     Automated analysis with R script. The core results of our study (differences in
variance between peak times of PER and TIM proteins) have been tested independently
by three of us (KS, BH, LL). The most rigorous analysis is presented below and can
easily be reproduced by running the script

     `FlyClockbase_PER_TIM_Methods_PeakValley_Comparisons_2016.txt`
which is provided in the Supplemental Material along with input files and the
Supplemental Statistical Analysis, which is an annotated PDF, collating all pdf output
from our plotting and analysis script. We executed the script on R version 3.2.4 (as of
2016-03-10, https://www.r-project.org). It requires the package "`data.table`" and the
library of robust statistical testing functions implemented by this script

     `http://dornsife.usc.edu/assets/sites/239/docs/Rallfun-v30.txt`
All statistical analyses used data from *PeakValleyTables*, where our final results are

taken from *ObsMod6* (see Figure 3 and text above). All corresponding input files are

provided next to the R script and are denoted as *ObsMod7* and *ObsMod8* as described

in the code. Times are given in *CZT* and have been linearized as described above.

　　　While the script contains numerous comments, it does not attempt to be elegant

code. Much of its over 12,000 lines appear at first glance to be repetitive with small

variations. It is currently not clear how to simplify the documentation of this script or

whether the time required for substantial code improvements would be well invested.

The trade-off between readability and coding time is further discussed in the R-code

and Supplemental Material under the approach to documenting code denoted as

'DISCOVARCY'-style.

　　　Outlier analysis. We addressed above those irregular values in *ObsOdd* that

were demonstrably due to human errors from data processing, removing them from

considerations below. Due to the substantial variability of the reported time series and

the diversity of measurement methods used to collect them, we were concerned that a

few substantially different outliers might obscure a robust trend exhibited by the majority

of observations. Thus, we used the following three different approaches for testing the

impact of outliers when analyzing Trait X by removing as outliers all values X_i, where

(i)　　X_i is outside of the range of non-outliers given by

　　　$(q_1 - 1.5 * IQR) < X_i < (q_3 + 1.5 * IQR)$, where IQR = $(q_3 - q_1)$ is the

　　　Inter-Quartile Range, and q_i are the corresponding quartiles,

(ii)　　X_i is identified an 'extreme value' by close visual inspection and its extreme

　　　difference to equivalent observations. This manual approach removed Protein

time series 14.1.1 for TIM, and 43.2.1, 43.3.1, 43.5.1 for PER, but none for the

corresponding mRNAs (see the *BestNoXtrem* input for the R-script above),

(iii) X_i is identified as an outlier by Carling's modification to the standard boxplot

approach (Carling 2000; Wilcox 2012). This technique uses sample size to

adjust the range of outliers to account for the tendency to identify a greater

number of outliers at smaller sample sizes. We used the implementation

described by Wilcox (2012, see section 3.13.3 and 3.13.5 on p.97-98 as

implemented in his R script "`Rallfun-v30.txt`" as function "`outbox`" when

called with parameters "`mbox=T, gval=NA`", so that his eq. 3.45 on p.97 is

applied). This method is applied to the data analyzed by our R script described

above.

Testing differences in variance. To test whether differences in variance are

statistically significant at the level of alpha = 0.05, we ran 100,000 bootstraps of the

percentile bootstrap method implemented by the function "`comvar2`" in the R script

"`Rallfun-v30.txt`", as described by Wilcox (2012) on p.175, section 5.5.2 and

elsewhere (Wilcox 2002). This function provides a 0.95 confidence interval for an

estimate of the difference between the variances of two groups, but was implemented in

a way that only detects significance at the 5% level (without giving *P* values). We used

this newer, more robust method to avoid problems associated with older methods such

as Levene's test (Nordstokke and Zumbo 2007).

Testing differences in mean. To test for 95% confidence in significantly different locations when comparing distributions, we used (i) the Mann−Whitney−U test as implemented in R (`wilcox.test`, 2 sided, unpaired), (ii) calculated 100,000 bootraps of the "`bootdpci`" difference as described on p.202 in section 5.9.12 of WILCOX (2012), and (iii) 100,000 bootstraps of the "`medpb2`" difference of medians as described on p.174 in section 5.4.3. of WILCOX (2012). Additional details of the function calls are easily found in the source code of our R script that performs these calculations; the results are given in the PDF output of this script, which is also available online as collated and annotated Supplemental Statistical Analysis.

Supplemental Statistical Analysis. The same R script produced the results plots shown in the main text below and the 81 pages of auto-generated plots shared with 6 additional pages for navigation as Supplemental Statistical Analysis in the Supplemental Material. It was generated by combining various snippets of code to test for all combinations of input data, outlier removal, observed *Traits*, molecule types and genes. This resulted in 32 distributions, combining the following features: input data (with and without the manually identified *BestNoXtrem* outliers), outlier removal (with and without applying Carling's outlier removal), and all combinations of *Traits* (peak, valley), molecule type (mRNA, Protein) and gene (*per*, *tim*). Method comparisons employed the same set of statistical tests for comparing locations and variances of two distributions (PCR vs non-PCR).

RESULTS

Experimental observations used in modeling

In addition to experimental results, our extensive literature review identified 66 studies

published since 1995 which focused on modeling the *D. melanogaster* core circadian

clock. Figure 4 shows that about 75% (50/66) of all modeling studies reuse parameters

reported in other modeling studies. Of the remaining studies, 13 relied exclusively on

abstract time series traits for estimating parameters, two used exclusively direct

experimental observations of time series data (KUCZENSKI *et al.* 2007; FATHALLAH-

SHAYKH *et al.* 2009), and one did both (LEISE AND MOIN 2007). In addition, one study re-

used parameters from one of the three studies mentioned above. Thus, only 6% (4/66)

of all simulation studies were based on direct experimental observations of time series.

This does not include the 13 studies that estimated parameters from abstract traits. The

substantially different numbers in each category might reflect difficulties inherent in

curating and incorporating experimental data into clock simulations (see above Section

on Models). We experienced first-hand many such difficulties. Although models will

never perfectly simulate reality and 'validation' is impossible on principal grounds

(ORESKES *et al.* 1994; BEERSMA 2005), we maintain that direct experimental evidence is

critical for increasing the relevance of models aiming to understand reality. Facilitating

1   the construction of more reliable models by including more direct observations

2   motivated us to build FlyClockbase. We found that FlyClockbase also enables

3   interesting retrospective meta-analyses, some of which we report below.

7   Place   FIGURE 4   about here.

8   *FIGURE 4. Where do models get their realism from?*

16   Place   FIGURE 5   about here.

17   *FIGURE 5. Overview of experimental studies available in*

18   *FlyClockbase and their use for parameter estimation over time.*

                    QQv1r1

FlyClockbase is a new resource enabling studies of circadian clock time series

FlyClockbase includes 403 time series covering 20 different mRNAs and proteins (13

and 7, respectively). An overview of its data model is given in Figure 2, relevant

abbreviations are summarized in Table 2. Table 3 shows its number of time series and

published studies corresponding to each type of mRNA or protein in FlyClockbase.

Figure 5 summarizes publication years from 1990 to 2015 for its 86 experimental

studies observing time series of clock components. Figure 5 highlights publication years

of the 16% (14/86) experimental wildtype observations used to directly inform 5% (3/66)

of all computational model studies of *D. melanogaster* circadian clocks (see Figure 4).

Place   TABLE 3   about here.

Hypothesis testing. FlyClockbase enables the study of many diverse biological

questions, making it impossible to present all corresponding biological results here (see

Discussion of using FlyClockbase for hypothesis testing). The analysis of most

biological questions, however, requires the computation of relevant *Traits* from the time

series stored in FlyClockbase. These *Traits* then need to be combined with the relevant

Attributes of the corresponding time series to form a row in the table of search results

(see Figure 2). We named these *PeakValleyTables*, since we constructed one for each

clock component in order to focus exclusively on analyses of the circadian timing of the

first peak and the first valley of each time series (defined by the respective maximal and

minimal amounts during the first day). After initially establishing FlyClockbase and

estimating the variability of all clock components, we limited the scope of this study to

the following two biological questions:

(i)   Comparing the variability of daily peaks and valleys of *per* and *tim* mRNA and

          protein, are there statistically significant differences in variance across

          independent time series?

(ii)   Which differences (if any) occur in the timing of *per* mRNA peak times based on

          different methods of observation?

We picked these specific questions because the initial release of FlyClockbase

contained the largest numbers of time series for these four components (ordered by

count we used 89 time series for *per*, 77 PER, 51 *tim,* 42 TIM). Thus, FlyClockbase

enables comparisons of variability of independently collected time series of circadian

clocks.

Error analysis. We conducted our initial retrospective meta-analysis of the

questions above after compiling FlyClockbase and refining it to correct all errors in data

handling that were known at that time (which produced *ObsMod5*, see Figure 3). Since

these initial results surprised us, we wanted to test their reliability. We had no estimates

for the frequency of errors that might be expected in FlyClockbase. It was not clear

whether there was a faster way of obtaining error estimates for such a complex,

manually compiled resource designed to organize scattered data that is heterogeneous,

diffuse, and characterized by many complexities, uncertainties, and gaps. We thus

opted for the most reliable approach available and rechecked every processing step

that could have affected the accuracy of our data sources or every peak and valley

timing used in our final analysis (producing *ObsMod6*, see Figure 3). These tests

improved our accuracy and generated realistic error rates for users of comparable

resources (see below).

Place   TABLE 4   about here.

Overview of *TimeSeries Traits*. The *PeakValleyTables* described above

allowed us to focus on efficient analyses of the timing of extreme amounts. These

prominent traits of time series have been used previously to estimate parameters and to

compare simulations and experiments (PETRI AND STENGL 2001; FATHALLAH-SHAYKH *et*

al. 2009). Table 4 provides summary statistics based on *ObsMod6 PeakValleyTables*

(see Figure 3) after applying outlier-removal method (i) described above. We will later

provide an independently computed overview of the mean and *SD* for the peak and

valley time of PER protein, TIM protein, *per* mRNA, and *tim* mRNA using outlier-removal

method (iii), see Figure 6 below. The amounts of *clk* and *cry* mRNA peak during the day

and reach a valley during the night. The components *per* mRNA, *tim* mRNA, *pdp1*

mRNA, PER protein, TIM, protein, and CRY protein display the opposite pattern: peaks

occur during the night and valleys during the day.

Variability of peak and valley times as measured by standard deviation varied

substantially from about one to four hours. This difference motivated us to determine

whether these differences in variability were statistically significant and might point to

mechanistic causes with implications for the inner workings of the clock. The most

frequently observed clock components were *per* and *tim* mRNA and protein, which

made them prime candidates for comparing their variability. However, one important

task remained before investigating our hypotheses: testing for errors.

Why estimate errors? We had to be reasonably certain that we could rely on

the values stored in our *PeakValleyTables.* This was particularly important for detecting

significant differences in variability, since random errors in the capture or processing of

data are known to easily generate spurious values that substantially impact observable

variability. As explored in the Discussion, attempts to detect and correct errors in

author-labs and distinguishing them from other potential sources of variability was

beyond the scope of our study. Thus, our goal was to be able to confidently exclude

errors during the capture and processing of data on our side. We defined as error in this

context any result that did not hold up to scrutiny, when rechecked using the rules we

had agreed on for processing time series data. These rules were set up after we gained

substantial experience with various counter-intuitive aspects of the data, as documented

in this study. Thus, our definitions of error were not arbitrary and therefore allow us to

contribute below a specific estimate of human error rates to the broader area of human

error analysis (see Discussion below).

Databases curated by human experts usually have substantially lower error rates

than those that were automatically compiled (SCHNOES *et al.* 2009; KOSKINEN *et al.*

2015). Still, the dangers of error accumulation in heterogeneous collections of data

contributed by humans are well known (CARTHEY *et al.* 2003; ZEEBERG *et al.* 2004;

PANKO AND AURIGEMMA 2010; PANKO 2016). It thus appears desirable to increase the

number of human error estimates available for biological information resources. The

analysis discussed next provided a unique point estimate of the error rates users might

want to expect when working with manually compiled resources of similar complexity. A

more thorough analysis of errors is beyond the scope of this study (and is likely to be

substantially affected by the compiler techniques discussed below). Nevertheless, our

results below support our claim that our main findings are probably not the result of a

unlucky confluence of statistical flukes generated by errors we could have detected

when reanalyzing the same data carefully.

Quality control in FlyClockbase and *TraitTables*. In order to detect and

reduce the number of errors in the *PeakValleyTables* used for computing variability, we

conducted four rounds of identifying values outside of the range defined by the mean ±

1 *SD* of a *Trait* (*Mod2*, *Mod3*, *Mod4*, *Mod5* in Figure 3); this was done for all clock

components in FlyClockbase and serves as the basic way of introducing all new data.

We used this as the starting point for obtaining the error estimates reported here. Since

time series *TraitTables* and search results tables are not strictly part of FlyClockbase

(see Figure 2), we separated our error estimates of FlyClockbase from those obtained

for *TraitTables* (see Results below) to improve the quality of our estimates. In a unique

effort we then re-examined in-depth all relevant content, attributes and traits of each

time series that informed the four most important *PeakValleyTables* (*per* mRNA, *tim*

mRNA, PER protein, and TIM protein). We scrutinized each value representing a peak

or valley time. This required a substantial effort, which aimed at a twofold goal:

(i)   removal of errors for improving our estimates of variability differences, and

(ii)   providing an approximate estimate for the numbers of similar errors that users of

FlyClockbase (or similarly complex data resources) might expect in other areas,

where such error rates have not yet been determined.

Before presenting our findings, we submit to the reader that the researchers compiling

FlyClockbase had been very conscientious, brought a high degree of expertise and

enthusiasm to the project, and did their very best to avoid mistakes. Thus we thought

conditions were favorable and we were not sure if we would find errors. Unfortunately,

aiming to avoid and correct mistakes does not protect against the inevitable occurrence

of mistakes at low but predictable rates (see Table 5 and Discussion).

Place   TABLE 5   about here.

Basic Null-Hypothesis H_{basic}. FlyClockbase includes a particularly large number

of repeatedly observed time series of *per* and *tim* mRNAs and proteins in wildtype (and

wildtype-like) circadian clocks of *D. melanogaster*. These time series were recorded as

wildtype control experiments while observing the effects of mutants in order to explore

circadian clock functions. We expect corresponding wildtype controls to produce similar

time courses that differ only by inevitable stochastic effects – in the absence of

experimental complications. Such complications would make flies non-comparable

across studies (see Discussion of potential causes, such as variable natural genetic

diversity across fly strains, developmental diversity, unknown environmental impact on

measurements, and others). We start by ignoring all such potential complications. Our

aim is to initially work with the simplest model that still appears somewhat useful. We

define a corresponding basic null-hypothesis H_{basic} to inform our background time series

expectations and enable a defined starting point for hypothesis testing with the help of

appropriately selected data in FlyClockbase. In light of the types of observations and

available calibrations, we define H_{basic} as:

The basic time series type in FlyClockbase is determined by the molecule type observed and its context, which for H_{basic} is defined as the central core clock of wildtype organisms of a defined taxonomic unit (like D. melanogaster) under standard experimental conditions. H_{basic} excludes time series that are (i) from non-LD 12:12 observations, (ii) affected by the presence of mutations known to or intended to alter the clock, (iii) measuring non-central circadian clocks that are independent or merely derived from an organisms' central clock, and (iv) any complications from genotypes, phenotypes, environments, methods, or more.

For these basic time series types defined above, FlyClockbase reports as H_{basic} the ensembles below containing either *dynamically changing amounts of a specified molecule type, or values of corresponding time series traits.*

These ensembles of the *most reliable observations in FlyClockbase are based on time series that satisfy the following conditions: (i) Timed amounts are reported as uncalibrated, relative measurements, which allow comparisons only within each time series, not between time series. (ii) Comparisons of observed amounts indicate all potential outcomes by drawing on as much evidence as possible and extracting as much quality as reasonable from the available data. This requires that comparisons define a method of incorporating such data appropriately. This data includes quantifications of uncertainty in methods of measurement, observational errors, replicates, accuracies of timed amounts, and methods of inference. (iii) When using H_{basic}, all relevant details need to be documented, including the data available at the time, its state of refinement*

(ObsMod+), and the methods used. Documentation requires specifying their

stabilizing versioning numbers as defined by the StabilizingZone of the POST

system (see Table P1 in Supplemental Material). (iv) Any observations included

in either ensemble type exclude outliers using Carling's method (2000) to enable

a focus on typical clocks.

Satisfying all these conditions, H_{basic} assumes that the remaining

variability of observed amounts or traits is only caused by the natural

stochasticity of discrete molecular events inside of individual cells.

H_{basic} is a powerful starting point for exploring clock biology. Next we will use H_{basic} to

compare typical H_{basic} behavior of time series traits observed in different clock

components. We then relax assumptions of H_{basic} to illustrate how FlyClockbase can

generate a variety of hypotheses about diverse subtleties that might be important for

generating high-quality observations of circadian clocks.

Hypothesis on Peak time variances: PER exceeds TIM

We initially screened for variability in the peak and valley times of all clock components

available (see Table 4). This analysis revealed substantial differences between the

standard deviations of peak times observed for PER and TIM respectively highlighted in

Figure 6. We hypothesized that these differences could have mechanistic causes and

might thus point to new insights into clock mechanisms (see Discussion). Before

exploring such mechanistic models, we wanted to know, whether our observations were

(i) statistically significant and (ii) not caused by low-level errors in data handling during

the construction of FlyClockbase in light of corresponding challenges with data quality

(MᴄCᴀʟʟᴜᴍ 2013) as faced by many biological information resources (see Discussion

and Supplemental Material for more details).

Place   FIGURE 6   about here.

Place   FIGURE 7   about here.

Data quality. Since variances are easily inflated by outliers caused by errors in

data handling, we aimed to obtain assessments of peak and valley times that were as

accurate as possible given the observations reported by the studies in FlyClockbase.

The desire for improving the overall rigor of our variability assessment motivated the

various procedures described above for identifying outliers and investigating potential

errors that might have artificially inflated estimates of variability. Thus, we manually re-

inspected all steps of inference for each peak and valley data point that contributed to

our final calculations of timing statistics. We started with the already highly refined

dataset *ObsMod5* and created *ObsMod6* by correcting all irregularities from data

handling or processing on our side that we could detect in FlyClockbase or our

TraitTables (see Figure 3). Starting from the figures of publications, we checked the

results of all our manual operations that could have modified traits, up to the final values

used as input for our R script that produced Figure 7. The statistics of the resulting

human error analysis is shown in Table 5.

Main analysis with outlier robustness. We decided to carefully investigate the role

of outliers when analyzing variance in order to arrive at robust conclusions. We were

motivated by the following considerations:

1. *Use of robust statistics.* We aimed to use state-of-the-art statistical methods

    designed for delivering robust conclusions that minimize the chances that a few

    odd values have an unduly large impact on the overall conclusions (see Wilcox

    (2012) for an introduction to robust estimation and hypothesis testing).

2. *Dealing with rare ObsOdd.* As described above, we checked and re-checked all

    observed peak and valley times that entered our final comparisons between *per*

    and *tim* components, aiming for the best interpretation of each reported time

    series. Despite this scrutiny, there were four observed time series that we could

    not interpret convincingly. All four extreme values pertain to protein peak time,

    with one extreme for TIM (time series ID 14.1.1, peak at 6.966h linearized *CZT*)

    and three extremes for PER (time series IDs 43.2.1, 43.3.1, and 43.5.1, all with a

peak at 28*h* linearized *CZT* and all from one study). All four also showed a clear

signal in their original figure that appeared to have been analyzed correctly

(based on our reading of the respective studies), yet all four reported times

appeared to be clearly distinct from the distribution of times reported by all other

similar studies. For example, TIM extreme outlier time series above observed a

peak almost exactly 12*h* away from the mean of the equivalent TIM distribution of

peak timings without outliers. While we could not find any indication that morning

and evening had been flipped in the corresponding study, it is very difficult to

exclude such human errors in light of the many challenges that easily frustrate

data analyses in spreadsheet software (see Discussion below). We stopped

further attempts of re-interpretation, since the original raw data was not publicly

available. As revisited in the Discussion, these values could represent any of the

following: genuine observations of typical behavior of rare fly clocks, rare

behavior of typical fly clocks (at least sustained for the length of the time series),

some rare combination of measurement protocol details that conspired to

systematically bias observations (which were correctly interpreted), or human

errors leading to misinterpretations in the complex chain of time series

observation and analysis. We could not find similarly extreme outliers in

corresponding observations of mRNA. We consider the overall evidence to be

too incomplete, contradictory and therefore inconclusive to determine which

process may contribute most.

Observing these extreme outliers might raise the distant possibility that

apparent wildtype circadian clock systems can exhibit extreme deviations from

their normal timing behavior in a few percent of occasions. However, exploring

the possibilities of such exotic behaviors is beyond the scope of this study and

not possible without a more elaborate error-management for observation and

analysis of time series in circadian clock studies. It would also require a

substantial number of individual time series observations in the *DetailSection* of

FlyClockbase to enable the independent computation of summary statistics.

To test whether these handpicked extreme outliers made a difference to

our conclusions, we grouped and denoted them in our Statistical Methods

Section as 'outlier removal approach (ii)'. We then created a parallel analysis

track in our R script that took every computation on the full dataset ('*Mod7*') and

repeated it on a manually created copy of the input files ('*Mod8*'), where this

manual outlier removal approach (ii) had been applied. To increase clarity,

'*Mod7*' or '*Mod8*' are also labeled, respectively, "*BestEachObs*" or "*BestNoXtrem*"

in our R code, and "with Extremes" or "no Extremes" in the titles of the

automatically generated plots. ('*Mod7*' is essentially identical with '*Mod6*' except

for trivial changes to facilitate automated reading from R; the *Mod7* and *Mod8*

files are only stored next to our R script that reads them as input, they are not

stored in the main time series trait folder).

As can be seen in the Supplemental Statistical Analysis online, the

removal of these four specific time series does not substantially change our

conclusions.

3. We were unsatisfied with the subjective nature of the decisions summarized

above as 'outlier removal method (ii)'. While extreme outliers are reasonably

easy to detect, there is a gradual transition to less extreme values, where

subjective decisions about the inclusion of particular time series could easily lead

to a new set of problems by creating ascertainment biases that are impossible to

correct for statistically. Thus, we decided to employ a principled method. After

some experimentation, we arrived at outlier removal approach (iii) which has

been described elsewhere (CARLING 2000; WILCOX 2012); see Statistical

Methods.

Our results in Table 6 report our current best analysis of the most reliable data on the

variability of *per* and *tim* protein and mRNA peak and valley timing accessible to us in

FlyClockbase. For the reasons given above, we decided to exclude outliers as identified

by our approach (iii) see Materials and Methods (CARLING 2000; WILCOX 2012).

Alternative handling of outliers. In the Supplemental Statistical Analysis we

compared results after removing outliers using approach (iii) as shown in Table 6 with

those obtained from the full dataset to investigate whether removing outliers affects

conclusions. Given the extraordinary range of timing variability for the peaks of PER and

TIM, it is unsurprising that the difference in variance reported in Table 6 loses statistical

significance when outliers are included. In Table 7 we summarize our results of

comparing different outlier approaches. Corresponding values for all other results in

Table 6 can be extracted from the Supplemental Statistical Analysis.

We conclude from Table 6 that the majority of circadian clocks in *D.*

melanogaster are significantly more variable in their timing of PER peaks in comparison

to TIM peaks (P<5%). In a minority of cases outliers can introduce such large amounts

of variability that indicators of significance are swamped and a loss of significance is

perceived (see Table 7 and Discussion). All other comparisons shown in the

Supplemental Statistical Analysis confirm this overview. Our observation of significant

differences in variability contrasts with the near absence of differences in the average

timing of these peaks.

Place   TABLE 6   about here.

Place   TABLE 7   about here.

Other peak comparisons. We find differences in the variability of mRNA peak

times, albeit interestingly with inverted sign, compared across *per-tim* equivalent parts.

This means that significantly lower variability in *per* mRNA peaks precedes significantly

larger variability in *PER* protein. Conversely, significantly larger variability in *tim* mRNA

precedes significantly lower variability in *TIM* protein. This flip indicates that the

differences in variability of the peaks for PER and TIM are not caused by corresponding

differences in the variance of their mRNA peaks. We therefore conclude that these

differences are caused by mechanisms affecting variances of peaks in the causal

reaction networks after the transcription of *per* and *tim*. These differences in variance

contrast with non-significant differences in the average peak timing of the same peak

time distributions.

Valley comparisons. The valleys of PER and TIM occur at significantly different average times, irrespective of how many outliers are included. However, the non-significant differences in variances reported in Table 6 become barely significant at the 5% level when adding the two outliers observed in TIM. When comparing the valleys of *per* and *tim* mRNAs, we found significant differences in average time; *per* valleys precede *tim* valleys, even though averages do not differ among peaks. Valleys of *per* and *tim* also showed significant differences in variance, with *per* being more variable than *tim*. We also report an inversion of variances when compared to their corresponding mRNA peaks. Our results for the valleys of *per* and *tim* mRNAs are not impacted by outliers, since none of our approaches to outlier analysis could identify any outliers among the 24 and 20 observations in FlyClockbase, respectively.

1 **Comparing methods for measuring *TimeSeries* of *per* mRNA**

2

3 In addition to comparing different clock components, FlyClockbase can contrast

4 experimental details that differ between independently observed time series of the same

5 clock component. These experimental details are stored as *TimeSeries Attributes* in

6 FlyClockbase and used for extracting corresponding sets of *TimeSeries* or *TimeSeries*

7 *Traits* for additional statistical analyses. Here we compared *per* mRNA time series

8 recorded by different measurement methods. We chose *per* mRNA because it is the

9 most common time series in FlyClockbase and thus provides the largest statistical

10 power for detecting potential systematic biases. Given the differences in variances

11 between clock components reported above, we wanted to know if such differences

12 could have been produced by using different methods of obtaining time series.

13

14 **Measurement methods:** The following five measurement methods were used to

15 collect at least three *per* mRNA time series in FlyClockbase: microarray, nascent-seq

16 and RNA-seq, PCR, RNase protection assays (RPAs), and Northern Blots. Two of the

17 four time series measured with microarrays were outliers, and all four of the time series

18 measured with RNA-seq or nascent-seq were from a single study. These time series

19 were consequently eliminated from our analyses. Many time series did not include a

20 valley time for *per* mRNA, so we focused solely on *per* peak time.

21 *Comparing means:* As expected, we could not find any significant differences in

22 the averages of peak times observed by different measurement methods when

23 comparing PCR vs Northern Blot, PCR vs RNAse Protection Assay (RPA), and

Northern Blot vs RPA (using the Mann-Whitney-U test at the 5% level, see

Supplemental Statistical Analysis).

   Comparing variances: We found significant differences in variance (using

comvar2) to compare measurement methods as above. Comparing the combined peaks

from time series measured with any PCR method to the pool of those measured with

RPA or Northern Blot resulted in significantly different variances for both. The 95% CI of

differences of variance reported by *comvar2* after 10^5 bootstraps for RPA and Northern

Blots are 0.4413 to 6.817 and 0.4764 to 10.586, respectively (for explanation on extra

digits, see Table 6). The histograms of the corresponding distributions are given in

Figures 8A and 8B.

                    Place   FIGURE 8 ABCD   about here.

   The differences in variances are strongest for peak times measured with PCR or

Northern Blots. Comparing the nine qPCR observations with the nine Northern Blots

showed significantly higher variance in qPCR observations (*comvar2*, 10^5 bootstraps,

95% CI for differences in variance: 0.353 to 7.598). Likewise, the 16 RT-PCR

observations in FlyClockbase are more variable than the nine Northern Blots (*comvar2*,

10^5 bootstraps, 95% CI for differences in variance: 0.46889 to 11.486). As reported

above, comparing peaks from PCR to those from RPAs showed significant differences

in variance. We expected this pattern to hold also for non-pooled PCR, but found results

to be no longer statistically significant. Each type of PCR had a very skewed distribution

of differences in variance with a substantial bias similar to that of Northern Blots (see

comvar2 results in the Supplemental Statistical Analysis). However, reduced sample

size diminished statistical power, so differences are no longer significant.

Method references. We were surprised by the larger variance in PCR results

compared to results from non-PCR methods. We speculated that this might not

necessarily be inherent to PCR but could be caused by a larger diversity of method

protocols. If this were general, we would expect studies with shared method protocol

references to report results with less variability than studies which did not share such

references. Therefore, in studies that did not cite any method references, we might

suspect a larger diversity of method protocol variants, and thus more diverse results.

Place   FIGURE 9   about here.

Figure 9 provides detailed information about the method protocol references

cited in the studies we used. As before, we focused on the peak times of individual time

series as observed by PCR, RPAs, or Northern Blots. Some authors largely followed the

methods outlined in other referenced studies, while others only incorporated protocol

references for specific aspects such as probe development, the use of controls, or RNA

extraction (see classification key in the caption of Figure 9 for a list of some

methodological aspects that might be of interest).

Table 8 provides summary counts of the time series represented in the various

broader categories of Figure 9 and the number of studies that observed them. Table 8

also provides summary counts of the time series from studies that either share or don't

share references with each other (along with the corresponding numbers of studies; this

classification excluded studies with no references). The underlying network of protocol

identities is represented in Figure 9, and histograms of the corresponding distributions

are given in Figure 8CD.

Place   TABLE 8   about here.

Presence or absence of method references: We grouped all analyzed time series

into two categories, which might be nicknamed "*Any Method Ref*" if their studies

provided at least one reference for details of the experimental protocols used to observe

time series using a given broad type of method, or "*No Method Ref*" if not a single

relevant experimental protocol reference was given. Comparing average peak times

among these groups showed no significant differences and no detectable indication of

bias (Mann-Whitney-U test, 95% CI: -1.00 to 1.00, p = 0.736). While differences in

variance were not significant, a clear bias in variance was noticeable in the distribution

of differences, possibly indicative of significance if higher sample sizes were available

(*comvar2*, 10^5 bootstraps, 95% CI for differences in variance: -1.385 to 5.6876).

Presence or absence of shared method references: We subdivided all available

time series with at least one method reference into those with and without shared

method references (see Figure 9 and Table 8). Average peak times did not change

significantly if method references were or were not shared, but some bias could be seen

(Mann-Whitney test, 95% CI for difference in location: -2.00 to 5.4208×10^{-5}, $P = 0.1829$).

Similarly, variances of peak times showed bias but were not significantly different

(*comvar2*, 10^5 bootstraps, 95% CI for differences in variance: -1.1045 to 6.325, see

Supplemental Statistical Analysis).

Reduction of statistical power by application of inappropriate logic? The

last groups above had both clear biases in their differences in variance as revealed by

comvar2, but neither test was significant at the 5% level. Combining both results in a

slightly different way might increase relevant sample sizes and significance. As shown

in Figure 10, we currently exclude studies with not a single method reference from our

analysis of studies with shared references. We initially justified this by assuming that

method references must be present for deciding whether studies could have used the

same methods at the lab-bench or not. It turns out that this line of reasoning might say

more about the need for greater clarify in logic formalisms than about the effects of

sharing references or not. We will next make explicit, what we unknowingly implied

above; we do so in order to demonstrate the difficulties of creating clear statements in

words chosen *adhoc* and aimed at clarifying statements in a limiting logic formalism.

Above we assume implicitly that a study without any method protocol references

may or may not have used the exact same methodological procedures as another study

without method references. Applying Boolean logic here attempts to force a

Yes   (OK)

No   (KO)

answer to the question about the use of a common protocol. This logic fails here,

because it does not allow us to specify that we cannot provide any certain answer

based on the absence of a reference (we cannot exclude the possibility that two studies

may have shared the same protocol). Instead, we might consider the BioBinary logic

that FlyClockbase is adopting from Evolvix (see Supplemental Material). This is a step

forward, because it can distinguish the cases of

> MIS   ('inapplicable' because 'no-method-given')

> OKO   ('somewhere between OK and KO')

from the other cases (OK/KO), where we can reach relatively more clear decisions

because method protocol references are given. Our statistical analysis above can now

be rephrased as stating that the absence of a reference is a BioBinary 'MIS' (since we

formally cannot answer the question due to a missing value). As a result, we would

exclude such values, since it is always easy to name multitudes of irrelevant values that

clearly should not have any influence on our analysis. This interpretation might appear

unsatisfactory at first, but it is still a step forward, because it is explicitly stated and

therefore more tangible, which might attract further scrutiny.

If explained clearly, most experimental biologists will probably not hesitate to

point out that the chances of using identical experimental protocols in the lab are

miniscule unless there is a shared reference to a common protocol (which would

probably be referenced). There are simply too many variations that are done most

easily. Thus, instead of assuming the absence of evidence (which might allow for

shared protocols, even if no references are given), we can assume evidence of absence

(since it is rather unlikely that two labs use the same procedures, without sharing

references to a common protocols). Given the many variations of PCR that are easily

created in the absence of further detailed instruction, it is difficult to see, how labs might

accidentally share procedures. Thus, technically, this case must be classified as a

BioBinary OKO; however, overwhelming evidence suggests that a KO (not sharing) is

much more likely.

Result: These considerations suggest that it would be more appropriate to add

the time-series with no references to the group of those that have only one (non-shared)

reference. This might increase statistical power enough to make tests significant that

compare "shared vs not shared references". Pivotally, these considerations

demonstrate the importance of using the right type of logic and type system. It is easy to

miss important points in Boolean logic without appropriate visualization. Since logic

formalisms are best automatically supported at the compiler level, a step change in

speed and quality of many biological analyses could be facilitated by a compiler that

correctly map BioBinary and more expressive logic formalisms to data – out of the box.

This conclusion is general for the construction of *VBIR*s, and independent from the

particular outcomes of any specific tests that might be performed, because the

existence of such unresolved difficulties in representing formal logic can easily create

bugs that bind large amounts of research time, which could otherwise be dedicated to

VBIR development.

Place   FIGURE 10   about here.

Comparing methods by study to reduce multiple comparisons. Figure 9

shows that a single study often reports multiple time series, which could bias the results

above (based on analyzing each time series individually without grouping into studies).

Thus, big studies contributing many time series might exert an unduly large influence on

the results. For example, nine of the 25 time series measured with PCR (36%) are from

a single study (MAJERCAK *et al.* 2004), but this study is only 1 of 13 (7.7%) that used

PCR to measure *per* mRNA. Given the many complexities of this system, the most

appropriate way to address this problem is unclear. Treating each time series

independently might give large studies too much influence. However, using only a single

value from each study (e.g. the mean), irrespective of how many time series it

represents, could give small studies too much weight and thus risk adding irrelevant and

noisy artificial variance. The latter approach is therefore an extreme approach of

countering potential problems with the former. We used this alterative study-based

approach to explore the robustness of our conclusions about measurement methods,

albeit with the added caution that the loss off statistical power might be too large for

reaching clear conclusions.

Indeed, using only a single average value from each study reduces the remaining

statistical power so much that no result remains significant (see Supplemental Statistical

Analysis). It is worth noting though that some comparisons of variance still showed such

a strong bias that the collection of additional observations might result in significance.

The comparisons that approached significance were these: Variance between PCR and

Northern Blots (*comvar2*, 95% CI: -0.281 to 13.340, with a point estimate for a

difference in variance of 4.926 h^2); Variance between PCR and RPA studies (*comvar2*,

95% CI: -0.161 to 8.426, with a point estimate for a difference in variance of 4.065 h^2).

Such comparisons of non-PCR methods vs PCR-methods were closer to significance

than the non-PCR method comparison of Northern Blots vs RPA studies (*comvar2*,

95% CI: -3.492 to 0.538, estimated difference -0.861 h^2). Differences in variance based

on shared references produced even weaker signals (but see Figure 10 and Discussion

of Logic for a potential boost of statistical power). All corresponding details are recorded

in the Supplemental Statistical Analysis.

                         QQv1r1

DISCUSSION

We discuss the more technical aspects of our work in the context of three broader aims of this study: (i) introducing FlyClockbase and connecting it to our current understanding of circadian clocks; (ii) using FlyClockbase to ask new questions about variability in circadian clock time series, possibly illuminating important aspects of clock mechanisms and methods of observation; (iii) improve and simplify how FlyClockbase, and by extension similar biological information resources (*VBIR*s), are constructed, expanded, and maintained.

Why learn lessons on data? It would be easy to finalize this study without the third aim. However, it is impossible to address the timely and relevant topic of organizing biological data without the concrete context of a specific resource like FlyClockbase. This discussion is relevant because the increasingly data-centric nature of biological discovery has resulted in calls for improved access to existing data (NIH *et al.* 2012; READ *et al.* 2015; WILKINSON *et al.* 2016), which is easier said than done (GOLDSTON 2008; DOAN *et al.* 2012; GITELMAN 2013; HUANG AND GOTTARDO 2013; MCCALLUM 2013). As physical access to data is increasing, the next frontier is defined by the ability to *efficiently* identify datasets of relevance for a given topic. The diversity of biological questions would make any one single resource for all biologists too cumbersome to use. Instead, this aim could be achieved more efficiently by

empowering research communities to construct resources for their own contexts, albeit

using a shared interoperable infrastructure. This infrastructure will be perceived as

useful to the degree it can provide standards that convincingly address common

challenges faced by all biologists aiming to construct *VBIRs* for organizing notoriously

uncertain biological data. Developing such standards is an enormous task that requires

the integration of lessons from many more studies than any single effort such as this

one could produce. Thus, our more modest aim here is to highlight lessons we learned

while constructing FlyClockbase, hoping they will be useful as the biological community

works towards finding more general solutions (NIH *et al.* 2012; READ *et al.* 2015;

WILKINSON *et al.* 2016).

Why we need many *VBIRs*? Efficiently constructing many *VBIRs* like

FlyClockbase is necessary for integrating biological information at the scale needed for

current research. The need is driven by immense challenges, such as mechanistically

understanding and curing cancer (NIH *et al.* 2016; SAMUELS *et al.* 2016), mapping

genotypes to phenotypes in personalized medicine and elsewhere (RODEN 2011;

MACKAY *et al.* 2012; KIRK *et al.* 2015; ASHLEY 2016), or the evolutionary systems biology

goal of mechanistically predicting realistic fitness landscapes (LOEWE 2016). Irrespective

of whether it is possible to realize these broader visions, any serious attempt will require

the diligent construction of many interoperable *VBIRs* that connect well to state of the

art expertise, and advance biological research in the relevant areas. Thus, we will next

examine FlyClockbase in this respect.

FlyClockbase is consistent with current hypotheses

Overall, time series in FlyClockbase are consistent with general published clock knowledge and with mechanisms currently thought to control the clock. Informative reviews of the clock that draw on previously published experimental studies are given elsewhere (HARDIN 2011; ÖZKAYA AND ROSATO 2012). Conclusions from these reviews are supported by FlyClockbase summary statistics given in Table 4.

CLK and other proteins. Increasing amounts of VRI protein between about ZT4 and ZT16 repress *clk* transcription, with an especially pronounced effect after ZT14 (CYRAN *et al.* 2003; GLOSSOP *et al.* 2003). Although CLK protein typically functions as a transcriptional activator for *vri*, the formation of the PER/DBT/CLK/CYC complex represses the activity of CLK between approximately ZT16 and ZT4 (HARDIN 2011). *vri* transcription is therefore decreased, causing lower levels of VRI and reducing the repression of *clk* transcription by VRI. This allows *clk* mRNA to increase and reach a maximum around dawn (ALLADA *et al.* 1998; ÖZKAYA AND ROSATO 2012). *clk* mRNA levels also increase due to the action of the transcriptional activator PDP1 protein, which becomes especially strong around ZT18 (CYRAN *et al.* 2003). Time series in FlyClockbase reflect this pattern of *clk* mRNA reaching a maximum in the morning (peak time = 2.79 *DZT* mean ± 3.16 h *SD*), decreasing throughout the day into the early night (valley time = 14.33 *DZT* mean ± 1.67 h *SD*) and then beginning to increase again in the late night. Although *clk* mRNA does show rhythmic expression, the amount of

CLK protein is not cyclic (Yᴜ *et al.* 2006). Data from FlyClockbase support this constant expression, as the n=5 peaks and n=9 valleys observed for CLK overlap almost completely. However, there might not be enough observations to fully settle the issue, since the variation for both CLK traits (±3.18h, ±4.30h *SD*) is currently lower than expected for peaks and valleys that are all drawn from one stable uniform distribution (±>6.8h; see: Materials and Methods Section, Peak-Valley Section, Randomizing Time Section) .

PER and TIM dynamics. PER protein, *per* mRNA, TIM protein, and *tim* mRNA have expression patterns which are generally opposite to that of *clk* mRNA. Transcription of *per* and *tim* begins mid-morning and is promoted by the transcriptional activator CLK (Aʟʟᴀᴅᴀ *et al.* 1998; Dᴀʀʟɪɴɢᴛᴏɴ *et al.* 1998). FlyClockbase shows the peak time of *per* mRNA at mean *DZT* = 14.61 ± 1.58 h *SD* and of *tim* mRNA at mean *DZT* = 14.59 ± 2.12 h *SD* (using outlier removal method (iii) as in Figure 6; for outlier removal method (i) see Table 4). This supports data suggesting mRNA levels increase through the day and into early evening (Aʟʟᴀᴅᴀ *et al.* 1998; Hᴀʀᴅɪɴ 2011). Protein accumulation reportedly lags behind that of mRNA by about six to eight hours (Zᴡɪᴇʙᴇʟ *et al.* 1991b; Hᴀʀᴅɪɴ 2011), though data from FlyClockbase support a shorter delay of around four to five hours (delay from mean *per* mRNA peak to mean PER peak = 4.03 or 4.84 h, delay from mean *tim* mRNA peak to mean TIM peak = 4.33 or 3.82 h, from Figure 6 or Table 4, respectively). Between approximately ZT18 and ZT4, the PER/DBT/CLK/CYC complex represses the activity of CLK (Lᴇᴇ *et al.* 1998; Lᴇᴇ *et al.* 1999; Bᴀᴇ *et al.* 2000; Hᴀʀᴅɪɴ 2011). This leads to decreased transcriptional activation

of *per* and *tim*, causing a decrease in *per* and *tim* mRNA levels, which is reflected in

FlyClockbase as a mean *per* mRNA valley at *DZT* = 3.61 ± 2.32 h *SD* and a mean *tim*

mRNA valley at *DZT* = 5.09 ± 1.15 h *SD* (Figure 6; difference in variance significant at

5% level). During the day, TIM protein is degraded in response to light (NAIDOO *et al.*

1999; BUSZA *et al.* 2004; OZTURK *et al.* 2011), as indicated in FlyClockbase by an early

mean valley at *DZT* = 5.84 ± 2.53 h *SD*. PER, which is typically stabilized by TIM, is

then destabilized and more prone to phosphorylation and subsequent degradation

(GEKAKIS *et al.* 1995; KLOSS *et al.* 2001; MERBITZ-ZAHRADNIK AND WOLF 2015). This

finding is consistent with FlyClockbase reports of a late PER mean valley = 9.41 ± 1.94

h *SD* (Figure 6). The PER valley is significantly different from the TIM valley (*P* =

1.489×10^{-7}) as determined by the Mann-Whitney-U test. Unless specified otherwise,

similar subsequent tests are two-sided and unpaired. To minimize search time for

readers, we added a few non-significant digits to many results in the main text to help

streamline searches for the context of such results in the Supplementary Statistical

Analyses. As the amount of PER decreases, inhibition of CLK-mediated transcription by

the PER/CLK/CYC/DBT complex also decreases, and CLK resumes promoting

transcription of *per* and *tim*.

FlyClockbase facilitates hypothesis-driven research

Circadian clocks in *D. melanogaster* are molecular systems of substantial complexity which have been inspiring generations of researchers to construct numerous hypotheses about how they work. FlyClockbase can substantially contribute to various life-stages of a hypothesis.

Starting with a blank slate. FlyClockbase can set the stage by integrating existing observations. It is beyond the scope of this paper to discuss the many ways of extending FlyClockbase beyond its current goals, for example, by providing the possibility to add time series from mutants. As FlyClockbase accumulates more observations, its power will increase to help researchers to put new observations into context by comparing them with already integrated data. Such comparisons can inspire new hypotheses and help evaluate them quickly. For example, we discuss below the hypothesis that significant differences in the timing variability of PER and TIM are the result of mechanistic interactions integral to the operation of this clock. This hypothesis grew out of our observation that the difference in variability of the peak time in these proteins was larger than we expected.

Using *Attributes* to compare timing variability. Another way of generating new hypotheses using FlyClockbase is to draw on the many attributes stored for time series. This structured information classifies time series in FlyClockbase in rich ways: many groups of column entries are easily combined into hypotheses for identifying

significant differences between different genotypes, strains, observation methods,

environmental conditions and more. Some hypotheses may not be tested easily, as

statistical significance may often require more data. However, FlyClockbase already has

enough data for testing some hypotheses, and insufficient data might inspire new

experiments for testing important ideas. In this way, FlyClockbase can even become a

tool for planning experiments.

Existing hypotheses on timing. Testing ideas against the data in FlyClockbase

will become increasingly efficient as increasing numbers of experimental results are

integrated into FlyClockbase. This results in a win-win for research productivity. The first

win is clear if sufficient data exists in FlyClockbase to test a hypothesis (saving time). If

the necessary data is not available in FlyClockbase, the second win is highlighting the

potential need for new experiments in areas with limited data. Researchers can then

decide whether to fill this gap with new experiments or prioritize other research. Again,

FlyClockbase can help propose experiments within its scope, which could broaden over

time.

Strength of FlyClockbase. Whether FlyClockbase will contain enough data

critically depends on (i) the ease of adding new studies in a consistent way, and (ii) the

effort required for checking the integrity of any data fragment. These two core

requirements drive our interests in usability and human error analysis as discussed

elsewhere throughout this study. Hence, work with FlyClockbase highlights how various

subtle yet time-consuming issues of data organization, automation, usability, and error

management that are usually classified as "non-biological" can easily become limiting

factors for advancing circadian clock research. Our work on FlyClockbase suggest that

it is more efficient to use the rather systematic approach of automating as much as

reasonable and produce corresponding *VBIRs* in batches. This enables efficiencies of

scale similar to those necessary for completing the human genome project (LANDER *et*

al. 2001; VENTER *et al.* 2001). In similar ways, *VBIRs* could help compare data, evaluate

methodologies, extend current knowledge, stimulate new ideas, test hypotheses, and

create new routes of inquiry. We will next illustrate how FlyClockbase improves

scientific productivity for testing hypotheses in its scope, before returning to practical

questions of usability, appropriate data models, and efficient implementation for *VBIR*s.

Peak timing hypotheses and more: PER variance exceeds TIM variance

We chose to compare PER protein, TIM protein, *per* mRNA, and *tim* mRNA because these components are integral to the circadian clock. They interact with many other clock parts (Figure 1), and null mutants for each gene (*per*[01] and *tim*[01]) lead to arrhythmicity (Konopka and Benzer 1971; Sehgal *et al.* 1994). Also, Table 4 shows that peak and valley observations of these four components were among the most abundant in FlyClockbase and thus best suited for testing differences for statistical significance.

Comparing averages. We first compared the mean peak and valley times for PER and TIM protein and for *per* and *tim* mRNA amounts. Neither the proteins nor the mRNA had significantly different mean peak times. The mean valley time for *tim* mRNA is significantly later than for *per* mRNA (Mann-Whitney-U test, $P = 0.012835$, see Supplementary Statistical Analyses for context; two related tests were shy of significance). It might be reasonable to expect this delay to propagate, such that first the peaks of mRNAs, then the peaks of proteins, and ultimately also the valleys of proteins might show a similar pattern of *tim* preceding *per*. However, this pattern is quickly broken, as the respective pairs of mRNA and protein share essentially mean peak times for TIM and PER that are statistically indistinguishable. Following the cycle to mean valley times for proteins even reverses into the opposite pattern: PER mean valley time is significantly later than TIM mean valley time (see Figure 6; $P = 1.489 \times 10^{-7}$ as reported above). Two related tests also indicated significance for all outlier removal approaches tested (see context in Supplementary Statistical Analyses). These timing patterns defy

the simplistic expectation of merely propagating delays and suggest mechanistic

causes. Demonstrating statistical significance with the help of FlyClockbase suggests

that these patterns might be worth simulating in stochastic models that capture causal

mechanisms and respect the discrete nature of molecules (and resulting variability).

Explaining averages. We propose that the earlier mean valley time for TIM can

be explained by the rapid degradation of TIM in response to light (BUSZA *et al.* 2004).

Still, TIM's peak time at mean *DZT* 18.92 ± 0.94 h *SD* might deserve a closer look. TIM

peaks in the middle of the dark period and not at its very end as might be expected if

light was solely responsible for degrading TIM.

The TIM degradation pathway is well characterized and begins with the activation

of CRY via a light-dependent conformational change (BERNDT *et al.* 2007; OZTURK *et al.*

2011; VAIDYA *et al.* 2013). This change allows CRY to bind TIM in the nucleus (CERIANI

et al. 1999; BUSZA *et al.* 2004). The F-box protein JET then ubiquitinates TIM to promote

degradation by the COP9 signalosome (KOH *et al.* 2006; KNOWLES *et al.* 2009).

Following TIM degradation, CRY is also degraded in response to JET-mediated

ubiquitination (PESCHEL *et al.* 2009). In-depth reviews of the TIM degradation pathway

are given elsewhere (HARDIN 2011; PESCHEL AND HELFRICH-FÖRSTER 2011).

While light-dependent TIM degradation could explain why TIM reaches its valley

before PER, it cannot account for the observation that *tim* mRNA reaches its valley a bit

after per mRNA. The discrepancy in valley time cannot be caused by differences in

mRNA production, as mean peak times for *per* and *tim* mRNA are not significantly

different. We found this irrespective of the test or outlier removal method. Differences in

variance barely exceeded 5% significance, albeit only if we remove outliers by approach

(iii). See Figure 6 for overview and Supplementary Statistical Analyses for details.

Therefore, we suggest considering differences in *per* and *tim* mRNA degradation. These

degradation patterns could be influenced by CURLED (see below).

Comparing variances. We also compared the variability of peak and valley

times for the proteins PER and TIM and for the mRNAs *per* and *tim*. Table 6 reports that

peak time is more variable for PER than TIM as indicated by differences in variance that

exceed $P = 0.05$, albeit only if we remove outliers using our approach (iii) as described

(CARLING 2000). Table 7 and the Statistical Methods present the necessary nuances. In

short, we are confident that the differences in variance that we observe in most time

series are significant, and not easily attributed to:

- *Statistical flukes* for the overwhelming majority of PER/TIM time series reported

  (see Table 6 and Supplementary Statistical Analyses for details on the robust

  bootstrap-based tests performed on the 84% relevant protein peak time series

  that were not removed as outliers by Carling's approach, i.e. 59 of 70 combined

  protein peak times of PER or TIM); or

- *Trivial data errors* or inappropriate data handling at our end (see Figure 3 for the

  substantial effort in checking FlyClockbase for errors that resulted in *Mod5*, which

  was used as starting point for manually re-checking every single time series that

  contributed to our observation of PER-TIM differences in variance and resulted in

  the correction of all errors in *Mod6* as reported in Table 5);

Thus, we reject the explanations above based on our work. In contrast, the potential explanations below for the origins of outliers are more difficult to reject and cannot be tested on a routine basis. It also appears unlikely to us that these explanations contribute more than occasional outliers to typical experimental observations. Therefore H_{basic}, the default basic null-hypothesis for data from FlyClockbase, recommends above that outliers are removed as described by CARLING (CARLING 2000). This method is 'approach (iii)' in our Statistical Methods and was chosen after comparing the features of related approaches for outlier removal as reviewed elsewhere (p. 97 in WILCOX, 2012).

Outliers. About 16% of all protein peak times (3/33 PER, 8/37 TIM) or 4.3% (14/325) of all peak valley traits in Table 6 have been identified as outliers by Carlings method. Including all outliers exhausts the statistical power that FlyClockbase can currently provide for investigating our differences of variance. As a result, statistical significance collapses (see Table 7; $P > 5\%$ in our tests). However, this observation is unlikely to affect our conclusion that some systematic biological mechanisms are probably responsible for producing the PER>TIM variance patterns we report. Indeed, the outliers raise intriguing questions about the sources of their variability. We cannot currently distinguish the following potential sources of variability that will be discussed separately.

Genetic background differences. A substantial minority of flies that are currently classified as wildtype could have circadian clocks with significant genetic differences. This hypothesis might not be as unlikely as it may appear when only considering core

clock genes as shown in Figure 1. Carefully listing *all* genes with potential impact on

clock timing quickly reveals much larger mutational targets. Clock models also depend

on specific rates of transcription, translation and degradation. These processes are

governed by huge molecular machines. Unless otherwise more harmful, mutations that

significantly delay or accelerate these machines will affect circadian rhythms.

Frequencies are unknown, but such mutations in the genetic background of a clock

might occur more often than mutations in core clock genes with similar effects on timing.

If true, these clock background mutations could contribute much to the natural genetic

variation in fly sleep patterns, which can be substantial (HARBISON *et al.* 2009a;

HARBISON *et al.* 2009b; HARBISON *et al.* 2013). In addition, selective pressure on

circadian clocks is substantial (BEAVER *et al.* 2002; SHARMA 2003; YERUSHALMI AND

GREEN 2009). It generates observable latitudinal clines of allele frequencies (COSTA *et*

al. 1992; ROSATO *et al.* 1996; SAWYER *et al.* 1997; SHARMA 2003; SANDRELLI *et al.* 2007;

HUT *et al.* 2013). Evolutionary importance of individual clock components has been

demonstrated for various clock genes (BEAVER *et al.* 2002), including *tim* (SANDRELLI *et*

al. 2007), and *per*, which contains a repetitive region that increases mutation rates for

length polymorphisms. The resulting mutational effects are apparently large enough to

maintain a latitudinal cline (COSTA *et al.* 1992; ROSATO *et al.* 1996; SAWYER *et al.* 1997;

KYRIACOU *et al.* 2007; WEEKS *et al.* 2007; KYRIACOU *et al.* 2008). Thus, mechanistic

differences between the circadian clocks of flies from the wild are likely to exist and may

resurface unexpectedly in clock studies.

   If relevant and substantial, such differences could greatly complicate construction

and parameter estimation in *the* "wildtype *D. melanogaster* circadian clock model".

While numerous models have contributed towards this aim (see Figure 4), there has not

yet been a *single* model that integrates *all* known data on the clock of a *single well-*

defined natural genotype. This ambitious aim becomes much more complicated if

natural variability in clock genes makes time series more variable. Such variability from

natural clock variants could undermine the statistical power of parameter estimation

methods for constructing a *single* clock model for a well-defined genotype.

Controlled observations of all data in a single line of fly descent is – in theory –

an easy way out. However, it might be difficult in practice to observe one fly well enough

to match the statistical power of results from many years of research by many labs.

Such focus on a single genotype could also generate a rather unusual clock model if

one of many rare mutants with large effects is present (EYRE-WALKER AND KEIGHTLEY

2007; HARBISON *et al.* 2013). Developing models of such precision could advance

methods for personalized medicine (HODSON 2016). However, most Drosophila clock

researchers will probably prefer less precise clock models that usually match more

observations in typical flies. Such general models could be inferred by parameter

estimation methods from sets of time series collected in many genotypes by various

methods but excluding outlier time series using the systematic approaches we

employed (CARLING 2000).

Environmental or developmental differences for measured flies. Unrecognized

environmental factors that vary among measurements might modulate genetically

identical circadian clocks. If true, experimental protocols for observing circadian rhythms

in Drosophila could be improved to increase accuracy of biological replicates. Given that

different authors do not necessarily report the same set of environmental attributes, a

first step towards improving experimental protocols might be to develop a standardized

set of *TimeSeriesAttributes* for FlyClockbase that improve the precision of reports from

ongoing studies. It has been demonstrated that age impacts the clock in flies (UMEZAKI

et al. 2012). Environmental factors that affect development in ways that strongly impacts

circadian rhythms could be a potential source of outliers.

Evolution in different lab environments. Consistent differences in selection can

cause flies to follow different evolutionary trajectories and sometimes the results can be

observed in the lab over a number of years (LEROI *et al.* 1994; SHABALINA *et al.* 1997;

HOFFMANN *et al.* 2001; HAAG-LIAUTARD *et al.* 2007; KEIGHTLEY *et al.* 2009). The flies

generating the data in FlyClockbase might have lived in environments with differences

significant enough to trigger some adaptive evolution over a number of years.

FlyClockbase does not yet have enough statistical power to detect significant

differences between strains – if they exist. For example, our initial internal screening

showed no differences between wildtype, color modified strains (*yw* or *w1118*) or other

strains. Currently, FlyClockbase does not have dedicated *TimeSeriesAttributes* for

characterizing the environmental history of flies from the decade leading up to the

measurements. FlyClockbase is ideally positioned for integrating such data, once it

becomes available, and the necessary *TimeSeriesAttributes* have been developed.

However, not all fly clock studies report the LD environment in which their strains

evolved for the previous 250 generations (KANNAN *et al.* 2012). We have no specific

evidence to support the claim that evolution in the lab produced the outliers we

observed. However, some statistically significant evolution of the *D. melanogaster*

circadian clock was observed after applying a relevant selective pressure for 80

generations in the lab (KANNAN *et al.* 2012). Also, note that *per* contains repetitive

nucleotides in its DNA, which result in high mutation rates for repeat polymorphisms

with adaptive significance (ROSATO *et al.* 1996; SAWYER *et al.* 1997).

Human errors. Setting up, performing, or analyzing clock experiments are

complex tasks, as are reporting experimental procedures, relevant labels, or analyzed

data. Such operations are error-prone (see discussion below) and can make

reproducibility a challenge (BAMMLER *et al.* 2005; FREEDMAN *et al.* 2015a). If all 16%

were the result of combining all human errors before publication, then the overall rate

would be surprisingly close to the 14% human error rate that we measured in Table 5,

and corrected before our final test of the hypothesis that PER variance exceeds TIM

variance. Our ability to detect human errors before publication is very limited. Hence, we

took published plots and their attributes at face value. We excluded time series that had

ambiguities we could not resolve (e.g. from poor plot quality), however, this does not

exclude human errors before publication (see Section on human errors in Supplemental

Material).

Conclusion on outliers. Thus, we have no reason to assume that errors before

publication could not have produced some of the 16%. It also seems unlikely that none

of the other causes above could have contributed. Distinguishing between the

hypotheses above is currently beyond the statistical power of FlyClockbase. However,

this might be irrelevant for many of the aims for which FlyClockbase was developed for:

a broader understanding of circadian clocks, often intentionally ignoring the details of

many special cases. Thus, even if we had perfect knowledge of all potential sources of

variability above, we might still want to exclude outliers using a systematic approach

such as the one we employed here (CARLING 2000).

Hypotheses on causes for differences in variance. An obvious explanation for

such differences in protein peak times between PER and TIM could be given by similar

differences in the mRNAs required for producing these proteins. This short-sighted

hypothesis is easily tested using FlyClockbase. It turns out to be demonstrably wrong.

As shown in Table 6 and Figure 6, the increased variance of PER relative to TIM cannot

be attributed to an overall greater variance of *per* mRNA, because the peak time of *per*

mRNA has a significantly lower variance than *tim* mRNA. Therefore, we can rule out

carry-over from similar patterns of variance in mRNA peaks.

Phosphorylation network size. Here we propose that the increased relative

variance of PER can be explained by the larger number of post- translational

modifications for PER (relative to those observed for TIM). Post-translational

modifications such as phosphorylation play a critical role in in the clock (WEBER *et al.*

2011; RISAU-GUSMAN AND GLEISER 2012). While the exact nature and mechanisms of

these modifications have yet to be fully resolved, there is strong evidence that PER

undergoes more post-translational modifications than TIM.

TIM protein is phosphorylated by SGG (KO *et al.* 2010), which promotes nuclear

accumulation of PER/TIM complexes (MARTINEK *et al.* 2001), perhaps by allowing

interaction with the nuclear import protein IMPalpha1 (JANG *et al.* 2015). SGG-

dependent TIM phosphorylation has also been implicated in light-induced TIM

degradation, likely in conjunction with CRY and JET (ROTHENFLUH *et al.* 2000a; BAE AND

EDERY 2006; KOH *et al.* 2006; PESCHEL *et al.* 2009). TIM protein therefore undergoes approximately two to three post-translational modifications.

PER protein, however, could be subject to ten or more post-translational modifications. PER is initially phosphorylated by NEMO, which then promotes additional phosphorylation by DBT (CHIU *et al.* 2011; YU *et al.* 2011). DBT phosphorylates PER multiple times and influences PER stability, nuclear translocation, and SLIMB-induced degradation (BAYLIES *et al.* 1992; EDERY *et al.* 1994; ROTHENFLUH *et al.* 2000a; MARTINEK *et al.* 2001; KO *et al.* 2002; KIM *et al.* 2007; CHIU *et al.* 2008; KIVIMÄE *et al.* 2008; KO *et al.* 2010; CHIU *et al.* 2011; MEZAN *et al.* 2013). PER is also phosphorylated by CK2A, which promotes nuclear import (LIN *et al.* 2002a; LIN *et al.* 2005; MEISSNER *et al.* 2008). PP1 and PP2A both work against these kinases to dephosphorylate and stabilize PER (HARMS *et al.* 2004; SATHYANARAYANAN *et al.* 2004; FANG *et al.* 2007; CHIU *et al.* 2008; GARBE *et al.* 2012).

Mechanistic phosphorylation network stochasticity hypotheses. Post-translational modifications could be opportunities for increasing the variability of timing. This is especially true if a required molecular type only exists in low copy numbers per cell at some relevant stages of a circadian cycle. As described above, PER protein is at the center of a large network of potential phosphorylation patterns and proteins, which also include dephosphorylations. This network of post-translational modifications dwarfs those observed in TIM protein. A large number of different types of potential modifications will break a large population of PER molecules into much smaller sub-populations, thereby greatly increasing stochasticity. The heterogeneity of this network and the relevance of antagonistic forces (dephosphorylation delays degradation)

increase the potential for stochasticity and complicate predictions without detailed stochastic simulations. In comparison, few rather large subpopulations for TIM probably result in copy numbers that are high enough to substantially reduce stochasticity.

Previous simulations have highlighted the possibility of additional variability in the time required for growing to a defined level, when amplification starts from smaller amounts. For example, biochemical systems like signal-transduction cascades that amplify very low molecular counts can easily generate differences in variance for times to reach a peak (LOEWE *et al.* 2009a; LOEWE *et al.* 2009b; AKMAN *et al.* 2010; EHLERT AND LOEWE 2014). The inherent stochasticity of circadian clocks might explain the observed variability via various mechanisms. Potential explanations could include systematic differences in the distributions of the low molecular counts at the start of the respective amplifications. If this does not cause all observed differences in variability of PER or TIM peak timing, differences could be further amplified by the nature of the different reaction networks that generate the peaks of PER or TIM.

Future simulations. While beyond the scope of this present study, we think that such mechanistic phosphorylation network stochasticity hypotheses are worth exploring in reasonably realistic stochastic simulation models.

Expanding hypotheses on CURLED. The inversion of variance differences seen when comparing mRNAs and proteins of PER and RIM suggests that the variability discussed above is probably governed by the post-translational processes described above. However, it is less clear how these processes might explain the significant differences in the variance of valley timing for *per* and *tim* mRNA.

Circadian mRNA degradation might be influenced by CURLED (CU), which is known to affect circadian rhythms. Although *curled* mutants have been known for decades, CU was only recently identified as dNOCTURNIN (NOC), the *D. melanogaster* homolog of the mammalian NOCTURNIN (GRÖNKE *et al.* 2009). NOC has been shown to associate with the CCR4-NOT complex, which promotes deadenylation (and subsequent degradation) of mRNA (TEMME *et al.* 2010). While NOC is thought to influence circadian gene control, specific NOC targets have yet to be identified (GODWIN *et al.* 2013). The gene *noc* produces three transcripts (*nocturnin-RD, nocturnin-RC, and nocturnin-RE*), and NOCTURNIN-RD is rhythmically expressed in DN3s (NAGOSHI *et al.* 2010), a subset of dorsal neurons which are part of the circadian circuit and contribute to evening activity (STOLERU *et al.* 2004). NOCTURNIN-RD knockdown mutants have abnormal responses to constant light exposure, suggesting that NOCTURNIN may play a role in circadian light responses (NAGOSHI *et al.* 2010). GREEN *et al.* (GREEN *et al.* 2007) also noted changes in gene expression in response to a high-fat diet for mutant *Noc*$^{-/-}$ mice, which could implicate NOC in circadian metabolic control. It would be premature to postulate an interaction between NOC and *per* or *tim* mRNA. Instead, we suggest here that NOC and other circadian proteins that influence mRNA degradation might be a fruitful area of investigation, particularly given the connection between NOC, light response, and metabolism.

      QQv1r1

Hypothesis: peaks from PCR methods are more variable

The *Attributes* collected for time series in FlyClockbase can be used to compare different groups of time series for a given clock component based on biological, methodological, or other factors. These comparisons can suggest sources of variability that affect future experiments and the interpretation of simulations. To illustrate this possibility, we compared different measurement methods for observing peaks and restricted our analysis to *per* mRNA time series, which produced 88 peaks, the largest number we could extract from FlyClockbase.

Methods background. Five methods were used to collect at least three independent time series: (i) Microarrays, (ii) RNA-seq and nascent-seq, (iii) PCR, (iv) RNase protection assays (RPAs), and (v) Northern Blots. Each method provides advantages and disadvantages. Historically, Northern Blots were the first of the five methods to be developed. Although they can provide information about transcript size (SHARKEY *et al.* 2004), they have low sensitivity (VANDENBROUCKE *et al.* 2001) and can only be used to analyze one gene at a time (FRYER *et al.* 2002). RPAs were developed after Northern Blots. They can analyze multiple transcripts (SHARKEY *et al.* 2004) and can be used to determine absolute RNA levels (VANDENBROUCKE *et al.* 2001). However, they might have low reproducibility (QU AND BOUTJDIR 2007), and RPA is time-intensive, typically requiring about four days (STREIT *et al.* 2009). All three of the newer techniques (PCR, microarrays, and RNA-seq) are high-throughput methods (BUSTIN 2002; SHARKEY *et al.* 2004; MORTAZAVI *et al.* 2008). PCR, RNA-seq and nascent-seq methods do not

require previous knowledge of specific genes or sequences to be identified (FRYER *et al.* 2002; MORTAZAVI *et al.* 2008). While automation can be difficult for PCR (FRYER *et al.* 2002), microarrays are typically automated. Although RNA-seq and nascent-seq are the most newly developed methods, technical variability may be a concern, particularly when using a low number of read counts (BULLARD *et al.* 2010; MCINTYRE *et al.* 2011).

Sensitivity vs reproducibility. When comparing measurement methods, it is important to consider both the sensitivity and reproducibility of each method; methods that have high sensitivity may or may not have high reproducibility. Northern Blots generally have low sensitivity (GILLILAND *et al.* 1990; WANG AND BROWN 1999; MALINEN *et al.* 2003). RPA is more sensitive than Northern Blots, but sensitivity remains a challenge, especially when using low amounts of mRNA (WANG AND BROWN 1999; VANDENBROUCKE *et al.* 2001). PCR is considerably more sensitive than either RPA or Northern Blots (WANG AND BROWN 1999; MALINEN *et al.* 2003). RNA-seq also shows good sensitivity (MORTAZAVI *et al.* 2008), but RNA-seq sensitivity depends on normalization techniques (BULLARD *et al.* 2010). There is mixed evidence for microarray sensitivity. For example, some researchers found that Northern Blots were slightly more sensitive than microarrays for 14 of 29 assayed genes (TANIGUCHI *et al.* 2001). Six of the remaining genes, however, were detected by microarrays and not by Northern Blots, suggesting microarrays were more sensitive to these genes (TANIGUCHI *et al.* 2001). Older microarrays might only have been able to detect changes reliably if they were at least two-fold (FRYER *et al.* 2002), but some newer Microarrays have become at least as sensitive as RNA-seq (WILLENBROCK *et al.* 2009).

Poor reproducibility may be masked by biological variability. In addition to differences in sensitivity, measurement methods vary in reproducibility, and the procedures used in each measurement method point to different potential sources of variability. For example, RNA-seq requires a small portion of the sample RNA to be used to construct a library, and PCR used to create this library can introduce variability through amplification bias (AIRD *et al.* 2011). However, biological variability typically outweighs methodological variability for RNA-seq (BULLARD *et al.* 2010). A detailed review of the reproducibility of RNA-seq is given elsewhere (SEQC/MAQC-III CONSORTIUM 2014). Reports of reproducibility for microarray studies have been mixed. The 2005 Toxicogenomics Research Consortium raised concerns of variability between laboratories and between platforms (BAMMLER *et al.* 2005), and cross-platform reproducibility issues were echoed elsewhere too (CANALES *et al.* 2006). However, simultaneously a large study by the MAQC Consortium found microarray experiments to be reproducible both across platforms and across laboratories (SHI *et al.* 2006). We expected time series measured with RNA-seq and microarrays to show the least variability, but there was not sufficient data to test this hypothesis.

Experimental causes for PCR variability. A number of different factors have been shown to influence the variability of PCR experiments (BUSTIN 2002). For example, different samples can have different amplification efficiencies (VANGUILDER *et al.* 2008), and, as noted above, the percentage of GC bases can introduce amplification bias (AIRD *et al.* 2011; ORPANA *et al.* 2012). Others noted that, although PCR is often

thought to be a "gold standard", extensive tests showed that calibration and selecting

appropriate primers and probes can be challenging (VANGUILDER *et al.* 2008;

SEQC/MAQC-III CONSORTIUM 2014). A low quantity of starting material can also

influence PCR variability (VANDENBROUCKE *et al.* 2001; BUSTIN AND NOLAN 2004).

PCR and Northern Blot accuracy. Despite these challenges, PCR was

developed more recently than RPA and Northern Blots, and the latter have largely have

largely fallen out of favor, at least partially due to the greater degree of sensitivity

afforded by PCR. Northern Blots are also less accurate than PCR (VANGUILDER *et al.*

2008) and are considered to have low reproducibility (QU AND BOUTJDIR 2007). We

therefore expected to see greater variability in time series measured with RPA and

Northern Blot and less variability in those measured with PCR. However, our analysis

revealed that peak values for time series collected with PCR were significantly more

variable than those from time series measured with RPA or Northern Blot.

Differences between qPCR and RT-PCT? We subdivided time series measured

with PCR into those measured with real-time PCR ("RT-PCR") and those not measured

with RT-PCR ("qPCR"). Neither type of PCR was more variable than the other.

Variability did not significantly differ between qPCR and Northern Blot or between qPCR

and RPAs, but careful inspection of the bootstrap distributions produced by *comvar2*

suggests that this could be merely an issue of statistical power (qPCR has fewer

samples than RT-PCR). Accordingly, RT-PCR was significantly more variable than

Northern Blot and RPA. Finding significantly higher variability for RT-PCR was also

surprising, given that real-time PCR was developed more recently than qPCR, and RT-

PCR is considered to be the standard for PCR, as it decreases experimental error by

requiring less data processing than qPCR (VANGUILDER *et al.* 2008).

Analysis of measurement protocol references using FlyClockbase. We

attempted to explain our observed increase of variability for time series traits observed

with PCR by examining the experimental protocol references cited by each type of

mRNA observation method. It is common for the methods section of studies in

FlyClockbase to reference the experimental protocol of previously published studies.

We hypothesized that differences in these protocol references could explain the

increased variability of time series measured with PCR. The time series identifier

structure, easy access to references, content of respective studies, and the overall

structure of FlyClockbase were instrumental for collecting and organizing information on

experimental protocol references, even though this data was not originally recorded.

The information on method references for *per* mRNA time series is shown in Figure 9.

Table 8 provides overview counts that translate into statistical power when analyzing

each method, either based on counts of time series (bold numbers in Table 8), or based

on counts of studies (non-bold numbers in Table 8).

Shared references are less frequent for PCR studies. Of the three methods

we analyzed, some methods generally cited more references than others. While all 45

time series using RPA cited at least one method reference, such references were cited

by only about 56% (5/9) of the studies using Northern Blots and about 72% (18/25) of

PCR based time series. Some of the method references were cited once, while others were cited more frequently. We defined "shared method references" as references cited by two or more studies which use the same measurement method. Studies using RPA had 93% (42/45) shared method references. 40% (2/5) of studies that used Northern Blots had at least one shared method reference, while only 17% (3/18) of PCR studies had a shared method reference. We hypothesized that decreased variability for time series measured with RPAs and Northern Blots could be attributed to increased number of method references and shared method references. However, our statistical tests found no significant differences between time series with or without method references or shared method references. We therefore suggest that increased variability in time series measured with PCR is not caused by a lack of properly documenting or a lack of using shared protocols but rather stems from actual differences in variability based on measurement method.

  QQv1r1

Explanation: large fluctuations from PCR stochasticity

Here we provide a mechanistic explanation for the increased variance of *per* mRNA peak times as measured by PCR (here brief for RT-PCR and qPCR) in comparison to non-PCR methods (here brief for Northern Blot and RPAs). Briefly, repeated replication required by PCR starts with substantial stochasticity at very low copy numbers before reaching its deterministic exponential growth phase. Non-PCR methods for observing *per* mRNA do not require replication and thus have less potential for variability. Thus, non-PCR methods cannot distort peak timings as much as PCR can.

Exponential growth causes for PCR variability. As indicated above, many different factors can influence PCR variability, including amplification bias, calibration, primers, probe selection, operator experience, and importantly the overall quantity of starting material. It is easy to compare at length potential reasons for variability in PCR and in other methods. We suggest the following simplified analysis that relies on the fundamentally different behavior of timing errors in exponential growth vs linear growth. Such errors generate the larger variance of PCR-measured peak timings. Our explanation requires three basic assumptions:

(i)   Real-world individuals cannot be divided without destroying them.

(ii)   Replication without resource limits inevitably leads to exponential growth.

(iii) Timing of later events in an exponentially growing system are easily affected by an earlier or later start of growth.

These assumptions define implicitly a theoretical model of exponential growth that can

explain the increased variability mechanistically. We then present evidence suggesting

the larger variability of PCR peak times should not come as a surprise.

Great sensitivity and poor reproducibility are linked. The goal of PCR is to

amplify rare nucleic acids by repeatedly replicating in well-defined rounds. During later

stages of growth many molecules are replicated simultaneously. Therefore, any

individual replication event will not significantly impact the overall population, as the

stochasticity of many individuals cancels out increasingly. Ordinary differential

equations work well for such large populations, because their constant violations of

basic assumption (i) are negligible here. The contrast of this precisely predictable

scenario could not be bigger when compared to the very early stages of a PCR reaction

designed to start with low copy numbers for maximal sensitivity. Here basic assumption

(i) severely constrains potentially parallel actions, because single molecules cannot be

broken up without affecting their functionality and are limited in what they can do

simultaneously. This limitation inevitably creates stochastic waiting times that lead to

larger or smaller growth delays that generate the initial timing differences at the root of

cascades of delay that propagate throughout the exponential growth phase due to

assumption (iii).

Amplifying single molecules is a hallmark of PCR's exceptional sensitivity. It also

causes PCR's reproducibility problems for the reasons just explained, making it

extremely sensitive to early rare template numbers. Here timing differences in

polymerase access and replication speed can quickly snowball into faster or slower

growth, and thereby, lower or higher amounts inferred for the original molecules

investigated. These problems are highly relevant for all forms of quantitative PCR,

which are designed to operate completely under exponential growth for better

quantitation.

Circadian clock cycles with small variations of initial amounts inside of cells,

stochastic timing differences, variations in extracted volume, and other factors can

easily conspire to modify final amounts inferred by PCR (if stopped before resources

become scarce). PCR time series measurements rely on these final amounts to define

the PCR end results used for inferring how much may have been present at the

beginning. For this to work as a quantitative method, PCR has to be stopped in the

middle of exponential growth, implying it will inevitably experience substantial noise

from slight variations in the starting conditions under a broad range of circumstances.

Observations of the theory. The strong impact of stochastic timing differences

in exponentially growing systems is easily demonstrated in stochastic simulations of a

very simple exponentially growing population (EHLERT AND LOEWE 2014). There the

initial amount is kept constant for all simulations, making timing differences the only

source of stochasticity. The same principles are responsible for translating the

stochasticity of low molecule counts at the input of sensitive signaling cascades into a

reliably transmitted signal, albeit with variation in the waiting time until the signal is

switched on completely (LOEWE *et al.* 2009a; LOEWE *et al.* 2009b). Thus it is not

surprising if experimental measurements show that different researchers with varying

PCR expertise can easily generate 100-fold differences in their inferred initial number of

molecules at the start of a PCR (BUSTIN 2002). Such variability might stem from small

changes introduced to factors that impact the exponential growth essential to PCR in

subtle, but powerful ways; see (BUSTIN 2002; BUSTIN AND NOLAN 2004).

How this applies to amount peak timing observed by PCR. As shown by growth

mechanism discussed above individual PCR reactions bring individual challenges,

which complicates observations of time series. The main reason is that each time point

measured by PCR will require an independent PCR reaction probably starting with a low

molecular count as obtained from sacrificing an independently running circadian clock.

Thus, observing mRNA clock oscillations by some quantitative PCR method will

inextricably intertwine two processes of variation that inevitably interfere with each

other's observation in these two ways:

(i)   Oscillations of the clock itself may exhibit substantial stochasticity depending

        on molecular amounts involved (AKMAN *et al.* 2010). This implies that the

        peak itself as measured by PCR may vary, even if PCR were perfectly

        precise.

(ii)   Low initial molecule counts of the templates that start any quantitative PCR

        reaction cause substantial inherent stochasticity that can substantially affect

        the final amount of PCR products measured. If this happens, researcher are

        likely to infer corresponding deterministic changes in the initial molecular

        counts (BUSTIN 2002).

Since every single time series observed by any quantifying PCR is inevitably impacted

by both, a substantial amount of random noise is added to each independently

  QQv1r1

observed time point. As a result, several time points might falsely appear to be peaks

(or the converse).

Summary. A low quantity of starting material can influence PCR variability to a

very large degree (VANDENBROUCKE *et al.* 2001; BUSTIN AND NOLAN 2004). Given the

systematically larger potential for measurement noise in PCR methods caused by the

low initial molecule count induced stochasticity, it might even be surprising that PCR is

not more noisy compared to methods like RPA or Northern Blots that do not require

exponential growth.

Model curation for integrating molecular systems biology data

The process of model curation inherently works towards integrating all data that is relevant and available for a given model of interest. Models may be broadly defined as systems, parts, processes or questions that are being represented from certain perspectives to efficiently find particular types of answers deemed to be interesting. We will next briefly discuss, how this view of model curation can facilitate the integration of knowledge-fragments from molecular systems biology in order to enable the emergence of expertise as represented by well-curated systems biology models (e.g. of circadian clocks) or corresponding relevant sets of real-world observations (e.g. of time series in FlyClockbase). We will first look at more specific followed by more general levels of abstraction before discussing other fundamental aspects of model curation.

Related concrete solutions. At a more specific level, there is no shortage of standards, formalisms, approaches, tools and other systems for supporting the application of more abstract categories (like ontologies or models) to concrete problem areas of interest. Examples include the Systems Biology Markup Language for constructing simulation models (HUCKA *et al.* 2003; KRAUSE *et al.* 2010), Systems Biology Graphical Notation for visually representing molecular reaction models (LE NOVERE *et al.* 2009; MOODIE *et al.* 2011), UMLS and SNOMED for defining and using medical reference terms with different approaches to synonyms (MAJOR *et al.* 1978; MERRILL 2009), and specific ontologies for listing existing entities such as 'all genes' in an area of interest (JONQUET *et al.* 2011; MUSEN *et al.* 2012).

At the most specific level are concrete collections of actual models implemented in one of the formalisms described above. For example, BioBase, which collects and curates published SBML models (LE NOVERE *et al.* 2006; CHELLIAH *et al.* 2015). This is closer to the level of FlyClockbase, which collects and curates published time series within its scope. The substantial number of different formalisms for describing models can be very confusing. To get a clearer understanding of essential, non-redundant aspects of model construction it can be useful to consider a more abstract perspective.

Related abstract frameworks. Several abstract perspectives exist. An ontology is a list of potentially existing things. A taxonomy is a list of potentially existing species. A type system is a classification of potentially existing types and how they could be used to compose new types. Similarly, a model is a specification of potentially existing elements in the world of the model. At the most general level, ontologies and taxonomies are fundamentally related (ARP *et al.* 2015). The same holds for type systems, the semantic web, and models in general. All these could be described as 'worlds', as each of these is like a small description of its world. Unless otherwise specified, they buy into the Closed World Assumption, which implies that nothing else exists or matters except for the details explicitly specified. At this abstract level, worlds are all equivalent to systems that encapsulate detailed statements about the conditional existence of sub-systems, items or events that may be nested or composed from defined structures, capabilities, and/or other properties. Such abstractions enable the detection of isomorphisms that can facilitate the transfer of equivalent solutions across problem domains and hence cut development costs by building on results obtained

elsewhere. For example, different elements or types can be grouped into a set in the

contexts of taxonomy, ontology, type system, or model construction. They each may use

different key words to describe this concept, but its core meaning, i.e. semantics, stays

the same. Each of these worlds comes with its own semantic formalism.

It can be challenging to navigate these abstract semantic formalisms for

representing the meaning of statements (VAN RENSSEN 2005). This resulted in the

paradoxical (non-expert) use of 'semantics' as synonym for 'meaningless' in common

language. A semantic model that is genuinely useful to its writer but incomprehensible to

its reader is not useful to that reader and thus appears 'meaningless, resulting in

semantic irreproducibility (LOEWE *et al.* 2016b). The resulting communication failure is a

substantial problem for modeling, programming, giving names and using names (LOEWE

et al. 2016b). FlyClockbase has not been spared; we encountered a broad range of

semantic problems caused by naming, from trivial spelling errors (with non-trivial

consequences in database searches) to profound research questions about the nature

of certain molecules (see discussion of CURLED above). Related questions of naming

and nomenclature are of critical importance for biomedical research; correspondingly

tools that efficiently map local nomenclature to standard nomenclature have been

identified as critically important (NIH *et al.* 2012), and would have made development of

FlyClockbase substantially faster (e.g. by helping to manage changes in local names).

Baseline: conceptual unity of reality despite diversity of experimental methods. Science builds on the physical unity of reality that is observed by different persons using different methods. This principle is usually so compelling that it is unconsciously assumed. It allows scientists to confidently assume the same conceptual unity for aspects of reality that are challenging to study because they may present a different view when investigated by different approaches. This principle of conceptual unity is the foundation of model curation. For example, let Q be the amount of a given protein type in a single cell specified by place and time. Then Q itself does not depend on the various methods subsequently used to measure Q. If results from different methods of observation contradict each other, we can confidently search for errors. The confidence is rooted in the conceptual unity of our world, or any cell, or Q.

Contradictory biological information. While developing FlyCockbase we repeatedly encountered situations where there were contradictions between different observations that appeared to be of equally high credibility. On some occasions, even substantial efforts on our part to check each credible source of confusion we could think of, did not identify any credible information on what could have gone wrong. Such situations are profoundly confusing and cost substantial amounts of time. Handling such difficult situations defines much of the quality of a *VBIR* and its underlying logic formalism (see also Discussion below and in Supplementary Text).

Debugging time limits. We found it important to limit the time we used for attempts to resolve such problems. In this we aimed to be generous yet responsible with our resources. We also started to search for more formal ways of signaling among

curators when a particular set of problematic time series already had been investigated

sufficiently. The implications of this question for the reproducibility debate are unclear.

Should a seemingly solid experimental study be declared 'irreproducible' because an

apparently rushed, ill-conceived experiment failed to reproduce results? Probably not.

Should a seemingly rushed, ill-conceived original study be defended as a valid original

observation, despite the apparent inability of thorough, time-consuming attempts to

reproduce results? Probably neither. However, where is the line between these two

rather extreme scenarios? It is not the task of biological model curators to assess the

credibility of experiments by repeating them. Hence, they need other means of

assessing the quality and relevance of reported observations for the model they curate.

A more differentiated formalized way of communicating various perceived problems

could greatly increase the efficiency of curation work by relieving curators of the implied

unrealistic obligations to always get to the bottom of all inconsistencies or to invent a

reliable taxonomy of resulting errors.

Communicating errors. Developing a formalism for communicating clearly how to

handle errors efficiently is a complicated problem more closely related to compiler

construction than to biological questions. It requires expertise in both areas. We started

to search for efficient ways of how to best represent outcomes of error analyses for

particular time series. We aimed to formalize such communications with the intent to

enable a compiler to exclude certain types of errors from the results of time series

searches. It became increasingly clear that binary choices like "error: yes/no" were

inappropriately simplistic for many real-world uses of data in biology.

Types of problems with data. The discussion above demonstrates that a differentiated approach is necessary for appropriately representing biological data. Not all trustworthy biological expertise is documented by directly observed data and not all data that is available has the quality most researchers would ideally aspire to. Statistical inference and logical deduction from experimental observations are also valuable tools of biological discovery. However, they can only yield conclusions that are as strong as the observations that support them. It is therefore of utmost importance for reliable and reproducible research in biology to represent specific experimental observations and general biological data as they are, including all known limitations and unforeseeable circumstances, confounding variables, or event. From this perspective, almost all data is imperfect to some degree. Imperfection in an imperfect world is not a problem, as long as we are aware of the nature of the imperfections. The current reproducibility crisis reflects in part the complicated nature of reporting the essence of research results concisely, yet without ignoring their limitations or omitting potentially undermining details (BAKER 2016). In our study, we have attempted to be as complete and open as possible, e.g. by conducting a human error analysis for the raw data of our most important conclusions and reporting multiple potential variations of our statistics (see Supplemental Statistical Analysis); this has both substantially increased the length of this report and the time to complete it. As can be seen in the overall structure of our R-script that computed our final analysis, such work often requires exploring various alternative analyses. These all initially appear to be equally valid ways of working around a given imperfection of the data. Substantial parts of calculating all useful tests can reasonably be delegated to a compiler for many frequently encountered scenarios –

assuming there is a formal way of communicating the type of data imperfection to the

compiler.

Using imperfect data. For the reasons above, imperfect biological data is

extremely valuable. Hence, no need to throw out baby hypotheses with imperfect data

bathwater. High-quality model curation considers what can reasonably be learned from

an imperfect dataset by describing as many quantitative aspects as reasonable,

reflecting ideas from the "New Statistics" (CUMMING 2013; CUMMING 2014). Often the

cutting edge of research is defined by situations where not enough high quality data is

available for a final interpretation. In fact, the value of resources like FlyClockbase is

precisely in their ability to synthesize the limits of what is known and highlight

hypotheses that merit further experimentation. Ignoring imperfect data in this context

would be inappropriate.

Imperfection spelled out. Hence, FlyClockbase will require better ways of

representing confusing and uncertain real-world data at the current biological cutting

edge of research. Data there can be aggregated, biased, contradictory, diffuse,

exception-prone, false, generated, gap-ridden, hidden, imprecise, jumbled, limited,

missing, modified, noisy, objectionable, problematic, questionable, redundant, scattered,

swapped, tangential, transformed, uncertain, veiled, washed, wobbly, xeroxed, or

otherwise imperfect. A hallmark of good biologists is their ability to intuitively navigate

these difficulties appropriately in their study systems. The rise of big data has led to

substantial experience in how to deal with imperfect data (MCCALLUM 2013).

Problem type repository. Developing *VBIRs* like FlyClockbase efficiently depends critically on the ability of biological model curators to describe these intuitions in ways that are formal enough, so that an automated solution can be developed eventually. Biological experiences with rates of identifying new species in an ecosystem where many of them exist (GROVE AND STORK 2000) indicate that eventually known species will be resampled. The same can be expected for the varied number of data problems that can be observed during the long-term development of *VBIR*s. *VBIR*s would greatly benefit from a central repository for the logic problems associated with imperfect data. Such a repository can substantially cut costs of identifying logic problems and would help in compiler construction, simply by documenting the extent of the problem. It is difficult, even for experienced biologists, to imagine many of the complications of real-world data in the absence of actual research interactions with real-world data. It is near impossible for complier constructers to do so without also being biologists who work with real-world data.

The role of logic. Communicating errors in clear ways fuels our interests in exploring logic formalisms beyond classic Boolean systems (see also below). Providing a full formal definition for a chosen logic formalism, alongside all appropriate proofs and consistency checks is clearly work best done by theoretical computer scientists with the corresponding formal training. For the reasons discussed above, we do not think that developing an appropriate logic can be reasonably delegated. As indicated above by the idea of problem type repository, we found that one of the biggest challenges was the need to realize the existence of a given problem. Typically, and trivially, biologists find it

easier to specify uncertainties and inconsistencies in real-world observations based on

their experience, while logicians find it easier to identify particular contradictions that

result from an inappropriately defined logic formalism. Our experience suggests that

their combined imagination and expertise will need to be complemented by a slow

careful collaborative review of the detailed problems in a sufficiently complex real-world

research scenario. To facilitate such collaboration, we have described elsewhere the

Flipped Programming Language Design approach (LOEWE *et al.* 2016b), which also

inspired our discussion below of Figure 11.

These complex efforts to develop a sufficiently expressive logic for problems with

biological observations contribute towards answering the next question that is in

principle very simple.

Simple question: how many molecules of a given type exist at a given time in a given cell? Modern biology has invested much effort into developing many diverse

approaches for investigating intracellular quantities of interest. Such quantities often

relate to the simple question of amounts in one of the myriads of forms in which it is

posed in biology today. Generating well-defined, credible answers that properly quantify

all relevant uncertainties would go a long way towards providing the data required for

algorithms aiming to solve the inverse problem (see Models Section). The answers to

this problem quantify the uncertainty of parameters, which are needed for simulating

models of molecular systems in cells. Such simulations can be seen as devices for

extending biologists' thinking capabilities and enable investigating new areas of biology

(see comments on evolutionary systems biology (LOEWE 2016; LOEWE *et al.* 2016b)).

Unfortunately, it is extremely challenging to answer the conceptually simple question above for real molecules in real cells with reasonable quantifications of uncertainty.

Observations in FlyClockbase. Many of the practical challenges of determining such amounts of molecules have been constant companions of our work with FlyClockbase. For example, consider the differences in techniques used to measure amounts of mRNAs or proteins produced by genes such as *per* or *tim*, (see Figure 10 or *TimeSeries AttributeTable* in FlyClockbase for details on methods). While each technique is limited in unique ways, a given quantity of interest can usually be measured in several ways that vary in trade-offs between precision, cost, and other method parameters. As a result, interesting quantities that can be measured in cells have often been measured by dozens of methods, each of which may be implemented by different independent experimental protocols and belong to one of several applicable broader methodological approaches. Each of these may provide different answers to the following practical questions that are highly relevant for model curation. Are amounts of molecules in a single relevant core clock cell of *D. melanogaster* …

- … *absolute counts* (our preferred ideal) or *relative* (usually reported)?
- … from a *single cell* (may be used to infer molecular noise) or from *averaging* over a population of cells (or some other aggregation difficult to disentangle)?
- … *complete and direct raw observations* (enabling independent statistical analyses) or *summaries* of "typical plots" of "the most relevant data" (that can introduce uncontrollable ascertainment biases as observed in other areas, e.g. (AMOS *et al.* 2003; CLARK *et al.* 2005; FOLL *et al.* 2008; LACHANCE AND TISHKOFF 2013; MINIKEL *et al.* 2014))?

- … *appropriately annotated* with all key details for maximizing long-term use in diverse meta-analyses (rare; authors have little guidance on what to report) or *missing annotations* for key details known to exist even if unreported (e.g. fly age, sex), or not reasonably knowable (e.g. fly clocks were disturbed by unexpected and unreported drastic changes in temperature)?

- … *reasonable approximations of reliable results* (as would be expected from diligent analyses of larger and higher-quality data sets if reproduced in the same system or *irreproducible* (recent observations may give reason to pause (IOANNIDIS *et al.* 2009a; SALANTI AND IOANNIDIS 2009; MOBLEY *et al.* 2013; FREEDMAN *et al.* 2015a; FREEDMAN *et al.* 2015b; HALSEY *et al.* 2015))?

The unity of reality implies that similar representational approaches can contribute towards rigorously assessing reproducibility and towards curating heterogeneous and imperfect datasets into an internally consistent *VBIR*: both efforts would benefit from explicitly stating all uncertainties and other problems associated with a given set of observations.

Curation efforts for circadian clock research. FlyClockbase is unique in its scope, datasets covered and many other aspects – as far as we can tell. In particular, we are not aware of other circadian clock time series repositories or meta-analyses that bring similar numbers of replicate time-series or studies together in order to answer questions about the differences in variances of peak times between different components of the core circadian clock of *D. melanogaster*. However, FlyClockbase is not the first biological information repository sharing observations about circadian

oscillations in gene expression. We will next discuss some examples of related efforts; a

review of additional bioinformatics resources relevant for clock research can be found

elsewhere (LOPES *et al.* 2013; LI *et al.* 2017).

　　　CircaDB (http://circadb.hogeneschlab.org) is a publicly accessible web

database storing time series observations that record how gene expression changes in

various mammalian tissues throughout the day (PIZARRO *et al.* 2013). It has been used

for documenting the large extent to which gene expression in mice follows circadian

patterns – with interesting implications for drugs that target the products of rhythmically

expressed genes and that might benefit from timed dosage (ZHANG *et al.* 2014a). While

many genes in the mouse clock are homologous to fly clock genes, there were no

observations of non-mammalian gene oscillations in *CircaDB* at the end of 2016. A

strength of CircaDB is the availability of detailed tissue specific data from mice.

　　　CGDB, the Circadian Gene DataBase (http://cgdb.biocuckoo.org) version 1.0

(as of 2017-01-14) contains information (i) on 1,382 instances where gene expression

followed circadian rhythms as observed by techniques like RT-PCR, Northern Blots or *in*

situ hybridization; (ii) on 26,582 observations of gene expression found in transcriptome

profiling studies to follow circadian rhythms; and (iii) on 44,836 potentially oscillating

genes as identified in a search for orthologs of oscillating genes (LI *et al.* 2017). A

strength of CGDB is its broad coverage of 148 different animals, plants, or fungi. Of the

27,964 genes with experimental evidence of oscillatory gene expression, 3166 have

been observed in *D. melanogaster*. Of these, 14 observations cover all isoforms of *per*

and *tim*, but only 5 of these were recorded in LD. The peak and valley times reported in

CGDB do not contradict those reported by us here; however, the reported sample size

does not have the statistical power to suggest new hypotheses on potential clock

mechanisms.

Deep (machine) learning approaches were investigated for their capacity to

predict time and to distinguish rhythmic from arrhythmic time series (AGOSTINELLI *et al.*

2016). To this end BioCycle$_{Real}$ was curated from 36 gene expression datasets,

including 32 from CircadiOmics (http://circadiomics.igb.uci.edu/) (PATEL *et al.* 2012).

Except for one from the plant *Arabidopsis thaliana*, all datasets came from mice

(AGOSTINELLI *et al.* 2016).

SCNseq (http://wgpembroke.com/shiny/SCNseq/   ) provides access to

temporal transcriptomics of circadian clock controlling cells in the suprachiasmatic

nucleus of the mouse brain at unprecedented precision (PEMBROKE *et al.* 2015).

Bioclock (http://www3.nd.edu/~bioclock/) is a repository of circadian

transcriptional profiling data from *Anopheles gambiae* and *Aedes aegypti*, mosquitoes

acting as vectors for malaria and yellow fever, respectively (RUND *et al.* 2011; RUND *et*

al. 2013; LEMING *et al.* 2014).

BioDare (http://biodare.ed.ac.uk/ ; http://millar.bio.ed.ac.uk/data.htm) is an

online service for data-sharing and analysis of circadian time series observations. It's 10

datasets from *A. thaliana* were used for comparing period estimation methods and other

clock research (ZIELINSKI *et al.* 2014).

dbCRY (http://www.dbcryptochrome.org/) facilitates comparative genomics of

crypotchromes, the light-sensing proteins in clocks (KIM *et al.* 2014); see Figure 1.

Diurnal 2.0 (http://diurnal.mocklerlab.org/) provides access to observations of circadian genome-wide gene expression patterns observed in several common model plants (MOCKLER *et al.* 2007).

EUCLIS (http://www.bioinfo.mpg.de/euclis/) is the 'EU Clock Information System'. It adapted an advanced database architecture from another systems biology project for circadian clock researchers in order to combine modules for experimental data, clock models, and a related digital library (BATISTA *et al.* 2007; LOPES *et al.* 2013).

Individual meta-analyses occasionally integrate different datasets in an effort to increase the statistical power and reliability of conclusions. For example, combining and curating data from five independent microarray studies in *D. melanogaster* confirmed the rhythmical expression of 81 transcripts while also identifying 133 new cycling transcripts (KEEGAN *et al.* 2007). To arrive at their conclusions, KEEGAN *et al*. had to obtain data directly from the authors of the microarray studies they analyzed as not all necessary data was available online (CLARIDGE-CHANG *et al.* 2001; MCDONALD AND ROSBASH 2001; CERIANI *et al.* 2002; LIN *et al.* 2002b; UEDA *et al.* 2002). In turn, the same happened with their results: "All data used to produce this report are available upon request. Files that contain the individually formatted results from each of the original reports were too numerous and large to be included with this manuscript …" (KEEGAN *et al.* 2007). Some (non meta-analysis) studies that generate substantial amounts of new data put in the substantial additional work necessary for making material available online (e.g. see http://biorhythm.rockefeller.edu/ (CLARIDGE-CHANG *et al.* 2001)). Merely *storing* complex data in one or more file archives online is usually easy, but organizing and documenting complex datasets for use by independent

researchers is not. This requires semantic reproducibility, which can quickly become

prohibitively complex (LOEWE *et al.* 2016b) if no existing conventions are shared with

users. Projects above have used database technology and/or web interfaces as shared

conventions facilitating communication; as argued elsewhere in our study, this is neither

ideal for all biologists nor for all work in biology. These problems are less acute for

studies that can fall back on using public repositories with an appropriate data format.

For example, a functional analysis study of fly genes expressed in response to the light-

induced resetting of the circadian clock (ADEWOYE *et al.* 2015) stored most data at the

NCBI-maintained Gene Expression Omnibus database (

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE39578); NCBI GEO offers

semantics particularly well suited for describing typical microarray datasets (BARRETT *et*

al. 2005; BARRETT *et al.* 2011; BARRETT *et al.* 2013; CLOUGH AND BARRETT 2016).

Generic model organism repositories. Resources like the individual-study

repositories above, meta-analyses, and circadian research specific repositories are

complemented by generic model organism resources such as FlyAtlas

(http://flyatlas.org) for gene expression information (CHINTAPALLI *et al.* 2007;

CHINTAPALLI *et al.* 2013a; CHINTAPALLI *et al.* 2013b; ROBINSON *et al.* 2013) and FlyBase (

http://flybase.org/) for genomic and other information (ASHBURNER AND DRYSDALE 1994;

ST PIERRE AND MCQUILTON 2009; GRAMATES *et al.* 2017).

Why curate circadian data? Without detracting from the important achievements

that continue to be enabled by the resources specified above, some points are worth

noting for further discussion. Scarcity is first. We attempted to be as inclusive as

possible and added resources far beyond the focus of our study on core clock genes in

flies; recent reviews (LOPES *et al.* 2013; LI *et al.* 2017) do not list much more. Yet

rhythmic gene expression is pivotal for fitness, health and more. This is underlined by

estimates of 10% or 43% of the expressed genome under rhythmic control (BOYLE *et al.*

2017) or of all protein coding genes showing circadian expression patterns somewhere

in the body (ZHANG *et al.* 2014a), respectively. Given this importance of circadian

biology, it seems surprising that not more circadian clock related repositories exist. One

potential reason is that the tools for managing data are inadequate and discourage

many biologists from getting involved. Secondly, the existing resources are scattered,

very heterogeneous with respect to their data structures resulting in poor

interoperability, and they are haphazardly reorganized (e.g. web addresses move,

internal structures are modified).

   Challenges. Circadian clock researchers have repeatedly stated over the years

that the arrival of new data due to experimental advances creates new challenges for

data management and data processing (ZIELINSKI *et al.* 2014). These could be

addressed by the development of a new infrastructure that standardizes data and

software to reduce re-implementation efforts, improve documentation, and increase

collaboration by sharing data (BATISTA *et al.* 2007; ZIELINSKI *et al.* 2014). This need for

improved and simplified infrastructure exists for systems biology in general (CASSMAN

2005) and similar ideas have been echoed in the debate about reproducibility (BUCK

2015). The underlying issues have not yet been solved on a broader scale (NIH *et al.*

2012; NIH 2015; NIH 2016; WILKINSON *et al.* 2016). A recent review of additional

bioinformatics resources pointed out that tools cannot replace researchers, because it is

"often necessary to conduct an evaluation of the results of a data mining effort to

determine the degree of reliability" (LOPES *et al.* 2013). Indeed, experiences at UniProt show that "expert curation is by far the most reliable method to report gold-standard information and provide an up-to-date knowledgebase containing experimental information" (THE UNIPROT CONSORTIUM 2017). We argue that much of the low-level work of ensuring reproducibility and adherence to formal standards could be handled reliably by a compiler that transparently executes well-defined recurring tasks (see Discussion below and in Supplemental Material).

Perspectives on biological model curation. Circadian clocks control rhythmic gene expression for a substantial and important fraction of the genome, approximating half of all genes in mice (ZHANG *et al.* 2014a). It is thus difficult to isolate clocks from the rest of the organism they govern. An overall assessment of the impact of circadian clocks might thus require simulating whole cells or even whole organisms. This perspective raises several questions.

Will this scale to real cells? The challenges of reproducibility for systems that are comparatively small are multiplied on much more complex systems, such as the molecular systems biology simulations of whole cells that have started recently at a larger scale (KARR *et al.* 2012; KARR *et al.* 2013; LEE *et al.* 2013; PURCELL *et al.* 2013; SANGHVI *et al.* 2013; KARR *et al.* 2014; KARR *et al.* 2015a; KARR *et al.* 2015b). The ultimate aim of such studies is to understand in detail, how a real cell work and evolves over time (LYNCH *et al.* 2014). However, the question is, whether our tools will be able to scale in such a way that errors can be kept down and our toolchain remains reliable. Hence, outstanding reproducibility of smaller models and datasets are a prerequisite for

any further integration. We have chosen to focus on the simplest possible

implementations when developing *VBIRs* to enable durability.

Will tool development overwhelm biological goals? The essential requirement of

tools that handle biological data more accurately and with more ease could continue to

bind disproportional amount of resources through a lack of coordination (see eg.

(CASSMAN 2005)). Usually only one excellent tool for a given task is needed, not several

that are usually good enough but break down in some special cases, which then require

completely new implementations. It is encouraging that the accuracy of computational

tools in some areas converges towards that of high-quality experiments (e.g.

(LEJAEGHERE *et al.* 2016)); likewise, the development of more precise higher-level

abstractions simplifies much of the lower-level programming (e.g. (ARP *et al.* 2015)).

However, the need for new and more precise tools is vast, and only few biologists can

program well enough to contribute. Thus, support from computer scientists and

professional programmers will certainly be needed. However, without the extraordinarily

close collaboration described here it will be extraordinarily difficult to develop tools that

are efficient enough in real biological research in order to drive adoption. Only then will

tool development start to contribute to the overall biological goals. In our analysis, the

development of a VBIR compiler is a particularly efficient way of tool development (see

Discussion below). The efficiencies from compiler development might help with raising

the funding for VBIRs on a more permanent footing (EMBER *et al.* 2013).

Simultaneously, experimental methods, their limits, and associated errors and biases

will require more rigorous analyses in order to contribute towards a more accurate

description of the precision associated with the actual observations (e.g. for sequencing errors see (ROBERTS *et al.* 2013; ROBERT AND WATSON 2015), for n-fold gene expression see (CANALES *et al.* 2006; CANALES 2016), for PCR see (BUSTIN 2002; BUSTIN AND NOLAN 2004; VANGUILDER *et al.* 2008), for tests of a parameter estimation method, e.g. see (DAIGLE *et al.* 2012); many more analyses for other methods are needed).

Will the biology try to advance too fast for its own good? The 1970's saw the rise of systems theory in ecology, albeit arguably too early (WOLKENHAUER 2001). Now systems biology has in principle a computational method at its disposal for every single step along the causality chain from genotype and environment to phenotype and fitness (LOEWE 2009; LOEWE 2012; LOEWE 2016). However, what does not exist at the moment is a rigorous and integrated problem management for the full causality chain. Clearly, more uncertain output at more causal calculations will combine with additional uncertainties at more consequential calculations. If this accumulation of uncertainties occurs on the long causal chain from genotype to phenotype, then it is presently not clear, which signal-to-noise ratio is to be expected. This question can only be resolved by an integral management of uncertainties similar to what we propose. Advancing simulations of whole cells or even organism too fast without allowing for appropriate precision to grow in method development and curation might cause rigorous scientists to lose patience, throw out the baby with the bathwater, and thereby cause unnecessary setbacks.

Balance. It is neither possible nor necessary to manage either of the extremes above beyond being aware of them, to avoid falling into either trap. The dynamic nature

1   of biological research will then run its course. However, any foreseeable scenario will

2   have a very large need for biological model curation, which will require many well-

3   equipped biologists, as high-quality model curation will always remain a human task.

4   Similarly, in any credible scenario, biological model curators will greatly benefit from

5   support by a well-equipped VBIRs compiler.

Towards a compiler for advancing FlyClockbase and biology

Working with FlyClockbase has given us ample opportunities to observe first-hand many

diverse problems that frequently complicate an otherwise efficient use of computers or

formal simulation methods for advancing biology. Below we highlight a few key

observations suggesting a greatly increased efficiency for integrating computers in the

workflow of FlyClockbase and similar *VBIR*s could be enabled by the construction of a

corresponding *VBIR* compiler. This requires paying the moderately increased cost of

constructing such a compiler only once. In return the whole biological community could

substantially cut the excessive costs of manually constructing or maintaining *VBIR*s.

The substantial costs of *VBIR*s construction conflict with the growing need of compiling

thousands of *VBIR*s that integrate in computable form the biological expertise

necessary for engaging the grand challenges of our time. Leveraging abstractions

developed in computer science for cutting through the complexities of data

management with the help of an appropriately designed compiler could greatly reduce

the costs of integrating biological expertise in order to address grand challenges more

efficiently.

    History. Similar thoughts about better computing for biological discovery have

been recurring since the dawn of computing (TURING 1936), fueling many discussions of

chances and challenges in diverse areas and applications, including the following

examples: harness the precision of logic for biological discovery (WOODGER *et al.* 1937),

simulate genetic systems (CROSBY 1973), open science (BARTLING AND FRIESIKE 2014),

improve reproducibility (IOANNIDIS 2005b; DONOHO 2009; HUANG AND GOTTARDO 2013;

LOEWE AND KEEL 2014; STODDEN *et al.* 2014; FREEDMAN *et al.* 2015a; JAMES *et al.* 2015;

STODDEN 2015; BARBA 2016; LOEWE *et al.* 2016a; LOEWE *et al.* 2016b), share data

(WILKINSON *et al.* 2016), and do so efficiently at larger scales (NIH *et al.* 2012; NIH

2015; NIH 2016). To contribute to this debate that also affects FlyClockbase, the next

sections distill the essence of selected key challenges we observed. We connect our

observations to relevant research in other disciplines to reduce rediscovery where

possible.

Unusual approaches to constructing an unusual compiler. We will conclude

that many or most of the problems below could be solved efficiently by carefully

constructing a corresponding compiler. Its specialty is to facilitate the implementation of

best-practice solutions for constructing VBIRs and addressing the many challenges

which biologists regularly face when they aim to use computers for advancing their

research. Our approach goes far beyond superficial reassignments of responsibility;

rather it proposes that broad classes of problems in biology could benefit from

computational solutions if the latter are designed with enough time and care for those

abstractions that matter for real-world biology. Figure 11 illustrates key aspects of the

development process we used. It represents a very unique informal blend of two

opposite extremes in software development. The extreme known as 'agile methods'

advocates for quick iterations that implement tangible improvements. Its successes

have made it popular, but it is not without a dark side that can stifle the development of

innovations and strategies needed for solving more complex problems (JANES AND SUCCI

2012; ANNOSI *et al.* 2016). The other extreme approach to software development is

known as the 'water-fall method'. It emphasizes thorough planning of various stages,

and clearly separates developing a design from implementing it. As captured in Figure

11, data integration in FlyClockbase followed faster (internal) release cycles. Questions

of fundamental importance with implications for formal or theoretical aspects of compiler

design followed a much slower timeline. This allows us to focus our implementation

resources on the most promising and efficient formalisms, and avoid the need for

implementing potential solutions that appear attractive for some time, but are replaced

by the need for further improvements. In such situations, overall speed of compiler

development probably benefits from manual VBIR curation, since this allows the

compiler design the time it needs to mature. Working in a research setting, as we did

with FlyClockbase, creates additional challenges, simply from the unpredictability of

research. Classifying potential bugs in compiler construction can be seen as a problem

similar to the development of the taxonomy of beetles: both exist in exceedingly larger

numbers, and continued random sampling eventually leads to re-encountering similar

bugs. Constructing a compiler that can deal with biology's uncertainty and complexity in

a stable and reliable way requires a very extensive sampling of these potential bugs (i.e.

logical program inconsistencies). We have developed the Flipped Programming

Language Design Approach in order to address this problem using repeated rounds of

rigorous review of proposed compiler designs from multiple usability and domain

experts (LOEWE *et al.* 2016b). Our work on FlyClockbase benefitted from this approach

and also contributed to its development. It illustrated for us, how developing good

abstractions can take a very long time, and how much finding them worth the effort.

Place   FIGURE 11   about here.

Cost of *not* constructing a *VBIR* compiler. Research on circadian clocks in

flies can be used to illustrate some of the cost to biology if no *VBIR* compiler is

available. As explained above, time series observations are extremely valuable for

inferring mechanistic models of clocks in flies. Yet, in the last 25 years, the vast majority

of models of the core *D. melanogaster* circadian clock have been based on *abstract*

clock features, such as the response to light, the period and presence of oscillations.

We conducted an extensive search for such models, and only three of the 66 models

identified used specific experimental time series to inform parameters (Figure 4). Even if

combined, these models only used a small fraction of all studies with time series data

they could have used (Figure 5). Specifically, parameters study A (FATHALLAH-SHAYKH *et*

al. 2009), B (LEISE AND MOIN 2007), and C (KUCZENSKI *et al.* 2007) were based on time

series from, respectively, one study (KADENER *et al.* 2007), three studies (LEE *et al.*

1998; BAE *et al.* 2000; SHAFER *et al.* 2002), and 11 studies (HARDIN *et al.* 1992; ZENG *et*

al. 1994; SEHGAL *et al.* 1995; SO AND ROSBASH 1997; BAE *et al.* 1998; LEE *et al.* 1998;

BLAU 1999; BAE *et al.* 2000; KIM *et al.* 2002; CYRAN *et al.* 2003; GLOSSOP *et al.* 2003).

The sparse use of directly observed experimental evidence such as time series is

understandable in light of the many challenges that complicate the integration of messy

biological real-world observations into the abstract mathematical models that are often

extremely simplified to facilitate their mathematical analysis. In addition to such

conceptual problems, deceptively simple problems such as the storage and

organization of very heterogeneous, imprecise, noisy and contradictory experimental

datasets can easily create insurmountable practical challenges for directly using

experimental time series data to inform parameters in models.

FlyClockbase substantially lowers this barrier by providing a nucleus for

collecting, organizing, and curating relevant time series and their many potentially

informative *Attributes*. If increasing numbers of experimental time series are deposited

in FlyClockbase and its organizational structures keep pace with this growth, then future

modeling studies could be structured in a way that enables the automated improvement

of some types of models in response to the submission of new data. Such data handling

capabilities are likely to enable the investigation of new biological aspects of circadian

clocks that are beyond practical limits of the complexity manageable by current tools. If

a reasonably well-working VBIRs compiler had been available for a long time, then the

substantially lower barrier to the development of a resource like FlyClockbase would

most likely have resulted in a more comprehensive use of hard won experimental data

in theoretical models. Even where datasets have been compiled and published under

open access, an unstructured way of storing them can very quickly make it prohibitively

complicated to keep them up-to-date on the longer-term (e.g. (WHITE *et al.* 2013; SUPP

et al. 2015a)). Such problems also pose challenges for citizen science projects (e.g.

(LOEWE 2007; SUPP *et al.* 2015b)). Even if computational results are fully structured from

one perspective, the lack of appropriate data structures for analysis from another

perspective, can create prohibitive barriers for some research (LOEWE 2002).

Counter intuitive challenges and other work. However, before such a vision of

biological research can move closer to reality, a number of counterintuitive challenges

will need to be addressed. Since an appropriate discussion is beyond the scope of this

paper and more details are given in the Supplemental Material, we will merely touch on

the tips of several icebergs below by discussing a few illustrative examples. Despite

great and sustained progress in logic research, much remains to be done to improve the

expressive power of known systems of logic (SMITH 2008).

More biological precision requires handling more imprecision more precisely. The

ultimate aim of FlyClockbase is to improve the precision of circadian clock models by

making all relevant data easily available for parameter estimation tool. Accomplishing

this goal goes far beyond compiling the data. It requires entirely new approaches for

dealing with uncertainty, imprecision, contradictions, gaps, and numerous exceptions

created by the astonishing diversity of methods used to observe biological systems.

These challenges belong to the many predictably unpredictable surprises that will be

encountered by any efforts for constructing *VBIR*s of substantial complexity, such as

FlyClockbase. Accurately describing biological observations made in this often-

confusing context is a substantial challenge we encountered while developing

FlyClockbase. We found that it is not enough to 'describe the issues in words'; this

would merely create additional free-text repositories with unstructured information,

maybe with a bit more focus than a corresponding collection of PDF-files with the full

text of the study. Such free texts could not have enabled us to search efficiently for time

series. We found it extraordinarily useful to have key information in a more structured

form, e.g., to compare mRNA measurement methods (see above). However, such

structure must not come at the expense of the ability to efficiently represent newly

encountered imprecision or data. We found that a working biologist with sufficient

domain expertise is the best expert for choosing how to handle newly encountered

information: ignore, describe in unstructured comments, or create corresponding

Columns in an *AttributeTable*. Without a substantially sophisticated and extendable

system for dealing with imprecisions, very little information will become available for

automated processing in more coordinated ways. Ignoring such problems may be

reasonable in some cases, but eventually, the inability to handle such imprecisions

correctly will artificially narrow distributions and create illusory precision that wrongly

rejects simulations as unrealistic and can unnecessarily complicate parameter

searches. Conversely, allowing for too broad a margin of error can easily result in a

misleading model caused by biologically unrealistic parameters. Thus, appropriately

managing errors and uncertainties in observed time series is one important key to

improving mechanistic models of circadian clocks informed by the real-world time series

in FlyClockbase.

Logic in gene regulatory networks. The classical Boolean logic of compilers and

gene regulatory networks share an unexpected connection if the input, output, and

every step in between are well approximated by just two states (KARLEBACH AND SHAMIR

2008). Thus, compilers could provide unexpected help for modeling gene regulatory

networks. If provided with the right details, compilers could also automatically detect

situations where gene regulation becomes stochastic due to low molecule counts in a

cell (MACNEIL AND WALHOUT 2011). The help of compilers that automatically analyze

complex logic constructs correctly could prove essential for understanding how

complicated binary gene regulatory networks behave if under the control of the daily

rhythms of a circadian clock (LOWREY AND TAKAHASHI 2004; DOHERTY AND KAY 2010;

ZHANG *et al.* 2014a). Logic modelling of various genetics problems has a long history

(COTTERMAN 1983; OPITZ 1983; CROW 2001) and a bright future helping us to

understand many diverse aspects of gene networks and their modular structures (MITRA

et al. 2013; LE NOVERE 2015; SAEZ-RODRIGUEZ *et al.* 2016).

Formal systems of logic are usually not logical enough for biology. This

paradox is easily resolved by contrasting the complexity of biology with the simplicity

typical for formal logic systems. Errors of omission in formal logic limit its ability to

express corresponding biological statements efficiently. Omissions have been found to

be among those types of errors that are most difficult to detect (PANKO 2016).

Our own work on FlyClockbase confirms the substantial frequency and cost

associated with errors of omission. Table 5 reports a substantial discrepancy between

two types of error rates observed during our exhaustive in-depth re-check of each time

series that could in principle affect our main conclusions regarding the variances of

peak timing that differ between PER and TIM. Some types of errors could be

characterized as 'simpler errors' like obvious swaps or typos in FlyClockbase itself. As

expected (PANKO 2016), these simpler errors occurred at much lower rates (affecting

cells of spreadsheets at rates just below 1%). In contrast, we detected a bit over 10% of

all time series when re-checking our meta-analysis for systematic errors such as

inadvertently omitting agreed-upon steps from routine analyses by trained curators. Much work in our study went into ensuring that important rules were indeed implemented in all applicable cases.

The unlimited potential of omissions to confound biological results repeatedly creates 'important biological investigations' aiming to determine whether a given biological conclusion might have been compromised by faulty logic (see our own examples above, where we excluded too much data). While these investigations can be essential for progress in biology, execution often involves excessive, tedious, 'non-biological' work towards finding elusive 'needles in haystacks.' This metaphor easily takes on ever more complicating levels of nesting when the logical 'needles' in question actually consist of errors of omission; finding them can be as challenging as identifying entirely new logical blind spots for the first time. These challenges were felt throughout the development of FlyClockbase from start to finish on numerous occasions, which were too many to track beyond a few illustrative examples.

Data quality, plot quality, and task completion. One specific time series figure seemed such a perfect interpretation nightmare that preventing publication of figures like it might be counted as a donation towards supporting FlyClockbase. This figure provided the initial illusion that it shouldn't be too difficult to unambiguously decide which data point belonged to each of its different time series. However, when actually attempting to extract the values, it slowly became clear that achieving unambiguity was impossible because the figure had been irreversibly degraded to the point where its semantic reproducibility was no longer complete (LOEWE *et al.* 2016b). The reason was an unlucky combination requiring *all* the following factors controlled by different entities:

(i) the authors' decision to combine all these time series into a single plot, (ii) miniscule, similar plot symbols, chosen by the authors or the plotting software, (iii) a very low scan quality for the figure, chosen by the publisher, and (iv) substantial overlaps of some of the time series, chosen by nature.

Cost. Aiming to collect all relevant data, we did not want to allow for arbitrary decisions that excluded plots due to a curator's fleeting perceptions of potential difficulties. It took several independent rounds of revisiting by the three most experienced curators and repeated discussions among them before all agreed on the irresolvable nature of this figure's ambiguities. In total, we spent more than four hours trying to resolve ambiguities (excluding time for finding and initial digitizing). In contrast, it might only have required 10 min of the person producing the figure to choose plot settings that would have completely eliminated more than four hours of work for us. From this we concluded: (i) FlyClockbase needs a reliable mechanism to help future curators avoid such known time killers (e.g.: 'KK' in POST LOEWE *et al.* 2016b) in the absence of substantial new information. (ii) Efforts such as FlyClockbase need to find principled ways for protecting their limited time resources against irresolvable ambiguous plots or dataset, without delegating decisions on the inclusion of data to a moment's fleeting perceptions of a single curator. (iii) It might pay huge dividends across all sciences if a targeted effort could improve the clarity of plots produced by typical default settings.

Implications for Logic. This example highlights the recurring observation that managing challenging tasks like the one discussed above might benefit from two different dedicated *BioBinary* values, one for 'progress of the work', one for 'results of

the work'. Using the *OKScale* in FlyClockbase, a 'Progress BioBinary' could store:

Progress: *KO* (not started), *OKO* (working), *OK* (done), *MIS* (incomplete because

problems occurred). Similarly, a 'Result BioBinary' could store: Result: *KO* (has errors),

OKO (intermediate), *OK* (completed), *MIS* (missing).

How to detect logic errors in FlyClockbase. The detection of logic errors can

be greatly accelerated by open discussions that invite outsiders to share their

observations freely. This could greatly improve the quality of FlyClockbase if this could

be made efficient. One of the most notorious bugs is error by omission. This is equally

true for omissions in typical program source code as it is in the analysis of biological

observations. Omissions are hard to find anywhere (PANKO 2016) and can affect the

reproducibility of results (HUANG AND GOTTARDO 2013) at great cost to science

(FREEDMAN *et al.* 2015a). Clearly, a well-defined formal system of logic that is capable of

handling biology's complexities would be a great asset for FlyClockbase and VBIRs in

general. It's formal axioms and rules would exclude many options as impossible,

thereby helping researchers to save time by avoiding many fruitless investigations and

focusing attention on options of likely interest. Pivotally, such a logic would achieve

these aims by providing a conceptual base-line for establishing the few potentially real

omissions that need to be considered, while keeping infinities of useless speculations

and contradictions from interfering with productive research. If correct, such a logic

would be extremely useful for FlyClockbase. However, to the degree that it is

misleading, its adoption would waste critical research capacity and lower the chances of

uncovering such problems, because people in general rarely question an adopted logic,

even if misleading. Thus, using *any* formal logic for understanding complex real-world

biology is a double-edged sword, which cannot be avoided when studying mechanisms, including those of clocks in flies.

The idea that a formal logic for biology could facilitate biological research was first expressed only a year after Turing defined the essence of computing (TURING 1936; WOODGER *et al.* 1937). Yet, general mutual inspiration aside, it has been difficult to develop a more general formal logic that "makes better logical sense in biology" Most researchers strongly prefer to collaborate on much more specific questions they understand comparatively well, and experts in logic do not often engage with the uncertainty of biological observations. Hence, it has been much easier for most researchers to produce successful special-purpose computing tools for biology, than to arrive at more general solutions. For example, a sorely needed general-purpose programming language designed by biologists for biologists is not available despite all research in bioinformatics, computational biology, and systems biology so far. A credible effort to produce such a language, requires experienced experimental biologists as prime partners on the very same table, where expert logicians design the formal aspects of a logic for biology in numerous iterations. The expressivity of such a logic needs to be tested by its ability to represent actual real-world wet-lab or field-expedition observations. Working on FlyClockbase as described in Figure 11 provided us with such a rare opportunity.

Identifying omissions in the logic of a complex system does not necessarily provide the right resolution and exceedingly many partial workaround solutions are usually found much faster. Such quick-fixes offer immediate relief, albeit at the cost of increasing accidental and historic complexity inessential to a system's function

(RAYMOND 2003). Without mechanisms for removal, the accumulation of such special case stop-gaps will eventually increase the complexity of a system until it collapses under its own rules. At this point, new potential users will no longer be able or willing to invest the time needed for learning how to navigate the system's idiosyncrasies. FlyClockbase will not be able to escape this eventual fate, if its data model is not carefully guarded against these problems. Inessential complexity creates numerous difficulties in many contexts, which include defining programming languages, logics, or type systems in computer science (PIERCE 2002), rules of operator precedence (RAZALI *et al.* 2015-10-26), designing data models, databases, or data integration frameworks (DE TRE *et al.* 2004; DOAN *et al.* 2012), maintaining a user-friendly organization for large libraries, information resources or hard drives, and constructing ontologies, taxonomies, modeling frameworks, or query languages (VAN RENSSEN 2005; ZEIGLER AND HAMMONDS 2007; RUBINSON 2014; ARP *et al.* 2015; HAZBER *et al.* 2015). With a bit of abstraction, the shared root of these problems might be summarized by asking: "What is the best description of a complex world with all its possibilities for nesting, linking, traveling, and communicating?" The relevance of such logic research for FlyClockbase, is that it greatly simplifies managing the complexities of consistently handling inconsistent biological data in FlyClockbase.

Error handling in the face of uncertainty. No *VBIR* of sufficient complexity will be free of errors. This certainly also applies to FlyClockbase. The question is, how to handle errors. Problems with tracing the identity, availability, accuracy, precision, and reliability of data have been the topic of numerous investigations in various contexts,

some of which involve big data (e.g. see (REASON AND MYCIELSKA 1982; REASON 1990;

REASON AND HOBBS 2003; GOLDSTON 2008; DOAN *et al.* 2012; GITELMAN 2013; GRIMES *et*

al. 2013; MCCALLUM 2013; REASON 2013; BLANKENBERG *et al.* 2014; REASON 2015)).

Opportunities. FlyClockbase presented us with excellent opportunities for

exploring numerous important issues for complex VBIRs aiming to integrate data that is

imperfect in some form, such as being incomplete, uncertain, contradictory, erroneous

or scattered across a wide range of sources. Any of these conditions occur frequently in

biology. It is beyond the scope of this study to explore all such challenges faced by

every biologist, whether she's aware or not; in the Supplemental Material we describe a

few of the insights gleaned from our work on FlyClockbase. These could be

summarized as follows.

Challenges. Any information resource of substantial biological interest will quickly

grow to a complexity at which it will inevitably accumulate a substantial amount of

human errors that are difficult to detect by human users. Many independent repeats of

biological information are typically associated with large amounts of genuine biological

variability. In many current biological databases, it can be difficult to distinguish such

genuine variability from artificial variability that is easily caused by human errors of

various well-known types. Such errors span a broad range of different complexities and

corresponding frequencies. For example, simple typos or label swaps usually occur at

low rates such as 1%, see (PANKO 2016). Simple logic errors occur at substantially

higher rates, especially in spreadsheets (PANKO 1998; PANKO AND AURIGEMMA 2010;

PANKO 2013; PANKO 2016). However, errors of omission are usually the hardest to find

(PANKO 2016). This is especially true, when an omission has become part of a logic

formalism. This is one reason, why it is so important to use good approaches to

represent *Null* (WHITE *et al.* 2013), and why it can be dangerous to confuse different

types of *Null* (WARAPORN AND PORKAEW 2008; HOARE 2009; THALHEIM AND SCHEWE

2011). FlyClockbase has experienced *Null*-confusion already. Entries in a *DZT* column

define an hour of the day. We initially allowed *DZT*=0*h* as a valid time, excluded 24*h*,

and defined 'absence' as '*NotGiven'*, a particular type of *'Nothing'*. However,

understanding '*Nothing*' correctly is difficult. Hence, it is unsurprising when biological

model curators occasionally allow well-known intuitive algebraic properties of addition to

affect their views of 'Null'. As a result, '*0 apples*' is correctly interpreted as '*no apples*';

yet it may fuel the erroneous idea of equating 'time not observed' and 'adding zero to a

list of hours. In this case, some *DZT*=0 values are correct and some are not, but

checking correctness is complicated and expensive. This test becomes trivial, if 24*h* is

included as a valid time, and 0*h* is defined as always invalid. For more details on such

challenges, see also discussion of the BioBinary data type in the Supplemental Material.

   Trans-disciplinary solutions. Several non-biological areas of research and

technology, such as computer science, space flight, and nuclear reactor safety have

developed sophisticated approaches for detecting and correcting potential human errors

(NASA *et al.* 2001-09-30; NASA *et al.* 2006-07; NASA *et al.* 2011; PANKO 2016). While

designers of biological information resources can learn much from the decades of

research that informed the development of human error analysis tools in those areas, it

is less straight forward how these insights could be applied to improve the quality of

biological information available to most biologists. A source of concern is the substantial

complexity of many human error analysis frameworks (REASON AND MYCIELSKA 1982;

REASON 1990; NASA *et al.* 2001-09-30; REASON AND HOBBS 2003; NASA *et al.* 2006-07;

GOLDSTON 2008; NASA *et al.* 2011; GITELMAN 2013; GRIMES *et al.* 2013; MCCALLUM

2013; REASON 2013; BLANKENBERG *et al.* 2014; REASON 2015; PANKO 2016). Most of

these frameworks will handle the complexity of biological data, but require near

prohibitive research and implementation efforts that make integration into grass roots

VBIR projects such as FlyClockbase not efficient if started by biologists. However, that

does not imply that sophisticated approaches cannot contribute to solutions, even if

VBIRs curators do not bring the expertise necessary for implementing a framework. To

see how this might work requires a look at an advanced area in computer science that

is not readily accessible to many: compiler construction.

Error analyses could be amortized across *VBIR*s by compilers. As argued

above, appropriate error analyses for a single *VBIR* are not feasible. However, our

experience with developing FlyClockbase suggests that a substantial number of

essential tasks are recurrent when compiling any *VBIR* of comparable complexity.

Efficiency. Thus, the most efficient solution to improving the quality of *VBIR*s

without exploding costs is to develop an automated compiler that can test for all known

VBIR problems and that supports a programming language that integrates biology

expertise (LOEWE *et al.* 2016b). Programmers frequently say that it is important to use

the right tool for a given programming task. Despite numerous biology-oriented libraries

for non-biological programing languages (e.g. (STAJICH *et al.* 2002)) no general-purpose

programming language exists yet for supporting typical complier-style consistency

analyses for general complex *biological* datasets like *VBIR*s. We will not repeat here the

substantial number reasons why such a language would be helpful and why current

(non-biological) programming languages are insufficient (see Supplemental Material

and additional reasons discussed in LOEWE *et al.* (2016b)).

Examples. Such a compiler could address tasks such as the following. There is a

need for handling missing data, inapplicable data and similar cases by choosing

appropriate representations that distinguish these cases instead of lumping them

together as 'NA' or the value zero (e.g. (CANDAN *et al.* 1997; WHITE *et al.* 2013)). All

biological measurements will always come as imprecise ranges, not as precise values.

Measurement methods for a given observation are usually heterogeneous and need

some description. Observations can be made in may be compared between various

MethodRealms, like *in vitro, in vivo*, or *in silico.* Comparisons between wildtypes and

mutants are frequent. Synonyms are almost ubiquitous. It is easy to continue this list

with many other aspects of biological interest. In addition, there are data processing

basics, such as the ability to read in all tables of FlyClockbase and produce a report of

all inconsistencies and errors that require human attention. The arrival of big data has

brought substantial experience with questions of data hygiene (GOLDSTON 2008; HOWE

et al. 2008; KRISHNAMURTHY *et al.* 2011; GITELMAN 2013; MCCALLUM 2013; SCHUTT AND

O'NEIL 2013; MAHMOOD 2016; ZWEIG 2016). Most of this expertise is also essential for

correctly and efficiently handling data in *VBIR*s. For all features like those above and all

error types detected, a solution only needs to be implemented once for simultaneously

improving the reliability of all *VBIR*s.

          QQv1r1

PopGen predictions on FlyClockbase survival and success

Most new versioned biological information resources (*VBIR*s) such as FlyClockbase

face a dizzying array of potential paths into the future, not unlike newly mutated alleles

in a population. As population geneticists have learned, all this complexity can be boiled

down to two essential outcomes (KIMURA 1962): all alleles are either kept or lost

eventually. To explore other useful aspects of this analogy, we will abstract a few brief

lessons from population genetics that also apply to collections of information.

The stage. If seen in such a general way, a newly arisen DNA-allele could be

compared to a newly published *VBIR* similar to FlyClockbase or a newly developed tool

in bioinformatics (thereby accessing a broader pool of historic precedents). Both alleles

and *VBIRs* contain new information, stored in DNA or on computer hard drives

respectively. Both are part of their ecosystems, which belong to different realms. An

allele exists in carbon-based organisms that compete for natural resources in a

population where the allele may be kept indefinitely. Omitting replication details allows

for simplification. One could think of alleles as replicators based on DNA; similarly,

memes were originally defined as replicators in mindspace (DAWKINS 1976). Thus like

FlyClockbase, each *VBIR,* can be seen as a meme that competes for the 'mindshare' of

humans potentially interested in a given topic. Technically, memes are units of

information that usually spread through communication and compete for the limited

attention of individuals and communities, irrespective of their success of replication

(DAWKINS 1976; LYNCH *et al.* 1989; GLEESON *et al.* 2014; DAWKINS 2016; HE *et al.* 2016). These generic features result in mechanisms similar to those of population genetics, which we use here to derive informal expectations for the future of FlyClockbase and similar *VBIRs* (a formal theory is beyond our scope). We do so hoping to avoid the most likely outcome, the complete loss of FlyClockbase, by aiming to increase the chances that FlyClockbase will be kept in the population of useful *VBIRs*. We next reinterpret concepts like aging, death, growth and reproduction from the perspective of *VBIR*s; Incomplete Fitness Traits (IFT) like these combine with a given environmental context to define fitness in biological evolution (LOEWE 2016). Even without a quantitative meme model, we expect qualitatively similar outcomes when translating IFTs to the realm of *VBIR*s memes. In many cases this will suffice to make decisions that increase the chances of survival for FlyClockbase.

Aging and death. *VBIRs* are aging if they degrade without the time and energy investments necessary for maintaining their semantic reproducibility (LOEWE *et al.* 2016b); they are on their deathbed when nobody wants to use them anymore, and are buried once nobody can remember them. Potential causes of death vary with age and include (i) being locked into remaining an exploratory toy 'too simple' for any real use, (ii) being 'too simple in comparison' from a lack of features that could have helped fight competing *VBIRs* and win over their human users, (iii) having become 'too complicated' for real-world users after years of accumulating inessential complexity (RAYMOND 2003), and (iv) many other causes from internal specifics to external generics (such as political decisions).

Growth and reproduction. *VBIRs* can grow in various respects, some helpful, some harmful, and some hard to assess. We use 'growth' here only in a narrow sense for helpful traits like features required by users. In contrast, we denote as 'aging' the growth of harmful traits like inessential complexity, whereas the reduction of such complexity can be seen as growth (e.g. by simplifying an interface to save user time). Likewise, the loss of useful features can be seen as aging caused by semantic irreproducibility. For example, this could be caused by incompatible changes in required software packages. Here growth always affects the quality of the best implementation of a *VBIR* type, in contrast to reproduction, which could be seen as increasing its mindshare through favorable communication and/or copying the *VBIR* data to new servers (presumably to win new voluntary users). Thus, growth and reproduction in this sense are likely to help a *VBIR* to spread and increase its fitness.

Speciation and merging. The same is not usually true for processes comparable to speciation. The 'forking' of a *VBIRs* repository into two independent lines of development is often perceived as an unwelcome increase in complexity by users without stake (e.g. Python 2 vs Python 3). This implies that the reduction of independent lines of development should be welcomed, but reality is more nuanced. A reduction from merging without loss of features is positive.

Extinction. However, sometimes it is impossible to save all features due to mutual incompatibility or other constraints; this might be comparable to extinction, where good features are irredeemably lost to global mindshare. If occurring to all development lines of a *VBIR* (e.g. due to catastrophic environmental changes such as 'loss of funding'), then the loss is usually tragic, even if the *VBIR* is preserved as a fossil on cutting-edge archives of its time (like floppy disks, CDs, bioinformatics journals, websites, and various open source repositories). As in real life, software fossils are rarely revived, an act that would require extra-ordinary semantic reproducibility as defined elsewhere (LOEWE 2016). Semantic reproducibility is very difficult to achieve, as seen and further discussed in the source code for the statistical analyses in this study and the discussion of the 'DISCOVARCY' documentation style (Supplemental Material). In both cases, it is much more likely to lose fossils to changing environments and random damage than to revive them successfully. Furthermore, chances of successful reactivation drop dramatically in both cases, as bacteria are easier to revive than dinosaurs, and old algorithms for merely sorting numbers are reused more easily than the software systems that put a man on the moon (though we do not wish to imply that either is possible). Extinction can happen to any *VBIR*, no matter how well known. Some planning can usually ensure preservation of a fossil form; ideally a tombstone will inform would-be users where the fossil is archived (see Supplementary Material).

Horizontal gene transfer. As we watched the evolution of FlyClockbase we witnessed a number of remarkable exchanges of information. Our experiences have played out in the conceptual arena defined by Figure 11: we started as initiators,

completed the substantial integration work presented here, and have used

FlyClockbase for research purposes (see Results). At the same time one of us has been

deeply engaged with developing the compiler architecture for the Evolvix modeling

language, aiming to meet particular requirements of biology. This combination of aims

has enabled a substantial flow of critical design information that has benefitted all sides.

Compiler architects benefit from first-hand exposure to challenging practical problems in

logic and data modeling in the domain of their target audience, while biologists are kept

from computationally short-sighted quick-fixes that otherwise could easily wreck a *VBIR*

on the longer-term. Such collaborations are powerful opportunities for uncovering and

clarifying misconceptions on all sides and at all levels; in our experience, they greatly

improve the conceptual quality and robustness of resulting solutions, but come at the

expense of the rate at which some more tangible results can be produced. Combining

these costs with those of human error analyses (see above) when developing reliable

VBIRs increases costs substantially, often prohibitively.

      Practically, we advocate that *VBIRs* do not reinvent the wheel of reliability

independently. This unnecessary reimplementation work is expensive and substantially

increases costs of developing and maintaining a *VBIR*. A thorough analysis of historic

sources of funding for various existing *VBIRs* has exposed a lack of support for this

critical work that integrates, consolidates, and checks the quality of data in *VBIR*s

(EMBER *et al.* 2013). An overview of these essential tasks in the context of FlyClockbase

is given in Figure 11. Here we suggest that much of these costs could disappear if the

initiators, integrators, and researchers working with a *VBIR* would have efficient means

of passing on their formal needs for data representation and analysis to the architects of

an integrative compiler. From their integrative perspective, these architects could then

provide solutions that are compatible and interoperable for many *VBIR*s. Support for

such a versatile open source compiler-building project that serves the *VBIRs* community

well would not nearly be as expensive as independently solving this problem repeatedly.

Experience indicates that well-maintained tools do get used; such a project could hence

substantially contribute towards closing the critical funding gap highlighted by a

thorough analysis elsewhere (EMBER *et al.* 2013). Here is not the space to provide a

reasonable overview of the many aspects of working towards an integrated compiler

architecture. Informed by experiences with FlyClockbase, the tips of several icebergs

are touched in Figure 11. It lists important needs of various contributors, and specifies

several types of lessons learned by *VBIR* contributors and services provided by the

compiler and its construction team envisioned here. This work generally occurs in three

broad stages of integration: combining fragmented insights gleaned from work on

FlyClockbase, investigating broader designs, and integrating solutions into a single

coherent architecture. The high-level analogy of aging and growth in *VBIRs* plays out on

the background summarized by Figure 11. The values of such *IFT*s governing the

evolutionary trajectory of *VBIR* meme evolution are determined by the hundreds of

small implementation decisions necessary for arriving at an overall coherent *VBIR*

organization perceived as elegant, expressive, useful, efficient, and overall simple

enough to be worth a user's while. Such simplicity is pivotal for engaging anonymous

users with a *VBIR* (or any other meme), especially since many are suffering from

information overload, data smog, and the resulting paradox of choice (SHENK 1997;

SCHWARTZ 2004). Physics is not the only discipline where theory should be "as simple

as possible, but not simpler" (EINSTEIN AND CALAPRICE 2011).

Potential predictions. So, what can we learn from population genetics to

improve the long-term usefulness of FlyClockbase? We do not aim to exhaustively list

all general lessons but rather present several possibilities of likely interest in the bigger

picture that we expect based on population genetics theory. This will allow readers to

connect additional dots between more detailed requirements and solutions presented

above as well as in the Supplemental Material. As we review the following potential

paths into the future we interchangeably use the terms 'FlyClockbase', 'new allele',

'bioinformatics tool', and *VBIR* to reduce repetition.

Loss is likely for all new information. Population genetics theory shows that

most newly arisen alleles are lost very quickly by the random sampling that occurs

between generations (KIMURA 1962). All alleles have to navigate this hurdle, regardless

of how beneficial they might otherwise be. Observing bioinformatics research quickly

reveals a similar pattern: on the web very many tools start out (and fizzle out),

professional researchers ensure that at least one peer-reviewed publication exists (but

lack the time to keep websites and tools from breaking), enthusiastic programmers will

keep tools working (but are happy with little documentation), good software engineers

understand the value of organization and documentation (but usually do not work in

biology). All new tools and resources face an intimidating phalanx of these and similar

dilemmas, which made us think hard about all possible avenues for simplifying the

overall system while increasing flexibility. First lesson: FlyClockbase is no exception and

faces the same challenges. It may sound strange to discuss death in the context of a

birth that we believe is to be celebrated. However, ignorance is not a good defense

against child mortality.

Loss is fast and 'child mortality' matters. Alleles that have just arisen by

mutation and new bioinformatics tools that have just been published also share another

important detail: they will probably be lost very soon. Except for extremely harmful

alleles, initial survival for good and bad alleles depends almost entirely on the individual

that carries them. Therefore, FlyClockbase must travel as light as possible if it is to

survive. Like other *VBIR*s, it must be able to fit it into the life of a single publicly known

person who can act as a synchronizing point of contact for coordinating further work

(even if not done by that person). Such public maintainers of *VBIR*s are probably

extremely busy and will have very little time and energy left for high-maintenance

solutions. This excludes the use of many great database technologies that unfortunately

shower their users regularly with recommended updates with various degrees of

compatibility and urgency. Without a highly automated process, such updates would

prohibitively increase the rate of aging for FlyClockbase or the energy required to

maintain it. Accordingly, we have been developing approaches to simplify life with

VBIRs like FlyClockbase, but much more remains to be done. Our various strategies for

simplifying are discussed above and in the Supplementary Material. Laurence Loewe

has agreed to be the first public maintainer of FlyClockbase and will post updates to the

GitHub website given at the beginning of our description of FlyClockbase.

                 QQv1r1

Fossilization is usually deadly. An easy way of avoiding the loss of a project from a broken hard drive is to submit it to a public repository such as Github. This ensures a form of travelling light, as everything stays in place, if a maintainer does nothing (maintenance cost is near zero, and mostly a thought and a password). This establishes a minimalistic baseline, as the mere existence of data (or ancient code, see above) does not differ much from fossils, which are awkward to access, dry and brittle to work with, and for all practical purposes impossible to revive. Lesson three: If nobody continues to work with the code, then chances are that it has already fossilized. Thus, we next review steps that are likely to facilitate future work with FlyClockbase.

Next practical steps for FlyClockbase

In order to raise the chances of survival and success as described above, we are

working towards implementing the following practical steps that improve the

organization of FlyClockbase and move it towards increased stability.

Reorganize files, define versioning policy and simplify folder structure. It is

very frustrating to work with a project where everything can move (and break) at a

moment's notice. Nascent resources never really know what awaits them, and

FlyClockbase has not been different. As a result, our time series data has seen more

profound reorganizations of its storage space than any of us had anticipated. Some of

this additional work was due to the fact that we were simultaneously developing crucial

technological underpinnings, such as TabFS (Figure 11) and the POST system (LOEWE

et al. 2016b). We also did not have a stepwise guide on *VBIR* construction with an

overview roadmap from an expert, which could have further reduced the work. However,

as indicated in Figure 11, the initial phases of a *VBIR* will always be special: each *VBIR*,

by definition, is ill-defined at its inception and negligibly small. As it starts growing, it is

restructured, renamed, and reorganized many times while in its 'embryonic' form. While

guidance helps, some messy aspects of initiating a *VBIR* are probably impossible to

avoid if the freedom is retained to develop any *VBIR* supporting any research. Figure 11

makes a clear distinction between this early more informal stage and the subsequent

iterations managed by integrators. While much of the work of integrators is also done at

          QQv1r1

the initiator stage, *VBIR* publication marks a milestone. It is an excellent opportunity for

internal restructuring and cleanup that should not be dismissed lightly, as reorganizing

will never be as easy again. This is particularly true if the versioning system changes.

Long-term resources require stabilizing versioning systems right from the start to reduce

inessential complexity and confusion. FlyClockbase will build on the *StablizingZone*

(LOEWE *et al.* 2016b) of the POST system. These needs motivated us to delay

publication of FlyClockbase while still under review.

Use a public distributed version control system to be efficient. The use of

Git for version control is rising and services like http://github.com allow open source

projects to be published 'at no cost'. Not having to pay for leaving code in a published

state increases chances of avoiding 'death by negligence' for many *VBIRs*. More

importantly, using Git allows *VBIR* collaborators to essentially cut the huge costs of

manually performing the search and merge operations regularly required for close

research collaborations. We have experienced enough of these complex operations to

appreciate the huge value provided by Git and have decided to use it for FlyClockbase

(currently in a closed repository using http://gitolite.com/gitolite/). However, using Git is

not free of costs. At first these seem reasonable: learn how to use Git and avoid

advanced moves that get 'the rest of us' into serious trouble (including loss of data).

However, in our experience, Git idiosyncrasies and the complexities of version trees

pose such formidable barriers for most biological users, that tool adoption requires a

large activation energy, even when using excellent graphical user interface software

(albeit developed for programmers). We have found an approach for getting biologists

to work with reasonably well with Git. It currently requires determination, detailed

instructions, an expert who performs all operations except the very simplest, and who

happily explains everything again until users follow the instructions (cleaning up the

mess, if they do not). Given the outstanding efficiency of Git, not just for FlyClockbase,

motivated us to explore how to hide our simplified Git workflow behind scripts called

when users 'hit a button'. While our design requires more development and testing, our

internal results so far suggest that it will be more than worth the effort develop this for

FlyClockbase (and reuse for other *VBIR*s). We highlight all this because many biologists

seem unable to imagine how much more efficient the development of a *VBIRs* can be if

Git works as it should. Conversely, many Git users seem unable to imagine why some

biologists prefer to explore every non-Git option first, irrespective of cost. We found that

some of these 'attractive' alternatives can easily turn into complexity traps or create

serious bottlenecks for development. This is in particular true for the prevalent mode of

distributing supporting material for journal articles, which allows reading data from files

without the ability to write back. Such immutability is good for ensuring well-defined

versions, but unless there are files that can actually be updated in place, it will be very

difficult to efficiently work with the data. Complications can abound when manually

merging two sets of text files, each with changes that accumulated in separately

evolving lines of revision descent. Since not all combinations have been tested together,

some could trigger prohibitive integration problems comparable to the severity of

Dobzhansky-Muller Incompatibilities known from evolutionary genetics (COYNE AND ORR

2004). In this analogy, using a system like Git for regularly merging all new changes into

a single main line of revision descent is comparable to keeping all individuals in one

large population. This efficiently prevents the accumulation of the source-code

equivalent of Dobzhansky-Muller Incompatibilities and is desirable for improving one

VBIR that serves a single purpose. Thus, typical archival data storage is not ideal for

FlyClockbase and similar *VBIRs*; we therefore chose Git and aim to mitigate its less

than ideal aspects.

Develop TabFS. For decades, there has been no shortage of databases, file

formats, file systems, and other types of storage - all with unique strengths and

weaknesses. There is no universal agreement on how to best store complex data

transparently. Text-based formats that distribute data across folders in a file system

provide instant and continuous access to content that is easy to read and write for

humans. This flexibility does not depend on any special tools that could break. However,

such transparency benefits are balanced by the need to ensure consistency in the

presence of notoriously inconsistent human users. While binary formats increase speed

and consistency, they complicate *VBIR* development and create costly dependencies on

special tools for reading or writing *any* data. As *VBIR* development requires biological

model curators to easily modify the data model of a *VBIR*, we decided against using

existing excellent binary technologies such as ProtocolBuffers

(https://developers.google.com/protocol-buffers/) and HDF5

(https://www.hdfgroup.org/hdf5/). For these and other reasons detailed in the

Supplemental Material we decided that *VBIR*s and TabFS require human readable text-

based file formats. Appropriately reviewing these is beyond the scope this paper, but

some recurring patterns provide food for thought. For example, the text-based

'eXtensible Mark-up Language' (XML) and the representation independent 'Abstract

Syntax Notation One' (ASN.1) are both widely used, formally defined

(https://www.w3.org/XML/ - https://www.ncbi.nlm.nih.gov/Structure/asn1.html) and

demonstrate the following possibilities for data storing file format standards:

1. it is possible to define broadly applicable standards that maintain a very simple and stable core set of features (encouraging simplicity in TabFS);

2. combining a few built-in data types with arbitrary nesting and repeating of user defined data types can inspire multitudes of specific extensions (suggesting TabFS will need to help users navigate diverse complex *VBIRs* code contributions to reduce complexity and unnecessary reinvention);

3. fierce competitors are capable of adopting shared standards resulting in win-win-win outcomes for both competitors and the general public; see the use of ASN.1 in telecommunication (and also by NCBI, see link above; the research benefits of a system that significantly simplifies the sharing of complex biological data are undisputed; developing FlyClockbase across the Win-Mac divide together with the Project Organization Stabilizing Tool (POST) system (LOEWE *et al.* 2016b) by using the process in Figure 11 led to major TabFS design ideas that could simplify current challenges enough to motivate grass-roots adoption; details beyond scope here);

4. the rise of simple text-based XML could not quench the need for even simpler text-based file formats that are popular for simplicity where it matters (suggesting TabFS needs to provide comparable ease of use);

5. simplistic file formats are insufficient for representing many types of biological complexity and will therefore never be adopted universally (suggesting TabFS must handle arbitrarily complex data in elegant ways).

                    QQv1r1

Comma Separated Value files (CSVs) and the equivalent tab-delimited table files of

TabFS are still particularly convenient file formats of choice due to their simplicity and

extraordinary broad interoperability. Research collaborations frequently share data

across very different systems. Thus, a file format that can easily be read and written

everywhere remains competitive against faster rivals that do not work everywhere.

Unfortunately, CSVs store only values, but cannot store types and cannot directly

describe arbitrary data structures. Therefore, all additional information requires

extensions that are rarely standardized. Recent text-based standards like JSON (see

http://json.org) or YAML (see http://yaml.org) cover many use-cases, but have not

replaced CSVs in many contexts.

Tables in their simplest form. The two-dimensional layout of CSVs is particularly

well suited for time series, arrays, and other frequent forms of biological data. CSVs are

easy to read and write with spreadsheet tools that are widely used among biologists.

Many experimental biologists would not hesitate to use such tools for modifying sets of

CSVs but would avoid equivalent tasks in SQL databases. This fundamental usability

advantage of text-based tables motivated our data storage choices for FlyClockbase.

The downside to this flexibility is the lack of formally defined computational expressivity

that is powerful enough to represent all the needs of *VBIR*s. Our numerous searches

have brought many interesting file formats to our attention, but none approaches the

simplicity and usability of CSVs while also providing a stable international standard with

the features necessary for efficient *VBIR*s development. This gap surprised us.

TabFS specification. We plan to fill this important *VBIR* tool gap by developing a

definition and implementation of TabFS. The TabFS specification aims to define

precisely a completely open and customizable, easily accessible and usable, extremely

simple and stable, maximally versatile and expressive storage system for long-term use

in *VBIR*s such as FlyClockbase. Here *long-term* indicates the requirement to be long-

term backwards compatible as defined by the '*TrustedTested'* (TT) level in the POST

system defined elsewhere (LOEWE *et al.* 2016b). To achieve these goals, some aspects

of computational speed will receive a lower priority in TabFS, as speed of *VBIR*

development is more important for TabFS than speed of execution. Practically, TabFS

builds on the stability of standard file systems and uses tables and other fragments in

files and folders to implement well-defined conventions for storing the necessary

nuances required for *VBIR*s development. A major design aim is to keep the raw

convenience and efficiency of tab-delimited table text files (hence the name TabFS).

Such files are easily edited by spreadsheet tools familiar to many biologists and readily

imported and exported by many other systems. TabFS is developed in the context of

VBIR development as described in Figure 11 and uses the flipped programming

language design approach presented elsewhere (LOEWE *et al.* 2016b). Many *VBIRs*

share similar problems, some of which are typical for biology. Solving them once in a

reusable way can greatly contribute to the reproducibility and the sustainability of

domain specific repositories for digital data (EMBER *et al.* 2013).

Define a type system for TabFS. Work towards defining each essential data

type for TabFS in general and *VBIRs* in particular will need to continue in parallel to

developing TabFS itself. Substantial overlap in development is essential for ensuring

that TabFS provides all important capabilities for making high-level *VBIRs* development

efficient while minimizing overall system complexity. Establishing a stable core of TabFS

first will greatly shrink the complexity of developing a stable and consistent type system

for recurrent tasks in both TabFS and general *VBIRs* development. The same

mechanisms will later be used by developers of any specific *VBIR* to define a type

system for their particular area that can then be enforced with the same mechanisms

that protect the integrity of TabFS or general *VBIR* types. Since type systems are

conceptually equivalent to ontologies at a high level (ARP *et al.* 2015), such work can be

structured in work-stages that are familiar to biologists since the start of taxonomy:

observe, describe, define. Practically:

1. *Observing* which types of folders, files, or fragments are useful for developing

    and maintaining a *VBIR* is only possible in the context of a real *VBIR* with real

    research problems, such as FlyClockbase. Pure thought or toy projects cannot

    reveal enough real-world nuisances and nuances for developing a high-quality

    VBIR type-system. The next step for the resulting list of observed entities is:

2. *Describing* at epic length in human readable text every detail about, why and

    how exactly each folder, file, or fragment is stored and used by expert biological

    model curators provides a solid foundation for the final step of explaining all this

    to computers:

3. *Defining* each type formally, which results in a checklist for determining the

    integrity of this *VBIR* type and for detecting all known errors.

Initially, such checklists are best developed and refined by expert users willing to accept

a temporary slowdown caused by the need to document and check every step of their

work, including the simplest ones (tedious for humans, essential preparation for

computers). Once sufficiently detailed, these checks can be automated, enabling

experts to focus their energy on complications that computers cannot currently handle

correctly. Such a style of collaboration with *VBIR* compilers would allow all parties to

focus on what they do best. Machines mindlessly repeat mind-numbing instruction

sequences. In contrast, experts focus on activities where humans excel: apply expertise

and common sense to check the integrity of computational results, think creatively about

new tasks for the *VBIR* compiler, and expand the *VBIR* by exploring interesting

hypotheses. Once this *VBIR* has matured enough to answer the interesting questions in

its field, start a new *VBIR*.

For the first several *VBIRs* most contributions to such a compiler will probably

focus on the basics of defining and referencing various types of memory devices, such

as folders, files, file names, tables and fragments of these. A well-defined type system

will greatly simplify the implementation of the consistency checks that are essential for

maintaining the integrity of FlyClockbase.

Automate TabFS checks to help expand the biology of FlyClockbase.

Developing FlyClockbase, TabFS and a *VBIR* compiler for ensuring the long-term

stability of *VBIR*s can be greatly facilitated by a code library implementing a storage

interface for TabFS instances. Detecting formal errors, enforcing rules and limitations,

ensuring the full execution of all aspects of a TabFS or *VBIR* task, and performing other

jobs can then be delegated to such a storage library and will no longer consume

precious development or research time. These new liberties can then be invested in

expanding the reach of FlyClockbase by adding the latest biological studies, new and

old mutants, and many other aspects. Additions require defining new columns or new

values for the controlled lists of existing columns. Carefully reviewing anticipated usage

reduces clutter in the name-spaces of FlyClockbase. This is pivotal, since column

names become immutable once pronounced '*TrustedTested'* as defined (LOEWE *et al.*

2016b). The ability of FlyClockbase to disentangle the long-term need for stability and

the short-term freedom required for *VBIR* innovation will critically depend on the early

introduction of a well-thought out stabilizing version number system for FlyClockbase,

lest it be killed by inessential complexity on the long run.

Conclusion

This study contributes important foundations to our overall goal of improving the reproducibility, reliability, and relevance of biological data analyses, starting with observations of the *D. melanogaster* circadian clock. To this end, we aim to automate as many repetitive tasks as possible by providing computational tools that can be efficiently used by experimental biologists. Ideally, this will inspire increased adoption of computational tools and empower biologists to expand their thinking capabilities to investigate new questions. This will be required to meet current grand challenges from personalizing medicine to predicting mechanistic fitness landscapes in evolutionary systems biology (Loewe 2016). Such types of problems often require the analysis of innumerable smaller computational models, which is impossible without highly automated information processing to cut through the associated cognitive complexity.

FlyClockbase as a *VBIR*. The resource we compiled might be able to serve as an example for a versioned biological information resource that is organized in a radically simple way by being completely accessible as tables of text. It also exemplifies what a 'small model' in a grand challenge context might look if comparable in size to our clock model (see Figure 1) with similar amounts of time series or other experimental data. We expect such data to be as scattered as it was for FlyClockbase. Experience with time series in FlyClockbase suggests that many other datasets are probably also likely to contain a mix of broad general trends and numerous statements that remain incomplete, imprecise and contradictory. To successfully handle this avalanche of

                 QQv1r1

challenges in biology, we have been analyzing observations and models of the fly

circadian clock. Simultaneously we have been collecting instances, where automation

by a compiler could greatly increase the efficiency of integrating biological knowledge-

fragments and maintaining the integrity of a VBIR in face of common uncertainties in

biological data.

Designing a compiler for biological data. The design of such a compiler is

greatly improved in our experience, when developed simultaneously and in close

collaboration with biological model curators who regularly expose compiler designers to

the many imperfections of biological data. The seemingly perfect abstractions of

compiler type systems need to meet the messy observations made in biology, and

conversely, biological observations need to become more organized by learning from

the abstraction techniques developed in computer science. Such trans-disciplinary

communication is possible in our experience (see Figure 11 for an overview of the

process). Consequently, our work in this study drills deep in distant areas from different

disciplines, both basic and applied. The volume of relevant material forced us

repeatedly to refer to Supplemental Material, the Evolvix BEST Names study (LOEWE *et*

al. 2016b), or simply limit scope (usually indicated). A brief overview of the relevant

research areas might illustrate these challenges for compiler construction.

Trans-disciplinary aspects. The seemingly disparate areas of enquiry in this

study are deeply connected by our desire to improve the reproducibility and reliability of

models in computational molecular systems biology. We study:

(i)   *the molecular genetics* of gene regulatory networks in *Drosophila* circadian

       clocks (reviewed in Figure 1),

(ii)   *the statistics of robust differences* in variance among observed time series traits

       (Figure 5),

(iii) *the applied mathematics of simulating* time series from Continuous Time Markov

       Chain models (Figure 4 lists models, leaving simulation for later),

(iv)   *the behavior of modelers*, namely how they prefer to parameterize their models

       (Figure 4),

(v)   *the human-computer interactions* that help to reduce data smog and information

       overload by improving visualization and organization in plots, in models, in and

       data structures (Figure 1,2,6,7,9),

(vi)   *the statistics of detecting human errors* in spreadsheets, data analysis, logic, and

       source code (Figure 3, Table 5, Discussion, Supplementary Material),

(vii) *the data science of reproducibility* for improving reliability, semantic, statistic, and

       other reproducibility of publishable research results from the early investigative

       stages (see Supplemental Material, Table P1 and 'DISCOVARCY' Documentation

       Style), and

(viii) *the computer science of compilers and programming languages* as needed for

       supporting the development of other biological information resources like

       FlyClockbase. This requires addressing a broad range of topics, including

       mathematical logic, type theory, arithmetic, syntax, semantics, memory

       organization, naming, and others. Figure 11 provides an overview of the types of

                     QQv1r1

interactions we have observed between biological model curators and a compiler

architect while developing FlyClockbase.

Thus, we touched the tips of many icebergs and often needed to limit our scope. Much

of this tension was caused by our desire to build a compiler that understands the

imprecisions and complexities of biology and supports the efficient construction of high-

quality *VBIR*s. We have pursued this goal by constructing such a *VBIR* and performing

manually all tasks that we would like to delegate; this gave us the opportunity to reflect

on the nature of the tasks and the quality of the outcome. This reduces the speed of

both: compiler construction and *VBIR* construction, but simultaneously greatly increases

quality. As argued by our analogy to aspects of population genetics theory, such

increases in quality can be pivotal for the survival of a *VBIR* like FlyClockbase, which

can easily be killed by small increases of inessential complexity. In this study we

provided a broad overview of this tandem work. We have removed from this paper all

aspects that can also stand on their own. For example, readers of this journal might be

less interested in a formally complete description of the data structures that comprise

Evolvix and the nuances of data models that contribute towards long-term stability. We

endeavored to keep in the main text only those computational aspects that are most

important for navigating the broader concepts used in FlyClockbase or that convey a

general overview of our approach to reducing the cost of maintaining digital repositories

with the help of a compiler designed to so. There is no reason why such a compiler

could not be used by individual researchers collecting their own data, some of which

they might want to share later. Therefore, our work presented here could also be seen

from the following points of view.

View on gene expression variability. The most direct purpose of our study is to

use FlyClockbase to generate and analyze hypotheses about circadian clocks in *D.*

melanogaster. We analyzed patterns of circadian variability across diverse independent

studies of fruit flies, accumulating the largest number of time series for this purpose to

date (to our knowledge). We have used the statistical power of FlyClockbase to detect

consistent differences in the variance of peak times for the important clock proteins PER

and TIM. This led us to hypothesize that these differences have mechanistic causes that

are worth investigating with the methods of computational molecular systems biology

(out of scope here). Our detailed analysis of variances in the peaks of PER and TIM

and the potential causes for outliers (see above) suggests the removal of outliers by

default using the method of Carling (2000) to focus more efficiently on estimating what

typical clocks usually do (without suppressing natural variability in time series). Similarly,

FlyClockbase can be used to compare the accuracy of different observation methods

(Figure 8) and many other *Attributes*. An important contribution of FlyClockbase towards

simulations of fly clock models of gene expression variability is its rich set of over 400

wildtype time series that can be used - in principle - to improve estimates for circadian

clock parameters. Such estimates might change the rather sobering observation that

most clock modelers do not use most experimental observations when deciding on the

parameter values for their simulations (see Figure 4). A study using state-of-the-art

inference methods for obtaining the best possible clock model has been moved beyond

the scope of this paper but could start immediately.

                    QQv1r1

View on simplifying *VBIR*s development. The broader purpose of our study is to develop, describe, and use FlyClockbase as a real-world testing ground for designing an extraordinarily reliable yet simple system for long-term backwards-compatible data integration. We also explored how to annotate, name, reference, identify, store, query, retrieve, and analyze the imperfect and complex biological data and its translation into well-defined computational concepts. Developing these capabilities is essential for the long-term mission of programming languages like Evolvix that aim to provide built-in support for biological research. This goal requires unusual amounts of direct user feedback from experimental biologists to the language designers, as described elsewhere (LOEWE *et al.* 2016b). Since computers and their computations are ultimately abstract, software engineers have come to value the input of so called 'domain experts' without whom it would be impossible to develop efficient and reliable non-trivial systems. Such feedback is easier to provide in engineering and other technical scenarios where domain experts and software engineers tend to speak a similar language. However, such a shared language does not usually exist in biology where the 'domain experts' are experimental biologists who often are not used to expressing their expertise in a form easily understood by software engineers. It is an important goal of Evolvix to fill that gap and enable the best experimental biologists to express their expertise in a form that is readily translatable into computable models. Simplifying the construction of VBIRs is an essential component of this larger goal and critically important for evolutionary systems biology (LOEWE 2016; LOEWE *et al.* 2016b).

View on Evolutionary System Biology. The ultimate long-term purpose of FlyClockbase is to substantially contribute towards implementing the vision of mechanistic simulations in evolutionary systems biology as detailed elsewhere (LOEWE 2009; LOEWE 2012; LOEWE 2016). Evolutionary systems biology aims to quantify fitness landscapes by mapping genotypes (via realistic fitness causality networks) to phenotypes and ultimately fitness. Since circadian clocks have a large impact on fitness, their behavior is of direct evolutionary importance (BEAVER *et al.* 2002; BEAVER *et al.* 2003; DODD *et al.* 2005; LOEWE AND HILLSTON 2008; AKMAN *et al.* 2010; BEAVER *et al.* 2010). Constructing a high-quality model of a circadian clock in *D. melanogaster* could thus provide the opportunity to explore many mutant options *in silico* (LOEWE AND HILLSTON 2008) and thus bring us closer to the goal of quantifying fitness landscapes of interest (LOEWE 2009; LOEWE 2012; LOEWE 2016). To enable this vision, myriads of models on the scale of FlyClockbase will need to be constructed, connected and analyzed both individually and in various combinations. Most of today's tools do not manage imprecision with the high degree of precision that is needed for integrating models at such a scale. To address these problems, we need the *VBIR*s automation discussed above and other new approaches to biological model curation.

Biological model curation. The substantial needs for biological model curation illustrated in this study highlight a challenge faced by biology as a discipline. Researchers have accumulated very large amounts of biological data that is currently scattered across the scientific literature in forms that are difficult to access efficiently (or become completely inaccessible as lab notebooks are being thrown out or primary data

is lost from hard drives). In FlyClockbase we integrated scattered data from across the

literature. The substantial amount of work involved forced us to acknowledge, that it is

not possible to engage in the integration of biological information at this scale without a

substantial investment of time. Even if *VBIR*s construction is eventually simplified to the

highest possible degree by the most user-friendly compiler and *VBIR*s construction

environment imaginable, the need for model curation in biology will not become trivial.

On the contrast, such a compiler could motivate a new generation of biologists to

actually revisit and integrate data that has long been ignored, because using it without

compiler support would have been too tedious. This possibility will likely boost interest in

a currently unusual avenue to biological research that is not well represented in the

biological job market of today.

   Status quo. For a long time, most biology undergraduates have been aiming to

work at the bench in a wet-lab. Biologists overly focused on wet-lab work might

undervalue the importance of biological model curation by underestimating the

intellectual efforts it requires. However, what use is experimental data if it remains

inaccessible? While biological model curation does not generate new data *per se*, it

makes existing experimental observations accessible in integrated forms. The resulting

information repositories, such as GeneBank, are prime sources of data used by

computational biologists. The rising importance of computational modeling and

bioinformatics in biology is now recognized well enough so that students in these areas

can readily self-identify and point to labs, role models and career paths. Such

computational professions require substantial training in formal methods, quantitative

approaches and computational tools – usually not easily understood by experimental

biologists who dedicate their career to investigating a particular system in great detail.

Conversely, many computational, mathematical, and other programming biologists

struggle to develop enough dedication for a career committed to studying a single

biological system. The time they take to develop their computational expertise takes

away from the time they have to develop their biological intuitions to the level required

for high-quality biological model curation.

A growing avenue to biological research. Work on biological model curation

which was integral to obtaining the results we presented alerted us to a rising need for

the integration of biological data. As shown by the new biological insights presented in

this study, biological model curation is as essential to biological research as

bioinformatics algorithm development, original lab observations, and field data

collecting. It does not stand behind lab experiments or computational work in its

potential for contributing new biological insights. The low entry bar to model curation

should not be mistaken for a lacking ability to advance the cutting edge of science. Each

major avenue of biological research has trivial activities that do not speak to its potential

for biological innovation. Pipetting samples into tubes does not reflect the complexities

of experimental biology. Defining the initial values for a few variables in a program does

not reflect the potential for innovations from computational biology. Similarly, the simple

activity of comparing a few numbers from a few studies in a spreadsheet does not

reflect the importance of biological model curation for progress towards addressing

grand scientific challenges. In our experience, in depth biological model curation for

non-trivial questions requires a substantial amount of attention that will not realistically

leave much room for additional work on the side, whether in wet-lab or in computation.

The FlyClockbase work present here demanded the undivided attention of several researchers and integrators. Model curation work is easy to scale up or down, but significant new findings still require dedicated resources – as everywhere in research.

What it takes to do biological model curation. While biological model curators are still rare, their work has more history that commonly known (see Introduction on biocurators). Biological model curators must have sufficient interests in the wet-lab work necessary for generating the observations they curate to know about typical pitfalls, but they typically do not work at the bench. They must be sufficiently aware of the strengths and weaknesses of relevant modeling approaches and extract the most relevant information from the scientific literature, but they do not need to be expert programmers. Most importantly, they need a passion for 'their' system to the point where they want to know everything about it, irrespective of the method used to observe it. This will enable them to accumulate enough expertise for learning about the strengths and weaknesses of different methods of observation and for developing an intuition about the quality of a given data set. Such expertise is essential for helping to improve the overall reproducibility of statistical processing pipelines by improving quality of relevant input data, as recently called for (LEEK AND PENG 2015).

On the shoulders of giants. We aimed to stand on the shoulders of giants in fly clock research. This would have been impossible without the biological contributions from the high-quality model curation work that resulted in FlyClockbase. To enable more biologists to stand on the shoulders of their giants we have been working towards capturing our experiences with FlyClockbase in the definitions of *VBIR*s. We expect that constructing a corresponding *VBIR*s complier will greatly accelerate the integration of

the biological expertise required to meet the grand challenges of our time. One of these

is to understand the long causality chain that starts with the daily rhythms of core clocks

and ends with detailed mechanisms for the changes in health and fitness caused by the

daily rhythms of the thousands of genes under circadian control.

Acknowledgements

We thank Jerry Yin, John Hogenesch and Chris Bradfield for stimulating discussions of circadian clocks, methods of how to observe them in mice and flies, and how to interpret aspects of these observations and of our findings. We thank John Yin, Millard Susman and Anthony D. Pietsch for many discussions on more general aspects of this work and numerous comments on earlier versions of this manuscript. We thank Cecilia Moog, Jocelyn Meyer, Sarah L. Northey, Ginger Ann Contreras, Coco Contreras. Kimberli Ward, and Martha Loewe for helping to refine some sections of the text, as well as reviewing BioBinary definitions and suggesting substantial relevant improvements.

We thank Alyssa Hotz, Claire Nusbaum, Devanie Tucker, Noah Waters, and Keegan Reilly for stimulating discussions of various aspects of circadian clocks and their importance and all other members of the Loewe Lab who have in various ways contributed to innumerable discussions of requirements for a programming language that can handle the uncertainties and complexities of biology.

We are very grateful to Ben Liblit, David Page, Ines Dutra, David Anderson, and Michael Ferris for numerous comments over the years on a wide range of technical, formal, and mathematical details that have shaped in several ways the design of FlyClockbase and/or aspects of analyses performed in this study.

We are particularly grateful to Seth A. Keel for setting up the initial Git repository of FlyClockbase and helping with numerous Git-troubles. Most importantly, he contributed critically to the stability of FlyClockbase by highlighting the flexibility-vs-performance trade-off in databases-vs-filesystems: he made us realize how databases

usually compromise flexibility, long-term stability, and low maintenance costs in order to

gain performance in searching for answers to prepared types of questions. In contrast,

for research we need maximal flexibility, reliability, accessibility, recoverability – all at

minimal complexity and costs. Seth showed that these traits are more important than

faster answers to more restricted, recurrent questions, since question types in research

are unpredictable.

Funding. We thank the Welton Family Foundation for an early grant to K.S. that

helped to start this work, the National Science Foundation for NSF CAREER Award

1149123 to L.L. that supported most of this work, and the Wisconsin Institute for

Discovery at the University of Wisconsin-Madison for general support. Any opinions,

findings, and conclusions or recommendations expressed in this material reflect those

of the corresponding author and do not necessarily reflect the views of NSF or other

funders or contributors to this work.

Author contributions

Conception and design. K.S. created, designed, developed, tested, and repeatedly

improved most aspects of FlyClockbase, *PeakValleyTables*, and their initial statistical

pilot analyses that indicated which results were interesting for our first analysis,

eventually leading to the analyses presented here. K.S. was supported by J. Dr. and

E.N. in many practical aspects, including substantial re-organizations of FlyClockbase;

she was also supported by L.L. in theoretical, modeling, and various data layout design

aspects, as well as in questions of computational modeling, and managing connections

to Evolvix. After L.L. set the initial stage by formulating the overarching goal to work

towards creating the best possible circadian clock model for *D. melanogaster,* K.S. took

the idea and in several iterations developed the biological model in Fig. 1 which drove

many biological aspects of FlyClockbase and of this paper; L.L. conceived the approach

for investigating error rates and potential blind spots of poor reliability in FlyClockbase.

L.L. conceived, designed, and has been continually developing Evolvix, TabFS,

POST, and various other concepts in Evolvix that also underpin FlyClockbase and

governed its technology choices. In his design work he was substantially supported by

K.S. who conducted countless usability reviews and provided numerous suggestions for

improving, clarifying and simplifying many diverse aspects of Evolvix, TabFS, logic

systems and operator precedence problems from a user's perspective. Her feedback

included conceptual suggestions, architectural aspects of item-type-context interplays

with BioBinaries to indicate TS analysis integrity, numerous ideas for syntax and name

choices that improve the consistency and visual elegance of many important

punctuation symbol choices for Evolvix, TabFS, and FlyClockbase. L.L. conceived and

designed the overall strategy of mixing disciplines by combining in-depth molecular

system biology model curation work integrating all expertise and observations about a

system, with research relevant for compiler construction. The latter includes in-depth

analyses of the limits of some formal logic systems and the practical aspects of

developing an architecture that better supports biology by developing a readable syntax

and naming-support for facilitating VBIR curatoration.

Experiments and acquisition of data. K.S. collected and organized all data

(as described above, supported by J.Dr. and E.N.). J.Da wrote a script that greatly

facilitated many early explorations by K.S. To measure human error rates, L.L.

developed a strategy in close collaboration with K.S. and J.Dr., who both executed this

strategy (with assistance from E.N.) by thoroughly searching for human errors that –

despite all caution – had persisted into *Mod5* (resulting in *Mod6* after correction; see

text).

Analysis and interpretation of data. Substantial statistical consulting and

numerous very helpful discussions came from B.H. and J.Da. The initial exploratory

statistical analysis was done by K.S. (with consulting support from B.H.). B.H. provided

an important independent statistical analysis that was instrumental in prioritizing

questions while designing the final statistical analysis. L.L. independently developed

and implemented the final statistical analysis performed by over 12K lines of R source

code available as Supplemental Material with the input files used. L.L. collated the

output with comments into a single PDF denoted as Supplemental Statistical Analysis in

the text. B.H. also contributed towards various practical aspects of exploring the data

and helped defining the approach that turned into the "linearization" of *CZT* discussed in

this study; B.H.'s regular practical and statistical input has been extremely helpful for

K.S. and L.L. by providing innumerable valuable pointers that helped navigate many

complex decisions about how to best analyze the data presented. B.H. thereby

substantially shaped the work and increased the overall statistical realism and rigor of

this study. K.S. designed and implemented the workflow for increasingly refining the

PeakValleyTables.

Drafting of manuscript, revising content. K.S. wrote the first draft of the whole

paper, created Figures 1-6+9, substantially refined many sections (Introduction, all

sections directly discussing biological parts, Materials and Methods, Results,

Discussion) and provided detailed feedback on most parts written or revised by L.L.. In

addition, K.S. helped adjust the overall paper structure and figure representations based

on discussions with L.L.. K.S. created several items of Supplemental Material,

summarizing various results in tables, and worked with L.L. to streamline the overall

presentation of results in order to reduce complexity for readers.

L.L. wrote the sections describing the data model of FlyClockbase (including its

Supplemental Materials and substantial revisions to the FlyClockbase overview Fig. 2).

L.L. greatly expanded K.S.'s texts on statistics and modeling, as used in this study

(robust statistics, see R script, Supplemental Statistical Materials; parameter estimation

for solving inverse problems). LL substantially edited the presentation of the Results and

Discussion, and added Figures 7,8,10,11, and Tables 2,8. L.L. motivated and greatly

improved the error analysis within the limitations of this study. L.L. wrote all sections on

biological model curators, and computational aspects that build links to VBIRs, TabFS,

Evolvix, reproducibility, logic, complier construction, the 'DISCOVARCY' documentation

style, and broader links to biological modeling and evolutionary systems biology. L.L.

provided biological consulting thought-out this project.

Watching the clock for 25 years in FlyClockbase:
Variability in circadian clocks of *Drosophila melanogaster* as uncovered by biological model curation

Katherine S. **Scheuer**[*,†], Bret **Hanlon**[‡], Jerdon W. **Dresel**[*,†],
Erik D. **Nolan**[*,†], John C. **Davis**[‡], Laurence **Loewe**[*,†]

[*]Systems Biology Theme, Wisconsin Institute for Discovery,[†]Laboratory of Genetics,
[‡]Department of Statistics, University of Wisconsin-Madison

Appendix: Supplemental Material

Table of Contents

1.   FlyClockbase design processes in more detail

An overview of biological and data integrity aspects of FlyClockbase, as well as a description of relevant data types and their units is given in the Models section of the main text. Here we present practical 'behind the scenes' aspects of information management work. Although information management work is not "biological", it is of substantial importance for the usability, reproducibility, and maintainability of FlyClockbase. Our paper stands by itself without these details. However, given the importance of reproducibility in computational biology; the enormous struggle to design systems that facilitate a desirable level of reproducibility over longer periods of time; as well as our interest in this topic, we decided to take the opportunity to reflect on some of the choices we made, and our subsequent experiences.

FlyClockbase development is intertwined with Evolvix. Many aspects of FlyClockbase are linked with our work on developing reliable general-purpose programming capabilities for Evolvix. Evolvix is a modeling language that aims to facilitate accurate modeling for investigating biological systems and to provide long-term backwards compatibility for computational biologists to build on each other's work in a more direct way (reducing time on re-implementing models). A prototype of Evolvix is available for download at (http://evolvix.org). The prototype download efficiently simulates pure mass-action models described in a declarative programming paradigm, and efficiently collects the most important time series data points during a specified simulation run (thwarting the origin of potentially large simulation results datasets before they become difficult to handle). This Evolvix prototype has been used for simulating models in research (EHLERT AND LOEWE 2014). Our practical experiences with this prototype in the context of a more complex modeling study (EHLERT AND LOEWE 2014) demonstrated the need for adding general-purpose programming capabilities to further simplify the construction of many modeling scenarios (debugging across several different languages as done currently is not an efficient use of research time). To overcome these challenges, we are developing an extension for Evolvix that

implements a general-purpose programming language, designed to simplify general programming as much as possible, in order to stay true to the original vision of Evolvix, i.e. to make accurate modeling easier. This effort turned out to be highly unusual and required us to develop a new approach to programming language design, which we described elsewhere as Flipped Programing Language Design approach (LOEWE *et al.* 2016b). Briefly, this approach ensures that unexperienced users of Evolvix can provide useful input in how the language is designed in order to ensure overall simplicity is maximized too, while experts work to maximize expressivity.

FlyClockbase and its challenges to programming languages. Many real-world modeling problems in biology share characteristics that simultaneously characterize corresponding general-purpose programming challenges. The biggest computational obstacles to efficient computation in biology might be in its unique mix of diversity, complexity, uncertainty and vastness. In combination, these efficiently frustrate many computational abstractions that work well for less intense combinations of these complicating traits. A particular challenge is often the unpredictability and subtleness of such problems and exceptions; thus, there is no substitute for covering ground "on the ground" by walking detail by detail through specific biological examples while paying particular attention to cognitive dissonances that indicate a poor fit between the biological realm and the realm of current programming approaches. When certain types of programming bugs are eventually observed repeatedly, a programming ecologist might take solace in the observation that continued sampling in complex ecosystems will eventually yield a finite count for the number of types in a species richness problem (DOPFER *et al.* 2008; ZHANG AND STERN 2009; MAGNUSSEN *et al.* 2010), even if that number is very large as in the case of bugs (GROVE AND STORK 2000; HAMILTON *et al.* 2010). FlyClockbase has provided us with the outstanding opportunity to observe at close range many instances that frustrate the efficient use of biological observations for computational purposes. Taken alone these issues are rarely large or complicated, but combined can be challenging ("glass of water" vs "tsunami"). This is a characteristic of scattered big data that often become relevant only in much larger datasets (to accumulate enough exceptions for triggering complexity problems). The

data collected for FlyClockbase presented numerous challenges regarding the diversity and uncertainty of time series data. These challenges of FlyClockbase have inspired various important requirements for Evolvix development, aiming to simplify such work in the future (see Discussion of main text for some examples, but reporting most results would be beyond the scope of this study).

Evolvix development follows a unique approach. The unique programming challenges of biology have inspired the rather unconventional Flipped Programming Language Design approach for developing Evolvix (LOEWE *et al.* 2016b). This design process for Evolvix is heavily front-loaded and emphasizes the coherent, user-friendly integration of functionally mature sets of features that are chosen with a view to long-term stability. The aim of these priorities is to reduce idiosyncrasies and confusion for users, as well as inessential complexity and costly busy-work for developers, especially on the long term. The slightly higher short-term development efforts required for facilitating the necessary review processes are negligible in comparison to the benefit of developing Evolvix in a way that generates a long-term backwards-compatible language.

Long-term goals of Evolvix. Evolvix aims to provide general-purpose programming capabilities designed by biologists for biologists to help with facing the complexities of biology. Given that inessential complexity is at the heart of many inefficiencies and problems in computational biology, it is mandatory for Evolvix to find the most efficient way of cutting as much programming complexity as possible. Efficiency will increase the ease of handling additional biological complexity. Architects of languages or compiler designs can only consciously address problems that they have noticed or that have been brought to their attention. Once such problems are openly visible, it is much easier to propose solutions. Once a solution has been proposed, it is much easier to critique it. The Flipped Programming Language Design approach used by Evolvix facilitates a communication line between a language architect that can still change the design of a language, and potential future users, who would otherwise just keep to themselves the many reasons that motivate them to classify a programing

language as too complicated for them. Evolvix aims to break that cycle, where
(otherwise clever) users prefer to not acquire computational skills that are well within
their reach, simply because they are confused by some language-specific idiosyncrasy.
To get passed this barrier to semantic reproducibility requires work towards debugging
what has been termed the Code2Brain Interface (LOEWE et al. 2016b).

How FlyClockbase helped Evolvix. In programming language design, as
elsewhere, problems that remain hidden rarely go away and accidentally solving them is
unlikely. We have used this perspective to turn these problems into a treasure trove of
inspiration for designing innovative Evolvix programming language features. Working
through the fly clock time series data that we integrated into FlyClockbase has
presented us with a rich set of subtleties, semi-regularities, exceptions, uncertainties,
contradictions, and other imperfections of real-world biological data. The decision to
face these difficulties has enabled the development of designs for general-purpose
programming language features that fit biological research problems much more
naturally than solutions developed in 'non-biological' general-purpose programming
languages (e.g. BEST Names for handling synonyms as described elsewhere (LOEWE et
al. 2016b)). Many of the general results from our substantial engagement in this trans-
disciplinary process remain beyond the scope of this study. However, the study that
introduces FlyClockbase is the most natural place for presenting some important
aspects of the process that helped us develop key concepts for representing the
confusing complexity of biological data that we encountered in our work with
FlyClockbase. An overview is described in Figure 11 and respective comments in the
main text. We have included illustrative examples of insight gained from this processes
below and in the Discussion.

How Evolvix helped FlyClockbase. Conversely, the intertwined development
process has motivated the adoption by FlyClockbase of numerous approaches originally
developed to meet Evolvix requirements. An unduly short list of examples include the
BEST Names concept, (LOEWE AND KEEL 2014; LOEWE et al. 2016a), the Project
Organization Stabilizing Tool (POST) system (LOEWE et al. 2016b), the insights we used

to assign a special role to standard filesystems for storing data (which eventually stimulated the development of TabFS as described in the main text Discussion as the storage structure underpinning FlyClockbase), BioBinaries, and the stabilizing versioning system briefly described next.

Stabilizing Versioning. To facilitate development, version variants of FlyClockbase are numbered according to the stabilizing versioning system developed for Evolvix and described as part of the "*StabilizingZone*" of the POST system (see online material of LOEWE *et al.* (2016b)). The term "version variants" is meant to highlights the diversity of types of variants that might change an instance of FlyClockbase in the stabilizing versioning system that it uses. For example, let us consider the version variants of this document when initially submitted to reviewers, and at the moment of initial publication as peer-reviewed study.

QQv1. The version variant associated with this submission is QQv1. This is the Brief Evolvix way of pointing to the meaning of the *StabilityCode* for "*QualityQuest*" and indicate that a substantial milestone has been reached that is known in *QQ* as *Version 1* and is now waiting for evaluation by various types of reviewers (including usability-reviewers and subject-matter expert reviewers for all relevant disciplines). There might be many or few, small or big, new versions, releases, and/or patches on the QQ level. How many depends on the number necessary for incorporating the feedback that is necessary for taking the next (more public) step in a responsible manner.

RRv1. The version variant RRv1 is associated with the first reviewed release that is intended for some productive use (albeit without particular stability guarantees yet). This is the Brief Evolvix way of saying "*ReviewedRelease*" and indicate that someone with the authority to release this variant has reviewed all pertinent issues and deemed the overall maturity to be sufficient for public release. For this study, this point will be reached, when FlyClockbase accompanies the final publication of this study in the form of a corresponding public release of FlyClockbase on Github.

TABLE P1. The *StablizingZone* of the *Project Organization Stabilizing Tool* (POST) system is defined by the *StabilityCodes MM* to *TT*, presented by their BEST Names.

Brief	Explicit	Summarizing Name
MM	*MockupModel*	MockupModel_UsedFor_RapidPrototyping_InformalLearning____ __ExpertimentsToBeThrownAway_StabilizingDesignNotCode
NN	*NewNonfunctional*	NewNonfunctional_UsedFor_NotYetFunctioning_DeepFoundations____ __ForLargerStableDesigns_ThatDoNotYetWorkForUsers
OO	*OperatesOften*	OperatesOften_UsedFor_Systems_PartiallyWorkingForEndUsers____ __while_StillMissing_ImportantFeatures_ToBeImplemented
PP	*PreProbing*	PreProbing_UsedFor_Preparing_PeerReviewAndPublicProbing____ __by_PolishingExistingFeatures_UntilSubmissionFor_Questioning
QQ	*QualityQuest*	QualityQuest_UsedFor_Questioning_AxiomsDataScienceAccuracy____ __RigorClarityUsability_InMany_ExpertBeginnerReviewRounds
RR	*ReviewedRelease*	ReviewedRelease_UsedFor_NewReleasesRecommended_by____ __QualityQuestEditors_after_AnsweringAllReviewerQuestions
SS	*StableSource*	StableSource_UsedFor_StunningSoftware_RunningInProduction____ __with_LongTermSuccess_and_VeryRareRevisionRequests
TT	*TrustedTested*	TrustedTested_UsedFor_Marking_VeryLongTermStableDesigns_in____ __WellUnderstoodDomains_AllowingBackwardsCompatibleGrowth

Both have been developed for the general-purpose programming features of Evolvix with the goal of facilitating the development of long-term backwards compatibility. More details have been discussed as the Evolvix BEST Names concept was introduced (LOEWE et al. 2016b).

Why stabilizing versioning? This version variant numbering system was selected to encourage responsible steps towards enabling long-term backwards compatibility for FlyClockbase and to enable it to be a *VBIR*s, i.e. a truly Versioned Biological Information Repository (vs being only a BIRs, where "Current" is the only well-defined state or versioning is incomplete). Stabilizing versioning alleviates the tension between repeated rounds of rigorous review as required for long-term stability and the flexibility to quickly experiment with risky ideas as required for innovation. Some of these ideas have to be tried out before it is possible to reasonably decide what to do with them (many of them should never appear in a system aiming for stability). The concept of a *StablizingZone* has been developed to alleviate a similar tension in Evolvix. It builds on the *StabilityCodes* that are part of the POST system and are shown in Table P1.

The role of TabFS in the stability of FlyClockbase. FlyClockbase depends on the stability of its underpinning storage infrastructure. As argued below, we decided against traditional databases, because the constant stream of upgrades easily imposes prohibitive burdens of IT administration work on those biologists who are likely to initiate and maintain *VBIR*s. To resolve this, FlyClockbase development has also been intertwined with the development of what we call 'TabFS', an extremely thin and transparent file system that sits on top of a standard file system. This study is not the place to appropriately describe and define TabFS; suffice to say it is being designed to have the abilities of a fully functional file system with additional features that simplify some work in computational biology. These features are combined with a priority on radical openness allowing the local user in the host operating system to change anything and everything (e.g. this allows a biological model curator to easily change the type system of a VBIR). These desirables of TabFS come at the price of having to deal with inconsistent states that are more likely to be caused by users in TabFS than in other file systems. Resolving this will require additional checks and redundancies to enable recovery of consistency when users generate errors. Thus, the biggest trade-off in this setup is a slower performance. However, a fundamentally important and conscious design decision for TabFS is to trade short-term performance for increased long-term durability. Practically, TabFS stores everything in tab-delimited tables of text

stored in the folders of a standard file system. It can be thought of as an organized

equivalent of CVS files, albeit with strict formal rules that enable compiler checks.

How does stabilizing versioning benefit FlyClockbase? The decision to buy into

the stability of major standard file systems does not mean that problems of storage

incompatibility cannot affect FlyClockbase. To efficiently organize FlyClockbase we

need to abstract recurring themes, and these abstractions need to be developed, which

takes time. FlyClockbase bundles these storage abstractions into TabFS; hence any

changes in TabFS that are not backward compatible will threaten the data stored in

FlyClockbase if not handled appropriately. This is where stabilizing versioning is

expected to be extremely helpful: it indicates to users how reliable a potential upgrade is

likely to be. It also indicates to developers and reviewers shared expectations about the

quality of the code they write or read. Thus, proper versioning is essential for stability of

FlyClockbase as a VBIR. Reporting more on our progress towards this goal is beyond

the scope of this study.

Types in FlyClockbase. In addition to using TabFS, FlyClockbase has also been

developing its own type system and content that requires coordinated changes and can

hence benefit from versioning.

Test case for Evolvix. Without detracting from its biological focus, FlyClockbase

provides an important real-world testing ground for developing approaches to enable

long-term backwards compatibility in Evolvix (LOEWE *et al.* 2016b).

Why not a conventional database? FlyClockbase is designed at a time when

data science is of growing importance for biomedical research (see (NIH *et al.* 2012)).

We have considered the many virtues of diverse well-developed database management

systems (from SQL to NoSQL to NewSQL). These guarantee consistency of data,

simplify complex searches via special query languages, and increase speed of access

for growing amounts of data that is appropriately organized (e.g. (JURNEY 2013;

MAHMOOD 2016)).

Unfortunately, these advantages usually come at a price that includes the hidden

or deferred costs of:

(i)   becoming increasingly dependent on a specific stack of software that is often growing in complexity,

(ii)   complicating installation or transport across platforms,

(iii) becoming too difficult for non-specialists to tentatively add new data and datatypes,

(iv) complicating the migration from an old (often rigid) database schema to one with new features (and complexity, but usually equally fragile).

Combined with poor documentation and sloppy naming, such problems can easily degrade the semantic reproducibility (LOEWE et al. 2016b) and, hence, the long-term usability of any data collection that builds on top of such tools. The Introduction of the main text discusses more aspects. As a result, pre-clinical biomedical research in the US has been estimated to invest about $7Bn/yr in studies that were deemed to contain irreproducible data analysis or reporting (FREEDMAN et al. 2015). Such an environment is hardly conducive to efficient biology, computational or otherwise; as a consequence, the critical importance for reproducibility of research results has been recognized across many disciplines (DONOHO 2009; GOECKS et al. 2010; KARR et al. 2012; NIH et al. 2012; STODDEN et al. 2014; JAMES et al. 2015; KARR et al. 2015a; KARR et al. 2015b; KENALL et al. 2015; POLDRACK AND POLINE 2015; STODDEN 2015; LOEWE 2016). As argued in the Discussion of the main text, simple text files have many desirable features. We decided to carefully build on those.

Logic challenges posed by missing data. Another notorious difficulty in many "standard" databases is how to indicate the absence of particular data, which can be interpreted as different types of zero. The difference between "not available", "not applicable" and various other types of "null" is widely recognized (ZANIOLO 1984; CANDAN et al. 1997; DE TRE et al. 2004; WARAPORN AND PORKAEW 2008; BOSC AND PIVERT 2010; HERNICH et al. 2011; THALHEIM AND SCHEWE 2011; HARTMANN AND LINK 2012; LIFSCHITZ et al. 2012; MARTINEZ et al. 2013; MIRZA 2015), yet standard databases struggle to provide automated support beyond an unspecified, potentially ambiguous 'NA' (may stand for *NotAvailable*, *NotApplicable*, *NotAllowable*, …). Further, they build on the Closed World Assumption (that all relevant data is always fully specified) and aim

to represent everything in the stark true-false dichotomy of Boolean logic. Current research in logic has provided many examples for the crippling effects of such black-white limitations when confronted with a more nuanced world of color (SMITH 2008). These problems surface very quickly, when attempting to reconcile the crisp true-false dichotomies of a world of Boolean variables with biological observations with a certainty that is less than clear-cut. To quickly allow for the identification of such biologically important cases without losing the advantages of the crisp semantics of Boolean logic, the following "OKScale" of four alternative states for a value of the type 'BioBinary' has been designed for Evolvix (see online material of LOEWE *et al.* (2016b), p. 16). FlyClockbase is adopting the *BioBinary* type to facilitate clearly marking the states described in Table P2.

The basic features of the *BioBinary* do not eliminate the need to find a way of representing the infinitely many more detailed types of OKO, KO, and MIS, which might be used to represent details about the infinitely many stages of incompleteness, zero, null-hypotheses, or potential mistakes and contradictions that could be encountered when analyzing biological data. It is extremely difficult to anticipate which of the more nuanced types of information beyond these four will be encountered while actually analyzing biological observations. Maximizing the flexibility for describing such exceptions, while providing useful guidelines to educate FlyClockbase users about important subtleties, improves the quality of the reported data and reduces the frequency of misleadingly strong statements that overstate their claim. Such statement can easily be generated by if the logic formalism of a system does not allow recording a weaker statement. Our most important goal is to have a logical formalism that is neither overly restrictive (which would prohibit the representation of certain observations), nor as unstructured and flexible as a blank page (which would make it very difficult to conduct any useful logic analyses). We found it helpful to work with lists of controlled vocabulary and only to add freeform comments for special cases, which could eventually add to controlled vocabulary if repetitive. The decisions about which columns are controlled, which are free, which entries make it into a controlled vocabulary list, and when a free-form comment will be extracted in order to populate a new column, and many more such decisions strongly depend on the subject matter and the content of a

VBIRs. We therefore decided that biological model curators would need the full ability to change any of the types in a VBIRs as they see fit to represent the observations they curate. Since VBIRs are Versioned, all these decisions can and should be properly reviewed to improve consistency.

TABLE P2. The *BioBinary* data type for cases in biology that are less than clear-cut.

OK	Indicates the full agreement with a specified test that has been completely executed (equivalent to Boolean "true");
KO	Indicates the full disagreement in all points with the specified test that has been completely executed (equivalent to Boolean "false");
OKO	Indicates any intermediate between OK and KO for cases that are well known not to be clear cut, albeit without storing any other information, such as a probability that could specify the distance from OK or KO (it is up to users to determine what they do want to store); No Boolean equivalent exist for OKO;
MIS	Indicates any mistake or problem that made it impossible to reach any of the other three conclusions, irrespective of the nature of this problem (it is up to users to decide how much about the potentially infinite complexities of such problems they wish to store; no Boolean equivalent exists for MIS).

This new data type has been designed for efficiently reasoning about biological observations with a certainty that is neither perfectly true nor false. More details elsewhere (LOEWE et al. 2016b).

Permissions, backups and the reliability of data storage. Beginners and experts, all have full administrative ("root") access to almost every aspect in FlyClockbase. The very open approach of FlyClockbase raises the question of how to guarantee the integrity of its use. While in theory a user can compromise FlyClockbase quickly, the chances of such an event with lasting consequences are much smaller than these unrestricted settings might suggest. When viewed in context:

(i) few users of FlyClockbase *aim* to compromise their installation; moderate hiding of critical infrastructure or clear indications warning about the potential of breaking sensitive parts is usually sufficient to prevent mistakes;

(i) the latest working release of FlyClockbase is never in danger if a release manager keeps backups;

(ii) distributed development of FlyClockbase poses the same challenges as all distributed information processing, namely, the merge/cache-invalidation/naming problem. It can be solved in principle by distributed version control systems (like Git), or by using cloud synchronized folders in combination with strictly following a version variant naming scheme to prevent loss of data (see Fig. 1BC in LOEWE *et al.* 2016b). We tested both approaches (Git and cloud-style automatic synchronization). We found both to be less inviting than they could be if a set of reliable scripts were to automate repetitive tasks in a way that reliably excludes loss of data (and without adding to the cognitive load of the user).

While it may seem that widely known database technologies are free from these problems, they in-turn come with independent problems of their own. For example, FlyClockbase may give ordinary users too much administrative power. Many other databases resolve this problem by locking down access. Yet, in turn these databases require special expertise or permissions for exploring new avenues of recording a richer spectrum of biologically relevant difficulties with contradictory scattered big data. Even experienced database architects may use natural language for the very early initial stages of database development. We aim to give model curators the ability to do exactly this '*in situ'*.

On some challenges in logic work. Capturing details important for describing biological observations in a consistent way can be challenging research in logic formalisms, because of the vagueness of some biological data (SMITH 2008). Most biological model curators are not likely to have the training in formal methods to define the new formalisms that remain at the end of this research. As argued above and in Figure 11 (main text), curators can still contribute a lot by providing examples and detailed expertise on how to interpret their various challenging aspects, which is impossible for non-biologists to do. To facilitate the practical work necessary for enabling this type of discussion, a number of smaller problems need to be resolved:

- *Form of recording.* Any persistent record of observations in logic needs to encode the observations of principles or bugs in some formal way that can also be read by other persons. In a world of infinite resources, it does not matter how these observations are encoded, as long as their semantics is completely reproducible (LOEWE *et al.* 2016b). However, for most biologists engaged with model curation, patience with logic formalisms is a limited resource. Approaches used in practice for communicating logic problems ought to be as simple to use as possible. A blank page or an arbitrary text description work well, written or spoken; it is the task of a person with more understanding in formalisms to translate such fragmented insights into a more coherent formal picture. Additional burdens, such as transforming descriptions of logic problems into a particular logic formalism, are probably counterproductive. Many curators might be tempted to indicate that they have nothing to report, simply to avoid the difficult task to re-encode their observations, e.g. in a relational database formalism. While certainly possible, it does not mean that this is advisable; in principle, these problems could also be expressed in the form of Gödel numbers (GÖDEL 1931), but this would only add huge amounts of inessential complexity. Thus, the simplest possible storage medium is probably best.

- *Durability of recording.* For the sake of argument, let us assume that there was a very straightforward way of recording all such statements in a database. Then such work can become unnecessarily complicated, if the database system requires frequent complex updates that are triggered by external database developers aiming

to deliver ongoing improvements like new features, security, performance or more. If such upgrading becomes excessive, paper and pencil might become increasingly attractive.

- *Size and complexity.* Documenting progress in debugging or developing logic does not usually generate large amounts of data, but the few cases that are there are important and complicated. Hence, an uncomplicated, distraction free storage medium is desirable.

- *Speed of copying.* For efficient communication, it is desirable to find a storage medium that is easily copied through communication channels. This is where non-electronic approaches have their greatest drawbacks.

Trade-offs. To solve these challenges in a practical way, let us consider the following advantages and disadvantages of paper, file systems and databases.

- *Paper*, for example, does not suddenly refuse to operate as expected because it needs an upgrade – a plus for stability. Its durability is legendary and generally measured on the scale of centuries. However, the speed of writing, reading, and copying is slow and error prone.

- *Dedicated databases*, relational or otherwise, when designed for fast retrieval of certain types of answers are hard to beat in speed for that particular application. Yet that speed will drop to zero, if the latest essential upgrade is not applied and all data in the system becomes inaccessible (see tombstone example below).

- *Filesystems* in contrast, offer the best of both worlds: almost the reliability of paper (assuming that there will always be electricity and hardware that enables access to the file system and perpetuation of backups); almost the flexibility of paper (assuming corresponding types of files); almost the speed of databases (assuming the data is organized so that the speed of reading or writing approaches the limit supported by the hardware). The big question for file systems is whether a way of organization can be found that keeps all important data appropriately organized.

Long-term stability in VBIRs data structures will enable new biology.

We suggest that a compiler could help curators to efficiently maintain long-term stable VBIRs. It would do so by guaranteeing that VBIRs are easily returned to a consistent standardized state. This standardization would have to be stable enough so that these VBIRs can serve as long-term foundations, which will eventually make it feasible to develop more consequential independent VBIRs on top of the more causal ones.

If foundational VBIRs describe causal genotypic or environmental information and more advanced VBIRs describe more consequential molecular, cellular, physiological, or other phenotypic information, then VBIRs enable the implementation of a full fitness-causality network that maps the genotypes and environments of an organism to its phenotypes in a transparent mechanistic way that connects well to the latest updates of all data that is available – if all relevant data and data structures are well curated and appropriately versioned for stability.

Tombstone example. Here we illustrate with an example the very real long-term danger to biological research posed by database technology that is not long-term backwards compatible. The TIGR Gene Indices were first published at http://www.tigr.org/tdb/tgi (QUACKENBUSH *et al.* 2000). After their introduction, the TIGR Gene Index databases quickly became a well-known tool for biological discovery (LEE AND QUACKENBUSH 2003; PERTEA *et al.* 2003; LEE *et al.* 2005). They are now no longer available online as documented on their tombstone: http://compbio.dfci.harvard.edu/tgi/ In case the tombstone itself disappears eventually, here we paraphrase its 2017 report of some vital statistics about TGI.

Supported by NIH, DOE and NSF 1998-2010, the relevant TGI papers were cited >2000 times. When the tombstone was written the TGI website still received >7 million hits per year (assuming the actual number on the tombstone was a typo). When funding ended in 2010, the team continued to maintain the website, but the hardware and software required behind the scenes began to fail. Effective July 15, 2014 operations had to be suspended, because there were not sufficient funds to maintain it properly. The software powering TGI (DFCI Gene Indices Software Tools) was 'fossilized' to

ftp://occams.dfci.harvard.edu/pub/bio/tgi/software/ and the data sets to

ftp://occams.dfci.harvard.edu/pub/bio/tgi/data/.

It is not up to us to comment on TGI's science or its funding history. Neither is relevant to our main point: extinction is a real risk for *VBIR*s and fossilization to some archive is not a real life-saver. The haphazard nature of funding for biological information repositories is well known and a significant source of concern (EMBER *et al.* 2013). Less obvious is the impact of a stable VBIRs compiler for TGI. Imagine the software behind TGI would use appropriate abstractions and thus not fail. Imagine it could continue to operate reliably on different hardware, including that of users. Imagine the software would be long-term backwards compatible. Imagine it could help many biologists to contribute to curation of TGI. Imagine a whole community would annotate, improve, deprecate, or otherwise edit various aspects of TGI in order to preserve its benefits or point to improved successor tools. Imagine other biological research codes had built on long-term stable parts of TGI and could all continue to operate simply by copying TGI to a local hard drive. Would that make a difference?

2.   Human error analyses: approaches, challenges, efficiencies

Problems with tracing the identity, availability, accuracy, precision, and reliability of data have been the topic of numerous investigations and recently received renewed attention in the context of assessing potential uses of scattered biological data sources and big data (e.g. see (REASON AND MYCIELSKA 1982; REASON 1990; REASON AND HOBBS 2003; GOLDSTON 2008; GITELMAN 2013; GRIMES *et al.* 2013; MCCALLUM 2013; REASON 2013; BLANKENBERG *et al.* 2014; REASON 2015)).

FlyClockbase presented us with an excellent opportunity to explore numerous issues related to the quality, accuracy, and reliability of complex collections of information that aim to integrate data that can be incomplete, uncertain, contradictory, erroneous and scattered across a wide range of sources. Much expertise in biology currently exists in such less-than-well-defined states that can be difficult or impossible to process computationally. Thus, we expect that some of the lessons we learned while compiling FlyClockbase become increasingly important as biologists expand their use of computational resources throughout their work. While it is beyond the scope of this study to appropriately review all corresponding challenges and potential solutions we encountered, a few observations of common challenges of general interest are reviewed next.

Finding capable curators. Compiling nontrivial resources of biological information requires a substantial and sustained effort from researchers who bring a certain set of skills. They need to (i) possess the background expertise necessary to understand the importance of the information they compile, and (ii) be motivated to collect information from various scattered representations into a single, less idiosyncratic representation that constitutes the new resource. New resource creation, as listed above, usually requires more effort than experimental or computational biologists are willing to dedicate. This might require researchers with a special interest in high-quality information collection regarding the respective system. Like airplane pilots, curators will need training in appropriate error handling.

Errors are inevitable. A broad body of research has shown that human errors in any non-trivial data processing are inevitable and that error rates increase with the complexity of a task (REASON 1990; REASON AND HOBBS 2003; REASON 2013; REASON 2015; PANKO 2016). This is true in particular for the use of spreadsheet software (PANKO 1998; PANKO AND SPRAGUE 1998; PANKO AND AURIGEMMA 2010), for which rates for various types of error have been measured. Our own results confirm that some errors (e.g. simpler data entry) occur with lower rates than other more complex errors, that stem from occasionally forgetting non-obvious steps in a complicated procedure (see Table 4). We can confirm previous findings, showing that even the most motivated researchers with substantial training and background expertise will inevitably err when transferring information in complicated scenarios where many complex details could go awry (see also (LESGOLD *et al.* 1988; GALLETTA *et al.* 1993; GALLETTA *et al.* 1997; PANKO AND SPRAGUE 1998)).

Errors are hard to detect for humans. Proofreading software is well known for not catching all spelling errors. It is also well known, that not all humans catch all spelling errors. Yet spelling errors are the simplest errors. All others are more difficult to detect in comparison. Errors in logic are particularly difficult; however, the worst category of errors are errors of omission (PANKO 2016).

Given the various complex operations required for integrating and analyzing data with FlyClockbase, FlyClockbase has competitive error rates, as found when re-visiting parts of the database (see Table 5). However, manually extracting even simple time series traits (e.g. peak-valley times) has proven more error prone than we expected. The complexity of traits can rise substantially as increasing numbers of important conditions and special cases need to be observed for extracting a valid trait. Eventually complex decisions, when integrating less-than-perfectly fitting new observations into biological information resources, might easily generate very complex error scenarios. This is mainly because such work requires many decisions about which types in the data and in a VBIR are functionally equivalent and when effective differences are large enough to justify the introduction of new categories. The complexity of such offline

decisions could rival the complexity (after preparation and training) of online decisions in mission-critical scenarios that have been analyzed elsewhere, from cardiac surgery (CARTHEY *et al.* 2001; REASON 2005) to space flight (NASA *et al.* 2011; HOOEY *et al.* 2014). In many of these scenarios, human failure is catastrophic; correspondingly sophisticated human reliability analysis methods have been developed to ensure that human error is exceedingly unlikely - if not preventable, in principle (REASON 1990; NASA *et al.* 2001-09-30; REASON AND HOBBS 2003; NASA *et al.* 2006-07; NASA *et al.* 2011; REASON 2013; REASON 2015).

Human error analysis and prevention methods are difficult to develop.
Many of the methods that are appropriate for investigating and preventing human error in contexts such as space flight, nuclear power plants, or aviation (NASA *et al.* 2006-07; NASA *et al.* 2011) are much too complex and costly for applying them towards improving the reliability of biological information resources like FlyClockbase. This is true even if the complexity of problems generated by resources such as FlyClockbase rivals or exceeds the complexity of navigating a space ship.

The huge immediate and catastrophic danger and costs of errors in navigating space ships and running nuclear power stations make it obvious and easy to justify the large effort required for in-depth analyses of human errors. In contrast, adding data to VBIRs seems harmless; it certainly is if only the immediate impact is considered, and it probably also is on the long term. However, in biomedical research there is a small chance that errors in VBIRs are not as harmless as they seem if they remain uncorrected. The following scenario is difficult to exclude categorically in bio-medical research. Imagine that some researcher integrates data into some medically relevant VBIR and makes an unnoticed error that remains in that VBIR for a long time and even replicates across repositories. If that error prevents the discovery an important new medical cure by leading a researcher to the wrong conclusions, then cost and risk calculations are no longer trivial.

It is not clear how to resolve this problem. If each VBIR produced its own error analyses, costs would quickly spiral out of control and few resources would remain for actual biological research.

Error analyses require a substantial level of biological expertise, yet are perceived as 'tedious' and 'non-biological' (by experimentalists) or 'non-automatable' (by computationally oriented biologists). Thus, recruiting experts for such error analyses can be exceedingly difficult or prohibitive, despite their importance for modern biology. The following potential 'solutions' are also likely to be counter-productive on the long term:

(i)   *Omitting error analyses.* While cost-effective during the initial stages of compiling a new resource, ignoring the potential for errors by not engaging in error analyses will result in a growing number of databases with questionable quality. Whenever databases have been analyzed, some rate of errors has been found (e.g. (JONES *et al.* 2007; HUTCHINS *et al.* 2010; WEIL *et al.* 2015)). It should not usually require open letters or other drastic measures (e.g. (DRAKE *et al.* 2008; ALLISON *et al.* 2016)) to possibly get errors corrected. The current state of software tools makes post-publication changes very difficult and most databases do not offer an easy way to submit proposed changes for review and efficient inclusion by administrators. While this explains the reluctance to update the latest available information, it is not conducive to advancing the integrity of scientific data. If no resolution to this error crisis can be found on the long-term, the resulting loss of quality will lead to a loss of reliability, reproducibility, and ultimately to a loss of trust in 'data-based' results. These costs are not immediately obvious to the authors of a new resource, but they will eventually manifest somewhere in the scientific community, either as irreproducible results, misguided research directions, or unnecessary 'confusion-tax' for the users of a resource.

(ii) *Automating database operations*. Some types of errors are greatly reduced by the use of automated database technology (e.g. leveraging SQL to keep indices up to date and reduce redundancy,). This is particularly effective for types of errors that occur in large numbers, thus making it easy to automate detection.

      Heterogeneity. However, the highly heterogeneous nature of many advanced biological information resources can easily generate as many potentially relevant exceptions as entries. Relevant exceptions not addressed by

the built-in logic of a database management system are at high-risk of causing errors, since errors by omission (of a special condition) are particularly difficult to detect (e.g. see discussions of *BioBinary*).

Such errors can be even more difficult to detect in automated systems, if they are not already detected automatically. Automation might tempt researchers to become over-reliant on computational results they no longer understand (especially if implemented by non-biological experts). The substantial computer science literature on compiler construction demonstrates that the use of automation via software does not protect against the innumerable types of errors that can be added to source code and can corrupt results (if not caught by a compiler; see (PARSONS 1992; AHO 2007; COOPER 2012; GRUNE 2012)). This suggests the following interesting conclusion:

Automation trades some human errors for others. For example, inconsistency errors that often occur in data that is entered manually are avoided by automation through programming; get programming can easily generate inconsistencies of its own that humans would not produce. Since about 2% - 5% of all lines of source code have been estimated to contain some errors (PANKO 2016), it is far from obvious how automatically handling the many special cases in resources like FlyClockbase can make additional analyses of various human errors superfluous.

(iii)   *No retrospective meta-analyses.* Not engaging with data analyses that trigger the need for error analyses robs biology of important perspectives. This would be equivalent to a call for stopping such otherwise non-controversial biological research. The impact on integrative work and relevant modeling efforts could be devastating. Hence, this options is not attractive.

While currently no ideal solutions exist, raised awareness of the problem is likely to eventually contribute to the development of appropriate solutions. As argued above in the context of compiler design: seeing the problem in the first place is often the most difficult step. Maybe compilers could also help by automating many tedious aspects of these error analyses.

Error analyses could be amortized across resources by compilers. As argued above, developing appropriate error analyses for a single *VBIR* is not feasible. However, our experience with developing FlyClockbase suggests that a substantial number of essential tasks are recurrent and of comparable complexity across the development of a groups of *VBIR*s. Examples include the need for handling missing data, inapplicable data, imprecise ranges (rather than precise values), heterogeneous method descriptions, comparisons of wild types to mutants, and many other aspects of biological interest. Thus:

> *The most efficient solution to improving the quality of VBIRs without*
> *exploding costs is to develop an automated compiler that can*
> *automatically test for all known and formally described cases of*
> *errors and supports a programming language that integrates biology*
> *expertise to simplify the description of test cases by biologists.*

Programmers frequently say that it is important to use the right tool for a given programming task, yet no such tool exists for the compilation of complex *biological* datasets. We will not repeat here the substantial number of reasons why such a language would be helpful and why current (non-biological) programming languages are insufficient (see Online Material and additional reasons discussed in Loewe et al. (2016a)).

Efficiencies of scale. In the long term, it would be more cost efficient to construct a compiler that can read in all tables of FlyClockbase (or other *VBIR*s) and produce a report of all inconsistencies and errors that require human attention (as well as produce tables with all requested search results). Thus, certain types of errors would be detected automatically. Error detection modules and other solutions would be implemented only once and would simultaneously improve the reliability of all VBIRs construction efforts.

For those who wonder why we did not directly implement such a compiler, we would like to point out that - as so often is the case in computational biology - the overall time for obtaining a given biological result by manual 'compilation' is usually much shorter than the time it would take to implement a corresponding compiler. This certainly was the case here. For example, compiling the time series data of wildtype flies and checking the integrity of PER and TIM data to reasonable reliability was completed *much* faster manually, then if we had attempted to automate it by constructing a compiler for this. This cost calculation changes very quickly if large amounts of integration work are anticipated over long amounts of time in FlyClockbase or other *VBIR*s. There seems to be little debate over the need for curating biological data, especially as it keeps growing (BOURNE AND MCENTYRE 2006; BURKHARDT *et al.* 2006; SALIMI AND VITA 2006; HOWE *et al.* 2008; BURGE *et al.* 2012).

Tool development and funding strategy development are connected tasks. Some development of tools beyond the scope of a single project already started, addressing the large amount of text mining done by biocurators (WEI *et al.* 2013; RAK *et al.* 2014; SINGHAL *et al.* 2016). However, developing clear and successful funding strategies has remained a challenge (EMBER *et al.* 2013; ORCHARD AND HERMJAKOB 2015; RODRIGUEZ-ESTEBAN 2015; REISER *et al.* 2016). The inefficiencies of present-day curation work in biology are substantial and might be sufficient to convince most funders to invest elsewhere. Our experiences have shown that many of these inefficiencies could be alleviated in principle by appropriately constructed tools. By today's standards it was unimaginably laborious to sequence a few genes in 1977, or a single human genome in 2001 (SANGER *et al.* 1977a; SANGER *et al.* 1977b; LANDER *et al.* 2001; VENTER *et al.* 2001; HAYDEN 2014; SHERIDAN 2014; TELENTI *et al.* 2016). Given the lack of appropriate tools, the lack of funding for corresponding work might be less surprising. Funding for sequencing human genomes in 1977 was negligible, if not $0.

3. DISCOVARCY questions on source code documentation style

The attitude of some programmers towards documentation can be summarized as:

The source code is the documentation.

Indeed, the agile approach to software development tries to avoid documentation, which is costly to write, and for fast developing code bases it is constantly out of date unless maintained with a large amount of effort (e.g. see comments in (HENRY 2013)). As a corollary, one might conclude that documenting scientific research code is a waste of time, since by definition such code is moving fast[1]. Unfortunately, such practice results in semantic irreproducibility (LOEWE *et al.* 2016b), which hampers research (FREEDMAN *et al.* 2015). On the other extreme, again, it is neither reasonable nor efficient to demand that all research codes are refactored and documented excessively. For the most part, research codes are indeed only used very few times.

For example, consider the highly specialized 'unique research application' R-script code, which was written especially for this study. It is likely to serve only the purpose of re-running the analyses of its paper, facilitating the computational reproducibility of the study's results. While writing that code, there was no mandate to use best practices for software engineering – unless it would help get the job done faster or more reliably. Hence, no refactoring for longer-term use done for increasing the quality of the source code. Good writing requires multiple rounds of revision; good code requires substantial refactoring. Depending on the occasion, this may be effort well spent, or time wasted. Our supplementary code for this study lives in this tension, and so do many other research computing codes written for specific unique analyses. How can such code be made more readable in order to help other researchers benefit from the work of the authors, and not waste their time on inessential complexities? This question is at the heart of semantic reproducibility (LOEWE *et al.* 2016b).

The following table introduces the DISCOVARCY questions on documentation style. The acronym DISCOVARCY highlights typical problem areas that can sometimes

---

[1] It is important to distinguish such fast-moving research code from other, more mature scientific code that implements important algorithms which have become important research tools and are thus expected to behave reliably.

easily be improved once awareness increases. The acronym is on purpose designed to illustrate the many transitions small and large that are between worst and best practice. Some of the distinguishing characteristics are easy to change while programming, but prohibitively costly later (e.g. choosing summarizing variable names). On other occasions, it is prohibitively costly to refactor a whole code base merely to make it easier to understand.

Such cost-quality related tensions highlighted by DISCOVARCY are linked to a fundamental trade-off faced by every innovator aiming to produce features reproducible by others. The trade-off forces numerous choices between the following ideals:

- *Reproducibility* (much time for tests and documenting, little time for new features);

- *New features* (little time for tests and documenting, much time for new features).

Every scientist needs to navigate the contrast between these ideals that is particularly sharp in research, where the goal is to make innovative new discoveries, while enabling others also to reproduce these innovations.

The same tensions exist for software developers and programmers, only that reproducibility comes with a human and a machine aspect. For machines, reproducibility is the repeated execution on machines; this links the diversity of machines included in this claim directly to the cost of reproducibility. The human aspect of reproducibility implies that a human being other than the developer can read and understand the code correctly, an ability defined by the quality of the Code2Brain interfaces involved (LOEWE *et al.* 2016b). As with machines, producing something that is more widely understood is usually more expensive.

It is worth noting that much of the work needed for improving clarity and guaranteeing the documentation of research codes could be delegated to a compiler that can easily translate between brief code preferred by developers and the more verbose alternatives more easily understood by newcomers. The Evolvix BEST Names concept has been developed in order to facilitate such translation (LOEWE AND KEEL 2014; LOEWE *et al.* 2016a; LOEWE *et al.* 2016b). Documenting code has a long history (KNUTH AND LEVY 1994) and requires more attention than this study can offer here. Hence, more details will have to be presented elsewhere.

1
2
3

TABLE D1: The DISCOVARCY questions on documentation style help highlight source code that is hard to read and might offer ideas for improving it efficiently.

Brief dNames	Quicker to produce, Costlier to consume Explicit Name for **Coder centric view**	**discovarcy vs DISCOVARCY** **Questions and comments on tension**	Costlier to produce, Quicker to consume Explicit Name for **Reader centric view** Brief Names
d	**Documentation**	Does documentation help the writer or the reader or neither? Supporting readers requires good writing, which may require prohibitive amounts of time (and writing skills) on part of programmers.	**Documentation D**
i	Is	"The Documentation *IS* the Source Code"? "The Documentation *Includes* Source Code"?	**I**ncludes **I**
s	**Source** Secrecy	How much advanced rare expertise is assumed? Are coding tricks explained? Are they marked?	**Source S** Simplicity
c	**Code** Complexity	Essential or inessential complexity? Complexity is easier than clarity. Simplicity is hard.	**Code C** Clarity
o	**Online** Obligatory Odysseys	Is all info included in one place for open access downloading for offline work? Are relevant links included or many open-ended searches needed?	**Online O** Optionally Offline
v	**Variable** View Variety	Is the meaning of variables and functions obvious from using explicit or summarizing names? Are they at long enough in the age of copy&paste to exclude random reuse of e.g. "x"?	**Variable V** Validation Verification
a	Arbitrary **Assumptions**	Problems can be solved in many different ways. Is explained why a given way was chosen?	Argued **A**ssumptions **A**
r	Random **Restrictions** Rare **R**eferences	Are meaningful automatic restrictions enforced? Did developers include links + refs they used?	Rigorous **Restrictions R** Relevant References
c	Chance Contradictory Chaotic **Comments**	Keeping comments updated may be too much work. Does a marker for 'retired text' exist, which indicates mixed quality and potential inconsistencies?	Complete **C** Clear Compelling **Comments**
y	**Your** Years	Understanding confusing code can quickly take impossibly long. Reading clear code is learning from a teacher. Would you enjoy reading your code if you hadn't written it? Will you in 10 years?	**Your** Yield **Y**

4 *Properties that make documentation poor ('discovarcy') usually exist on a continuum with those*
5 *that make documentation great ('DISCOVARCY').*

4. Example of an error a compiler could have caught by type checking: Effect of linearizing clock times on distributions of *Peaks* and *Valleys*

Please see Methods in the main text for motivating the linearization of times and for details on how it works. To illustrate the effect in the context of real data analysis, we selected types of mRNA and protein from FlyClockbase that are affected to varying degrees. Their observed peak and valley times are in Table S1, which compares their respective *ObsRaw* (non-linearized) and *ObsMod* (linearized) times.

Recall. For observing the impact of linearizing time, we remember that the time measurements in *ObsRaw TimeSeries* belong to the time type *DailyZT*, or *DZT*. If this information about types is understood by a compiler, then the compiler will know that such time measurements are recorded in time units of decimalized *Hours* and that they are reset to 0*h* at the dawn of each day, which will reoccur every 24*h* and increase a separate day counter by +1. If we ignore the day counter as usually done in circadian biology research, then *DZT* is a cyclical time type, which simultaneously is also a cyclical real number type. In contrast, all *ObsMod TimeSeries* are measured in linear time without resets, as defined by *ContinuousZT, or CZT*. A compiler that understands this type information will know how to convert *CZT* to *DZT* and back (by adjusting the additional number of days that have occurred; see main text). If the compiler understands the assumptions required for computing valid summary statistics, it can check if those assumptions are met.

For example, using the well-known equations for inferring the mean of a Normal distribution requires that all values exist on a linear real number line, before attempting to calculate averages. Using values of a fundamentally different type (e.g. with the value `'MyText'`) would result in a compiler error (“`Cannot use 'MyText' when averaging`”). For time measurements of type *CZT*, the decision is easy, because the definition $0 < CZT \leq$ infinity can never generate cyclical time. For DZT time types, the decision is not as easy, because definitions like $0 < DZT \leq 24h$ do not imply the end of time once the 24*h* mark has passed. Flies, clocks and the rest of the world simply move on and will generate new data points. A sloppy *DZT* implementation will ignore the day-counter and act as if controlled by a time-loop – albeit with no effect on the linear time

experienced by flies. The logical contradiction between the existence of a time-loop as assumed by such programs absence of such a time-loop in real life does not always matter. For example, if comparing only events from a *single* day, nothing can go wrong. However, some time series in FlyClockbase are longer than 24*h*, which can generate profound confusion, as explained next.

Problem in principle. To illustrate the problem, let us consider an artificial dataset of *DZT* peak times for which a mean of 23*h* has been inferred for the first day and where the observed min and max has been 22.1*h* and 23.9*h*, respectively. Despite a circular number type, calculating a mean of 23*h* is appropriate; no data is included from another day; hence, all times are effectively linear. Let us assume that the next peak added occurs 90 minutes after 23*h*. If measured in *CZT*, this new peak would add 24.5*h* to an expected mean of 23*h*; the time remains linear and the mean remains valid. But expressed as *DZT*, this new peak would appear as a big outlier at 0.5*h* in simplistic comparisons with a mean of 23h (which implicitly drop a day). This type mismatch would misrepresent a peak of +1.5*h* after 23*h* as -22.5*h,* an erroneously large distance with the wrong sign. Such errors will bias inferred means in the wrong direction and greatly inflate inferred variances, possibly obscuring genuine biological signals. If undetected, these errors may substantially bias biological conclusions at unspecified cost to both reproducibility and biology.

Problem in practice. Initially we were not aware of these logical subtleties in handling different types of time. Calculating averages and variances requires a linear scale, a condition that is *completely* met out-of-the-box for all *Peak* or *Valley* times of some core clock components reported in Table S1. In these cases, time-linearizing does not affect summary statistics, since all observations already belong to the same day (e.g. TIM Protein *Peak*, 18.41 ± 2.54*h*). However, if some observations belong to the previous or next day, a change in summary statistics can be observed (e.g. PER protein *Peak: Raw* 16.77 ± 5.48 to *Mod* 19.51 ± 3.40 *h SD*; see also the other average values in bold). To interpret SDs, it is instructive to consider random times of a day.

TABLE S1. The impact of linearizing circadian time on summary statistics like the *Peak* (A) and *Valley* (B) traits of time series.

A		Peak Median				Peak *Avg ± SD*				
Clock Component	n_{Raw} n_{Mod}	**Raw** Circ	**Mod** Linear	**Mod** **Maj**	**Mod** **Odd**	**Raw** Circ	**Mod** Linear	**Mod** **Maj**	**Mod** **Odd**	n_{Maj} n_{Odd}
clk mRNA	12 12	6.00	3.56	3.56	5.03	**8.79** **± 8.59**	2.79 ± 3.16	3.56 ± 1.12	3.02 ± 3.73	4 8
per mRNA	88 88	14.9	14.9	15.0	12.35	14.62 ± 1.68	14.64 ± 1.65	14.72 ± 1.01	14.48 ± 2.87	68 22
tim mRNA	48 48	14.83	14.83	15.0	12.0	14.67 ± 2.09	14.54 ± 1.92	14.85 ± 0.98	14.14 ± 2.66	30 22
PER protein	36 36	18.93	19.0	19.0	16.87	**16.61** **± 5.99**	17.72 ± 4.79	18.61 ± 2.10	19.71 ± 5.90	30 10
TIM protein	39 39	19.0	19.0	19.0	19.0	18.03 ± 3.50	18.65 ± 2.83	18.50 ± 1.35	18.65 ± 4.29	35 11

B		Valley Median				Valley *Avg ± SD*				
Clock Component	n_{Raw} n_{Mod}	**Raw** Circ	**Mod** Linear	**Mod** **Maj**	**Mod** **Odd**	**Raw** Circ	**Mod** Linear	**Mod** **Maj**	**Mod** **Odd**	n_{Maj} n_{Odd}
clk mRNA	30 30	14.97	14.97	14.97	14.50	14.33 ± 1.67	14.33 ± 1.67	14.58 ± 0.69	14.15 ± 2.12	17 17
per mRNA	24 24	5.0	4.5	5.0	4.75	**6.61** **± 6.22**	3.61 ± 2.60	4.20 ± 1.36	3.65 ± 3.40	20 10
tim mRNA	21 21	5.0	5.0	5.0	5.0	4.86 ± 1.52	4.86 ± 1.52	5.13 ± 0.81	4.84 ± 1.37	18 11
PER protein	47 47	9.0	9.0	8.68	12.0	9.01 ± 2.17	9.27 ± 2.10	8.68 ± 0.92	10.08 ± 3.06	32 15
TIM protein	36 36	4.0	4.0	4.0	10	6.19 ± 3.26	6.19 ± 3.26	4.83 ± 1.51	9.66 ± 3.98	29 10

Bold values indicate suspiciously high SD suggestive of excessive noise. ObsRaw are in circular time and not yet linearized; ObsMod are in linearized time. Mod refers to data from the Mod6 PeakValleyTables; see main text Figure 3. To compare summary statistics of distributions that mostly come from usual or extreme values, statistics are also calculated for Maj (Majority) or Odd values. The counts of these values

are given by n_{Maj} or n_{Odd} for usual majority or odd extreme values; these were collected from the middle of ObsMod or from its margins, respectively. For further explanations see text above. This table also draws on summary statistics of Odd6 to show the increased variability of odd values. The other results in this table are from a comparative TS tables analysis completed on 2016-08-16 for Obs Raw, Mod, and Maj. No outliers were removed for calculating Raw, Mod, or Odd. The number of Obs included is given by n_{Raw}, n_{Mod}, n_{Maj}, and n_{Odd}, for Raw, Mod, Maj, and Odd, respectively. For related analyses see also Table 4 in the main text, as well as Table 6, Table 7, Figure 6, Figure 7, and parts of the Supplemental Statistical Analysis.

Uninformative DZTs. As reported in the main text, $SD > 6h$ is on the order of the SD of a uniform distribution across the day (deterministic 1h intervals from 0 to $24h$ result in $SD \pm 7.36h$ around $12h$ and drawing 1,000 corresponding uniformly distributed values does not fundamentally change the results (SDs: 6.81, 6.92, 7.02 h). Informal tests reducing the sample size to 20 presented a similar picture, albeit with substantially more noise (as we would expect for sample sizes comparable to the number of time series for a component in FlyClockbase).

Detecting excessive noise. We can use the random *DZT* results to detect some cases, where a clock signal is known to exist, but measurement or data handling problems have obscured the signal dramatically (so it looks like random noise). Indeed, Table S1 presents such cases. Peaks of Raw *clk* mRNA ($SD \pm 8.59h$), PER protein ($SD \pm 5.99h$), and *per* mRNA valley ($SD \pm 6.22h$) all show this problem, but there are also enough other cases to clearly demonstrate that not all time series are affected.

Solution by linearization. Close inspection indicated the problem (see above). We used these insights to devise a linearization procedure. The manual effort required for detecting the problem in the first place and then developing a solution consumed substantial amounts of research time. Other researchers will have to retrace or re-invent our work at substantial cost if they aim to avoid this problem. If a solution could be made available in the form of a compiler, then others could use and build on our results with almost no effort.

How a simple VBIR compiler could help. Compilers can easily detect type-mismatches such as "*DZT*" vs "*CZT*" and plainly refuse any operations that require any mix of types. While this would mean that our data could not be analyzed with automated help, it would also guarantee that the root problem of type mismatch is brought to the attention of researchers.

How a more advanced VBIR compiler could help. Sophistication beyond plain checks of type labels requires dedicated code addressing specific problem using domain specific biological expertise. For example, a *VBIR* compiler with definitions for a well-developed type system for circadian time series analysis could automatically check for potential *DZT* vs *CZT* confusion when calculating statistics beyond day 1, while allowing day 1 to be calculated if all data points are indeed from that day. Such a compiler would use a test for matching types to deduce the implicit drop of days from inspecting the type mismatch in combination with the actual data available at days beyond the first and explore remaining safe options. If user-friendly enough, it would create an understandable error diagnosis and alert the biological model curator in charge. If intelligent enough, such a compiler will offer a list of potential solutions from which the biologist in charge can choose efficiently how to address this problem and newcomers could request additional information. If analyses could only be performed by lucky accident (because *this* dataset allowed it, but others might not), then this compiler could produce a warning to that effect, so curators can anticipate potential problems as new data arrives.

Other examples. It would be easy to create a long and detailed list of cases where appropriate support by a compiler could have speeded up our work with FlyClockbase. Examples include the lack of support for expressing ranges or three-point-estimates for parameter values or measurements with substantial uncertainty, the lack of warnings when spreadsheet parsing or calculating algorithms come across values of the wrong type, the lack of appropriate number systems that guarantee correct handling of cyclical and linear numbers and many more.

Compilers that directly assist in biological research. A sufficiently advanced compiler would be like a helpful assistant[2], finding errors like needles in haystacks of potential problems, cutting debugging time, and preventing untold additional problems on the way. Such a compiler will be useful to the degree that it can provide diagnoses and abstractions for detailed and difficult challenges. These challenges need to be solved first by a human, who can then build/teach a compiler how to solve the problem.

Once the compiler will then be able to routinely perform the corresponding work the cost savings for *VBIR* development and teaching become substantial (the compiler can easily point to additional resources and more as demonstrated by the integrated help function of the R statistical programming language. If statistics has its language in R, why does biology not have its own language? The efficiencies gained would allow the scientists of today to enable the next generation of scientists tomorrow to get much faster to stand on the shoulders of giants. The compiler would be like an aerial cableway, which gets others faster to a place of uncluttered overviews.

---

[2] See http://elm-lang.org/blog/compilers-as-assistants for an exposition of what it might mean if compilers actually operated as assistants. This problem is inextricably linked to questions of syntax design. For example, see the importance of operator priorities (Razali, *et al.* 2015) and parsing which impacts the flexibility and quality of error messages that a compiler can produce (see Grune & Jacobs, 2008).

LITERATURE CITED IN SUPPLEMENTAL MATERIAL

Aho, A. V. L., Monica S.; Sethi, Ravi; Ullman, Jeffrey D., 2007 *Compilers: Principles, Techniques, and Tools*. Pearson Education, Boston, MA.

Allison, D. B., A. W. Brown, B. J. George and K. A. Kaiser, 2016 Reproducibility: A tragedy of errors. Nature 530**:** 27-29.

Blankenberg, D., J. E. Johnson, J. Taylor, A. Nekrutenko and G. Team, 2014 Wrangling Galaxy's reference data. Bioinformatics 30**:** 1917-1919.

Bosc, P., and O. Pivert, 2010 Modeling and Querying Uncertain Relational Databases: A Survey of Approaches Based on the Possible Worlds Semantics. International Journal of Uncertainty Fuzziness and Knowledge-Based Systems 18**:** 565-603.

Bourne, P. E., and J. McEntyre, 2006 Biocurators: Contributors to the world of science. Plos Computational Biology 2**:** 1185-1185.

Burge, S., T. K. Attwood, A. Bateman, T. Z. Berardini, M. Cherry *et al.*, 2012 Biocurators and Biocuration: surveying the 21st century challenges. Database-the Journal of Biological Databases and Curation.

Burkhardt, K., B. Schneider and J. Ory, 2006 A biocurator perspective: Annotation at the research collaboratory for structural bioinformatics protein data bank. Plos Computational Biology 2**:** 1186-1189.

Candan, K. S., J. Grant and V. S. Subrahmanian, 1997 A unified treatment of null values using constraints. Information Sciences 98**:** 99-156.

Carthey, J., M. R. de Leval and J. T. Reason, 2001 The human factor in cardiac surgery: errors and near misses in a high technology medical domain. Ann Thorac Surg 72**:** 300-305.

Cooper, K. D. T., Linda, 2012 *Engineering A Compiler*. Morgan Kaufmann, Burlington, MA.

de Tre, G., R. de Caluwe and H. Prade, 2004 Null values revisited in prospect of data integration. Semantics of a Networked World: Semantics for Grid Databases 3226**:** 79-90.

Donoho, D. M., Arian; Rahman, Inam; Shahram, Morteza; Stodden, Victoria, 2009 15 Years of Reproducible Research in Computational Harmonic Analysis. Computing in Science and Engineering 2009**:** 8-18.

Dopfer, D., W. Buist, Y. Soyer, M. A. Munoz, R. N. Zadoks *et al.*, 2008 Assessing genetic heterogeneity within bacterial species isolated from gastrointestinal and environmental samples: How many isolates does it take? Applied and Environmental Microbiology 74**:** 3490-3496.

Drake, V. J., B. Frei and L. M. Bouley, 2008 An error in the US Department of Agriculture nutrient database results in vitamin A values that are 6 times too high. Am J Clin Nutr 87**:** 1067-1068; author reply 1068-1069.

Ehlert, K., and L. Loewe, 2014 Lazy Updating of hubs can enable more realistic models by speeding up stochastic simulations. J Chem Phys 141**:** 204109.

Ember, C., R. Hanisch, G. Alter, H. Berman, M. Hedstrom *et al.*, 2013 Sustaining Domain Repositories for Digital Data: A White Paper, pp. http://datacommunity.icpsr.umich.edu/sites/default/files/WhitePaper_ICPSR_SDRDD_121113.pdf in *Workshop on Sustaining Domain Repositories for Digital Data, June 24-25, 2013*, ICPSR, University of Michigan, Alfred P. Sloan Foundation.

Freedman, L. P., I. M. Cockburn and T. S. Simcoe, 2015 The economics of reproducibility in preclinical research. PLoS Biol 13**:** e1002165.

Galletta, D. F., D. Abraham, M. El Louadi, W. Lekse, Y. A. Pollailis *et al.*, 1993 An Empirical Study of Spreadsheet Error Performance. Journal of Accounting, Managment, and Information Technology 3**:** 79-95.

Galletta, D. F., K. S. Hartzel, S. Johnson and J. L. Joseph, 1997 Spreadsheet Presentation and Error Detection: An Experimental Journal. Journal of Management Information Systems 13**:** 45-63.

Gitelman, L., 2013 *"Raw data" is an oxymoron*. The MIT Press, Cambridge, Massachusetts ; London, England.

Gödel, K., 1931 Über formal unentscheidbare Sätze der Principia Mathematica und verwandter Systeme, I. Monatshefte für Mathematik und Physik 38**:** 173–198.

Goecks, J., A. Nekrutenko, J. Taylor and T. Galaxy, 2010 Galaxy: a comprehensive approach for supporting accessible, reproducible, and transparent computational research in the life sciences. Genome Biol 11**:** R86.

Goldston, D., 2008 Big data: Data wrangling. Nature 455**:** 15-15.

Grimes, M. L., W. J. Lee, L. Van der Maaten and P. Shannon, 2013 Wrangling Phosphoproteomic Data to Elucidate Cancer Signaling Pathways. Plos One 8.

Grove, S. J., and N. E. Stork, 2000 An inordinate fondness for beetles. Invertebrate Taxonomy 14**:** 733-739.

Grune, D., and C. J. H. Jacobs, 2008 *Parsing techniques : a practical guide*. Springer, New York.

Grune, D. e. a., 2012 *Modern Compiler Design*. Springer Science+Business Media, New York, NY.

Hamilton, A. J., Y. Basset, K. K. Benke, P. S. Grimbacher, S. E. Miller *et al.*, 2010 Quantifying Uncertainty in Estimation of Tropical Arthropod Species Richness. American Naturalist 176**:** 90-95.

Hartmann, S., and S. Link, 2012 The Implication Problem of Data Dependencies over SQL Table Definitions: Axiomatic, Algorithmic and Logical Characterizations. Acm Transactions on Database Systems 37.

Hayden, E. C., 2014 Technology: The $1,000 genome. Nature 507**:** 294-295.

Henry, P. L., 2013 "Ultimately, the source code is the documentation for developers" quoted in: "Lessons from the Agile trenches: The ups and downs of Agile development". ibm.com/developerWorks**:** http://www.ibm.com/developerworks/library/a-agiletrenches/a-agiletrenches-pdf.pdf.

Hernich, A., L. Libkin and N. Schweikardt, 2011 Closed World Data Exchange. Acm Transactions on Database Systems 36.

Hooey, B. L., M. Aurisicchio, R. Bracewell and D. C. Foyle, 2014 Evidence-Based Error Analysis: Supporting the Design of Error-Tolerant Systems. Human-Computer Interaction: Applications and Services, Pt Iii 8512**:** 401-412.

Howe, D., M. Costanzo, P. Fey, T. Gojobori, L. Hannick *et al.*, 2008 Big data: The future of biocuration. Nature 455**:** 47-50.

Hutchins, L. N., Y. Ding, J. P. Szatkiewicz, R. Von Smith, H. Yang *et al.*, 2010 CGDSNPdb: a database resource for error-checked and imputed mouse SNPs. Database (Oxford) 2010**:** baq008.

James, D., N. Wilkins-Diehr, V. Stodden, D. Colbry, C. Rosales *et al.*, 2015 Standing Together for Reproducibility in Large-Scale Computing: Report on reproducibility@XSEDE, pp. http://arxiv.org/abs/1412.5557 in *Arxiv: Distributed, Parallel, and Cluster Computing (cs.DC)*, arXiv:1412.5557v2 [cs.DC].

Jones, C. E., A. L. Brown and U. Baumann, 2007 Estimating the annotation error rate of curated GO database sequence annotations. BMC Bioinformatics 8**:** 170.

Jurney, R., 2013 *Agile data science*. O'Reilly Media, Beijing ; Sebastopol, CA.

Karr, J. R., J. C. Sanghvi, D. N. Macklin, M. V. Gutschow, J. M. Jacobs *et al.*, 2012 A whole-cell computational model predicts phenotype from genotype. Cell 150**:** 389-401.

Karr, J. R., K. Takahashi and A. Funahashi, 2015a The principles of whole-cell modeling. Curr Opin Microbiol 27**:** 18-24.

Karr, J. R., A. H. Williams, J. D. Zucker, A. Raue, B. Steiert *et al.*, 2015b Summary of the DREAM8 Parameter Estimation Challenge: Toward Parameter Identification for Whole-Cell Models. PLoS Comput Biol 11**:** e1004096.

Kenall, A., S. Edmunds, L. Goodman, L. Bal, L. Flintoft *et al.*, 2015 Better reporting for better research: a checklist for reproducibility. Gigascience 4**:** 32.

Knuth, D. E., and S. Levy, 1994 *The CWEB system of structured documentation : version 3.0*. Addison-Wesley Pub. Co., Reading, Mass.

Lander, E. S., L. M. Linton, B. Birren, C. Nusbaum, M. C. Zody *et al.*, 2001 Initial sequencing and analysis of the human genome. Nature 409**:** 860-921.

Lee, Y., and J. Quackenbush, 2003 Using the TIGR gene index databases for biological discovery. Curr Protoc Bioinformatics Chapter 1**:** Unit 1 6.

Lee, Y., J. Tsai, S. Sunkara, S. Karamycheva, G. Pertea *et al.*, 2005 The TIGR Gene Indices: clustering and assembling EST and known genes and integration with eukaryotic genomes. Nucleic Acids Res 33**:** D71-74.

Lesgold, A., H. Rubinson, P. Feltovich, R. Glaser, D. Klopfer *et al.*, 1988 Expertise in a Complex Skill: Diagnosing X-Ray Pictures. in *The Nature of Expertise.*, edited by R. G. Micheline Chi and M. Farr. Lawrence Erlbaum, Hillsdale, NJ.

Lifschitz, V., K. Pichotta and F. K. Yang, 2012 Relational theories with null values and non-herbrand stable models. Theory and Practice of Logic Programming 12**:** 565-582.

Loewe, L., 2016 Systems in Evolutionary System Biology, pp. pp. 297-318 (http://evolutionarysystemsbiology.org/pdf/Loewe-2016-evosysbio.pdf) in *Encyclopedia of Evolutionary Biology*, edited by Richard M. Kliman (Chief) and Hiroshi Akashi (Section). Academic Press (Elsevier), Oxford.

Loewe, L., and S. Keel, 2014 BEST Names for Semantic Units to Support Reproducible Modeling, pp. 4 pages, https://www.xsede.org/documents/659353/703287/xsede659314_loewe.pdf in *2014-07-14 Workshop: "reproducibility@XSEDE", XSEDE14 Annual Conference (* https://www.xsede.org/web/reproducibility *)*, Atlanta, GA, USA. .

Loewe, L., K. S. Scheuer and S. Keel, 2016a Evolvix Concept: BEST Names for Semantic Units Increase Semantic Reproducibility and Help Combine Complex Biological Models, pp. https://sites.google.com/site/dslabhs2016/proceeding-papers in *Workshop on Data Science, Learning, and Applications to Biomedical & Health Sciences, January 7 - 8, 2016*. NY Academy of Sciences, New York Academy of Sciences.

Loewe, L., K. S. Scheuer, S. A. Keel, V. Vyas, B. Liblit *et al.*, 2016b Evolvix BEST Names for semantic reproducibility across code2brain interfaces. Annals of the New York Academy of Sciences**:** http://dx.doi.org/10.1111/nyas.13192 ; http://evolvix.org/naming ; http://evolvix.org/post.

Magnussen, S., B. Smith, C. Kleinn and I. F. Sun, 2010 An urn model for species richness estimation in quadrat sampling from fixed-area populations. Forestry 83**:** 293-306.

Mahmood, Z., 2016 *Data science and big data computing : frameworks and methodologies*. Springer Berlin Heidelberg, New York, NY.

Martinez, M. V., C. Molinaro, J. Grant and V. S. Subrahmanian, 2013 Customized Policies for Handling Partial Information in Relational Databases. Ieee Transactions on Knowledge and Data Engineering 25**:** 1254-1271.

McCallum, Q. E., 2013 *Bad Data Handbook: Mapping the world of data problems*. O'Reilly, Sebastopol, CA.

Mirza, G. A., 2015 Null Value Conflict: Formal Definition and Resolution. 2015 13th International Conference on Frontiers of Information Technology (Fit)**:** 132-137.

NASA, F. Chandler, T., Y. H. Chang, James, A. Mosleh, J. Marble, L. *et al.*, 2006-07 *Human Reliability Analysis Methods Selection Guidance for NASA*. NASA/OSMA, NASA, Washington, DC.

NASA, K. Leiden, K. R. Laughery, J. Keller, J. French *et al.*, 2001-09-30 *A Review of Human Performance Models for the Prediction of Human Error*. National Aeronautics and Space Administration, System-Wide Accident Prevention Program, Ames Research Center, Moffett Field, CA, http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.468.6969&rep=rep1&type=pdf.

NASA, M. Stamatelatos and H. Dezfuli, 2011 *Probabilistic Risk Assessment Procedures Guide for NASA Managers and Practitioners (2nd Edition)*, NASA Headquarters, Washington, DC.

NIH, D. DeMets, L. Tabak, R. Altman, D. Botstein *et al.*, 2012 Data and Informatics Working Group Report to the Advisory Committee to the Director, pp. National Institutes of Health,, http://acd.od.nih.gov/Data and Informatics Working Group Report.PDF.

Orchard, S., and H. Hermjakob, 2015 Shared resources, shared costs--leveraging biocuration resources. Database (Oxford) 2015.

Panko, R. R., 1998 Applying code inspection to spreadsheet testing. Effective Utilization and Management of Emerging Information Technologies**:** 358-362.

Panko, R. R., 2016 Human Error Website. Accessed 2016-10-31**:** http://panko.shidler.hawaii.edu/HumanErr/Index.htm

Panko, R. R., and S. Aurigemma, 2010 Revising the Panko-Halverson taxonomy of spreadsheet errors. Decision Support Systems 49**:** 235-244.

Panko, R. R., and R. H. Sprague, 1998 Hitting the wall: errors in developing and code inspecting a 'simple' spreadsheet model. Decision Support Systems 22**:** 337-353.

Parsons, T. W., 1992 *Introduction to Compiler Construction*. Computer Science Press, New York.

Pertea, G., X. Huang, F. Liang, V. Antonescu, R. Sultana *et al.*, 2003 TIGR Gene Indices clustering tools (TGICL): a software system for fast clustering of large EST datasets. Bioinformatics 19**:** 651-652.

Poldrack, R. A., and J. B. Poline, 2015 The publication and reproducibility challenges of shared data. Trends Cogn Sci 19**:** 59-61.

Quackenbush, J., F. Liang, I. Holt, G. Pertea and J. Upton, 2000 The TIGR gene indices: reconstruction and representation of expressed gene sequences. Nucleic Acids Res 28**:** 141-145.

Rak, R., R. T. Batista-Navarro, A. Rowley, J. Carter and S. Ananiadou, 2014 Text-mining-assisted biocuration workflows in Argo. Database (Oxford) 2014.

Razali, N., J. Noble and S. Marshall, 2015-10-26 Operators and Precedence in Programming Languages. PLATEAU'15**:** 53-56 (http://dx.doi.org/10.1145/2846680.2846690).

Reason, J., 2005 Safety in the operating theatre - Part 2: human error and organisational failure. Qual Saf Health Care 14**:** 56-60.

Reason, J. T., 1990 *Human error*. Cambridge University Press, Cambridge England ; New York.

Reason, J. T., 2013 *A life in error : from little slips to big disasters*. Ashgate, Surrey, UK England.

Reason, J. T., 2015 *Organizational accidents revisited*. Ashgate, Burlington, VT.

Reason, J. T., and A. Hobbs, 2003 *Managing maintenance error : a practical guide*. Ashgate, Aldershot, Hampshire, England ; Burlington, VT, USA.

Reason, J. T., and K. Mycielska, 1982 *Absent-minded? : the psychology of mental lapses and everyday errors*. Prentice-Hall, Englewood Cliffs, N.J.

Reiser, L., T. Z. Berardini, D. Li, R. Muller, E. M. Strait *et al.*, 2016 Sustainable funding for biocuration: The Arabidopsis Information Resource (TAIR) as a case study of a subscription-based funding model. Database (Oxford) 2016.

Rodriguez-Esteban, R., 2015 Biocuration with insufficient resources and fixed timelines. Database (Oxford) 2015.

Salimi, N., and R. Vita, 2006 The biocurator: Connecting and enhancing scientific data. Plos Computational Biology 2**:** 1190-1192.

Sanger, F., G. M. Air, B. G. Barrell, N. L. Brown, A. R. Coulson *et al.*, 1977a Nucliotide sequence of bacteriophage phi X174 DNA. Nature 265**:** 687-695.

Sanger, F., S. Nicklen and A. R. Coulson, 1977b DNA sequencing with chain-terminating inhibitors. Proceedings of the National Academy of Sciences of the U.S.A. 74**:** 5463-5467.

Sheridan, C., 2014 Illumina claims $1,000 genome win. Nat Biotechnol 32**:** 115.

Singhal, A., R. Leaman, N. Catlett, T. Lemberger, J. McEntyre *et al.*, 2016 Pressing needs of biomedical text mining in biocuration and beyond: opportunities and challenges. Database (Oxford) 2016.

Smith, N. J. J., 2008 *Vagueness and Degrees of Truth*. Oxford University Press, Oxford, Great Britain.

Stodden, V., 2015 Reproducing Statistical Results. Annu. Rev. Stat. Appl. 2**:** 1-19.

Stodden, V., F. Leisch and R. D. Peng (Editors), 2014 *Implementing Reproducible Research*. CRC Press, Boca Raton, FL.

Telenti, A., L. C. Pierce, W. H. Biggs, J. di Iulio, E. H. Wong *et al.*, 2016 Deep sequencing of 10,000 human genomes. Proc Natl Acad Sci U S A 113**:** 11901-11906.

Thalheim, B., and K. D. Schewe, 2011 NULL 'Value' Algebras and Logics. Information Modelling and Knowledge Bases Xxii 225**:** 354-367.

Venter, J. C., M. D. Adams, E. W. Myers, P. W. Li, R. J. Mural *et al.*, 2001 The sequence of the human genome. Science 291**:** 1304-1351.

Waraporn, N., and K. Porkaew, 2008 Semantics of null values in subqueries. Imecs 2008: International Multiconference of Engineers and Computer Scientists, Vols I and Ii**:** 400-405.

Wei, C. H., H. Y. Kao and Z. Lu, 2013 PubTator: a web-based text mining tool for assisting biocuration. Nucleic Acids Res 41**:** W518-522.

Weil, G., C. Motamed, A. Eghiaian, M. L. Guye and J. L. Bourgain, 2015 The use of a clinical database in an anesthesia unit: focus on its limits. J Clin Monit Comput 29**:** 163-167.

Zaniolo, C., 1984 Database Relations with Null Values. Journal of Computer and System Sciences 28**:** 142-166.

Zhang, H. M., and H. Stern, 2009 Sample Size Calculation for Finding Unseen Species. Bayesian Analysis 4**:** 763-791.

LITERATURE CITED

2016 *Advances in mathematical modeling, optimization and optimal control*. Springer Science+Business Media, New York, NY.

Aarts, A. A., J. E. Anderson, C. J. Anderson, P. R. Attridge, A. Attwood *et al.*, 2015 Estimating the reproducibility of psychological science. Science 349.

Abruzzi, K., X. Chen, E. Nagoshi, A. Zadina and M. Rosbash, 2015 RNA-seq profiling of small numbers of Drosophila neurons. Meth Enzymol 551**:** 369-386.

Abruzzi, K. C., J. Rodriguez, J. S. Menet, J. Desrochers, A. Zadina *et al.*, 2011 Drosophila CLOCK target gene characterization: implications for circadian tissue-specific gene expression. Genes & development 25**:** 2374-2386.

Ades, A. E., N. J. Welton, D. Caldwell, M. Price, A. Goubar *et al.*, 2008 Multiparameter evidence synthesis in epidemiology and medical decision-making. J Health Serv Res Policy 13 Suppl 3**:** 12-22.

Adewoye, A. B., C. P. Kyriacou and E. Tauber, 2015 Identification and functional analysis of early gene expression induced by circadian light-resetting in Drosophila. BMC Genomics 16**:** 570.

Agostinelli, F., N. Ceglia, B. Shahbaba, P. Sassone-Corsi and P. Baldi, 2016 What time is it? Deep learning approaches for circadian rhythms. Bioinformatics 32**:** i8-i17.

Aimo, L., R. Liechti, N. Hyka-Nouspikel, A. Niknejad, A. Gleizes *et al.*, 2015 The SwissLipids knowledgebase for lipid biology. Bioinformatics 31**:** 2860-2866.

Aird, D., M. G. Ross, W. S. Chen, M. Danielsson, T. Fennell *et al.*, 2011 Analyzing and minimizing PCR amplification bias in Illumina sequencing libraries. Genome Biol 12**:** R18.

Akman, O., M. L. Guerriero, L. Loewe and C. Troein, 2010 Complementary approaches to understanding the plant circadian clock. Proceedings 3rd Workshop "From Biology To Concurrency and back" - FBTC 2010 published in Electronic Proceedings in Theoretical Computer Science 19**:** 1-19 (http://arxiv.org/abs/1002.4661v1001).

Akten, B., E. Jauch, G. K. Genova, E. Y. Kim, I. Edery *et al.*, 2003 A role for CK2 in the Drosophila circadian oscillator. Nature neuroscience 6**:** 251-257.

Allada, R., N. E. White, W. V. So, J. C. Hall and M. Rosbash, 1998 A mutant Drosophila homolog of mammalian Clock disrupts circadian rhythms and transcription of period and timeless. Cell 93**:** 791-804.

Allison, D. B., A. W. Brown, B. J. George and K. A. Kaiser, 2016 Reproducibility: A tragedy of errors. Nature 530**:** 27-29.

Altman, D. G., S. M. Gore, M. J. Gardner and S. J. Pocock, 1983 Statistical guidelines for contributors to medical journals. Br Med J (Clin Res Ed) 286**:** 1489-1493.

Amos, W., C. M. Hutter, M. D. Schug and C. F. Aquadro, 2003 Directional evolution of size coupled with ascertainment bias for variation in Drosophila microsatellites. Mol Biol Evol 20**:** 660-662.

Anderson, D. F., 2007 A modified next reaction method for simulating chemical systems with time dependent propensities and delays. J Chem Phys 127**:** 214107.

Anderson, D. F., A. Ganguly and T. G. Kurtz, 2011 Error Analysis of Tau-Leap Simulation Methods. Annals of Applied Probability 21**:** 2226-2262.

Annosi, M. C., M. Magnusson, A. Martini and F. P. Appio, 2016 Social Conduct, Learning and Innovation: An Abductive Study of the Dark Side of Agile Software Development. Creativity and Innovation Management 25**:** 515-535.

Arp, R., B. Smith and A. D. Spear, 2015 *Building ontologies with Basic Formal Ontology*. Massachusetts Institute of Technology, Cambridge, Massachusetts http://ontology.buffalo.edu/smith/.

Ascher, U. M., and L. R. Petzold, 1998 *Computer Methods for Ordinary Differential Equations and Differential-Algebraic Equations*. SIAM, Philadelphia.

Ashburner, M., and R. Drysdale, 1994 FlyBase--the Drosophila genetic database. Development 120**:** 2077-2079.

Ashley, E. A., 2016 Towards precision medicine. Nat Rev Genet 17**:** 507-522.

Bae, K., and I. Edery, 2006 Regulating a Circadian Clock's Period, Phase and Amplitude by Phosphorylation: Insights from Drosophila. Journal of Biochemistry 140**:** 609-617.

Bae, K., C. Lee, P. Hardin and I. Edery, 2000 dCLOCK is present in limiting amounts and likely mediates daily interactions between the dCLOCK-CYC transcription factor and the PER-TIM complex. Journal of Neuroscience 20**:** 1746-1753.

Bae, K., C. Lee, D. Sidote, K. Y. Chuang and I. Edery, 1998 Circadian regulation of a Drosophila homolog of the mammalian Clock gene: PER and TIM function as positive regulators. Mol Cell Biol 18**:** 6142-6151.

Bagheri, N., M. J. Lawson, J. Stelling and F. J. Doyle, 2008a Modeling the Drosophila melanogaster Circadian Oscillator via Phase Optimization. Journal of biological rhythms 23**:** 525-537.

Bagheri, N., J. Stelling and F. J. Doyle, 2007 Circadian phase entrainment via nonlinear model predictive control. International Journal of Robust and Nonlinear Control 17**:** 1555-1571.

Bagheri, N., S. R. Taylor, K. Meeker, L. R. Petzold and F. J. Doyle, 2008b Synchrony and entrainment properties of robust circadian oscillators., pp. S17-28 in *J R Soc Interface*.

Baker, M., 2016 1,500 scientists lift the lid on reproducibility. Nature 533**:** 452-454.

Bammler, T., R. P. Beyer, S. Bhattacharya, G. A. Boorman, A. Boyles *et al.*, 2005 Standardizing global gene expression analysis between laboratories and across platforms. Nat Methods 2**:** 351-356.

Banko, Z., and J. Abonyi, 2015 Mixed dissimilarity measure for piecewise linear approximation based time series applications. Expert Systems with Applications 42**:** 7664-7675.

Bao, S., J. Rihel, E. Bjes, J. Y. Fan and J. L. Price, 2001 The Drosophila double-timeS mutation delays the nuclear accumulation of period protein and affects the feedback regulation of period mRNA. Journal of Neuroscience 21**:** 7117-7126.

Barba, L. A., 2016 The hard road to reproducibility. Science 354**:** 142-142.

Barkai, N., and S. Leibler, 2000 Circadian clocks limited by noise. Nature 403**:** 267-268.

Barrett, T., T. O. Suzek, D. B. Troup, S. E. Wilhite, W. C. Ngau *et al.*, 2005 NCBI GEO: mining millions of expression profiles--database and tools. Nucleic Acids Res 33**:** D562-566.

Barrett, T., D. B. Troup, S. E. Wilhite, P. Ledoux, C. Evangelista *et al.*, 2011 NCBI GEO: archive for functional genomics data sets--10 years on. Nucleic Acids Res 39**:** D1005-1010.

Barrett, T., S. E. Wilhite, P. Ledoux, C. Evangelista, I. F. Kim *et al.*, 2013 NCBI GEO: archive for functional genomics data sets--update. Nucleic Acids Res 41**:** D991-995.

Bartling, S., and S. Friesike, 2014 *Opening science*. Springer, New York, http://book.openingscience.org/

Bateman, A., 2010 Curators of the world unite: the International Society of Biocuration. Bioinformatics 26**:** 991.

Batista, G. E. A. P. A., X. Wang and E. J. Keogh, 2011 A Complexity-Invariant Distance Measure for Time Series.

Batista, R. T., D. B. Ramirez, R. D. Santos, M. C. del Rosario and E. R. Mendoza, 2007 EUCLIS--an information system for circadian systems biology. IET Syst Biol 1**:** 266-273.

Bauer, P., A. Thorpe and G. Brunet, 2015 The quiet revolution of numerical weather prediction. Nature 525**:** 47-55.

Baylies, M. K., L. B. Vosshall, A. Sehgal and M. W. Young, 1992 New Short-Period Mutations of the Drosophila Clock Gene Per. Neuron 9**:** 575-581.

Beaver, L. M., B. O. Gvakharia, T. S. Vollintine, D. M. Hege, R. Stanewsky *et al.*, 2002 Loss of circadian clock function decreases reproductive fitness in males of Drosophila melanogaster., pp. 2134-2139 in *P Natl Acad Sci Usa*.

Beaver, L. M., L. A. Hooven, S. M. Butcher, N. Krishnan, K. A. Sherman *et al.*, 2010 Circadian clock regulates response to pesticides in Drosophila via conserved Pdp1 pathway., pp. 513-520 in *Toxicol. Sci.*

                 QQv1r1

Beaver, L. M., B. L. Rush, B. O. Gvakharia and J. M. Giebultowicz, 2003 Noncircadian regulation and function of clock genes period and timeless in oogenesis of Drosophila melanogaster., pp. 463-472 in *J Biol Rhythms*.

Beersma, D. G. M., 2005 Why and how do we model circadian rhythms? Journal of Biological Rhythms 20**:** 304-313.

Bell-Pedersen, D., V. M. Cassone, D. J. Earnest, S. S. Golden, P. E. Hardin *et al.*, 2005 Circadian rhythms from multiple oscillators: Lessons from diverse organisms. Nature Reviews Genetics 6**:** 544-556.

Berndt, A., T. Kottke, H. Breitkreuz, R. Dvorsky, S. Hennig *et al.*, 2007 A Novel Photoreaction Mechanism for the Circadian Blue Light Photoreceptor Drosophila Cryptochrome. Journal of Biological Chemistry 282**:** 13011-13021.

Bishop, C. M., 2006 *Pattern recognition and machine learning*. Springer, New York.

Blankenberg, D., J. E. Johnson, J. Taylor, A. Nekrutenko and G. Team, 2014 Wrangling Galaxy's reference data. Bioinformatics 30**:** 1917-1919.

Blau, J., 1999 Cycling vrille Expression Is Required for a Functional Drosophila Clock. Cell 99**:** 661-671.

Boos, D. D., and L. A. Stefanski, 2011 P-Value Precision and Reproducibility. American Statistician 65**:** 213-221.

Bourne, P. E., and J. McEntyre, 2006 Biocurators: Contributors to the world of science. Plos Computational Biology 2**:** 1185-1185.

Boyle, G., K. Richter, H. D. Priest, D. Traver, T. C. Mockler *et al.*, 2017 Comparative Analysis of Vertebrate Diurnal/Circadian Transcriptomes. PLoS One 12**:** e0169923.

Brandes, C., J. D. Plautz, R. Stanewsky, C. F. Jamison, M. Straume *et al.*, 1996 Novel features of drosophila period Transcription revealed by real-time luciferase reporting. Neuron 16**:** 687-692.

Brodland, G. W., 2015 How computational models can help unlock biological systems. Seminars in Cell & Developmental Biology 47-48.

Broman, K. W., 1999 Cleaning genotype data. Genet Epidemiol 17 Suppl 1**:** S79-83.

Broman, K. W., M. P. Keller, A. T. Broman, C. Kendziorski, B. S. Yandell *et al.*, 2015 Identification and Correction of Sample Mix-Ups in Expression Genetic Data: A Case Study. G3 (Bethesda) 5**:** 2177-2186.

Brooksbank, C., G. Cameron and J. Thornton, 2005 The European Bioinformatics Institute's data resources: towards systems biology. Nucleic Acids Res 33**:** D46-53.

Buck, S., 2015 Solving reproducibility. Science 348**:** 1403.

Bullard, J. H., E. Purdom, K. D. Hansen and S. Dudoit, 2010 Evaluation of statistical methods for normalization and differential expression in mRNA-Seq experiments. Bmc Bioinformatics 11.

Burge, S., T. K. Attwood, A. Bateman, T. Z. Berardini, M. Cherry *et al.*, 2012 Biocurators and Biocuration: surveying the 21st century challenges. Database-the Journal of Biological Databases and Curation.

Burkhardt, K., B. Schneider and J. Ory, 2006 A biocurator perspective: Annotation at the research collaboratory for structural bioinformatics protein data bank. Plos Computational Biology 2**:** 1186-1189.

Bussieck, M. R., and A. Meeraus, 2004 General algebraic modeling system (GAMS). Modeling Languages in Mathematical Optimization 88**:** 137-157.

Bustin, S. A., 2002 Quantification of mRNA using real-time reverse transcription PCR (RT-PCR): trends and problems. Journal of Molecular Endocrinology 29.

Bustin, S. A., and T. Nolan, 2004 Pitfalls of quantitative real-time reverse-transcription polymerase chain reaction. J Biomol Tech 15**:** 155-166.

Busza, A., M. Emery-Le, M. Rosbash and P. Emery, 2004 Roles of the two Drosophila CRYPTOCHROME structural domains in circadian photoreception. Science 304**:** 1503-1506.

Buzbas, E. O., and N. A. Rosenberg, 2015 AABC: approximate approximate Bayesian computation for inference in population-genetic models. Theor Popul Biol 99**:** 31-42.

Canales, B. K., 2016 Comment On: Prospective evaluation of urinary metabolic indices in severely obese adolescents after weight loss surgery. Surg Obes Relat Dis 12**:** 367-368.

Canales, R. D., Y. Luo, J. C. Willey, B. Austermiller, C. C. Barbacioru *et al.*, 2006 Evaluation of DNA microarray results with quantitative gene expression platforms. Nat Biotechnol 24**:** 1115-1122.

Candan, K. S., J. Grant and V. S. Subrahmanian, 1997 A unified treatment of null values using constraints. Information Sciences 98**:** 99-156.

Carling, K., 2000 Resistant outlier rules and the non-Gaussian case. Computational Statistics & Data Analysis 33**:** 249-258.

Carter, S. J., H. J. Durrington, J. E. Gibbs, J. Blaikley, A. S. Loudon *et al.*, 2016 A matter of time: study of circadian clocks and their role in inflammation. Journal of Leukocyte Biology 99**:** 549-560.

Carthey, J., M. R. de Leval, D. J. Wright, V. T. Farewell, J. T. Reason *et al.*, 2003 Behavioural markers of surgical excellence. Safety Science 41**:** 409-425.

Cassman, M., 2005 Barriers to progress in systems biology. Nature 438**:** 1079.

Cathala, G., J. F. Savouret, B. Mendez, B. L. West, M. Karin *et al.*, 1983 A Method for Isolation of Intact, Translationally Active Ribonucleic-Acid. DNA-a Journal of Molecular & Cellular Biology 2**:** 329-335.

Ceriani, M. F., T. K. Darlington, D. Staknis, P. Más, A. A. Petti *et al.*, 1999 Light-dependent sequestration of TIMELESS by CRYPTOCHROME. Science 285**:** 553-556.

Ceriani, M. F., J. B. Hogenesch, M. Yanovsky, S. Panda, M. Straume *et al.*, 2002 Genome-wide expression analysis in Drosophila reveals genes controlling circadian behavior. Journal of Neuroscience 22**:** 9305-9319.

Chelliah, V., N. Juty, I. Ajmera, R. Ali, M. Dumousseau *et al.*, 2015 BioModels: ten-year anniversary. Nucleic Acids Res 43**:** D542-548.

Chen, K. F., N. Peschel, R. Zavodska, H. Sehadova and R. Stanewsky, 2011 QUASIMODO, a Novel GPI-Anchored Zona Pellucida Protein Involved in Light Input to the Drosophila Circadian Clock. Current biology : CB 21**:** 719-729.

Chen, W.-F., J. Majercak and I. Edery, 2006 Clock-gated photic stimulation of timeless expression at cold temperatures and seasonal adaptation in Drosophila., pp. 256-271 in *J Biol Rhythms*.

Cheng, Y., B. Gvakharia and P. E. Hardin, 1998 Two alternatively spliced transcripts from the Drosophila period gene rescue rhythms having different molecular and behavioral characteristics., pp. 6505-6514 in *Mol Cell Biol*.

Chintapalli, V. R., M. Al Bratty, D. Korzekwa, D. G. Watson and J. A. Dow, 2013a Mapping an atlas of tissue-specific Drosophila melanogaster metabolomes by high resolution mass spectrometry. PLoS One 8**:** e78066.

Chintapalli, V. R., J. Wang and J. A. Dow, 2007 Using FlyAtlas to identify better Drosophila melanogaster models of human disease. Nat Genet 39**:** 715-720.

Chintapalli, V. R., J. Wang, P. Herzyk, S. A. Davies and J. A. Dow, 2013b Data-mining the FlyAtlas online resource to identify core functional motifs across transporting epithelia. BMC Genomics 14**:** 518.

Chiu, J. C., H. W. Ko and I. Edery, 2011 NEMO/NLK phosphorylates PERIOD to initiate a time-delay phosphorylation circuit that sets circadian clock speed., pp. 357-370 in *Cell*.

Chiu, J. C., J. T. Vanselow, A. Kramer and I. Edery, 2008 The phospho-occupancy of an atypical SLIMB-binding site on PERIOD that is phosphorylated by DOUBLETIME controls the pace of the clock. Genes & development 22**:** 1758-1772.

Chomczynski, P., and N. Sacchi, 1987 Single-Step Method of Rna Isolation by Acid Guanidinium Thiocyanate Phenol Chloroform Extraction. Analytical Biochemistry 162**:** 156-159.

Chouhan, N. S., R. Wolf, C. Helfrich-Forster and M. Heisenberg, 2015 Flies remember the time of day. Curr Biol 25**:** 1619-1624.

Chylek, L. A., L. A. Harris, J. R. Faeder and W. S. Hlavacek, 2015 Modeling for (physical) biologists: an introduction to the rule-based approach. Physical Biology 12.

Cirelli, C., 2009 The genetic and molecular regulation of sleep: from fruit flies to humans. Nature Reviews Neuroscience 10**:** 549-560.

Citri, Y., H. V. Colot, A. C. Jacquier, Q. Yu, J. C. Hall *et al.*, 1987 A family of unusually spliced biologically active transcripts encoded by a Drosophila clock gene., pp. 42-47 in *Nature*.

Claridge-Chang, A., H. Wijnen, F. Naef, C. Boothroyd, N. Rajewsky *et al.*, 2001 Circadian regulation of gene expression systems in the Drosophila head. Neuron 32**:** 657-671.

Clark, A. G., M. J. Hubisz, C. D. Bustamante, S. H. Williamson and R. Nielsen, 2005 Ascertainment bias in studies of human genome-wide polymorphism. Genome Res 15: 1496-1502.

Clough, E., and T. Barrett, 2016 The Gene Expression Omnibus Database. Methods Mol Biol 1418: 93-110.

Collins, F. S., and L. A. Tabak, 2014 Policy: NIH plans to enhance reproducibility. Nature 505: 612-613.

Coogan, A. N., A. L. Baird, A. Popa-Wagner and J. Thome, 2016 Circadian rhythms and attention deficit hyperactivity disorder: The what, the when and the why. Progress in Neuro-Psychopharmacology & Biological Psychiatry 67: 74-81.

Cooper, K. D. T., Linda, 2012 *Engineering A Compiler*. Morgan Kaufmann, Burlington, MA.

Costa, R., A. A. Peixoto, G. Barbujani and C. P. Kyriacou, 1992 A latitudinal cline in a Drosophila clock gene., pp. 43-49 in *Proc Biol Sci*.

Cotterman, C. W., 1983 Relationship and probability in Mendelian populations. Am J Med Genet 16: 393-440.

Coyne, J. A., and H. A. Orr, 2004 *Speciation*. Sinauer Associates, Sunderland, Mass.

Crosby, J. L., 1973 *Computer simulation in genetics*. Wiley, London, New York,.

Crow, J. F., 2001 Shannon's brief foray into genetics. Genetics 159: 915-917.

Csillery, K., M. G. Blum, O. E. Gaggiotti and O. Francois, 2010a Approximate Bayesian Computation (ABC) in practice. Trends Ecol Evol 25: 410-418.

Csillery, K., M. G. B. Blum, O. E. Gaggiotti and O. Francois, 2010b Approximate Bayesian Computation (ABC) in practice. Trends in Ecology & Evolution 25: 410-418.

Cumming, G., 2013 The New Statistics: A How-To Guide. Australian Psychologist 48: 161-170.

Cumming, G., 2014 The New Statistics: Why and How. Psychological Science 25: 7-29.

Curran-Everett, D., 2009 Explorations in statistics: confidence intervals. Advances in Physiology Education 33: 87-90.

Cusick, M. E., H. Yu, A. Smolyar, K. Venkatesan, A. R. Carvunis *et al.*, 2009 Literature-curated protein interaction datasets. Nat Methods 6: 39-46.

Cyran, S. A., A. M. Buchsbaum, K. L. Reddy, M.-C. Lin, N. R. J. Glossop *et al.*, 2003 vrille, Pdp1, and dClock form a second feedback loop in the Drosophila circadian clock. Cell 112: 329-341.

Daigle, B. J., Jr., M. K. Roh, L. R. Petzold and J. Niemi, 2012 Accelerated maximum likelihood parameter estimation for stochastic biochemical systems. BMC Bioinformatics 13: 68.

Darlington, T. K., K. Wager-Smith, M. F. Ceriani, D. Staknis, N. Gekakis *et al.*, 1998 Closing the circadian loop: CLOCK-induced transcription of its own inhibitors per and tim. Science 280: 1599-1603.

Dawkins, R., 1976 *The selfish gene*. Oxford University Press, Oxford.

Dawkins, R., 2016 *The selfish gene : 40th anniversary edition*. Oxford University Press, New York, NY.

de Tre, G., R. de Caluwe and H. Prade, 2004 Null values revisited in prospect of data integration. Semantics of a Networked World: Semantics for Grid Databases 3226: 79-90.

Debray, T. P., K. G. Moons, G. van Valkenhoef, O. Efthimiou, N. Hummel *et al.*, 2015 Get real in individual participant data (IPD) meta-analysis: a review of the methodology. Res Synth Methods 6: 293-309.

Demidenko, E., 2016 The p-Value You Can't Buy. American Statistician 70: 33-38.

Deng, Z., S. Arsenault, C. Caranica, J. Griffith, T. Zhu *et al.*, 2016 Synchronizing stochastic circadian oscillators in single cells of Neurospora crassa. Sci Rep 6: 35828.

Diambra, L., and C. P. Malta, 2012 Modeling the emergence of circadian rhythms in a clock neuron network. PLoS One 7: e33912.

Ding, H., G. Trajcevski, P. Scheuermann, X. Wang and E. J. Keogh, 2008 Querying and Mining of Time Series Data Experimental Comparison of Representations and Distance Measures.

Dissel, S., V. Codd, R. Fedic, K. J. Garner, R. Costa *et al.*, 2004 A constitutively active cryptochrome in Drosophila melanogaster. Nature neuroscience 7: 834-840.

DiStefano, J. J., 2013 *Dynamic systems biology modeling and simulation*. Elsevier, Academic Press, Amsterdam.

Doan, A., A. Halevy and Z. G. Ives, 2012 *Principles of data integration*. Morgan Kaufmann, Waltham, MA.

Dodd, A. N., N. Salathia, A. Hall, E. Kevei, R. Toth *et al.*, 2005 Plant circadian clocks increase photosynthesis, growth, survival, and competitive advantage. Science 309: 630-633.

Doherty, C. J., and S. A. Kay, 2010 Circadian control of global gene expression patterns., pp. 419-444 in *Annu Rev Genet*.

Doi, S. A., 2014 Evidence synthesis for medical decision making and the appropriate use of quality scores. Clin Med Res 12**:** 40-46.

Donoho, D. M., Arian; Rahman, Inam; Shahram, Morteza; Stodden, Victoria, 2009 15 Years of Reproducible Research in Computational Harmonic Analysis. Computing in Science and Engineering 2009**:** 8-18.

dos Santos, G., A. J. Schroeder, J. L. Goodman, V. B. Strelets, M. A. Crosby *et al.*, 2015 FlyBase: introduction of the Drosophila melanogaster Release 6 reference genome assembly and large-scale migration of genome annotations. Nucleic Acids Research 43**:** D690-D697.

Drager, A., and B. O. Palsson, 2014 Improving collaboration by standardization efforts in systems biology. Front Bioeng Biotechnol 2**:** 61.

Drummond, G. B., and S. L. Vowler, 2011 Show the data, don't conceal them. Journal of Physiology-London 589**:** 1861-1863.

Drummond, G. B., and S. L. Vowler, 2013 Statistical Reporting Guidelines. Type I: families, planning and errors. Exp Physiol 98**:** 3-6.

Dubruille, R. R., A. A. Murad, M. M. Rosbash and P. P. Emery, 2009 A constant light-genetic screen identifies KISMET as a regulator of circadian photoresponses. Audio, Transactions of the IRE Professional Group on 5**:** e1000787-e1000787.

Dumbell, R., O. Matveeva and H. Oster, 2016 Circadian Clocks, Stress, and Immunity. Frontiers in Endocrinology 7.

Edery, I., L. J. Zwiebel, M. E. Dembinska and M. Rosbash, 1994 Temporal phosphorylation of the Drosophila period protein., pp. 2260-2264 in *P Natl Acad Sci Usa*.

Edwards, A. W. F., 1992 *Likelihood*. John Hopkins University Press.

Efthimiou, O., T. P. Debray, G. van Valkenhoef, S. Trelle, K. Panayidou *et al.*, 2016 GetReal in network meta-analysis: a review of the methodology. Res Synth Methods 7**:** 236-263.

Ehlert, K., and L. Loewe, 2014 Lazy Updating of hubs can enable more realistic models by speeding up stochastic simulations. J Chem Phys 141**:** 204109.

Einstein, A., and A. Calaprice, 2011 *The ultimate quotable Einstein*. Princeton University Press, Princeton, N.J.

Ember, C., R. Hanisch, G. Alter, H. Berman, M. Hedstrom *et al.*, 2013 Sustaining Domain Repositories for Digital Data: A White Paper, pp. http://datacommunity.icpsr.umich.edu/sites/default/files/WhitePaper_ICPSR_SDRDD_121113.pdf in *Workshop on Sustaining Domain Repositories for Digital Data, June 24-25, 2013*, ICPSR, University of Michigan, Alfred P. Sloan Foundation.

Emery, P., W. V. So, M. Kaneko, J. C. Hall and M. Rosbash, 1998 CRY, a Drosophila clock and light-regulated cryptochrome, is a major contributor to circadian rhythm resetting and photosensitivity. Cell 95**:** 669-679.

Erguler, K., and M. P. Stumpf, 2011 Practical limits for reverse engineering of dynamical systems: a statistical analysis of sensitivity and parameter inferability in systems biology models. Mol Biosyst 7**:** 1593-1602.

Eyre-Walker, A., and P. D. Keightley, 2007 The distribution of fitness effects of new mutations. Nat Rev Genet 8**:** 610-618.

Fang, Y., S. Sathyanarayanan and A. Sehgal, 2007 Post-translational regulation of the Drosophila circadian clock requires protein phosphatase 1 (PP1). Genes & development 21**:** 1506-1518.

Fathallah-Shaykh, H. M., 2010 Dynamics of the Drosophila Circadian Clock: Theoretical Anti-Jitter Network and Controlled Chaos. Plos One 5.

Fathallah-Shaykh, H. M., J. L. Bona and S. Kadener, 2009 Mathematical Model of the Drosophila Circadian Clock: Loop Regulation and Transcriptional Integration. Biophysical Journal 97**:** 2399-2408.

Foll, M., M. A. Beaumont and O. Gaggiotti, 2008 An approximate Bayesian computation approach to overcome biases that arise when using amplified fragment length polymorphism markers to study population structure. Genetics 179**:** 927-939.

                                     QQv1r1

Forger, D., M. Drapeau, B. Collins and J. Blau, 2005 A new model for circadian clock research? Molecular systems biology 1**:** 2005.0014.

Freedman, L. P., I. M. Cockburn and T. S. Simcoe, 2015a The economics of reproducibility in preclinical research. PLoS Biol 13**:** e1002165.

Freedman, L. P., M. C. Gibson, S. P. Ethier, H. R. Soule, R. M. Neve *et al.*, 2015b Reproducibility: changing the policies and culture of cell line authentication. Nat Methods 12**:** 493-497.

Fryer, R. M., J. Randall, T. Yoshida, L. L. Hsiao, J. Blumenstock *et al.*, 2002 Global analysis of gene expression: Methods, interpretation, and pitfalls. Experimental Nephrology 10**:** 64-74.

Garbe, D. S., Y. Fang, X. Zheng, M. Sowcik, R. Anjum *et al.*, 2012 Cooperative interaction between phosphorylation sites on PERIOD maintains circadian period in Drosophila. PLoS Genetics 9**:** e1003749-e1003749.

Gekakis, N., L. Saez, A. M. Delahaye-Brown, M. P. Myers, A. Sehgal *et al.*, 1995 Isolation of timeless by PER protein interaction: defective interaction between timeless protein and long-period mutant PERL., pp. 811-815 in *Science*.

Gerard, C., D. Gonze and A. Goldbeter, 2009 Dependence of the period on the rate of protein degradation in minimal models for circadian oscillations. Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences 367**:** 4665-4683.

Gibson, R., B. Alako, C. Amid, A. Cerdeno-Tarraga, I. Cleland *et al.*, 2016 Biocuration of functional annotation at the European nucleotide archive. Nucleic Acids Res 44**:** D58-66.

Gillespie, D. T., 1977 Exact Stochastic Simulation of Coupled Chemical-Reactions. Abstracts of Papers of the American Chemical Society 173**:** 128-128.

Gillespie, D. T., 2007 Stochastic simulation of chemical kinetics. Annu Rev Phys Chem 58**:** 35-55.

Gillespie, D. T., 2008 Simulation Methods in Systems Biology. Formal Methods for Computational Systems Biology 5016**:** 125-167.

Gillespie, D. T., A. Hellander and L. R. Petzold, 2013 Perspective: Stochastic algorithms for chemical kinetics. Journal of Chemical Physics 138**:** 170901.

Gilliland, G., S. Perrin, K. Blanchard and H. F. Bunn, 1990 Analysis of Cytokine Messenger-Rna and DNA - Detection and Quantitation by Competitive Polymerase Chain-Reaction. Proceedings of the National Academy of Sciences of the United States of America 87**:** 2725-2729.

Gitelman, L., 2013 *"Raw data" is an oxymoron*. The MIT Press, Cambridge, Massachusetts ; London, England.

Glaser, F. T., and R. Stanewsky, 2005 Temperature synchronization of the Drosophila circadian clock. Current biology : CB 15**:** 1352-1363.

Gleeson, J. P., J. A. Ward, K. P. O'Sullivan and W. T. Lee, 2014 Competition-induced criticality in a model of meme popularity. Phys Rev Lett 112**:** 048701.

Glossop, N., J. Houl, H. Zheng, F. Ng, S. Dudek *et al.*, 2003 VRILLE feeds back to control circadian transcription of Clock in the Drosophila circadian oscillator. Neuron 37**:** 249-261.

Glynn, E. F., J. Chen and A. R. Mushegian, 2006 Detecting periodic patterns in unevenly spaced gene expression time series using Lomb-Scargle periodograms. Bioinformatics 22**:** 310-316.

Godwin, A. R., S. Kojima, C. B. Green and J. Wilusz, 2013 Kiss your tail goodbye: The role of PARN, Nocturnin, and Angel deadenylases in mRNA biology. Biochimica Et Biophysica Acta-Gene Regulatory Mechanisms 1829**:** 571-579.

Goldbeter, A., 1995 A Model for Circadian Oscillations in the Drosophila Period Protein (Per). Proceedings. Biological sciences / The Royal Society 261**:** 319-324.

Goldbeter, A., 2002 Computational approaches to cellular rhythms. Nature 420**:** 238-245.

Goldston, D., 2008 Big data: Data wrangling. Nature 455**:** 15-15.

Gonze, D., 2011 Modeling circadian clocks: From equations to oscillations. Central European Journal of Biology 6**:** 699-711.

Gonze, D., J. Halloy and A. Goldbeter, 2002a Deterministic versus stochastic models for circadian rhythms. Journal of Biological Physics 28**:** 637-653.

Gonze, D., J. Halloy and A. Goldbeter, 2002b Robustness of circadian rhythms with respect to molecular noise. Proceedings of the National Academy of Sciences of the United States of America 99**:** 673-678.

Gonze, D., J. Halloy and A. Goldbeter, 2004 Emergence of coherent oscillations in stochastic models for circadian rhythms, pp. 221-233 in *Physica a-Statistical Mechanics and Its Applications*.

Gonze, D., J. Halloy, J.-C. Leloup and A. Goldbeter, 2003 Stochastic models for circadian rhythms: effect of molecular noise on periodic and chaotic behaviour. Comptes Rendus Biologies 326**:** 189-203.

Goodwin, B. C., 1964 A statistical mechanics of temporal organization in cells. Symp Soc Exp Biol 18**:** 301-326.

Goodwin, B. C., 1965 Oscillatory behavior in enzymatic control processes. Adv Enzyme Regul 3**:** 425-438.

Gramates, L. S., S. J. Marygold, G. D. Santos, J. M. Urbano, G. Antonazzo *et al.*, 2016 FlyBase at 25: looking to the future. Nucleic Acids Res.

Gramates, L. S., S. J. Marygold, G. D. Santos, J. M. Urbano, G. Antonazzo *et al.*, 2017 FlyBase at 25: looking to the future. Nucleic Acids Res 45**:** D663-D671.

Green, C. B., N. Douris, S. Kojima, C. A. Strayer, J. Fogerty *et al.*, 2007 Loss of Nocturnin, a circadian deadenylase, confers resistance to hepatic steatosis and diet-induced obesity. Proceedings of the National Academy of Sciences of the United States of America 104**:** 9888-9893.

Grima, B., A. Dognon, A. Lamouroux, E. Chélot and F. Rouyer, 2012 CULLIN-3 Controls TIMELESS Oscillations in the Drosophila Circadian Clock. PLoS biology 10**:** e1001367.

Grima, B., A. Lamouroux, E. Chélot, C. Papin, B. Limbourg-Bouchon *et al.*, 2002 The F-box protein Slimb controls the levels of clock proteins Period and Timeless. Nature 420**:** 178-182.

Grimes, M. L., W. J. Lee, L. Van der Maaten and P. Shannon, 2013 Wrangling Phosphoproteomic Data to Elucidate Cancer Signaling Pathways. Plos One 8.

Grönke, S., I. Bickmeyer, R. Wunderlich, H. Jäckle and R. P. Kühnlein, 2009 Curled encodes the Drosophila homolog of the vertebrate circadian deadenylase Nocturnin., pp. 219-232 in *Genetics*.

Grove, S. J., and N. E. Stork, 2000 An inordinate fondness for beetles. Invertebrate Taxonomy 14**:** 733-739.

Grune, D. e. a., 2012 *Modern Compiler Design*. Springer Science+Business Media, New York, NY.

Haag-Liautard, C., M. Dorris, X. Maside, S. Macaskill, D. L. Halligan *et al.*, 2007 Direct estimation of per nucleotide and genomic deleterious mutation rates in Drosophila. Nature 445**:** 82-85.

Halsey, L. G., D. Curran-Everett, S. L. Vowler and G. B. Drummond, 2015 The fickle P value generates irreproducible results. Nature Methods 12**:** 179-185.

Hamblen, M. J. M., N. E. N. White, P. T. P. Emery, K. K. Kaiser and J. C. J. Hall, 1998 Molecular and behavioral analysis of four period mutants in Drosophila melanogaster encompassing extreme short, novel long, and unorthodox arrhythmic types. Genetics 149**:** 165-178.

Harbison, S. T., M. A. Carbone, J. F. Ayroles, E. A. Stone, R. F. Lyman *et al.*, 2009a Co-regulated transcriptional networks contribute to natural genetic variation in Drosophila sleep. Nat Genet 41**:** 371-375.

Harbison, S. T., T. F. Mackay and R. R. Anholt, 2009b Understanding the neurogenetics of sleep: progress from Drosophila. Trends Genet 25**:** 262-269.

Harbison, S. T., L. J. McCoy and T. F. Mackay, 2013 Genome-wide association study of sleep in Drosophila melanogaster. BMC Genomics 14**:** 281.

Hardin, P., and J. C. Hall, 1992 Circadian cycling in the levels of protein and mRNA from Drosophila melanogaster's period gene, pp. 155-169 in *Molecular Genetics of Biological Rhythms*, edited by M. W. Young.

Hardin, P. E., 1994 Analysis of period mRNA cycling in Drosophila head and body tissues indicates that body oscillators behave differently from head oscillators., pp. 7211-7218 in *Mol Cell Biol*.

Hardin, P. E., 2011 Molecular Genetic Analysis of Circadian Timekeeping in Drosophila, pp. 141-173. Elsevier.

Hardin, P. E., J. C. Hall and M. Rosbash, 1990 Feedback of the Drosophila period gene product on circadian cycling of its messenger RNA levels. Nature 343**:** 536-540.

Hardin, P. E., J. C. Hall and M. Rosbash, 1992 Circadian oscillations in period gene mRNA levels are transcriptionally regulated., pp. 11711-11715 in *P Natl Acad Sci Usa*.

Harms, E., S. Kivimae, M. W. Young and L. Saez, 2004 Posttranscriptional and posttranslational regulation of clock genes. Journal of Biological Rhythms 19**:** 361-373.

 QQv1r1

Hazber, M. A. G., R. X. Li, X. W. Gu, G. D. Xu and Y. H. Li, 2015 Semantic SPARQL query in a relational database based on ontology construction. 2015 11th International Conference on Semantics, Knowledge and Grids (Skg)**:** 25-32.

He, S., X. Zheng and D. Zeng, 2016 A Model-Free Scheme for Meme Ranking in Social Media. Decis Support Syst 81**:** 1-11.

Hermann-Luibl, C., T. Yoshii, P. R. Senthilan, H. Dircksen and C. Helfrich-Förster, 2014 The Ion Transport Peptide Is a New Functional Clock Neuropeptide in the Fruit Fly Drosophila melanogaster. Journal of Neuroscience 34**:** 9522-9536.

Heyde, C. C., 1997 *Quasi-likelihood and its application : a general approach to optimal parameter estimation*. Springer, New York.

Hirschman, L., G. A. Burns, M. Krallinger, C. Arighi, K. B. Cohen *et al.*, 2012 Text mining for the biocuration workflow. Database (Oxford) 2012**:** bas020.

Hoare, C. A. R., 2009 Null References: The Billion Dollar Mistake. Presentation at QCon 2009-08-25**:** https://www.infoq.com/presentations/Null-References-The-Billion-Dollar-Mistake-Tony-Hoare ; https://www.lucidchart.com/techblog/2015/2008/2031/the-worst-mistake-of-computer-science/ ; https://valbonne-consulting.com/computing-pioneers/tony-hoare-1934/

Hodson, R., 2016 Precision medicine. Nature 537**:** S49.

Hoffmann, A. A., R. Hallas, C. Sinclair and L. Partridge, 2001 Rapid loss of stress resistance in Drosophila melanogaster under adaptation to laboratory culture. Evolution 55**:** 436-438.

Hong, C. I., and J. J. Tyson, 1997 A proposal for temperature compensation of the circadian rhythm in Drosophila based on dimerization of the PER protein. Chronobiology International 14**:** 521-529.

Howe, D., M. Costanzo, P. Fey, T. Gojobori, L. Hannick *et al.*, 2008 Big data: The future of biocuration. Nature 455**:** 47-50.

Huang, E. P., X. F. Wang, K. R. Choudhury, L. M. McShane, M. Gonen *et al.*, 2015 Meta-analysis of the technical performance of an imaging procedure: guidelines and statistical methodology. Stat Methods Med Res 24**:** 141-174.

Huang, Y., and R. Gottardo, 2013 Comparability and reproducibility of biomedical data. Brief Bioinform 14**:** 391-401.

Hucka, M., A. Finney, H. M. Sauro, H. Bolouri, J. C. Doyle *et al.*, 2003 The systems biology markup language (SBML): a medium for representation and exchange of biochemical network models. Bioinformatics 19**:** 524-531.

Hughes, M. E. M., G. R. G. Grant, C. C. Paquin, J. J. Qian and M. N. M. Nitabach, 2012 Deep sequencing the circadian and diurnal transcriptome of Drosophila brain. Genes & development 22**:** 1266-1281.

Hurley, J. M., J. J. Loros and J. C. Dunlap, 2016 The circadian system as an organizer of metabolism. Fungal Genetics and Biology 90**:** 39-43.

Hut, R. A., S. Paolucci, R. Dor, C. P. Kyriacou and S. Daan, 2013 Latitudinal clines: an evolutionary view on biological rhythms. Proceedings of the Royal Society B-Biological Sciences 280.

Ince, D. C., L. Hatton and J. Graham-Cumming, 2012 The case for open computer programs. Nature 482**:** 485-488.

Ioannidis, J. P., 2005a Molecular bias. Eur J Epidemiol 20**:** 739-745.

Ioannidis, J. P., 2005b Why most published research findings are false. PLoS Med 2**:** e124.

Ioannidis, J. P., D. B. Allison, C. A. Ball, I. Coulibaly, X. Cui *et al.*, 2009a Repeatability of published microarray gene expression analyses. Nat Genet 41**:** 149-155.

Ioannidis, J. P., G. Thomas and M. J. Daly, 2009b Validating, augmenting and refining genome-wide association signals. Nat Rev Genet 10**:** 318-329.

Ishikawa, T., A. Matsumoto, T. Kato, S. Togashi, H. Ryo *et al.*, 1999 DCRY is a Drosophila photoreceptor protein implicated in light entrainment of circadian rhythm., pp. 57-65 in *Genes Cells*.

Itoh, T. Q., T. Tanimura and A. Matsumoto, 2011 Membrane-bound transporter controls the circadian transcription of clock genes in Drosophila., pp. 1159-1167 in *Genes Cells*.

James, D., N. Wilkins-Diehr, V. Stodden, D. Colbry, C. Rosales *et al.*, 2015 Standing Together for Reproducibility in Large-Scale Computing: Report on reproducibility@XSEDE, pp.

http://arxiv.org/abs/1412.5557 in *Arxiv: Distributed, Parallel, and Cluster Computing (cs.DC)*, arXiv:1412.5557v2 [cs.DC].

Janes, A. A., and G. Succi, 2012 The dark side of agile software development, pp. 215-228 in *Proceedings of the ACM international symposium on New ideas, new paradigms, and reflections on programming and software*. ACM, Tucson, Arizona, USA.

Jang, A. R., K. Moravcevic, L. Saez, M. W. Young and A. Sehgal, 2015 Drosophila TIM binds importin alpha1, and acts as an adapter to transport PER to the nucleus. PLoS Genet 11**:** e1004974.

Janssens, A. C., A. M. Gonzalez-Zuloeta Ladd, S. Lopez-Leon, J. P. Ioannidis, B. A. Oostra *et al.*, 2009 An empirical comparison of meta-analyses of published gene-disease associations versus consortium analyses. Genet Med 11**:** 153-162.

Jasny, B. R., G. Chin, L. Chong and S. Vignieri, 2011 Data replication & reproducibility. Again, and again, and again .... Introduction. Science 334**:** 1225.

Jaynes, E. T., and G. L. Bretthorst, 2003 *Probability theory : the logic of science*. Cambridge University Press, Cambridge, UK ; New York, NY.

Jin, L., 2011 A data-driven test to compare two or multiple time series. Computational Statistics & Data Analysis 55**:** 2183-2196.

Johnson, R. T., and K. Dickersin, 2007 Publication bias against negative results from clinical trials: three of the seven deadly sins. Nature Clinical Practice Neurology 3**:** 590-591.

Johnson, V. E., 2013 Revised standards for statistical evidence. Proceedings of the National Academy of Sciences of the United States of America 110**:** 19313-19317.

Jones, C. E., A. L. Brown and U. Baumann, 2007 Estimating the annotation error rate of curated GO database sequence annotations. BMC Bioinformatics 8**:** 170.

Jonquet, C., P. Lependu, S. Falconer, A. Coulet, N. F. Noy *et al.*, 2011 NCBO Resource Index: Ontology-Based Search and Mining of Biomedical Resources. Web Semant 9**:** 316-324.

Kadener, S., J. S. Menet, K. Sugino, M. D. Horwich, U. Weissbein *et al.*, 2009 A role for microRNAs in the Drosophila circadian clock. Genes Development 23**:** 2179-2191.

Kadener, S., D. Stoleru, M. McDonald, P. Nawathean and M. Rosbash, 2007 Clockwork Orange is a transcriptional repressor and a new Drosophila circadian pacemaker component. Genes & development 21**:** 1675-1686.

Kamp, P.-H., 2011 The most expensive one-byte mistake: Did Ken, Dennis, and Brian choose wrong with NUL-terminated text strings? ACM Queue - The Bikeshed 9**:** 1-4 (comments at http://queue.acm.org/detail.cfm?id=2010365).

Kannan, N. N., K. M. Vaze and V. K. Sharma, 2012 Clock accuracy and precision evolve as a consequence of selection for adult emergence in a narrow window of time in fruit flies Drosophila melanogaster., pp. 3527-3534 in *J Exp Biol*.

Karlebach, G., and R. Shamir, 2008 Modelling and analysis of gene regulatory networks. Nature Reviews Molecular Cell Biology 9**:** 770-780.

Karr, J. R., N. C. Phillips and M. W. Covert, 2014 WholeCellSimDB: a hybrid relational/HDF database for whole-cell model predictions. Database (Oxford) 2014.

Karr, J. R., J. C. Sanghvi, D. N. Macklin, A. Arora and M. W. Covert, 2013 WholeCellKB: model organism databases for comprehensive whole-cell models. Nucleic Acids Res 41**:** D787-792.

Karr, J. R., J. C. Sanghvi, D. N. Macklin, M. V. Gutschow, J. M. Jacobs *et al.*, 2012 A whole-cell computational model predicts phenotype from genotype. Cell 150**:** 389-401.

Karr, J. R., K. Takahashi and A. Funahashi, 2015a The principles of whole-cell modeling. Curr Opin Microbiol 27**:** 18-24.

Karr, J. R., A. H. Williams, J. D. Zucker, A. Raue, B. Steiert *et al.*, 2015b Summary of the DREAM8 Parameter Estimation Challenge: Toward Parameter Identification for Whole-Cell Models. PLoS Comput Biol 11**:** e1004096.

Katakura, Y., and H. Ohmori, 2006 Synchronized Control for Circadian Rhythm Based on the Central Clock Model of Drosophila. Audio, Transactions of the IRE Professional Group on**:** 2358-2361.

Keegan, K. P., S. Pradhan, J. P. Wang and R. Allada, 2007 Meta-analysis of Drosophila circadian microarray studies identifies a novel set of rhythmically expressed genes. PLoS Comput Biol 3**:** e208.

                         QQv1r1

Keightley, P. D., U. Trivedi, M. Thomson, F. Oliver, S. Kumar *et al.*, 2009 Analysis of the genome sequences of three Drosophila melanogaster spontaneous mutation accumulation lines. Genome Res 19**:** 1195-1201.

Khodor, Y. L., J. Rodriguez, K. C. Abruzzi, C.-H. A. Tang, M. T. Marr *et al.*, 2011 Nascent-seq indicates widespread cotranscriptional pre-mRNA splicing in Drosophila. Genes & development 25**:** 2502-2512.

Kidd, P. B., M. W. Young and E. D. Siggia, 2015 Temperature compensation and temperature sensation in the circadian clock. Proc Natl Acad Sci U S A 112**:** E6284-6292.

Killeen, P. R., 2005 An alternative to null-hypothesis significance tests. Psychol Sci 16**:** 345-353.

Kim, E. Y., K. Bae, F. S. Ng, N. R. J. Glossop, P. E. Hardin *et al.*, 2002 Drosophila CLOCK protein is under posttranscriptional control and influences light-induced activity., pp. 69-81 in *Neuron*.

Kim, E. Y., H. W. Ko, W. Yu, P. E. Hardin and I. Edery, 2007 A DOUBLETIME kinase binding domain on the Drosophila PERIOD protein is essential for its hyperphosphorylation, transcriptional repression, and circadian clock function. Molecular and cellular biology 27**:** 5014-5028.

Kim, S., R. I. Dogan, A. Chatr-Aryamontri, C. S. Chang, R. Oughtred *et al.*, 2016 BioCreative V BioC track overview: collaborative biocurator assistant task for BioGRID. Database-the Journal of Biological Databases and Curation.

Kim, Y. M., J. Choi, H. Y. Lee, G. W. Lee, Y. H. Lee *et al.*, 2014 dbCRY: a Web-based comparative and evolutionary genomics platform for blue-light receptors. Database (Oxford) 2014**:** bau037.

Kimura, M., 1962 On the probability of fixation of mutant genes in a population. Genetics 47**:** 713-719.

Kirk, P. D., A. C. Babtie and M. P. Stumpf, 2015 Systems biology (un)certainties. Science 350**:** 386-388.

Kivimäe, S., L. Saez and M. W. Young, 2008 Activating PER repressor through a DBT-directed phosphorylation switch. PLoS biology 6**:** e183.

Kloss, B., J. L. Price, L. Saez, J. Blau, A. Rothenfluh *et al.*, 1998 The Drosophila clock gene double-time encodes a protein closely related to human casein kinase Iepsilon. Cell 94**:** 97-107.

Kloss, B. B., A. A. Rothenfluh, M. W. M. Young and L. L. Saez, 2001 Phosphorylation of PERIOD Is Influenced by Cycling Physical Associations of DOUBLE-TIME, PERIOD, and TIMELESS in the Drosophila Clock. Neuron 30**:** 8-8.

Knowles, A., K. Koh, J.-T. Wu, C.-T. Chien, D. A. Chamovitz *et al.*, 2009 The COP9 Signalosome Is Required for Light-Dependent Timeless Degradation and Drosophila Clock Resetting. Journal of Neuroscience 29**:** 1152-1162.

Knuth, D. E., 1974 Structured programming with go to statements. Computing Surveys 6**:** 261-301.

Ko, H. W., J. Jiang and I. Edery, 2002 Role for Slimb in the degradation of Drosophila Period protein phosphorylated by Doubletime. Nature 420**:** 673-678.

Ko, H. W., E. Y. Kim, J. Chiu, J. T. Vanselow, A. Kramer *et al.*, 2010 A hierarchical phosphorylation cascade that regulates the timing of PERIOD nuclear entry reveals novel roles for proline-directed kinases and GSK-3beta/SGG in circadian clocks. Journal of Neuroscience 30**:** 12664-12675.

Koh, K., X. Zheng and A. Sehgal, 2006 JETLAG resets the Drosophila circadian clock by promoting light-induced degradation of TIMELESS. Science 312**:** 1809-1812.

Konopka, R. J., and S. Benzer, 1971 Clock mutants of Drosophila melanogaster., pp. 2112-2116 in *P Natl Acad Sci Usa*.

Koskinen, P., P. Toronen, J. Nokso-Koivisto and L. Holm, 2015 PANNZER: high-throughput functional annotation of uncharacterized proteins in an error-prone environment. Bioinformatics 31**:** 1544-1552.

Kotsifakos, A., V. Athitsos and P. Papapetrou, 2016 Query-sensitive distance measure selection for time series nearest neighbor classification. Intelligent Data Analysis 20**:** 5-27.

Kraft, P., E. Zeggini and J. P. Ioannidis, 2009 Replication in genome-wide association studies. Stat Sci 24**:** 561-573.

Krause, F., J. Uhlendorf, T. Lubitz, M. Schulz, E. Klipp *et al.*, 2010 Annotation and merging of SBML models with semanticSBML. Bioinformatics 26**:** 421-422.

Krishnamurthy, B., W. Willinger, P. Gill and M. Arlitt, 2011 A Socratic method for validation of measurement-based networking research. Computer Communications 34**:** 43-53.

Kuczenski, R. S., K. C. Hong, J. Garcia-Ojalvo and K. H. Lee, 2007 PERIOD-TIMELESS interval timer may require an additional feedback loop., pp. e154 in *PLoS Comput Biol*.

Kula-Eversole, E., E. Nagoshi, Y. Shang, J. Rodriguez, R. Allada *et al.*, 2010 Surprising gene expression patterns within and between PDF-containing circadian neurons in Drosophila. Proceedings of the National Academy of Sciences of the United States of America 107**:** 13497-13502.

Kulasiri, D., and Z. Xie, 2008 Modelling Circadian Rhythms in Drosophila and Investigation of VRI and PDP1 Feedback Loops Using a New Mathematical Model. Mathematical Modelling of Natural Phenomena 3**:** 1-26.

Kumar, S. S., D. D. Chen and A. A. Sehgal, 2012 Dopamine acts through Cryptochrome to promote acute arousal in Drosophila. Genes & development 26**:** 1224-1234.

Kurata, H., T. Tanaka and F. Ohnishi, 2007 Mathematical identification of critical reactions in the interlocked feedback model. PLoS ONE 2**:** e1103-1103.

Kurosawa, G., A. Mochizuki and Y. Iwasa, 2002 Comparative study of circadian clock models, in search of processes promoting oscillation. Journal of theoretical biology 216**:** 193-208.

Kurtz, T. G., 1972 Relationship between Stochastic and Deterministic Models for Chemical Reactions. Journal of Chemical Physics 57**:** 2976-&.

Kyriacou, C. P., A. A. Peixoto and R. Costa, 2007 A cline in the Drosophila melanogaster period gene in Australia: neither down nor under. Journal of Evolutionary Biology 20**:** 1649-1651.

Kyriacou, C. P., A. A. Peixoto, F. Sandrelli, R. Costa and E. Tauber, 2008 Clines in clock genes: fine-tuning circadian rhythms to the environment. Trends in Genetics 24**:** 124-132.

Lachance, J., and S. A. Tishkoff, 2013 SNP ascertainment bias in population genetic analyses: why it is important, and how to correct it. Bioessays 35**:** 780-786.

Laibe, C., and N. Le Novere, 2007 MIRIAM Resources: tools to generate and resolve robust cross-references in Systems Biology. BMC Syst Biol 1**:** 58.

Lamaze, A. A., A. A. Lamouroux, C. C. Vias, H.-C. H. Hung, F. F. Weber *et al.*, 2011 The E3 ubiquitin ligase CTRIP controls CLOCK levels and PERIOD oscillations in Drosophila. EMBO Reports 12**:** 549-557.

Lamba, P., D. Bilodeau-Wentworth, P. Emery and Y. Zhang, 2014 Morning and Evening Oscillators Cooperate to Reset Circadian Behavior in Response to Light Input. CellReports 7**:** 601-608.

Lander, E. S., L. M. Linton, B. Birren, C. Nusbaum, M. C. Zody *et al.*, 2001 Initial sequencing and analysis of the human genome. Nature 409**:** 860-921.

Law, A. M., and W. D. Kelton, 2000 *Simulation modeling and analysis*. McGraw-Hill, Boston.

Le Novere, N., 2015 Quantitative and logic modelling of molecular and gene networks. Nat Rev Genet 16**:** 146-158.

Le Novere, N., B. Bornstein, A. Broicher, M. Courtot, M. Donizelli *et al.*, 2006 BioModels Database: a free, centralized database of curated, published, quantitative kinetic models of biochemical and cellular systems. Nucleic Acids Res 34**:** D689-691.

Le Novere, N., M. Hucka, H. Mi, S. Moodie, F. Schreiber *et al.*, 2009 The Systems Biology Graphical Notation. Nature Biotechnology 27**:** 735-741.

Lear, B. C., L. Zhang and R. Allada, 2009 The neuropeptide PDF acts directly on evening pacemaker neurons to regulate multiple features of circadian behavior. PLoS biology 7**:** e1000154-e1000154.

Lebiedz, D., M. Rehberg and D. Skanda, 2012 Robust optimal design of synthetic biological networks. Methods Mol Biol 813**:** 45-55.

Lee, C., K. Bae and I. Edery, 1998 The Drosophila CLOCK protein undergoes daily rhythms in abundance, phosphorylation, and interactions with the PER-TIM complex. Neuron 21**:** 857-867.

Lee, C., K. Bae and I. Edery, 1999 PER and TIM inhibit the DNA binding activity of a Drosophila CLOCK-CYC/dBMAL1 heterodimer without disrupting formation of the heterodimer: a basis for circadian transcription., pp. 5316-5325 in *Mol Cell Biol*.

Lee, E., E. H. Jeong, H.-J. Jeong, E. Yildirim, J. T. Vanselow *et al.*, 2014a Phosphorylation of a Central Clock Transcription Factor Is Required for Thermal but Not Photic Entrainment. PLoS Genetics 10**:** e1004545.

Lee, R., J. R. Karr and M. W. Covert, 2013 WholeCellViz: data visualization for whole-cell models. BMC Bioinformatics 14**:** 253.

Lee, X. J., C. C. Drovandi and A. N. Pettitt, 2014b Model choice problems using approximate Bayesian computation with applications to pathogen transmission data sets. Biometrics.

Leek, J. T., and R. D. Peng, 2015 Statistics: P values are just the tip of the iceberg. Nature 520**:** 612.

Leise, T. L., and E. E. Moin, 2007 A mathematical model of the Drosophila circadian clock with emphasis on posttranslational mechanisms. Journal of theoretical biology 248**:** 48-63.

Lejaeghere, K., G. Bihlmayer, T. Bjorkman, P. Blaha, S. Blugel *et al.*, 2016 Reproducibility in density functional theory calculations of solids. Science 351**:** aad3000.

Leloup, J.-C., 2009 Circadian clocks and phosphorylation: Insights from computational modeling. Central European Journal of Biology 4**:** 290-303.

Leloup, J. C., and A. Goldbeter, 1997 Temperature compensation of circadian rhythms: Control of the period in a model for circadian oscillations of the per protein in Drosophila. Chronobiology International 14**:** 511-520.

Leloup, J. C., and A. Goldbeter, 1998a A model for circadian rhythms in Drosophila incorporating the formation of a complex between the PER and TIM proteins. Journal of biological rhythms 13**:** 70-87.

Leloup, J. C., and A. Goldbeter, 1998b Modeling circadian oscillations of the PER and TIM proteins in Drosophila. Biological Clocks 1152**:** 81-88.

Leloup, J. C., and A. Goldbeter, 2000 Modeling the molecular regulatory mechanism of circadian rhythms in Drosophila. Bioessays 22**:** 84-93.

Leloup, J. C., and A. Goldbeter, 2001 A molecular explanation for the long-term suppression of circadian rhythms by a single light pulse., pp. R1206-1212 in *Am J Physiol Regul Integr Comp Physiol*.

Leming, M. T., S. S. Rund, S. K. Behura, G. E. Duffield and J. E. O'Tousa, 2014 A database of circadian and diel rhythmic gene expression in the yellow fever mosquito Aedes aegypti. BMC Genomics 15**:** 1128.

Lerner, I., O. Bartok, V. Wolfson, J. S. Menet, U. Weissbein *et al.*, 2015 Clk post-transcriptional control denoises circadian transcription both temporally and spatially. Nat Commun 6**:** 7056.

Leroi, A. M., A. K. Chippindale and M. R. Rose, 1994 Long-term laboratory evolution of a genetic life-history trade-off in Drosophila melanogaster. 1. The role of genotype-by-environment interaction. Evolution 48**:** 1244-1257.

Leugering, G. n., 2012 *Constrained optimization and optimal control for partial differential equations*. Birkhäuser, Basel Switzerland ; New York.

Lew, M. J., 2012 Bad statistical practice in pharmacology (and other basic biomedical disciplines): you probably don't know P. British Journal of Pharmacology 166**:** 1559-1567.

Lewis, J., C. E. Breeze, J. Charlesworth, O. J. Maclaren and J. Cooper, 2016 Where next for the reproducibility agenda in computational biology? BMC Syst Biol 10**:** 52.

Li, Q., and X. Lang, 2008 Internal noise-sustained circadian rhythms in a Drosophila model., pp. 1983-1994 in *Biophys J*.

Li, S., K. Shui, Y. Zhang, Y. Lv, W. Deng *et al.*, 2017 CGDB: a database of circadian genes in eukaryotes. Nucleic Acids Res 45**:** D397-D403.

Li, Y., and M. Rosbash, 2013 Accelerated degradation of perS protein provides insight into light-mediated phase shifting., pp. 171-182 in *J Biol Rhythms*.

Lim, C., B. Y. Chung, J. L. Pitman, J. J. McGill, S. Pradhan *et al.*, 2007 clockwork orange encodes a transcriptional repressor important for circadian-clock amplitude in Drosophila. Current biology : CB 17**:** 1082-1089.

Lim, C., J. Lee, C. Choi, V. L. Kilman, J. Kim *et al.*, 2011 The novel gene twenty-four defines a critical translational step in the Drosophila clock., pp. 399-403 in *Nature*.

Lin, J.-M., V. L. Kilman, K. Keegan, B. Paddock, M. Emery-Le *et al.*, 2002a A role for casein kinase 2alpha in the Drosophila circadian clock. Nature 420**:** 816-820.

Lin, J.-M., A. Schroeder and R. Allada, 2005 In vivo circadian function of casein kinase 2 phosphorylation sites in Drosophila PERIOD. Journal of Neuroscience 25**:** 11175-11183.

Lin, Y., M. Han, B. Shimada, L. Wang, T. M. Gibler *et al.*, 2002b Influence of the period-dependent circadian clock on diurnal, circadian, and aperiodic gene expression in Drosophila melanogaster.

Proceedings of the National Academy of Sciences of the United States of America 99**:** 9562-9567.

Linné, C. v., 1758 *Caroli Linnæi ... Systema naturæ per regna tria naturæ, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis*. impensis L. Salvii, Holmiæ,.

Loewe, L., 2002 Global computing for bioinformatics. Briefings in Bioinformatics 3**:** 377-388.

Loewe, L., 2007 Evolution@home: observations on participant choice, work unit variation and low-effort global computing. Software Practice & Experience 37**:** 1289-1318.

Loewe, L., 2009 A framework for evolutionary systems biology. BMC Systems Biology 3**:** 27.

Loewe, L., 2012 How evolutionary systems biology will help understand adaptive landscapes and distributions of mutational effects, pp. 399-410 in *Evolutionary Systems Biology*, edited by O. Soyer. Springer, New York, NY.

Loewe, L., 2016 Systems in Evolutionary System Biology, pp. pp. 297-318 (http://evolutionarysystemsbiology.org/pdf/Loewe-2016-evosysbio.pdf) in *Encyclopedia of Evolutionary Biology*, edited by Richard M. Kliman (Chief) and Hiroshi Akashi (Section). Academic Press (Elsevier), Oxford.

Loewe, L., and J. Hillston, 2008 The distribution of mutational effects on fitness in a simple circadian clock. Lecture Notes in Bioinformatics 5307**:** 156-175.

Loewe, L., and S. Keel, 2014 BEST Names for Semantic Units to Support Reproducible Modeling, pp. 4 pages, https://www.xsede.org/documents/659353/703287/xsede659314_loewe.pdf in *2014-07-14 Workshop: "reproducibility@XSEDE", XSEDE14 Annual Conference (* https://www.xsede.org/web/reproducibility *)*, Atlanta, GA, USA. .

Loewe, L., S. Moodie and J. Hillston, 2009a Quantifying the implicit process flow abstraction in SBGN-PD diagrams with Bio-PEPA. EPTCS 6**:** 93-107 http://arxiv.org/abs/0910.1410.

Loewe, L., S. Moodie and J. Hillston, 2009b Technical Report: Defining a textual representation for SBGN Process Diagrams and translating it to Bio-PEPA for quantitative analysis of the MAPK signal transduction cascade. Technical Report, School for Informatics, University of Edinburgh EDI-INF-RR-1334**:** http://www.inf.ed.ac.uk/publications/report/1334.html

Loewe, L., K. S. Scheuer and S. Keel, 2016a Evolvix Concept: BEST Names for Semantic Units Increase Semantic Reproducibility and Help Combine Complex Biological Models, pp. https://sites.google.com/site/dslabhs2016/proceeding-papers in *Workshop on Data Science, Learning, and Applications to Biomedical & Health Sciences, January 7 - 8, 2016*. NY Academy of Sciences, New York Academy of Sciences.

Loewe, L., K. S. Scheuer, S. A. Keel, V. Vyas, B. Liblit *et al.*, 2016b Evolvix BEST Names for semantic reproducibility across code2brain interfaces. Annals of the New York Academy of Sciences**:** http://dx.doi.org/10.1111/nyas.13192 ; http://evolvix.org/naming ; http://evolvix.org/post.

Loftus, G. R., 1993 A Picture Is Worth 1000 P-Values - on the Irrelevance of Hypothesis-Testing in the Microcomputer Age. Behavior Research Methods Instruments & Computers 25**:** 250-256.

Long, D. M., M. R. Blake, S. Dutta, S. D. Holbrook, J. Kotwica-Rolinska *et al.*, 2014 Relationships between the circadian system and Alzheimer's disease-like symptoms in Drosophila. PLoS One 9**:** e106068.

Lopes, R. d. S., N. M. Resende, A. C. Honorio-Franca and E. L. Franca, 2013 Application of bioinformatics in chronobiology research. ScientificWorldJournal 2013**:** 153839.

Lorenz, L. J., J. C. Hall and M. Rosbash, 1989 Expression of a Drosophila mRNA is under circadian clock control during pupation., pp. 869-880 in *Development*.

Lorsch, J. R., F. S. Collins and J. Lippincott-Schwartz, 2014 Cell Biology. Fixing problems with cell lines. Science 346**:** 1452-1453.

Lowrey, P. L., and J. S. Takahashi, 2004 Mammalian circadian biology: elucidating genome-wide levels of temporal organization. Annu Rev Genomics Hum Genet 5**:** 407-441.

Lynch, A., G. M. Plunkett, A. J. Baker and P. F. Jenkins, 1989 A Model of Cultural-Evolution of Chaffinch Song Derived with the Meme Concept. American Naturalist 133**:** 634-653.

Lynch, M., M. C. Field, H. V. Goodson, H. S. Malik, J. B. Pereira-Leal *et al.*, 2014 Evolutionary cell biology: Two origins, one objective. Proceedings of the National Academy of Sciences of the United States of America 111**:** 16990-16994.

  QQv1r1

Ma, L., J. Ma and K. Xu, 2015 Effect of Spaceflight on the Circadian Rhythm, Lifespan and Gene Expression of Drosophila melanogaster., pp. e0121600 in *PLoS ONE*.

Mackay, T. F., S. Richards, E. A. Stone, A. Barbadilla, J. F. Ayroles *et al.*, 2012 The Drosophila melanogaster Genetic Reference Panel. Nature 482**:** 173-178.

Macneil, L. T., and A. J. Walhout, 2011 Gene regulatory networks and the role of robustness and stochasticity in the control of gene expression. Genome Res 21**:** 645-657.

Mahmood, Z., 2016 *Data science and big data computing : frameworks and methodologies*. Springer Berlin Heidelberg, New York, NY.

Majercak, J., W.-F. Chen and I. Edery, 2004 Splicing of the period gene 3'-terminal intron is regulated by light, circadian clock factors, and phospholipase C., pp. 3359-3372 in *Mol Cell Biol*.

Majercak, J., D. Kalderon and I. Edery, 1997 Drosophila melanogaster deficient in protein kinase A manifests behavior-specific arrhythmia but normal clock function.

Majercak, J., D. Sidote, P. E. Hardin and I. Edery, 1999 How a circadian clock adapts to seasonal decreases in temperature and day length., pp. 219-230 in *Neuron*.

Major, P., B. J. Kostrewski and J. Anderson, 1978 Analysis of the semantic structures of medical reference languages: part 2. Analysis of the semantic power of MeSH, ICD and SNOMED. Med Inform (Lond) 3**:** 269-281.

Malinen, E., A. Kassinen, T. Rinttila and A. Palva, 2003 Comparison of real-time PCR with SYBR Green I or 5'-nuclease assays and dot-blot hybridization with rDNA-targeted oligonucleotide probes in quantification of selected faecal bacteria. Microbiology 149**:** 269-277.

Mandrioli, D., and M. Pradella, 2015 Programming Languages shouldn't be "too Natural". ACM SIGSOFT Software Engineering Notes 40**:** 1--4.

Marrus, S. B., H. Zeng and M. Rosbash, 1996 Effect of constant light and circadian entrainment of perS flies: evidence for light-mediated delay of the negative feedback loop in Drosophila. EMBO Journal 15**:** 6877-6886.

Martinek, S., S. Inonog, A. S. Manoukian and M. W. Young, 2001 A role for the segment polarity gene shaggy/GSK-3 in the Drosophila circadian clock. Cell 105**:** 769-779.

Marygold, S. J., M. A. Crosby, J. L. Goodman and C. FlyBase, 2016 Using FlyBase, a Database of Drosophila Genes and Genomes. Methods Mol Biol 1478**:** 1-31.

Masri, S., K. Kinouchi and P. Sassone-Corsi, 2015 Circadian clocks, epigenetics, and cancer. Current Opinion in Oncology 27**:** 50-56.

Matsumoto, A., M. Ukai-Tadenuma, R. G. Yamada, J. Houl, K. D. Uno *et al.*, 2007 A functional genomics strategy reveals clockwork orange as a transcriptional regulator in the Drosophila circadian clock. Genes & development 21**:** 1687-1700.

Mavelli, F., 2012 Stochastic simulations of minimal cells: the Ribocell model. BMC Bioinformatics 13 Suppl 4**:** S10.

Mazumdar, M., S. Banerjee and H. L. Van Epps, 2010 Improved reporting of statistical design and analysis: guidelines, education, and editorial policies. Methods Mol Biol 620**:** 563-598.

McCallum, Q. E., 2013 *Bad Data Handbook: Mapping the world of data problems*. O'Reilly, Sebastopol, CA.

McClung, C. A., 2013 How Might Circadian Rhythms Control Mood? Let Me Count the Ways..... Biological Psychiatry.

McDonald, M. J., and M. Rosbash, 2001 Microarray analysis and organization of circadian gene expression in Drosophila. Cell 107**:** 567-578.

McDonald, M. J., M. Rosbash and P. Emery, 2001 Wild-type circadian rhythmicity is dependent on closely spaced E boxes in the Drosophila timeless promoter. Mol Cell Biol 21**:** 1207-1217.

McIntyre, L. M., K. K. Lopiano, A. M. Morse, V. Amin, A. L. Oberg *et al.*, 2011 RNA-seq: technical variability and sampling. Bmc Genomics 12.

McKenzie, J. E., G. Salanti, S. C. Lewis and D. G. Altman, 2013 Meta-analysis and The Cochrane Collaboration: 20 years of the Cochrane Statistical Methods Group. Syst Rev 2**:** 80.

McNutt, M., 2014 Journals unite for reproducibility. Science 346**:** 679.

Meissner, R.-A., V. L. Kilman, J.-M. Lin and R. Allada, 2008 TIMELESS is an important mediator of CK2 effects on circadian clock function in vivo. Journal of Neuroscience 28**:** 9732-9740.

Menet, J. S., K. C. Abruzzi, J. Desrochers, J. Rodriguez and M. Rosbash, 2010 Dynamic PER repression mechanisms in the Drosophila circadian clock: from on-DNA to off-DNA. Genes & development 24**:** 358-367.

Merbitz-Zahradnik, T., and E. Wolf, 2015 How is the inner circadian clock controlled by interactive clock proteins?: Structural analysis of clock proteins elucidates their physiological role. FEBS Lett 589**:** 1516-1529.

Merrill, G. H., 2009 Concepts and synonymy in the UMLS Metathesaurus. J Biomed Discov Collab 4**:** 7.

Mesnard, O., and L. A. Barba, 2016 Reproducible and replicable CFD: it's harder than you think. Computing in Science and Engineering**:** https://arxiv.org/abs/1605.04339.

Mezan, S., R. Ashwal-Fluss, R. Shenhav, M. Garber and S. Kadener, 2013 Genome-wide assessment of post-transcriptional control in the fly brain., pp. 49 in *Front Mol Neurosci*.

Millis, S. R., 2003 Statistical practices: The seven deadly sins. Child Neuropsychology 9**:** 221-233.

Minikel, E. V., I. Zerr, S. J. Collins, C. Ponto, A. Boyd *et al.*, 2014 Ascertainment bias causes false signal of anticipation in genetic prion disease. Am J Hum Genet 95**:** 371-382.

Mitchell, C. S., A. Cates, R. B. Kim and S. K. Hollinger, 2015 Undergraduate Biocuration: Developing Tomorrow's Researchers While Mining Today's Data. J Undergrad Neurosci Educ 14**:** A56-65.

Mitra, K., A. R. Carvunis, S. K. Ramesh and T. Ideker, 2013 Integrative approaches for finding modular structure in biological networks. Nature Reviews Genetics 14**:** 719-732.

Miura, S., T. Shimokawa and T. Nomura, 2008 Stochastic simulations on a model of circadian rhythm generation. Bio Systems 93**:** 133-140.

Mobley, A., S. K. Linder, R. Braeuer, L. M. Ellis and L. Zwelling, 2013 A Survey on Data Reproducibility in Cancer Research Provides Insights into Our Limited Ability to Translate Findings from the Laboratory to the Clinic. Plos One 8.

Mockler, T. C., T. P. Michael, H. D. Priest, R. Shen, C. M. Sullivan *et al.*, 2007 The DIURNAL project: DIURNAL and circadian expression profiling, model-based pattern matching, and promoter analysis. Cold Spring Harb Symp Quant Biol 72**:** 353-363.

Molina-Rodríguez, M., and V. Álvarez, 2016 Los ritmos circadianos en cancer y la cronoterapia. Iatreia 29.3**:** 301.

Moodie, S., S. Moodie, N. Le Novere, E. Demir, H. Mi *et al.*, 2011 Systems Biology Graphical Notation: Process Description language Level 1. Nature Precedings.

Mori, U., A. Mendiburu and J. A. Lozano, 2016 Similarity Measure Selection for Clustering Time Series Databases. Ieee Transactions on Knowledge and Data Engineering 28**:** 181-195.

Mortazavi, A., B. A. Williams, K. Mccue, L. Schaeffer and B. Wold, 2008 Mapping and quantifying mammalian transcriptomes by RNA-Seq. Nature Methods 5**:** 621-628.

Moura Neto, F. D., and A. n. J. Silva Neto, 2013 *An introduction to inverse problems with applications*. Springer, Heidelberg ; New York.

Murtaugh, P. A., 2014 In defense of P values. Ecology 95**:** 611-617.

Musen, M. A., N. F. Noy, N. H. Shah, P. L. Whetzel, C. G. Chute *et al.*, 2012 The National Center for Biomedical Ontology. J Am Med Inform Assoc 19**:** 190-195.

Muskus, M. J., F. Preuss, J.-Y. Fan, E. S. Bjes and J. L. Price, 2007 Drosophila DBT lacking protein kinase activity produces long-period and arrhythmic circadian behavioral and molecular rhythms., pp. 8049-8064 in *Mol Cell Biol*.

Myers, M. P., K. Wager-Smith, A. Rothenfluh-Hilfiker and M. W. Young, 1996 Light-Induced Degradation of TIMELESS and Entrainment of the Drosophila Circadian Clock. Science, New Series 271**:** 1736-1740.

Nagoshi, E., K. Sugino, E. Kula, E. Okazaki, T. Tachibana *et al.*, 2010 Dissecting differential gene expression within the circadian neuronal circuit of Drosophila., pp. 60-68 in *Nat Neurosci*.

Naidoo, N., W. Song, M. Hunter-Ensor and A. Sehgal, 1999 A role for the proteasome in the light response of the timeless clock protein. Science 285**:** 1737-1741.

Najafi, A., G. Bidkhori, J. H. Bozorgmehr, I. Koch and A. Masoudi-Nejad, 2014 Genome scale modeling in systems biology: algorithms and resources. Curr Genomics 15**:** 130-159.

Nakagawa, S., and I. C. Cuthill, 2007 Effect size, confidence interval and statistical significance: a practical guide for biologists. Biological Reviews 82**:** 591-605.

NASA, F. Chandler, T., Y. H. Chang, James, A. Mosleh, J. Marble, L. *et al.*, 2006-07 *Human Reliability Analysis Methods Selection Guidance for NASA*. NASA/OSMA, NASA, Washington, DC.

NASA, K. Leiden, K. R. Laughery, J. Keller, J. French *et al.*, 2001-09-30 *A Review of Human Performance Models for the Prediction of Human Error*. National Aeronautics and Space Administration, System-Wide Accident Prevention Program, Ames Research Center, Moffett Field, CA, http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.468.6969&rep=rep1&type=pdf.

NASA, M. Stamatelatos and H. Dezfuli, 2011 *Probabilistic Risk Assessment Procedures Guide for NASA Managers and Practitioners (2nd Edition)*, NASA Headquarters, Washington, DC.

NCI-NHGRI Working Group on Replication in Association Studies, S. J. Chanock, T. Manolio, M. Boehnke, E. Boerwinkle *et al.*, 2007 Replicating genotype-phenotype associations. Nature 447**:** 655-660.

Ng, A., B. Bursteinas, Q. Gao, E. Mollison and M. Zvelebil, 2006 Resources for integrative systems biology: from data through databases to networks and dynamic system models. Brief Bioinform 7**:** 318-330.

NIH, 2015 National Institutes of Health Plan for Increasing Access to Scientific Publications and Digital Scientific Data from NIH Funded Scientific Research. 2015-02**:** https://grants.nih.gov/grants/NIH-Public-Access-Plan.pdf.

NIH, 2016 NIH Request for Information (RFI): Strategies for NIH Data Management, Sharing, and Citation . RFI Notice Number NOT-OD-17-015**:** http://grants.nih.gov/grants/guide/notice-files/NOT-OD-17-015.html.

NIH, Blue Ribbon Panel, Tyler Jacks, E. Jaffee, D. Singer *et al.*, 2016 Cancer Moonshot Blue Ribbon Panel Report 2016-10-17, pp. https://www.cancer.gov/research/key-initiatives/moonshot-cancer-initiative/blue-ribbon-panel/blue-ribbon-panel-report-2016.pdf. National Institutes of Health,.

NIH, D. DeMets, L. Tabak, R. Altman, D. Botstein *et al.*, 2012 Data and Informatics Working Group Report to the Advisory Committee to the Director, pp. National Institutes of Health,, http://acd.od.nih.gov/Data and Informatics Working Group Report.PDF.

Nordstokke, D. W., and B. D. Zumbo, 2007 A Cautionary Tale About Levene's Tests for Equal Variances. Journal of Educational Research & Policy Studies 7**:** 1-14.

Nuzzo, R., 2014 Scientific method: statistical errors. Nature 506**:** 150-152.

Oconnell, P., and M. Rosbash, 1984 Sequence, Structure, and Codon Preference of the Drosophila Ribosomal Protein-49 Gene. Nucleic Acids Research 12**:** 5495-5513.

Ogawa, Y., K. Arakawa, K. Kaizu, F. Miyoshi, Y. Nakayama *et al.*, 2008 Comparative study of circadian oscillatory network models of Drosophila. Artificial life 14**:** 29-48.

Oh, Y., S.-E. Yoon, Q. Zhang, H.-S. Chae, I. Daubnerová *et al.*, 2014 A homeostatic sleep-stabilizing pathway in Drosophila composed of the sex peptide receptor and its ligand, the myoinhibitory peptide., pp. e1001974 in *PLoS Biol.*

Okada, T., T. Sakai, T. Murata, K. Kako, K. Sakamoto *et al.*, 2001 Promoter analysis for daily expression of Drosophila timeless gene. Biochemical and Biophysical Research Communications 283**:** 577-582.

Oliver, D., and J. Phillips, 1970 Technical Note. Drosophila Information Service.

Onitilo, A. A., 2014 Is it time for the Cochrane Collaboration to reconsider its meta-analysis methodology? Clin Med Res 12**:** 2-3.

Opitz, J. M., 1983 Cotterman and combinatorial genetics. Am J Med Genet 16**:** 389-392.

Orchard, S., and H. Hermjakob, 2015 Shared resources, shared costs--leveraging biocuration resources. Database (Oxford) 2015.

Oreskes, N., K. Shraderfrechette and K. Belitz, 1994 Verification, Validation, and Confirmation of Numerical-Models in the Earth-Sciences. Science 263**:** 641-646.

Orpana, A. K., T. H. Ho and J. Stenman, 2012 Multiple Heat Pulses during PCR Extension Enabling Amplification of GC-Rich Sequences and Reducing Amplification Bias. Analytical Chemistry 84**:** 2081-2087.

Özkaya, Ö., and E. Rosato, 2012 The circadian clock of the fly: a neurogenetics journey through time., pp. 79-123 in *Adv Genet*.

Ozturk, N. N., C. P. C. Selby, Y. Y. Annayev, D. D. Zhong and A. A. Sancar, 2011 Reaction mechanism of Drosophila cryptochrome. Proceedings of the National Academy of Sciences of the United States of America 108**:** 516-521.

Packer, M., 2016 Data sharing: lessons from Copernicus and Kepler. BMJ 354**:** i4911.

Panko, R. R., 1998 Applying code inspection to spreadsheet testing. Effective Utilization and Management of Emerging Information Technologies**:** 358-362.

Panko, R. R., 2013 Users and Uses Research: Understanding "The Familiar Unknown". Journal of Organizational and End User Computing 25**:** Iv-Xi.

Panko, R. R., 2016 Human Error Website. Accessed 2016-10-31**:** http://panko.shidler.hawaii.edu/HumanErr/Index.htm

Panko, R. R., and S. Aurigemma, 2010 Revising the Panko-Halverson taxonomy of spreadsheet errors. Decision Support Systems 49**:** 235-244.

Parekh, P. K., A. R. Ozburn and C. A. McClung, 2015 Circadian clock genes: effects on dopamine, reward and addiction. Alcohol 49**:** 341-349.

Patel, V. R., K. Eckel-Mahan, P. Sassone-Corsi and P. Baldi, 2012 CircadiOmics: integrating circadian genomics, transcriptomics, proteomics and metabolomics. Nature Methods 9**:** 772-773.

Patil, P., P. O. Bachant-Winner, B. Haibe-Kains and J. T. Leek, 2015 Test set bias affects reproducibility of gene signatures. Bioinformatics 31**:** 2318-2323.

Pegoraro, M., S. Noreen, S. Bhutani, A. Tsolou, R. Schmid *et al.*, 2014 Molecular evolution of a pervasive natural amino-acid substitution in Drosophila cryptochrome. PLoS ONE 9**:** e86483.

Pembroke, W. G., A. Babbs, K. E. Davies, C. P. Ponting and P. L. Oliver, 2015 Temporal transcriptomics suggest that twin-peaking genes reset the clock. Elife 4.

Peschel, N., 2008 New insights into circadian photoreception and the molecular regulation of the resetting of Drosophilas circadian clock. 1-163.

Peschel, N., K. F. Chen, G. Szabo and R. Stanewsky, 2009 Light-dependent interactions between the Drosophila circadian clock factors cryptochrome, jetlag, and timeless. Curr Biol 19**:** 241-247.

Peschel, N., and C. Helfrich-Förster, 2011 Setting the clock – by nature: Circadian rhythm in the fruitfly Drosophila melanogaster. Febs Letters 585**:** 1435-1442.

Petri, B., and M. Stengl, 2001 Phase response curves of a molecular model oscillator: implications for mutual coupling of paired oscillators., pp. 125-141 in *J Biol Rhythms*.

Pierce, B. C., 2002 *Types and programming languages*. MIT Press, Cambridge, Mass.

Pierce, B. C., 2005 *Advanced topics in types and programming languages*. MIT Press, Cambridge, Mass.

Pittendrigh, C., and G. Victor, 1957 An Oscillator Model for Biological Clocks in *Rhythmic and Synthetic Processes in Growth*, edited by D. Rudnick. Princeton University Press, Priceton, NJ.

Pizarro, A., K. Hayer, N. F. Lahens and J. B. Hogenesch, 2013 CircaDB: a database of mammalian circadian gene expression profiles. Nucleic Acids Res 41**:** D1009-1013.

Plautz, J. D., M. Straume, R. Stanewsky, C. F. Jamison, C. Brandes *et al.*, 1997 Quantitative analysis of Drosophila period gene transcription in living animals. Journal of biological rhythms 12**:** 204-217.

Plowman, A. B., 2008 BIAZA statistics guidelines: toward a common application of statistical tests for zoo research. Zoo Biol 27**:** 226-233.

Pogue-Geile, K. L., J. Lyons-Weiler and D. C. Whitcomb, 2006 Molecular overlap of fly circadian rhythms and human pancreatic cancer. Cancer Letters 243**:** 55-57.

Poldrack, R. A., and J. B. Poline, 2015 The publication and reproducibility challenges of shared data. Trends Cogn Sci 19**:** 59-61.

Preussner, M., and F. Heyd, 2016 Post-transcriptional control of the mammalian circadian clock: implications for health and disease. Pflugers Archiv-European Journal of Physiology 468**:** 983-991.

Price, J. L., J. Blau, A. Rothenfluh, M. Abodeely, B. Kloss *et al.*, 1998 double-time Is a Novel Drosophila Clock Gene that Regulates PERIOD Protein Accumulation. Cell 94**:** 83-95.

Purcell, O., B. Jain, J. R. Karr, M. W. Covert and T. K. Lu, 2013 Towards a whole-cell modeling approach for synthetic biology. Chaos 23**:** 025112.

Qiu, J., and P. E. Hardin, 1996 per mRNA cycling is locked to lights-off under photoperiodic conditions that support circadian feedback loop function., pp. 4182-4188 in *Mol Cell Biol*.

        Updates may exist at bioRxiv https://doi.org/10.1101/099192   QQv1r1

Qu, Y., and M. Boutjdir, 2007 RNase Protection Assay for Quantifying Gene Expression Levels in *Methods in Molecular Biology - Cardian Gene Expression: Methods and Protocols*, edited by J. Zhang and G. Rokosh. Humana Press Inc., Totowa, N.J.

Raymond, E. S., 2003 *The Art of Unix Programming*. Addison-Wesley (Pearson), Boston, MA, USA. http://www.catb.org/~esr/writings/taoup/.

Razali, N., J. Noble and S. Marshall, 2015-10-26 Operators and Precedence in Programming Languages. PLATEAU'15: 53-56 (http://dx.doi.org/10.1145/2846680.2846690).

Read, K. B., J. R. Sheehan, M. F. Huerta, L. S. Knecht, J. G. Mork *et al.*, 2015 Sizing the Problem of Improving Discovery and Access to NIH-Funded Data: A Preliminary Study. PLoS One 10: e0132735.

Reason, J. T., 1990 *Human error*. Cambridge University Press, Cambridge England ; New York.

Reason, J. T., 2013 *A life in error : from little slips to big disasters*. Ashgate, Surrey, UK England.

Reason, J. T., 2015 *Organizational accidents revisited*. Ashgate, Burlington, VT.

Reason, J. T., and A. Hobbs, 2003 *Managing maintenance error : a practical guide*. Ashgate, Aldershot, Hampshire, England ; Burlington, VT, USA.

Reason, J. T., and K. Mycielska, 1982 *Absent-minded? : the psychology of mental lapses and everyday errors*. Prentice-Hall, Englewood Cliffs, N.J.

Refinetti, R., G. C. Lissen and F. Halberg, 2007 Procedures for numerical analysis of circadian rhythms. Biol Rhythm Res 38: 275-325.

Reiser, L., T. Z. Berardini, D. Li, R. Muller, E. M. Strait *et al.*, 2016 Sustainable funding for biocuration: The Arabidopsis Information Resource (TAIR) as a case study of a subscription-based funding model. Database (Oxford) 2016.

Richier, B., C. Michard-Vanhee, A. Lamouroux, C. Papin and F. Rouyer, 2008 The clockwork orange Drosophila protein functions as both an activator and a repressor of clock gene expression. J Biol Rhythms 23: 103-116.

Risau-Gusman, S., and P. M. Gleiser, 2012 Modelling the effect of phosphorylation on the circadian clock of Drosophila. Journal of theoretical biology 307: 53-61.

Risau-Gusman, S., and P. M. Gleiser, 2014 A Mathematical Model of Communication between Groups of Circadian Neurons in Drosophila melanogaster. Journal of biological rhythms 29: 401-410.

Robert, C., and M. Watson, 2015 Errors in RNA-Seq quantification affect genes of relevance to human disease. Genome Biol 16: 177.

Robert, C. P., J. M. Cornuet, J. M. Marin and N. S. Pillai, 2011 Lack of confidence in approximate Bayesian computation model choice. Proc Natl Acad Sci U S A 108: 15112-15117.

Roberts, A., L. Schaeffer and L. Pachter, 2013 Updating RNA-Seq analyses after re-annotation. Bioinformatics 29: 1631-1637.

Robinson, S. W., P. Herzyk, J. A. Dow and D. P. Leader, 2013 FlyAtlas: database of gene expression in the tissues of Drosophila melanogaster. Nucleic Acids Res 41: D744-750.

Roden, D. M., 2011 Personalized medicine and the genotype-phenotype dilemma. J Interv Card Electrophysiol 31: 17-23.

Rodriguez, J., C.-H. A. Tang, Y. L. Khodor, S. Vodala, J. S. Menet *et al.*, 2013 Nascent-Seq analysis of Drosophila cycling gene expression. Proceedings of the National Academy of Sciences of the United States of America 110: E275-E284.

Rodriguez-Esteban, R., 2015 Biocuration with insufficient resources and fixed timelines. Database (Oxford) 2015.

Roenneberg, T., and M. Merrow, 1998 Molecular circadian oscillators: An alternative hypothesis. Journal of Biological Rhythms 13: 167-179.

Rosato, E., A. A. Peixoto, A. Gallippi, C. P. Kyriacou and R. Costa, 1996 Mutational mechanisms, phylogeny, and evolution of a repetitive region within a clock gene of Drosophila melanogaster., pp. 392-408 in *J Mol Evol*.

Rosato, E., E. Tauber and C. P. Kyriacou, 2006 Molecular genetics of the fruit-fly circadian clock. European Journal of Human Genetics 14: 729-738.

Rothenfluh, A., M. W. Young and L. Saez, 2000a A TIMELESS-independent function for PERIOD proteins in the Drosophila clock. Neuron 26: 505-514.

Rothenfluh, A. A., M. M. Abodeely, J. L. J. Price and M. W. M. Young, 2000b Isolation and analysis of six timeless alleles that cause short- or long-period circadian rhythms in Drosophila. Genetics 156**:** 665-675.

Rouyer, F., M. Rachidi, C. Pikielny and M. Rosbash, 1997 A new gene encoding a putative transcription factor regulated by the Drosophila circadian clock., pp. 3944-3954 in *EMBO J.*

Rubinson, C., 2014 Ontological Considerations When Modeling Missing Data With Relational Databases. Social Science Computer Review 32**:** 769-780.

Rund, S. S., J. E. Gentile and G. E. Duffield, 2013 Extensive circadian and light regulation of the transcriptome in the malaria mosquito Anopheles gambiae. BMC Genomics 14**:** 218.

Rund, S. S., T. Y. Hou, S. M. Ward, F. H. Collins and G. E. Duffield, 2011 Genome-wide profiling of diel and circadian gene expression in the malaria vector Anopheles gambiae. Proc Natl Acad Sci U S A 108**:** E421-430.

Ruoff, P., M. K. Christensen and V. K. Sharma, 2005 PER/TIM-mediated amplification, gene dosage effects and temperature compensation in an interlocking-feedback loop model of the Drosophila circadian clock. Journal of theoretical biology 237**:** 41-57.

Ruoff, P., and L. Rensing, 1996 The temperature-compensated goodwin model simulates many circadian clock properties. Journal of Theoretical Biology 179**:** 275-285.

Ruoff, P., L. Rensing, R. Kommedal and S. Mohsenzadeh, 1997 Modeling temperature compensation in chemical and biological oscillators. Chronobiology International 14**:** 499-510.

Ruoff, P., M. Vinsjevik, C. Monnerjahn and L. Rensing, 1999 The Goodwin Oscillator: On the Importance of Degradation Reactions in the Circadian Clock. Journal of biological rhythms 14**:** 469-479.

Rutila, J. E., V. Suri, M. Le, W. V. So, M. Rosbash *et al.*, 1998 CYCLE is a second bHLH-PAS clock protein essential for circadian rhythmicity and transcription of Drosophila period and timeless. Cell 93**:** 805-814.

Saez-Rodriguez, J., J. C. Costello, S. H. Friend, M. R. Kellen, L. Mangravite *et al.*, 2016 Crowdsourcing biomedical research: leveraging communities as innovation engines. Nature Reviews Genetics 17**:** 470-486.

Salanti, G., and J. P. Ioannidis, 2009 Synthesis of observational studies should consider credibility ceilings. J Clin Epidemiol 62**:** 115-122.

Salavaty, A., 2015 Carcinogenic effects of circadian disruption: an epigenetic viewpoint. Chinese Journal of Cancer 34.

Salimi, N., and R. Vita, 2006 The biocurator: Connecting and enhancing scientific data. Plos Computational Biology 2**:** 1190-1192.

Salsburg, D. S., 1985 The Religion of Statistics as Practiced in Medical Journals. American Statistician 39**:** 220-223.

Sambrook, J., E. Fritsch and T. Maniatis, 1989 *Molecular Cloning: A Laboratory Manual.* Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.

Samuels, S., B. Balint, H. von der Leyen, P. Hupe, L. de Koning *et al.*, 2016 Precision medicine in cancer: challenges and recommendations from an EU-funded cervical cancer biobanking study. Br J Cancer.

Sandrelli, F., E. Tauber, M. Pegoraro, G. Mazzotta, P. Cisotto *et al.*, 2007 A molecular basis for natural selection at the timeless locus in Drosophila melanogaster. Science 316**:** 1898-1900.

Sanghvi, J. C., S. Regot, S. Carrasco, J. R. Karr, M. V. Gutschow *et al.*, 2013 Accelerated discovery via a whole-cell model. Nat Methods 10**:** 1192-1195.

Sathyanarayanan, S., X. Zheng, R. Xiao and A. Sehgal, 2004 Posttranslational regulation of Drosophila PERIOD protein by protein phosphatase 2A. Cell 116**:** 603-615.

Savalei, V., and E. Dunn, 2015 Is the call to abandon p-values the red herring of the replicability crisis? Frontiers in Psychology 6.

Sawyer, L. A., J. M. Hennessy, A. A. Peixoto, E. Rosato, H. PARKINSON *et al.*, 1997 Natural variation in a Drosophila clock gene and temperature compensation., pp. 2117-2120 in *Science.*

Scheper, T. O., D. Klinkenberg, C. Pennartz and J. van Pelt, 1999a A mathematical model for the intracellular circadian rhythm generator. Journal of Neuroscience 19**:** 40-47.

                     QQv1r1

Scheper, T. O. T., D. D. Klinkenberg, J. J. van Pelt and C. C. Pennartz, 1999b A model of molecular circadian clocks: multiple mechanisms for phase shifting and a requirement for strong nonlinear interactions. Journal of biological rhythms 14**:** 213-220.

Schnoes, A. M., S. D. Brown, I. Dodevski and P. C. Babbitt, 2009 Annotation error in public databases: misannotation of molecular function in enzyme superfamilies. PLoS Comput Biol 5**:** e1000605.

Schutt, R., and C. O'Neil, 2013 *Doing data science*. O'Reilly Media, Beijing ; Sebastopol.

Schwartz, B., 2004 *The paradox of choice : why more is less*. Ecco, New York.

Scribner, E. Y., and H. M. Fathallah-Shaykh, 2011 Modeling of regulatory networks: theory and applications in the study of the Drosophila circadian clock. Methods in enzymology 487**:** 39-71.

Sehgal, A., J. L. Price, B. Man and M. W. Young, 1994 Loss of Circadian Behavioral Rhythms and per RNA Oscillations in the Drosophila Mutant timeless. Science, New Series 263**:** 1603-1606.

Sehgal, A., A. Rothenfluh-Hilfiker, M. Hunter-Ensor, Y. Chen, M. P. Myers *et al.*, 1995 Rhythmic expression of timeless: a basis for promoting circadian cycles in period gene autoregulation., pp. 808-810 in *Science*.

Sephton, S., and D. Spiegel, 2003 Circadian disruption in cancer: a neuroendocrine-immune pathway from stress to disease? Brain Behavior and Immunity 17**:** 321-328.

SEQC/MAQC-III Consortium, 2014 A comprehensive assessment of RNA-seq accuracy, reproducibility and information content by the Sequencing Quality Control Consortium. Nat Biotechnol 32**:** 903-914.

Shabalina, S. A., L. Y. Yampolsky and A. S. Kondrashov, 1997 Rapid decline of fitness in panmictic populations of Drosophila melanogaster maintained under relaxed natural selection. Proceedings Of the National Academy Of Sciences Of the United States Of America 94**:** 13034-13039.

Shafer, O. T., M. Rosbash and J. W. Truman, 2002 Sequential nuclear accumulation of the clock proteins period and timeless in the pacemaker neurons of Drosophila melanogaster. Journal of Neuroscience 22**:** 5946-5954.

Sharkey, F. H., I. M. Banat and R. Marchant, 2004 Detection and quantification of gene expression in environmental bacteriology. Applied and Environmental Microbiology 70**:** 3795-3806.

Sharma, A., S. Tiwari and M. Singaravel, 2016 Circadian rhythm disruption: health consequences. Biological Rhythm Research 47**:** 191-213.

Sharma, V. K., 2003 Adaptive significance of circadian clocks. Chronobiology International 20**:** 901-919.

Sharpe, D., 2013 Why the Resistance to Statistical Innovations? Bridging the Communication Gap. Psychological Methods 18**:** 572-582.

Shenk, D., 1997 *Data smog : surviving the information glut*. Harper Edge, San Francisco, Calif.

Shi, L. M., L. H. Reid, W. D. Jones, R. Shippy, J. A. Warrington *et al.*, 2006 The MicroArray Quality Control (MAQC) project shows inter- and intraplatform reproducibility of gene expression measurements. Nature Biotechnology 24**:** 1151-1161.

Shi, M., Z. Yue, A. Kuryatov, J. M. Lindstrom, A. Sehgal *et al.*, 2014 Identification of Redeye, a new sleep-regulating protein whose expression is modulated by sleep amount. eLife 3**:** e01473.

Sidote, D., J. Majercak, V. Parikh and I. Edery, 1998 Differential effects of light and heat on the Drosophila circadian clock proteins PER and TIM. Molecular and Cellular Biology 18**:** 2004-2013.

Simoni, A., W. Wolfgang, M. P. Topping, R. G. Kavlie, R. Stanewsky *et al.*, 2014 A mechanosensory pathway to the Drosophila circadian clock., pp. 525-528 in *Science*.

Singhal, A., R. Leaman, N. Catlett, T. Lemberger, J. McEntyre *et al.*, 2016 Pressing needs of biomedical text mining in biocuration and beyond: opportunities and challenges. Database (Oxford) 2016.

Smith, N. J. J., 2008 *Vagueness and Degrees of Truth*. Oxford University Press, Oxford, Great Britain.

Smolen, P., 2002 A detailed predictive model of the mammalian circadian clock.

Smolen, P., D. Baxter and J. Byrne, 2002 A Reduced Model Clarifies the Role of Feedback Loops and Time Delays in the Drosophila Circadian Oscillator. Biophysical Journal 83**:** 2349-2359.

Smolen, P., D. A. Baxter and J. H. Byrne, 2000a Mathematical modeling of gene networks. Neuron 26**:** 567-580.

Smolen, P., D. A. Baxter and J. H. Byrne, 2000b Modeling transcriptional control in gene networks - Methods, recent results, and future directions. Bulletin of Mathematical Biology 62**:** 247-292.

Smolen, P., D. A. Baxter and J. H. Byrne, 2001 Modeling circadian oscillations with interlocking positive and negative feedback loops. Journal of Neuroscience 21**:** 6644-6656.

Smolen, P., P. E. Hardin, B. S. Lo, D. A. Baxter and J. H. Byrne, 2004 Simulation of Drosophila circadian oscillations, mutations, and light responses by a model with VRI, PDP-1, and CLK. Biophysical Journal 86**:** 2786-2802.

So, W. V., and M. Rosbash, 1997 Post-transcriptional regulation contributes to Drosophila clock gene mRNA cycling. EMBO Journal 16**:** 7146-7155.

Sonawani, A., P. Nilawe, R. S. Barai and S. Idicula-Thomas, 2010 ClotBase: a knowledgebase on proteins involved in blood coagulation. Blood 116**:** 855-856.

Soranno, P. A., E. G. Bissell, K. S. Cheruvelil, S. T. Christel, S. M. Collins *et al.*, 2015 Building a multi-scaled geospatial temporal ecology database from disparate data sources: fostering open science and data reuse. Gigascience 4**:** 28.

Soubeyrand, S., F. Carpentier, F. Guiton and E. K. Klein, 2013 Approximate Bayesian computation with functional statistics. Statistical Applications in Genetics and Molecular Biology 12.

St Pierre, S., and P. McQuilton, 2009 Inside FlyBase: biocuration as a career. Fly (Austin) 3**:** 112-114.

Stainforth, D. A., T. Aina, C. Christensen, M. Collins, N. Faull *et al.*, 2005 Uncertainty in predictions of the climate response to rising levels of greenhouse gases. Nature 433**:** 403-406.

Stajich, J. E., D. Block, K. Boulez, S. E. Brenner, S. A. Chervitz *et al.*, 2002 The Bioperl toolkit: Perl modules for the life sciences. Genome Res 12**:** 1611-1618.

Stanewsky, R., 2003 Genetic analysis of the circadian system in Drosophila melanogaster and mammals. Journal of neurobiology 54**:** 111-147.

Stanewsky, R., M. Kaneko, P. Emery, B. Beretta, K. Wager-Smith *et al.*, 1998 The cryb mutation identifies cryptochrome as a circadian photoreceptor in Drosophila., pp. 681-692 in *Cell*.

Stanewsky, R., K. S. Lynch, C. Brandes and J. C. Hall, 2002 Mapping of elements involved in regulating normal temporal period and timeless RNA expression patterns in Drosophila melanogaster., pp. 293-306 in *J Biol Rhythms*.

Stanewsky, R. R., C. F. C. Jamison, J. D. J. Plautz, S. A. S. Kay and J. C. J. Hall, 1997 Multiple circadian-regulated elements contribute to cycling period gene expression in Drosophila. EMBO Journal 16**:** 5006-5018.

Stanton-Geddes, J., C. G. de Freitas and C. D. Dambros, 2014 In defense of P values: comment on the statistical methods actually used by ecologists. Ecology 95**:** 637-642.

Sterne, J., 2003 Commentary: Null points - has interpretation of significance tests improved? International Journal of Epidemiology 32**:** 693-694.

Stodden, V., 2015 Reproducing Statistical Results. Annu. Rev. Stat. Appl. 2**:** 1-19.

Stodden, V., F. Leisch and R. D. Peng (Editors), 2014 *Implementing Reproducible Research*. CRC Press, Boca Raton, FL.

Stodden, V., M. McNutt, D. H. Bailey, E. Deelman, Y. Gil *et al.*, 2016 Enhancing reproducibility for computational methods. Science 354**:** 1240-1241.

Stoleru, D. D., Y. Y. Peng, J. J. Agosto and M. M. Rosbash, 2004 Coupled oscillators control morning and evening locomotor behaviour of Drosophila. Nature 431**:** 862-868.

Streit, S., C. W. Michalski, M. Erkan, J. Kleeff and H. Friess, 2009 Northern blot analysis for detection and quantification of RNA in pancreatic cancer cells and tissues. Nat Protoc 4**:** 37-43.

Stumpf, M. P., 2014 Approximate Bayesian inference for complex ecosystems. F1000Prime Rep 6**:** 60.

Subramanian, P., V. Prasanna, J. J. Jayapalan, P. S. Abdul Rahman and O. H. Hashim, 2014 Role of Bacopa monnieri in the temporal regulation of oxidative stress in clock mutant (cryb) of Drosophila melanogaster., pp. 37-44 in *J Insect Physiol*.

Sullivan, D. E., J. L. Gabbard, Jr., M. Shukla and B. Sobral, 2010 Data integration for dynamic and sustainable systems biology resources: challenges and lessons learned. Chem Biodivers 7**:** 1124-1141.

Sun, Y. Q., J. Y. Li, J. X. Liu, B. Y. Sun and C. Chow, 2014 An improvement of symbolic aggregate approximation distance measure for time series. Neurocomputing 138**:** 189-198.

Sunnaker, M., A. G. Busetto, E. Numminen, J. Corander, M. Foll *et al.*, 2013 Approximate Bayesian Computation. PLoS Computational Biology 9.

Supp, S. R., D. N. Koons and S. K. M. Ernest, 2015a Using life history trade-offs to understand core-transient structuring of a small mammal community. Ecosphere 6.

Supp, S. R., F. A. La Sorte, T. A. Cormier, M. C. W. Lim, D. R. Powers *et al.*, 2015b Citizen-science data provides new insight into annual and seasonal variation in migration patterns. Ecosphere 6.

Taniguchi, M., K. Miura, H. Iwao and S. Yamanaka, 2001 Quantitative assessment of DNA microarrays - Comparison with Northern blot analyses. Genomics 71**:** 34-39.

Tarantola, A., 2005 *Inverse problem theory and methods for model parameter estimation*. Society for Industrial and Applied Mathematics, Philadelphia, PA.

Tarantola, A., 2006 Popper, Bayes and the inverse problem. Nature Physics 2**:** 492-494.

Tarantola, A., and B. Valette, 1982 Inverse Problems = Quest for Information. Journal of Geophysics-Zeitschrift Fur Geophysik 50**:** 159-170.

Taroni, F., A. Biedermann and S. Bozza, 2016 Statistical hypothesis testing and common misinterpretations: Should we abandon p-value in forensic science applications? Forensic Science International 259.

Temme, C., L. B. Zhang, E. Kremmer, C. Ihling, A. Chartier *et al.*, 2010 Subunits of the Drosophila CCR4-NOT complex and their roles in mRNA deadenylation. Rna-a Publication of the Rna Society 16**:** 1356-1370.

Thalheim, B., and K. D. Schewe, 2011 NULL 'Value' Algebras and Logics. Information Modelling and Knowledge Bases Xxii 225**:** 354-367.

Tharyan, P., 1998 The relevance to meta-analysis, systematic reviews and the cochrane collaboration to clinical psychiatry. Indian J Psychiatry 40**:** 135-148.

The UniProt Consortium, 2017 UniProt: the universal protein knowledgebase. Nucleic Acids Res 45**:** D158-D169.

Theurkauf, W. E., H. Baum, J. Y. Bo and P. C. Wensink, 1986 Tissue-Specific and Constitutive Alpha-Tubulin Genes of Drosophila-Melanogaster Code for Structurally Distinct Proteins. Proceedings of the National Academy of Sciences of the United States of America 83**:** 8477-8481.

Toni, T., D. Welch, N. Strelkowa, A. Ipsen and M. P. Stumpf, 2009a Approximate Bayesian computation scheme for parameter inference and model selection in dynamical systems. J R Soc Interface 6**:** 187-202.

Toni, T., D. Welch, N. Strelkowa, A. Ipsen and M. P. H. Stumpf, 2009b Approximate Bayesian computation scheme for parameter inference and model selection in dynamical systems. Journal of the Royal Society Interface 6**:** 187-202.

Trafimow, D., and M. Marks, 2015 Editorial: banning null hypothesis significance testing procedures. Basic and Applied Social Psychology 37**:** 1-2.

Tressoldi, P. E., D. Giofre, F. Sella and G. Cumming, 2013 High Impact = High Statistical Standards? Not Necessarily So. Plos One 8.

Turing, A. M., 1936 On computable numbers, with an application to the Entscheidungsproblem. Proc. Lond. Math. Soc. 42**:** 230-265.

Tyson, J. J., C. I. Hong, C. D. Thron and B. Novak, 1999 A simple model of circadian rhythms based on dimerization and proteolysis of PER and TIM., pp. 2411-2417 in *Biophys J*.

Ueda, H. R., M. Hagiwara and H. Kitano, 2001 Robust oscillations within the interlocked feedback model of Drosophila circadian rhythm. Journal of theoretical biology 210**:** 401-406.

Ueda, H. R., A. Matsumoto, M. Kawamura, M. Iino, T. Tanimura *et al.*, 2002 Genome-wide transcriptional orchestration of circadian rhythms in Drosophila. Journal of biological chemistry 277**:** 14048-14052.

Umezaki, Y., T. Yoshii, T. Kawaguchi, C. Helfrich-Förster and K. Tomioka, 2012 Pigment-dispersing factor is involved in age-dependent rhythm changes in Drosophila melanogaster., pp. 423-432 in *J Biol Rhythms*.

Vaidya, A. T., D. Top, C. C. Manahan, J. M. Tokuda, S. Zhang *et al.*, 2013 Flavin reduction activates Drosophila cryptochrome. Proceedings of the National Academy of Sciences of the United States of America 110**:** 20455-20460.

Van Gelder, R. N., H. Bae, M. J. Palazzolo and M. A. Krasnow, 1995 Extent and character of circadian gene expression in Drosophila melanogaster: identification of twenty oscillating mRNAs in the fly head., pp. 1424-1436 in *Curr Biol*.

Van Gelder, R. N., and M. A. Krasnow, 1996 A novel circadianly expressed Drosophila melanogaster gene dependent on the period gene for its rhythmic expression., pp. 1625-1631 in *EMBO J*.

van Gend, C., and J. L. Snoep, 2008 Systems biology model databases and resources. Essays Biochem 45**:** 223-236.

van Renssen, A. S. H. P., 2005 *Gellish: A Generic Extensible Ontological Language - Design and Application of a Universal Data Structure*. Delft University Press, Delft, Netherlands.

Vandenbroucke, I. I., J. Vandesompele, A. De Paepa and L. Messiaen, 2001 Quantification of splice variants using real-time PCR. Nucleic Acids Res 29.

VanGuilder, H. D., K. E. Vrana and W. M. Freeman, 2008 Twenty-five years of quantitative PCR for gene expression analysis. Biotechniques 44**:** 619-626.

Vaux, D. L., 2012 Know when your numbers are significant. Nature 492**:** 180-181.

Venter, J. C., M. D. Adams, E. W. Myers, P. W. Li, R. J. Mural *et al.*, 2001 The sequence of the human genome. Science 291**:** 1304-1351.

Vilar, J. M. G., H. Y. Kueh, N. Barkai and S. Leibler, 2002 Mechanisms of noise-resistance in genetic oscillators. Proceedings of the National Academy of Sciences of the United States of America 99**:** 5988-5992.

Wager-Smith, K., and S. A. Kay, 2000 Circadian rhythm genetics: from flies to mice to humans. Nature Genetics 26**:** 23-27.

Wang, J., and T. Zhou, 2010 A computational model clarifies the roles of positive and negative feedback loops in the Drosophila circadian clock. Physics Letters A 374**:** 2743-2749.

Wang, T., and M. J. Brown, 1999 mRNA quantification by real time TaqMan polymerase chain reaction: Validation and comparison with RNase protection. Analytical Biochemistry 269**:** 198-201.

Waraporn, N., and K. Porkaew, 2008 Semantics of null values in subqueries. Imecs 2008: International Multiconference of Engineers and Computer Scientists, Vols I and Ii: 400-405.

Wasserstein, R. L., and N. A. Lazar, 2016 The ASA's Statement on p-Values: Context, Process, and Purpose. American Statistician 70**:** 129-131.

Weber, F. F., D. D. Zorn, C. C. Rademacher and H.-C. H. Hung, 2011 Post-translational timing mechanisms of the Drosophila circadian clock. Audio, Transactions of the IRE Professional Group on 585**:** 1443-1449.

Weeks, A. R., S. W. McKechnie and A. A. Hoffmann, 2007 Robust clines and robust sampling: a reply to Kyriacou et al., pp. 1652-1654 in *J Evol Biol*.

Weiss, R., O. Bartok, S. Mezan, Y. Malka and S. Kadener, 2014 Synergistic interactions between the molecular and neuronal circadian networks drive robust behavioral circadian rhythms in Drosophila melanogaster. PLoS Genet 10**:** e1004252.

White, E. P., E. Baldridge, Z. T. Brym, K. J. Locey, D. J. McGlinn *et al.*, 2013 Nine simple ways to make it easier to (re)use your data. Ideas in Ecology and Evolution 6**:** 1–10.

Wierling, C., R. Herwig and H. Lehrach, 2007 Resources, standards and tools for systems biology. Brief Funct Genomic Proteomic 6**:** 240-251.

Wijnen, H., F. Naef, C. Boothroyd, A. Claridge-Chang and M. W. Young, 2006 Control of daily transcript oscillations in Drosophila by light and the circadian clock. PLoS Genetics 2**:** e39-e39.

Wilcox, R. R., 2002 Comparing the variances of two independent groups. Br J Math Stat Psychol 55**:** 169-175.

Wilcox, R. R., 2012 *Introduction to Robust Estimation & Hypothesis Testing 3rd Edition*. Academic Press (Elsevier), Amsterdam.

Wilkinson, D. J., 2012 *Stochastic modelling for systems biology*. CRC Press/Taylor & Francis, Boca Raton.

Wilkinson, M. D., M. Dumontier, I. J. Aalbersberg, G. Appleton, M. Axton *et al.*, 2016 The FAIR Guiding Principles for scientific data management and stewardship. Sci Data 3**:** 160018.

Wilkinson, R. D., 2013 Approximate Bayesian computation (ABC) gives exact results under the assumption of model error. Stat Appl Genet Mol Biol 12**:** 129-141.

Willenbrock, H., J. Salomon, R. Sokilde, K. B. Barken, T. N. Hansen *et al.*, 2009 Quantitative miRNA expression analysis: comparing microarrays with next-generation sequencing. RNA 15**:** 2028-2034.

Wolkenhauer, O., 2001 Systems biology: the reincarnation of systems theory applied in biology? Brief Bioinform 2**:** 258-270.

Woodger, J. H., A. Tarski and W. F. Floyd, 1937 *The axiomatic method in biology*. The University Press, Cambridge Eng.

Wooley, J. C., and H. S. Lin, 2005 Computational Modeling and Simulation as Enablers for Biological Discovery in *Catalyzing Inquiry at the Interface of Computing and Biology*, edited by J. C. Wooley and H. S. Lin, Washington (DC).

Woolston, C., 2015 Psychology journal bans P values. Nature 519**:** 9-9.

Wu, Q. Q., K. Smith-Miles and T. H. Tian, 2014 Approximate Bayesian computation schemes for parameter inference of discrete stochastic models using simulated likelihood density. Bmc Bioinformatics 15.

Wulbeck, C., E. Grieshaber and C. Helfrich-Forster, 2009 Blocking endocytosis in Drosophila's circadian pacemaker neurons interferes with the endogenous clock in a PDF-dependent way. Chronobiol Int 26**:** 1307-1322.

Xie, Z., and D. Kulasiri, 2007 Modelling of circadian rhythms in Drosophila incorporating the interlocked PER/TIM and VRI/PDP1 feedback loops. Journal of theoretical biology 245**:** 290-304.

Xie, Z., D. Kulasiri, S. Samarasinghe and J. Qian, 2010 An Unbiased Sensitivity Analysis Reveals Important Parameters Controlling Periodicity of Circadian Clock. Biotechnology and Bioengineering 105**:** 250-259.

Xu, K., X. Zheng and A. Sehgal, 2008 Regulation of feeding and metabolism by neuronal and peripheral clocks in Drosophila., pp. 289-300 in *Cell Metab.*

Yerushalmi, S., and R. M. Green, 2009 Evidence for the adaptive significance of circadian rhythms. Ecology Letters 12**:** 970-981.

Yi, M., Y. Jia, Q. Liu, J. Li and C. Zhu, 2006 Enhancement of internal-noise coherence resonance by modulation of external noise in a circadian oscillator. Phys Rev E Stat Nonlin Soft Matter Phys 73**:** 041923.

Yin, H. S., H. G. Qi, J. W. Xu, W. N. N. Hung and X. Y. Song, 2014 Generalized Framework for Similarity Measure of Time Series. Mathematical Problems in Engineering.

Yoshii, T., T. Todo, C. Wülbeck, R. Stanewsky and C. Helfrich-Förster, 2008 Cryptochrome is present in the compound eyes and a subset of Drosophila's clock neurons. The Journal of comparative neurology 508**:** 952-966.

Yoshii, T., S. Vanin, R. Costa and C. Helfrich-Förster, 2009 Synergic entrainment of Drosophila's circadian clock by light and temperature., pp. 452-464 in *J Biol Rhythms*.

Young, M. W., and S. A. Kay, 2001 Time zones: a comparative genetics of circadian clocks. Nature reviews. Genetics 2**:** 702-715.

Yu, W., H. Zheng, J. H. Houl, B. Dauwalder and P. E. Hardin, 2006 PER-dependent rhythms in CLK phosphorylation and E-box binding regulate circadian transcription. Genes & development 20**:** 723-733.

Yu, W. W., J. H. J. Houl and P. E. P. Hardin, 2011 NEMO Kinase Contributes to Core Period Determination by Slowing the Pace of the Drosophila Circadian Oscillator. Current biology : CB 21**:** 6-6.

Zak, D. E., F. J. Doyle, D. G. Vlachos and J. S. Schwaber, 2001 Stochastic kinetic analysis of transcriptional feedback models for circadian rhythms. Proceedings of the 40th Ieee Conference on Decision and Control, Vols 1-5**:** 849-854.

Zeeberg, B. R., J. Riss, D. W. Kane, K. J. Bussey, E. Uchio *et al.*, 2004 Mistaken identifiers: gene name errors can be introduced inadvertently when using Excel in bioinformatics. BMC Bioinformatics 5**:** 80.

Zeigler, B. P., 2012 *Guide to modeling and simulation of systems of systems*. Springer, New York.

Zeigler, B. P., and P. E. Hammonds, 2007 *Modeling & simulation-based data engineering : introducing pragmatics into ontologies for net-centric information exchange*. Academic, Burlington, MA.

Zeigler, B. P., H. Praehofer and T. G. Kim, 2000 *Theory of modeling and simulation : integrating discrete event and continuous complex dynamic systems*. Academic Press, San Diego.

Zeng, H., P. E. Hardin and M. Rosbash, 1994 Constitutive overexpression of the Drosophila period protein inhibits period mRNA cycling., pp. 3590-3598 in *EMBO J*.

Zerr, D. M., J. C. Hall, M. Rosbash and K. K. Siwicki, 1990 Circadian fluctuations of period protein immunoreactivity in the CNS and the visual system of Drosophila., pp. 2749-2762 in *J Neurosci*.

Zhang, R., N. F. Lahens, H. I. Ballance, M. E. Hughes and J. B. Hogenesch, 2014a A circadian gene expression atlas in mammals: implications for biology and medicine. Proc Natl Acad Sci U S A 111**:** 16219-16224.

Zhang, Z., W. Zhu and J. Luo, 2014b Bringing biocuration to China. Genomics Proteomics Bioinformatics 12**:** 153-155.

Zheng, X., K. Koh, M. Sowcik, C. J. Smith, D. Chen *et al.*, 2009 An isoform-specific mutant reveals a role of PDP1 epsilon in the circadian oscillator., pp. 10920-10927 in *J Neurosci*.

Zheng, X., M. Sowcik, D. Chen and A. Sehgal, 2014 Casein kinase 1 promotes synchrony of the circadian clock network., pp. 2682-2694 in *Mol Cell Biol*.

Zielinski, T., A. M. Moore, E. Troup, K. J. Halliday and A. J. Millar, 2014 Strengths and Limitations of Period Estimation Methods for Circadian Data. Plos One 9.

Ziemann, M., Y. Eren and A. El-Osta, 2016 Gene name errors are widespread in the scientific literature. Genome Biol 17**:** 177.

Zinn, K., D. Dimaio and T. Maniatis, 1983 Identification of 2 Distinct Regulatory Regions Adjacent to the Human Beta-Interferon Gene. Cell 34**:** 865-879.

Zordan, M. A., and F. Sandrelli, 2015 Circadian clock dysfunction and psychiatric disease: could fruit flies have a say? Frontiers in Neurology 6.

Zweig, K. A., 2016 Section 11.8 Data responsibility and Data hygiene, pp. 350-361 in *Network analysis literacy: a practical approach to the analysis of Networks*, edited by K. A. Zweig. Springer Berlin Heidelberg, New York, NY.

Zwiebel, L. J., P. E. Hardin, J. C. Hall and M. Rosbash, 1991a Circadian oscillations in protein and mRNA levels of the period gene of Drosophila melanogaster., pp. 533-537 in *Biochem Soc Trans*.

Zwiebel, L. J., P. E. Hardin, X. Liu, J. C. Hall and M. Rosbash, 1991b A Posttranscriptional Mechanism Contributes to Circadian Cycling of a Per-Beta-Galactosidase Fusion Protein. Proceedings of the National Academy of Sciences of the United States of America 88**:** 3882-3886.

                     QQv1r1

TABLES OVERIEW – MAIN TEXT

TABLE 1. **Circadian clock components** referenced in this study as introduced by their
Brief, Explicit, Summarizing and Technical Name equivalents defined by
using the Evolvix BEST Names concept.

TABLE 2. **FlyClockbase and VBIR concepts and related keywords** as introduced by their Brief, Explicit,
and Summarizing Name equivalents defined by using the Evolvix BEST Names concept.

TABLE 3. **Data in FlyClockbase** counting independently observed D. melanogaster time series
summaries available for each clock component in the Summary Section of FlyClockbase.

TABLE 4. **Time series overview:** average (*Avg*), standard deviation (*SD)*, and number of time series
available (*n*) for peak and valley times of some mRNA and proteins in FlyClockbase.

TABLE 5. **Human error rates** for four *Mod5 PeakValleyTables*.

TABLE 6: **Timing distribution differences** tested between *per* and *tim* to compare averages and
variances of peak and valley times for proteins and mRNAs after removing outliers.

TABLE 7: **Impact of outlier removal** approaches on differences
in variance of protein peak times (PER – TIM).

TABLE 8. **Standardization proxies**: overview of potential approximations for quantifying method detail
standardization and method detail similarity for independent observations of *per* mRNA.

FIGURES OVERIEW – MAIN TEXT

FIGURE 1. **Systems biology overview** of the circadian clock in *D. melanogaster*.

FIGURE 2. **Overview of FlyClockbase** organization and data model.

FIGURE 3. **Workflow for refining** the time series traits reported in peak-valley-tables.

FIGURE 4. **Where do models get their realism from?** Overview of data sources reported.

FIGURE 5. **Data accumulation over time**: overview of experimental studies available in FlyClockbase
and their use for parameter estimation over time.

FIGURE 6. **Hypotheses on timing variances in PER and TIM** as tested by FlyClockbase data.
FlyClockbase extends our understanding of circadian clocks in flies by enabling observations of
significant differences in variance between peaks and valleys of different clock components.

FIGURE 7. **Histograms for *per* and *tim* mRNA and protein** time series traits like peaks and valleys as
observed in independent experiments show how FlyClockbase enables comparing variability.

FIGURE 8. **Comparisons of PCR vs non-PCR methods** for measuring amounts of *per* mRNA in time
series as enabled by FlyClockbase.

FIGURE 9. **Referenced protocol use similarity matrix** for studies reporting *per* mRNA time series,
grouping protocols by measurement approach.

FIGURE 10: **Logic of datasets in statistics**: example for an interaction of statistics and formal logic
where the logic for selecting datasets unnecessarily reduced the power of a statistical test.

FIGURE 11: **Biological model curator and compiler architect collaboration model** for improving the
integration of biological data into *VBIR*s while constructing the type system required by a
VBIR compiler for handling the ubiquitous uncertainty of biological data more precisely.

RELATED ANALYSES

Hypotheses comparing timing in PER and TIM: data and analyses in Table 4, Table S1, and Table 6,
Table 7, Figure 6, Figure 7, as well as parts of the Supplemental Statistical Analysis are related.

Comparing PCR vs non-PCR methods for measuring time series in per mRNA: data and analyses in
Table 8, Figure 8, Figure 9, Figure 10, and parts of the Supplemental Statistical Analysis are related.

2 **TABLE 1. Circadian clock components referenced in this study**

3 **as introduced by their Brief, Explicit, Summarizing and Technical Name**

4 **equivalents defined by using the Evolvix BEST Names concept.**

Brief Name	Explicit Name	Summarizing Name (description)	Technical Name FlyBase ID
clk	clock	Promotes transcription of *per*, *tim, pdp1, vri, cwo*	FBgn0023076
per	period	Binds TIM, inhibits CLK action	FBgn0003068
tim	timeless	Binds PER, inhibits CLK action, degraded in response to light	FBgn0014396
cry	cryptochrome	Activated by light, promotes TIM degradation	FBgn0025680
cwo	clockwork orange	Stabilizes "jitters" (minor variations) in the clock	FBgn0259938
pdp1	PAR-domain protein 1	Promotes transcription of *clk*	FBgn0016694
vri	vrille	Inhibits action of PDP1	FBgn0016076
pdf	pigment-dispersing factor	Synchronizes different clock neurons	FBgn0023178
sgg	shaggy	Phosphorylates TIM, promotes nuclear localization	FBgn0003371
jet	jetlag	F-box protein, contributes to TIM degradation	FBgn0031652

 QQv1r1

ck2a	casein kinase 2 alpha	Phosphorylates PER, promotes nuclear localization	FBgn0264492
dbt	doubletime	Phosphorylates PER, promotes degradation	FBgn0002413
nmo	nemo	Phosphorylates PER, promotes further phosphorylation	FBgn0011817
slmb	slimb	F-box protein, contributes to PER degradation	FBgn0283468
cyc	cycle	Binds to CLK	FBgn0023094
cu, noc	curled, nocturnin	Implicated in light response, metabolism, mRNA degradation	FBgn0261808

Brief Names and Explicit Names represent abbreviations and full common names, respectively. Summarizing Names provide a brief description, and Technical Names correspond to hyperlinked FlyBase Identifiers. In the text, we follow the convention of denoting the proteins in CAPITAL letters and mRNAs or genes in *italics* (hence the use of italics for the brief names here). More about the way synonymous dialects (defined by the Evolvix BEST Names concept) can help navigate biological naming complexity can be found elsewhere (LOEWE et al. (2014; 2016a) and LOEWE et al. (2016b) for a more in-depth discussion of naming).

TABLE 1 END

TABLE 2.   FlyClockbase and *VBIR* concepts and related keywords

as introduced by their Brief, Explicit, and Summarizing Name

equivalents defined by using the Evolvix BEST Names concept.

Overview of concepts by naming gives the *Brief* and *Explicit* Names used in the main text for important

FlyClockbase data structures (always in *italics*, usually capitalizing the beginning of significant words).

Their *Summarizing* Name gives a description that could be turned into a rather long very descriptive

name. The nature of these names is defined by the Evolvix BEST Names concept, see

http://evolvix.org/naming and LOEWE *et al*. (2016b). Order of entries loosely follows conceptual relations.

See main text for more details.

Brief Name	Explicit Name	Summarizing Name (description)
	Italicized ProperNames	*Brief* and *Explicit* names of FlyClockbase components are treated both as variables (hence *italicized*, indicating flexible content) and as proper nouns (hence Capitalized to contrast their special role with the generic role given by their name). In program code *ConcatenateAllWords* without spaces; our main text may add *SomeSpaces BackIn* for readability.
VBIR	*Versioned Biological Information Resource*	Systematic collection of biological information for a specific purpose, irrespective of technology used to implement it. Like FlyClockbase, *VBIRs* usually need to cope with much uncertainty, contradictions, gaps, and numerous exceptions in biological data. Their versioning indicates that they store biological information using controlled immutable increasing versioning numbers that label each publicly released *VBIR* variant and help ensure that previously released data remains accessible under that label.

POST	*POSTsystem*	Project Organization Stabilizing Tool system is at the very foundations of organizing files in FlyClockbase, Evolvix, and TabFS and Evolvix. Stable online reference: http://evolvix.org/post
StabZ	*StablizingZone*	Backbone of a stabilizing versioning number system defined by the POST system of Evolvix; represented by the double capital letters ranging from *MM* to *TT* (from *MockupModel* to *TrustedTested*). See Supplemental Material Table P1.
TabFS	*TabFS*	Tab-delimited table text file system for organizing the basic file layout of FlyClockbase and Evolvix
Sec	*Section*	Topical area in a *VBIR*.
SumS	*SummarySection*	Stores summarized statistics of published time series in FlyClockbase
DetS	*DetailsSection*	Stores all details of non-summarized individual raw time series submissions in FlyClockbase to enable independent statistical analyses.
H_{basic}	*BasicNullHypothesis*	See main text Results for a full definition.
Obs	*Observation*	*Obs* in FlyClockbase refers to biological observations that are closer to real-world biology than derived analyses (e.g. estimates, results). *Obs* stands in direct contrast to *Simulations* (*Sim*, may try to mirror a given *Obs*; see http://evolvix.org/post for more details).
ObsRaw	*ObservationRaw*	*Obs* in the form of *Raw* data as received (use versioning to correct trivial errors if needed; no additional modifications allowed).
ObsMod	*ObservationModified*	Any raw *Obs* modified before further processing, including the removal of obvious outliers (or any changes to input data that could be questioned). *ObsMods* start all with *ObsRaw*, and distinguish each modification by appropriate versioning, such as *ObsMod6*, originating from *ObsMod5*.
ObsOdd	*ObservationOdd*	Any *Obs* classified as unusual, extreme or unexpected. Conceptual opposite of *ObsMaj*. Where possible store processing status as in *ObsRawOdd*, or *ObsMod1Odd*.
ObsMaj	*ObservationMajority*	Any *Obs* classified as usual, typical or expected. Conceptual opposite of *ObsMaj*. Where possible, store processing status as in *ObsMod6Maj* (*Majority* of *Mod6* of *Obs*).

Tbl	*Table*	A fundamental data structure in FlyClockbase; each *Tbl* is to be stored as a separate tab-delimited file that defines *Rows* and *Columns* and can be edited by spread-sheet software (see FlyClockbase for format details). *Tbls* for *Content*, *Attributes*, and *Traits* are very different.
Col	*Column*	Minimal vertical selection of a *Table*, including each *Row*; first *Row* is *ColumnHeader*.
Row	*Row*	Minimal horizontal selection of a *Table*, including each *Col*; first *Col* may be *RowHeader* or *Time*, or an immutable *TableIndex* for unambiguously identifying each *Row*.
IDX	*IDIndex,* *TableIndex*	Identifier index column in a table of TabFS or FlyClockbase, where the type of index is usually specified by a sequence of additional suffixes.
ID	*Identifier*	An unambiguous label (= name) that can be used in its context to refer to specific content it denotes.
IDL	*IDLocal*	Atomic identifier, local and thereby relative to its local context, as defined elsewhere. This shortest of all IDs is given by a single positive integer.
IDF	*IDFragment*	An *ID* fragment requires context (for its first=top *IDL*), but also specifies contexts for each *IDL* it contains in its path of local IDs (separated by ".")
IDM	*IDMemory*	Identifier guaranteed to refer to a unique item within a defined memory area, such as FlyClockbase. It can easily be made globally unique by prefixing it with the *IDG* of its memory area.
IDG	*IDGlobal*	Identifier that is as globally unique as possible (see figure 3 in LOEWE *et al.* 2016b).

Itm	*Item*	Any entity that provides a *Context* for its *Content*, and that can itself be *Content* in some other *Context*. This makes *Items* nestable. *Items* can also have Attributes and Traits, which complement *Content*, albeit without being *Content.*
Cxt	*Context*	*Context* belongs to an *Item* and contains *Content*, *Attributes*, and *Traits.*
Cnt	*Content*	Any container for directly storing 'the data itself', describing data item(s) of primary interest; contains neither *Attributes* nor *Traits*. *Content* is always stored in some *Context.*
Att	*Attribute*	Any container for storing 'data about data (meta-data)', describing the non-computable type and value of a fragment of information inherent to the content of a given item; contains no *Traits.*
Tra	*Trait*	Any container for storing 'data derived from data', describing a defined type and computable value that capture an emergent feature derived from *Content* and *Attributes.*
Ref	*Reference*	*Refs* in FlyClockbase, *TabFS* or Evolvix are always bibliographic references ready for citing in a paper.
CntTbl	*ContentTable*	Frequently used *Type* of exclusively *Content*-storing *Table* (e.g. for *TimeSeries*); there exists always a related *AttributeTable* for storing newly learned Attributes.
AttTbl	*AttributeTable*	Set of *Attributes* for a given *ContentTable*. This Type of *Table* stores exclusively *Attributes* and always refers to a *ContentTable.*
TraTbl	*TraitTable*	*Table* storing exclusively *Traits.* Always refers to a *ContentTable* (and its *Attributes* and code as source).
	SearchResult	Set of *Tables* with *Content* and *Attributes* copied and *Traits* computed from FlyClockbase

BBin	*BioBinary*	A *BBin* stores 2 bits to distinguish 4 states enumerated by the *OKScale* (*OK*, *OKO, KO, MIS*), designed for statements in biology, where *FullyTrue* (*OK*) or *FullyFalse* (*KO*) are less appropriate than *Intermediate* (*OKO*), or assuming applicability is a *Mistake* (*MIS*).
OK, OKO	*FullyTrue, Intermediate*	see *BBin*
KO, MIS	*FullyFalse, Mistake*	see *BBin*
OKS	*OKScale*	see *BBin*
TS	*TimeSeries*	*Content* data type central to FlyClockbase; stored in a *Table* with at least one *Column* for *Time* and one for an *Obs* of the amount of molecules specified in the *ColumnHeader*
ZT	ZeitgeberTime	A time measure used in circadian clock research to denote the time that passed since the last dawn (i.e. Lights went "on"). Any event during the day is then measured in terms of decimalized hours after ZT = 0*h*. Range: 0*h* < ZT ≤ 24*h.*
DZT	*DailyZeitgeberTime*	Identical to ZT. *DZT* replaces ZT in FlyClockbase and in this study. Its shows a decimalized *hour* of the day but hides its counter of days. *DZT* was defined to unmistakably refer to the circular nature of *hours* / day. Range: 0*h* < *DZT* ≤ 24*h.*
CZT	*ContinuousZeitgeberTime*	Almost identical to *DZT*, but not quite. Both appear the same for the first 24*h*. They start diverging at the beginning of day 2 = end of day 1, when *DZT* time is "reset" to 0*h*, while *CZT* progresses steadily: …, 23*h*, 24*h*, 25*h*, … as measured in decimalized *hours.* Range: 0*h* < *CZT* < unlimited *h*.
	Peak	Daily time of highest amount of a component.
	Valley	Daily time of lowest amount of a component.
	PeakValleyTables	The specific *SearchResultTable* compiled manually from FlyClockbase for testing hypotheses explored in this study.
h	*hours*	Default unit for measuring time in this study. Applies to *Peak*, *Valley*, *CZT*, *DZT*, and ZT. Here a day is defined by 24*h*. Fractions of hours are decimal; FlyClockbase does not recognize minutes (e.g.: 0.9*h* = 54min; 1.5*h* = 90 min; etc).
-	*comvar2*	Robust statistical test for measuring the significance of differences in variance using bootstrap methods (albeit only for a *P*-value of 5%; see Statistical Methods for details).

Avg	*Average*	Averages default to the arithmetic mean
SD	*StandardDeviation, StDev*	Typically shown in plots of this study: the standard deviation inferred from *n* data points which are assumed to follow a Normal distribution; *Var* = SD^2
Var	*Variance*	Tested for significant differences by *comvar2*: the variance inferred using the same assumptions as for SD. *Var* = SD^2
n	*NumberObserved*	Number of observations in a given set.
SEM	*StandardError_of_Mean*	Sometimes used in confusing ways to obscure variability. Ideally, all individual replicates are reported to enable reanalyzes; else quantiles or other summary statistics may aggregate values of the actual distribution of interest. SEM is not a measure of the data, but estimates errors of a summary statistic of the data; giving it a '3rd hand quality'. Some authors further degrade meta-analysis power by confusing SD and SEM (SALSBURG 1985; VAUX 2012).

CI	*ConfidenceInterval*	Denotes a credible range of values for a point estimate. A *CI* is defined by repeating a point estimate procedure (infinitely) many times and creating a histogram of the results. In that histogram the range of values of the given point estimates that a given percentage of values is expected to take, is defined as a *CI*. A *CI* is defined incompletely without
		  - the percentage of all estimates it is to contain (e.g. 95% or 0.95 when rejecting 2.5% of correct values on both sides),
		  - type information about the point estimate (e.g. *Avg,* or *Var*),
		  - specifying what is being replicated while measuring the *CI* (experiments, bootstrap samples, etc.).

Min	*ValueMinimum*	Guaranteed minimum of all values of a type.
Lo	*ValueCredibleLow*	Lower limit for a value that is still credible.
Mi	*ValueCredibleMiddle*	Denotes the most credible *Middle* point value. The type of this value defines how to compute credible values for *Lo*, *Mi* or *Hi*. Hence, this trio can generally be reused for quantifying distributions of any type, including those implicitly defined by the output of a computational procedure or human intuition.
Hi	*ValueCredibleHigh*	Higher limit for a value that is still credible.
Max	*ValueMaximum*	Guaranteed maximum of all values of a type. The type defines whether observations or possibilities are to be considered.

EMU	*ErrorMeasureUnknown*	An abstract error statement type for cases where nothing is known that could possibly limit potential errors. Used as default for unique *Obs* without error estimates. *EMU* is the equivalent to *Nothing* when it comes to making a statement about the *limits* of errors. *EMU* states that no information is available that might reduce the range of credible errors, not circumstantially, statistically, contextually, conceptually, numerically, or in any other way. *EMU* errors may be infinite (since unknown!), but they instantly collapse when estimating or observing actual errors in the real world. Under this logic, values with *no* attempt to quantify uncertainty are infinitely uncertain (not infinitely certain as often assumed). *EMUs* reward attempts to engage with uncertainty by estimating errors indirectly or by observing them directly when recording as many independent *Obs* replications as feasible.
-	*Nothing*	Abstract type that is foundational to all other types that indicate the absence of a particular thing, idea or activity.
-	*NotGiven*	A special type of *Nothing* dedicated to representing that (i) a certain observation must have existed and (ii) could in principle have been recorded, but (iii) for whatever reason, the information was *NotGiven*. To reduce ambiguity in FlyClockbase, we redefined symbols that are commonly used in ambiguous ways to represent the resulting contradiction in type. Banning such errors simplifies recognizing such problems when importing data. For example, does 'NA' stand for 'NotAvailable' or 'NotApplicable'? Does '0' indicate absence or is it a measure of time using cyclical or linear numbers? What type is implied in 'NULL'?
-	*MethodRealm*	Specifies in which of the broad biological realms of observation a given time series was measured *(in vitro, in vivo, ex vivo, post mortem)*

TABLE 2 END

1 **TABLE 3.**   **Data in FlyClockbase counting independently observed**

2   ***D. melanogaster* time series summaries available for each clock**

3   **component in the Summary Section of FlyClockbase.**

Clock Component	Time Series [*n*]	Studies [*n*]
per mRNA	89	42
PER protein	77	26
tim mRNA	51	27
TIM protein	42	16
clk mRNA	33	18
vri mRNA	31	14
cry mRNA	18	10
CRY protein	16	6
pdp1 mRNA	15	10
CLK protein	10	5
cwo mRNA	5	3
PDF protein	4	2
VRI protein	2	2
jet mRNA	2	1
dbt mRNA	2	2
CYC protein	2	1
cyc mRNA	1	1
sgg mRNA	1	1
ck2 mRNA	1	1
PDP1 protein	1	1
Total	**403**	**86**

5   TABLE 3 END

1   **TABLE 4.**   **Time series overview:**

2   **average (*Avg*), standard deviation (*SD), and number of time series available (*n*)**

3   **for peak and valley times of some mRNA and proteins in FlyClockbase.**

Component	Peak		Valley	
of the Clock	*Avg* ± *SD* [*h*]	*n*	*Avg* ± *SD* [*h*]	*n*
per mRNA	14.64 ± 1.65	88	3.61 ± 2.60	24
tim mRNA	14.54 ± 1.92	48	4.86 ± 1.52	21
TIM protein	18.41 ± 2.54	37	6.33 ± 3.18	35
PER protein	19.50 ± 3.40	33	9.27 ± 2.10	40
clk mRNA	2.79 ± 3.16	12	14.33 ± 1.67	30
CLK protein	11.57 ± 3.18	5	12.51 ± 4.30	9
CRY protein	23.02 ± 1.59	15	9.96 ± 3.73	16
cry mRNA	6.77 ± 2.96	11	16.23 ± 2.54	17
pdp1 mRNA	14.70 ± 1.81	15	4.68 ± 1.88	5

6   A clock component is listed if at least four different time series values exist for both peak and valley.

7   Statistics were computed for *Mod6* modified *PeakValleyTables* (see Figure 3) without removing outliers.

8   All times in decimal fractions of hours have been linearized such that *CZT* (given) is effectively equivalent

9   to a corresponding *DZT* = ZT. For related analyses see also Table S1, Table 6, Table 7, Figure 6,

10   Figure 7, and parts of the Supplemental Statistical Analysis.

12   TABLE 4 END

TABLE 5.   **Human error rates for four *Mod5 PeakValleyTables*.**

Clock Component TimeSeries counted	Errors counted in FlyClock-base	*Peak* TraitTable		*Valley* TraitTable	
		All *Peaks* counted	**Errors** counted	**All *Valleys*** counted	**Errors** counted
per mRNA (88)	2	87	7	24	6
tim mRNA (49)	1	48	2	20	0
TIM protein (43)	1	29	4	34	6
PER protein (57)	1	30	12	39	8

To approximate an error rate for *PeakValleyTables*, we chose the four *Mod5* tables of the most abundant clock components and re-examined the timing of each peak and valley in-depth. All errors were corrected in the *Mod6 PeakValleyTables*. Not all time series in FlyClockbase were long enough to include both a peak and a valley for day 1 (474 expected vs 311 observed peaks as shown in Figure 6). While the error rate for FlyClockbase is reasonably low for the columns tested (5 errors in at least about 5 effective columns in the big *TimeSeries AttributeTable* suggest an overall rate of at most about 5 / (5 * (88+49+57+43)) = 0.42% per cell; rule of thumb: ca 1% for trivial errors), manually exporting and computing *TraitTables* resulted in much higher error rates. Most of these are caused by the inconsistent manual application of technical procedures that had been agreed upon when analyzing the traits of highly heterogeneous time-series in their broader context (the 25 peak errors and the 20 valley errors resulted in about 45 / 311 = 14% errors; rule of thumb: > 10% for errors from forgetting one of a few rules). Ultimately, it is desirable to construct *PeakValleyTables* automatically via script. However, it is not clear whether this necessarily reduces the number of errors or the amount of time to achieve a given accuracy goal, unless data structures are rather uniform and/or the same errors reoccur often with only minor variations. A large number of procedural errors and their extreme heterogeneity can greatly complicate the (manual) implementation of code that needs to recognize each type of error in order to address it.

TABLE 5 END

TABLE 6.   **Timing distribution differences tested between *per* and *tim***

to compare averages and variances of peak and valley times

for proteins and mRNAs after removing outliers.

	Point distribution CZT Observed [*h*]		Differences Distribution of *CZT* point distributions [*h*]			Test
	per	*tim*	*Lo* 2.5%	*Middle* 50%	*Hi* 97.5%	*P*-Value
Protein peak						
Var	4.1377	0.89211	0.97136	3.2456	5.6033	*** (*P* < 5%)**
Avg	18.644	18.915	−0.99996	−1.2345x10^{-5}	0.99994	0.98753
n	91% =30/33	78% =29/37				
mRNA peak						
Var	2.4877	4.4906	−4.1225	−2.0029	−0.0035509	*** (*P* < 5%)**
Avg	14.606	14.587	−0.83733	4.3138x10^{-6}	0.99996	0.66505
n	99% =87/88	100% =48/48				
Protein valley						
Var	3.7568	6.3773	−6.8437	−2.6205	1.3534	(*P* > 5%)
Avg	9.4058	5.8421	3.00	4.00	5.00	**1.489x10^{-7}**
n	100% =39/39	94% =34/36				
mRNA valley						
Var	5.3608	1.3215	0.17418	4.0393	8.3148	*** (*P* < 5%)**
Avg	3.606	5.0894	−2.0611	−1.00	−1.8909x10^{-5}	**0.012835**
n	100% =24/24	100% =20/20				

Removal procedures have been described elsewhere (CARLING 2000; WILCOX 2012), see our approach

(iii) in Statistical Methods, and *Mod6* in Figure 3 (this table and Figure 6 are based on the same data).

Abbreviations: *Var* = *Variance* calculated for data sampled from a given distribution to quantify spread;

Avg = *Average* = arithmetic mean calculated for a sample to quantify the location of a distribution; *n* =

numbers of observations without/with outliers; *P* = probability of rejecting the null-hypothesis that the point

distributions observed for *tim* and *per* are drawn from the same underlying distribution according to an

unpaired, two-sided Mann-Whitney-U test for *Avg*, and the "*comvar2*" method with 10^5 bootstraps from

Wilcox (2012) for *Var*, as described in Materials and Methods (significant differences are highlighted in

bold). The additional digits in this table do not necessarily reflect accuracy. They were kept to simplify

searching for their context in the Supplemental Statistical Analysis (searches are fast, even though not all

mappings are unique; trailing zeroes were omitted by R and thus added back to four of the numbers;

some digits are greyed out to reduce visual clutter). See main text for discussion and consequences of

adding outliers back into this table. For related analyses see also Table 7, Figure 6, Figure 7, Table 4,

Table S1 and parts of the Supplemental Statistical Analysis.

TABLE 6 END

1 **TABLE 7.**   **Impact of outlier removal approaches on**

2   **differences in variance of protein peak times (PER – TIM).**

Outlier removal approach	Outliers counted / All data	PER-TIM Variance *Lo* 2.5% CI	PER-TIM Variance *Mi* 50%	PER-TIM Variance *Hi* 97.5% CI	Significant at *P* = 0.05
No outliers removed	0/33 (PER) 0/37 (TIM)	**-8.308**	4.7564	15.755	*no*
Removal type (ii): Manual extremes	3/30 (PER) 1/36 (TIM)	**-1.7526**	1.3432	4.2201	*no*
Removal type (iii): Carling (2000)	3/30 (PER) 8/29 (TIM)	0.97136	3.2456	5.6033	*yes*

Manual removal of four extremely different time series (row (ii)) substantially narrows the distribution of

differences in variance seen prior to removal. However, this does not yet achieve the statistical

significance seen when applying the Carling (2000) outlier removal approach (iii) as used in Table 6.

Each row reports one comparison that calculates *Var*(PER)-*Var*(TIM) using "*comvar2*", 10^5 bootstraps;

see Statistical Methods Section. As seen in rows 1-2, if the *CI* includes 0, the test indicates that the

variances of both distributions are drawn from a common distribution with a probability of 5% or more. For

related analyses see also Table 6, Figure 6, Figure 7, Table 4, Table S1 and parts of the Supplemental

Statistical Analysis.

13   TABLE 7 END

TABLE 8.   Standardization proxies: overview of potential approximations for

quantifying method detail standardization and method detail similarity

for independent observations of *per* mRNA.

Measurement method	Count of time series / studies	Any method refs ≥ 1 given for time series / given for studies	Shared method refs ≥ 1 given for time series / given for studies
PCR	**25** / 13	**72% (18 / 25)** / 54% (7 / 13)	**17% (3 / 18)** / 29% (2 / 7)
RPA	**45** / 24	**100% (45 / 45)** / 100% (24 / 24)	**93% (42 / 45)** / 88% (21 / 24)
Northern Blot	**9** / 5	**56% (5 / 9)** / 60% (3 / 5)	**40% (2 / 5)** / 67% (2 / 3)
Microarray	**4** / 3	**0% (0 / 4)** / 0% (0 / 3)	*not applicable* / *not applicable*
RNA-Seq	**4** / 1	**100% (4 / 4)** / 100% (1 / 1)	*not applicable* / *not applicable*

The count column specifies the number of time series or studies that used the general method type

specified for the row. We approximated method standardization with a binary variable indicating whether

or not any method references were cited in the paper. Here we assumed that papers who did not cite any

method paper either adapted an existing method or did not closely follow a given set of instructions for

other reasons (see text for implications). To approximate the similarity of methods that produced time

series, we used another binary variable to indicate whether or not a given study shared at least one

method reference with any of the other studies that cited at least one method reference. We assumed

that shared references might indicate the use of more similar methods, which would result in more similar

observations compared to 'time series from papers that had only non-shared method references' (but still

required at least one method reference to be included). Note the subtle difference in logic when

considering a comparison to 'time series from papers that did not give shared method references' (which

includes those with no method references at all). See Figure 10 for a visualization of the corresponding

sets and Figure 9 for the raw data on protocol use, which underpins this analysis.

TABLE 8 END

FIGURE 1

FIGURE 1. Systems biology overview of the circadian clock in *D. melanogaster*. This simplified

diagram shows the basic components of the *D. melanogaster* circadian clock. In one feedback loop, CLK

binds CYC and promotes the transcription of *per* and *tim*. The proteins PER and TIM then form a complex

and repress this effect of CLK. Light activates CRY by inducing a conformational change (denoted by *),

and CRY and SGG work with JET to promote the degradation of TIM. Without TIM, PER is

phosphorylated, interacts with SLMB, and is degraded. In a second feedback loop, the CLK/CYC complex

promotes transcription of *pdp1, vri, and cwo*. PDP1 promotes transcription of *clk*, and VRI inhibits the

action of PDP1. CWO weakly inhibits CLK-promoted transcription of *per*, *tim*, *vri*, *pdp1,* and itself. The

notation in this diagram uses Systems Biology Graphical Notation Process Description level 1 version 1.3

(Moodie *et al.* 2011), with minor modifications.

1   **FIGURE 2**

2
3
FIGURE 2. Overview of FlyClockbase organization and data model. We extracted as much raw data

and as many reported experimental details as possible from plotted time series summary figures and their

publications. We stored bibliographic information in the *References AttributeTable* (Step S3) and stored

figures as files ready for plot digitizing (Step S1). For convenient reanalysis, we included all files of open

access studies in FlyClockbase. From each figure with appropriate Attributes we extracted the

FlyClockbase *TimeSeries Content* (*TS*) itself as *RawObservations* (*ObsRaw*, Step S2). *TS* report some

measure of the number of molecules of a given clock component as observed at a given time and in a

given volume. The volume and other scaling factors are usually unknown and may be non-linear on

occasion. Here we assume that amounts of *Obs TS* indicate a measure of relative abundance among

neighboring amounts of the same given *TS*. To place each *TS* in its appropriate context, we collected as

many *TS Attributes* as we could reasonably extract from the experimental details given in the original

study (Step S4). These *Attributes* are stored in the *SumS TS AttributeTables.* To obtain the more refined

TS ContentTables ObsModified (*ObsMod*), we rechecked the data based on information in its associated

publication (such as exact times of data collection, Step S5). Time series in *SumS ObsRaw* were not

refined into *ObsMod* if they were shorter than six hours or if more than half of their values were missing or

unreadable. All other *TS* were copied and modified into *ObsMod* and used with their respective *Attributes*

for further analyses, ultimately aiming to produce a retrospective meta-analysis. To export the relevant

data from FlyClockbase requires a *SearchResult*, which includes the specified *Content*, *Attributes*, and

computed *Traits* (stored in *TS TraitTables*)*.* Such *SearchResults* could be compiled manually or

automatically and are stored in a *TableSet*, which can include *TS TraitTables*. For the retrospective meta-

analysis presented here we manually compiled *TS TraitTables* reporting one peak and one valley from

each *ObsMod TS* measured in *CZT* (Steps S6, S7, S8). These peak-valley *TS TraitTables* are further

processed in Figure 3. *Content* and *Attributes* for >400 *TS* derived from published studies are stored in

the *Summary Section* (*SumS*), which does not include raw experimental data of separately observed *TS.*

These will be stored in *DetS*, the *Detail Section*, which is planned as an extension that will enable the

submission and processing of more detailed and less summarized *ObsRaw* TS *Content* in order to enable

researchers to compute their own independent summary statistics.

FIGURE 3

FIGURE 3. Workflow for refining the time series traits reported in peak-valley-tables. These tables

represent a *SearchResult* (as in Figure 2) and contain a special set of *Traits* and *Attributes* extracted from

FlyClockbase. In this study we extracted the times associated with the maximum ("*Peak*") and minimum

("*Valley*") amounts of the first each day of each time series. Each row of a peak-valley table contains a

peak observation and a valley observation for one day of one time series, along with *Attributes* of interest.

One peak-valley table was produced for each clock component of interest.

11   *Refinement of outliers* (*Mod1-Mod5*). The tables for each component formed a set that was

refined together (from *Raw➔Mod1➔*…etc.). *Raw PeakValleyTables* show data for every day of the time

series, while all *Modified* ("*Mod*") *PeakValleyTables* only show data for the first day of the time series. To

start identifying outliers and potential errors in these tables, we searched for clock components in which

the *SD* for peak, valley, or both exceeded 6 hours. We examined these clock components more closely

and corrected errors to create the first collection of modified *PeakValleyTables* ("*Mod1*"). We chose *SD* =

6 as initial cutoff value, since it approximates a uniform random distribution over all 24 hours (to us a sign

of problems in correctly identifying the right values. For each modified peak-valley table in *Mod1*, we then

calculated the *SD* and mean for peaks and valleys. Individual time series observations outside of the

mean $\pm$ 1 *SD* were noted for further investigation and recorded in a peak-valley table reserved for these

"*Odd*" values ("*Odd1*"). We examined *Odd* values for potential errors that we knew could be caused by

data handling in FlyClockbase and corrected them appropriately to create the next set of modified

PeakValleyTables ("*Mod2*"). We repeated this process until we created *Odd5* and *Mod5*. Refinement up to

this point was geared towards checking statistical outliers for errors in handling data or determining traits

(in contrast to our error analysis below).

Human Error Analysis Context. When the refinement process was first mapped out, the plan was

to stop at *Mod5* and use this repeatedly refined dataset for calculating the final analyses of PER and TIM.

Since the curators were diligent motivated researchers aware of potential errors and carefully working to

avoid them, there was no expectation that substantial numbers of errors could still lurk beneath the

surface. Due to the importance of our conclusions for modeling the differences in stochasticity of PER and

TIM we decided to still review all FlyClockbase data before finalizing the statistical analysis of the main

result of this paper. We started our review with several random tests, manually conducted on the spot.

These detected enough errors to justify a more systematic approach. We decided to avoid shortcuts and

focus on an exclusive complete in-depth re-review of all those time series in FlyClockbase where errors in

our data handling or trait computation could have affected our conclusions about the peak or valley timing

of *per* mRNA, *tim* mRNA, *PER* protein, and *TIM* protein.

Errors tested included the following notable cases. (i) Peaks and valleys from maxima and

minima at the margins of brief time series were probably not genuine, as they would have been unlikely to

coincide with the start or stop of time series observations. Excluding these substantially reduced the

variance we observed. (ii) The first day was non-representative in some important respect. If peak or

valley times during day 1, our usually monitored interval, were noticeably different from subsequent days,

we concluded that measurements probably started too early and used the peak and valley from a day that

followed closely. (iii) On rare occasions fields in FlyClockbase had been swapped or misprocessed in

some way.

Benefits. While laborious, this effort payed off in multiple ways (see Results and Discussion): (i) It

greatly increased confidence in our PER-TIM peak time variance analyses; see Results. (ii) We manually

identified particularly extreme outliers not caused by errors on our side. This motivated us to analyze their

impact by defining 'outlier removal approach (ii)' and choose more systematic outlier removal approaches;

see Statistical Methods; see Discussion for potential implications. (iii) FlyClockbase now has an internal

human error analysis, see Results. Error rates might be extrapolated to non-*per*/*tim* components or other

VBIRs of comparable complexity. Currently not many opportunities exist for measuring real-world human

error rates in advanced data organization scenarios. (iv) Valuable lessons in automation and *VBIR*

compiler construction were learned as a compiler architect worked closely with expert curators during

manual error identification. Such manual work is necessarily the first step towards automation of such

checks in a corresponding *VBIR* compiler infrastructure as envisioned in the Discussion.

Datasets of interest. Mod6 is the most refined set of observations with the highest quality

produced by this study and was used as the basis for all biological analyses presented here. It was

produced by correcting in *Mod5* all errors in the data handling or trait calculation of *per*, *tim*, PER or TIM,

while copying all data for other clock components unchanged from *Mod5*. The dataset *ObsMod7* is

essentially identical to *ObsMod6,* except for replacing a few cells in the table that were not analyzed here,

but caused problems when R parsed the file. *ObsMod8* is identical to *ObsMod7*, except for manual

removal of the manually identified outliers (see outlier removal approach (ii) in Statistical Methods).

ObsMod7 and ObsMod8 are provided with the R-source-code archive in the Supplemental Material.

FIGURE 4

FIGURE 4. Where do models get their realism from? Overview of data sources reported.

Surprisingly few experimentally observed time series are used for estimating parameters in *D.*

melanogaster circadian clock simulation studies. We reviewed 66 modeling studies identified here by

AuthorYear. Only three of these did estimate parameters directly from full time series observed in

experiments (using only 14 of the 86 experimental studies included in FlyClockbase). These three studies

(and one that used their results) are denoted on the left by rectangles and rhombuses. The pie chart

above them illustrates how many experimental studies were used (blue, solid line, 14) or ignored (pink,

dashed line, 72) by modelers. The 13 studies represented by pentagons chose the simpler (but not

simple) approach of searching for clock parameter combinations that satisfied some of the abstract time

series requirements such as period length, see gray inset box. The 49 ovals represent models with

parameters mostly or entirely based on the work of other models. Colored shapes represent models that

informed the parameters of other models, where arrows of the same color connect the original and the

subsequent models it informs (gray indicates that a model was not yet used for parameterizing other

models).

  QQv1r1

FIGURE 5

FIGURE 5. Data accumulation over time: overview of experimental studies available in

FlyClockbase and their use for parameter estimation over time.

(A) All studies in FlyClockbase are displayed based on the year they were published. Most studies have

not yet been used to inform model parameters (pink dashed line bars).

(B) Summary of studies that were later used for parameter estimation are shown as blue solid line bars.

These 14 studies are given (HARDIN *et al.* 1992; ZENG *et al.* 1994; SEHGAL *et al.* 1995; MARRUS *et al.*

1996; SO AND ROSBASH 1997; BAE *et al.* 1998; LEE *et al.* 1998; BLAU 1999; BAE *et al.* 2000; KIM *et al.*

2002; SHAFER *et al.* 2002; CYRAN *et al.* 2003; GLOSSOP *et al.* 2003; KADENER *et al.* 2007) and come from

the richer collection of 86 studies reported in version variant QQv1 of FlyClockbase: (HARDIN *et al.* 1990;

ZERR *et al.* 1990; ZWIEBEL *et al.* 1991b; HARDIN *et al.* 1992; HARDIN 1994; SEHGAL *et al.* 1994; ZENG *et al.*

1994; SEHGAL *et al.* 1995; VAN GELDER *et al.* 1995; BRANDES *et al.* 1996; MARRUS *et al.* 1996; QIU AND

HARDIN 1996; VAN GELDER AND KRASNOW 1996; MAJERCAK *et al.* 1997; ROUYER *et al.* 1997; SO AND

ROSBASH 1997; STANEWSKY *et al.* 1997; BAE *et al.* 1998; CHENG *et al.* 1998; DARLINGTON *et al.* 1998;

EMERY *et al.* 1998; HAMBLEN *et al.* 1998; KLOSS *et al.* 1998; LEE *et al.* 1998; STANEWSKY *et al.* 1998; BLAU

1999; ISHIKAWA *et al.* 1999; BAE *et al.* 2000; ROTHENFLUH *et al.* 2000b; BAO *et al.* 2001; CLARIDGE-CHANG

et al. 2001; MCDONALD *et al.* 2001; OKADA *et al.* 2001; KIM *et al.* 2002; SHAFER *et al.* 2002; STANEWSKY *et*

al. 2002; UEDA *et al.* 2002; AKTEN *et al.* 2003; CYRAN *et al.* 2003; GLOSSOP *et al.* 2003; DISSEL *et al.* 2004;

MAJERCAK *et al.* 2004; GLASER AND STANEWSKY 2005; CHEN *et al.* 2006; KOH *et al.* 2006; WIJNEN *et al.*

2006; YU *et al.* 2006; FANG *et al.* 2007; KADENER *et al.* 2007; MATSUMOTO *et al.* 2007; MUSKUS *et al.* 2007;

KIVIMÄE *et al.* 2008; MEISSNER *et al.* 2008; PESCHEL 2008; YOSHII *et al.* 2008; DUBRUILLE *et al.* 2009;

KADENER *et al.* 2009; LEAR *et al.* 2009; PESCHEL *et al.* 2009; WULBECK *et al.* 2009; YOSHII *et al.* 2009;

ZHENG *et al.* 2009; KULA-EVERSOLE *et al.* 2010; CHEN *et al.* 2011; CHIU *et al.* 2011; ITOH *et al.* 2011;

LAMAZE *et al.* 2011; LIM *et al.* 2011; GRIMA *et al.* 2012; HUGHES *et al.* 2012; KUMAR *et al.* 2012; LI AND

ROSBASH 2013; RODRIGUEZ *et al.* 2013; HERMANN-LUIBL *et al.* 2014; LAMBA *et al.* 2014; LEE *et al.* 2014a;

OH *et al.* 2014; PEGORARO *et al.* 2014; SHI *et al.* 2014; SIMONI *et al.* 2014; SUBRAMANIAN *et al.* 2014;

WEISS *et al.* 2014; ZHENG *et al.* 2014; ABRUZZI *et al.* 2015; MA *et al.* 2015).

1   **FIGURE 6**

FIGURE 6. Hypotheses on timing variances in PER and TIM as tested by FlyClockbase data,

which extends our understanding of circadian clocks in flies by enabling observations of

significant differences in variance between peaks and valleys of different clock components. Here

we summarize results from comparing mRNAs and proteins of *per* and *tim*. Yellow funnels highlight

changes in respective variances between timing traits of *per* and *tim* that are not negligible (over 5% of

10^5 bootstraps of available data reject the correspondingly customized null-hypothesis H_{basic} that

variances are equal). For related analyses see also Table 6, Table 7, Figure 7, Table 4, Table S1 and

parts of the Supplemental Statistical Analysis. Overall, the summary statistics for the timing of peaks and

valleys of the mRNAs and proteins of *per* and *tim* are consistent with our current understanding of the *D.*

melanogaster circadian clock.

Open circles with dashed curves indicate mRNAs, and filled triangles with solid curves represent proteins; *per* (green) is shown in the top two, and *tim* (red) in the bottom two curves, all adjusted to fit locations of peaks and valleys within one cycle. Averages and *SD*s were calculated for *CZT* times, linearized as described in Materials and Methods, and then back-transformed to *DZT* (if differences existed as in the rare cases where day 1 of a time series was unusable).

The statistics shown report summaries for the *n* remaining time series after removing outliers by applying the method of Carling (2000) to the data in the *TraitTables* compiled from the time series observations in FlyClockbase and refined as described for '*Mod6'* in Figure 3. Histograms of the data summarized here are shown in Figure 7 as 'CoreData' and statistical results testing the significance of differences are reported in Table 6, the text of the Results Section, and in the Supplemental Statistical Analysis (automatically produced by the R script provided as Supplemental Material). Table 4 and Table S1 present related results from an independent statistical analysis of the same data that was completely implemented using only spread-sheet software. While the results shown in this figure and the underlying outlier analysis were performed in R, the input dataset *Mod6* analyzed by the R script was produced using spread-sheets.

2   **FIGURE 7:**

5   **FIGURE 7. Histograms for *per* and *tim* mRNA and protein time series traits like peaks and valleys**

6   **as observed in independent experiments show how FlyClockbase enables comparing variability.**

7   Here we compare the variability of timing for peaks (A, C) and valleys (B, D) of the proteins (A, B) and

8   mRNAs (C, D) for the genes *per* (green) and *tim* (yellow). For all panels (A-D), we provide overlapping

transparent histograms of all data (more transparent, dashed or dotted line) and of the core dataset from

which outliers were removed by a boxplot method with Carling's modification (less transparent, solid or no

line). Descriptive summary statistics of the 'CoreData' distributions (excluding outliers) are given in Figure

6 and measures of statistical significance of differences are reported in Table 6 of the Results Section,

and in the Supplemental Statistical Analysis (the annotated PDF of plots that were auto-generated by the

script that also produced the plots in this figure; for an overview of calculations in the script, see Statistical

Methods; for details, see R code in Supplemental Material). We assessed the impact of manually

identified rare extreme outliers by excluding them from a copy of the dataset (termed "NoXtremes") for

which we repeated all analyses, including R's automated outlier detection using Carling's modification; the

results are not substantially different and are reported in parts of the the Supplemental Statistical

Analysis. For other related analyses see also Table 6, Table 7, Figure 6, Table 4, and Table S1.

  QQv1r1

FIGURE 8:

FIGURE 8. Comparisons of PCR vs non-PCR methods for measuring amounts of per mRNA in

time series as enabled by FlyClockbase. Here we compare the variability of peak timing in all available

time series of *per* mRNA without grouping by study, in order to contrast distributions of peak times

obtained from time series (A) observed by all PCR methods combined (red) vs (blue) all non-PCR

methods combined (blue); (B) observed by different PCR methods like qPCR or RT-PCR individually (red)

vs (blue) all non-PCR methods combined (blue); (C) observed in studies citing at least one method

reference (green) vs (purple) studies with no method reference (purple); and (D) observed in studies

sharing two or more method references (green) vs (purple) studies with method references unique among

studies contributing to *per* mRNA observations in FlyClockbase (*and* citing at least one method reference,

purple). For all panels (A-D), we provide overlapping transparent histograms. Measures of statistical

significance of differences are reported in the Results Section and parts of the Supplemental Statistical

Analysis. For related data and analyses see also Table 8, Figure 9, and Figure 10.

            QQv1r1

1   **FIGURE 9:**

FIGURE 9. Referenced protocol use similarity matrix for studies reporting *per* mRNA time series,

grouping protocols by measurement approach. The vertical axis shows time series IDs of *per* mRNA

time series grouped by broader measurement method, and the horizontal axis indicates the experimental

protocols references cited. The numbers given in the table for each time series and reference refer to a

classification key indicating which of several broader types of methods had been used (1 – RNA

extraction; 2 – Expression analysis; 3 – SYBR green detection protocol; 4 – RNA quantification; 5 – RNA

purification; 6 – serial dilutions; 7 – controls; 8 – general protocol; 9 – probes; 10 – hybridization, RNase

digestion, electrophoretic product separation; 11 – dissection; 12 – transfer to gel, labeling probes,

hybridization, washing; 13 – PCR for cDNA library). Boxes shaded in gray represent the references cited

by each time series, and the numbers in the gray boxes correspond to the broader types of methods to

which this specific part of a protocol reference belongs. Gray boxes shaded in the "no reference" column

indicate time series from studies without method references. All time series from the same study

measured with a given method are outlined in dark grey. For related data and analyses see also Table 8,

Figure 8, Figure 10, and parts of the Supplemental Statistical Analysis. Table 8 provides summaries of

counts of time series and studies based on information in this table. The details of experimental protocols

are described in their respective references (OLIVER AND PHILLIPS 1970; KONOPKA AND BENZER 1971;

CATHALA *et al.* 1983; ZINN *et al.* 1983; OCONNELL AND ROSBASH 1984; THEURKAUF *et al.* 1986;

CHOMCZYNSKI AND SACCHI 1987; CITRI *et al.* 1987; LORENZ *et al.* 1989; SAMBROOK *et al.* 1989; HARDIN *et al.* 1990; ZWIEBEL *et al.* 1991a; HARDIN AND HALL 1992; HARDIN 1994; SEHGAL *et al.* 1994; ZENG *et al.*

1994; SEHGAL *et al.* 1995; VAN GELDER *et al.* 1995; MARRUS *et al.* 1996; MYERS *et al.* 1996; MAJERCAK *et al.* 1997; STANEWSKY *et al.* 1997; ALLADA *et al.* 1998; BAE *et al.* 1998; CHENG *et al.* 1998; EMERY *et al.*

1998; PRICE *et al.* 1998; SIDOTE *et al.* 1998; STANEWSKY *et al.* 1998; MAJERCAK *et al.* 1999; CERIANI *et al.*

2002; GRIMA *et al.* 2002; UEDA *et al.* 2002; MENET *et al.* 2010; NAGOSHI *et al.* 2010; ABRUZZI *et al.* 2011;

KHODOR *et al.* 2011; RODRIGUEZ *et al.* 2013).

  QQv1r1

Figure 10

A: [-------------------------------] vs [----------------]
B: [-------------] vs [-------------]
C: [-------------] vs [-----------------------------------]

FIGURE 10: Logic of datasets in statistics: example for an interaction of statistics and formal logic

where the logic for selecting datasets unnecessarily reduced the power of a statistical test.

Here we illustrate the logic we used for constructing the datasets that group observed time series into the

sets used for approximating the degree of method standardization and method similarities that produced

the per mRNA time series observations in FlyClockbase. We denote as 'any' an individual time series

from an individual study with a count of references to experimental protocols, where the count is ≥1; 'no'

stands for a count of 0. Grouping **A** and **B** indicate test groupings performed by the R script in the

Supplemental Material. Grouping **C** indicates the grouping suggested by a more thorough logical analysis

performed after completing the statistical tests presented in the Results Section. For related data and

analyses see also Table 8, Figure 8, Figure 9, and parts of the Supplemental Statistical Analysis.

FIGURE 11

FIGURE 11: Biological model curator and compiler architect collaboration model for improving the integration of biological data into *VBIR*s while constructing the type system required by a *VBIR* compiler for handling the ubiquitous uncertainty of biological data more precisely.

Here we show how biological model curators and compiler builders can collaborate by depicting important aspects of the trans-disciplinary interactions that led to the construction of FlyClockbase, a versioned biological information resource (*VBIR*). We have been using our compiler expertise to inform fundamental decisions about conceptual data structures and file formats in FlyClockbase and TabFS (designed simultaneously, all formal definitions are beyond the scope of this paper).

Division of work. We divided work among domains as indicated by the dotted line. Above, biological model curators worked with a focus on constructing and using FlyClockbase in order to

understand the circadian clock of *Drosophila*. On the other side was the compiler architect focusing on

TabFS data structures and the logic necessary for representing the biological data collected by the

curators for FlyClockbase.

Mutual benefits. Designs were chosen such that implementing the TabFS compiler logic remained

as simple as possible, while enabling formal tests of the integrity of *VBIR* data. In the present study, we

developed many of the designs necessary for automation by manually pioneering important tasks. Doing

so in a real-life research system of non-trivial complexity was important; this allowed us to explore many

real-world details with substantial anticipated impacts on future use. The biological model curators

focused on integrating and analyzing time series; this often enabled them to readily provide the compiler

architect with detailed advance information about important use-cases and likely problems. The compiler

architect could then evaluate potential design options long before implementation could start. This

allowed for iterative reviews from several perspectives and with substantial time for analysis. It also

resulted in a *VBIR* design and compiler logic that substantially complement each other and are thus

prepared for automated testing of *VBIR* data integrity by following a number of simple rules.

Collaborative exchanges of insights at the dotted *VBIR-TabFS* compiler interface are exemplified

as broad arrows, indicating some important discussion topics (in terms closer to computer science for

brevity; our actual discussions were informal and used vocabulary closer to biology). Our trans-

disciplinary communication interface required both the high-level aspects of human collaboration and the

low-level aspects of technical information exchange. The information flows shown here critically depend

on biological model curators with a deep understanding of, and passion for, the details of state-of-the-art

biology. Their expertise is essential for defining the boundaries of the system that is being modeled. While

working to quantify its uncertainties properly in the *VBIR*, the biologists must be committed to bringing all

potential problems of interpretation to the attention of the architect of a compiler, which is being designed

to meet the formal needs of the *VBIR*. The architect must be able to select an appropriate mathematical

logic formalism for representing the relevant biological problems in meaningful ways. The team needs

excellent trans-disciplinary communication skills for efficiently describing, checking, and negotiating the

uncounted decisions that collectively generate the systematic organization of an efficient *VBIR*. This

requires a high sensitivity for the diffuse difficulties of accurately capturing computationally the

uncertainties and contradictions in biological observations encountered while constructing a *VBIR*. Such

biological problems require an appropriate logic formalism and thus need to be seen by the compiler

architect, irrespective of perceived severity.

Traps. Avoiding do-it-yourself analyses of 'simple problems' by non-computational biologists is

important for protecting against deceptive simplicity. It may not be possible to solve such problems on

desirable timelines; still, compiler developers have better chances of spotting the dangerous costly bugs

they can cause, which helps to identify solutions (that may already exist for other reasons). For example,

quantifying uncertainty with the cutting-edge logic formalism known as 'fuzzy plurivaluationism' enables

the representation of both semantic indeterminacy *and* various degrees of truth (SMITH 2008);

'BioBinaries' extend Boolean logic in similar ways (see Supplemental Material). It often takes a trained

eye to spot the need for a richer logic and it can be particularly hard to hunt logic errors that omit the

possibility of some options (PANKO 2016). Finding an appropriate logic formalism is pivotal, because it is

impossible to compute without assumptions about logic. Computers always produce logical answers, no

matter how flawed their logic might be when interpreted by humans in a real-world context. Such

problems are most efficiently addressed at the compiler level by enforcing an 'appropriate' type system.

Naturally, related discussions revolve around how to define 'appropriate common sense'.

Progress in defining type systems used for representing a biological system must be driven by

those biologists who study the respective biology; to help them capture type defining ideas, we use the

TabFS storage system of nested folders and tables which biologists can easily navigate using common

operating systems and spreadsheet tools (see main text).

Cost of constructing VBIRs without an appropriate compiler. In contrast, without the appropriate

logic and the equivalent of TabFS built into a corresponding *VBIR* compiler, it will still be necessary for

each *VBIR* to construct, find or develop corresponding tools if quality is important. Unfortunately,

economies of scale will be missing in this case, resulting in greatly reduced quality at greatly increased

cost. Thus, developing *VBIRs* on a shoe-string budget will either limit *VBIRs* to comparatively simple

datasets (thereby excluding much of biology) or is likely to trigger many rushed decisions about

underpinning logic. We found that correspondingly simplistic type systems easily frustrate non-computing

biologists by expecting them to coerce observed data and its uncertainties into an inappropriate logic

(demanding a precision not provided by the data). This results in biases that are often near-impossible to

detect, quantify, or exclude (e.g. by testing whether coercion indeed occurred). Thus, representing

biological data in programming languages that do not readily support an appropriate logic can easily

generate misleading code. It is possible in theory to work around such deficiencies. However, in practice,

such logic mismatches often trigger huge costs to be paid after indeterminate times. This is particularly

true if such logic has to be developed without the necessary time, logic expertise and compiler tools. As

many professional programmers know from their own experience, "premature optimization is the root of

all evil" (KNUTH 1974). This principle readily translates to premature choices of a logic formalism for

representing a biological system, prematurely chosen type systems, the premature optimization of

implementation speed and others that create numerous problems in computational biology. It is thus

conceivable that a substantial contribution towards the notoriously difficult task of funding *VBIR*

development might come from the construction of a compiler that simplifies these complex decisions by

offering solutions that can represent biological information and uncertainty more easily.

Conclusions for FlyClockbase. Accordingly, we have been designing the FlyClockbase type

system for maximizing simplicity and radical openness, allowing all users to easily suggest necessary

expansions (simply by changing cells in table text files of TabFS). Such suggestions still require review by

a *VBIR* architect with expertise in formal type systems. However, more eyes are likely to identify more

type-system challenges and might thus inspire better solutions. As indicated in the figure, many solutions

are fragmented at first, and larger designs emerge later. Still, the ultimate integration goal for each *VBIR*

is to find a single coherent architecture that is "as simple as possible, but not simpler" (EINSTEIN AND

CALAPRICE 2011).

Models based on Direct Data

Experimental studies with directly observed time series data that were
- Used in parameter estimation
- Ignored in parameter estimation

Shapes denote models with parameters based on:
- Direct experimental data
- Experimental data and time series abstractions
- Time series abstractions
- Other modeling studies

Models based on Abstract Time Series Requirements

Time series traits abstracted from experimental observations
- Has a period of about 24 hours
- Includes feedback loops
- Is robust to minor changes
- Appears physiologically realistic
- Has appropriate light response, including phase shifts
- Oscillates with reasonable amplitude
- Simulates mutant behavior as expected
- Components show nuclear localization
- Shows expected mRNA to protein delay
- Has appropriate amount of PER protein
- Has expected time of *per* mRNA peak

A   **Peaks for Proteins PER, TIM**

TIM

Protein
CoreData
solid line,
area less
transparent

Protein
AllData
dashed line,
area more
transparent

PER

Count of observed time series

Continuous Zeitgeber Time of peak

B   **Valleys for Proteins PER, TIM**

Protein
CoreData
solid line,
area less
transparent

PER

Protein
AllData
dashed line,
area more
transparent

TIM

Count of observed time series

Continuous Zeitgeber Time of valley

C   **Peaks for mRNAs *per, tim***

mRNA
CoreData
blank line,
area less
transparent

per

tim

mRNA
AllData
dotted line,
area more
transparent

Count of observed time series

Continuous Zeitgeber Time of peak

D   **Valleys for mRNAs *per, tim***

mRNA
CoreData
blank line,
area less
transparent

=(in this case)=

mRNA
AllData
dotted line,
area more
transparent

tim

per

Count of observed time series

Continuous Zeitgeber Time of valley

Method References Cited by the Study Reporting a given TimeSeries

Measurement Method	Time Series	Abruzzi et al 2011	Allada et al 1998	Bae et al 1998	Brandes et al 1996	Cathala et al 1983	Ceriani et al 2002	Cheng et al 1998	Chomczynski & Sacchi 1987	Citri et al 1987	Emery et al 1998	Grima et al 2002	Hardin et al 1990	Hardin et al 1992	Hardin 1994	Khodor et al 2011	Konopka & Benzer 1971	Lorenz et al 1989	Majercak et al 1997	Majercak et al 1999	Marrus et al 1996	Menet et al 2010	Myers et al 1996	Nagoshi et al 2010	O'Connell & Rosbash 1984	Oliver & Phillips 1970	Price et al 1998	Rodriguez et al 2012	Sambrook et al 1989	Sehgal et al 1995	Sehgal et al 1994	Sidote et al 1998	Stanewsky et al 1997	Stanewsky et al 1998	Theurkauf et al 1986	Ueda et al 2002	van Gelder et al 1995	Zeng et al 1994	Zinn et al 1983	Zwiebel et al 1991	No reference
PCR	6.3.1																																								■
	7.1.1																																								■
	11.1.1																																								
	11.1.2																																								
	13.1.1								1																																
	15.1.1																																								■
	20.1.1																					2																			
	27.1.1																																				1				
	28.1.1											3																													
	31.1.2																								4																
	31.2.2																								4																
	31.15.1																								4																
	40.2.1																																								■
	42.5.1																																								
	48.1.2																																			5					
	48.2.1																																			5					
	56.1.1		6									7												8																	
	56.2.1		6									7												8																	
	56.3.1		6									7												8																	
	56.15.1		6									7												8																	
	56.16.1		6									7												8																	
	56.17.1		6									7												8																	
	56.17.2		6									7												8																	
	56.18.1		6									7												8																	
	56.18.2		6									7												8																	
RNAse Protection Assay	56.4.1												7, 8											9																	
	60.2.1												7, 8																												
	62.4.1										10		7																					8							
	62.5.1										10		7																					8							
	62.6.1										10		7																					8							
	62.6.2				9						10		7, 9																					8						9	
	64.1.1																												9												
	69.1.1			9									7, 8					1																9			8				
	70.1.1																				8																				
	74.1.1												7, 8, 9					1																9			8				
	75.1.1												7, 8, 9	11																							11				
	75.1.2												7, 8, 9	11																							11				
	75.2.1												7, 8, 9	11																							11				
	75.2.2												7, 8, 9	11																							11				
	79.1.1																																9	9							
	82.2.1												9		9																										
	84.3.1																						8																		
	85.6.1												7, 9				1, 10																								9
	85.6.2												7, 9				1, 10																								9
	85.7.1												7, 9				1, 10																								9
	85.7.2												7, 9				1, 10																								9
	85.8.1												7, 9				1, 10																								9
	85.8.2												7, 9				1, 10																								9
	86.3.1												7, 9				1, 10																								9
	86.3.2												7, 9				1, 10																								9
	87.4.1																																			1, 9					
	88.1.1												8																												
	88.2.1												8																												
	88.3.1												8																												
	92.1.1												8				1											11													
	92.1.2												8				1											11													
	92.2.1												8				1											11													
	92.3.1												8				1											11													
	93.1.1																	9															9						8		
	94.1.1												8, 9																												
	94.2.1												8, 9																												
	95.1.1												1, 7, 9, 10																												
	95.2.1												1, 7, 9, 10																												
	95.2.2												1, 7, 9, 10																												
	95.3.1												1, 7, 9, 10																												
	96.1.1									9			1, 7				8																								
	96.1.2									9			1, 7				8																								
	96.2.1									9			1, 7				8																								
	96.2.2									9			1, 7				8																								
	97.1.1																	1, 8							9	11															
Northern Blot	41.5.1																																								■
	65.2.1																																								
	65.2.2																																								
	65.2.3																																								
	83.1.1																												12												
	89.1.1									9														9														8			
	89.2.1									9														9														8			
	89.3.1									9														9														8			
	91.1.1							1																				5, 8													
Microarray	2.7.1																																								■
	2.7.2																																								
	47.5.1																																								
	63.1.1																																								■
RNA-seq	17.5.1	13															1											1													
	17.5.2	13															1											1													
	17.5.3	13																																							
	17.5.4	13																																							