Title: MmpL3 is the flippase for mycolic acids in mycobacteria

Authors: Zhujun Xu[a], Vladimir A Meshcheryakov[a], Giovanna Poce[c], Shu-Sin Chng[a,b,*]

Affiliations:

[a]Department of Chemistry, National University of Singapore, Singapore 117543.

[b]Singapore Center on Environmental Life Sciences Engineering (SCELSE), National

University of Singapore, Singapore 117456.

[c]Dipartimento di Chimica e Tecnologie del Farmaco, Sapienza University of Rome, Rome

00185, Italy

[*]To whom correspondence should be addressed.   E-mail: chmchngs@nus.edu.sg

Author contributions: Z.X. and S.-S.C. designed research; Z.X. performed all experiments

described in this work; G.P. synthesized and provided BM212; V.A.M. expressed and purified

MmpL3-CTD; Z.X., and S.-S.C. analyzed and discussed data; Z.X., and S.-S.C. wrote the

paper.

The authors declare no conflict of interest.

Classification

Biological Sciences - Biochemistry

Abstract

The defining feature of the mycobacterial outer membrane (OM) is the presence of mycolic acids (MAs), which in part render the bilayer extremely hydrophobic and impermeable to external insults, including many antibiotics. While the biosynthetic pathway of MAs is well studied, the mechanism(s) by which these lipids are transported across the cell envelope is(are) much less known. MmpL3, an essential inner membrane (IM) protein, is implicated in MA transport, but its exact function has not been elucidated. It is believed to be the cellular target of several anti-mycobacterial compounds; however, evidence for direct inhibition of MmpL3 activity is also lacking. Here, we establish that MmpL3 is the MA flippase at the IM of mycobacteria, and is the molecular target of BM212, a 1,5-diarylpyrrole compound. We develop assays that selectively access mycolates on the surface of *Mycobacterium smegmatis* spheroplasts, allowing us to monitor flipping of MAs across the IM. Using these assays, we establish the mechanism-of-action of BM212 as a potent MmpL3 inhibitor, and employ it as a molecular probe to demonstrate the requirement for functional MmpL3 in the transport of MAs across the IM. Finally, we show that BM212 binds MmpL3 directly and inhibits its activity. Our work provides fundamental insights into OM biogenesis and MA transport in mycobacteria. Furthermore, our assays serve as an important platform for accelerating the validation of small molecules that target MmpL3, and their development as future anti-tuberculosis drugs.

Keywords

Mycobacterium tuberculosis; membrane biogenesis; lipid transport; trehalose monomycolate;

Mycobacterial membrane protein Large; drug binding and inhibition

Significance statement

Biological membranes define cellular boundaries, allow compartmentalization, and represent a prerequisite for life; yet, our understanding of membrane biogenesis remains rudimentary. Mycobacteria, including the human pathogen *Mycobacterium tuberculosis*, are surrounded by a double-membrane cell envelope that makes them intrinsically resistant to many antibiotics. Specifically, the outer membrane contains unique lipids called mycolic acids, whose transport mechanism across the envelope is unknown. In this study, we established the role of an essential membrane protein as the flippase for mycolic acids, and demonstrated that this protein is a direct target of an anti-mycobacterial compound. Our work provides insights into outer membrane biogenesis and lipid transport in mycobacteria, and the means to evaluate drugs that disrupt mycolic acid transport at the inner membrane.

\body

The outer membrane (OM) of *Mycobacterium tuberculosis*, the causative agent of

tuberculosis (TB), is distinctively characterized by the abundance of mycolic acids (MAs),

C_{60}-C_{90} long chain, branched fatty acids, packed together to produce a bilayer with markedly

reduced fluidity and permeability (1). These MAs come in the forms of trehalose

monomycolates (TMMs), trehalose dimycolates (TDMs), and mycolates covalently attached

to arabinogalactan (AG) polysaccharides, which are in turn linked to the peptidoglycan and

collectively known as the mAGP complex (Fig. 1*A*). MAs are synthesized at the inner

membrane (IM) as TMMs via a highly-conserved and well-characterized pathway (2), which

is the target of first line anti-TB drug, isoniazid (3). How MAs are transported across the cell

envelope and assembled into the OM, however, is less understood; proteins mediating TMM

flipping across the IM and transit across the periplasm have not been identified and/or

characterized (Fig. 1*A*). At the OM, the Ag85 complex transfers a mycolate chain from one

TMM molecule to another to form TDM, or to the AG polysaccharides to form the mAGP

complex (4). Tethering the OM to the cell wall via the AG polysaccharides further rigidifies

the membrane, making it extremely impermeable to a wide range of compounds, including

many antibiotics (1). The OM, and hence MAs, are essential for mycobacterial growth.

Recently, a conserved essential IM protein, MmpL3 (Mycobacterial membrane protein

Large 3), has been implicated in MA transport. Depletion of *mmpL3* in *Mycobacterium*

smegmatis results in accumulation of TMMs and reduced formation of TDMs and AG-linked

mycolates (5, 6), suggesting an impairment in TMM transport to the OM. Consistent with this,

MmpL3, like other MmpL proteins, belongs to the resistance, nodulation, and cell division

(RND) protein superfamily, and is believed to be a proton motive force (pmf)-dependent

transporter (7). Based on its cellular localization, MmpL3 is likely involved in either TMM

flipping across the IM, TMM release from the IM into the periplasm, or both (Fig. 1*A*). Yet,

its exact role has not been clearly defined, due largely to the lack of functional assays for its

putative transport activity. Treatment of mycobacteria with a few structurally-distinct small

molecule scaffolds, including ethylenediamines (e.g. SQ109) (8), 1,5-diarylpyrroles (e.g.

BM212) (9,10), adamantyl ureas (e.g. AU1235) (5), and others (11-15), result in similar

changes in mycolate species as in *mmpL3* depletion. These compounds inhibit growth, and

select for resistance mutations in *mmpL3*; however, there is limited evidence that they are

direct MmpL3 inhibitors. The lack of activity assays for MmpL3 made it impossible to test

the proposed mechanism-of-action of these putative inhibitors.

Here, we report that MmpL3 is the TMM flippase at the IM. Using a spheroplast model,

we developed assays to monitor IM topology of TMM. We found that 1,5-diarylpyrrole

BM212 inhibits TMM flipping across the IM in wild-type spheroplasts. Furthermore, we

established that specific MmpL3 variants confer resistance against this inhibition, indicating

that MmpL3 is required for flipping TMM across the IM. Finally, we demonstrated that

BM212 binds MmpL3 in vitro in a specific manner, and therefore directly targets MmpL3.

Our work represents the first demonstration of lipid transport activity of a key member of the

MmpL protein family, and highlights the importance of using small molecule probes to

interrogate protein function. Our assays have great utility in the validation and development

of MmpL3-targeting small molecules as future anti-TB drugs.

Results

Spheroplasts serve as a viable system to monitor TMM topology

To develop a functional assay for TMM flipping across the IM, we sought a system where TMM topology in the IM can be monitored. Mycobacterial spheroplasts are ideal for this purpose because they lack the OM and cell wall (16), and are bound only by the IM (17, 18), providing easy access to molecules-of-interest at this membrane. Due to the loss of periplasmic contents upon the formation of spheroplasts, we also expect the transport pathway(s) for TMM to the OM to be disrupted, thereby resulting in accumulation of TMM at the IM. *M. smegmatis* spheroplasts were successfully generated via sequential treatment with glycine and lysozyme (SI Appendix, Fig. S1), as previously reported (16). To examine whether MA synthesis is intact in spheroplasts, we profiled newly-synthesized lipids metabolically labelled with $[^{14}C]$-acetate. Thin layer chromatographic (TLC) analysis of lipids extracted from whole cells revealed a few major species whose syntheses are inhibited by isoniazid, indicating that these are mycolate-based lipids (Fig. 1*B*). We assigned two of these species as TDM and TMM on the basis of reported retention factors of these lipids on TLC plates developed under the same solvent system (19). We showed that mycolates are still produced in *M. smegmatis* spheroplasts; however, the extracted lipids only contain TMM, but not TDM. Furthermore, we can no longer detect newly-synthesized mAGP in the form of liberated mycolic acid methyl esters (MAMEs) in these spheroplasts (SI Appendix, Fig. S2). These results are consistent with the loss of Ag85 enzymes and the OM, where TDM and mAGP syntheses occur, and also with the lack of TMM transport to any possible remnants of

the OM. Given the extreme hydrophobicity of mycolates, we conclude that

newly-synthesized TMMs accumulate in the IM of spheroplasts, thus establishing a platform

for monitoring TMM flipping across the bilayer.

TMMs accumulated in spheroplasts reside in the outer leaflet of the IM

We next examined whether newly-synthesized TMMs accumulated in the inner or outer

leaflet of the IM in spheroplasts, by monitoring its accessibility to degradation by

recombinant LysB, a lipolytic enzyme. LysB is a mycobacteriophage-encoded esterase that is

specific for mycolates and plays the role of an endolysin important for the release of phage

particles from infected cells (20, 21). Substantial amounts (~77%) of newly-synthesized

TMMs in spheroplasts are readily and specifically hydrolyzed by purified LysB with the

concomitant release of MAs (Fig. 2*A*). Part of this hydrolysis can be attributed to the

background exposure of TMMs to LysB in a subset of spheroplasts that lysed during the

experiment (~30-50% cell lysis irrespective of the addition of LysB; Fig. 2*B*, and SI

Appendix, Fig. S3). The rest of newly-synthesized TMMs that were cut by LysB are likely

accessible on the surfaces of intact spheroplasts. We showed that an inactive LysB variant

does not result in the same effect (Fig. 2*A*). In addition, we demonstrated that LysB does not

enter intact spheroplasts (Fig. 2*B*), nor does it induce additional cell lysis compared to

controls (Fig. 2*B*, and SI Appendix, Fig. S3). Taken together, these results establish that most

of newly-synthesized TMMs have been translocated across the IM in intact spheroplasts, and

therefore reside in the outer leaflet of the membrane.

MmpL3 is responsible for flipping TMM across the IM

Several compounds, including SQ109, BM212 and AU1235, are believed to affect MmpL3-mediated TMM transport because mutations in *mmpL3* confer resistance against these small molecules (5, 8, 9). While it is not yet clear if these compounds directly inhibit MmpL3, they may be useful as probes to determine if MmpL3 is responsible for TMM flipping across the IM. Specifically, we asked whether these small molecules are able to inhibit TMM flipping in wild-type spheroplasts, and whether they would become less effective in doing so in spheroplasts expressing MmpL3 variants that confer resistance against them. We first tested the effects of these compounds in our LysB assay in wild-type spheroplasts. Remarkably, BM212 and AU1235 are able to reduce LysB-mediated hydrolysis of newly-synthesized TMMs in *M. smegmatis* spheroplasts (Fig. 3*A* and *C*), at concentrations 2x and 4x above their reported minimal inhibitory concentrations (MICs) (SI Appendix, Table S1) (5, 9). In contrast, SQ109 has no effect (Fig. 3*B* and *C*). For BM212, hydrolysis by LysB is strongly reduced, close to the level of background attributed to non-specific spheroplast lysis (SI Appendix, Fig. S3). We showed that the effects of BM212 and AU1235 are not due to direct inhibition of LysB activity, since TMMs are still hydrolyzed in detergent-solubilized samples (SI Appendix, Fig. S4). Instead, significant amounts of newly-synthesized TMMs are no longer accessible to LysB in the presence of either compound in intact spheroplasts, indicating inhibition of TMM flipping across the IM. As an alternative method to assess TMM topology in the IM and the effects of these drugs, we also examined the ability of membrane-impermeable fluorophore-conjugated streptavidin to bind to newly-synthesized TMMs engineered to contain biotin (biotin-TMMs) (Fig. 4*A*). Here, we

metabolically labelled TMMs with 6-azido-trehalose (22), which allowed us to covalently

attach an alkyne-containing biotin probe to TMM via the bio-orthogonal click reaction (23).

In wild-type spheroplasts, biotin-TMMs can be detected on the surface, indicating that

6-azido-TMMs have been translocated across the IM (Fig. 4*C* and *F*). Biotin labelling of

6-azido-trehalose groups, and therefore detection with streptavidin, is largely prevented if

LysB was added (SI Appendix, Fig. S5), confirming that we are visualizing trehalose-linked

mycolates (i.e. TMMs) at the IM in spheroplasts. We demonstrated that both BM212 and

AU1235 drastically reduce the amounts of 6-azido-TMMs, and hence biotin-TMMs, that can

be labelled with streptavidin (Fig. 4*D, E* and *F*). Again, SQ109 is not effective in this assay

(SI Appendix, Fig. S6). Consistent with the LysB accessibility assay, these results establish

that BM212 and AU1235, but not SQ109, inhibit TMM flipping across the IM.

To establish whether MmpL3 is responsible for flipping TMM across the IM, we

employed specific MmpL3 variants that render *M. smegmatis* cells less sensitive to BM212,

and tested if TMM flipping in spheroplasts expressing these variants would be more resistant

to the effects of BM212. The growth of cells expressing $MmpL3_{V197M}$ or $MmpL3_{A326T}$

variants is only fully inhibited in the presence of 4-8 times the concentration of BM212 that

inhibits wild-type growth (SI Appendix, Table S1) (9). We showed that BM212 does not

affect MmpL3 levels in wild-type spheroplasts (SI Appendix, Fig. S7); yet it strongly inhibits

TMM flipping in a dose-dependent manner (Fig. 5*A* and *D*). We further demonstrated that

BM212 is less effective at reducing LysB accessibility to TMM in spheroplasts expressing

$MmpL3_{V197M}$ or $MmpL3_{A326T}$ variants (Fig. 5*B, C* and *D*). In fact, BM212 is also unable to

inhibit the display of 6-azido-TMMs on the surface of spheroplasts expressing $MmpL3_{V197M}$

(Fig. 5*E*, *F*, *G* and *H*). Since TMM is only accessible on the outer leaflet of the IM in the

presence of functional MmpL3 (i.e. not inhibited by BM212), we conclude that MmpL3 is

the TMM flippase.

BM212 binds and directly inhibits MmpL3 function

MmpL3 function is believed to require the pmf, specifically the proton gradient (7).

Consistently, we showed that proton gradient uncouplers, such as CCCP and nigericin (Fig.

3), but not membrane potential disruptors, such as valinomycin-K$^+$ (SI Appendix, Fig. S8),

can inhibit LysB accessibility to TMMs in spheroplasts. Whether BM212 inhibits TMM

flipping by directly targeting MmpL3 is not clear. Contrary to previous reports (24), we did

not observe effects on the proton gradient (SI Appendix, Table S2) nor the membrane

potential (SI Appendix, Fig. S10) in spheroplasts treated with BM212, at concentrations that

inhibited TMM flipping. Therefore, BM212 inhibits MmpL3 not via indirect effects on the

pmf. To determine if BM212 directly targets MmpL3, we examined the ability of BM212 to

bind physically to purified MmpL3 (SI Appendix, Fig. S11) in vitro. We demonstrated that

[^{14}C]-BM212 binds purified wild-type MmpL3 in a saturable manner (Fig. 6*A*). Furthermore,

we showed that this interaction can be competed away by excess non-labelled BM212 (Fig.

6*B*), indicating that BM212 can bind specifically to MmpL3. In fact, mutations in *mmpL3*

that confer resistance to BM212 mostly cluster in a specific region when mapped onto a

structural model of MmpL3 (Fig. 6*C*), revealing a possible binding site. We conclude that

BM212 inhibits TMM flipping across the IM by binding MmpL3 directly.

Discussion

How the mycobacterial OM is assembled is not well understood. Many members of the MmpL protein family are believed to be transporters that contribute to the assembly of various OM lipids (25-28); however, their specific roles have not been clearly defined. MmpL3 is the only member of this family essential for growth (5-7). Using two independent assays that allow determination of TMM topology in the IM of mycobacterial spheroplasts, and employing putative inhibitors as molecular probes to modulate protein function, we have provided strong biochemical evidence that establish MmpL3 as the TMM flippase. Whether MmpL3 is the only protein mediating TMM flipping, or whether it is also involved in TMM release from the IM is not clear (Fig. 1*A*). One can posit that a second yet identified protein may be necessary for extracting TMM from the IM and handing it to a putative chaperone. This scenario would be comparable to the transport of lipoproteins across the cell envelope in Gram-negative bacteria (29). Alternatively, this step may also be mediated by MmpL3, in which case, flipping and release of TMM might be coupled; this would suggest MmpL3 could interact with the putative chaperone. Extending from this idea, a third scenario may be possible where TMM flipping and release are essentially one single step in a mechanism similar to RND efflux pumps, whereby TMM never really resides in the outer leaflet of the IM. This latter model is, however, less like since we have been able to decouple these steps by observing TMM translocation across the IM in our spheroplasts, which are effectively devoid of any putative chaperones. Moreover, despite being in the same RND protein superfamily, the structure of MmpL3 differs substantially from canonical RND efflux pumps (30). MmpL3 does form trimers like RND pumps (31), but the periplasmic domains of

MmpL3 are much smaller (30), and it contains a large C-terminal cytoplasmic domain.

Therefore, MmpL3 may not export TMM via an efflux mechanism. Further characterization

of this system would be necessary to tease apart these models.

TB is one of the leading causes of death by infectious disease, and remains a major

health problem worldwide (32). With the rapid emergence of multi- and extensive-drug

resistant (MDR/XDR) TB, there is an urgent need to develop anti-TB drugs with novel

mechanisms-of-action. In this regard, drugs inhibiting MmpL3, which has been shown to be

an ideal target (33), would be especially important. While many small molecules are thought

to inhibit MmpL3, it is puzzling how molecules with different molecular scaffolds may bind

and target the same protein. We have now developed assays that measure the topology of

TMM in the IM of mycobacterial spheroplasts, allowing the validation of true MmpL3

inhibitors. As a start, we have established that BM212 binds MmpL3 directly and inhibits its

function. Furthermore, we have shown that SQ109, a molecule that has reached phase IIb

clinical trials (Sequella), does not actually inhibit TMM flipping. In fact, it is likely that many

of these molecules do not inhibit MmpL3, and have other targets, as has been shown for

tetrahydropyrazo[1,5-*a*]pyrimidine-3-carboxamides (34). Our assays will help to select and

advance small molecules currently under development as MmpL3-targeting drugs.

Materials and Methods

Detailed descriptions of all experimental procedures can be found in the *SI Appendix*.

Assessing TMM accessibility to degradation by purified LysB in spheroplasts. *M.*

smegmatis spheroplasts were metabolically labelled with sodium [1-^{14}C]-acetate for 2 h,

followed by addition of purified LysB for 30 min at 37˚C. Lipids were extracted directly after

the LysB treatment, analyzed by TLC, and visualized via phosphor imaging. Where indicated,

putative MmpL3 inhibitors were added 15 min prior to addition of [^{14}C]-acetate.

6-azido-TMM surface display assay. *M. smegmatis* spheroplasts were metabolically

labelled with 6-azido-trehalose for 2 h at 37˚C to synthesize 6-azido-TMMs, which were then

reacted with Click-IT® Biotin DIBO alkyne to generate biotin-TMMs. Surface-exposed

biotin-TMMs were detected on spheroplasts using Alexa Fluor® 488-conjugated streptavidin,

and visualized by fluorescence microscopy. Where indicated, putative MmpL3 inhibitors

were added 15 min prior to addition of 6-azido-trehalose, and included in all wash buffers.

BM212-MmpL3 binding assay. Purified MmpL3 was incubated with indicated

concentrations of [^{14}C]-BM212 and/or cold BM212 for 30 min at room temperature. Samples

are analyzed using Clear Native-PAGE, followed by phosphor imaging and CB staining.

Acknowledgments

We thank Graham Hatfull (U Pittsburgh) for providing the pLAM3 plasmid for LysB

overexpression, and Derek Lin (NUS) for constructing the inactive LysB$_{S82A}$ variant. We are

grateful to Benjamin Swarts (Central Michigan U) for his generous gift of 6-azido-trehalose.

We also thank Eric Rubin (Harvard School of Public Health) for providing strains, and for

critical discussion and comments on the manuscript. We acknowledge Jasmine Chen

(Mechanobiology Institute) for help with microscopy. This work was supported by the

National University of Singapore Start-up funding, the Singapore Ministry of Education

Academic Research Fund Tier 1 and Tier 2 (MOE2014-T2-1-042) grants (to S.-S.C.).

References

References

1. Brennan PJ, Nikaido H (1995) The envelope of mycobacteria. *Annu Rev Biochem*

64(1):29-63.

2. Takayama K, Wang C, Besra GS (2005) Pathway to synthesis and processing of mycolic

acids in *Mycobacterium tuberculosis*. *Clin Microbiol Rev* 18(1):81-101.

3. Banerjee A, et al (1994) *inhA*, a gene encoding a target for isoniazid and ethionamide in

Mycobacterium tuberculosis. *Science* 263:227-229.

4. Jackson M, et al (1999) Inactivation of the antigen 85C gene profoundly affects the

mycolate content and alters the permeability of the *Mycobacterium tuberculosis* cell

envelope. *Mol Microbiol* 31(5):1573-1587.

5. Grzegorzewicz AE, et al (2012) Inhibition of mycolic acid transport across the

Mycobacterium tuberculosis plasma membrane. *Nat Chem Biol* 8(4):334-341.

6. Varela C, et al (2012) MmpL genes are associated with mycolic acid metabolism in

mycobacteria and corynebacteria. *Chem Biol* 19(4):498-506.

7. Domenech P, Reed MB, Barry CE (2005) Contribution of the *Mycobacterium*

tuberculosis MmpL protein family to virulence and drug resistance. *Infect Immun*

73(6):3492-3501.

8. Tahlan K, et al (2012) SQ109 targets MmpL3, a membrane transporter of trehalose

monomycolate involved in mycolic acid donation to the cell wall core of *Mycobacterium*

tuberculosis. *Antimicrob Agents Chemother* 56(4):1797-1809.

9. La Rosa V, et al (2012) MmpL3 is the cellular target of the antitubercular pyrrole

derivative BM212. *Antimicrob Agents Chemother* 56(1):324-331.

10. Poce G, et al (2013) Improved BM212 MmpL3 inhibitor analogue shows efficacy in

acute murine model of tuberculosis infection. *PLoS One* 8(2):e56980.

11. Stanley SA, et al (2012) Identification of novel inhibitors of *M. tuberculosis* growth using

whole cell based high-throughput screening. *ACS Chem Biol* 7(8):1377-1384.

12. Remuiñán MJ, et al (2013) Tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide and

N-benzyl-6′,7′-dihydrospiro[piperidine-4,4′-thieno[3,2-c]pyran]   analogues   with

bactericidal efficacy against *Mycobacterium tuberculosis* targeting MmpL3. *PLoS One*

8(4):e60933.

13. Rao SP, et al (2013) Indolcarboxamide is a preclinical candidate for treating

multidrug-resistant tuberculosis. *Sci Transl Med* **5**(214):214ra168.

14. Lun S, et al (2013) Indoleamides are active against drug-resistant *Mycobacterium*

tuberculosis. *Nat Commun* 4:2907.

15. Dupont C, et al (2016) A new piperidinol derivative targeting mycolic acid transport in

Mycobacterium abscessus. *Mol Microbiol* 101(3):515-529.

16. Dhiman RK, et al (2011) Lipoarabinomannan localization and abundance during growth

of Mycobacterium smegmatis. *J Bacteriol* 193(20):5802-5809.

17. Udou T, Ogawa M, Mizuguchi Y (1982) Spheroplast formation of *Mycobacterium*

smegmatis and morphological aspects of their reversion to the bacillary form. *J Bacteriol*

151:1035-1039.

18. Udou T, Ogawa M, Mizuguchi Y (1983) An improved method for the preparation of

mycobacterial spheroplasts and the mechanism involved in the reversion to bacillary form:

electron microscopic and physiological study. *Can J Microbiol* 29:60-68.

19. Bansal-Mutalik R, Nikaido H (2014) Mycobacterial outer membrane is a lipid bilayer and

the inner membrane is unusually rich in diacyl phosphatidylinositol dimannosides. *Proc*

Natl Acad Sci USA 111(13):4958-4963.

20. Payne KM, Hatfull GF (2012) Mycobacteriophage endolysins: diverse and modular

enzymes with multiple catalytic activities. *PLoS One* 7(3):e34052.

21. Gil F, et al (2010) Mycobacteriophage Ms6 LysB specifically targets the outer membrane

of *Mycobacterium smegmatis*. *Microbiology* 156(5):1497-1504.

22. Swarts BM, et al (2012) Probing the mycobacterial trehalome with bioorthogonal

chemistry. *J Am Chem Soc* 134(39):16123-16126.

23. Jewett JC, Sletten EM, Bertozzi CR (2010) Rapid Cu-free click chemistry with readily

synthesized biarylazacyclooctynones. *J Am Chem Soc* 132(11):3688-3690.

24. Li W, et al (2014) Novel insights into the mechanism of inhibition of MmpL3, a target of

multiple pharmacophores in *Mycobacterium tuberculosis*. *Antimicrob Agents Chemother*

58(11):6413-6423.

25. Converse SE, et al (2003) MmpL8 is required for sulfolipid-1 biosynthesis and

Mycobacterium tuberculosis virulence. *Proc Natl Acad Sci USA* 100(10):6121-6126.

26. Jain M, Cox JS (2005) Interaction between polyketide synthase and transporter suggests

coupled synthesis and export of virulence lipid in *M. tuberculosis*. *PLoS Pathog* 1(1):e2.

27. Pacheco SA, Hsu FF, Powers KM, Purdy GE (2013) MmpL11 protein transports mycolic

acid-containing lipids to the mycobacterial cell wall and contributes to biofilm formation

in *Mycobacterium smegmatis*. *J Biol Chem* 288(33):24213-24222.

28. Belardinelli JM, et al (2014) Biosynthesis and translocation of unsulfated acyltrehaloses

in *Mycobacterium tuberculosis*. *J Biol Chem* 289(40):27952-27965.

29. Okuda S, Tokuda H (2011) Lipoprotein sorting in bacteria. *Annu Rev Microbiol*

65:239-259.

30. Chim N, et al (2015) The structure and interactions of periplasmic domains of crucial

MmpL membrane proteins from *Mycobacterium tuberculosis*. *Chem Biol*

22(8):1098-1107.

31. Belardinelli JM, et al (2016) Structure-function profile of MmpL3, the essential mycolic

acid transporter from *Mycobacterium tuberculosis*. *ACS Infect Dis* 2(10):702-713.

32. Global actions and investments fall far short of those needed to end the global TB

epidemic: WHO report 2016. *World Health Organization, Geneva*

(WHO/HTM/TB/2016.13).

33. Li W, et al (2016) Therapeutic potential of the *Mycobacterium tuberculosis* mycolic acid

transporter, MmpL3. *Antimicrob Agents Chemother* 60(9):5198-5207.

34. Cox JA, et al (2016) THPP target assignment reveals EchA6 as an essential fatty acid

shuttle in mycobacteria. *Nat Microbiol* 1:15006.

35. Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE (2015) The Phyre2 web

portal for protein modeling, prediction and analysis. *Nat Protocols* 10(6):845-858.

Figures

Fig. 1. TMM biosynthesis is intact in mycobacterial spheroplasts. (*A*) A schematic diagram

illustrating the processes important for MA transport across the cell envelope. Following

synthesis, TMMs must be flipped across the IM, released from the IM and then transported

across the periplasm (presumably via a chaperone). MmpL3 is implicated in TMM transport

at the IM, but its exact role has not been elucidated. At the OM, the Ag85 complex transfers

the mycolate chain from TMM to cell wall-linked AG polysaccharides, or to another TMM to

form TDM. Other known lipid species found in the OM and IM are omitted for simplicity. PL,

phospholipid. (*B*) TLC analysis of newly-synthesized [^{14}C]-labelled lipids extracted from

wild-type *M. smegmatis* cells (WC) and spheroplasts (SP), visualized by phosphor imaging.

Lipids were radiolabelled in the presence or absence of isoniazid as indicated. The

developing solvent system comprises chloroform-methanol-water (30:8:1). A mycolate-based

species that appears only in the presence of glucose is indicated with an asterisk (*). PE,

phosphatidylethanolamine; CL, cardiolipin; PI, phosphatidylinositol; PIM,

phosphatidylinositol mannoside.

Fig. 2. Newly-synthesized TMMs in mycobacterial spheroplasts are accessible to degradation

by LysB, indicating that these TMMs reside in the outer leaflet of the IM. (*A*) TLC analyses

of newly-synthesized [^{14}C]-labelled lipids extracted from *M. smegmatis* spheroplasts treated

with functional or non-functional (S82A) LysB. Lipids were resolved on TLCs developed

using solvent systems comprising either chloroform-methanol-water (30:8:1) (*left*) or

hexane-diethylether-acetic acid (70:30:1) (*right*), followed by phosphor imaging. In addition

to MA, treatment with functional LysB also resulted in the release of an unidentified apolar

lipid, annotated with an asterisk (*). TAG, triacylglycerol. (*B*) α-GroEL2 and α-His

immunoblot analyses of pellet and supernatant fractions obtained from sedimentation of *M.*

smegmatis spheroplasts treated with functional or non-functional (S82A) LysB.

Fig. 3. Anti-mycobacterial compounds, BM212 and AU1235, reduce TMM accessibility to

LysB in spheroplasts, indicating inhibition of TMM flipping across the IM. (*A, B*)

Representative TLC analyses of [^{14}C]-labelled lipids newly-synthesized in the presence of

indicated concentrations of (*A*) BM212, AU1235, and (*B*) SQ109, and extracted from *M.*

smegmatis spheroplasts following treatment with or without purified LysB. The effects of

pmf disruptors, carbonyl cyanide *m*-chlorophenyl hydrazone (CCCP) and nigericin, were also

tested. At higher concentrations, these uncouplers affected lipid synthesis, consistent with the

depletion of ATP. DMSO and methanol were used to dissolve the respective compounds and

thus serve as negative controls. Equal amounts of radioactivity were spotted for each sample.

The developing solvent system comprises chloroform-methanol-water (30:8:1). (*C*) A

graphical plot showing the effects of various compounds on the amounts of LysB-accessible

TMMs in spheroplasts. The percentage of TMMs accessible to LysB is given by the

difference in TMM levels between samples with or without LysB treatment, normalized

against that in control samples without LysB treatment. TMM levels in each sample were

quantified as a fraction of total mycolates (TMM+MA). Average percentages and standard

deviations from three biological replicates are plotted. The average background of TMM

hydrolysis due to random cell lysis during the experiment (~40%) is indicated. Student's

t-test: *, $p < 0.05$ compared to the corresponding DMSO or methanol controls.

Fig. 4. Anti-mycobacterial compounds, BM212 and AU1235, reduce surface display of 6-azido-TMMs in spheroplasts, indicating inhibition of TMM flipping across the IM. (*A*) A schematic diagram illustrating the 6-azido-TMM surface display assay. Spheroplasts were incubated with 6-azido-trehalose to allow synthesis of 6-azido-TMMs (22), which were subsequently labelled with alkyne-containing biotin (DIBO-biotin) via click chemistry (23). Surface-exposed biotin-TMMs were recognized by Alexa Fluor 488-conjugated streptavidin, and visualized by fluorescence microscopy. (*B-E*) Representative bright-field and fluorescence microscopy images following DIBO-biotin/Alexa Fluor 488-streptavidin labelling of spheroplasts synthesizing (*B*) TMM, or 6-azido-TMM in the presence of (*C*) DMSO, (*D*) BM212 (2xMIC), and (*E*) AU1235 (2xMIC). Scale bars = 3 μm. (*F*) The fluorescence intensity per unit area for individual spheroplasts (N=100) in each condition (*B-E*) is plotted, with the medians and interquartile ranges indicated. Mann-Whitney test: ****, p < 0.0001 compared to the "no drug treatment" control.

Fig. 5. Mutations in MmpL3 render BM212 less effective in the inhibition of TMM flipping

across the IM. (*A-C*) Representative TLC analyses of [^{14}C]-labelled lipids newly-synthesized

in the presence of indicated concentrations of BM212, and extracted from (*A*) WT, (*B*)

mmpL3$_{V197M}$, and (*C*) *mmpL3$_{A326T}$ M. smegmatis* spheroplasts following treatment with or

without purified LysB. Equal amounts of radioactivity were spotted for each sample. The

developing solvent system comprises chloroform-methanol-water (30:8:1). (*D*) A graphical

plot showing the dose-dependent effects of BM212 on the percentage of TMMs accessible to

LysB in the respective spheroplasts (quantification as per Fig. 3). Average percentages and

standard deviations from three biological replicates are plotted. The average background of

TMM hydrolysis due to random cell lysis during the experiment (~40%) is indicated.

Student's t-test: *, $p < 0.05$; **, $p < 0.01$ compared to the corresponding DMSO controls for

each respective strain. (*E-G*) Representative bright-field and fluorescence microscopy images

442   following DIBO-biotin/Alexa Fluor 488-streptavidin labelling of *mmpL3*$_{V197M}$ spheroplasts

443   synthesizing (*E*) TMM, or 6-azido-TMM in the presence of (*F*) DMSO, and (*G*) BM212

444   (2xMIC). Scale bars = 3 μm. (*H*) The fluorescence intensity per unit area for individual

445   spheroplasts (N=100) in each condition (*E-G*) is plotted, with the medians and interquartile

446   ranges indicated. Mann-Whitney test: N.S., $p > 0.5$ compared to the "no treatment" control.

Fig. 6. BM212 binds MmpL3 in vitro in a specific manner. (*A*) Clear Native-PAGE analyses

of purified MmpL3-His samples incubated with increasing concentrations of [^{14}C]-BM212,

visualized separately by phosphor imaging and Coomassie blue (CB) staining. The amount of

[^{14}C]-BM212 bound to MmpL3 at each concentration was quantified and plotted. (*B*) Clear

Native-PAGE analyses of purified MmpL3-His samples incubated with a fixed concentration

of [^{14}C]-BM212, but in the presence of increasing concentrations of cold BM212. Gels were

visualized separately by phosphor imaging and CB staining. (*C*) Mutations that confer

resistance against BM212 cluster on a structural model of MmpL3, suggesting a possible

binding site. A Phyre2 (35) structural model for *M. smegmatis* MmpL3 without its C-terminal

cytoplasmic domain is shown in side (*left*) and top (*right*) views. For clarity, periplasmic

domains are removed from the top view image. Residues important for passage of protons are

460 highlighted in black. Residues that conferred resistance against BM212 (10) when mutated in

461 MmpL3 from *M. smegmatis*, *M. bovis BCG* and *M. tuberculosis* are highlighted in red, purple

462 and cyan, respectively.