Evaluation of Oxford Nanopore MinION^{TM} Sequencing for 16S rRNA Microbiome Characterization

Xiao Ma[1], Elyse Stachler[1], Kyle Bibby[1, 2*]

[1]Department of Civil and Environmental Engineering, University of Pittsburgh, Pittsburgh, PA

15261,

[2]Department of Computational and Systems Biology, University of Pittsburgh Medical School,

Pittsburgh, PA 15261, USA

*Corresponding author

*Corresponding Author: Kyle Bibby, 709 Benedum Hall, Pittsburgh, PA 15261

BibbyKJ@pitt.edu, 412-624-9207

Keywords: Nanopore, microbiome, 16S rRNA, sequencing

Abstract

In this manuscript we evaluate the potential for microbiome characterization by sequencing of near-full length 16S rRNA gene region fragments using the Oxford Nanopore MinION (hereafter 'Nanopore') sequencing platform. We analyzed pure-culture *E. coli* and *P. flourescens*, as well as a low-diversity mixed community sample from hydraulic fracturing produced water. Both closed and open reference operational taxonomic unit (OTU) picking failed, necessitating the direct use of sequences without OTU picking. The Ribosomal Database Project classifier against the Green Genes database was found to be the optimal annotation approach, with average pure-culture annotation accuracies of 93.8% and 82.0% at the phyla and genus levels, respectively. Comparative analysis of an environmental sample using Nanopore and Illumina MiSeq sequencing identified high taxonomic similarity when using a weighted metric (Bray-Curtis), and significantly reduced similarity when using an unweighted metric (Jaccard). These results highlight the great potential of Nanopore sequencing to analyze broad microbial community trends, and the challenge of applying Nanopore sequencing to discern rare taxa in mixed microbial communities. Finally, we observed that between-run carryover accounted for >10% of sequence reads, necessitating future development to either prevent carryover or filter sequences of interest (e.g. barcoding).

Introduction

Interest in studying the microbiome, microbiota associated with various environments, has exploded in recent years largely due to the rapid expansion in 'next-generation sequencing' capabilities and subsequent reduction in cost. It has been recognized that the human microbiome plays an important role in many different clinical outcomes, including obesity (Turnbaugh et al.

2006), immune state (Kumar et al.), and infection (Khoruts et al. 2010). The human microbiome

is comprised of a diverse array of commensal microorganisms in and on the human body, and

emerging research has suggested a clinical role for the microbiome in either therapeutic

development (e.g. probiotics) (Good et al. 2014) or diagnostics (Pflughoeft and Versalovic 2012).

Concurrently, significant interest has emerged in the microbiomes of various other environments,

such as buildings (Adams et al. 2016, Lax and Gilbert 2015, Prussin et al. 2016) and water

systems (Baron et al. 2014).

Currently, the most common microbiome analysis approach is high-throughput amplicon

sequencing of the 16S rRNA gene region. The most widely used sequencing technologies are

Illumina sequencing platforms (e.g. the MiSeq and HiSeq). These platforms are accurate and

generate a large amount of data, but are limited by capital costs, the necessity to pool samples to

reduce per-sample costs, sequence read length, and a turnaround time of days to weeks that may

be inadequate for many applications. Recently, Oxford Nanopore has released a small and

inexpensive sequencing platform called the MinION^{TM}, which has been previously reviewed

(Laver et al. 2015, Mikheyev and Tin 2014). Capital costs are reduced to per-run costs that are

comparable with the current Illumina platforms, and data analysis is possible in near-real time,

enabling investigators to generate sequence data as-needed. For example, samples were correctly

assigned to the *Salmonella* species in 20 minutes and serotype in 40 minutes (Quick et al. 2015).

Amplicon (non-metagenomic) sequencing has also been successfully employed to identify both

bacterial and viral origin (Kilianski et al. 2015, Quick et al. 2016). Additionally, this technology

routinely produces sequences >10kb in length, enabling sequencing of the full 16S rRNA gene

region and more reliable taxonomic placement. Despite the benefits, the primary drawback to

this technology is relatively high error rates (currently reported to be ~8%), hindering

metagenomic analysis of highly diverse microbiome samples and requiring additional development and validation. Previous investigations have demonstrated successful 16S rRNA microbial community classification via sequencing of a single mouse gut microbiome (Shin et al. 2016) and a single mock microbial community (Benítez-Páez et al. 2016), including species-level assignment (Benítez-Páez et al. 2016); however, a formal assessment of 16S rRNA sequencing on the Nanopore platform, including analysis of pure-culture samples for annotation validation, is currently lacking. Development of Nanopore technology for microbiome analysis would enable rapid (<12 hours from sample to results) and low cost microbiome characterization that would be applicable to both the clinic and other applications.

In the current study we evaluate the potential for microbiome characterization via Nanopore sequencing of near full-length 16S rRNA PCR amplicons. First, we evaluate the accuracy and annotation strategies of sequences from pure culture *E. coli* and *P. flourescens* to determine the most appropriate sequence analysis approach. We then compare performance of Nanopore sequencing against the current state of the art Illumina sequencing using a sample from hydraulic fracturing wastewater. Finally, we investigate an apparent between-run carryover phenomena, and propose necessary future investigations to enable 16S rRNA microbiome characterization on the Nanopore platform.

Materials and Methods

Overview of DNA samples and sequencing libraries

Three pure culture bacterial DNA samples and an environmental DNA sample were analyzed in

the current study. The three pure culture bacterial DNA samples were extracted from pure

cultures of: (1) *Escherichia coli* (ATCC 15597), (2) *Pseudomonas fluorescens* (ATCC 13525),

and (3) *Mycobacterium smegmatis* str. mc2 155, respectively. The environmental DNA samples,

which had previously undergone 16S rRNA gene sequencing using the Illumina MiSeq platform,

were a hydraulic fracturing produced water sample (Lipus et al. In Revision) and a river water

sample (unpublished).

All Nanopore sequencing runs were conducted using the MinION Mk IB platform following

recommended sequencing protocols (Oxford Nanopore Technologies). The *E. coli* 16S rRNA

amplicon 2D library and the hydraulic fracturing produced water 16S rRNA amplicon 2D library

were sequenced individually using a Nanopore MIN-105 flow cell and a Nanopore MIN-106

flow cell, respectively. The remaining libraries were sequenced on a MIN-106 flow cell

following a sequential order: (1) *P. fluorescens* 2D library; (2) *M. smegmatis* 2D library; (3) river

water sample 2D library; (4) *M. smegmatis* 1D library. For the sequential sequencing runs, flow

cell washing was conducted immediately following the completion of the previous sequencing

run using a Nanopore washing kit WSH002 (Oxford Nanopore). The Oxford Nanopore

recommended washing protocol was used between runs, namely 150 µL of WSH002 solution A

was loaded to the flow cell through priming port and incubated at room temperature for 10

minutes, then 150 µL of WSH002 solution B was loaded through the priming port before the

next sequencing run and incubated for another 10 minutes at room temperature.

Nanopore sequencing library preparation

Previously described universal primers targeting the 16S rRNA gene region (S-D-bact-0008-c-
S20 and S-D-bact-1391-a-A-17) (Klindworth et al. 2013) were used for PCR. Each PCR was
conducted in a total volume of 50 µL, containing 5 µL 10x buffer, 5 µL dNTP mix, 2.5 µL of
each forward and reverse primer, 0.25 µL DreamTaq, 1 µL template DNA, and 33.75 uL
nuclease free molecular grade water. The temperature condition for the PCR was 3 minutes at
95°C; 30 cycles composed of 20 seconds at 95 °C, 30 seconds at 47 °C for annealing, 1 minute at
72 °C; and a final elongation at 72 °C for 15 minutes. All PCR products were purified using
Ampure XP beads and normalized to 45 µL containing 1 µg of purified PCR products. Negative
controls were used for all PCR reactions and DNA extractions, and all controls were negative.

2D libraries were prepared using a Nanopore NSK007 sequencing kit and recommended protocol
(Oxford Nanopore Technologies). The end repair step of the purified PCR products was
conducted by adding 7 µL Ultra II End-Prep buffer, 3 µL Ultra II End-Prep enzyme mix (New
England Biolabs), and 5 µL control DNA provided with Nanopore NSK007 sequencing kit. The
end repair reaction mix was incubated at 20°C for 5 minutes and 65°C for 5 minutes. The end-
repaired PCR products were further purified using AMPure XP beads and ligated to the
sequencing adapters by adding 8 µL molecular grade water, 10 µL Nanopore NSK007 adapter
mix, 2 µL Nanopore NSK007 HPA solution, and 50 µL Blunt/TA Master Mix (New England
Biolabs), and then incubated at room temperature for 10 minutes. 1 µL HPT solution from the
NSK007 kit was added and incubated for an additional 10 minutes at room temperature. The
ligated and tethered 2D libraries were purified by using MyOne C1 beads (Thermo Scientific)
and eluted in 25 µL elution buffer (Oxford Nanopore Technologies). A description of the *M.*
smegmatis 1D library preparation is included in the Supplementary Information.

All sequencing flow cells were primed using 500 µL Running Buffer Fuel Mix diluted in 500 µL

molecular grade water following the recommended priming protocol (Oxford Nanopore

Technologies). After priming, 6 µL of each 2D sequencing library was mixed with 37.5 µL

Running Buffer Fuel Mix (Oxford Nanopore Technologies) and 31.5 µL molecular grade water,

then loaded to Nanopore flow cell for sequencing.

Sequence data processing

Base-calling and initial format conversion

The raw FAST5 files were base-called using Metrichor v2.42.2 with 2D Basecalling for FLO-

MIN106 250bps workflow and 1D Basecalling for FLO-MIN106 450bps workflow. Passed 2D

reads of 2D sequencing libraries and passed template reads of 1D sequencing libraries were

converted to FASTA files for downstream analysis using Poretools (Loman and Quinlan 2014).

Operational Taxonomic Unit Evaluation

Operational taxonomic unit (OTU) picking was conducted by using both closed-reference and *de*

novo picking strategies implemented in QIIME 1.9.2 (Caporaso et al. 2010a). Closed-reference

OTU picking was conducted by using Greengenes 13.8 as the reference database (DeSantis et al.

2006).

Taxonomy Assignment

For pure culture *E. coli* and *P. fluorescens* sequencing data, taxonomy was assigned to each

sequence read within QIIME 1.9.0 (Caporaso et al. 2010b) using the RDP classifier (Wang et al.

2007) against Greengenes 13.8 (DeSantis et al. 2006) and RDP 16S rRNA training set (Wang et

al. 2007) as the reference database, respectively; as well as using BLASTn (Altschul et al. 1990)

against Greengenes 13.8 (DeSantis et al. 2006) as a reference database. For subsequent analyses,

taxonomy was assigned to each individual sequence read using the RDP classifier (Wang et al.

2007) against Greengenes 13.8 (DeSantis et al. 2006) as this approach was found to achieve the

highest taxonomy assignment accuracy for the pure culture *E. coli* and *P. fluorescens* sequence

data.

Illumina 16S rRNA amplicon sequencing data of the produced water and river water samples

were re-processed using the same approach as Nanopore sequence data. The Illumina data were

clustered into OTUs using 100% similarity threshold, i.e. each identical Illumina sequencing

read was assigned taxonomy using RDP classifier (Wang et al. 2007) against Greengenes 13.8

(DeSantis et al. 2006).

Jaccard and Bray-Curtis distances, which are dissimilarity distances measuring level of

dissimilarity between two microbial communities, were calculated using QIIME 1.9.2 (Caporaso

et al. 2010a) to measure the degree of similarity of the produced water microbial community

similarity between Nanopore and Illumina platforms. Significance of correlation between the two

technical replicates of the produced water sample across sequencing platforms was conducted

using Pearson's correlation implemented in Minitab 16.

Results and Discussion

Sequencing

Sequencing was performed on eight 16S rRNA libraries; however, due to apparent between-run

carryover (discussed below), only the initial run on each of three flow cells was used for

subsequent analyses of annotation approach and accuracy. These sequencing libraries were

derived from two pure-culture samples, *E. coli* and *P. fluorescens*, and a low-diversity sample

from hydraulic fracturing produced water. Detailed sequencing results are shown in Table 1.

Raw   sequence   data   can   be   found   on   Figshare   using   DOI:

https://dx.doi.org/10.6084/m9.figshare.4515752.v3.

Pure Culture Analyses

We first evaluated the suitability of Operational Taxonomic Unit (OTU) picking methods using

Nanopore sequence data. OTU picking via comparison with a reference library, i.e. closed-

reference OTU picking, failed, with no sequences being assigned to an OTU (i.e. all sequences

were excluded). *De novo* OTU clustering, i.e. OTU clustering by determining between-sequence

similarity, was then evaluated using similarity thresholds between 90-100%. At the typically

used similarity threshold of 97%, all sequences from both pure-culture samples were assigned to

unique OTUs (i.e. the ratio of OTUs to sequences was one). Results from *de novo* OTU

clustering evaluation are shown in Figure 1. These results highlighted the challenge of clustering

reads from long, error-prone sequences, and necessitated analyzing the taxonomy of sequences

individually without OTU picking.

We next evaluated the ability to accurately annotate the taxonomy of pure-culture Nanopore 16S

rRNA sequences using three different annotation approaches: the naïve Bayesian Ribosomal

Database Project (RDP) classifier with the Green Genes database; the RDP classifier against the

RDP database; and BLAST against the Green Genes database. Results from this evaluation are

shown in Figure 2. The RDP classifier against the Green Genes database was found to be the best

performing annotation strategy. Using this approach, the annotation accuracy for *E. coli* was

96.7% and 81.9% at the phyla and genus level, respectively, and was 90.9% and 82.0% for *P.*

fluorescens at the phyla and genus level, respectively.

Comparison with Illumina Sequencing

We next evaluated Nanopore sequencing to characterize an environmental sample from

hydraulic fracturing produced water (Lipus et al. In Revision). This sample was selected as it

exhibited low alpha-diversity in previous analyses. The same DNA extract was used for both

analyses.

By assigning taxonomy to each individual sequence, five phyla were identified by the Nanopore

platform, and eleven phyla were identified by the Illumina platform. Among them, four phyla

that together accounted for greater than 99% of sequence relative abundance were identified by

both platforms (Table 2). Nine shared genera were detected by both platforms, accounting for

relative abundances of 98.3% on the Nanopore platform and 81.6% on the Illumina platform

(Table 2). Both the Nanopore and Illumina platforms revealed similar microbial community

structure of the produced water sample. The Firmicutes Phylum dominated the microbial

community with relative abundance higher than 90% with both platforms (Table S1). Phyla

unique to the Nanopore and Illumina platforms accounted for less than 0.5% of relative

abundance (Table S1). At the genus level, the produced water microbial community was

dominated by the genus *Halanaerobium* (Table S2), with 95.6% of Nanopore sequences and

76.8% of Illumina sequences being assigned to *Halanaerobium*. In addition, 14.6% of Illumina

16S rRNA sequence reads were assigned to Clostridiales, which is within the same class with

Halanaerobium (Clostridia) (Table S2).

We also calculated the Jaccard distance (solely based on presence/absence of each taxa) and

Bray-Curtis distance (based on both presence/absence and relative abundance of each taxa)

between the produced water microbial community revealed by Nanopore and Illumina platforms

at different taxonomic levels. Jaccard and Bray-Curtis distances measure the degree of microbial

community structure difference between two samples, with a value of one indicating no

community structure overlap and a value of zero indicating identical microbial communities. We

adopted these measures to evaluate the level of difference between the technical replicates of the

same produced water sample between the Nanopore and Illumina sequencing platforms. The

Jaccard distance increased from 0.62 at the phylum level to 0.89 at genus level; the Bray-Curtis

distance increased from 0.04 at phylum level to 0.22 at genus level (Figure 3). Jaccard distances

were higher than Bray-Curtis distances at all phylogenetic levels, because more taxa were

assigned using short sequence reads by Illumina sequencing (Table 2) and Jaccard distance only

accounts for the presence and absence of each assigned taxa whereas Bray-Curtis distances take

relative abundance into account.

Pearson correlation of the relative abundance of each taxon between Nanopore and Illumina data

was conducted to further evaluate the reproducibility of sequencing results between the two

platforms. Significant correlation was found at all phylogenetic levels from phylum to genus (R

values > 0.98, p values < 0.001), indicating reproducible taxonomic assignment results can be

obtained between Nanopore and Illumina platforms.

It should be noted in these comparisons that different primer sets were used for the two analysis

approaches, which has previously been shown to bias microbiome community structure as

analyzed by 16S rRNA sequencing (Pinto and Raskin 2012). Despite this additional source of

bias, the above analyses imply that the weighted community structure is comparable between the

two platforms, encouraging future development.

Between-Run Sample Carryover

We noted an apparent carryover of sequences between pure culture runs of *P. fluorescens* and *M.*

smegmatis. The potential for sequence carryover has been anecdotally reported in the literature

(Greninger et al. 2015). We subsequently excluded all runs except the first run on each flow cell

from earlier analyses, and undertook a formal evaluation of sequence carryover.

Results from analysis of the *Mycobacterium smegmatis* run are shown in Figure 4. In this run,

76.5% of sequences were assigned to the correct Actinobacteria phyla, 11.0% of sequences were

incorrectly assigned to another domain or unable to be assigned, and 12.4% of sequences were

incorrectly assigned to the Proteobacteria phyla, presumptively resulting from sequence

carryover. 54.5% of sequences were assigned to the correct *Mycobacterium* genus whereas

10.8% of sequences were assigned to the *Pseudomonas* genus.

Following the *P. fluorescens* and *M. smegmatis* runs, we subsequently ran an additional

environmental sample derived from river water and re-ran a *M. smegmatis* sample using 1D

technology. In the second *M. smegmatis* run, we observed only 0.5% of sequences to be assigned

to the correct Mycobacteriaceae family, compared with 54.5% in the first *M. smegmatis* run.

These results imply that continued carry-over serves to significantly decrease output quality;

however, additional validation with controlled microbial community composition is necessary to

confirm this observation.

Areas of Future Development

This investigation has identified multiple necessary areas of future development to enable 16S

rRNA microbiome characterization on the Nanopore platform. First, strategies to exclude

between-run carryover, either via improved washing between runs or a barcode approach, would

enable multiple runs on the same flow cell, significantly reducing per-run costs. In this

investigation we performed six runs on the same flow cell while observing minimal output loss.

Second, improved bioinformatics strategies are necessary to exclude poor quality sequences. In

the *E. coli* and *P. fluorescens* runs, 3.3 and 9.0% of sequences, respectively, were not assigned to

any phyla, suggesting poor sequence quality and confounding both alpha- and beta-diversity

analyses. Finally, it would be beneficial to develop a 16S rRNA annotation pipeline based upon

optimized analysis strategies that provides output in near real-time, facilitating field and clinic

applications and alleviating current bioinformatics challenges from interested investigators.

Ultimately, the development of a rapid and low-cost microbiome approach will facilitate the

application of clinical and environmental microbiome technologies.

Acknowledgements

Support for this project was provided by the University of Pittsburgh Central Research

Development fund. KB was a member of the Oxford Nanopore Early Access Program, which

initially provided access to the sequencing platform at reduced cost.

References

Adams, R.I., Bhangar, S., Dannemiller, K.C., Eisen, J.A., Fierer, N., Gilbert, J.A., Green, J.L.,
Marr, L.C., Miller, S.L. and Siegel, J.A. (2016) Ten questions concerning the microbiomes of
buildings. Building and Environment 109, 224-234.
Altschul, S.F., Gish, W., Miller, W., Myers, E.W. and Lipman, D.J. (1990) Basic local alignment
search tool. Journal of Molecular Biology 215, 403-410.
Baron, J.L., Vikram, A., Duda, S., Stout, J.E. and Bibby, K. (2014) Shift in the Microbial
Ecology of a Hospital Hot Water System following the Introduction of an On-Site
Monochloramine Disinfection System. PLoS ONE 9(7), e102679.
Benítez-Páez, A., Portune, K.J. and Sanz, Y. (2016) Species-level resolution of 16S rRNA gene
amplicons sequenced through the MinION™ portable nanopore sequencer. Gigascience 5(1), 4.
Caporaso, J., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F., Costello, E., Fierer, N.,
Pena, A., Goodrich, J., Gordon, J., Huttley, G., Kelley, S., Knights, D., Koenig, J., Ley, R.,
Lozupone, C., McDonald, D., Muegge, B., Pirrung, M., Reeder, J., Sevinsky, J., Turnbaugh, P.,
Walters, W., Widmann, J., Yatsunenko, T., Zaneveld, J. and Knight, R. (2010a) QIIME allows
analysis of high-throughput community sequencing data. Nature Methods 7, 335 - 336.
Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K.,
Fierer, N., Pena, A.G., Goodrich, J.K. and Gordon, J.I. (2010b) QIIME allows analysis of high-
throughput community sequencing data. Nature methods 7(5), 335-336.
DeSantis, T.Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E.L., Keller, K., Huber, T., Dalevi,
D., Hu, P. and Andersen, G.L. (2006) Greengenes, a Chimera-Checked 16S rRNA Gene
Database and Workbench Compatible with ARB. Applied and Environmental Microbiology
72(7), 5069-5072.
Good, M., Sodhi, C.P., Ozolek, J.A., Buck, R.H., Goehring, K.C., Thomas, D.L., Vikram, A.,
Bibby, K., Morowitz, M.J., Firek, B., Lu, P. and Hackam, D.J. (2014) Lactobacillus rhamnosus
HN001 decreases the severity of necrotizing enterocolitis in neonatal mice and preterm piglets:
evidence in mice for a role of TLR9. American Journal of Physiology - Gastrointestinal and
Liver Physiology 306(11), G1021-G1032.
Greninger, A.L., Naccache, S.N., Federman, S., Yu, G., Mbala, P., Bres, V., Stryke, D., Bouquet,
J., Somasekar, S., Linnen, J.M., Dodd, R., Mulembakani, P., Schneider, B.S., Muyembe-Tamfum,
J.-J., Stramer, S.L. and Chiu, C.Y. (2015) Rapid metagenomic identification of viral pathogens in
clinical samples by real-time nanopore sequencing analysis. Genome medicine 7(1), 99.
Khoruts, A., Dicksved, J., Jansson, J.K. and Sadowsky, M.J. (2010) Changes in the composition
of the human fecal microbiome after bacteriotherapy for recurrent Clostridium difficile-
associated diarrhea. Journal of clinical gastroenterology 44(5), 354-360.

Kilianski, A., Haas, J.L., Corriveau, E.J., Liem, A.T., Willis, K.L., Kadavy, D.R., Rosenzweig,
C.N. and Minot, S.S. (2015) Bacterial and viral identification and differentiation by amplicon
sequencing on the MinION nanopore sequencer. Gigascience 4(12), 10.1186.
Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M. and Glöckner, F.O.
(2013) Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-
generation sequencing-based diversity studies. Nucleic Acids Research 41(1), e1-e1.
Kumar, P., Monin, L., Castillo, P., Elsegeiny, W., Horne, W., Eddens, T., Vikram, A., Good, M.,
Schoenborn, Alexi A., Bibby, K., Montelaro, Ronald C., Metzger, Dennis W., Gulati, Ajay S.
and Kolls, Jay K. Intestinal Interleukin-17 Receptor Signaling Mediates Reciprocal Control of
the Gut Microbiota and Autoimmune Inflammation. Immunity 44(3), 659-671.
Laver, T., Harrison, J., O'Neill, P., Moore, K., Farbos, A., Paszkiewicz, K. and Studholme, D.J.
(2015) Assessing the performance of the Oxford Nanopore Technologies MinION. Biomolecular
detection and quantification 3, 1-8.
Lax, S. and Gilbert, J.A. (2015) Hospital-associated microbiota and implications for nosocomial
infections. Trends in molecular medicine 21(7), 427-432.
Lipus, D., Vikram, A., Ross, D., Bain, D., Gulliver, D., Hammack, R. and Bibby, K. (In Revision)
Predominance and Metabolic Potential of Halanaerobium in Produced Water from Hydraulically
Fractured Marcellus Shale Wells. Applied and Environmental Microbiology.
Loman, N.J. and Quinlan, A.R. (2014) Poretools: a toolkit for analyzing nanopore sequence data.
Bioinformatics 30(23), 3399-3401.
Mikheyev, A.S. and Tin, M.M. (2014) A first look at the Oxford Nanopore MinION sequencer.
Molecular ecology resources 14(6), 1097-1102.
Pflughoeft, K.J. and Versalovic, J. (2012) Human microbiome in health and disease. Annual
Review of Pathology: Mechanisms of Disease 7, 99-122.
Pinto, A.J. and Raskin, L. (2012) PCR Biases Distort Bacterial and Archaeal Community
Structure in Pyrosequencing Datasets. PLoS ONE 7(8), e43093.
Prussin, A.J., II, Vikram, A., Bibby, K.J. and Marr, L.C. (2016) Seasonal Dynamics of the
Airborne Bacterial Community and Selected Viruses in a Children?s Daycare Center. PLoS ONE
11(3), e0151004.
Quick, J., Ashton, P., Calus, S., Chatt, C., Gossain, S., Hawker, J., Nair, S., Neal, K., Nye, K.
and Peters, T. (2015) Rapid draft sequencing and real-time nanopore sequencing in a hospital
outbreak of Salmonella. Genome Biol 16(114.2015), 10.1186.
Quick, J., Loman, N.J., Duraffour, S., Simpson, J.T., Severi, E., Cowley, L., Bore, J.A.,
Koundouno, R., Dudas, G., Mikhail, A., Ouédraogo, N., Afrough, B., Bah, A., Baum, J.H.J.,
Becker-Ziaja, B., Boettcher, J.P., Cabeza-Cabrerizo, M., Camino-Sánchez, Á., Carter, L.L.,
Doerrbecker, J., Enkirch, T., Dorival, I.G., Hetzelt, N., Hinzmann, J., Holm, T., Kafetzopoulou,
L.E., Koropogui, M., Kosgey, A., Kuisma, E., Logue, C.H., Mazzarelli, A., Meisel, S., Mertens,
M., Michel, J., Ngabo, D., Nitzsche, K., Pallasch, E., Patrono, L.V., Portmann, J., Repits, J.G.,
Rickett, N.Y., Sachse, A., Singethan, K., Vitoriano, I., Yemanaberhan, R.L., Zekeng, E.G.,
Racine, T., Bello, A., Sall, A.A., Faye, O., Faye, O., Magassouba, N.F., Williams, C.V.,
Amburgey, V., Winona, L., Davis, E., Gerlach, J., Washington, F., Monteil, V., Jourdain, M.,
Bererd, M., Camara, A., Somlare, H., Camara, A., Gerard, M., Bado, G., Baillet, B., Delaune, D.,
Nebie, K.Y., Diarra, A., Savane, Y., Pallawo, R.B., Gutierrez, G.J., Milhano, N., Roger, I.,
Williams, C.J., Yattara, F., Lewandowski, K., Taylor, J., Rachwal, P., J. Turner, D., Pollakis, G.,
Hiscox, J.A., Matthews, D.A., Shea, M.K.O., Johnston, A.M., Wilson, D., Hutley, E., Smit, E.,
Di Caro, A., Wölfel, R., Stoecker, K., Fleischmann, E., Gabriel, M., Weller, S.A., Koivogui, L.,

Diallo, B., Keïta, S., Rambaut, A., Formenty, P., Günther, S. and Carroll, M.W. (2016) Real-time,
portable genome sequencing for Ebola surveillance. Nature 530(7589), 228-232.
Shin, J., Lee, S., Go, M.-J., Lee, S.Y., Kim, S.C., Lee, C.-H. and Cho, B.-K. (2016) Analysis of
the mouse gut microbiome using full-length 16S rRNA amplicon sequencing. Scientific Reports
6, 29681.
Turnbaugh, P.J., Ley, R.E., Mahowald, M.A., Magrini, V., Mardis, E.R. and Gordon, J.I. (2006)
An obesity-associated gut microbiome with increased capacity for energy harvest. Nature
444(7122), 1027-1131.
Wang, Q., Garrity, G.M., Tiedje, J.M. and Cole, J.R. (2007) Naïve Bayesian Classifier for Rapid
Assignment of rRNA Sequences into the New Bacterial Taxonomy. Applied and Environmental
Microbiology 73(16), 5261-5267.

Table 1. Sequencing results and output from Oxford Nanopore MinION runs.

Sample	Sequencing Chemistry	Total Reads	Passed Reads (%)	Average Q-Score of Pass Reads (±SD)	Average Read Length (±SD)	Run Time (Min)
E. coli	2D	4093	1447 (35.4%)	11.95 ± 1.44	1304.23± 302.70	15
P. fluorescens	2D	7374	2671(36.2%)	13.73 ± 1.67	1277.99 ± 321.87	20
M. smegmatis[*]	2D	11627	4502(38.7%)	13.56 ± 1.7	1263.30 ± 300.29	20
Produced Water	2D	16525	5461 (33.0%)	13.79 ± 1.66	1320.96 ± 215.06	20
River Water	2D	9745	3143 (32.3%)	13.0 ± 1.76	1233.62 ± 385.52	20
M. smegmatis	1D	10383	7463 (71.9%)	7.82 ± 0.78	1414.44 ± 697.89	20

[*]*M. smegmatis* 2D, river water sample, and *M. smegmatis* 1D library were excluded from primary analyses due to apparent carryover from previous runs.

Table 2. Comparison of number of taxa at different phylogenetic levels for the produced water sample

	Phylum	Class	Order	Family	Genus

Number of taxa assigned using Nanopore data	5	9	13	17	20
Number of taxa assigned using Illumina data	11	21	36	63	84
Number of shared taxa	4	8	11	13	9
Total relative abundance of shared taxa (Nanopore)	99.9%	100.0%	99.9%	98.5%	98.3%
Total relative abundance of shared taxa (Illumina)	99.4%	98.0%	96.3%	81.8%	81.6%

Figure 1. Number of observed *de novo* operational taxonomic units (OTUs) per number of
sequences at different similarity thresholds for *E. coli* and *P. fluorescens* pure culture DNA
sample sequenced by Nanopore.

405   in

407   Figure 2. Accuracy of taxonomy assignment at different phylogenetic levels for (A) *E. coli*; and
408   (B) *P. fluorescens*

Figure 3. Jaccard (unweighted) and Bray-Curtis (weighted) dissimilarity values for a hydraulic
fracturing produced water sample analyzed by Nanopore and Illumina sequencing; distance
value of one indicating no community structure overlap, and a value of zero indicating identical
community structure

Figure 4. Taxonomy assignment of *Mycobacterium smegmatis* 16S rRNA pure culture
sequencing following *Pseudomonas fluorescens* sequencing.