Title: Histone 4 acetylation regulates behavioral individuality in zebrafish

Angel-Carlos Román[1,2,*,#], Julián Vicente-Page[1,2,*], Alfonso Pérez-Escudero[2,3], Jose M. Carvajal-González[4], Pedro M. Fernández-Salguero[4], Gonzalo G. de Polavieja[1,2,#]

[1]Champalimaud Neuroscience Programme, Champalimaud Centre for the Unknown, Avenida Brasília s/n, 1400-038 Lisbon, Portugal.

[2]Instituto Cajal, Consejo Superior de Investigaciones Científicas, Av. Doctor Arce, 37, 28002 Madrid, Spain.

[3]Physics Department, MIT, Cambridge, Massachusetts, USA

[4]Departamento de Bioquímica y Biología Molecular y Genética, Universidad de Extremadura. Av. de Elvas s/n, 06071 Badajoz, Spain.

[*] Contributed equally

[#]Correspondence to: angel.roman@neuro.fchampalimaud.org and gonzalo.polavieja@neuro.fchampalimaud.org

Keywords: Epigenetics, HDAC, YY1, behavioral variability, zebrafish

20
21
22
23 **ABSTRACT**
24

Behavioral individuality arises even in isogenic populations under identical environments,

but its underlying mechanisms remain elusive. We found that inbred and isogenic zebrafish

(*Danio rerio*) larvae showed consistent behavioral individuality when swimming freely in

identical wells or in reaction to stimuli. We also found that behavioral individuality

depends on the histone acetylation levels. Individuals with high levels of histone 4

acetylation behaved similar to the average of the population, but those with low levels

deviated and showed behavioral individuality. More precisely, we found behavioral

individuality to be related to individuality in histone 4 acetylation of a set of genomic

regions related to neurodevelopment. We found evidence that this modulation depends on a

complex of Yin Yang 1 (YY1) and histone deacetylase 1 (HDAC1). We suggest, using

stochastic modelling, that this complex is part of the molecular machinery giving

individuality in histone acetylation in neurodevelopmental genes ultimately responsible for

behavioral individuality.

INTRODUCTION

Classically, the phenotypic diversity of a population is considered to be generated by the genetic differences between its members and the disparity of their environmental influences[1]. A simple prediction from this view alone would then be that isogenic populations would not show variability when the environment is constant. Nevertheless, a pioneering study showed that there was variability independent of genetic differences in some morphological traits in mice raised in identical environments[2]. In recent years, similar results have been obtained for behavioral variability in mice and flies[3,4]. There are several mechanisms that might contribute to this effect, including developmental noise[5], maternal and paternal effects[6], or the different experiences the individuals obtain by interacting with the environment or other animals[4], among others.

Our knowledge about behavioral variability independent of genetic differences has increased substantially, but its underlying mechanisms remain unclear. Neuronal mechanisms such as neurogenesis, or serotonin signaling have been shown to be final targets of behavioral individuality[3,4], but the molecular mechanisms remain elusive. Chromatin modifications could be a promising mechanism to encode stable differences among individuals and they have been hypothesized as a potential mechanism for the generation of experience-dependent behavioral individuality[4]. DNA methylation differences have been associated to behavioral castes in honeybees[7], and they are necessary and sufficient to mediate social defeat stress[8]. Histone acetylation is another of the main epigenetic modifications[9] and it has been shown to regulate different behaviors such as mating preference in prairie voles[10] or cast-mediated division of labor in ants[11]. We thus reasoned that molecular mechanisms linked to epigenetic modifications could lead to behavioral individuality.

We used zebrafish from 5 to 8 days post fertilization (dpf) to dissect the molecular

substrates of behavioral individuality. Laboratory zebrafish larvae show individuality in

behavior[12] and they present some advantages such as its wide genomic information, the

simplicity of its pharmacological treatments and the possibility to do large-scale behavioral

analysis. Additionally, it is relevant to use a species in which we can observe directly

developmental changes, as some of the mechanisms responsible for behavioral individuality are

likely accumulated during development[13]. Here we established zebrafish larvae as a model for

the analysis of behavioral individuality to study individuality in free-swimming behavior. We

found that in our experimental tests behavioral individuality of zebrafish larvae is independent of

the genetic variability of the population but it is linked to histone acetylation differences.

RESULTS

Behavioral individuality in larval zebrafish is stable for days

We used three steps to establish zebrafish larvae as a model to study behavioral

individuality using a high-throughput setup (see **Methods and Extended Data Figure 1a-k** for

the custom-built video tracking software, downloadable from **www.multiwelltracker.es**. We

first obtained that each larvae showed differences in their spontaneous behavior. Simple eye

inspection of trajectories reveals this behavioral individuality (**Figure 1a-b,** left, 7-8 dpf,

respectively; **Extended Data Figure 2a-b,** left, 5-6 dpf). This can be quantified using two

parameters: overall activity (percentage of time in movement) and radial index (average relative

distance from the border towards the center of the well). These two parameters were chosen

because they are independent of each other (**Figure 1c**, P=0.98), while others like speed,

bursting or tortuosity correlated with activity (**Methods, Extended Data Figure 1l**, P<0.006).

We performed several control experiments to show that the observed individuality was not

affected by potential artifacts in the set-up (**Methods, Extended Data Figure 2c-e**). Also, we

tested that these behavioral parameters describe individuality also in response to stimuli like light

flashes, mechanical perturbation or at a novel tank (**Methods, Extended Data Figure 2f).**

In a second step, we showed that individual differences were robust along several days

(**Figure 1d,** R=0.69 and R=0.58, *P*<0.001 for linear correlation of activity and radial index,

respectively, 7 vs. 8 dpf; **Extended Data Figure 2g,** R=0.48 and R=0.41, *P*<0.01, 5 vs. 6 dpf).

The third step consisted in showing that inter-individual variability is larger than intra-individual

variability. In the two-dimensional phenotypic space defined by activity and radial index, it can

be directly seen that the area covered by the behavior of one individual is smaller than the area of

the whole population (**Figure 1e-f,** left, 7-8 dpf; **Extended Data Figure 2a-b** right, 5-6 dpf, see

Methods). Measuring variability by the Coefficient of Variation (CV), we found this difference

to be significant (**Methods, Figure 1g,** 8 dpf; **Extended Data Figure 2h-j**, 5-7 dpf respectively;

P<0.001 in all cases). Intra-individual and inter-individual variability levels remained stable

from 5 to 8 dpf (**Extended Data Figure 2k**, *P*<0.01 in all comparisons**).**

The degree of individuality may be measured by variability in the population. This can be

visualized using the probability density of finding an individual in a population with a given

mean activity and radial index (**Methods, Figure 1e-f**, right, 7-8 dpf; **Extended Data Figure 2a-**

b, right, 5-6 dpf). It can be quantified using *generalized variance*[14], a single parameter that

summarizes this two-dimensional variability, that we used to compare two populations

(**Methods**; **Extended Data Table 1** for summary of all results; other variability measures in

Extended Data Table 2).

Sources of behavioral variability in zebrafish

Our setup allowed us to perform high-throughput tests to study the possible origins of

behavioral individuality. Behavioral individuality could in principle depend on environmental

manipulations and the genetic differences across the population. Our experiments minimized

environmental influences by isolating eggs in plates at pharyngula stage (24 hpf) and by keeping

them at a controlled temperature (27-28ºC). Manual changes in water (24 hours before the

experiment) or feeding did not affect individuality (**Figure 2a,** P=0.42 and **Figure 2b**, P=0.38,

respectively).

We found that behavioral variability of a population did not depend on the genetic

variability of its individuals. Our control inbred WIK zebrafish population (F1) resulted from a

single batch of eggs retrieved from two adults with at least three cycles of inbreeding. We

obtained the same behavioral variability after two more inbreeding cycles (WIK F3, **Figure 2c**,

P=0.33) and in an isogenic population[15] (CG2, **Figure 2c**, P=0.44). Also, we did not find

changes in the behavioral variability using groups of siblings from genetically diverse outbred

parents (LPS line, **Figure 2c**, P=0.38).

Changes in chromatin acetylation alter behavioral variability

The absence of effects from genetic variability prompted us to test whether behavioral

individuality could be modified by different epigenetic factors. To test the contribution of DNA

methylation we used 5-azacytidine (AZA), an inhibitor of DNA-methyltransferases. We found

that AZA added to the water did not alter the behavioral individuality of a population (15 mM

AZA, **Figure 3a**, P=0.44) even if it reduced 3-methyl DNA in larval zebrafish (**Extended Data**

Figure 3a). We then studied the role of histone deacetylation, a reversible molecular process in

which an acetyl functional group is removed from specific residues of Histone 3 and 4[9]. This

system is regulated by a group of enzymes called Histone Deacetylases (HDACs). To test the

effect of HDACs on behavioral individuality, we used sodium butyrate (NaBu, a class I HDAC

inhibitor) at the standard concentration of 2 mM[16]. We first confirmed that NaBu increases the

level of total acetyl-histone 4 in larval zebrafish (**Extended Data Figure 3b**). We then found that

this treatment reduced the behavioral variability of a WIK F3 sibling population after 24 hours (2

mM NaBu, **Figure 3b**) compared to control PBS-treated larvae (PBS, **Figure 3b**, $P<0.001$).

Note that this treatment only altered variability and not the mean of the population parameters

($P=0.63$). When we retired the treatment, behavioral variability was recovered after additional

24h (**Extended Data Figure 3c**, $P=0.71$). Similarly to the behavior, the total levels of acetyl-

histone 4 increased with the treatment and were recovered 24 hours after retiring the treatment

(**Extended Data Figure 3b**). We also studied transgenic larvae with alterations in the

deacetylation pathway as an alternative more specific than the use of drugs. We found that

heterozygotic mutant populations of the histone deacetylase *hdac1* (*hdac1 +/-*) showed a reduced

behavioral variability compared to their AB controls, mirroring the results obtained with the

drugs (**Figure 3c**, $P=0.008$). Our results suggest that histone deacetylation pathway modulates

the behavior of zebrafish larvae without affecting its average behavior.

High acetylation levels result in a behavior close to the population average

We have shown that the degree of behavioral variability of a population depends on its

average acetylation levels. Since an increment in the global acetylation decreased the behavioral

variability without changing the average behavior, we reasoned that the individuals with higher

mean acetylation should be placed near the average population behavior in the phenotypic space.

To test this hypothesis, we performed an experiment with 90 individuals to obtain their

acetylation state depending on their distance to the average behavior of the population. As we

needed at least five larvae in order to get enough tissue for the experiment, we pooled 5 larvae

with very similar behavior and measured their acetylation state. We found that larvae whose

behavior was placed near the average of the population had higher mean acetylation values

(**Figure 4a**, left). To quantify the dependence between the average acetylation and the position in

the phenotypic space of the individuals, we first defined a coordinate system (centered on the

average behavior of the population) and then obtained two magnitudes for each pool of fish: their

distance to the center (r) and their angle with the horizontal axis (θ), **Figure 4a**, right. We found

that the acetylation levels of the individuals highly correlated with their phenotypic distance to

the average, while we found no correlation with their angular position (see **Figure 4b**, blue dots,

P<0.001, P=0.53, respectively). We found similar correlations when we analyzed the distance to

the mean of each behavioral parameter separately (**Extended Data Figure 4a**, P<0.001).

We have seen that individuals with higher acetylation levels display a behavior similar to

the average of the population, while the variability of the population behavior increases at lower

acetylation levels. This is compatible with our previous experiments that reduced the behavioral

variability of a population by increasing its acetylation levels. In fact, if we use fish treated with

NaBu to perform the same analysis, we find that the individuals not only have a behavior similar

to the average of the non-treated population, but also that their acetylation is at the same level of

the non-treated individuals with more acetylation (**Figure 4b**, red dots). This shows that the

treated animals present acetylation levels within the physiological range of the animals,

consistent with NaBu having the global effect of increasing the acetylation levels of the

population by bringing them close to the animals with more acetylation.

Differences in the acetylation of a set of genes are linked to behavioral variability

Our results link acetylation level of the individuals and their behavioral individuality.

However, fish with similar (low) acetylation levels also can show very different behavior, so

there must be other factors contributing to behavioral variability. We hypothesized that these

factors could be the acetylation differences in specific genomic regions associated to behavior.

To explore this possibility, we compared the acetylation variability between two groups of WIK

zebrafish, one with high and the other with low behavioral variability. For the first population,

we built four clusters of five sibling fish with low intra-cluster behavioral variability and high

inter-cluster variability, so that they covered the complete phenotypic space of the population

(clusters c1-c4 in **Extended Data Figure 4b**, see **Methods** for details). For the second

population, we built four behavioral clusters each made of up of five individuals selected at

random from the phenotypic space. This process eliminates any systematic difference in

behavior across clusters. We then retrieved the acH4 epigenomic profiles of the clusters in each

group using chIP-seq and computed the acetylation variability of each genomic region across

behavioral clusters using techniques adopted from gene expression analysis (see **Methods** for

details).

We found that there were more genomic regions with higher variability in acetylation

across clusters in the first population, suggesting that acetylation variability in the regions

correlated with behavioral variability (**Figure 5a,** $P<0.0001$). We then identified the epigenomic

regions with high variability in histone acetylation as they are potentially related to behavioral

variability (**Methods**, **Extended Data Table 3,** $P<0.01$). We found that genes located near these

regions are enriched in different Gene Ontology (GO) terms ($P<0.001$) mainly related to

neurobiological processes (**Figure 5b**). This chIP-seq analysis predicts that acetylation

variability in specific regions can be associated to behavioral differences. In order to assess the

specificity of these hypervariable regions in behavioral individuality, we checked that our

previous relation between histone acetylation levels and average behavior (**Figure 4b**) was

maintained in the hypervariable regions but not in the rest (see **Figure 5c**). To find if these

regions could have a causal action in behavioral individuality, we decided to affect them by

impairing DNA-interacting proteins that significantly bind near these regions. We found that

several DNA motifs that were enriched near the hypervariable regions were Yin-Yang binding

sites (*P*<0.0001, **Extended Data Figure 5c**). Yin-Yang 1 (YY1) is a transcription factor that can

activate or repress the same target gene depending on recruited co-factors[17], like HDAC1[18]. To

study the impact of YY1 on hypervariable regions, we selected eight of these regions. We first

confirmed that the acetylation variability seen in the chIP-seq experiment was maintained in

these regions using conventional chIP (**Extended Data Figure 5d**). Remarkably, the mRNA

expression of the genes located near these regions also showed a high variability (**Extended**

Data Figure 5d). Then, we studied the interaction between YY1 and these eight regions, and we

found that YY1 is significantly bound to them (**Figure 5d**). This is consistent with YY1 affecting

our hypervariable regions, so we decided to analyze their acetylation variability in a *yy1* +/-

mutant population. We found that *yy1* +/- fish showed a significant reduction in the acetylation

variability (**Figures 5e,** *P*<0.001). In addition, mRNA expression variability in near genes was

also reduced in *yy1* +/- fish compared to controls (**Figure 5f,** *P*<0.001). This reduction in

molecular variability led us to analyze the behavioral variability of the *yy1* +/- population, and

we found that the individuality was significantly reduced (**Figure 5g,** *P*=0.003). We thus

confirmed that hypervariable genes are related to behavioral and acetylation individuality by

using YY1 impairment as a method to alter the regulation of hypervariable regions.

YY1 and HDAC1 regulate histone acetylation individuality

We investigated further how YY1 acts on the hypervariable regions in search of a

possible mechanism for molecular variability. HDAC1 is a known partner of YY1[18] and we thus

tested how this partnership can be related to hypervariable regions. We first confirmed that

HDAC1 binds to the eight selected regions (**Figure 6a**). We then tested, using rechIP

experiments, that YY1 and HDAC1 bind together to these regions (**Figure 6b**), suggesting that

these two proteins form a complex that interacts with the hypervariable regions. As YY1 is

known to be regulated through deacetylation by HDAC1[18] , we assessed the effect of sodium

butyrate (NaBu) on YY1 acetylation in zebrafish larvae. Using co-immunoprecipitation, we

found that YY1 was acetylated in control conditions and that this acetylation was impaired after

NaBu treatment (**Figure 6c**). Furthermore, YY1 binding to the selected hypervariable regions

was decreased in NaBu-treated larvae (**Figure 6d**), suggesting that a functional YY1 is needed to

bind the hypervariable regions. A potential mechanism for the acetylation modulation consists in

YY1 guiding the YY1/HDAC1 complex to the hypervariable regions, which would then be

deacetylated by HDAC. Using a simple model that simulates stochastic binding of this complex

to the hypervariable regions (see **Figure 6e** and **Methods** for the details of the model), we found

that the stochasticity of the interactions is sufficient to generate variability in a population

composed by identical individuals. The model gives individuality at the level of mean acetylation

level and also in the concrete patterns of acetylation, here measured simply by the acetylation

variability of the genome of each individual. Specifically, there is not only individuality at the

average acetylation level but also at the acetylation variability level, considering it as a proxy of

the differences between individual acetylation profiles (**Figure 6f**).

DISCUSSION

In this paper, we have found that a histone acetylation pathway modulates individual behavior in a genetic-independent manner without affecting the global average behavior of the population. Histone acetylation levels of an individual correlated with and its individual behavior compared to the average of the population. Therefore, while the average behavior might depend more strongly on genetic background (as seen for different strains in **Figure 2**) or environmental changes (as seen for different responses in **Extended Data Figure 2f**), behavioral individuality could result from histone acetylation.

Several stochastic mechanisms can underlie behavioral individuality, such as paternal and maternal effects, differences in the experience received by individuals, developmental noise or stochastic DNA binding, among others. We propose the stochastic action of the complex formed by YY1 and HDAC1 as part of the molecular machinery that translates these factors into acetylation differences.

Another open question is how acetylation variability could lead to behavioral variability. A possibility is that histone acetylation is functionally transformed into changes in gene expression, as we have shown for eight of the hypervariable regions. Genes located near the hypervariable regions are significantly related to several neurodevelopmental processes, so differences in their expression might result in differences in brain development and then ultimately in behavior.

METHODS

Zebrafish lines and care

Zebrafish (*Danio rerio*) WIK strain[19] was kindly provided by Dr. Bovolenta (CBM-

UAM) and inbred in our laboratory for at least three generations before the experiments.

Afterwards, WIK F1 population was generated from a single batch of embryos from a single

couple of adult fish. Two additional cycles of inbreeding were carried out, crossing a couple of

siblings from the former generation. CG2 clone population, generated by double gymnogenetic

heat-shock, and characterized by being pure isogenic zebrafish was kindly provided by Dr.

Revskoy (Univ Northwestern) as a control of reduced genetic differences between siblings. The

outbred LPS (Local Pet Store) strain was recently described[20], and used as a model of genetic

heterogeneity. Heterozygotic *hdac1* and *yy1* mutant strains with wild-type counterparts were

obtained from ZIRC.

Care and breeding of the zebrafish strains were as described[20], with specific additional

details. Eggs were isolated after 24 hours post-fertilization, and maintained in custom multiwell

plates until 10 days post-fertilization (dpf). They were fed (JBL NovoBaby) from 6 dpf and

water was changed daily if it is not indicated specifically in the experiment.

All the experiments using animals were approved and performed following the guidelines

of the CSIC (Spain) and the Fundaçao Champalimaud (Portugal) for animal bioethics.

Free-swimming setup and recording

The setup consists of a monochrome camera located over the wells at a distance of 70 cm

and pointing downwards. The camera used was a 1.2 MPixel camera (Basler A622f, with a

Pentax objective of focal length 16 mm). The wells are circular, carved on transparent PMMA

(24 wells per plate, and typically two plates are recorded simultaneously), and have their walls

tilted so that even in the most lateral wells the wall never hides the larva from the camera. Each

well is 15 mm deep, and has a diameter of 1.8 mm at the bottom and a diameter of 30 mm at the

top (**Extended Data Figure 1a**). For the experiments, each well is filled with a volume of 3 ml.

The dishes are supported by a white PMMA surface that is only partially opaque. Behind this

white surface we place two infrared led arrays (830nm, TSHG8400 Vishay Semiconductors)

pointing outwards (**Extended Data Figure 1a**). Two paper sheets stand between the lights and

the central space that lies directly under the wells. With this disposition we ensure that only

diffuse indirect light reaches the wells, so that the illumination is roughly uniform (most of the

light comes from below the wells through the white surface). All the set-up is surrounded by

white curtains. Video camera recorded at a 25 fps rate (**Extended Data Figure 1b-c** for

examples of a single frame and final trajectories).

A larval population (5-8 dpf) consisted of at least 24 fish siblings from the same batch of

embryos. After five minutes of acclimation to the new environment, the larvae were recorded for

20 minutes. Water temperature was maintained in a strict range (27-28 ºC) during each

experiment.

Custom-built software tracking larvae

We developed multiwellTracker, a software to automatically track zebrafish larva in

wells. The software is available at http://www.multiwelltracker.es.

Detection of wells

The program is prepared to auto-detect circular wells, regardless of their spatial

arrangement. To detect the wells we use the circular Hough transform (we have modified the

code of Tao Peng distributed by Matlab Central under BSD license). In order to estimate the

diameter of the wells, it computes the image's Hough transform for 100 radii different in 5 pixels

and a rough estimate of the largest possible radius (length of the longer side of the image divided

by the square root of the number of wells) (**Extended Data Figure 1d**). The system selects the

highest point of this measure as an estimate of the radius of the wells (r_{est}). It is possible to skip

this first step and instead specify manually a value for r_{est}. This may be advisable when many

videos are recorded with the same set-up and the same wells.

In the second step the system locates the centers of the wells. To do this it performs a

Hough transform of the original image, this time with radii only in the range between $0.8r_{est}$ and

$1.2r_{est}$. The transformed image usually has clear peaks in the centers of the wells. Then it filters

the transformed image with a Gaussian filter to increase its smoothness (the resulting

transformed image is shown in **Extended Data Figure 1e**). Then, it selects the maximum of the

transformed image as the center of the first well. To prevent selecting the same well twice, the

system discards all the pixels of the transformed image that are within radius r_{est} of the selected

center (**Extended Data Figure 1f**). It selects the new maximum as the center of the second well,

and repeats the procedure until all wells have been found (**Extended Data Figure 1g**). The

experimenter can correct the result by manually clicking on the center of the wells that have not

been correctly located (<1% of cases).

Pre-processing of images

In order to control for fluctuations in illumination, each frame is normalized by dividing

the intensity of each pixel by the average intensity across all pixels of the frame. After

normalizing the frame, a 2D Gaussian filter is used to smooth the image (**Extended Data Figure**

1h shows the image before and **Extended Data Figure 1i** after filtering).

Background subtraction and detection of the larva

In order to extract the image of the larva from the background, the system finds the

average of 1,000 frames equi-spaced along the whole video. This average image is what we will

call "static background". By subtracting each frame by the static background, we obtain an

image in which the larvae correspond to dark regions (**Extended Data Figure 1j**). However,

because of relatively slow changes in the set-up over time, the system uses the static background

in combination with a dynamic background, which is computed as the average of the previous 5

frames. The difference between the current frame and the dynamic background will only show

larvae that are moving in that precise moment (**Extended Data Figure 1k**).

The specific algorithm to detect the larva is as follows. First, the difference between the

current frame and the static background is thresholded keeping only pixels for which the

difference is below -0.5. We then find connected components ("blobs") in this thresholded

image, keeping those that are larger than 1 pixel. Because these blobs come from the difference

with the static background, both static and moving larvae will be detected. But at this stage some

blobs come simply from noise. In order to filter out noisy blobs, the system accepts a blob if it

fulfills at least one of these two conditions: (a) It contains at least one pixel that was identified as

part of the larva in the previous frame or (b) it contains at least one pixel for which the difference

between the current frame minus the dynamic background is below the same threshold as before

(-0.5).

Removal of reflections

In most cases only one blob is obtained after the process described in previous sections.

But when the larva is close to the wall of the well, its reflection on the wall may also be selected.

The system considers that a blob A is a reflection of blob B when all of the following conditions

are met: (a) Blob B is bigger than blob A, (b) blob B is closer to the center of the well than blob

A and (c) the lines between the center of the well and the two blobs form an angle $< 10°$. When

these three conditions are met, the system removes blob A.

Acquisition of the position of the larvae

If more than one blob remains in the same well after the previous steps, the system

selects the one with highest contrast. For the selected blob, the system takes the position of its

most contrasted pixel, and adds this position to the trajectory of the larva. If in a well no blob

remains after the previous steps, the trajectory is left with a gap. When this happens, the program

will not re-track the larva until it moves.

Behavioral Parameters

Different parameters reflecting the behavior of individual larvae were measured, and

finally two of them were used through the paper: (i) activity (percentage of time in movement)

and (ii) radial index (average position from the border towards the center of the well). We also

studied three additional parameters: (i) Tortuosity in the trajectory was calculated as the scalar

product of the velocity vectors between two consecutive frames and the value in **Extended Data**

Figure 1l was obtained by averaging this parameter through the whole video, excluding the

frames where the animal was immobile. (ii) Speed was calculated as the average distance (in

pixels) travelled per frame, in those frames where the fish was active. (iii) Bursting was obtained

as the total number of frames where fish changed from immobility to motion. We found that

these three parameters correlated with activity.

The average of each individual parameter was tested from 5 to 8 days post-fertilization

(dpf) to assess if individual behavior was significantly stable along the days using Pearson

coefficient of correlation.

Additional validation of the experimental setup

Several controls were performed for possible experimental artifacts affecting wells differently. Behavioral parameters were robust to 90 degrees counterclockwise rotations of the multi-well plate (**Extended Data Figure 2c,** left**,** R=0.73, *P*<0.001, and R=0.68, *P*<0.001; for linear correlation tests of individual behavior) or to interchanging the larvae between outer and inner wells (**Extended Data Figure 2c,** right**,** R=0.65, *P*<0.001, and R=0.61, *P*<0.001; using the same correlation test). Also, we found no correlation between the small differences in illumination across wells and behavior (**Extended Data Figure 2d**). We further corroborated using a significance test that the differences in behavior did not have an origin in systematic differences across wells. For this, we found that the average behavioral parameters obtained in fifteen individual experiments were not different between wells (**Extended Data Figure 2e**, typically *P*=0.4 and always *P*>0.19 for both parameters).

Stimulus response tests

We studied the influence that our free-swimming behavioral parameters could have on the performance of the individuals when they respond to three different stimuli.

Response to mechanical disturbance

We applied mechanical perturbations to each larva by pipetting up and down the water content of the well for four times. Perturbations were applied at 6 dpf to previously recorded animals, and the 20-minute recording was done at 7 dpf. The recording was performed in the usual setup.

Response to strong Light Pulse

In complete darkness, we applied three different light flashes to the larvae and study their

behavior in the 90 subsequent seconds. The flashes and the recording were performed in the

usual setup. Pre-recording behavioral parameters were obtained the day before.

Novel tank with light/dark preference

In order to study the effect that a novel setup could have on the behavior of larvae we

built a rectangular setup, which changed the geometry of the previous circular wells. The setup

dimensions were 84 mm x 21 mm and it was built in transparent acrylic. To try to see if our

parameters had any effect on the light-dark preference, half of the floor of the setup (42 mm x 21

mm) was white while the other half was black. The height of the setup was 5 mm. Larvae were

placed in the center of the white part and recorded for 10 minutes. Activity was calculated as

previously described and distance to the wall was represented by the average distance to the

longest walls, normalized to 1 in the middle point of both walls and to 0 in the exact position of

the walls.

The effect of our behavioral tests resulted in a decrease (increase) in mean activity (radial

index), but maintaining the same individuality of the pre-recorded free-swimming experiments

(**Figure S2f;** *P*<0.04 for changes in mean activity and radial index compared to control larvae of

the same age; *P*<0.02 for linear correlation of activity and radial index. In the case of novel tank,

radial index cannot be applied because the wells are elongated and was then replaced by the

minimum distance to the longer walls. We note, however, that this parameter showed no

correlation with the radial index of pre-recorded experiments in the same animals.

Inter-individual vs. Intra-individual differences

The behavioral parameters (activity and radial index) were also obtained from

consecutive fragments of 30 seconds for each 20-minute experiment for each larva. This was

fitted to a two-dimensional Gaussian, but for clarity when representing many animals (like in

Figure 1e-f, left) an isocontour of the Gaussian for each animal was used. An isocontour is an

ellipse with principal axes given by the eigenvectors of the covariance matrix. We chose the

isocontour with length of each semiaxis given by the square root of the eigenvalue of the

covariance matrix, as this reduces to the standard deviation in each direction for cases with no

correlation between the two variables. Intra-individual variation distribution was obtained using

the coefficients of variations (CVs) for activity and radial index separately. Inter-individual

variation was calculated the same way but using fragments from different fish.

Comparing the behavioral variability between two animal groups

A simple visual method to characterize the variability in a population is to plot the bi-

dimensional distribution of the activity and radial index of individuals (like in **Figure 1e-f,**

right). To do so, we used Gaussian kernel smoothing that consists in adding up Gaussians

centered at the data points as

$$P(x,y) = \frac{1}{N} \frac{1}{2\pi\sigma_x\sigma_y} \sum_{i=1}^{N} \exp\left(-\frac{1}{2}\left[\frac{(x-x_i)^2}{\sigma_x^2} + \frac{(y-y_i)^2}{\sigma_y^2}\right]\right)$$

with x_i and y_i the mean activity and radial index values of individual i of a total of N members of

the population. An optimal smoothing uses standard deviations of each Gaussian given by

$\sigma_x = N^{-1/6}\alpha_x$ with α_x the standard deviation in the x_i data values, and similarly for σ_y using the

y_i values   (see   B.E.   Hansen,   unpublished   manuscript,

http://www.ssc.wisc.edu/~bhansen/718/NonParametrics1.pdf). The volume under the probability

surface has a value of 1, even when the values of the probability density are already up to 90.

The probability surface sits on an area on the x-y plane of approximately 0.4x0.4, making the

total volume under the surface to be 1.

While this distribution gives a visual and intuitive characterization of behavioral

variability, an even simpler characterization is achieved using, for each group, a single parameter

summarizing its two-dimensional variability. We used *generalized variance* as this single

parameter, measured as the determinant of the covariance matrix (**Extended Data Table 1**),

while other parameters like the standard deviation for each parameter gave similar statistical

results (**Extended Data Table 2**).

ChIP-seq, conventional chIP, and rechIP analyses

Clusters of four fish were built for studying the molecular variability of a population, as

the amount of tissue in a single larva was not sufficient to measure histone acetylation levels. In

addition, the clustering also reduced the noise that could arise from the molecular analysis of a

single larva. In the first (behavioral) cluster, a Hierarchical Clustering analysis using Euclidean

distance as the metric and the average linkage clustering as the linkage criteria was used the

clustering in the chIP-seq, as the total population consisted of 72 larvae. The selection in the

NaBu experiment used the same algorithm. In the random experiment, fish were randomly

selected from the population.

The samples were crosslinked with 1.8% formaldehyde for 30' and then quenched with

1% glycine for 5'. Extracts were lysed using a SDS Lysis buffer (50 mM Tris-HCl pH 8.1, 1%

SDS, 10 mM EDTA) for 30' at 4ºC, and then diluted with a Dilution buffer (6.7 mM Tris-HCl

pH 8.1, 0.01% SDS, 1.2 mM EDTA, 1.1% Triton X-100, 167 mM NaCl). 2 mM sodium butyrate

was added to avoid histone deacetylation activity during the preparation. Then, the fish were

sonicated with two pulses (30" ON / 30" OFF) of 15' each with the Diagenode Bioruptor. Before

pre-clearing the samples with protein A/G beads, an input sample was obtained. Then, the

extracts were immunoprecipitated overnight using 1 μg of the anti- acetyl-Histone 4, anti-

GAPDH, anti-HDAC1 or anti-YY1 antibodies. Bound DNA was recovered with protein A/G

beads, then washed with Low-Salt (120 mM Tris-HCl pH 8, 0.1% SDS, 2 mM EDTA, 1% Triton

X-100, 150 mM NaCl), High-Salt (120 mM Tris-HCl pH8, 0.1% SDS, 2 mM EDTA, 1% Triton

X-100, 500 mM NaCl), LiCl (10 mM Tris-HCl pH 10, 1 mM EDTA, 0.25 M LiCl, 1% NP40,

1% sodium deoxycholate) and two times with 1X TE (10 mM Tris-HCl pH 8, 1 mM EDTA)

buffers, and recovered with Elution (1% SDS, 0.1 M NaHCO3). DNA purified samples were de-

crosslinked using sodium chloride, and cleared with Qiagen spin columns. In the case of rechIP,

the samples were reincubated with the second antibody after elution, and another round of

washes were performed.

The final samples were processed at the Genomics Unit at the Scientific Park of Madrid

in the case of chIP-seq experiments. Libraries were built, and the samples were sequenced using

an Illumina GAII. Raw data of the experiment can be obtained in the NCBI GEO repository

(GSE IDXXX). Reads were aligned to *Danio rerio* genome sequence (Zv7) with BWA, and final

reads in a 25-bp window were mapped to the reference genome using custom Perl scripts. The

repetitive regions were removed from the analysis and the results were also normalized using the

average number of reads in each sample. The standard deviation of every 25 bp region for each

experiment (Behavior, Random and NaBu) was calculated, and then we quantified the

probability for each region of being more variable than in the Random experiment. We selected

the regions with $P<0.01$ and with the highest standard deviation (top 25%). Nearest genes were

retrieved, and their human orthologs were analyzed for Gene Ontology. In the case of

conventional chIP or rechIP, qPCR analyses for amplification differences using specific primers

were performed. In addition, hypervariable acetylated regions with their flanking sequences (300

bp) were used to predict enriched DNA motifs and their potential biological activity with MEME

suite[21].

Reagents and antibodies

Sodium butyrate and AZA (Sigma-Aldrich) was dissolved in Phosphate-buffered saline

(PBS), and used in a final 2 mM and 15 mM concentration of fish water, respectively. PBS alone

was used as vehicle control. The pharmacological treatment lasted for 24 hours from 7 dpf to

8dpf. Acetyl-Histone 4 antibody was obtained from Promega, anti-HDAC1 from Epigentek, anti-

YY1 antibody from SantaCruz, anti-GAPDH from Sigma-Aldrich, and McrBC enzyme from

New England Biolabs.

Western Immunoblotting

Treated and untreated groups of fish (5-10) were frozen at different times, and then

protein extracts were isolated from tissue using an extraction buffer (80 mM Tris-HCl pH 7.5, 2

mM EDTA, 2 mM EGTA, 0.27 M Saccharose, 10 mM β-glycerolphosphate, 5 mM Sodium

pyrophosphate, 50 mM Sodium Fluoride, 1% Triton X-100, 0.1 mM Sodium vanadate, 0.1% β-

Mercaptoethanol, 1X Complete protease inhibitor cocktail) during 30' at 4ºC with vortexing.

After centrifugation, debris was removed, and the protein content was measured with Coomasie

reagent (Pierce). Anti- acetyl-Histone 4, anti- ß-Actin and secondary antibodies were used

following manufacturer recommendations.

RNA isolation and qPCR quantification

Total RNA was isolated using homogenized extracts from three fish per sample by Trizol

(Life Technologies) extraction and RNAeasy (Qiagen) purification. Retrotranscription was done

with iScript (Bio-Rad) following manufacturer recommendations. Finally, quantification of the

target genes was measured using qPCR with specific primers, and p-values obtained by using

Student's T-test.

Quantification of Histone 4 acetylation levels

Eighteen clusters of five fish from a total population of 90 were obtained from the

behavioral space (activity/radial index) using an *ad hoc* algorithm. First, 18 centroids were

randomly chosen, and 5 individuals were assigned to the nearest (not occupied) centroid. Then,

centroids were redefined using the average values of the new clusters, and a new round of

assignment of the fish to the centroids was done. This iteration was repeated until the centroids

were stable. Then, acetyl-H4 levels were quantified using Epigentek kit and following

manufacturer recommendations.

Quantification of methylated DNA

DNA methylation was quantified using larval DNA digested by MCrBC enzyme as

previously done[22] following kit instructions.

Simulation of YY1/HDAC1 complex activity

We simulated how the YY1/HDAC1 complex could stochastically deacetylate

epigenomic regions. Regions can be considered as transcription factor binding sites (we used

Nreg=1,000 regions **Figure 6f**). Each region starts with an initial acetylation value given by a

Gaussian distribution with a fixed mean acetylation and a standard deviation. The value of the

mean μ is arbitrary as long as the relation of the rest of the parameters with this one is fixed and

the value of the standard deviation is *μ/10* (in **Figure 6f** we used *$\mu=10$*). We fixed the number

molecules of the complex to be the same for each fish (*Complex molecules=3,000*). The complex

binds to randomly selected regions and, once bound, the complex decreases the acetylation of

each region by an amount of $0.3 \cdot \mu$. We allowed several complexes to bind at each region. The

complex has an error rate of 10% so, once bound, the probability of deacetylating is 0.9. This

error rate has the final consequence of generating different mean and variance acetylation values

for the individuals of a population (**Figure 6f**). The minimum acetylation value for each region is

0 and, once reached, the complex cannot select that region for further binding. We repeated the

whole process for 100 fish, each one representing a point in **Figure 6f**.

The pseudocode of the simulation would be as follows:

560   **while** *Complex molecules > 0*

561   *Complex binds to a region r, p(binding)=1/Nregions*

562   *Complex deacetylates region r by an amount of $0.3 \cdot \mu$, with a probability p=0.9*

563   *YY1 molecules=YY1 molecules-1*

564   **endwhile**

Statistical analysis

All statistical tests to compare the differences between two distributions were conducted by calculating the value of their representative parameter, shuffling randomly the data of both distributions for 1,000 - 10,000 times and computing a P-value given by the proportion of times in which the difference in the representative parameter of the random distributions was higher than their original value. All the experiments were done at least three times with different biological datasets, and P-values were calculated using the three replicas. Figures show a representative experiment of the triplicate. MATLAB was used for all the computations and the statistical analysis.

AUTHOR CONTRIBUTIONS

A.-C.R. designed project, performed experiments and analysis and wrote the paper, J.V.-P. performed experiments and analysis and wrote the paper, A.P.-E. developed the tracking software, J.M.C.-G. performed experiments, P.M.F.S provided reagents, and G.G.d.P. designed and supervised project, performed analysis and wrote the paper.

ACKNOWLEDGEMENTS

We thank Rui Costa, Carlos Ribeiro, Antonia Groneberg, and other members of the Champalimaud Neuroscience Programme for discussions, and Isidro Dompablo (CSIC) and Ana Catarina Certal (Champalimaud Foundation) for technical assistance and animal care. CG2 and WIK lines were kind gifts from Sergei Revskoy and Paola Bovolenta, respectively. This work was supported by a Juan de la Cierva and a FEBS Long-Term fellowship (to A.-C.R.), a JAE-CSIC Predoctoral Fellowship (to J.V.P.), Spanish Plan Nacional Ministerio de Economia y Competitividad grants BFU2014-54699-P (to J.M.C.-G.), BFU2011-22678 (to P.M.F.S),

595
REFERENCES:

1.   Galton, F. *English men of science, their nature and nurture*. (1874).

2.   Gärtner, K. A third component causing random variability beside environment and
genotype. A reason for the limited success of a 30 year long effort to standardize
laboratory animals? *Lab. Anim.* **24,** 71–77 (1990).

3.   Kain, J. S., Stokes, C. & de Bivort, B. L. Phototactic personality in fruit flies and its
suppression by serotonin and white. *Proc. Natl. Acad. Sci. U. S. A.* **109,** 19834–9 (2012).

4.   Freund, J. *et al.* Emergence of individuality in genetically identical mice. *Science* **340,**
756–9 (2013).

5.   Waddington, C. H. The strategy of the genes. A discussion of some aspects of theoretical
biology. With an appendix by H. Kacser. *Strateg. genes A Discuss. some ...* ix +-262 .
(1957). doi:10.1007/3-540-32786-X_7

6.   Seong, K. H., Li, D., Shimizu, H., Nakamura, R. & Ishii, S. Inheritance of stress-induced,
ATF-2-dependent epigenetic change. *Cell* **145,** 1049–1061 (2011).

7.   Herb, B. R. *et al.* Reversible switching between epigenetic states in honeybee behavioral
subcastes. *Nat. Neurosci.* **15,** 1371–1373 (2012).

8.   Laplant, Q. *et al.* Dnmt3a regulates emotional behavior and spine plasticity in the nucleus
accumbens. *Nat. Neurosci.* **13,** 1137–1143 (2010).

9.   Grunstein, M. Histone acetylation in chromatin structure and transcription. *Nature* **389,**
349–352 (1997).

10. Wang, H., Duclot, F., Liu, Y., Wang, Z. & Kabbaj, M. Histone deacetylase inhibitors
facilitate partner preference formation in female prairie voles. *Nat. Neurosci.* **16,** 919–24
(2013).

11. Simola, D. F. *et al.* Epigenetic (re)programming of caste-specific behavior in the ant
Camponotus floridanus. *Science (80-.).* **351,** (2015).

12. Pantoja, C. *et al.* Neuromodulatory Regulation of Behavioral Individuality in Zebrafish.
Neuron **91,** 587–601 (2016).

13. Fraga, M. F. *et al.* Epigenetic differences arise during the lifetime of monozygotic twins.
Proc. Natl. Acad. Sci. U. S. A. **102,** 10604–10609 (2005).

14.   Wilks, S. S. Certain generalizations in the analysis of variance. *Biometrika* **24,** 471–494
626   (1932).

627   15.   Mizgirev, I. V. & Revskoy, S. A new zebrafish model for experimental leukemia therapy.
628   *Cancer Biol. Ther.* **9,** 895–903 (2010).

629   16.   Heruth, D. P., Zirnstein, G. W., Bradley, J. F. & Rothberg, P. G. Sodium butyrate causes
630   an increase in the block to transcriptional elongation in the c-myc gene in SW837 rectal
631   carcinoma cells. *J. Biol. Chem.* **268,** 20466–20472 (1993).

632   17.   Shi, Y., Seto, E., Chang, L. S. & Shenk, T. Transcriptional repression by YY1, a human
633   GLI-Krüppel-related protein, and relief of repression by adenovirus E1A protein. *Cell* **67,**
634   377–388 (1991).

635   18.   Yao, Y. L., Yang, W. M. & Seto, E. Regulation of transcription factor YY1 by acetylation
636   and deacetylation. *Mol. Cell. Biol.* **21,** 5979–5991 (2001).

637   19.   Nechiporuk, A., Finney, J. E., Keating, M. T. & Johnson, S. L. Assessment of
638   polymorphism in zebrafish mapping strains. *Genome Res.* **9,** 1231–1238 (1999).

639   20.   Pérez-Escudero, A., Vicente-Page, J., Hinz, R. C., Arganda, S. & de Polavieja, G. G.
640   idTracker: tracking individuals in a group by automatic identification of unmarked
641   animals. *Nat. Methods* **11,** 743–8 (2014).

642   21.   Bailey, T. L. *et al.* MEME Suite: Tools for motif discovery and searching. *Nucleic Acids*
643   *Res.* **37,** (2009).

644   22.   Torres-Núñez, E. *et al.* Matricellular protein SPARC/osteonectin expression is regulated
645   by DNA methylation in its core promoter region. *Dev. Dyn.* **244,** 693–702 (2015).

652
653
Figure 1. **Behavioral individuality in a population of 48 larval zebrafish.** **(a)** Example 20-

minute trajectories for the group at 7 dpf (left), zoomed in for four fish (right). **(b)** The same

group at 8 dpf. **(c)** Radial index *vs*. activity at 7 dpf. **(d)** Correlation of activity (blue) and radial

index (red) between 7 dpf and 8 dpf for the same group. **(e)** Left: Population variability in

activity and radial index of the same group at 7 dpf. Each ellipse represents the behavioral

variability for each single fish as described in **Methods**. Colors as in **a**. Right: Probability

density of finding an individual with a given mean activity and radial index. **(f)** Same as **e** but at

8 dpf. **(g)** Smoothed histogram of the intra-individual variability (red) and inter-individual

662 variability (blue) of the same group at 8 dpf, measured as CV of activity (left) and radial index

663 (right).

Figure 2. **Environmental changes and genetic background do not have an impact on behavioral variability. (a)** Probability density of finding an individual with a given mean activity and radial index for additional larval groups with and without daily water changes, at 7 dpf. **(b)** Same as **a**, but for additional daily fed and non-fed groups. **(c)** Same as **a** for additional groups with different genetic backgrounds: WIK F1 (three inbreeding cycles), WIK F3 (five inbreeding cycles), CG2 (gymnozygotic fish clones) and LPS (outbred parents).

Figure 3. **Histone acetylation modulates behavioral variability.** **(a)** Probability density map

for fish treated with a PBS solution as control and AZA. (**b**) The same for PBS and NaBu. **(c)**

Probability density map for *hdac1 +/+* and *hdac1 +/-* larvae.

Figure 4. Relation between acetylation levels and behavior. (**a**) Average acetylation levels of fish depending on their behavior (left). Schematic representation of the two parameters used to analyze the dependence between acetylation and behavior (right). (**b**) Relation between the values of histone 4 acetylation and the two parameters of the coordinate system centered on the average behavior of the population: the distance to the average (left) and the angle with the horizontal axis (right).

Figure 5. **Hypervariable acetylation regions related to behavioral individuality**. **(a)** Probability distribution of the variability (SD) in the acetylation of the behavioral (blue) and random control (red) clusters, obtained with chIP-seq. Horizontal line, $P<10^{-4}$. **(b)** Gene ontology of the hypervariably acetylated regions. **(c)** Relation between normalized number of reads in each cluster of the chIP-seq experiment and their distance to average behavior, as in **Figure 4b**. Blue dots represent the average of the hypervariable regions and red dots represent the average of the rest of the regions. **(d)** YY1 and GAPDH binding to eight selected hypervariable regions.

697   **(e)** Comparison between acH4 variability of the eight regions in *yy1* +/+ and *yy1* +/-

698   populations. **(f)** The same as **e**, but for mRNA expression variability of genes located near the

699   regions. **(g)** Probability density map for *yy1* +/+ and *yy1* +/- larvae.

Figure 6. **YY1 and HDAC1 regulate acetylation individuality**. (**a**) YY1 and GAPDH binding

to eight selected hypervariable regions. (**b**) RechIP binding of YY1/HDAC1 and YY1/GAPDH

to the same regions. (**c**) Co-immunoprecipation of acetyl-lysine and YY1 in larvae treated with

PBS and NaBu, respectively. (**d**) YY1 binding to the eight regions in larvae treated with PBS and

NaBu. (**e**) Schematic model of the stochastic action of YY1/HDAC1 complex on a set of

epigenomic regions. All regions have the same binding probability, given by $p(b)=1/n_r$, where n_r

is the total number of regions .(**f**) Correlation between the acetylation variability (measured as

the coefficient of variation of the acetylation across all regions) and the average acetylation of

the individuals of a population.