The RAG transposon is active through the deuterostome evolution and domesticated in jawed

vertebrates.

Jose Ricardo Morales Poole [1], Sheng Feng Huang [2], Anlong Xu [2,3], Pierre Pontarotti [1]

Jose Ricardo Morales Poole and Sheng Feng Huang contributed equally to the work.

1 Aix Marseille Université, CNRS, Centrale Marseille, I2M UMR 7373, équipe évolution

biologique modélisation, 13453, Marseille, France.

2 State Key Laboratory of Biocontrol, Guangdong Key Laboratory of Pharmaceutical Functional

Genes, School of Life Sciences, Sun Yat-sen University, Guangzhou, 510275, People's Republic of

China.

3 Beijing University of Chinese Medicine, Dong San Huan Road, Chao-yang District, Beijing,

100029, People's Republic of China.

Abstract

RAG1 and RAG2 are essential subunits of the V(D)J recombinase required for the generation of the

variability of antibodies and T-cell receptors in jawed vertebrates. It was demonstrated that the

amphioxus homologue of RAG1-RAG2 is encoded in an active transposon, belonging to

transposase DDE superfamily. We show here that the RAG transposon has been active through the

deuterostome evolution and is still active in several lineages. The RAG transposon corresponds to

several families present in deuterostomes. RAG1-RAG2 V(D)J recombinase evolved from one of

them, partially due to the new ability of the transposon to interact with the cellular reparation

machinery. Considering the fact that the RAG transposon survived millions of years in many

different lineages, in multiple copies and that DDE transposases evolved many times their

association with proteins involved in repair mechanisms, we propose that the apparition of V(D)J

recombination machinery could be a predictable genetic event.

Introduction

The recombination-activating gene products known as RAG1 and RAG2 proteins constitute the

enzymatic core of the V(D)J recombination machinery of jawed vertebrates. The RAG1-RAG2

complex catalyzes random assembly of variable, diverse and joining gene segments that are present

in the jawed vertebrate genome in numerous copies and together with hyper-mutation generate the

enormous diversity of the assembled antibodies and T-cell receptors. Therefore, RAG1-RAG2 role

in the V(D)J rearrangement of antigen receptors is crucial for the jawed vertebrate adaptive

immunity [1]. Concerning the origins of RAG1-RAG2 remains elusive for more than 30 years as the

genes were only found in jaw vertebrates [2]. On the other hand, striking similarities between RAG1

and DDE transposase has been noted: common reaction chemistry for DNA cleavage, similar

organization of protein domain structure and sequence similarities between recombination signal

sequences (RSSs) and terminal inverted repeat (TIRs) targeted by transposases [3, 4]. The hypothetical

transposon ancestry of RAG was further supported upon the demonstration of RAG1-RAG2

mediated transposition *in vitro* [5, 6] and *in vivo* [7-10], thought the efficiency of such reactions in vivo is

highly disfavored comparing to recombination.

A next step in the understanding of RAG1-RAG2 recombinase evolution was the discovery of a

RAG1-RAG2-like locus in purple sea urchin genome, which genes for both proteins are oriented in

close proximity in a head-to-head manner as RAG1-RAG2 locus in vertebrates. However this locus

lacks TIR and thus does not show typical features of a transposon [11].

Due to the similarity between RAG1 and *Transib* transposon (a family from the DDE transposon

superfamily) and the fact that RAG2 lacks similarity to any known transposon protein, even though

harbors Kelch-like repeats and PHD domain as other eukaryotic proteins, led several authors to

propose that a *Transib*-like transposon joined the deuterostomian ancestor genome followed by

exon shuffling events bringing *Transib* and the ancestor of RAG2 together [4]. As a result, the RAG1-

RAG2 locus was then recruited for an unknown function. A second, much more recent recruitment

as RAG1-RAG1 V(D)J recombinase most likely occurred at the base of the jawed vertebrate

evolution. Kapitonov and Koonin [12] went a step further and provided *in silico* evidences that RAG1

and RAG2 subunits of the V(D)J recombinase evolved from two proteins encoded in a single

transposon as they found three sequences that could correspond to fossilized RAG1-RAG2

transposon (including TIRs) in one starfish genome. A major step in the understanding of the

RAG1-RAG2 evolution was reported by our group [13] showing for the first time the presence of an

active RAG transposon in the cephalochordate *Branchiostoma belcheri* named ProtoRAG. The full

length ProtoRAG transposon is bound by 5 bp target site duplications (TSDs) and a pair of terminal

inverted repeats (TIRs) resembling V(D)J recombination signal sequences (RSSs). Between the

TIRs reside tail-to-tail oriented, intron-containing and co-transcribed, RAG1-like and RAG2-like

genes. The RAG transposon has been recently active in amphioxus as shown by indel

polymorphisms. Furthermore the amphioxus RAG1-RAG2-like proteins together could mediate

TIR-dependent transposon excision, host DNA recombination, transposition and even signal joint

formation at low frequency, using reaction mechanisms similar to those used by vertebrate RAGs [13].

Here we bring more information about the evolution of RAG transposons. We show that beside *B.*

belchieri, an active RAG transposon is found in the hemichordate *Ptychodera flava,* that several

fossilized transposons are found in several deuterostomes species suggesting that RAG transposon

has been active through the history of the deuterostome lineage.

Results

Description of an active RAG transposon in *P. flava* and many fossilized transposons in

deuterostomes

Due to the discovery of an active RAG transposon in amphioxus *B. belchieri*, we screened all the

available genome and EST projects using the query sea urchin RAG1L and RAG2L sequences.

Many hits in several deuterostomians species were found, hits are found in protosomians but they

show low similarity and correspond to the transib transposons [14] and the chapaev transposon family

[15]. The family reported by Panchin and Moroz as well as many other families were found during our

survey. However the connection between these families and the RAG1-RAG2 is not clear even if

they are related.

Among the hits found in deuterostomes, one of them corresponds to a complete transposon and

other several fossilized transposons (see Supplementary Table 1) in the hemichordate *P. flava*. In

other deuterostome species, we evidenced RAG1L-RAG2L structures without TIRs but with many

fragment copies of the RAG1L-RAG2L locus, some species with an incomplete transposon with

TIR and RAGL sequences and many other copies of RAG1L-RAG2L fragments. The presence of

TIR on many of these copies might indicate that they correspond to fossilized transposons.

Transcribed sequences database are available for several deuterostomes and in most of the case

RAG1L and RAG2L transcripts are found, complete or incomplete, thus revealing the

domestication of the transposon or their activity.

Based on the phylogeny of the RAG1L and RAG2L protein sequences, we can find several RAG

families in *P. flava*. Among them, B and C families have unambiguous TIR and TSD structure. The

phylogenetic analysis is shown in Figure 2 and supplementary Table 1 and described in the

phylogenetic relation between the RAG families session. Two copies of B family show a TSD-

5TIR-RAG1L-RAG2L-3TIR-TSD structure. While one of these copies encodes a complete RAG1L

and RAG2L protein, the other one has suffered different level of pseudogenization. But its presence

confirm beyond doubt that the authentic RAG transposon appears in this family. In the other hand,

the C family has four copies with 5TIR-(RAG1L-RAG2L)-3TIR structure, two of them even

containing the correct TSD in addition of 12 5/3TIR. All this copy seems to be inactivated

(Supplementary Table 1). We failed to find TSD-TIR structure for other RAG-like families (A,

unclassified families) in *P. flava*, this could be due to the poor genome assembly or to the fact that

some families have become inactive. Anyway, this finding is sufficient to prove that multiple

families of RAG transposon have been and are thriving in *P. flava*. Moreover, we found several

fossilized transposons in the case of *Patiria minata* as partially described in 2015 [12], a 5TIR-

RAG1L_fragment-3TIR structure containing TSD and no RAG2L protein, a 5TIR adjacent to

RAG1L structure (TSD-5TIR-RAG1L) and other 12 5/3TIR sequences. These structures indicate

that RAG was an active transposon during the echinoderm evolution. Afterwards a comparative

sequence analysis was made in *B. belcheri*, *Branchiostoma floridae*, *P. flav*a (Pfl) and *P. minata*

(Pmi) TIR sequences (Figure 2) showing no identity between different *Transib*, vertebrate RSS and

amphioxus, Pmi and Pfl species if we exclude the first CAC nucleotides. Nonetheless both

sequences analyzed in amphioxus, shares TIR similarity, suggesting a possibly common origin of

RAG transposon between this two species of amphioxus. However, there is no identity between B

and E RAG transposon families in *P. flav*a, suggesting despite the similarity between RAG-like

proteins of both families, no TIR similarity between each other, as they may be not reactive or

functionally compatible. Previously, an equivalent of RSS nonamer, a stretch of nine highly

conserved nucleotides has been found in the amphioxus ProtoRAG TIR, though this ProtoRAG

nonamer have no similarity with the nonamer found in RSS [13]. However, there are no such nonamer

or equivalently conserved oligomer found in *P. minata* and *P. flava* B and C ProtoRAG family. All

this suggests that the nonamer structure is not important in echinoderms and hemichordates phyla,

but became important in amphioxus and vertebrates.

The species tree in Figure 1 (see also Supplementary Table 1) shows a summary of RAG1L-RAG2L

sequences distribution in deuterostome according to the available data. When genomic and

transcription data are available species name appears in red while when only genomic data are

available the species names are colored in blue and by last only available expressed sequence data

corresponds to the species name colored in black. We predict that the transposon is active if bona

fide sequences are present in the genome in several copies and fragments and if the putative

transposons are transcribed as in the case for *P. flav*a RAGL-B and *B. belcheri*. In the other hand *P.*

miniata seems not to be transcribed since only fossilized transposons are found in the genome. In

two species of sea urchin, *Eucidaris tribuloides* and *Lytechinus variegatus,* no transcribed

sequences are found, but many copies of RAG1L-RAG2L are present on the genome without TIRs

indicating that might be fossilized transposons that became inactivated by the loss of the TIR

sequences.

The case of *S. purpuratus* is more difficult to understand: the published RAG1L-RAG2L locus [11]

renamed here RAG1L-RAG2L B1, was believed to be domesticated as the coding sequence is not

interrupted by stop codons and therefore RAG1L and RAG2L could be functional proteins has no

TIR sequences and both RAG1L and RAG2L are transcribed [11]. However, we found many

fragments highly similar to this sequence in the *S. purpuratus* genome which could reveal a recent

transposition event followed by the domestication of one of its copies (see supplementary data and

figure 2). We find another RAGL copy arose from a duplication event occurred at the origin of the

echinoderms, named RAG1L-B2. The RAG1L-B2 copy is only found fragmented with multiple

recent copies in the genome whereas is complete as RAG1L transcript. A possible explanation for

this second locus could be the existence of an active transposon with the genome sequence not well

assembled or otherwise a domesticated or recent fossilized transposon. For most of the species we

do not have information at the genomic level, but if we find RAGL transcript, this sequence could

correspond to an active transposon, domesticated transposon or recent pseudogene. Anyway, this

shows that the transposon has been present in their ancestors.

Features of the proteins encoded by the RAG-like proteins

In ambulacraria (echinoderm and hemichordate) the deuterostome RAG1-like, 816-1136 aa-long

shares around 26.52% sequence identity between RAG1L-B family and vertebrate RAG1, around

33.21% between the orthologous RAGL-A family and the vertebrate RAG1 and only 27.79%

between RAG1L-A and RAG1L-B, while inside RAG1L-B family are sharing 48.75% of sequence

identity and only 20.13% respect to *Transib* transposase in terms of core region. As regards to

RAG1 lancelet, 30.47% and 37.62% sequence identity are shared with A and B families

respectively and only 27.45% with RAG1 vertebrate (see Supplementary Figure 2A). Clusters of

high identity are found between RAG1L and vertebrate RAG1 along much of their length,

suggesting conservation of multiple functional elements. Vertebrate RAG1 uses four acidic residues

to coordinate critical divalent cations at the active site [16] and all four are conserved in RAG1L

(Supplementary Figure 1A, red highlight). In addition, many cysteine and histidine residues that

coordinate zinc ions and play a critical role in proper folding of RAG1 [17], are conserved between

RAG1L and vertebrate RAG1 (Supplementary Figure 1A, * and # symbols). However, RAG1L

does not share much identity with vertebrates RAG1 in the region corresponding to the nonamer

binding domain, consistent with the fact that RAG transposons TIRs have no clear similarity to the

RSS nonamer. In fact, different families of RAG1-like have little similarity to each other in the

putative nonamer binding domain, consistent with the fact that different ProtoRAG families have

very different TIR sequences and no obvious nonamer regions, excluding the amphioxus TIR.

Finally, there are also some RAG1-like specific conserved regions (see Supplementary Figure 1,

underlined by *). Should be noted that PflRAG1L-A share many sites with jawed vertebrate RAG1.

RAG2L 366-535aa long, shares weak sequence identity between B family and vertebrate RAG2

(18.69%) and between B family and lancelet RAG2L (25.02%). In the other hand RAG2L-B family

shares around 45.90% while lancelet RAG2L shares only 20.24% identity with RAG2 vertebrate

(supplementary Figure 2B). However, the N-terminal six-bladed β-propeller domain (six Kelch-like

repeats), which is conserved in both vertebrate RAG2 and ProtoRAG RAG2L, can be discerned in

RAG2L. Strikingly, amphioxus RAG2L lacks the entire RAG2 C-terminal region, including the

PHD domain as shown previously [13]. However, this PHD domain is present in all other echinoderm

and hemichordate RAG2 proteins (Supplementary Figure 1B). Thus, the absence of this region in

amphioxus RAG transposon might be a secondary loss.

Phylogenetic relation between the RAG families

The phylogenetic analysis with the complete RAG sequences from the available deuterostome

species are shown in Figure 3A and 3B and synthesized in Table 1. At least two sub-families have

been present in the ancestral deuterostome, named RAGL-B and RAGL-A. Other families as

RAGL-C have not been included in the phylogenetic history as they are found only in one species

(Table 1).

In the case of the orthologous relation found between RAG1L-A of *P. flava* (hemichordate) and

vertebrates RAG1 recombinase, we can observe that RAGL-A was lost in many lineages excluding

hemichordate and jawed vertebrates. RAGL-B conversely, is lost in tunicates and in vertebrate

lineage but conserved in several lineages as cephalochordates, hemichordates and echinoderms.

Furthermore the phylogenetic analysis shows that RAGL-B has been duplicated in the echinoderm

ancestor after the hemichordate/echinoderm split and both copies have been kept (even if most of

them have been inactivated) in most of the echinoderm species (Table 1 and Figure 3C).

RAG transposon has been active during the deuterostome evolution

The Figure 4 shows the RAG transposon activity summary during deuterostome evolution. The

transposon has been active in the deuterostome ancestor and in the branch that leads to the common

ancestor of chordate, still active in cephalochordate and domesticated as a RAG1-RAG2 V(D)J

recombinase in the common ancestor of jawed vertebrate. The transposon has been lost in the

Petromyzon lineage (the class Myxini genome is not available and therefore we cannot state for this

important phylogenetic phyla). The transposon has been active in the branch originated from the

node between deuterostome and ambulacraria antecesors. It remains active in hemichordates inside

the subphylum of Enteropneusta (at least on the *P. flava* lineage) but is lost in the other

enteropneusts as *S. kowalevskii.* Unfortunately we do not have genome information for the other

hemichordate subphyla: Pterobranchia. In the case of the echinoderm lineage, the transposon has

been present in the echinoderm common ancestor, in the branch leading to the common ancestor of

crinoid, in the clade formed by the sea urchin and holothuroids and in the clade formed by

starfishes/ophiures. It has been then lost in the crinoid lineage. The transposon has been active in

the branch that goes from the common ancestor of echinoderm to the common ancestor of sea

urchin/Holothuroids and starfishes/brittle stars. Concerning the Asteroidea/Ophiuroidea group, the

transposon has been active in their common ancestor and has been active in the Ophiure lineage in

particular in *O. spicalatus* where the transposon is likely to be active or has lost its activity recently.

The transposon seems to have been inactivated in the starfish lineage but fragments showing

similarities to RAG1-L and/or RAG2-L transposons are found in this species. Furthermore

transposons are clearly found fossilized in *P. miniata*. In the case of sea urchin/holothuroids group,

it seems that the transposon has been active in their common ancestor and inactivated in the

holothurian lineage. By last transposon seems to be active in some sea urchin lineages as in *E.*

tribuloide but much less in others.

Discussion

In this report we show that a RAG transposon has been present in the deuterostome common

ancestors and was active since then in some lineages, fossilized later during evolution and

domesticated at least in the case of jawed vertebrates. The structural and regulatory features that

cause the jawed vertebrate RAG V(D)J recombinase to favor deletional/inversional recombination

over transposition as in the case of the RAG transposase [13] is not yet resolved. It could be explained

by how the cleaved ends and particularly the signals ends are processed. The RAG V(D)J

recombinase binds signal ends tightly as excepted for a transposase but it has acquired the

possibility to give up these end efficiently to the non-homologous end joining machinery. This

allows recombination and prevents the propagation [1]. Thus the jawed vertebrate V(D)J recombinase

differs from the current RAG transposon, as well as its transposon precursor, in how it interfaces

with the DNA repair apparatus. This new property occurred likely in the jawed vertebrate common

ancestor.

DDE transposases have been shown to interact with repair proteins. For example the Sleeping

Beauty transposase interacts directly with the Ku70 repair protein [18] and the pogo transposase of *D.*

melanogaster interacts with the profilirating cell nuclear antigen (PCNA), a key protein for DNA

replication and repair [19]. Therefore, the associations of DDE transposon with DNA repair and

replication factors appear to evolve in a convergent manner [20]. This characteristic and the fact that

the transposon survived during millions of years in multiple copies in different lineages increased

the probability of the co-option of the RAG transposon as V(D)J recombinase. Therefore the

apparition of V(D)J recombination machinery in the jawed vertebrate phyla could be labeled as a

predictable genetic events.

Our results could also explain better the origins of the T-cell receptor and B-cell receptor gene

organization. The earlier proposed scenario [4, 21] involved an insertion of the RAG transposon into

the ancestral IG/TCR V-gene, prior to the externalization of the RAG1-RAG2 complex while

leaving the RSS-like TIR within the IG/TCR V-gene. This was followed by duplication of this new

genetic structure: **V**RSS-RSS**J**. The RAG transposon was then co-opted as V-J recombinase and the

system started to work. However, this scenario explains the V-J structure IG light chain, TCR alpha

and gamma chain but not the VDJ organization of IG heavy chain or TCR beta and delta chains.

This is how we explain this: while one RAG was domesticated (likely RAGL-A orthologue), other

RAG transposons (likely RAGL-B orthologue) were still active as one of them split the **V**RSS-

RSS**J** copy and gave rise to **V**RSS-RSS**D**RSS-RSS**J**. RAG transposase became then extinct and

finally was lost during vertebrate evolution.

Material and methods

Identification of RAG1 and RAG2-like sequence in different data bases

RAG1-RAG2-like locus identified in the echinoderm *Strongylocentrotus purpuratus* and in the

vertebrate genome were used as a protein sequence to perform a TBLASTN-based search against

the NCBI nr protein, transcriptome shotgun assembly (TSA) and the WGS database (as of June

2016) [22]. These retrieved sequences were extracted and translated by ExPASy Translate tool.

Potential open reading frames of RAG1-RAG2 elements used in this study were predicted using

FGENESH [23] with the sea urchin organism specific gene-finding parameters. The mRNA sequences

were then assembled into contigs by CAP3 [24].

Phylogenetic analysis

The alignment and trees were constructed using MEGA6 (complete deletion, WAG with Freqs. (+F)

correction model, 1000 bootstrap replicates [25]). Thus, whether are active, fossilized or domesticated

were classified into families. Short sequence copies, were analyzed one by one with the reference

data set as it is not informative to compare sequences that do not overlap.

Sequence searches for TIR and TSD motifs

We used three methods to search target site duplication (TSD) and terminal invert repeat (TIR)

sequences. In the first method, the upstream and downstream 20 Kb of sequence flanking the

RAG1-RAG2-like sequences were extracted and separated into a set of small fragments (using a

window size of 60 bp and a step size of 1 bp). In the first method, each upstream fragment was

compared with each downstream fragment for 4-6 bp TSDs and possible TIRs using a custom Perl

script. We required 40% identity for potential TIR pairs, and allowed only one mismatch for TSD

pairs. In the second method, all upstream fragments were compared against all downstream

fragments using BLAST. We required a minimum e-value of 100 and sequence identity of 40% in

the BLAST search. However, these two methods failed to work well and provided no reliable

results. Therefore, we turn to the e third method. In this method, we posited that if there are multiple

copies of ProtoRAG transposons in the genome assembly, comparison between these copies could

help to determine their terminal sequences (TIR, etc.).

We focused on finding more complete elements that contain both TIR and RAG gene fragments,

such as "5TIR-RAGs-3TIR", "5TIR-RAGs" and "RAGs-3TIR".

Here is our procedure:

1. First we identified all genomic regions containing RAG1/2 fragments by using TBLASTN

and the amphioxus and vertebrate RAG1/2 proteins as queries;

2. The region containing RAG1/2 plus upstream 20kb and downstream 20kb was extracted,

which we called the RAG region;

3. Because there should a clear border between the ProtoRAG and the host DNA, we can

determine the potential 5' and 3'-terminal of the ProtoRAG transposon by comparing RAG

regions with each other by using BLASTN (see the figure below);

4. Finally, we examine the potential 5/3-terminal sequences of the RAG regions. Most of them

have been destroyed and therefore no detectable TIRs, but there are several of them show

clear and intact TIR structure.

5. And the TSD if presents, should be right next to the TIR sequences.

297
298
Therefore, the sequences containing the RAG1/2-like fragments and the 20 Kb flanking regions

were compared to each other and also to the whole genome assembly using BLAST. The terminal

sequences were analyzed using a custom Perl script and then subjected to manual inspection.

Summary of data availability

In order to detect the absence or presence of a given structure in the genome or transcriptome, we

need to extract all the available taxonomic information from the NCBI database. It has to be noted

that even if the sequence for a given genome is not complete, when RAG1L-RAG2L seems to be an

active transposon, we should find an active or at least a fossilized transposons (in several copies).

Focusing on the genome database we can find species as *Parastichopus parvimensis*, *Acanthaster*

planci, *Ophiothrix spiculata, Petromyzon marinus, Branchiostoma belcheri, Oikopleura dioica,*

Botryllus schlosseri & Ciona savignyi. Transcript sequences can be provided for *Saccoglossus*

kowalevskii, Anneissia japonica, Psathyrometra fragilis, Abyssocucumis albatrossi, Sclerodactyla

briareus, *Apostichopus japonicus*, *Parastichopus californicus*, *Echinarachnius parma*, *Evechinus*

chloroticus, *Paracentrotus lividus*, *Sphaerechinus granularis*, *Arbacia punctulata*, *Henricia sp.* AR-2014, *Echinaster spinulosus*, *Peribolaster folliculatus*, *Leptasterias sp.* AR-2014, *Pisaster ochraceus*, *Marthasterias glacialis*, *Asterias rubens*, *Asterias forbesi*, *Asterias amurensis*, *Luidia clathrata*, *Patiria pectinifera* & *Ophiocoma echinata*. Finally, together with genomic information and transcript expression we have *Ptychodera flava*, *Eucidaris tribuloides*, *Strongylocentrotus purpuratus*, *Lytechinus variegatus*, *Patiria miniata*, *Homo sapiens*, *Mus musculus*, *Gallus gallus*, *Xenopus tropicalis*, *Latimeria chalumnae*, *Danio rerio*, *Carcharhinus leucas*, *Carcharhinus plumbeus*, *Branchiostoma floridae* and *Ciona intestinalis*.

References

1.   Teng G., Schatz D.G., 2015. Regulation and Evolution of the RAG Recombinase. *Adv. Immunol.* **128**, 1-39.

2.   Danchin E. *et al.,* 2004. The major histocompatibility complex origin. *Immunol. Rev.* **198**, 216-232.

3.   Kapitonov V.V., Jurka J., 2005. RAG1 core and V(D)J recombination signal sequences were derived from Transib transposons. *PLoS Biol.* **3**, 998-1011.

4.   Fugmann S.D., 2010. The origins of the Rag genes from transposition to V(D)J recombination. *Semin. Immunol.* **22**, 10-16.

5.   Agrawal A. *et al.,* 1998. Transposition mediated by RAG1 and RAG2 and its implications for the evolution of the immune system. Nature **394**, 744-751.

6.   Hiom K. *et al.,* 1998. DNA transposition by the RAG1 and RAG2 proteins: a possible source of oncogenic translocations. *Cell* **94**, 463-470.

7.   Chatterji M. *et al.,* 2006. Mobilization of RAG-generated signal ends by transposition and insertion in vivo. *Mol. Cell. Biol*. **26**, 1558-1568.

8.   Curry, J.D. *et al.*, 2007. Chromosomal reinsertion of broken RSS ends during T cell

development. J. Exp. Med., 204, 2293-2303.

9.   Ramsden D.A. *et al*., 2010. Weed, B.D., Reddy, Y.V. V(D)J recombination: Born to be wild.

Semin Cancer Biol. **20**, 254-260.

10.   Vanura K. *et al*., 2007. In vivo reinsertion of excised episomes by the V(D)J recombinase: a

potential   threat   to   genomic   stability.   PLoS   Biol.   5,

http://journals.plos.org/plosbiology/article?id=10.1371/journal.pbio.0050043

11.   Fugmann S.D. *et al*., 2006. An ancient evolutionary origin of the Rag1/2 gene locus. *Proc.*

Natl. Acad. Sci. USA **103**, 3728-3733.

12.   Kapitonov V.V., Koonin E.V., 2015. Evolution of the RAG1-RAG2 locus: both proteins

came   from   the   same   transposon.   *Biol.*   *Direct*,

http://biologydirect.biomedcentral.com/articles/10.1186/s13062-015-0055-8.

13.   Huang S. *et al*., 2016. Discovery of an Active RAG Transposon Illuminates the Origins of

V(D)J Recombination. *Cell.* **166**, 102-114.

14.   Panchin Y., Moroz L.L., 2008. Molluscan mobile elements similar to the vertebrate

recombination-activating genes Biochem Biophys Res Commun. 369, 818–823.

15.   Kapitonov V.V., Jurka J., 2007. Chapaev-a novel superfamily of DNA transposons. Repbase

Reports. 7, 777–777.

16.   Ru H. *et al.,* 2015. Molecular Mechanism of V(D)J Recombination from Synaptic RAG1-

RAG2 Complex Structures. *Cell.* **163**, 1138-1152.

17.   Kim M.S. *et al.,* 2015. Crystal structure of the V(D)J recombinase RAG1-RAG2. *Nature*

518, 507-511.

18.   Izsvák Z. *et al*., 2004. Healing the wounds inflicted by sleeping beauty transposition by

double-strand break repair in mammalian somatic cells. *Mol Cell.* **13**, 279-290.

19.   Warbrick E. *et al*., 1998. PCNA binding proteins in Drosophila melanogaster: the analysis

of a conserved PCNA binding domain. *Nucleic Acids Res.* 26, 3925-3932.

20.   Feschotte C., Pritham E.J, 2007. DNA transposons and the evolution of eukaryotic genomes.

Annu. Rev. Genet. **41**, 331-368.

21.   Koonin E.V., Krupovic M., 2015. Evolution of adaptive immunity from transposable

elements combined with innate immune systems. *Nat. Rev. Genet.* **16**, 184-192.

22.   Altschul S.F. *et al*., 1990. Basic local alignment search tool. J. Mol. Biol. 215:403-410.

23.   Solovyev V. *et al*., 2006. Automatic annotation of eukaryotic genes, pseudogenes and

promoters. *Genome Biol.* **7,** S10.1-S10.12.

24.   Huang X., Madan A., 1999. CAP3: A DNA sequence assembly program. *Genome Res.* **9**,

868-877.

25.   Tamura K., *et al*., 2013. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0.

Mol. Biol. Evol. **30**, 2725-2729.

We thank the EBM laboratory for advice and Olivier Loison for editing the manuscript.

Author's contribution: JRMP, PP and SFH conceived the project and design the study. JRMP,

PP and SFH analyzed the results. JRMP, PP, SFH and ALX wrote the manuscript.

The authors declare no competing financial interest

Correspondence and requests for materials should be addressed to **pierre.pontarotti@univ-amu.fr**

and **morales.poole@gmail.com**

Figures and tables

Figure 1 | Distribution of the RAG1-RAG2 sequences in deuterostomes. Only species for which

the genomic and/or transcription data are available are represented in the phylogenetic tree. Species

are colored in red when genomic and transcription data are available, in blue when only genomic

data are available and species are colored in black only when expressed sequence data are available.

```
                        *
ccgCTGCGtgc : 11 : Pfl_B_BCFJ01017854_BCFJ01052781
gccCAATGtgc : 11 : Pfl_B_BCFJ01094280
cacTGTGG--- :  8 : Pfl_B_BCFJ01052780
cacCATCCgta : 11 : Pfl_C_BCFJ01036631
aacCCCGGctg : 11 : Pfl_C_BCFJ01107546
gtcTGCCA--- :  8 : Pfl_C_BCFJ01102604
ctcGGGTG--- :  8 : Pfl_C_BCFJ01084502
ttaCCTGC--- :  8 : Pfl_C_BCFJ01046932
tgcTGCCA--- :  8 : Pfl_C_BCFJ01016857
tcgCAGTG--- :  8 : Pfl_C_BCFJ01107167
gtgCATTG--- :  8 : Pfl_C_BCFJ01047137
tgcGGCCG--- :  8 : Pfl_C_BCFJ01031953
---CGCATtcg :  8 : Pfl_C_BCFJ01070588
---GGGTGcca :  8 : Pfl_C_BCFJ01150129
---GGCCGgtc :  8 : Pfl_C_BCFJ01103012
---CAGTGtgg :  8 : Pfl_C_BCFJ01107168
---CGCGCtcg :  8 : Pfl_C_BCFJ01287958
---CACGGgta :  8 : Pfl_C_BCFJ01048214
---CGCATtcg :  8 : Pfl_C_BCFJ01083599
----------- :  - : blank
cgtCCAGGgtc : 11 : Pmi_JH779599
aacCCATAccg : 11 : Pmi_JH774215
ctcTGTTAtat : 11 : Pmi_JH780459
gagTTTAG--- :  8 : Pmi_JH769343
gatTTTAG--- :  8 : Pmi_JH774292
tgtTCATG--- :  8 : Pmi_JH782081
gcgATGTG--- :  8 : Pmi_JH775549
aagCGCGA--- :  8 : Pmi_JH781149
---CCTGGcat :  8 : Pmi_AKZP01156453
---TTCAGcaa :  8 : Pmi_JH771625
---CATATgca :  8 : Pmi_AKZP01165822
---GTCCGgga :  8 : Pmi_AKZP01162400
```

Figure 2 | Full length alignment of the TSD and TIR sequences. (A) Alignment of RAG

transposon TSDs and flanking sequences from the *P. flava*, *P. miniata*, *B. belcheri*, *B. floridae*

genome. (B) Alignment of *ProtoRAG* TIR sequences with the consensus RSS and *Transib* TIR.

IUPAC codes used in the alignment: N=A, C, G or T; K=G or T; W=A, T; V=A, C or G. Lower case

indicates an undetermined nucleotide. Shading indicates sequence conservation, with darker gray

indicating a higher degree of conservation. Bb: *B. belcheri*; Bf: *B. floridae*. Pfl: *P. flava*, Pmi: *P.*

minata RAG *transposon* copy identification numbers correspond to those listed in Table S1.

Figure 3 | Phylogenetic tree with RAG1 and RAG2 complete sequence and outline of the duplication (D) and lost (⊗). Phylogenetic tree with RAG1 (A) and RAG2 (B) complete sequence. Outline of the duplication and loss of the RAG transposon (C).

Figure 4 | Evolution of the RAG transposon. Transposon activity is indicated in bold pink and

V(D)J recombinase activity is indicated in bold green.

Table 1 | Presence of RAG subfamilies in the different species. Species columns (left to right): *P. flava*, *L. variegatus*, *E. tribuloides*, *P. lividus*, *S. purpuratus*, *E. parma*, *P. pectinifera*, *A. amurensis*, *Leptasterias sp.*, *A. forbesi*, *E. chloroticus*, *P. miniata*, *O. spiculata*, *Henricia sp. AR-2014*, *S. granularis*, *E. spinulosus*, *O. echinata*, *A. japonica*, *P. fragilis*, *A. albanassi*, *S. briareus*, *A. japonicus*, *P. parvimensis*, *P. californicus*, *A. punculata*, *P. folliculatus*, *P. ochraceus*, *M. glacialis*, *A. rubens*, *L. clathrata*, *A. planci*.

Row labels (top to bottom): RAG1B1-like; RAG1B2-like; RAG1B-like (*P. flava* = *); Other families (*P. flava* = **); Not phylogenetically assigned; Not found; RAG2B1-like; RAG2B2-like; RAG2B-like (*P. flava* = ***); Other families (*P. flava* = ****); Not phylogenetically assigned; Not found.

Table 1 | Presence of RAG subfamilies in the different species. Sequences were classified through phylogenetic analysis. Short sequence copies, were analyzed one by one against the reference data described in Figure 3A and 3B as it is not informative to compare sequences that do not overlap. The classification as B family (or A family labeled with "**") is straightforward as it is based on orthologous relationships between different phyla (differences between echinoderms and hemichordates for example). Inside B family, two groups named B1 and B2 are found in several echinoderms. If an echinoderm sequence is classify as B family, but not as B1 or B2 we call it B-like (RAG1Bd-like is labeled with "*" while RAG2Bb-like and RAG2Bc-like are labeled with "***"). We have two specific cases, C family only found in *P. flava* (RAG1 labeled with "**" and RAG2 labeled with "****") and X family in *Ophiotrix spiculata*. The rest of species if they do not belong to A or B family, are not phylogenetically assigned due to the fact that none enough phylogenetic signals are available.

Supplementary figure and table

434
Figure S1 | Features of the proteins encoded by the RAG and RAG-like protein. (A) Protein

alignment of RAG1L with vertebrate RAG1. Repeat motifs in amphioxus and the purple sea urchin

RAG1L were removed and replaced with an "X" and highlighted in yellow. Three regions of

conserved cysteine and histidine residues that might bind zinc are underlined with green bars. The

N-terminal zinc binding dimerization domain is underlined with dark-red bars. The subdomains of

the RAG1 core region are indicated with colored bars. The conserved acidic catalytic residues are

highlighted with red shading (D600, E662, D708 and E962 on mouse RAG1). The PflRAG1LA

show slightly more similar to vertebrate RAG1, and those regions were labeled with "*". GenBank

accessions for mouse RAG1, shark RAG1, lancelet RAG2L and sea urchin RAG1L are NP_033045,

XP_007886047, KJ748699 and NP_001028179, respectively.

(B) Protein alignment of RAG2L with vertebrate RAG2. Color shading shows the conservation of

physiochemical properties. The N-terminal amino acid sequence can be grouped into Kelch-like

repeats. The central conserved GG motifs of the six Kelch-like repeats are underlined in red. The

plant homeodomain (PHD) is also underlined below the alignment. GenBank accessions for mouse

RAG2, shark RAG2, lancelet RAG2L and sea urchin RAG2L are NP_033046, XP_007885835,

KJ748699 and NP_001028184, respectively.

452
```
Percent Identity  Matrix - created by Clustal2.1

 1: transib-1_HM  100.00   18.97   21.85   19.39   19.54   19.60   19.93   18.49   18.40   19.83   21.75
 2: BbeRAG1L        18.97  100.00   39.96   35.03   39.49   37.53   40.20   37.88   30.47   27.55   27.35
 3: PflRAG1LB       21.85   39.96  100.00   35.39   40.37   40.84   41.04   39.00   28.49   26.54   27.30
 4: SpuRAG1LB1      19.39   35.03   35.39  100.00   62.11   43.13   46.87   42.06   27.34   24.74   24.69
 5: EtrRAG1LB1      19.54   39.49   40.37   62.11  100.00   45.70   47.65   43.71   28.79   26.60   26.77
 6: AfoRAG1LB2      19.60   37.53   40.84   43.13   45.70  100.00   56.55   54.70   26.78   26.87   25.86
 7: EclRAG1LB2      19.93   40.20   41.04   46.87   47.65   56.55  100.00   58.77   27.55   28.35   26.75
 8: SpuRAG1LB2      18.49   37.88   39.00   42.06   43.71   54.70   58.77  100.00   26.80   27.03   25.79
 9: PflRAG1LA       18.40   30.47   28.49   27.34   28.79   26.78   27.55   26.80  100.00   33.26   33.15
10: HsaRAG1         19.83   27.55   26.54   24.74   26.60   26.87   28.35   27.03   33.26  100.00   64.41
11: CleRAG1         21.75   27.35   27.30   24.69   26.77   25.86   26.75   25.79   33.15   64.41  100.00
```

453
```
Percent Identity  Matrix - created by Clustal2.1

 1: HsaRAG2     100.00   55.34   19.93   17.10   19.21   16.34   19.74   18.62   16.75   18.54
 2: CplRAG2      55.34  100.00   20.54   17.05   20.21   16.75   20.16   19.05   19.90   21.04
 3: BbeRAG2L     19.93   20.54  100.00   29.50   28.61   21.05   23.91   27.46   20.53   25.07
 4: PflRAG2LBc   17.10   17.05   29.50  100.00   29.80   28.67   29.26   30.77   27.63   27.82
 5: PflRAG2LB    19.21   20.21   28.61   29.80  100.00   29.50   29.61   34.13   33.33   29.48
 6: StrRAG2LB1   16.34   16.75   21.05   28.67   29.50  100.00   64.16   41.29   37.20   37.01
 7: EtrRAG2LB1   19.74   20.16   23.91   29.26   29.61   64.16  100.00   44.74   42.64   39.91
 8: PmiRAG2LB2   18.62   19.05   27.46   30.77   34.13   41.29   44.74  100.00   53.79   49.33
 9: EchRAG2LB2   16.75   19.90   20.53   27.63   33.33   37.20   42.64   53.79  100.00   59.09
10: PliRAG2LB2   18.54   21.04   25.07   27.82   29.48   37.01   39.91   49.33   59.09  100.00
```

Figure S2 | Percent Identity Matrix of RAG1 (S2A) and RAG2 (S2B). In order to provide a

multiple alignment, Clustal-Omega requires a guide tree which defines the order in which

sequences/profiles are aligned. A guide tree in turn is constructed, based on a distance matrix.

Conventionally, this distance matrix is comprised of all the pair-wise distances of the sequences.

The distance measure Clustal-Omega uses for pair-wise distances of un-aligned sequences is the k-

tuple measure. By default, the distance matrix is used internally to construct the guide tree and is

then discarded. By specifying, the internal distance matrix can be written to file.

Table S1 | RAGL distribution in non-chordate genome and expressed sequence. Distribution in

the cephalordate phyla: *B. belcheri* and *B. floridae* available in [13].

464