A Bayesian Perspective on Accumulation in the Magnitude System

Benoît Martin[1], Martin Wiener[2], Virginie van Wassenhove[1*]

[1]CEA, DRF/I2BM, NeuroSpin; INSERM, U992, Cognitive Neuroimaging Unit; Université Paris-Sud; Université Paris-Saclay, F-Gif/Yvette, France.[2]Department of Psychology, George Mason University, Fairfax, VA USA

ABSTRACT

Theoretical and empirical work posits the existence of a common magnitude system in the brain. Such a proposal implies that manipulating stimuli in one magnitude dimension (e.g. time) should interfere with the subjective estimation of another magnitude dimension (e.g. space). Here, we asked whether a generalized Bayesian magnitude estimation system would sample sensory evidence using a common, amodal prior. Two psychophysical experiments separately tested participants on their perception of duration, surface, and numerosity when non-target magnitude dimensions and the rate of sensory evidence accumulation were manipulated. First, we found that duration estimation was resilient to changes in surface or numerosity, whereas lengthening (shortening) duration yielded under- (over-) estimations of surface and numerosity. Second, the perception of numerosity and surface were affected by changes in the rate of sensory evidence accumulation, whereas duration was not. Our results suggest that a generalized magnitude system based on Bayesian computations would minimally necessitate multiple priors.

Keywords: space, time, number, decision-making

INTRODUCTION

INTRODUCTION

The representation of space, time, and number is foundational to the computational brain [1-3],

yet whether magnitudes share a common (conceptual or symbolic) format in the brain is unclear. Walsh's

A Theory Of Magnitude (ATOM) [2] proposes that analog quantities are mapped in a generalized

magnitude system which entails that space, time, and number may share a common neural code. One

additional implication for the hypothesis of a common representational system for magnitudes is that the

estimation of a target magnitude dimension should be affected by the manipulation of another, non-target

magnitude dimension, such that the larger the magnitude of the non-target feature, the larger one should

perceive the target magnitude to be (Figure 1A). Such predictions can be formalized in Bayesian terms

[04] so that each dimension magnitude yields a likelihood estimate subsequently informed by an amodal

prior common to all magnitude dimensions (Figure 1B). In line with ATOM and the common magnitude

system hypothesis, a growing body of behavioral evidence ([5-26], for review see [27-29]) suggests the

existence of interferences across magnitude dimensions. Several neuroimaging studies also suggest the

possibility of a common neural code for quantity estimations mostly implicating parietal cortices (Bueti

and Walsh, 2009; Dormal, et al. 2012; Pinel et al., 2004 ; Hayashi et al., 2015; Hubbard et al., 2005;

Riemer et al., 2016; Nieder, 2013 but see [37-38]). However, while a variety of interactions between time,

space and number has been reported, the directionality of these interactions is not always consistent in the

literature (e.g. [13-14]): for instance, manipulating the duration of events has seldom been reported to

affect numerical and spatial magnitudes [12, 13, 26] and size [5, 7] typically influence duration. Yet, if

time, number and space share a common representational system in the brain and a common amodal prior,

all magnitude dimensions should interact with each other in a bi-directional manner (Figure 1A).

Recent discussions in the field suggest that a common magnitude system would evaluate and

combine quantities in the external world on the basis of Bayesian computations [13, 39]. Convergent with

this proposal, recent examinations of Bayesian processing in magnitude estimation have demonstrated a

number of distinct effects [4]. One primary example is the so-called central tendency effect, wherein

magnitude estimates regress to the mean of the stimulus set, such that large (small) magnitudes are under

(over) estimated. Crucially, central tendency effects have been demonstrated across a number of different

magnitude judgments, including time (historically known as Vierordt's law [39, 40]), number [41],

distance and angle [42]. Further, correlations between the degree of central tendency have been found

between different magnitude dimensions [42], suggesting the existence of "global" priors for magnitude

[4]. The notion of global priors is compatible with Walsh's ATOM model by suggesting that, though

differences in the initial processing of different magnitude dimensions may exist, the representation of

these magnitudes is amodally stored. Further, the existence of global priors would provide an explanation for congruency effects between different magnitude dimensions. However, a single global prior for magnitude estimations would not explain why congruency effects may be inconsistent between dimensions or why the directionality of interferences may differ across magnitude dimensions.

To address this first working hypothesis, we used a paradigm in which stimuli consisted of clouds of dynamic dots characterized by the total *duration* of the trial (D: time magnitude), the total *number* of dots (N: number magnitude) and the overall *surface* filled by the dots (S: space magnitude). Two experiments were conducted (Figure 2A-B). In a first experiment (Experiment 1), while participants estimated a target magnitude dimension (e.g. D), we independently manipulated the non-target magnitude dimensions (e.g. S and N). This design allowed us to test all possible combinations and investigate possible interactions between magnitudes (Table 1). If magnitude dimensions interact, increasing or decreasing the N or the S should lead to an under/overestimation of D (Figure 2D). These effects should be bidirectional so that when participants estimate N or S, increasing or decreasing the non-target dimensions D should lead to under or overestimation of the target magnitude dimension N or S. On the other hand, if dimensions are independent, manipulating the number of events in a given trial should not affect duration or surface estimates. Similarly, decreasing or increasing the duration should not affect numerical or spatial judgments if magnitudes are independent.

In a second working hypothesis, we manipulated the accumulation regime of sensory evidence for N and S estimation (Figure 2B). The accumulation of sensory evidence in time for space and number has seldom been controlled for or manipulated during magnitude estimations. In a prior experiment [13], constraining the duration of sensory evidence accumulation in the S and N dimensions, the estimation of duration remained resilient to changes in the other dimensions, whereas D affected the estimation of S and N: curiously, the longer (shorter) durations decreased (increased) the estimation of S and N. These results were discussed in the context of a possible Bayesian integration of magnitude dimensions. Similarly, here, using a dynamic paradigm in which N and S accumulate over time raises the question of the implications of varying the speed or rate of sensory evidence delivery: for a given N or S, if D increases, the speed of presentation decreases, and vice versa. Hence, while the number of dots and the cumulative surface accumulated linearly in time in the Experiment 1 (Figure 1B), in the second experiment (Experiment 2) we investigated whether changes in the rate of presentation of visual information affected the estimation of D, N, and S. Two evidence accumulation regimes were tested: a fast-slow (FastSlow) and a slow-fast distribution (SlowFast), see Stimuli part in Material & Methods section.

In a third question, we wished to investigate the extent to which Bayesian models could explain

the behavioral results obtained in magnitude estimation, independent of the means by which participants

provided their estimates. Thus far, studies demonstrating central tendency effects [39, 42, 43] have all

relied on continuous estimation procedures, wherein participants estimated a particular magnitude value

with a motor response. In particular, these studies utilized reproduction tasks, which require participants to

demarcate where (when) a particular magnitude matched a previously presented standard. In contrast, the

majority of studies demonstrating congruency effects in magnitude estimation have all employed two-

alternative forced choice (2AFC) designs. This difference may be particularly relevant as recent studies

have demonstrated that the size-time congruency effect, one of the most heavily studied and replicated,

depends on the type of decision being made ([44] but see [25,45] for congruency effects with temporal

reproduction). As such, for both Experiment 1 and 2, we provide systematic quantifications of the

magnitude estimates as categorical estimations together with analysis of continuous reports.

MATERIALS & METHODS

Participants

A total of 45 participants were tested. 3 participants did not come to the second session and 10 were disregarded for poor performance. Hence, 17 healthy volunteers (7 males, mean age 24.9 ± 5.8 y.o.) took part in Experiment 1, and 15 participants (8 males, mean age 26.5 ± 7 y.o.) took part in Experiment 2. All had normal or corrected-to-normal vision. Both experiments took place in two sessions one week apart. Prior to the experiment, participants gave a written informed consent. The study was conducted in agreement with the Declaration of Helsinki (2008) and was approved by the Ethics Committee on Human Research at Neurospin (Gif-sur-Yvette, France). Participants were compensated for their participation.

Stimuli

The experiment was coded using Matlab 8.4 with Psychtoolbox (v 3.0.12) and built on a published experimental design (see [13]). Visual stimuli were clouds of grey dots which appeared dynamically on a black computer screen (1024 x 768 pixels, 85 Hz refresh rate). Dots appeared within a virtual disk of diameter 12.3 to 15.2 degrees of visual angle; no dots could appear around the central fixation protected by an invisible inner disk of 3.3 degrees. Two dots could not overlap, the duration of each dot varied between 35 ms to 294 ms, and the diameter between 0.35 to 1.14 degrees. A cloud of dots was characterized by its duration (D: total duration of the trial during which dots were presented), its numerosity (N: cumulative number of dots presented on the screen in a given trial) and its surface (S: cumulative surface covered by the dots during the entire trial). On any given trial, D, N and S could each take 6 possible values corresponding to 75, 90, 95, 105, 110 and 125 % of the mean value. We fixed D to 800 ms ($D_{mean} = 800$ ms) and initially picked $N_{mean} = 30$ dots and $S_{mean} = 432$ mm² which were individually calibrated in the first session of the experiment (see Procedure). To ensure that luminance was not used as a cue to perform the task, the relative luminance of dots varied randomly across all durations among 57, 64, 73, 85, 102 and 128 in the RGB-code. In Experiment 1, the total number of dots accumulated linearly over time, 2 to 7 dots at a time in steps of 9 to 13 iterations (Figure 2A). In Experiment 2, the total number of dots accumulated in a fast-to-slow or in a slow-to-fast progression (FastSlow or SlowFast, respectively): in FastSlow, $75\% \pm 10\%$ of the total number of dots in the trial were presented in the first 25% of the duration of the trial, whereas in SlowFast, $25\% \pm 10\%$ of the total number of dots was shown in 75% of the total duration of the trial.

Procedure

Participants were seated in a quiet room ~60 cm away from the computer screen with their head maintained on a chinrest. The main task consisted in estimating the magnitude of the trial along one of its three possible dimensions (D, N, or S). Each experiment consisted of two sessions: in the first session, stimuli were calibrated to elicit an identical discrimination threshold in all three dimensions on a per individual basis (see [13]); the second session consisted in the experiment proper.

In the first session of Experiment 1 and 2, the task difficulty across magnitudes was individually calibrated by computing the participant's Point of Subjective Equality (PSE: 50% discrimination threshold) and the Weber Ratio (WR) for each dimension D, N, and S. Specifically, participants were passively presented with exemplars of the minimum and maximum value for each dimension and were then required to classify 10 of these extremes as minimum '-' or maximum '+' by pressing 'h' or 'j' on an AZERTY keyboard. Participants then received feedback indicating the actual number of good answers that they provided. Subsequently, the PSE and the WR were independently assessed for each magnitude by varying one dimension and keeping the other two dimensions at their mean values (e.g. if D varied among its 6 possible values, S was S_{mean} and N was N_{mean}). 5 trials per value of the varying dimension were collected yielding a total of 30 trials per dimension from which the individual's PSE and WR were computed and compared. This process (~15 min) was iterated until the PSE and the WR were similar and stabilized across magnitudes. The final mean values (mean +/- SD) for Experiment 1 were $N_{mean} = 32$ +/- 3 dots and $S_{mean} = 476$ +/- 58 mm². The final mean values for Experiment 2, $N_{mean} = 32$ +/- 2 dots and $S_{mean} = 490$ +/- 41 mm².

In the second session, and for both Experiment 1 and 2, participants first performed 30 trials/magnitude dimension similar to the first session to ensure that their PSE and WR remained identical. Only two participants in Experiment 1 required a recalibration procedure.

Subsequently, participants performed the magnitude estimation task *proper* in which participants were asked to provide a continuous estimation of a given magnitude by moving a cursor on a vertical axis whose extremes were the minimal and maximal values of the dimension. In a given trial, participants were provided with the written word 'Durée' (Duration), 'Nombre' (Number) or 'Surface' (Surface) which indicated to participants which dimension they had to estimate (Figure 2A). At the end of a trial, the vertical axis appeared on the screen with the relative position of '+' and '-' pseudo-randomly assigned. The cursor was always initially set in the middle position on the axis. Participants used the mouse to vertically move the slider along the axis and made a click to validate their response. They were asked to emphasize accuracy over speed. Trials were pseudo-randomized across dimensions and conditions.

In Experiment 1, five experimental conditions were tested per dimension: in the control condition, the two

non-target dimensions were kept to their mean values, and in the four remaining conditions, one of the

other non-target dimension was minimal or maximal while the other was kept to its mean (Table 1). A

total of 1080 trials were tested (3 dimensions x 5 conditions x 6 values x 12 repetitions).

In Experiment 2, there were two main sensory accumulation regimes (FastSlow, and SlowFast) and the

emphasis was on the effect of duration on surface and number. Hence, the main control condition

consisted in assessing the estimation of duration with S_{mean} and N_{mean} and whether the rate of evidence

delivery affected duration estimates. Similarly for N and S, two control conditions investigated the effect

of the rate of stimulus presentation, without varying non-target dimensions. Due to results obtained in

Experiment 1, Experiment 2 neither investigated interactions of N or S on D, nor interactions between N

and S. Ten experimental blocks alternated between FastSlow and SlowFast presentations counterbalanced

across participants. 12 repetitions of each possible combination were tested yielding a total of 144 trials

for D (2 distributions x 6 durations x 12 repetitions), 432 trials for N (2 distributions x 3 conditions x 6

numerosities x 12 repetitions) and 432 trials for S (2 distributions x 3 conditions x 6 surfaces x 12

repetitions) for a grand total of 1008 trials. (Table 2).

Analyses

To analyze the point-of-subjective equality (PSE) and the Weber Ratio (WR), participants' continuous

estimates were first transformed into categorical values: a click between the middle of the axis and the + (-

) was considered as a + (-) response. Proportions of + were computed on a per individual basis and

separately for each dimension and each experimental condition. Proportions of + responses were fitted

using the logit function (Matlab 8.4) on a per individual basis. Goodness-of-fits were individually assessed

and participants for whom the associated p-values in the control conditions were >.05 were excluded from

the analysis. Per condition, PSE and WR that were 2 standard deviations away from the mean were

disregarded and replaced by the mean of the group (max of 2 values / condition across all individuals).

Statistics were run using R (Version 3.2.2). PSE and WR were defined as:

$$\text{PSE} = f^{-1}(0.5)$$

$$\text{WR} = 0.5 \times (f^{-1}(0.75) - f^{-1}(0.25)) \times \frac{1}{\text{PSE}}$$

Additionally, continuous estimates were analyzed to interrogate central tendency effects. For each

magnitude dimension, continuous estimates were expressed as the relative position on the slider that

participants selected on each given trial, with higher percentages indicating closer proximity to '+'. To
measure the central tendency effects, continuous estimates were plotted against the corresponding
magnitude for each condition, also expressed as a percentage – where 0 indicated the smallest magnitude
and 100 indicated the largest – and fits with a linear regression, and the slope and y-intercept of the best
fitting line were extracted [46, 47]. Slope values closer to 1 indicated veridical responding (participants
responded with perfect accuracy), whereas values closer to 0 indicated a complete regression to the mean
(participants provided the same estimate for every magnitude). In contrast, intercept values of these
regressions could indicate an overall bias for over- or under-estimation. To compare central tendency
effects between magnitude dimensions, correlation matrices between slope values for D, N, and S were
constructed. Bonferroni corrections were applied to control for multiple comparisons.

RESULTS

To examine Bayesian effects in the magnitude system, we evaluated both choice and continuous
judgments in two magnitude estimation experiments using variations of the same paradigm (Figure 2A).
To evaluate choice responses, continuous estimates were binned according to which end of the scale they
were closer to. Previous work has demonstrated that bisection tasks and continuous estimations are
compatible and provide similar estimates of duration [43, 48]. Our intention was thus to first replicate the
effects of Lambrechts and colleagues [13] with a modified design, and second, to measure central
tendency effects in our sample to examine whether these effects correlated between magnitude
dimensions, which would suggest the existence of global priors ([4], Figure 2D).

Control conditions: matching task difficulty across magnitude dimensions

Two independent repeated-measures ANOVAs with the PSE or WR as dependent variables using
magnitude dimensions (3: D, N, S) and control conditions (3: Linear (Experiment 1), SlowFast and
FastSlow (Experiment 2) distributions) as within-subject factors did not reveal any significant differences
(all $p > .05$). This suggested that participants' ability to discriminate the different values presented in the
tested magnitudes was well matched across magnitude dimensions (Figure 2C).

Experiment 1: Duration affects Number and Surface estimates

We first analyzed the data of Experiment 1 as categorical choices. Figure 3A illustrates the grand average

estimations of duration, numerosity, and surface for all experimental manipulations (colored traces) along

with changes of PSE (insets).

Separate 2x2 repeated-measures ANOVAs were run on PSE for each target magnitude dimension using

the non-target magnitude dimensions (2) and their magnitude values (2: min, max) as within-subject

factors. No main effects of non-target magnitude dimension ($F[1,16] = 0.078$, $p = 0.780$), magnitude value

($F[1,16] =0.025$, $p =0.875$) or their interaction ($F[1,16] = 0.003$, $p =0.957$) were found on duration (D)

indicating that manipulating N or S did not change participants' estimation of duration (Fig. 3A, left

panel). In the estimation of N, main effects of non-target dimensions ($F[1,16] = 7.931$, $p = 0.0124$), their

magnitude value ($F[1,16]=25.53$, $p =0.000118$) and their interaction ($F[1,16] = 23.38$, $p =0.000183$) were

found. Specifically, when the non-target dimensions were at their minimal value, the PSE obtained in the

estimation of N was lower than when the non-target dimensions were at their maximal value.

Additionally, in N estimation, D_{min} lowered the PSE more than S_{min}, and D_{max} raised the PSE more than

S_{max}. Paired t-tests were run contrasting the PSE obtained in the estimation of N during the control (D_{mean}

S_{mean}) and other experimental conditions: D_{min} significantly increased [$PSE(D_{min}) < PSE(D_{mean})$: $p = 4.1e^{-5}$]

whereas D_{max} significantly decreased [$PSE(D_{max}) > PSE(D_{mean})$: $p = 0.0032$] the perceived number of dots

(Fig. 3A, middle panel, inset). There were no significant effects of S on the estimation of N. Altogether,

these results suggest that the main effect of non-target dimension on numerosity estimation was driven by

the duration of the stimuli.

In the estimation of S, we found no main effect of non-target magnitude dimension ($F[1,16] = 1.571$, $p =

0.228$) but a significant main effect of magnitude value ($F[1,16]=22.63$, $p =0.000215$). The interaction

was on the edge of significance ($F[1,16] = 3.773$, $p =0.0699$) suggesting that, as for N, only one non-target

magnitude dimension may be the main driver of the significant results observed in the magnitude effect.

Paired t-tests contrasting the PSE obtained in the estimation of S during the control (D_{mean} N_{mean}) and other

conditions showed that D_{min} significantly increased ($PSE(D_{min}) < PSE(D_{mean})$: $p = 8.7e^{-4}$), whereas D_{max}

significantly decreased ($PSE(D_{max}) > PSE(D_{mean})$: $p = 0.035$) the perceived surface (Fig. 3A, right panel).

No significant effects of N on S were found. As observed for the estimation of N, these results suggest that

the main effect of non-target magnitude dimension on the estimation of S was entirely driven by the

duration.

Overall, the analysis of PSE indicated that participants significantly overestimated N and S when dots

were presented over the shortest duration, and underestimated N and S when dots accumulated over the

longest duration. Additionally, manipulating N or S did not significantly alter the estimation of duration.

No significant interactions between N and S were found. To ensure that these results could not be

accounted for by changes in participants' perceptual discriminability in the course of the experiment,

repeated-measures ANOVA were conducted independently for each target dimension (3: D, N, S) with the

Weber Ratio (WR) as dependent variable and experimental conditions (5) as main within-subject factors.

No significant differences (all $p > .05$) were found suggesting that the WRs were stable over time, and that

task difficulty remained well matched across dimensions in the course of the experiment.

For the analysis of continuous estimates (Figure 3B), we first examined the effect of central tendency for

each target magnitude dimension, collapsing across the non-target magnitudes (Figure 4A). A repeated

measures ANOVA of slope values with magnitude as a within-subjects factor revealed a main effect of the

target magnitude [$F(2,32) = 13.284$, $p = 0.000063$]. Post-hoc paired t-tests identified this effect as driven

by a lower slope, indicating a greater regression to the mean for S as compared to D [$t(16) = 3.495$, $p =$

0.003] and N [$t(16) = 5.773$, $p = 0.000029$], with no differences in slope values between D and N [$t(16) =$

0.133, $p = 0.896$] (Figure 4B).

Further analyses revealed comparable findings as the categorical analysis: separate 2x3 repeated measures

ANOVAs were run for each magnitude dimension, with the non-target magnitude dimension and its

magnitude value as within-subject factors. Analysis of slope values revealed no significant main effects

or interactions for any of the tested magnitudes (all $p > .05$), indicating no change in central tendency, as a

function of the non-target magnitudes. However, an analysis of intercept values demonstrated a

significant main effect of the non-target dimension for S [$F(2,32) = 24.571$, $p < 0.00001$] and N [$F(2,32) =$

39.901, $p < 0.00001$], but not for D [$F(2,32) = 0.010$, $p = 0.99$]. Specifically, intercept values were shifted

higher (lower) when D_{max} (D_{min}) was the non-target dimension in both the S and N tasks (Figure 3B, red

hues), but not when the non-target dimension was N for S, S for N, or for either S or N when D was the

target magnitude dimension.

To examine the central tendency effects across magnitude dimensions, we correlated the slope values

between target magnitude dimensions (Figure 4C). Collapsing across the non-target dimensions, we

found that all three slope values significantly correlated with one another [D to S: Pearson $r = 0.594$; D to

N: $r = 0.896$; S to N: $r = 0.662$]. Given that S exhibited a greater central tendency than D or N, we

compared the Pearson correlation coefficients with Fisher's z-test for the differences of correlations; this

analysis revealed that the D to N correlation was significantly higher than the D to S correlation [$Z = 2.03$,

$p = 0.04$], and marginally higher than the S to N correlation [$Z = 1.73$, $p = 0.083$], suggesting that D and N

dimensions, which had similar slope values, were also more strongly correlated with each other than with

S. To further explore this possibility, we conducted partial Pearson correlations of slope values; here, the

only correlation to remain significant was D to N, when controlling for S [$r = 0.8352$], whereas D to S,

controlling for N, and N to S, when controlling for D were no longer significant [$r = 0.0018$ and 0.3627,

respectively].

The results of the correlation analysis revealed that D and N tasks were highly correlated in slope,

indicating that individual subjects exhibited a similar degree of central tendency for these two magnitude

dimensions (Figure 4A). To explore this at a more granular level, we expanded our correlation analysis to

include all non-target dimensions (Figure 4C). The result of this analysis, with a conservative Bonferroni

correction ($r > 0.8$) for multiple comparisons confirmed the above results, demonstrating that D and N

dimensions were correlated across most non-target dimensions, but that D and N dimensions were weakly

and not significantly correlated with S. This finding suggests that D and N estimation may rely on a

shared prior, that is separate from S; however, a shared (D, N) prior would not explain why D estimates

were unaffected by changes in N, nor would it explain why S estimates are affected by changes in D.

Lastly, we sought to compare the quantifications based on continuous data with those from the categorical

analysis. Previous work has demonstrated that the WR on a temporal bisection task correlates with the

central tendency effect from temporal reproduction [43]. To confirm this, we measured the correlations

between the slope values of continuous magnitude estimates with the WR from the categorical analysis.

As predicted, we found a significant negative correlation between slope and WR for D ($r = -0.69$) and N (r

$= -0.57$); however, the correlation for S failed to reach significance ($r = -0.41$, $p = 0.1$), indicating that

greater central tendency (lower slope values) were associated with increased variability (larger WR). This

finding is notable, as the analysis of WR values did not reveal any difference between magnitude

dimensions. This suggests that the slope of continuous estimate judgments may be a better measure of

perceptual uncertainty than the coefficient of variation derived from categorical responses.

Experiment 2: Duration is robust to accumulation rate, not N and S

In Experiment 2, participants estimated D, N or S while the accumulation regime was manipulated as

either FastSlow or SlowFast (Fig. 2B, Table 2). As previously, we systematically analyzed the categorical

and the continuous reports. First, we tested the effect of the accumulation regime on the estimation of each

magnitude dimension by using a 2x3 repeated-measures ANOVA with PSE measured in control

conditions (Table 2, 1st row) as independent variable and distribution (2: FastSlow, SlowFast) and

magnitude dimension (3: N, D, S) as within-subject factors. Marginal main effects of accumulation regime

($F[1,28] = 2.872$, $p = 0.0734$) and magnitude dimensions ($F[1,14] = 4.574$, $p = 0.0506$) were observed.

Their interaction was significant (F[2,28] = 10.54, p = 0.0004). A post-hoc t-test revealed no significant

effects of accumulation regime on the estimation of D (p = 0.23), but significant effects of accumulation

regime in the estimation of N (p =0.016) and S (p = 0.0045) (Fig. 5A).

Second, we tested the effect of D and accumulation regime on the estimation of N and S (Figure 5A, top

insets). We conducted a 2x2x2 repeated-measures ANOVA with PSE as an independent variable and

magnitude dimension (2: N, S), accumulation regime (2: FastSlow, SlowFast), and duration (3: D_{min}, D_{max})

as within-subject factors. Main effects of accumulation regime (F[1,14] = 22.12, p = 0.000339) and

duration (F[1,14] = 27.65, p = 0.000121) were found, suggesting that both N and S were affected by the

distribution of sensory evidence over time, and by the duration of the sensory evidence accumulation. No

other main effects or interactions were significant although two interactions trended towards significance,

namely the two-way interaction between accumulation regime and duration (F[1,14] = 3.482, p = 0.0831)

and the three-way interaction between dimension, accumulation regime, and duration (F[1,14] = 3.66, p =

0.0764). These trends were likely driven by the SlowFast condition as can be seen in Figure 5A.

For the analysis of continuous data (Figure 5B), we first examined any overall differences in slope values

for different accumulation regimes (FastSlow *vs.* SlowFast) across all three target magnitude dimensions

(D, N, S). A (3x2) repeated measures ANOVA with the above as within-subjects factors revealed a main

effect of magnitude [$F(2,32)$ = 7.878, p = 0.002], with S once again demonstrating the largest slope value,

but no effect of accumulation regime or interaction (both p >.05), suggesting that the rate of accumulation

did not influence the central tendency effect. However, on the basis of our *a priori* hypothesis, post-hoc

tests revealed a significantly lower slope value for N in SlowFast compared to FastSlow [$t(17)$ = 3.067, p

= 0.007], suggesting that participants exhibited more central tendency for numerosity when the

accumulation rate was slow in the first half of the trial (Figure 5B). The analysis of intercept values did

not reveal any effects of accumulation regime or magnitude (all p >.05). However, on the basis of our *a*

priori hypothesis, post-hoc tests demonstrated that S exhibited a significantly lower intercept for SlowFast

compared to FastSlow [$t(17)$ = 3.609, p = 0.002], with no changes for either D or N (both p >.05),

indicating that participants underestimated surface when the rate of evidence accumulated slowly in the

first half of the trial.

For S and N, further examination of slope values for the three possible durations using a 2x2x3 repeated

measures ANOVA with magnitude (S, N), accumulation regime (FastSlow,SlowFast), and duration (D_{min},

D_{mean}, D_{max}) as within-subjects factors, revealed a significant main effect of magnitude [$F(2,32)$ = 7.717, p

= 0.013] and of accumulation regime [$F(2,32)$ = 11.345, p = 0.004], but not of duration [$F(2,32)$ = 1.403,

p = 0.261]. Using the same analysis for intercept values, we found no main effects of magnitude [$F(2,32)$

= 1.296, p = 0.272], but a significant effect of accumulation regime [$F(2,32)$ = 5.540, p = 0.032] and of

duration [$F(2,32) = 21.103$, $p = 0.000001$]. More specifically, we found that intercept values were lower for longer durations, indicating greater underestimation when the interval tested was longer. No other effects reached significance (all $p > .05$).

Overall, these findings indicate that duration estimations were immune to changes in the rate of accumulation of non-target magnitudes, similar to the findings of Experiment 1. Also similar, we found that estimates of S and N were affected by duration as non-target magnitude, with longer durations associated with greater underestimation of S and N (Figure 6). In addition, our results demonstrate a difference between accumulation regimes for S and N, with SlowFast regimes associated with greater underestimation than FastSlow, regardless of duration. Lastly, we observed that SlowFast accumulation regimes led to an increase in the central tendency effect, suggesting that slower rates of accumulation may increase reliance on the magnitude priors.

Discussion

In this study, we report that when sensory evidence accumulates equally over time and when task difficulty is equated across magnitude dimensions (space, time, number), duration estimates are resilient to manipulations of number and surface, whereas number and surface estimates are biased by the temporal properties of sensory evidence accumulation. Specifically, we replicated the findings of Lambrechts and colleagues [13] by demonstrating that number and surface estimates are under- and over-estimated when presented for long and short durations, respectively. As we did not find robust bidirectional interactions between dimensions, these findings do not support the idea of a common magnitude system in the brain. However, we do not argue that time, number and space do not interact under certain constraints. Specifically, by considering a Bayesian model relying on multiple priors (one for each dimension), magnitudes may interact when providing conflicting sensory cues. Recent hypotheses suggest that a Bayesian framework can provide a general explanation for the variety of behavioral features observed in magnitude estimations independently applied to distance, loudness, numerical or temporal judgments [4]. The proposed Bayesian framework combines an estimate of the likelihood (sensory input) with a prior representation (memory). One major goal of our study was thus to determine the degree to which different magnitude dimensions might rely on an amodal global prior representation of magnitude as

would be expected in a generalized magnitude system such as ATOM [2]. To accomplish this,
participants took part in two experiments independently manipulating the congruence across magnitude
dimensions (Experiment 1) and the rate of sensory evidence provided to participants (Experiment 2).

A first prediction was that if different magnitude dimensions rely on a single amodal prior, then
magnitude estimates should exhibit similar levels of central tendency across magnitude dimensions
(duration, surface, number; Figure 2D). Instead, in Experiment 1, our results demonstrated that surface
estimates exhibited greater central tendency than either duration or number, and further was not correlated
with the degree of central tendency for either dimension. However, duration and number did exhibit
correlated central tendency effects. This finding suggests that estimates of surface are distinct from
estimates of duration and number, but that duration and number may be more similar to one another.
Indeed, neural recording studies in the prefrontal and parietal cortex of non-human primates have revealed
overlapping, yet largely separate, representations of duration and size [49, 50], and number and size [51,
52], respectively. Further, while number, size, and time exhibit common activations of the right parietal
cortex, they each engage larger networks of regions beyond this area [37, 53, 54]. For size estimates,
recent work suggests that comparisons of object size draw on expectations from prior experience in other
brain regions [55]. Yet, as no strong bidirectional effects were observed between duration and number, it
is unlikely that duration and number share neuronal populations with similar tuning features.

Another possible interpretation of the results obtained in Experiment 1 is to consider multiple
priors in magnitude estimations. When participants make temporal judgements, the combination of prior
knowledge $P(\pi)$ and noisy sensory inputs $P(D|\pi)$ (duration of the given trial) enables participants to make
an accurate posterior estimate, represented by $P(\pi|D) \propto P(D|\pi) \bullet P(\pi)$ (see [4], Box 3 for more details),
which explains the regression to the mean. Neither numerosity nor surface priors are present in this
equation, which could explain why duration estimates are robust to numerosity or surface manipulations.
Because numerosity and surface accumulate over time, one possible strategy for the participants is to
estimate numerosity and surface based on the speed of presentation of stimuli, and on the duration of the
trial (i.e. a high (low) speed and a long (short) duration of presentation correspond to a large (small) value
of numerosity or surface). Under this hypothesis, the uncertainty related to the temporal dimension may
add noise in the decision or the accumulation process, so that the perceived duration of the trial can bias
numerosity and surface estimates (Figure 6). When numerosity and surface accumulate over a given
duration, if that duration is short (long) it will affect the accuracy of participants' estimations. Because the
short (long) duration was overestimated (underestimated) it may explain why participants overestimated
(underestimated) numerosity and surface in the D_{min} (D_{max}) condition. This explanation would be
compatible with the hypothesized effect of duration as introducing noise on sensory accumulation.

Indeed, one noteworthy aspect unique to the time dimension is that the objective rate of
presentation is fixed [56]. That is, objective time by conventional measurements proceeds at a single mean
rate. In contrast, we can experimentally manipulate the rate at which we present information for number
and surface. In Experiment 1, in order to keep the values of surface and number fixed when duration was
manipulated, we necessarily had to change the rate of accumulation for these values. For example,
between short and long durations with the same value of number, we had to change the rate of
accumulation for number so that the same total value was reached at the end of the duration. This may
explain the incongruent effects of duration on surface and number; shorter (longer) durations may
engender larger (smaller) estimates of surface and number because the rate of accumulation is faster
(slower). In this sense, surface and number are not being influenced by the magnitude of time as a
dimension *per se*, but rather time is interfering with the rate of accumulation, and so the effect of duration
is an epiphenomenon of the experimental design. Hence, to test this hypothesis, we modulated the
accumulation rate of the presentation of numerosity and surface in Experiment 2.

In Experiment 2, where the rate of accumulation for number and surface experienced a rate-
change a little less (more) than halfway through the presentation time from fast-to-slow (slow-to-fast), we
replicated and extended our findings of Experiment 1. Specifically, we again found that shorter (longer)
durations led to longer (shorter) estimates of surface and number, regardless of the rate-change in
accumulation regime. However, we also found a difference in accumulation regimes: when the rate of
accumulation was slower in the beginning of the session, estimates of surface and number were smaller
than when the rate of accumulation was faster. It is important to remember that the ultimate value of the
presented surface and number was the same, regardless of the accumulation regime. As such, participants
were biased in their estimates by the rate of evidence accumulation in the first-half of the given trial,
regardless of how long that trial lasted. This strongly suggests that human observers are biased by the rate
of accumulation at the start of a trial, and are resistant to changes in rate throughout the trial.

This observation is important in the context of ongoing discussions on drift-diffusion processes in
which the accumulation of evidence following the first end-point depends on terminated processes and
guess probability ([57] case study 1 and Figure 1;[58] Figure 2, 4A) and on the importance of change
points during the accumulation process [59,60]. Additionally, this finding strengthens the hypothesis that
the effect of duration on surface and numerosity may occur as a result of the impact on the implicit timing
or accumulation rate, and not as a function of the explicitly perceived duration. This would be consistent
with recent findings suggesting that noise memory - known to scale with duration - was not the primary
factor of errors in decision-making but that noise in sensory evidence was instead a major contributor [61].

Our results suggest that speeding up the rate of evidence and lengthening the duration of a trial may be

equivalent to increasing noise in sensory accumulation of other magnitude dimensions (Figure 6).

In Experiment 2, we also investigated the effect of accumulation regime on central tendency as

participants again provided continuous magnitude estimates on a vertical sliding scale. Previous

magnitude studies using continuous estimates have demonstrated a central tendency effect, with

over(under)-estimations for small (large) magnitudes [4], the degree to which depends on the uncertainty

inherent in judging the magnitude in question [43, 62]. The result of this analysis revealed that, when the

rate of accumulation was slow (fast) in the beginning of the trial, the degree of central tendency was

greater (lesser); further, the objective duration of the trial did *not* impact central tendency. This finding

suggests that slower accumulation regimes engender greater uncertainty in magnitude estimates, and that

this uncertainty may be present before the ultimate decisional value is reached. Previous work in

decision-making with evidence accumulation has suggested that the objective duration of a trial leads to

greater reliance on prior estimates, as longer presentation times are associated with greater uncertainty

[63]. Specifically, Hanks and colleagues [63] found that a drift-diffusion model that incorporated a bias

signal to rely on prior evidence that grows throughout the trial could explain reaction time differences in a

dots-motion discrimination task. Notably, the bias signal is incorporated into the drift-diffusion process,

such that longer trials push the accumulation rate towards a particular value, depending on the prior. A

critical manipulation in this study was the emphasis on speed or accuracy for subjects; increased emphasis

for accuracy led to longer decision times and greater reliance on the prior, as explained by the model. Our

results suggest otherwise – the duration of the trial alone cannot determine reliance on the prior. If the

effect of duration solely led to greater reliance on the prior, then we should have seen central tendency

effects increase with longer durations, which did not occur in either Experiments 1 or 2. Instead, the rate

of evidence accumulation determined reliance on the prior(s), regardless of duration, with slower rates

leading to greater reliance.

Additional studies using neuroimaging techniques such as M/EEG need to investigate the neural

correlates underpinning accumulation processes in the brain when estimating magnitudes (Centro-Parietal

Positivity, for example, see [64,65]), to fully explain the behavioral results obtained in these two

experiments. Further, fMRI studies must be conducted to elucidate the neural circuits for memory

representations of different magnitudes [47]. Bayesian approaches may provide interesting perspectives on

magnitudes estimations, and additional studies need to be performed to understand to which extent these

models can be applied to explain the variety of results observed in the literature. One intriguing

observation is the finding that duration estimates were not only resilient to changes in numerosity or

surface, but also to the rate of sensory evidence. This finding is unexpected and runs counter-intuitive to

various findings in time perception. In this task, these robust findings suggest that unlike surface and
numerical estimates, duration may not rely on the accumulation of discretized sensory evidence.

504 DATA ACCESSIBILITY

506 AUTHORS' CONTRIBUTIONS

507 B.M., M.W. and V.v.W. designed the study. B.M. collected the data. B.M., M.W. and V.v.W. analyzed
508 the data. All authors co-wrote the manuscript and gave final approval for publication.

509 COMPETING INTERESTS

510 The authors declare having no competing interests.

511 FUNDING

512 This work was supported by an ERC-YStG-263584 to VvW.

513 ACKNOWLEDGMENTS

514 We thank members of UNIACT at NeuroSpin for their help in recruiting and scheduling participants, and
515 Baptiste Gauthier for discussions.

REFERENCES

1. Gallistel, C. R. Animal cognition: the representation of space, time and number. *Annu Rev Psychol* **40,** 155–189 (1989).

2. Walsh, V. A theory of magnitude: common cortical metrics of time, space and quantity. *Trends Cogn. Sci. (Regul. Ed.)* **7,** 483–488 (2003).

3. Gallistel, C. R. in *Space, Time and Number in the Brain* 3–12 (Academic Press, 2011).

4. Petzschner, F. H., Glasauer, S. & Stephan, K. E. A Bayesian perspective on magnitude estimation. *Trends Cogn. Sci. (Regul. Ed.)* **19,** 285–293 (2015).

5. Xuan, B., Zhang, D., He, S. & Chen, X. Larger stimuli are judged to last longer. *J Vis* **7,** 2.1-5 (2007).

6. Dormal, V., Seron, X. & Pesenti, M. Numerosity-duration interference: A Stroop experiment. *Acta Psychologica* **121,** 109–124 (2006).

7. Casasanto, D. & Boroditsky, L. Time in the mind: Using space to think about time. *Cognition* **106,** 579–593 (2008).

8. Oliveri, M. *et al.* Perceiving numbers alters time perception. *Neurosci. Lett.* **438,** 308–311 (2008).

9. Bottini, R. & Casasanto, D. Implicit Spatial Length Modulates Time Estimates, But Not Vice Versa. *Spatial Cognition VII* 152–162 (2010).

10. Stavy, R. & Tirosh, D. How Students (mis-) Understand Science and Mathematics: Intuitive Rules. (2000).

11. Dormal, V. & Pesenti, M. Numerosity-Length Interference. *Experimental Psychology* **54,** 289–297 (2007).

12. Javadi, A. H. & Aichelburg, C. When time and numerosity interfere: the longer the more, and the more the longer. *PLoS ONE* **7,** e41496 (2012).

13. Lambrechts, A., Walsh, V. & van Wassenhove, V. Evidence Accumulation in the Magnitude System. *PLoS ONE* **8,** e82122 (2013).

14. Wiener, M. & Thompson, J. C. The Effect of Background Context on the Size–Time Illusion. *Timing & Time Perception* **4,** 167–186 (2016).

15. Yamamoto, K., Sasaki, K. & Watanabe, K. The number–time interaction depends on relative magnitude in the suprasecond range. *Cogn Process* **17,** 59–65 (2016).

16. Cai, Z. G. & Wang, R. Numerical Magnitude Affects Temporal Memories but Not Time Encoding. *PLOS ONE* **9,** e83159 (2014).

17. Xuan, B., Chen, X.-C., He, S. & Zhang, D.-R. Numerical magnitude modulates temporal comparison: An ERP study. *Brain Research* **1269,** 135–142 (2009).

18. Roitman, J. D., Brannon, E. M., Andrews, J. R. & Platt, M. L. Nonverbal representation of time and number in adults. *Acta Psychol (Amst)* **124,** 296–318 (2007).

19. Calabria, M. & Rossetti, Y. Interference between number processing and line bisection: a methodology. *Neuropsychologia* **43,** 779–783 (2005).

20. Fischer, M. H., Castel, A. D., Dodd, M. D. & Pratt, J. Perceiving numbers causes spatial shifts of attention. *Nature Neuroscience* **6,** 555 (2003).

21. Viarouge, A. & de Hevia, M. D. The role of numerical magnitude and order in the illusory perception of size and brightness. *Front. Psychol.* **4,** (2013).

22. Dehaene, S., Bossini, S. & Giraux, P. The mental representation of parity and number magnitude. *Journal of Experimental Psychology: General* **122,** 371–396 (1993).

23. Ishihara, M., Keller, P. E., Rossetti, Y. & Prinz, W. Horizontal spatial representations of time: Evidence for the STEARC effect. *Cortex* **44,** 454–461 (2008).

24. Rammsayer, T. H. & Verner, M. The effect of nontemporal stimulus size on perceived duration as assessed by the method of reproduction. *Journal of Vision* **14,** 17–17 (2014).

25. Rammsayer, T. H. & Verner, M. Larger visual stimuli are perceived to last longer from time to time: The internal clock is not affected by nontemporal visual stimulus size. *Journal of Vision* **15,** 5–5 (2015).

26. Vallesi, A., McIntosh, A. R. & Stuss, D. T. How time modulates spatial responses. *Cortex* **47,** 148–156 (2011).

27. Newcombe, N. S., Levine, S. C. & Mix, K. S. Thinking about quantity: the intertwined development of spatial and numerical cognition. *WIREs Cogn Sci* **6,** 491–505 (2015).

28. Winter, B., Marghetis, T. & Matlock, T. Of magnitudes and metaphors: Explaining cognitive interactions between space, time, and number. *Cortex* **64,** 209–224 (2015).

29. Bonato, M., Zorzi, M. & Umiltà, C. When time is space: Evidence for a mental time line. *Neuroscience & Biobehavioral Reviews* **36,** 2257–2273 (2012).

30. Bueti, D. & Walsh, V. The parietal cortex and the representation of time, space, number and other magnitudes. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* **364,** 1831–1840 (2009).

31. Dormal, V., Dormal, G., Joassin, F. & Pesenti, M. A common right fronto-parietal network for numerosity and duration processing: An fMRI study. *Hum. Brain Mapp.* **33,** 1490–1501 (2012).

32. Pinel, P., Piazza, M., Le Bihan, D. & Dehaene, S. Distributed and overlapping cerebral representations of number, size, and luminance during comparative judgments. *Neuron* **41,** 983–993 (2004).

33. Hayashi, M. J. *et al.* Time Adaptation Shows Duration Selectivity in the Human Parietal Cortex. *PLoS Biol* **13,** e1002262 (2015).

34. Hubbard, E. M., Piazza, M., Pinel, P. & Dehaene, S. Interactions between number and space in parietal cortex. *Nat Rev Neurosci* **6,** 435–448 (2005).

35. Riemer, M., Diersch, N., Bublatzky, F. & Wolbers, T. Space, time, and numbers in the right posterior parietal cortex: Differences between response code associations and congruency effects. *NeuroImage* **129,** 72–79 (2016).

36. Nieder, A. Coding of abstract quantity by 'number neurons' of the primate brain. *J Comp Physiol A* **199,** 1–16 (2013).

37. Coull, J. T., Charras, P., Donadieu, M., Droit-Volet, S. & Vidal, F. SMA Selectively Codes the Active Accumulation of Temporal, Not Spatial, Magnitude. *Journal of Cognitive Neuroscience* **27,** 2281–2298 (2015).

38. Javadi, A. H., Brunec, I. K., Walsh, V., Penny, W. D. & Spiers, H. J. Transcranial electrical brain stimulation modulates neuronal tuning curves in perception of numerosity and duration. *NeuroImage* **102, Part 2,** 451–457 (2014).

39. Jazayeri, M. & Shadlen, M. N. Temporal context calibrates interval timing. *Nat Neurosci* **13,** 1020–1026 (2010).

40. Shi, Z., Church, R. M. & Meck, W. H. Bayesian optimization of time perception. *Trends Cogn. Sci. (Regul. Ed.)* **17,** 556–564 (2013).

41. Agrillo, C., Ranpura, A. & Butterworth, B. Time and numerosity estimation are independent: Behavioral evidence for two different systems using a conflict paradigm. *Cogn Neurosci* **1,** 96–101 (2010).

42. Petzschner, F. H. & Glasauer, S. Iterative Bayesian Estimation as an Explanation for Range and Regression Effects: A Study on Human Path Integration. *J. Neurosci.* **31,** 17220–17229 (2011).

43. Cicchini, G. M., Arrighi, R., Cecchetti, L., Giusti, M. & Burr, D. C. Optimal encoding of interval timing in expert percussionists. *J. Neurosci.* **32,** 1056–1060 (2012).

44. Yates, M. J., Loetscher, T. & Nicholls, M. E. R. A generalized magnitude system for space, time, and quantity? A cautionary note. *Journal of Vision* **12,** 9–9 (2012).

45. Rammsayer, T. H. & Verner, M. Evidence for different processes involved in the effects of nontemporal stimulus size and numerical digit value on duration judgments. *Journal of Vision* **16,** 13–13 (2016).

46. Vilares, I., Howard, J. D., Fernandes, H. L., Gottfried, J. A. & Kording, K. P. Differential representations of prior and likelihood uncertainty in the human brain. *Curr Biol* **22,** 1641–1648 (2012).

47. Wiener, M., Michaelis, K. & Thompson, J. C. Functional correlates of likelihood and prior representations in a virtual distance task. *Hum. Brain Mapp.* **37,** 3172–3187 (2016).

48. Levy, J. M., Namboodiri, V. M. K. & Hussain Shuler, M. G. Memory bias in the temporal bisection point. *Front Integr Neurosci* **9,** (2015).

49. Genovesio, A., Tsujimoto, S. & Wise, S. P. Neuronal Activity Related to Elapsed Time in Prefrontal Cortex. *Journal of Neurophysiology* **95,** 3281–3285 (2006).

50. Marcos, E., Tsujimoto, S. & Genovesio, A. Independent coding of absolute duration and distance magnitudes in the prefrontal cortex. *Journal of Neurophysiology* jn.00245.2016 (2016). doi:10.1152/jn.00245.2016

51. Nieder, A., Diester, I. & Tudusciuc, O. Temporal and Spatial Enumeration Processes in the Primate Parietal Cortex. *Science* **313,** 1431–1435 (2006).

52. Tudusciuc, O. & Nieder, A. Neuronal population coding of continuous and discrete quantity in the primate posterior parietal cortex. *PNAS* **104,** 14513–14518 (2007).

53. Dormal, V., Andres, M. & Pesenti, M. Dissociation of numerosity and duration processing in the left intraparietal sulcus: A transcranial magnetic stimulation study. *Cortex* **44,** 462–469 (2008).

54. Dormal, V. & Pesenti, M. Common and specific contributions of the intraparietal sulci to numerosity and length processing. *Hum. Brain Mapp.* **30,** 2466–2476 (2009).

55. Konkle, T. & Oliva, A. A Real-World Size Organization of Object Responses in Occipitotemporal Cortex. *Neuron* **74,** 1114–1124 (2012).

56. Riemer, M., Trojan, J., Kleinböhl, D. & Hölzl, R. A 'view from nowhen' on time perception experiments. *Journal of Experimental Psychology: Human Perception and Performance* **38,** 1118–1124 (2012).

57. Ratcliff, R. A note on modeling accumulation of information when the rate of accumulation changes over time. *Journal of Mathematical Psychology* **21,** 178–184 (1980).

58. Ratcliff, R., Smith, P. L., Brown, S. D. & McKoon, G. Diffusion Decision Model: Current Issues and History. *Trends in Cognitive Sciences* **20,** 260–281 (2016).

59. Churchland, A. K. & Kiani, R. Three challenges for connecting model to mechanism in decision-making. *Current Opinion in Behavioral Sciences* **11,** 74–80 (2016).

60. Barack, D. L. & Gold, J. I. Temporal trade-offs in psychophysics. *Current Opinion in Neurobiology* **37,** 121–125 (2016).

61. Brunton, B. W., Botvinick, M. M. & Brody, C. D. Rats and Humans Can Optimally Accumulate Evidence for Decision-Making. *Science* **340,** 95–98 (2013).

62. Murai, Y. & Yotsumoto, Y. Timescale- and Sensory Modality-Dependency of the Central Tendency of Time Perception. *PLOS ONE* **11,** e0158921 (2016).

63. Hanks, T. D., Mazurek, M. E., Kiani, R., Hopp, E. & Shadlen, M. N. Elapsed Decision Time Affects the Weighting of Prior Probability in a Perceptual Decision Task. *J. Neurosci.* **31,** 6339–6352 (2011).

64. O'Connell, R. G., Dockree, P. M. & Kelly, S. P. A supramodal accumulation-to-bound signal that determines perceptual decisions in humans. *Nat. Neurosci.* **15,** 1729–1735 (2012).

65. Kelly, S. P. & O'Connell, R. G. Internal and external influences on the rate of sensory evidence accumulation in the human brain. *J. Neurosci.* **33,** 19434–19441 (2013).

750 **FIGURES**

FIGURE 1

Figure 1: A Bayesian Magnitude System. A: Proposals for a common magnitude system in the brain suggest that
estimating one magnitude dimension (e.g. space) should be affected by the manipulation of any other magnitude
dimensions (e.g. time) (Walsh, 2003). Such interactions should show bidirectional interferences so that, in our example,
manipulating the spatial dimension of an event should affect the estimation of its duration in a comparable manner as
manipulating its time dimension would affect the estimation of its spatial dimension. **B:** To account for bidirectional
interferences across magnitudes in a Bayesian framework for magnitude estimation, a common global or amodal prior can
be posited to constrain the estimations of all magnitude dimensions [4]. Under such model, increasing (decreasing) the
value of one magnitude should increase (decrease) the estimation of another magnitude.

Figure 2: Experimental Design. A: On a given trial, participants were presented with a word ("durée", "nombre" or "surface") indicating the dimension to estimate. In Experiment 1, one magnitude dimension could vary +/- 25 %, 10 %, and 5% of its mean value, while the second one was set to its minimal or maximal value, and the third one to its mean value (Table 1). At the end of the stimulus presentation, participants used a vertical scale to estimate the target magnitude. **B:** Three distributions were used for evidence accumulation: while D linearly accumulates over time (black trace), the rate of dot presentation could be manipulated to control N and S. Experiment 1 tested a linear distribution (filled black trace); Experiment 2 tested two distributions: a fast-slow (filled grey trace) and a slow-fast (dotted grey trace) distribution. **C:** Equated task difficulty across magnitudes. For illustration purposes, the psychometric curve captures the grand average performance obtained for the estimation of Duration, Number and Surface when all non-target dimensions were kept at their mean value. The task difficulty was equated across magnitude dimensions so that no differences in discriminability (PSE and WR) existed between the tested dimensions. Bars are 2 s.e.m. **D:** Predictions for the effect of non-target manipulations on the estimation of the target magnitude dimension. Left panel: varying the target magnitude while keeping the non-target dimensions to their mean values provided the control central tendency and intercept. In a common Bayesian magnitude estimation system [4], comparable tendency and intercept should be predicted pending controlled matching between magnitudes and task difficulty (panel C). Right panel: estimation of D while N is set to its maximal value (in green, Nmax). Maximal value in non-target magnitude may affect the central tendency and the intercept if an amodal global prior common to D and N is implicated in the estimation of duration. In this example, N_{max} would bias the lowest (highest) duration values towards larger (smaller) values and lead the intercept to move upwards.

FIGURE 3

Figure 3: Duration affects the estimation of S and N (Experiment 1). A: Categorical quantifications and PSE. The percentage of « + » responses as a function of the target magnitude dimension (D: left panel, N: middle panel, S: right panel) were fitted on a per individual basis. The inset in the top left of each figure depicts the shift in PSE for each experimental condition compared to the control condition represented by the black vertical line. No effects of N or S on D were found; no effects of S on N or of N on S were found; both N and S were significantly overestimated when presented during the shortest duration (Dmin, red) and underestimated when presented with the longest duration (Dmax, orange). **B**: Continuous judgments. Individual performances (transparent dots) and mean performances (filled dots) for the estimation of D (left panels) with changing N (green) and S (blue); of N (middle panels) with changing D (red) and S (blue); and of of S (left panels) with changing D (red) and N (green). The continuous scale was mapped from 0 to 100. The dotted line is the ideal observer's performance. All experimental conditions showed a central tendency. No effects (intercept or central tendency) of N or S on D were found (left top and bottom graphs, respectively); no effects of S on N (middle bottom graph) or of N on S (left bottom graph) were found. Significant main effects of D were found on the central tendency and the intercept of N (middle top) and S (left top). N_{min}: miminal numerosity value; N_{max}: maximal numerosity value; S_{min}: minimal surface value; S_{max}: maximal surface value; D_{min}: minimal duration value; D_{max}: maximal duration value. *** $p < 0.001$; ** $p < 0.01$; * $p < 0.05$; bars are 2 s.e.m.

FIGURE 4

Figure 4: Central Tendency Effects in Magnitude Estimation (Experiment 1). A: Average continuous estimates for all three magnitude dimensions as target, collapsed across all non-target magnitude dimensions. Magnitudes were normalized as a percentage of the maximum presented magnitude value, with zero representing the smallest and 100 representing the largest magnitude presented. Continuous estimates were similarly normalized as a percentage of the sliding scale, with zero representing a minimal estimate of "-" and 100 representing a maximal estimate of "+". The dashed identity line indicates where estimates should lie for veridical performance. Deviations from this identity line (arrows) exhibited central tendency, wherein smaller and larger magnitudes were over- and under-estimated, respectively. **B:** Slope values extracted from a best fitting linear regression in A quantify the degree of central tendency, with smaller values indicating greater regression to the mean. Similar slope values were observed for duration and number, but significantly lower estimates were found for surface estimates than either duration or number. **C:** Correlation matrix of slope values for every target and non-target magnitude trial type. Each target magnitude dimension was tested in the presence of three possible values (min, mean, max) of a non-target dimension. Outlined pixels represent those Pearson correlation coefficients that survived a multiple comparison correction. Slope values across non-target dimensions were correlated within each target dimension (sections along the diagonal); crucially, slope values for duration and number were correlated with each other when each one was the target magnitude (lower left and upper right sections). Further, no correlation between surface and either duration or number as target dimensions were observed (middle top and bottom sections). * indicates p<.05; bars are 2 s.e.m.

848   **FIGURE 5**

Figure 5 : **The Accumulation Regime affects Numerosity and Surface but not Duration (Experiment 2). A:** Psychometric curves illustrate the grand average proportion of "+" as a function of duration (left panel), number (middle panel) or surface (right panel) when the sensory evidence accumulation was manipulated. The SlowFast (SF) results are reported in light gray, the FastSlow (FS) results are reported in black. Left panel: top inset reports the mean PSE of duration estimation observed for each distribution as compared to the ideal observer (vertical black line). Middle panel: top inset reports the mean PSE of number estimation observed for each distribution (SF, FS) and manipulation of D (D_{mean}: black, D_{min}: red, D_{max}: pink). Right panel: top inset reports the mean PSE of surface estimation observed for each distribution (SF, FS) and manipulation of D (D_{mean}: black, D_{min}: red, D_{max}: pink). The estimation of D was not affected by the accumulation regime of sensory evidence whereas N and S were overestimated in the FastSlow as compared to the SlowFast distribution. N and S were also affectd by manipulating of the non-target duration. **B:** Individual (transparent dots) and mean (filled dots) continuous judgments for the estimation of D (left panel), N (middle panel) and S (right panel) with SlowFast (top row) and FastSlow (bottom row) regimes

of sensory evidence accumulation. The dotted line is the ideal observer's performance. D_{min}: minimal duration value; D_{max}: maximal duration value. *** $p < 0.001$; ** $p < 0.01$; * $p < 0.05$; bars are 2 s.e.m.

FIGURE 6

Figure 6: Accumulation regime influences Bayesian estimates. Our results suggest that the speed or rate of sensory evidence accumulation early in the trial (shaded region) and the duration of the trial affects surface and numerosity estimations. Additionally, slower rates of sensory evidence are associated with greater uncertainty; greater uncertainty results in increased reliance on the priors. To accommodate these findings, we suggest that the rate of sensory evidence is effectively estimated independently of the total duration although the duration may regulate the noise level in sensory evidence accumulation of quantities.

898 **TABLES**

901 **TABLE 1**

	DURATION [.75; .9; .95; 1.05; 1.1; 1.25] **x** D_{mean}	SURFACE [.75; .9; .95; 1.05; 1.1; 1.25] **x** S_{mean}	NUMEROSITY [.75; .9; .95; 1.05; 1.1; 1.25] **x** N_{mean}
CONTROL	$\mathbf{S_{mean}}$, $\mathbf{N_{mean}}$	$\mathbf{D_{mean}}$, $\mathbf{N_{mean}}$	$\mathbf{S_{mean}}$, $\mathbf{D_{mean}}$
MIN	$\mathbf{S_{min}}$, N_{mean}	$\mathbf{D_{min}}$, N_{mean}	$\mathbf{S_{min}}$, D_{mean}
	S_{mean} , $\mathbf{N_{min}}$	D_{mean} , $\mathbf{N_{min}}$	S_{mean} , $\mathbf{D_{min}}$
MAX	$\mathbf{S_{max}}$, N_{mean}	$\mathbf{D_{max}}$, N_{mean}	$\mathbf{S_{max}}$, D_{mean}
	S_{mean} , $\mathbf{N_{max}}$	D_{mean} , $\mathbf{N_{max}}$	S_{mean} , $\mathbf{D_{max}}$

Table 1: Full design of Experiment 1 testing the interactions across magnitudes with a linear sensory evidence
accumulation regime. The tested magnitude dimension could take 6 possible values corresponding to 75, 90, 95, 105, 110 and
125 % of its mean value. In CONTROL trials (first row) of the duration condition (D, second column), participants estimated the
duration of the trial when D varied between the 6 possible values while the surface and number dimensions were kept to their
mean values (S_{mean} and N_{mean}, respectively). In MIN trials (second row), participants estimated the magnitude of a given
dimension (e.g. D) varying between the 6 possible values while one of the two non-target magnitude dimensions was kept at its
mean value (e.g. N_{mean}) and the other was set to its minimal value (e.g. S_{min}). In MAX trials (third row), participants estimated the
magnitude of a given dimension varying between the 6 possible values while one of the two non-target dimensions was kept at its
mean value and the other was set to its maximal value. In total, 72 trials per experimental condition were collected (i.e. 12 trials
per tested magnitude value in all possible combinations). D: duration; S: surface; N: number; min = minimal; max = maximal.

TABLE 2

	DURATION $[.75; .9; .95; 1.05; 1.1; 1.25]$ **x** D_{mean}	SURFACE $[.75; .9; .95; 1.05; 1.1; 1.25]$ **x** S_{mean}	NUMEROSITY $[.75; .9; .95; 1.05; 1.1; 1.25]$ **x** N_{mean}
CONTROL	S_{mean} , N_{mean}	D_{mean} , N_{mean}	S_{mean} , D_{mean}
MIN		D_{min} , N_{mean}	
			S_{mean} , D_{min}
MAX		D_{max} , N_{mean}	
			S_{mean} , D_{max}

Table 2: Full design of Experiment 2 testing the effects of duration and FastSlow and SlowFast sensory evidence accumulation regime on magnitude estimation. In CONTROL trials, as in Experiment 1, participants estimated the value of each target magnitude dimension when non-target dimensions were kept to their mean values. In MIN and MAX trials, participants estimated S or N when D was the shortest (D_{min}) or the longest (D_{max}) and the other dimension was kept to its mean value (N_{mean} and S_{mean}, respectively). Importantly in Experiment 2, the clouds of dots could accumulate over time according to two accumulation regimes (FS:FastSlow, and SF:SlowFast). A total of 144 trials per experimental condition was collected (i.e. 12 trials for a given magnitude value in a specific condition x 2 distributions).

932