WikiGenomes: an open Web application for community consumption and curation of gene
annotation data in Wikidata.

Tim E. Putman[1], Sebastien Lelong[1], Sebastian Burgstaller-Muehlbacher[1] , Andra Waagmeester[2],
Colin Diesh[4], Nathan Dunn[3], Monica Munoz-Torres[3] Gregory S. Stupp[1], Andrew I. Su[1] and
Benjamin M. Good[1]

1. Department of Molecular and Experimental Medicine, The Scripps Research Institute, La
Jolla, USA
2. Micelio, Antwerp, Belgium
3. Environmental Genomics and Systems Biology Division, Lawrence Berkeley National
Laboratory, Berkeley, California.
4. Division of Animal Sciences, University of Missouri, Columbia, MO 65211, USA

{tputman, bgood, asu}@scripps.edu

Abstract

With the advancement of genome sequencing technologies, new genomes are being
sequenced daily. While these sequences are deposited in publicly available data warehouses,
their functional and genomic annotations (beyond genes which are predicted automatically)
mostly reside in the text of primary publications. Professional curators are hard at work
extracting those annotations from the literature for the most studied organisms and depositing
them in structured databases. However, the resources don't exist to fund the comprehensive
curation of the thousands of newly sequenced organisms in this manner. Here, we describe
WikiGenomes (wikigenomes.org), a web application that facilitates the consumption and
curation of genomic data by the entire scientific community. WikiGenomes is based on
Wikidata, an openly editable knowledge graph with the goal of aggregating published
knowledge into a free and open database. WikiGenomes empowers the individual genomic
researcher to contribute their expertise to the curation effort and integrates the knowledge into
Wikidata, enabling it to be accessed by anyone without restriction.

Introduction

Sequencing an organism's genome has become a routine procedure in the life sciences, but until that genome is annotated, it tells us very little about the organism. The knowledge captured in genomic and functional annotations provides the 'blueprint' of the biology of the organism that can be leveraged to drive all manner of scientific inquiry. The knowledge represented by annotations is mostly published in the free text of scientific journal articles. In order to make this type of knowledge computable and accessible as structured annotations, we typically rely on centralized curation efforts like those supported by model organism databases such as the Zebrafish Model Organism database (ZFIN) (1) and Mouse Genome Informatics (MGI) (2). Unfortunately, the vast majority of sequenced genomes do not have a dedicated curation team, hence new curation models need to be explored.

Wikidata is a recently developed project of the Wikimedia Foundation that enables the collaborative construction of a centralized graph database (3). It was initially created as a central hub for structured data across all the language-specific Wikipedias, but its potential applications are much broader. Wikidata follows the Resource Description Framework (RDF) (https://www.w3.org/RDF/ (4)) model, the W3C standard for data interchange. RDF links related entities together into 'triples', defined by a subject concept, an object concept, and a predicate that describes the relationship between them. This semantic structure allows the modeling of complex systems in a queryable knowledge graph (3,5). The contents of Wikidata currently represent 28 million concepts, from all domains of knowledge, spanning over 1.3 billion triples (grafana.wikimedia.org/dashboard/db/wikidata-query-service).

While other resources have also mapped their data onto an RDF data model (6–8), Wikidata is unique in that it allows both read and write access. Wikidata content can be both queried and edited programmatically via an application programming interface (API) (www.**wikidata**.org/w/**api**.php) and can be queried using the Wikidata Query Service (WDQS) (query.wikidata.org) via the SPARQL Protocol and RDF Query Language (4,9,10). SPARQL is a query language for graph databases similar in syntax to SQL for relational databases.

Notably, all Wikidata content is in the public domain, eliminating the potential for problematic restrictions on downstream reuse and redistribution (11). It is populated through various programs ("bots") (12) en masse, and through the discrete contributions of individual users through Wikidata's own web interface or through a handful of tools developed by Wikimedia Foundation's WMF Labs (tools.wmflabs.org/).

There is growing appreciation within the bioinformatics community for the potential of Wikidata as an open resource that can be populated, queried and edited by anyone. However, Wikidata has not yet widely reached the biological domain experts who have much to gain and contribute. Taking advantage of the querying capabilities of Wikidata can be a challenge for biologists as structured query languages like SPARQL are not commonly part of a researcher's toolkit. Further, the domain-specific data models (e.g. the patterns of connections between genes, proteins, and their annotations (www.wikidata.org/wiki/Wikidata:WikiProject_Molecular_biology)) underlying the content in the Wikidata graph are not readily apparent nor automatically enforced by the generic Wikidata web interface. This makes providing meaningful contributions to the graph a challenge for the researcher and a roadblock for community curation in general. While Wikidata offers a powerful technical platform for constructing an open community knowledge base of unprecedented scope, it cannot reach this potential without the large-scale participation of experts in the domains of knowledge that it seeks to represent. To attract specialists, e.g. life scientists, its

82   content must be presented in compelling applications that are more useful than the applications
83   that these specialists currently use to do their work.
84
Here, we describe WikiGenomes (www.wikigenomes.org), the first domain-specific application
built on Wikidata, tailored to the needs of biomedical researchers. WikiGenomes is a Web
application that is designed and built to allow the genomic researcher to contribute to curating
the knowledge they are discovering. It facilitates both the consumption and community curation
of genomic data in Wikidata, extending the reach of the biocuration effort deeper into the long
tail of sequenced genomes.

Results

Wikidata
Our group and others have been working to populate Wikidata with a comprehensive and
centralized knowledge graph that represents biomedical knowledge in a structured, queryable,
and computable format (5,12). Previously, we loaded genomic data from a variety of organisms
including *Homo sapiens* "human" (www.wikidata.org/wiki/Q15978631), *Mus musculus* "mouse"
(www.wikidata.org/wiki/Q83310), *Rattus norvegicus* "brown rat"
(www.wikidata.org/wiki/Q184224), *Saccharomyces cerevisiae S288c* "baker's yeast"
(www.wikidata.org/wiki/Q27510868), and *Macaca nemestrina* "Southern pig-tailed macaque"
(www.wikidata.org/wiki/Q618026). In this work, we systematically expanded this list to include
all genes (390,719) and proteins (372,178) from the 120 NCBI prokaryotic reference genomes
(https://www.ncbi.nlm.nih.gov/genome/browse/reference/) (**Figure 1**). These organisms are all
reasonably well-studied, but in general did not have community-maintained genome databases.
In addition to organism/genetic data, we contribute to and maintain data for proteins, chemical
compounds and diseases (5,12,13).
106
To ensure that the work we are doing in Wikidata is trusted as high quality by the scientific
community, our group references the source and provenance of all claims in a consistent and
structured way. Direct links to the data sources allow the user to see for themselves the source
of an annotation and read further in the referenced publication to come to their own
conclusions. Standards for representing the evidence and provenance of any claim made using
our infrastructure is outlined in a detailed project wiki
(https://www.wikidata.org/wiki/User:ProteinBoxBot/evidence). References can be accessed
either manually by the web interface or programmatically via the Wikidata API
(https://www.wikidata.org/w/api.php).
116
In addition to recording data provenance, Wikidata claims can be qualified by how the initial
claim was made. For example a Gene Ontology (GO) annotation can be qualified using the
Gene Ontology Evidence Codes to indicate the determination method for this claim. The
combination of references and qualifiers provides the necessary context to the stated claim,
making it more trustworthy and useful. This model allows the researcher to, for example, treat
electronically curated annotations differently from those that were manually curated.
123
Data Aggregation and Maintenance in Wikidata
As a foundation, we loaded key information for each genome about each gene and gene
product into Wikidata. Wikidata is not simply an open database into which data can be
dumped. It is a structured knowledge graph that requires careful consideration and community
consensus in its design. Currently, these entity models in Wikidata are not comprehensive
mirrors of those models in their sources (e.g. NCBI, UniProt, etc...). We have chosen to

aggregate key data from a variety of sources, creating a minimal viable model that can be
expanded on by us or others.   The basic data in Wikidata for all the organisms we have loaded
includes genomic position and orientation, Gene Ontology annotations, Enzyme Commission
Numbers, InterPro domains and external database identifiers. The data model that we have
created in Wikidata, and the sources of the annotations and entities are illustrated and reported
in Figure 1.
136
137   <FIGURE 1 HERE>
138
Data retrieval, standardization, loading and maintenance are done routinely (currently monthly)
via our publically available WikidataIntegrator Python package
(https://github.com/SuLab/WikidataIntegrator and https://pypi.python.org/pypi/wikidataintegrator)
as well as M, and we are working towards custom update schedules that would follow each
source's update schedule. WikidataIntegrator interacts with the Wikidata API to create new or
edit existing Wikidata items when appropriate. The data points are mapped to the appropriate
"properties" in Wikidata (**Supplemental Table 1**). Mapping these properties to a new semantic
model in Wikidata is key to aggregation of data from different sources into a single cohesive
graph that represents our collective knowledge of each concept and the relationships between
those concepts.
149
An example of other data we have loaded and maintain is the Gene Ontology
(http://www.geneontology.org/), which we have loaded to Wikidata to provide semantic
framework for gene product annotations. The structure was represented as a hierarchical tree
using the Wikidata 'Subclass of' (P279) property. We programmatically gathered professionally
curated Gene Ontology annotations from the European Bioinformatics Institute's QuickGo API
(https://www.ebi.ac.uk/QuickGO/), and NCBI's Gene Database FTP resources
(ftp.ncbi.nlm.nih.gov/gene/DATA/gene2go.gz) for all of the annotated gene products in Wikidata
(5,12,13). Any gene product item in Wikidata that is annotated with a specific GO term will be
linked to that GO term's Wikidata item via the appropriate Wikidata GO property (e.g. *molecular*
function P680) (**Figure 1**). Our methods for loading and maintaining other domains, such as
disease and chemical data, are described in our previous works (12,13).
161
WikiGenomes
To allow non-developer biologists to interface with the genomic data knowledge graph that we
developed in Wikidata, we built WikiGenomes. WikiGenomes currently supports the 120 NCBI
Bacterial reference genomes (https://www.ncbi.nlm.nih.gov/genome/browse/reference/) and
thier genes and gene products.   We designed WikiGenomes to allow a user to consume
genomic data from any organism in Wikidata in a single web application. The user begins by
selecting an organism using a type-ahead search feature on the application's landing
page. Once an organism has been selected, the application is directed to the primary interface
page where the data for the selected organism is loaded and displayed. In addition to providing
access to genetic data for these organisms, WikiGenomes also allows users to directly
contribute genomic and protein annotations to the Wikidata knowledge graph (**Figure 2**).
173
174   <FIGURE 2 HERE >
175
The primary page of the interface is divided into four windows that render the data for the
selected organism: its genome, genes/proteins, and annotations for the currently displayed
gene/protein. To select and load data, there are two forms that allow organism and gene/protein
selection (**Figure 3**). Each of the data elements are drawn dynamically from the Wikidata

knowledge graph using the WDQS SPARQL query system. The genome data is rendered in the
open source JBrowse genome browser (14,15). The WebApollo and JBrowse development
teams expanded on already existing code in the JBrowse framework to allow JBrowse to load
annotations gathered from Wikidata SPARQL queries to the WDQS endpoint (16). The DNA
sequences themselves are curated by the RefSeq Project (16) and because bulk primary data is
not suitable for storing in Wikidata, are collected monthly from GenBank's FTP web service
(See **Supplementary File 1** for all Wikidata queries that drive the system and **Supplementary**
Table 2 for the FTP paths of data collected from NCBI). The open source code for
WikiGenomes is available at https://github.com/putmantime/CMOD.Django.
189
190   <FIGURE 3 HERE>
191
While bioinformaticians can programmatically gather data and upload it to Wikidata,
WikiGenomes provides a domain specific editing interface for non-programmers. WikiGenomes
distills editing tasks into straightforward forms that guide the user through the process of
building annotations. Internally, the server translates the data entered through these forms into
the structures needed to populate the Wikidata knowledge graph and submits them via the
Wikidata API (**Figure 2**). WikiGenomes uses the WikiMedia OAuth extension to allow users to
edit Wikidata using their own Wikidata account (www.mediawiki.org/wiki/Extension:OAuth). This
makes it possible to utilize the Wikidata edit history to track the contributions of individual
editors, potentially offering ways to reward good editors as well as mechanisms for detecting
vandalism automatically (Good 2012).
202
203   *Community Curation Forms*
204   There are currently two editing forms in WikiGenomes, one for adding functional annotations
205   and one for adding operon annotations. The forms are designed to allow a user to curate data
206   from a publication by prompting the user to provide the necessary information and references to
207   make a useful contribution to Wikidata (**Figure 4**).
208
209   <FIGURE 4 HERE>
210
Gene Ontology Form
Molecular function, cellular component and biological process annotations types are added to
Wikidata (and subsequently WikiGenomes) through the Gene Ontology Annotation Form
(**Figure 4A**). The WikiGenomes GO form allows users to make structured GO annotations to
gene products by identifying an annotation from a primary publication, mapping the annotation
to the proper GO term, and then pushing it to Wikidata referencing the publication. We have
chosen to adhere to the GO annotation model, rather than user generated free-text, to guide the
annotation process in a consistent manner. This is accomplished by using the form to search
Wikidata by typeahead for the proper GO term (as mentioned we maintain the GO ontology in
Wikidata), selecting the proper GO evidence code (http://geneontology.org/page/guide-go-
evidence-codes) to provide a determination method (i.e. EXP: Inferred from Experiment
http://geneontology.org/page/exp-inferred-experiment), and select the publication as a reference
using the PubMed Identifier (PMID) (**Figure 4A**). When the annotation is submitted, the
annotation data is pushed to Wikidata using the Wikidata API and the annotation will be
viewable in WikiGenomes shortly after (there is ~5 min delay as the Wikidata SPARQL endpoint
updates).
227
Operon Form

Regulatory annotations are rare in large data warehouses like NCBI and regulatory annotation
curation would greatly benefit from community involvement. To this end we created a form to
allow user curation of prokaryotic operons (**Figure 4B**). The WikiGenomes Operon Form works
similarly to the GO form in using typeahead search boxes to retrieve relevant data from
Wikidata, as well as input boxes to upload novel data. The user can create a new operon or
add genes to an existing operon, input the determination method and reference the PMID, then
submit that annotation and (as in the GO Form) our backend framework will do the work of
creating the structured and standardized annotation in Wikidata.
237
Discussion
WikiGenomes was built as a proof a concept that Wikidata can serve as a universal database
backend for domain specific applications. WikiGenomes is in the stage of functional prototype
and usage stats are thus not yet available. WikiGenomes provides a working demonstration of
a new technical pattern that, we posit, offers the potential to fundamentally change how
biocuration is organized in ways that, over time, will result in a more efficient movement of
knowledge throughout the scientific community. While this is not yet a tool in common usage,
we believe that it shows the potential of centralized, community owned knowledge aggregation
projects like Wikidata, and hope it will ultimately help stimulate community curation at both the
level of individual researchers and bioinformatics laboratories.

Distributed Curation and Access, Centralized and Integrated Content

The incorporation of operon information for *Chlamydia pneumoniae* into WikiGenomes
illustrates the pattern of centralizing content while decentralizing control. Despite prokaryotic
operons being heavily studied, the actual annotations are rarely included in genomic assemblies
and often reside in supplemental tables of primary publications. *Chlamydia pneumoniae* is a
pathogen affecting animals and humans causing lung infections
(www.cdc.gov/pneumonia/atypical/cpneumoniae/index.html). The operons of the *C.*
pneumoniae strain *CWL029* genome were annotated in a 2011 study (17) but access to this
information was only provided in the form of a supplementary Excel file. To our knowledge,
none of those ~200 experimentally derived annotations have been curated in GenBank or in
operon-specific databases such as DOOR (18) or ODB3 (19). Illustrating the 'small data to big
data' approach of WikiGenomes, we loaded this content programmatically via the Wikidata API.
Individual domain experts could also have loaded these data via the operon form (**Figure 4b**).
Using either method, these annotations are now fully-linked data, viewable as a track in the
WikiGenomes browser and in any other application drawing content from the open Wikidata
knowledge base. Integrating this knowledge into the central Wikidata knowledge graph creates
queryable connections to other organisms, diseases, drugs and small molecules. The more
extensive the knowledge graph becomes, the more it can be used to explore biological research
questions (5),(12,13).

Stepping Towards Community Curation

So far, the great majority of content accessible through WikiGenomes has been added by our
research group. While this content is valuable, it is only a starting point on the path towards an
open, community-curated knowledge graph for biology. The fundamental logic of the
community curation pattern has been well-described many times (20–22). By incorporating the
scientific community into the process of curating the knowledge that they are generating and
consuming, we could, in theory, dramatically increase the scale at which biocuration can
collectively operate.   Here, as usual, the devil is in the details. With certain notable exceptions,
most attempts to stimulate community curation fail to attract enough editors to achieve their

goals. With the WikiGenomes application, we make no claim that we have solved this challenge
of incentives and recruitment. To realize its vision, WikiGenomes will need to transition to a
model where the content consumers become the content producers. To make this transition, a
number of problems must be addressed. In facing them, some inspiration can be taken from
one of WikiGenomes' most important and successful predecessors, the Gene Wiki effort to
organize functional gene information in the context of Wikipedia articles (23).
282
The key distinction between the Gene Wiki and other projects with very similar goals but less
success in recruiting contributors (20,24), was that it was embedded directly in Wikipedia. The
Wikipedia context provided the Gene Wiki with immediate discoverability (e.g. Gene Wiki
articles rank highly in Google search results), connection to a large community of editors, and a
proven social and technical framework for supporting large-scale community content
creation. Wikidata now offers most of these same characteristics for structured information. Its
content is highly discoverable, with hyperlinks coming in from nearly every Wikipedia article and
a high performance query engine that provides a single point of entry into all of its data. It has
already attracted thousands of editors from the Wikipedia communities as well as external
groups more focused on structured data. It offers a user interface for human editing of content
as well as an effective API for programmatic updates. Finally, communities are forming around
it in similar ways to Wikipedia. There are social structures in place for administration of the
higher-level data structures it utilizes (e.g. the allowed properties) and emerging domain-
focused communities such as WikiProject Medicine and WikiProject Molecular and Cellular
Biology that work to build and maintain domain-specific content within the larger knowledge
graph.
299
With all of those similarities, one key distinction is that structured information is much more
difficult to edit and to present effectively than the hypertext of Wikipedia. While the
Wikidata interface (www.wikidata.org) provides a useful, general-purpose foundation for
navigating and editing its content, it is not nearly sufficient as a way to connect end-users with
the information they need. WikiGenomes provides a genome-focused, editable view of
biological knowledge in Wikidata; it offers the large community of scientists an easy way to both
access and share knowledge in a structured way, thus opening a door towards open community
curation.
308
Conclusions and Next Steps
While tools like WikiGenomes are a step in the right direction for converting 'small data' to 'big
data' several key problems remain to be solved in the community curation model. Interfaces
like WikiGenomes that provide access to content drawn from openly editable public sources
should consider ways to help users assess the trustworthiness of content based on automatic
inspection of the provenance information provided in associated references as well as in the
histories of the editors who generated the content. In addition to making trust computable, such
provenance information will also be of value in attributing the work of the curators who devote
time to these open resources - potentially providing an aspect of the solution to the incentive
problem. Aside from presenting trust-related information, the forms for adding content should
be designed in ways that help domain experts generate high-quality annotations in the first
place. The addition of documentation, logical constraints based on underlying ontologies, and
helper interfaces are areas where the experience of the existing biocuration resources would be
very valuable.
323
WikiGenomes is a first foray into the world of applications that could be built with the content
that is growing in Wikidata. As that content expands and diversifies, many new applications will

need to be created to bring that information most effectively to the people that need it and who
can contribute to it. Wikidata is unlikely to be the end-all and be-all of open biological
knowledge bases. While it has a number of advantages that make it an ideal platform for
initiating the endeavor of linking all biological knowledge, it has limitations in terms of content
scope that will eventually need to be overcome through the provision of equally open systems
that are tailored more specifically to biological content.
332
Immediate further development of WikiGenomes is focused on teaming up with biocuration
groups to help refine the user experience and identify important areas where the data model
needs to be expanded. One example would be to work with further with the WebApollo team to
integrate WebApollo (25) into WikiGenomes, taking advantage of WebApollo's annotation
features. Another useful feature we will work towards implementing is the ability to export data
from Wikidata via WikiGenomes in standard formats (e.g. GFF3). We hope to eventually create
a tool in WikiGenomes that brings the biocuration and basic research communities together,
creating a more efficient system that will allow for more comprehensive reference genome
curation of biological knowledge than is currently possible. Eventually we would make
WikiGenomes a deployable toolkit that smaller research communities can customize and deploy
while easily sharing their work with the global community.
344
345

Acknowledgements
We would like to thank the Wikimedia foundation for supporting the Wikidata project, and
especially the Wikidata community for their efforts in populating it. This project was supported
by the National Institutes of General Medical Sciences (R01GM089820) and NIH Common Fund
programs for Big Data to Knowledge (U54GM114833) and Extracellular RNA Communication
(U54DA036134).

References

1. Howe, D. G., Bradford, Y. M., Conlin, T., et al. (2013) *Nucleic Acids Res.*, **41**, D854–60, ZFIN, the Zebrafish Model Organism Database: increased support for mutants and transgenics.
2. Bult, C. J., Eppig, J. T., Blake, J. A., et al. (2016) *Nucleic Acids Res.*, **44**, D840–7, Mouse genome database 2016.
3. Vrandečić, D. and Krötzsch, M. (2014) *Commun. ACM*, **57**, 78–85, Wikidata: a free collaborative knowledgebase.
4. Quilitz, B. and Leser, U. In *The Semantic Web: Research and Applications*; Bechhofer, S.; Hauswirth, M.; Hoffmann, J.; Koubarakis, M., Eds.; Lecture Notes in Computer Science; Springer Berlin Heidelberg, 2008; pp. 524–538.
5. Putman, T. E., Burgstaller-Muehlbacher, S., Waagmeester, A., et al. (2016) *Database*, **2016**, Centralizing content and distributing labor: a community model for curating the very long tail of microbial genomes.
6. Quest, D. J., Land, M. L., Brettin, T. S., et al. (2010) *BMC Bioinformatics*, **11 Suppl 6**, S15, Next generation models for storage and representation of microbial biological annotation.
7. Miles, A., Zhao, J., Klyne, G., et al. (2010) *J. Biomed. Inform.*, **43**, 752–761, OpenFlyData: an exemplar data web integrating gene expression data on the fruit fly Drosophila melanogaster.
8. Cheung, K.-H., Yip, K. Y., Smith, A., et al. (2005) *Bioinformatics*, **21 Suppl 1**, i85–96, YeastHub: a semantic web use case for integrating data in the life sciences domain.
9. Prud'Hommeaux, E., Seaborne, A. and Others (2008) *W3C recommendation*, **15**, SPARQL query language for RDF.
10. Pérez, J., Arenas, M. and Gutierrez, C. In *The Semantic Web - ISWC 2006*; Cruz, I.; Decker, S.; Allemang, D.; Preist, C.; Schwabe, D.; Mika, P.; Uschold, M.; Aroyo, L. M., Eds.; Lecture Notes in Computer Science; Springer Berlin Heidelberg, 2006; pp. 30–43.
11. Himmelstein, D. Integrating resources with disparate licensing into an open network https://thinklab.com/discussion/integrating-resources-with-disparate-licensing-into-an-open-network/107#1.
12. Burgstaller-Muehlbacher, S., Waagmeester, A., Mitraka, E., et al. (2016) *Database*, **2016**, Wikidata as a semantic framework for the Gene Wiki initiative.
13. Elvira Mitraka, Andra Waagmeester, Sebastian Burgstaller, Lynn M. Schriml, Benjamin M. Good, Andrew I. Su *Proceedings of the 2015 Swat4LS International Conference in Cambridge England*, Wikidata: A platform for data integration and dissemination for the life sciences and beyond.
14. Skinner, M. E., Uzilov, A. V., Stein, L. D., et al. (2009) *Genome Res.*, **19**, 1630–1638, JBrowse: a next-generation genome browser.
15. Buels, R., Yao, E., Diesh, C. M., et al. (2016) *Genome Biol.*, **17**, 66, JBrowse: a dynamic web platform for genome visualization and analysis.
16. O'Leary, N. A., Wright, M. W., Brister, J. R., et al. (2016) *Nucleic Acids Res.*, **44**, D733–45, Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation.
17. Albrecht, M., Sharma, C. M., Dittrich, M. T., et al. (2011) *Genome Biol.*, **12**, R98, The transcriptional landscape of Chlamydia pneumoniae.
18. Mao, F., Dam, P., Chou, J., et al. (2009) *Nucleic Acids Res.*, **37**, D459–63, DOOR: a database for prokaryotic operons.
19. Okuda, S. and Yoshizawa, A. C. (2011) *Nucleic Acids Res.*, **39**, D552–5, ODB: a database for operon organizations, 2011 update.
20. Mons, B., Ashburner, M., Chichester, C., et al. (2008) *Genome Biol.*, **9**, R89, Calling on a million minds for community annotation in WikiProteins.

21. Howe, D., Costanzo, M., Fey, P., et al. (2008) *Nature*, **455**, 47–50, Big data: The future of
biocuration.
22. Pico, A. R., Kelder, T., van Iersel, M. P., et al. (2008) *PLoS Biol.*, **6**, e184, WikiPathways:
pathway editing for the people.
23. Huss, J. W., 3rd, Orozco, C., Goodale, J., et al. (2008) *PLoS Biol.*, **6**, e175, A gene wiki for
community annotation of gene function.
24. Hoffmann, R. (2008) *Nat. Genet.*, **40**, 1047–1051, A wiki for the life sciences where
authorship matters.
25. Lee, E., Helt, G. A., Reese, J. T., et al. (2013) *Genome Biol.*, **14**, R93, Web Apollo: a web-
based genomic annotation editing platform.

Figures

Figure 1

415
Figure 1 Wikidata Data Model and Sources Schematic of the basic structure of the data
model in Wikidata. Entities include purple: Organism data sourced from NCBI's taxonomy
database (www.ncbi.nim.nih/taxonomy), light pink: gene data sourced from NCBI's Gene
Database (www.ncbi.nim.nih/gene), green: Gene product data sourced from UniProt
(www.uniprot.org), red: Gene Ontology annotations sourced from EBI's QuickGO API
(www.ebi.ac.uk/QuickGo), orange: InterPro domain annotations sourced from the InterPro
project (www.ebi.ac.uk/interpro) and yellow: Operon data currently sourced from primary
publications hosted in PubMed (www.ncbi.nim.nih.gov/pubmed). The names of the Wikidata
properties that construct the data model are included in the upper left hand corner.
425

426 **Figure2**

428
429
430 **Figure 2 Process of contributing to and consuming data in WikGenomes via Wikidata.**
431 WikiGenomes retrieves data from Wikidata via the WDQS SPARQL query service, and allows
432 contribution of defined edits back into Wikidata via a guided annotation curation process. The
433 data then becomes available to any web application that uses Wikidata in a similar way.
434

Figure 3

Figure 3 Overview of the WikiGenomes Interface. A. The 'Organism Search' form selects
the organism and populates the main page of the application with data specific to that
organism. **B.** The 'Gene Search' form loads data for a selected gene/protein. **C. Genome** The
WikiGenomes genome browser is an instance of JBrowse, a high-performance, web-based,
client-side genome browser that currently displays gene and operon tracks. **D. Organism** The
"Organism Box" displays the name of the selected organism and basic core identifiers
(hyperlinked to their respective database entries). These include the Wikidata 'QID', and the
taxonomy ID from NCBI's Taxonomy Database. This content window is where any type of
metadata about the organism will be added as it becomes available in Wikidata (e.g.,
morphology, lifestyle, Gram staining, disease associations, drugs that have action against it). **E.**
Gene/Protein The 'Gene/Protein' content window displays the gene name and basic identifiers
for the currently loaded gene and gene product in a tabbed view. The 'Gene' tab includes the
NCBI Entrez ID, the Wikidata QID, and the NCBI Locus Tag (a core identifier for bacterial
genes). The 'Protein' tab contains the UniProt ID, NCBI RefSeq Protein ID and Wikidata
QID. These are the core identifiers required for the current data model in Wikidata, but as more
and more mapped identifiers from different resources are added to Wikidata, they will be
displayed here. **F. Annotations** The annotations that have been curated in Wikidata for the
currently loaded Gene/Protein are displayed in the content window in the bottom right corner of
the application, separated and organized by navigation tabs. The tabs represent the domains of
annotations that are pulled from various resources into Wikidata for bacterial genes and
proteins. Currently, annotation types include Molecular Function, Biological Process, Cellular
Component, InterPro Domain, Operon, Enzyme Commission Number and Genomic Position.
Each annotation type displays its own relevant information consisting of hyperlinked database
identifiers, the hyperlinked Wikidata identifier, the annotation itself, and (if appropriate) genomic
coordinates of the annotation (which are also rendered in the genome browser). All annotations
are linked to references retrieved from their underlying Wikidata statements. **G. Authorize to**

Edit Button. This button redirects to WikiMedia.org login where the user can user login to their
Wikidata account and authorize WikiGenomes to make edits using their credentials.
466

467   **Figure 4**
468

470
Figure 4 Editing Forms A. Gene Ontology Form The Gene Ontology Form prompts the user
to supply 3 pieces of information: 1) the Wikidata item of the GOterm, 2) the method that was
used for determination (GO evidence code *i.e. derived from experiment, sequence similarity,*
etc...) and 3) the PMID of the publication that the statement originated from. The GO term
selection box incorporates type-ahead autocomplete allowing the user to find and select the
proper GO term to describe the annotation. Upon user submission, WikiGenomes submits the
API call to write the new annotation to Wikidata. **B**. **Operon Form** The Operon Form prompts
the user to enter the name of the operon; if the operon already exists, WikiGenomes will find it
in Wikidata and add to it. If the operon does not exist in Wikidata, the new name will be set as
the label of a new Wikidata item for that operon.   The user then provides the genes whose
expression is regulated by the operon. The input field for genes also provides type-ahead
functionality, allowing the user to quickly search for and select genes from that genome. Like all

WikiGenomes forms, the user must also provide the publication that the statement was derived
from.
485
486