***Medicago truncatula* MOT1.3 is a plasma membrane molybdenum**

transporter required for nitrogenase activity in root nodules.

Manuel Tejada-Jiménez[1,a,b], Patricia Gil-Díez[1], Javier León-Mediavilla[1], Jiangqi

Wen[2], Kirankumar S. Mysore[2], Juan Imperial[1,3], Manuel González-Guerrero[1,a]

[1]Centro de Biotecnología y Genómica de Plantas (UPM-INIA). Campus de

Montegancedo. Universidad Politécnica de Madrid. Crta. M-40 km 38. 28223 Pozuelo

de Alarcón (Madrid). Spain.

[2]Plant Biology Division, The Samuel Roberts Noble Foundation, Ardmore, Oklahoma

73401.

[3]Consejo Superior de Investigaciones Científicas. Madrid. Spain.

[a]Corresponding authors: Manuel Tejada-Jiménez (q62tejim@uco.es), phone: +34

957218591; and Manuel González-Guerrero (manuel.gonzalez@upm.es), phone: +34

918334558

[b]Present address: Department of Biochemistry and Molecular Biology. Universidad de

Córdoba. Campus de Rabanales. Edificio Severo Ochoa. Córdoba. Spain.

Total word count for the main text: 5932

Introduction word count:1033

Materials and Methods word count: 1245

Results word count: 1679

Discussion word count: 1776

Acknowledgements word count: 103

Number of Figures: 7 (6 in colour)

Number of Supporting information files: 1 Materials and Methods, 1 Figure Legends, 2

Tables, 12 Figures

Summary

• Molybdenum, as a component of the iron-molybdenum cofactor of nitrogenase,

is essential for symbiotic nitrogen fixation. This nutrient has to be provided by

the host plant through molybdate transporters.

• Members of the molybdate transporters family MOT1 were identified in the

model legume *Medicago truncatula* and their expression in nodules determined.

Yeast toxicity assays, confocal microscopy, and phenotypical characterization of

a *Tnt1* insertional mutant line were carried out in the one *M. truncatula* MOT1

family member expressed specifically in nodules.

• Among the five MOT1 members present in *M. truncatula* genome, *MtMOT1.3* is

the only one uniquely expressed in nodules. MtMOT1.3 shows molybdate

transport capabilities when expressed in yeast. Immunolocalization studies

revealed that MtMOT1.3 is located in the plasma membrane of nodule cells. A

mot1.3-1 knockout mutant showed an impaired growth concomitant with a

reduction in nitrogenase activity. This phenotype was rescued by increasing

molybdate concentrations in the nutritive solution, or upon addition of an

assimilable nitrogen source. Furthermore, *mot1.3-1* plants transformed with a

functional copy of *MtMOT1.3* showed a wild type-like phenotype.

• These data are consistent with a model in which MtMOT1.3 would be

responsible for introducing molybdate into nodule cells, which will be later used

to synthesize functional nitrogenase.

Keywords: symbiotic nitrogen fixation, molybdate transport, *Medicago truncatula*,

nitrogenase, metal transport.

Introduction

Molybdenum is one of the scarcest oligonutrients in the biosphere (Esteifel, 2002). Plants use it as part of the molybdenum cofactor (Moco) in just five enzymes involved in nitrate assimilation, purine metabolism, phytohormone production, and sulfite detoxification (Tejada-Jimenez et al., 2013). This nutrient, unlike other transition metals, is recovered from soil as the oxyanion molybdate instead of a cationic form. This determines that the transporters involved in this process are not the classical ones required for iron, copper, or zinc uptake, but members of other families. Molybdate shares some physicochemical characteristics with sulfate leading to cross-inhibition of sulfate transport by molybdate, probably due to non-specific molybdate transport through sulfate transporters in plants (Stout et al., 1951). Therefore, until recently it was thought that sulfate transporters might mediate molybdate transport in eukaryotic systems (Mendel & Hansch, 2002; Kaiser et al., 2005). The only known plant-type specific molybdate transporters belong to the Molybdate Transporter type 1 (MOT1) family and were identified in parallel in the green alga *Chlamydomonas reinhardtii* (Tejada-Jimenez et al., 2007) and in the higher plant *Arabidopsis thaliana* (Tomatsu et al., 2007). They share a high degree of homology with the sulfate transport family SULTR, but lack the conserved STAS domain (Tejada-Jimenez et al., 2007). In *C. reinhardtii*, CrMOT1 is responsible for high-affinity molybdate uptake, a process that is not severely affected by sulfate, indicating that MOT1 proteins are molybdate-specific transporters (Tejada-Jimenez et al., 2007). In Arabidopsis two members of the MOT1 family have been identified. One of them has been proposed to play a role in efficient Mo uptake from the soil (Tomatsu et al., 2007), although this function is not clear given the conflicting subcellular localizations reported for this transporter in plasma membrane or in mitochondria (Tomatsu et al., 2007; Baxter et al., 2008); while a second member of the MOT1 family is located in the vacuole of leaves and seems to be involved in intracellular and inter-organ Mo transport (Gasber et al., 2011). More recently, a MOT1 protein (LjMOT1) has been identified in *Lotus japonicus*, with a role in molybdate uptake from soil and translocation to the shoots, being located in the plasma membrane when expressed in tobacco leaves (Gao et al., 2016). In addition, another molybdate transporter family, MOT2, belonging to the major facilitator superfamily has been identified in *C. reinhardtii* (Tejada-Jimenez et al., 2011), indicating that molybdate transporters have appeared at least twice in evolution. However, its functionality as molybdate transporters has only been proved in this alga.

While all the plants employ molybdenum for Moco biosynthesis, legumes have

an additional use for it. These organisms also require molybdenum for the assembly of

the iron-molybdenum cofactor (FeMoco) of nitrogenase (Georgiadis et al., 1992; Rubio

& Ludden, 2008), the enzyme responsible for nitrogen fixation in their root nodules. In

legumes, FeMoco is assembled by diazotrophic bacteria living within differentiated root

organs, the nodules. These organs are developed in a complex process starting with the

detection of rhizobial nodulation factors (Nod) by the host plant that leads to root hair

curling, bacteria trapping, hydrolysis of the plant cell wall and bacteria delivery to the

root nodule primordium through an infection thread (Kondorosi et al., 1984; Brewin,

1991; Oldroyd, 2013). Once in the plant cytoplasm, rhizobia together with a plant-

derived membrane result in organelle-like structures, the symbiosomes, where nitrogen

fixation takes place. Two different types of nodules can be found in legumes (Sprent,

2007): determinate and inderterminate nodules. In inderterminate nodules, as those

found in *Medicago truncatula*, the continuous meristem growth results in the formation

of at least four different zones in mature nodules: the meristematic zone that drives

nodule growth; the infection/differentiation zone, where rhizobia are released through

the infection thread and differentiates to bacteroids; the fixation zone where nitrogenase

carries out its enzymatic activity; and the senescent zone where bacteroids are degraded

(Vasse et al., 1990). An additional nodule zone, the interzone, can be defined as the

transition between infection/differentiation and fixation zone (Roux et al., 2014). Once

the oxygen tension drops in the interzone, endosymbiotic bacteroids express nitrogenase

and the machinery that allows them synthesize FeMoco. Among them, is the *modABC*

operon, responsible for molybdate uptake from the peribacteroid space (Maupin-Furlow

et al., 1995; Delgado et al., 2006; Hernandez et al., 2009). Consequently, for molybdate

to reach the bacteroids, it has to cross the plasma and the symbiosome membranes, a

process that has to be mediated by transporters belonging to two different families,

given the two different directions of transport required.

In spite of the essential role that molybdenum has in nitrogenase, and the

importance of nitrogenase in legume colonization of new environments and in

sustainable agriculture, the transporters involved in molybdenum supply to the nodule

cells are not known. It can be hypothesized that molybdate transfer could in some cases

be mediated by sulfate transporters. In this case, molybdate transfer across the

symbiosome membrane could be carried out by SST1-like proteins, that have been

previously associated with sulfate delivery to symbiosomes (Krusell et al., 2005). At the

plasma membrane, another sulfate transporter, such a SHST1 homologue could mediate

molybdate uptake by rhizobia-infected cells since this transporter from the legume

Stylosanthes hamata has already been shown to transport molybdate when it is

expressed in yeast (Fitzpatrick et al., 2008). However, *in planta* SHST1 is expressed in

the root and it is involved in sulfate uptake (Smith et al., 1995), while its relationship

with plant molybdate transport has not been determined yet. A more likely alternative

would be transporters from the MOT1 or MOT2 families, since they could finely tune

molybdate delivery for symbiotic nitrogen fixation, given their high specificity for this

anion.

In this work, we have identified a *M. truncatula* member of the MOT1 family

(MtMOT1.3) involved in molybdate transport to nodule cells. *MtMOT1.3* is specifically

expressed in nodules. Its protein product is located in the plasma membrane of infected

and non-infected cells in the fixation zone of the nodule, coinciding with the zone where

MtMOT1.3 is expressed. *M. truncatula* plants lacking a functional *MtMOT1.3* gene

show a reduction of nitrogenase activity connecting MtMOT1.3 function with

molybdenum supply for FeMoCo biosynthesis. This is the first molybdate transporter

known to be specific of legume symbiotic nitrogen fixation.

152   Materials and Methods

<u>Biological material and growth conditions</u>

M. truncatula R108 seeds were scarified in the presence of concentrated sulfuric

acid for 7 min. Then, seeds were washed several times with cold water, surface

sterilized with 50 % (v/v) bleach for 90 s and incubated overnight with sterile water in

the dark. After 48 h at 4 ºC, seeds were germinated in water-agar plates 0.8 % (w/v).

Seedlings were transferred to sterile perlite pots or to Jenner's solution for hydroponic

growth (Brito et al., 1994), and inoculated with *Sinorhizobium meliloti* 2011 or the same

bacterial strain transformed pHC60 (Cheng & Walker, 1998). Plants were grown in a

greenhouse under 16 h light and 22 ºC. In the case of perlite pots, plants were watered

every two days with Jenner's solution or water alternatively. Nodules were collected 28

dpi. Plants growing under non-symbiotic conditions were supplemented every two

weeks with 2 mM KNO_3. *Agrobacterium rhizogenes* strain ARqua1 having the

appropriate vector was used for *M. truncatula* hairy-root transformation (Boisson-

Dernier et al., 2001). Agroinfiltration for transitory expression experiments were

performed in *N. benthamiana* leaves using *A. tumefaciens* C58C1 carrying the

corresponding genetic construct.

Saccharomyces cerevisiae strain 31019b (MATa *ura3 mep1Δ mep2Δ::LEU2*

mep3Δ:: KanMX2) was used for heterologous expression assays (Marini et al., 1997).

Yeasts were grown in synthetic dextrose (SD) or yeast peptone dextrose medium

supplemented with 2 % glucose (Sherman et al., 1986).

<u>Quantitative real-time RT-PCR</u>

Transcriptional expression studies were carried out by real-time RT-PCR

(StepOne plus, Applied Biosystems) using the Power SyBR Green master mix (Applied

Biosystems). Primers used are indicated in Supporting Information, Table S1. RNA

levels were normalized by using the *ubiquitin carboxy-terminal hydrolase* gene as

internal standard. RNA isolation and cDNA synthesis were carried out as previously

described (Tejada-Jimenez et al., 2015).

<u>GUS Staining</u>

pMtMOT1.3::GUS construct was obtained by amplifying 1.1 kb upstream of the

MtMOT1.3 (Nakagawa et al., 2007) start codon using the primers 5MtMOT1.3pGW and

3MtMOT1.3pGW (Supporting Information, Table S1). The resulting fragment was

cloned by Gateway cloning technology (Invitrogen) into pGWB3 vector. Hairy-root

transformation was carried out as described above. GUS staining was performed in root

of 28-dpi plants as previously described (Vernoud et al., 1999). Nodules sections were

clarified with 50% bleach for 30 min.

<u>Immunohistochemistry and confocal microscopy</u>

A genomic region comprising *MtMOT1.3* full gene and 1.1 kb upstream of its

start codon was cloned into pGWB13 vector (Nakagawa et al., 2007) using Gateway

cloning technology (Invitrogen). The resulting genetic construct contains *MtMOT1.3*

gene under the control of its own promoter and with three C-terminal HA epitopes in

frame. Hairy-root transformation was carried out as indicated above. *M. truncatula*

transformed plants were inoculated with *S. meliloti* 2011 constitutively expressing GFP.

Nodules and roots were collected at 28-dpi and fixed at 4 ºC overnight in 4 % (w/v)

paraformaldehyde and 2.5 % (w/v) sucrose in phosphate-buffered saline (PBS). Fixed

plant material was sectioned, 100 μm wide, with a Vibratome 1000 Plus. Sections were

dehydrated by serial incubation with methanol (30 %, 50 %, 70 % and 100 % [v/v] in PBS) for 5 min and then rehydrated following the same methanol series in reverse order. Cell wall permeabilization was carried out by incubation with 2 % (w/v) cellulase in PBS for 1 h and 0.1 % (v/v) Tween 20 for 15 min. Sections were blocked with 5 % (w/v) bovine serum albumin in PBS and then incubated with 1:50 anti-HA mouse monoclonal antibody (Sigma) in PBS at room temperature for 2 h. Primary antibody was washed three times with PBS for 10 min and subsequently incubated with 1:40 Alexa594-conjugated anti-mouse rabbit monoclonal antibody (Sigma) in PBS at room temperature for 1 h. Secondary antibody was washed three times with PBS for 10 min, and then DNA was stained using DAPI. Images were obtained with a confocal laser-scanning microscope (Leica SP8).

Transient expression in *Nicotiana benthamiana* leaves

MtMOT1.3 transient expression in tobacco leaves was carried out as previously described (Voinnet et al., 2003). *MtMOT1.3* CDS was cloned into pGWB5 (Nakagawa et al., 2007) by Gateway cloning technology (Invitrogen), resulting in C-terminal fusion to GFP. *A. tumefaciens* C58C1 (Deblaere et al., 1985) cells independently transformed with this construct, with the plasma membrane marker pm-CFP pBIN (Nelson et al., 2007) or with the silencing suppressor p19 of *Tomato bushy stunt virus* (Voinnet et al., 2003) were injected into 4-week-old *N. benthamiana* leaves. Expression of the appropriate construct was analyzed after 3 d by confocal laser-scanning microscopy (Leica).

Nitrogenase activity

Acetylene reduction assay was used to measure nitrogenase activity (Hardy et al., 1968). Wild-type and mutant plants 28 dpi were separately introduced in 30-mL tubes. Each tube was sealed with rubber stoppers and contained at least four independently transformed plants. Three mililiters of air of each bottle was replaced by the same volume of acetylene, tubes were subsequently incubated for 30 min at room temperature. Produced ethylene was measured by analyzing 0.5 mL of gas from each bottle in a Shimadzu GC-8A gas chromatograph using a Porapak N column.

Metal content determination

Inductively coupled plasma optical emission spectrometry was carried out at the

Unit of Metal Analysis in the Scientific and Technology Centers of the Universidad de

Barcelona (Spain).

<u>Nitrate reductase activity</u>

Nitrate reductase activity was analyzed as described by Scheible et al. (1997)

with some modifications. We started from 100 mg of fresh leaves. The extraction buffer

used for crude extracts preparation contained 100 mM potassium phosphate pH 7.5, 5

mM magnesium acetate, 10 % glycerol (v/v), 10 % polyvinylpolypyrrolidone (w/v), 0.1

% Triton X-100, 1 mM EDTA, 0.05 % ß-mercaptoethanol, 1 mM PMSF. Plant material

was homogenized with liquid nitrogen and 1:6 extraction buffer (v/v). The crude extract

was centrifuged at 14,000 xg at 4 ºC for 15 min. The reaction was initiated by adding

0.05 mL of crude extract to 0.5 mL of reaction buffer and incubated at 30 ºC for 20 min.

The reaction buffer contained 50 mM potassium phosphate pH 7.5, 10 mM KNO_3, 5

mM EDTA and 0.5 mM NADH. Nitrate reduction reaction was stopped by adding 1

volume of 1 % sulfanilamide in 2.4 M HCl, and 1 volume of 0.02 % N-1-naphtyl-

ethylenediamine. After centrifugation absorbance at 540 nm was measured in

UV/visible spectrophotometer Ultrospect 3300 pro (Amersham Bioscience).

<u>Bioinformatics</u>

Members of the MOT1 family in *M. truncatula* were identified by the Basic

Local Alignment Search Tool in the *M. truncatula* Genome Project database

(http://www.jcvi.org/medicago/index.php), using as reference sequence a member of the

MOT1 family from Arabidopsis (NP_180139). Protein sequence alignment and

unrooted tree visualization were performed with MEGA7 package

(http://www.megasoftware.net/) using ClustalW software and neighbor joining

algorithm. Accession numbers: *Arabidopsis thaliana* AtMOT1.1, NP_180139.1;

AtMOT1.2, NP_178147.1; AtSULTR1.1, NP_192602.1; AtSULTR2.1, NP_196580.1;

AtSULTR3.1, ANM64117.1; AtSULTR4.1, AED91910.1; *Brassica napus* BnMOT1.1,

XP_013709878.1; BnMOT1.3, XP_013667816.1; BnSULTR1.1, XP_013715602.1;

BnSULTR2.1, NP_001302517.1; BnSULTR3.1, XP_013737126.1; BnSULTR4.1,

XP_013667920.1; *Glycine max* GmMOT1.1, XP_003545516.1; GmMOT1.4,

KRH05029.1; GmMOT1.6, XP_003527804.1; GmMOT1.7, XP_003523708.1;

GmSULTR1.3, XP_006593569.1; GmSULTR2.1, XP_003531538.1; GmSULTR3.1,

XP_003521258.1; GmSULTR4.2, XP_003552670.1; *Lotus japonicus* LjMOT1.1,

AFK43331.1; LjMOT1.2, AJE26312.1; LjSST1, CAL36108.1; *Medicago truncatula*

MtMOT1.1, XP_013465770.1; MtMOT1.2, XP_013465776.1; MtMOT1.3,

XP_013460259.1; MtMOT1.4, XP_013454709.1; MtMOT1.5, XP_003603486.1;

MtSULTR1.1, XP_003614968.1; MtSAT1, XP_003602002.1; *Oryza sativa*

OsMOT1.1, XP_015650610.1; OsMOT1.2, XP_015621613.1; OsSULTR1.2

XP_015650733.1; OsSULTR2.1, ABF94445.1; OsSULTR3.1, BAS82415.1;

OsSULTR4.1, XP_015612472.1; *Phaseolus vulgaris* PvMOT1.1, XP_007161270.1;

PvMOT1.2, XP_007137089.1; PvMOT1.3, XP_007161272.1; PvMOT1.4,

XP_007135722.1; PvSULTR1.1, XP_007141140.1; PvSULTR2.1, XP_007163633.1;

PvSULTR3.1, XP_007162459.1; PvSULTR4.1, XP_007139276.1; *Vitis vinifera*

VvMOT1.1, XP_002281989.2; VvMOT1.2, XP_002285217.1; VvSULTR1.1,

XP_010664070.1; VvSULTR2.1, XP_010652824.1; VvSULTR3.1, XP_003632327.1;

VvSULTR4.2, XP_002282491.2.

Statistical tests

Data were analyzed by Student's unpaired t test to calculate statistical

significance of observed differences. Test results with p-values less than 0.05 were

considered as statistically significant.

Results

MtMoT1.3 belongs to the molybdate transporter MOT1 family and is specifically

expressed in nodules.

Due to the possible role of MOT1 transporters in Mo supply for symbiotic

nitrogen fixation, we carried out a search for MOT1 members in the *M. truncatula*

genome. Five putative *M. truncatula* proteins were found (encoded by *Medtr1g010210,*

Medtr1g010270, Medtr3g464210, Medtr4g011600 and *Medtr3g108190*) showing a

sequence identity between 51.4 % and 65.7 % with an Arabidopsis member of the

MOT1 family. These proteins were annotated as sulfate transporter-like proteins;

however, sequence comparison showed that they cluster with plant MOT1 proteins and

are more distant to sulfate transporters (SULTR) (Fig. 1a). Additionally, these proteins

also contain the two sequence motifs conserved in all MOT1 proteins (Tejada-Jimenez

et al., 2007) (Fig. 1b). Therefore, we named these proteins MtMOT1.1 to MtMOT1.5,

respectively. The number of MOT1 members present in *M. truncatula* contrasts with the

situation reported for Arabidopsis where only two MOT1 transporters are present

(Tomatsu et al., 2007; Baxter et al., 2008; Gasber et al., 2011). Analysis of the number

of MOT1 members in already sequenced plants showed that these proteins are present,

in average, in a higher number in legumes than in non-legumes plants (Supporting

Information, Table S2). This finding suggests a particular role of MOT1 proteins in

legumes, where an additional molybdenum sink is present: diazotrophic bacteria in root

nodules.

In order to study the possible relationship of the five putative MOT1 members of

M. truncatula with symbiotic nitrogen fixation, a transcriptional analysis of these genes

was carried out by real-time RT-PCR in plants during symbiotic association with *S.*

meliloti or in uninoculated plants fertilized with nitrogen. We found that *MtMOT1.3*

expression is restricted to the nodule (Fig. 1c and 1d), while transcripts from the other

four *MtMOT1* genes were also detected at varying levels in other organs of the analyzed

plants (Supporting Information, Fig. S1). These results strongly suggest an important

role of MtMOT1.3 in the nodule, likely in molybdate transport connected to symbiotic

nitrogen fixation. *MtMOT1.3* expression levels were not significantly affected by

molybdate concentration in the nutritive solution (Supporting Information, Fig. S2).

MtMOT1.3 is a molybdate transporter

All the MOT1 proteins reported so far mediate molybdate transport, in form of

the oxyanion molybdate, to the cytosol (Tejada-Jimenez et al., 2013). The high

sequence similarity that MtMOT1.3 shares with the already characterized molybdate

transporters suggests that this protein could also mediate molybdate transport in the

same direction. To test this hypothesis, *MtMOT1.3* was heterologously expressed in the

yeast *S. cerevisiae*. This yeast is a good model to study of Mo transporters since it is one

of the few organisms that do not use molybdenum, excluding any effect of endogenous

specific molybdate transport activity (Mendel & Bittner, 2006). Toxicity studies showed

that yeasts expressing *MtMOT1.3* exhibit a defective growth in synthetic dextrose (SD)

solid medium in the presence of 50 μM molybdate, compared with yeast transformed

with the empty vector pDR196, but no growth differences were observed with the

control when no additional molybdate was added to the growth medium (Fig. 2a).

Similar results were obtained in SD liquid medium by monitoring yeast growth along

the time (Fig. 2b; Supporting Information, Fig. S3). The toxic effect observed in yeast is

the likely result of molybdenum being transported and accumulated in the cell as a

result of MtMOT1.3 activity (Fig. 2c). This result supports the functionality of
MtMOT1.3 as molybdate transporter towards the cytosol.

MtMOT1.3 is expressed in the plasma membrane of nodule cells of the interzone and
fixation zones

To investigate the role of MtMOT1.3 in molybdenum supply to the nodule, we
studied the tissue specific localization of _MtMOT1.3_ expression. This was assessed by
analyzing the expression of the ß-glucoronidase (_gus_) gene under the control of 1.1 kb
of genomic DNA directly upstream of the _MtMOT1.3_ starting codon, that was selected
as promoter. Using this genetic construct, GUS activity was found in the nodule
interzone, with the maximum in the area corresponding with the fixation zone (Fig. 3a
and 3b). This expression distribution within the nodule matches with the transcription
data available in the Symbimics database obtained by means of laser-capture
microdissection coupled to RNA sequencing (Roux et al., 2014), where _MtMOT1.3_
expression is mainly detected in the interzone and fixation zones (Fig. 3c).
Localization of MtMOT1.3 protein was also analyzed using an
immunohistochemical and confocal microscopy approach. _M. truncatula_ plants were
transformed with a genetic construct comprising the genomic region of _MtMOT1.3_
fused in frame with three hemagglutinin (HA) epitopes in C-terminus (MtMOT1.3-HA),
under the control of the same promoter used for the GUS assay. The chimeric
MtMOT1.3-HA protein was detected (in the red channel) with a mouse anti-HA
antibody and an Alexa594-conjugated anti-mouse secondary antibody. Its position
within the nodule was traced by DNA staining using 4',6-diamino-phenylindole (DAPI)
(blue) and a _S. meliloti_ strain constitutively expressing GFP (green). In agreement with
the GUS activity data, MtMOT1.3-HA was detected in the interzone and fixation zones
of the nodule (Fig. 4a to 4c). Particularly, MtMOT1.3-HA signal was found in the
periphery of infected and non-infected cells (Fig. 4d to 4f), fitting with plasma
membrane localization. This signal was not the result of autofluorescence, since
sections obtained from the same biological material, subjected to the same preparation
protocol with the exception of the incubation with the primary anti-HA antibody, did
not show any fluorescence in the Alexa594 emission spectrum (Supporting Information,
Fig. S4). Moreover, the specific peripheral distribution of MtMOT1.3-HA was also
observed using an Alexa488-conjugated secondary antibody in nodules containing m-
Cherry-expressing _S. meliloti_ (Supporting Information, Fig. S5). No MtMOT1.3-HA

signal was detected by Western blot analyses in roots from nodulated plants (Supporting

Information, Fig. S6), consistent with the nodule-specifc expression data obtained by

real-time PCR. To confirm the subcellular localization of MtMOT1.3 in the plasma

membrane, its coding sequence fused to the green fluorescent protein (GFP) in C-

terminus was transiently expressed in *N. benthamiana* leaves together with a plasma

membrane marker fused to cyan fluorescent protein (CFP). MtMOT1.3 signal and

plasma membrane marker signal co-localized in *N. benthamiana* leaves cells expressing

both genetic constructs (Fig. 4g to 4i), supporting the localization of MtMOT1.3 in the

plasma membrane. However, no GFP or CFP signals were found in cells expressing

only the plasma membrane marker or MtMOT1.3, respectively (Supporting

Information, Fig. S7), thus ruling out any non-specific signal when both constructs are

co-expressed.

<u>Lack of *MtMOT1.3* leads to a reduction of nitrogenase activity</u>

In order to investigate the role of MtMOT1.3 in symbiotic nitrogen fixation, the

mutant line NF10801 (*mot1.3-1*) was identified by a reverse genetics screening (Cheng

et al., 2011; Cheng et al., 2014) from a *Transposable Element from N. tabacum* (*Tnt1*)

insertion mutant library (Tadege et al., 2008). The mutant *mot1.3-1* carries a *Tnt1*

insertion in the second exon of the *MtMOT1.3* gene at position +1782 from the starting

codon (Fig. 5a). *MtMOT1.3* transcripts were not detected in the homozygous *mot1.3-1*

mutant (Fig. 5b), therefore MtMOT1.3 activity is not present in this mutant line.

The phenotype of *mot1.3-1* was evaluated in symbiotic conditions, with nitrogen

fixation as the sole nitrogen source, and under low Mo availability. In these conditions,

mot1.3-1 mutant showed a reduced growth rate compared with wild-type plants (Fig.

5c). Consequently, plant biomass was reduced in the *mot1.3-1* mutant by 70 % in shoots

and 55 % in roots (Fig. 5d). Nitrogenase activity was measured in *mot1.3-1* and wild-

type plants by the acetylene-reduction assay (Dilworth, 1966; Schöllhorn & Burris,

1966). *mot1.3-1* plants showed a reduction of 90 % in nitrogenase activity, as compared

to the activity measured in wild-type plants (Fig. 5e). On a per-nodule basis, *mot1.3-1*

plants had in average 24 % of the nitrogenase activity of the wild-type ones (Supporting

Information, Fig. S8). Nodulation was not affected in the *mot1.3-1* mutant, since

nodules were comparable to wild-type plants in terms of number and shape (Fig. 5f and

5g). Moreover, no differences in nodule anatomy, or nodulation kinetics were observed

between mutant and control plants (Supporting Information, Fig. S9). However, these

nodules were on average smaller than those from wild type plants (Fig. 5g; Supporting

Information, Fig. S10). In addition, nodule molybdenum content in *mot1.3-1* plants was

significantly higher than the control (Fig. 5h), indicative of a role on molybdenum

homeostasis in this organ. In order to investigate whether the phenotype of *mot1.3-1*

plants is caused by a shortage in molybdenum supply, these plants were watered with a

nutrient solution containing 5 µg/L ammonium heptamolybdate. Under molybdenum

sufficient conditions, *mot1.3-1* mutants exhibited a growth rate, biomass and

nitrogenase activity comparable to wild-type plants (Supporting Information, Fig. S10).

A similar result was obtained in hydroponics, a growth condition where molybdenum

concentrations can be better controlled (Supporting Information, Fig. S11).

Interestingly, no MOT1 family member was more highly expressed in *mot1.3-1* nodules

than in wild-type plants (Supporting Information, Fig. S12).

To validate that the phenotype observed was the result of the *Tnt1* insertion in

MtMOT1.3, mot1.3-1 plants were transformed with *MtMOT1.3-HA* construct under its

own promoter (same genetic construct used for MtMOT1.3 immunolocalization assay).

Insertion of the mutated gene in the *mot1.3-1* mutant under low molybdenum conditions

also restored wild-type growth (Fig. 5a to 5e), supporting that mutation of *MtMOT1.3* is

responsible for this abnormal phenotype. In addition, it also validates the

immunofluorescence data, since without a wild type-like expression profile and

localization, no complementation would have been observed.

To check whether the lack of MtMOT1.3 activity could affect other processes in

the plant, the phenotype of *mot1.3-1* mutant was assayed under non-symbiotic

conditions, with nitrate as nitrogen source. In this situation, plant growth relies on the

activity of the molybdo-enzyme nitrate reductase to reduce nitrate to nitrite that will be

subsequently converted to ammonia and incorporated to amino acids by the glutamine

synthetase/glutamine synthase pathway (Bernard & Habash, 2009). In these conditions

mot1.3-1 plants showed a phenotype similar to wild-type plants in terms of growth rate,

biomass and nitrate reductase activity, regardless of the Mo availability (Fig. 6).

Therefore, the function of MtMOT1.3 seems to be restricted to symbiotic conditions

when symbiotic nitrogen fixation has an important role in plant growth.

Discussion

Obtaining nitrogen from the atmosphere by biological nitrogen fixation is a

sustainable alternative to the intensive use of synthetic nitrogen fertilizers in agriculture

(Smil, 1999). Legumes are able to use atmospheric N_2 as nitrogen source by means of a

symbiotic association with nitrogen-fixing bacteria from the soil. They occupy 12-15%

of the world arable land (Graham & Vance, 2003). However, maintaining an adequate

nitrogen fixation rate involves a complex biological process that requires an efficient

nutrient supply from the plant to the rhizobia (Udvardi & Poole, 2013). Molybdenum

supply is critical for symbiotic nitrogen fixation since this oligonutrient is needed to

synthesize the enzyme nitrogenase, directly involved in the reduction of N_2 to produce

NH_4^+. In plants molybdenum is also present in other enzymes carrying out important

metabolic processes such as nitrate assimilation (nitrate reductase) or phytohormone

biosynthesis (aldehyde oxidase) (Tejada-Jimenez et al., 2013). Little is known about

how plants take up molybdate from the soil and redistribute it to the sink organs,

although members of the molybdate transport protein family MOT1 are very likely

involved in this process in Arabidopsis (Tomatsu et al., 2007; Baxter et al., 2008;

Gasber et al., 2011). However, their physiological role in molybdenum homeostasis is

still unclear. Less is known for molybdate nutrition in legumes, in spite of the

importance of molybdenum in symbiotic nitrogen fixation. Considering the renewed

interest and recent advances towards introducing nitrogen fixing abilities into non-

legumes (Charpentier & Oldroyd, 2010; Curatti & Rubio, 2014; Lopez-Torrejon et al.,

2016) and the need for a more sustainable agriculture (Foley et al., 2011), this gap of

our knowledge needs to be filled if we are to ensure proper delivery of molybdenum to

produce functional nitrogenase in new biological systems. The present work represents

a first but decisive step toward optimization of molybdate allocation for nitrogen

fixation, where we have identified a nodule-specific MOT1 member in *M. truncatula*

(MtMOT1.3) as responsible for molybdenum supply for symbiotic nitrogen fixation.

Legumes appear to have expanded the number of genes encoding MOT1

proteins in their genome in order to adapt to the presence of an additional molybdenum

sink, the root nodule, doubling the average copy number in monocots and other dicots.

This is in contrast to genes encoding Nramp transporters, some of which play an

essential role in iron transport in rhizobia-infected cells (Kaiser et al., 2003; Tejada-

Jimenez et al., 2015), where no significant increase in gene copy numbers were

observed. One possible explanation for this, based on the fact that few plant proteins use

molybdenum-based cofactors, is that MOT1 transporters are only expressed in very few

cell types, highly specialized in the physiological processes catalyzed by Moco or

FeMoco-dependent enzymes. If this hypothesis is right, nodule cells should express a

specific MOT1 transporter responsible for ensuring molybdenum supply for symbiotic

nitrogen fixation, as is the case for *MtMOT1.3* in *M. truncatula.*

MtMOT1.3 is expressed in the nodule interzone and in the early fixation zone.

This transcription pattern is also validated by the transcriptomic data obtained from the

Symbimics database (Roux et al., 2014). This expression profile is consistent with a

situation in which the rhizobia-infected cell is increasing its molybdenum content to

transfer it to the bacteroids, so that when the physiological conditions are right, they can

start synthesizing FeMoco. In contrast, iron-transporting *MtNramp1*, responsible for

iron uptake by rhizobia-infected cells, shows a different expression profile, with a

maximum of expression in the infection/differentiation zone (Tejada-Jimenez et al.,

2015). This difference for the expression of iron and molybdenum transporter genes

indicates that the uptake of these two elements occurs in two separate moments during

nodule development, probably reflecting an earlier need for certain ferroproteins other

than nitrogenase. It also indicates that molybdate release from the vasculature for uptake

by rhizobia-infected cells occurs at the interzone and early fixation zone, in contrast to

what has been proposed for iron and other transition metals (Rodriguez-Haas et al.,

2013; González-Guerrero et al., 2014; González-Guerrero et al., 2016). However,

neither *MtMOT1.3* nor *MtNramp1* are expressed in the older parts of the fixation zone.

This suggests that no additional metal uptake is taking place in this zone, indicating

either a high degree of protein stability in the fixation zone, or an effective recycling of

essential metallic nutrients.

MtMOT1.3 is a plasma membrane-bound protein, as indicated by both the

immunolocalization of a HA-tagged fusion in *M. truncatula* nodules and the

colocalization studies of MtMOT1-GFP with a plasma membrane marker in tobacco

leaves. In Arabidopsis, while AtMOT1.2 is clearly localized in the vacuole of leaf cells

(Gasber et al., 2011), contradicting localization data have been reported for AtMOT1.1,

claiming either plasma membrane or mitochondrial localizations (Tomatsu et al., 2007;

Baxter et al., 2008). These conflicting data seem to be caused by the GFP marker fused

to AtMOT1.1, since when it is fused to the N-terminus of AtMOT1.1 it leads to plasma

membrane localization, while the C-terminus fusion results in mitochondrial

localization. Our MtMOT1.3 subcellular localization data have been obtained using two

different methodologies: immunolocalization in *M. truncatula* cells by detecting HA

epitope fused to MtMOT1.3, and transient expression in *N. benthamiana* fused to GFP.

Both approaches yielded the same result reinforcing MtMOT1.3 localization in the

plasma membrane. The observed immunofluorescence data are not the result of any autofluorescence, since similar fluorescence profiles were observed using two different fluorescent probes. Moreover, when no probe was added to the sections, no fluorescence was detected in them, indicating that the signal was the result of the antibody binding. There is no evidence of unspecific recognition of any epitope other than the HA-tag in the Western blot analysis carried out using an anti-HA antibody. Furthermore, previous data showed that the Alexa594-conjugated antibody used was specific for HA-expressing proteins (Tejada-Jiménez et al., 2015). Finally, expressing *MtMOT1.3-HA* in a *mot1.3-1* mutant background resulted in complementation and recovery of the wild-type phenotype, which indicates that the tagged transporter is being targeted to the proper subcellular compartment.

Like MOT1 family members, MtMOT1.3 transports molybdate towards the cytosol, as indicated by the yeast toxicity assays. The phenotype characterization of a *Tnt1* mutant line strongly suggests that this activity is used by *M. truncatula* to introduce molybdate into nodule cells. The knockout mutant line *mot1.3-1* has lower nitrogenase activity than wild-type control plants, both when whole plants or nodules were compared. Consequently, plant growth is reduced due to lack of fixed nitrogen. Nodules from *mot1.3-1* are on average smaller than those from wild-type plants. This is not caused by a delayed nodulation, since nodulation kinetics are very similar to wild-type plants. Neither it is due to a defect of the nodulation process, as the nodule organization does not show any alteration in *mot1.3-1* nodules. Finally, the observed reduction in nodule size (around 75% of wild type) cannot account for the observed reduction in nodule nitrogenase activity (ca 25% of wild-type leves). Alternatively, it could be hypothesized that the reduction in nodule size could be caused by the reduced uptake of a nutrient essential for nodule functioning. The *mot1.3-1* phenotype is molybdenum-dependent, since molybdate fortification of the nutritive solution resulted in wild-type looking *mot1.3-1* plants, in apparent contradiction to the higher levels of molybdenum detected in *mot1.3-1* nodules. This accumulation pattern has also been observed when studying a nodule-specific plasma membrane copper transporter (unpublished data). One possible explanation is the existence of a signal indicating intracellular metal deficiency that would trigger more metal being transported to the nodule. Since the uptake transporter is not present, this would result in metal accumulation in the nodule apoplast and in the vasculature. In any case, the molybdate transport capabilities shown in yeast, the alteration of molybdate homeostasis in

nodules, the reduction of a molybdate-dependent enzymatic activity in nodules, its
plasma membrane localization, and the restoration of a wild-type phenotype by
increasing molybdate concentrations in the nutritive solution, point out to a role of
MtMOT1.3 in molybdate uptake by nodule cells.

The phenotype reversal by increasing the molybdate content of the nutritive
solution is the result of other membrane transporters that at higher molybdate
concentrations could counterbalance the absence of MtMOT1.3 activity. In *M.*
truncatula all the MOT1 proteins are expressed within the nodule, although only
MtMOT1.3 exhibited a nodule-specific transcription pattern. It could be argued that
another member of the MOT1 family could carry out this role. If this were the case, an
induction of its expression levels in *mot1.3-1* nodules would be expected. However, we
observe a slight reduction of their expression levels, suggesting a different role in
molybdenum homeostasis in nodules. Alternatively, sulfate transporters could also
compensate the lack of MtMOT1.3; for instance, the sulfate transporter SHST1 from the
legume *Stylosanthes hamata* is able to transport molybdate upon expression in *S.*
cerevisiae (Fitzpatrick et al., 2008). Furthermore, proteins of the MOT2 family have
been also related to molybdate transport in the green alga *C. reinhardtii* (Tejada-
Jimenez et al., 2011) and members of the MOT2 family are present in *M. truncatula*
genome. However according to the database Symbimics, none of SHST1 or MOT2 *M.*
truncatula orthologues are significantly expressed in nodules (Roux et al., 2014), which
indicates that either another MOT1 transporter carried out this role in spite of the
reduced expression levels, or a member of a yet-to-be-identified molybdate transporter
family would be supplementing MtMOT1.3 function.

Under non-symbiotic conditions, MtMOT1.3 does not play an essential role, as
indicated by its expression profile and by the lack of phenotype observed under these
conditions. Even when the plants require Moco-dependent nitrate reductase activity to
grow, no differences were observed between mutant and wild-type plants. These results
reinforce the hypothesis of specialization of a legume MOT1 transporter during
evolution to provide molybdenum for symbiotic nitrogen fixation.

In conclusion, molybdate transported by the host plant is released from the
vasculature into the apoplast of the interzone/early fixation zone (Fig. 7). From there,
MtMOT1.3 introduces molybdate into rhizobia-infected cells, as well as in the non-
infected ones within the nodule. The latter raises the question of why the non-infected
cells would require molybdenum. Possible explanations would be that they are being

prepared if they become infected, that they are buffering or storing molybdenum to be
used later, or that this molybdenum is required to synthesize active Moco-dependent
enzymes in these cells. Once in the cell cytosol, molybdate has to be transported across
the symbiosome membrane. Since no molybdate-specific efflux system is known, it
could be speculated that this role is carried out by a sulfate transporter specific of this
membrane such as SST1, whose mutation has a severe impact on nitrogenase activity
(Krusell et al., 2005). Once in the peribacteroid space, ModABC would introduce
molybdate into the bacteroid (Delgado et al., 2006; Cheng et al., 2016).

Acknowledgments
This research was funded by a European Research Council Starting Grant (ERC-2013-
StG-335284), to MGG. Development of *M. truncatula Tnt1* mutant population was, in
part, funded by the National Science Foundation, USA (DBI-0703285) to KSM. We
would like to thank Dr. T. Coque and Dr. A. SanMillán (Ramón y Cajal Health
Research Institute) for allowing us to use their Bioscreen equipment; and Dr. J. M.
Argüello for sending us the mCherry-expressing *S. meliloti*. We would also like to
acknowledge the other members of Laboratory 281 at Centro de Biotecnología y
Genómica de Plantas (UPM-INIA) for their support and feedback in preparing this
manuscript.

Author contribution
MTJ carried out most of the experiments. PGD performed the confocal microscopy
using the anti-HA Alexa488-conjugated antibody, the nodule development cross-
section, and the molybdenum content in nodules. JLM carried out the expression of
MOT1 genes in the *mot1.3-1* mutant, as well as studied the effecto of molybdate
concentrations in transcription levels. Both PGD and JLM studied the nodulation
process in wild type and *mot1.3-1* plants over time. JW and KSM performed *Medicago*
truncatula mutant screening and isolated the *mot1.3-1* allele. MTJ, JI, and MGG
designed the experiments, analyzed the data and wrote the article.

References

Baxter I, Muthukumar B, Park HC, Buchner P, Lahner B, Danku J, Zhao K, Lee

J, Hawkesford MJ, Guerinot ML, Salt DE. 2008. Variation in molybdenum content

across broadly distributed populations of *Arabidopsis thaliana* is controlled by a

mitochondrial molybdenum transporter (MOT1). *PLoS Genetics* **4**: e1000004.

Bernard SM, Habash DZ. 2009. The importance of cytosolic glutamine synthetase in

nitrogen assimilation and recycling. *New Phytologist* **182**: 608-620.

Boisson-Dernier A, Chabaud M, Garcia F, Becard G, Rosenberg C, Barker DG.

2001. *Agrobacterium rhizogenes*-transformed roots of *Medicago truncatula* for the

study of nitrogen-fixing and endomycorrhizal symbiotic associations. *Molecular Plant-*

Microbe Interactions **14**: 695-700.

Brewin NJ. 1991. Development of the legume root nodule. *Annual Review of Cell*

Bioloy **7**: 191-226.

Brito B, Palacios JM, Hidalgo E, Imperial J, Ruiz-Argueso T. 1994. Nickel

availability to pea (*Pisum sativum* L.) plants limits hydrogenase activity of *Rhizobium*

leguminosarum bv. *viciae* bacteroids by affecting the processing of the hydrogenase

structural subunits. *Journal of Bacteriology* **176**: 5297-5303.

Charpentier M, Oldroyd G. 2010. How close are we to nitrogen-fixing cereals?

Current Opinion in Plant Biology **13**: 556-564.

Cheng G, Karunakaran R, East AK, Poole PS. 2016. Multiplicity of sulfate and

molybdate transporters and their role in nitrogen fixation in *Rhizobium leguminosarum*

bv. *viciae* Rlv3841. *Molecular Plant- Microbe Interactions* **29**: 143-152.

Cheng HP, Walker GC. 1998. Succinoglycan is required for initiation and elongation

of infection threads during nodulation of alfalfa by *Rhizobium meliloti. Journal of*

Bacteriology **180**: 5183-5191.

Cheng X, Wen J, Tadege M, Ratet P, Mysore KS. 2011. Reverse genetics in

Medicago truncatula using *Tnt1 i*nsertion mutants. *Methods in Molecular Biology* **678**:

179-190.

Cheng X, Wang M, Lee HK, Tadege M, Ratet P, Udvardi M, Mysore KS, Wen J.

2014. An efficient reverse genetics platform in the model legume *Medicago truncatula*.

New Phytologist **201**: 1065-1076.

Curatti L, Rubio LM. 2014. Challenges to develop nitrogen-fixing cereals by direct

nif-gene transfer. *Plant Science* **225**: 130-137.

Deblaere R, Bytebier B, De Greve H, Deboeck F, Schell J, Van Montagu M,
Leemans J. 1985. Efficient octopine Ti plasmid-derived vectors for Agrobacterium-
mediated gene transfer to plants. *Nucleic Acids Research* **13**: 4777-4788.

Delgado MJ, Tresierra-Ayala A, Talbi C, Bedmar EJ. 2006. Functional
characterization of the *Bradyrhizobium japonicum modA* and *modB* genes involved in
molybdenum transport. *Microbiology* **152**: 199-207.

Dilworth MJ. 1966. Acetylene reduction by nitrogen-fixing preparations from
Clostridium pasteurianum. Biochimica et Biophysica Acta **127**: 285-294.

Esteifel EI. 2002. Molybdenum and tungsten: their roles in biological processes. In:
Sigel A, Sigel H, eds. *Metal ions in biological systems* New York: Marcel Dekker Inc.,
1-30.

Fitzpatrick KL, Tyerman SD, Kaiser BN. 2008. Molybdate transport through the
plant sulfate transporter SHST1. *FEBS Letters* **582**: 1508-1513.

Foley JA, Ramankutty N, Brauman KA, Cassidy ES, Gerber JS, Johnston M,
Mueller ND, O'Connell C, Ray DK, West PC, et al. 2011. Solutions for a cultivated
planet. *Nature* **478**: 337-342.

Gao JS, Wu FF, Shen SL, Meng Y, Cai YP, Lin Y. 2016. A putative molybdate
transporter LjMOT1 is required for molybdenum transport in *Lotus japonicus*.
Physiologia Plantarum. **158**: 331-340.

Gasber A, Klaumann S, Trentmann O, Trampczynska A, Clemens S, Schneider S,
Sauer N, Feifer I, Bittner F, Mendel RR, et al. 2011. Identification of an Arabidopsis
solute carrier critical for intracellular transport and inter-organ allocation of molybdate.
Plant Biology **13**: 710-718.

Georgiadis MM, Komiya H, Chakrabarti P, Woo D, Kornuc JJ, Rees DC. 1992.
Crystallographic structure of the nitrogenase iron protein from *Azotobacter vinelandii*.
Science **257**: 1653-1659.

González-Guerrero M, Matthiadis A, Saez A, Long TA. 2014. Fixating on metals:
new insights into the role of metals in nodulation and symbiotic nitrogen fixation.
Frontiers in Plant Science **5**: 45.

González-Guerrero M, Escudero V, Saez A, Tejada-Jimenez M. 2016. Transition
metal transport in plants and associated endosymbionts: Arbuscular mycorrhizal fungi
and rhizobia. *Frontiers in Plant Science* **7**: 1088.

Graham PH, Vance CP. 2003. Legumes: importance and constraints to greater use.
Plant Physiology **131**: 872-877.

Hardy RW, Holsten RD, Jackson EK, Burns RC. 1968. The acetylene-ethylene assay
for n(2) fixation: laboratory and field evaluation. *Plant Physiology* **43**: 1185-1207.

Hernandez JA, George SJ, Rubio LM. 2009. Molybdenum trafficking for nitrogen
fixation. *Biochemistry* **48**: 9711-9721.

Kaiser BN, Gridley KL, Ngaire Brady J, Phillips T, Tyerman SD. 2005. The role of
molybdenum in agricultural plant production. *Annals of Botany* **96**: 745-754.

Kaiser BN, Moreau S, Castelli J, Thomson R, Lambert A, Bogliolo S, Puppo A,
Day DA. 2003. The soybean NRAMP homologue, GmDMT1, is a symbiotic divalent
metal transporter capable of ferrous iron transport. *Plant Journal* **35**: 295-304.

Kondorosi E, Banfalvi Z, Kondorosi A. 1984. Physical and genetic analysis of a
symbiotic region of *Rhizobium meliloti*: identification of nodulation genes. *Molecular*
General Genetics **193**: 445-452.

Krusell L, Krause K, Ott T, Desbrosses G, Kramer U, Sato S, Nakamura Y,
Tabata S, James EK, Sandal N, et al. 2005. The sulfate transporter SST1 is crucial for
symbiotic nitrogen fixation in *Lotus japonicus* root nodules. *Plant Cell* **17,** 1625-1636.

Lopez-Torrejon G, Jimenez-Vicente E, Buesa JM, Hernandez JA, Verma HK,
Rubio LM. 2016. Expression of a functional oxygen-labile nitrogenase component in
the mitochondrial matrix of aerobically grown yeast. *Nature Communications* **7**: 11426.

Marini AM, Soussi-Boudekou S, Vissers S, Andre B. 1997. A family of ammonium
transporters in *Saccharomyces cerevisiae. Molecular and Cellular Biology* **17**: 4282-
4293.

Maupin-Furlow JA, Rosentel JK, Lee JH, Deppenmeier U, Gunsalus RP,
Shanmugam KT. 1995. Genetic analysis of the *modABCD* (molybdate transport)
operon of *Escherichia coli. Journal of Bacteriology* **177**: 4851-4856.

Mendel RR, Hansch R. 2002. Molybdoenzymes and molybdenum cofactor in plants.
Journal of Experimental Botany **53**: 1689-1698.

Mendel RR, Bittner F. 2006. Cell biology of molybdenum. *Biochimimica et*
Biophysica Acta **1763**: 621-635.

Nakagawa T, Kurose T, Hino T, Tanaka K, Kawamukai M, Niwa Y, Toyooka K,
Matsuoka K, Jinbo T, Kimura T. 2007. Development of series of gateway binary
vectors, pGWBs, for realizing efficient construction of fusion genes for plant
transformation. *Journal of Bioscience and Bioengineering* **104**: 34-41.

Nelson BK, Cai X, Nebenfuhr A. 2007. A multicolored set of in vivo organelle
markers for co-localization studies in Arabidopsis and other plants. *Plant Journal* **51**:
1126-1136.

Oldroyd GE. 2013. Speak, friend, and enter: signalling systems that promote beneficial
symbiotic associations in plants. *Nature Reviews Microbiology* **11**: 252-263.

Rodriguez-Haas B, Finney L, Vogt S, González-Melendi P, Imperial J, González-
Guerrero, M. 2013. Iron distribution through the developmental stages of *Medicago*
truncatula nodules. *Metallomics* **5**: 1247-1253.

Roux B, Rodde N, Jardinaud MF, Timmers T, Sauviac L, Cottret L, Carrere S,
Sallet E, Courcelle E, Moreau S, et al. 2014. An integrated analysis of plant and
bacterial gene expression in symbiotic root nodules using laser-capture microdissection
coupled to RNA sequencing. *Plant Journal* **77**: 817-837.

Rubio LM, Ludden PW. 2008. Biosynthesis of the iron-molybdenum cofactor of
nitrogenase. *Annual Review of Microbiology* **62**: 93-111.

Scheible WR, Gonzalez-Fontes A, Morcuende R, Lauerer M, Geiger M, Glaab J,
Gojon A, Schulze ED, Stitt M. 1997. Tobacco mutants with a decreased number of
functional nia genes compensate by modifying the diurnal regulation of transcription,
post-translational modification and turnover of nitrate reductase. *Planta* **203**: 304-319.

Schöllhorn R, Burris RH. 1966. Study of intermediates in nitrogen fixation.
Federation Proceedings **25**: 710.

Sherman F, Fink GR, Hicks JB. 1986. *Methods in yeast genetics.* Plainview, NY:
Cold Spring Harbor Lab Press.

Smil V. 1999. Nitrogen in crop production: An account of global flows. *Global*
Biogeochemical cycles **13**: 647-662.

Smith FW, Ealing PM, Hawkesford MJ, Clarkson DT. 1995. Plant members of a
family of sulfate transporters reveal functional subtypes. *Proceedings of the National*
Academy of Sciences, USA **92**: 9373-9377.

Sprent JI. 2007. Evolving ideas of legume evolution and diversity: a taxonomic
perspective on the occurrence of nodulation. *New Phytologist* **174**: 11-25.

Stout PR, Meagher WR, Pearson GA, Johnson CM. 1951. Molybdenum nutrition of
crop plants: I. The influence of phosphate and sulfate on the absortion of molybdenum
from soils and solution cultures. *Plant and Soil* **3**: 51-87.

Tadege M, Wen J, He J, Tu H, Kwak Y, Eschstruth A, Cayrel A, Endre G, Zhao
PX, Chabaud M, et al. 2008. Large-scale insertional mutagenesis using the Tnt1
retrotransposon in the model legume *Medicago truncatula*. *Plant Journal* **54**: 335-347.

Tejada-Jimenez M, Galvan A, Fernandez E. 2011. Algae and humans share a
molybdate transporter. *Proceedings of the National Academy of Sciences, USA* **108**:
6420-6425.

Tejada-Jimenez M, Llamas A, Sanz-Luque, E, Galvan A, Fernandez E 2007. A
high-affinity molybdate transporter in eukaryotes. *Proceedings of the National Academy*
of Sciences, USA **104,** 20126-20130.

Tejada-Jimenez M, Chamizo-Ampudia A, Galvan, A, Fernandez E, Llamas A.
2013. Molybdenum metabolism in plants. *Metallomics* **5,** 1191-1203.

Tejada-Jimenez M, Castro-Rodriguez R, Kryvoruchko I, Lucas MM, Udvardi M,
Imperial J, González-Guerrero M. 2015. *Medicago truncatula* natural resistance-
associated macrophage Protein1 is required for iron uptake by rhizobia-infected nodule
cells. *Plant Physiology* **168**: 258-272.

Tomatsu H, Takano J, Takahashi H, Watanabe-Takahashi A, Shibagaki N,
Fujiwara T. 2007. An *Arabidopsis thaliana* high-affinity molybdate transporter
required for efficient uptake of molybdate from soil. *Proceedings of the National*
Academy of Sciences, USA **104**: 18807-18812.

Udvardi M, Poole PS. 2013. Transport and metabolism in legume-rhizobia symbioses.
Annual Review of Plant Biology **64**: 781-805.

Vasse J, Billy F, Camut S, Truchet G. 1990. Correlation between ultrastructural
differentiation of bacteroids and nitrogen fixation in alfalfa nodules. *Journal of*
Bacteriology **172**: 4295-4306.

Vernoud V, Journet EP, Barker DG. 1999. MtENOD20, a Nod factor-inducible
molecular marker for root cortical cell activation. *Molecular Plant-Microbe Interactions*
12: 604-614.

Voinnet O, Rivas S, Mestre P, Baulcombe D. 2003. An enhanced transient expression
system in plants based on suppression of gene silencing by the p19 protein of tomato
bushy stunt virus. *Plant Journal* **33**: 949-956.

 Figure Legends

Figure 1. MtMOT1.3 is a member of the MOT1 protein and is expressed in the nodule.

(a) Unrooted tree of the plant SULTR and MOT1 families. (b) Alignment of the

conserved motifs of the MOT1 proteins present in all *M. truncatula* MOT1 members.

Sequences were aligned using ClustalW method and it is extracted from a larger

alignment including the full protein sequence of the *M. truncatula* MOT1 members. (c)

Determination of *MtMOT1.3* expression in nodulated and nitrogen-fertilized *M.*

truncatula plants relative to the internal standard gene Ubiquitin carboxyl-terminal

hydrolase. Data are the mean ± SD of two independent experiments with 4 pooled

plants. n.d., non-detected. (d) *MtMOT1.3* expression in nodulated and nitrogen-fertilized

M. truncatula plants. *Ubiquitin carboxyl-terminal hydrolase1* (*MtUb1*) was used as

control for PCR amplifications.

Figure 2. MtMOT1.3 transports molybdate towards the cytosol. (a) *S. cerevisiae* strain

31019b was transformed with PDR196 empty vector or PDR196 containing *MtMOT1.3*

coding sequence, and grown at 28 ºC for 3d in serial dilution (10X) on SD solid

medium medium without an added molybdenum source or containing 50 µM sodium

molybdate. (b) *S. cerevisiae* strains used in (a) were grown in SD liquid medium, with

and without 50µM sodium molybdate, at 28 ºC for 42 h. Yeast growth was monitored

by measuring optical density at 600 nm every 3h. (c) Molybdenum content in the yeast

grown in (b) after 42 h. Data are the mean ± SD of two independent experiments.

Figure 3. *MtMOT1.3* gene is expressed in the interzone and fixation zone of *M.*

truncatula nodules. (a) and (b) GUS staining of *M. truncatula* 28-dpi nodules

transiently expressing *gus* gene under the control of *MtMOT1.3* promoter. (a) Intact

nodule. (b) Longitudinal nodule section. Bars = 200 µm. (c) Expression of *MtMOT1.3*

in *M. truncatula* nodules by laser-capture microdissection coupled to RNA sequencing.

Data were obtained from the Symbimics database

(https://iant.toulouse.inra.fr/symbimics/). Meris, meristem; Infec, infection zone; Differ,

differentiation zone; Inter, interzone; Fix, fixation zone.

Figure 4. MtMOT1.3 is localized in the plasma membrane. (a) to (c) Cross section of a

28-dpi *M. truncatula* nodule transiently expressing MtMOT1.3-HA construct and

infected with a *S. meliloti* strain constitutively expressing GFP (green). MtMOT1.3-HA

was detected using an Alexa-594-conjugated antibody (red). (a) MtMOT1.3-HA signal.

(b) Overlay of MtMOT1.3-HA and *S. meliloti* signals. (c) Overlay of MtMOT1.3-HA

signal and differential interference contrast. a, meristem; b, infection/differentiation; c,

interzone; b, fixation. Bars = 200 μm. (d) to (f) Closer view in the fixation zone of M.

truncatula nodules. DNA was stained using DAPI (blue). (d) MtMOT1.3-HA signal. (e)

S. meliloti signal. (f) Overlay of MtMOT1.3-HA, *S. meliloti* and DNA signals. Bars =

50 μm. (g) to (i) Transient co-expression of MtMOT1.3-GFP and plasma membrane

marker-CFP in *N. benthamiana* leaves. (g) MtMOT1.3-GFP signal. (h) Plasma

membrane marker-CFP signal. (i) Overlay of MtMOT1.3-GFP, plasma membrane

marker-CFP signals and differential interference contrast. Bars = 25 μm.

Figure 5. *MtMOT1.3* mutation results in a reduced nitrogen fixation rate. (a) Position of

the *Tnt1* insertion within the *MtMOT1.3* genomic region. (b) RT-PCR amplification of

MtMOT1.3 transcript in 28 dpi nodules of *M. truncatula* wild type (WT) and mutant

(*mot1.3-1*) plants. *Ubiquitin carboxyl-terminal hydrolase1* (*MtUb1*) was used as control

for PCR amplifications. (c) Growth of representative plants of wild type, *mot1.3-1*

mutant and *mot1.3-1* mutant transformed with the *MtMOT13-HA* construct. Bar = 3 cm.

(d) Dry weight of shoots and roots. Data are the mean ± SD of at least 6 independently

transformed plants. Asterisk indicates significant differences: $**p < 0.01$, $***p < 0.005$.

(e) Nitrogenase activity in 28-dpi nodules. Acetylene reduction was measured in

duplicate from two sets of four pooled plants. Data are the mean ± SD. (f) Number of

nodules per plant. Data are the mean ± SD of at least 6 independently transformed

plants. (g) Representative nodules of each *M. truncatula* line. Bar = 500 μm. (h)

Molybdenum content in shoots, roots, and nodules of wild type, *mot1.3-1* mutant, and

mot1.3-1 mutant transformed with the *MtMOT13-HA* construct. Data are the mean ± SD

of 3 sets of 4 pooled plants.

Figure 6. *mot1.3-1* mutant phenotype is not present in non-symbiotic conditions. (a)

Growth of representative plants of wild type and *mot1.3-1* mutant. Bar = 3 cm. (b) Dry

weight of shoots and roots. Data are the mean ± SD of at least 8 independent plants. (c)

Nitrate reductase activity. Nitrate reduction was measured in duplicate. Data are the

mean ± SD.

Figure 7. Proposed model for MtMOT1.3 function in molybdenum homeostasis in *M.*
truncatula. Molybdenum, in form of molybdate would be released in the
infection/differentiation zone, as already described for iron, in a process that could be
assisted by a yet-to-be identified transporter that might belong to the MOT1 family
(MtMOT1.X). From this point MtMOT1.3 would introduce molybdate into the
cytoplasm of infected and non-infected cells within the nodule. The molybdate transport
across the symbiosome membrane could be mediated by an orthologue of *L. japonicus*
sulfate transporter SST1. Finally, molybdate would entry the bacteroids through the
membrane component of the bacterial high affinity molybdate transport system ModB.
IM, bacteroid inner membrane; OM, bacteroid outer membrane; SM, symbiosome
membrane; PM, nodule cell plasma membrane.

Figure 1

b

MtMOT1.1	97	PMPVQPMKSIAAAA	110	350 FGAVPTCHGAGGLAGQYKFGGRSG 373
MtMOT1.2	93	PMPVQPMKSISAEA	106	334 FGAMPSCHGAGGLAGQYKFGGRSG 357
MtMOT1.3	99	PMPVQPMKSIAASA	102	354 FGCVPTCHGAGGLAGQYKFGGRSG 377
MtMOT1.4	96	PMPVQPMKSIAAVA	109	337 FGAMPCCHGAGGLAGQYRFGGRSG 360
MtMOT1.5	90	PMPVQPMKSIAAVA	103	330 FGAMPCCHGAGGLAGQYKFGGRSG 353

Conserved MOT1 motifs   P-PVQPMKXIAA-A   FG-MP-CHGAGGLA-QY-FGGR-G

Figure 2

Figure 3

a
Meristem
Infection /
Differentiation
Interzone
Fixation

b
Meristem
Infection /
Differentiation
Interzone
Fixation

c

Relative expression (%)

Meris Infec Differ Inter Fix

Figure 4

Figure 5

a

Tnt-1

b

	WT	mot1-3.1
MtMOT1.3		
MtUb1		

c

WT

mot1.3-1

mot1.3-1:: MtMOT1.3

d

Dry weight (mg)

■ Shoots
□ Roots

WT   *mot1.3-1 mot1.3-1:: MtMOT1.3*

***

**

e

nmol ethylene / h·plant

WT   *mot1.3-1 mot1.3-1:: MtMOT1.3*

f

Nodules / plant

WT   *mot1.3-1 mot1.3-1:: MtMOT1.3*

g

WT   *mot1.3-1 mot1.3-1:: MtMOT1.3*

h

Mo conc. (µg/g)

■ WT
□ *mot1.3-1*
▨ *mot1.3-1::MtMOT1.3*

Shoot   Root   Nodule

Figure 6

Figure 7

Supplemental Methods

<u>Nodule development studies</u>

Sterile seedlings were transferred to plates with Fahreus agar medium (Sauviac et al., 2005) without added molybdate and inoculated with 100 μl of saturated culture of *S. meliloti*. These plants were grown for 28 days at 16 h light and 22 ºC and 8 h darkness and 16 ºC. Morphological analyses were carried out in 28 dpi nodules collected and fixed in 0.25 % glutaraldehyde, 4 % formaldehyde, 2.5 % sucrose in 50 mM potassium phosphate buffer (pH7.4) at 4 ºC. Samples were dehydrated in an ethanol series and embedded in LR-White resin (London Resin Company Ltd, UK). Finally, nodules were placed in gelatin capsules, filled with resin and polymerized at 60 ºC for 24 h. Serial thin sections (0.5 μm) were cut with a Reichert Ultracut microtome (Leica, Vienna, Austria) fitted with a glass knife at the National Center for Electron Microscopy (Universidad Complutense de Madrid, Spain). Sections were stained with 1 % toluidine blue. Direct observation of sections was performed under a Zeis Axiophot photomicroscope (Carl Zeiss, Oberkoche, Germany) with an attached digital camera (Leica DFC 420C, Heerburgg, Switzerland).

<u>Immunohistochemistry and confocal microscopy</u>

The localization of MtMOT1.3-HA was also determined using an anti-mouse Alexa488-conjugated antibody in sections from nodules colonized by a *S. meliloti* strain that constitutively expresses m-Cherry. Nodules and roots were collected at 28-dpi and fixed at 4 ºC overnight in 4 % (w/v) paraformaldehyde and 2.5 % (w/v) sucrose in phosphate-buffered saline (PBS). Fixed plant material was sectioned, 100 μm wide, with a Vibratome 1000 Plus. Cell wall permeabilization was carried out by incubation with 2 % (w/v) cellulase in PBS for 1 h and 0.1 % (v/v) Tween 20 for 15 min. Sections were blocked with 5 % (w/v) bovine serum albumin in PBS and then incubated with 1:50 anti-HA mouse monoclonal antibody (Sigma) in PBS at room temperature for 2 h. Primary antibody was washed three times with PBS for 10 min and subsequently incubated with 1:40 Alexa4884-conjugated anti-mouse rabbit monoclonal antibody (Sigma) in PBS at room temperature for 1 h. Secondary antibody was washed three times with PBS for 10 min, and then DNA was stained using DAPI. Images were obtained with a confocal laser-scanning microscope (Leica SP8).

<u>Western blot</u>

Total root and nodule protein extracts were obtained from *M. truncatula* wild-type and from transformants expressing *MtMOT1.3-HA* as described by Larrainzar et al. (Larrainzar et al., 2007). Briefly, 100 mg fresh weight of nodules or roots was homogeneized in a mortar with liquid nitrogen and resuspended in cold extraction buffer (25 mM MES, 450 mM mannitol, 7 mM EDTA, 7 mM $CaCl_2$, 5 mM $MgCl_2$, 20 mM ascorbic acid, 10 mM ditithreitol pH7.2). Homogenates were briefly centrifuged at 500 x g for 5 min to remove cell debris. Protein determination was performed in accordance to Bradford (Bradford, 1976). 25 μg of this crude extract was loaded in SDS-PAGE gels (Laemmli, 1970), and protein bands visualized with Coomassie Brilliant Blue. HA-labelled proteins were detected by electroblotting the gels onto nitrocellulose membranes and immunostaining with a primary anti-HA mouse antibody (1:1000, Sigma) and a horseradish peroxidase-conjugated anti HA rabbit antibody (1:1000, Agrisera).

Bradford MM. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* **7**: 248-254.

Laemmli UK. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**: 680-685.

Larrainzar E, Wienkoop S, Weckwerth W, Ladrera R, Arrese-Igor C, González EM. 2007. *Medicago truncatula* root nodule proteome analysis reveals differential plant and bacteroid responses to drought stress. *Plant Physiology* **144**: 1495-1507.

Sauvia L, Niebel A, Boisson-Dernier A, Barker DG, de Carvalho-Niebel F. 2005. Transcript enrichment of Nod factor-elicited early nodulin genes in purified hair fractions of model legume *Medicago truncatula*. *Journal of Experimental Botany* **56**: 2507-2513.

Table S1. Primers used in this study

Name	Sequence (5' - 3')	Use
5MtMOT1.1+2087QF	CCATTTGGAACCAGAAAAGTCCA	Quantitative expression of *MtMOT1.1*
3MtMOT1.1+2234QR	GAACCCTAGCAAGACCTCACA	Quantitative expression of *MtMOT1.1*
5MtMOT1.2+1689QF	GACCGAACTTTGCAGAAAGCA	Quantitative expression of *MtMOT1.2*
3MTMOT1.2+1813QR	TGGGGATTTGGCTTGTCATCA	Quantitative expression of *MtMOT1.2*
5MTMOT1.3+1077QF	TTGTCATGGTGCCGGTGGAC	Quantitative expression of *MtMOT1.3*
3MTMOT1.3+1237QR	ACACTCCTAAGATCCCGACCGACCG	Quantitative expression of *MtMOT1.3*
5MTMOT1.4QF	CACACGACAACATCTCCAACGACCAACC	Quantitative expression of *MtMOT1.4*
3MTMOT1.4QR	AAGGTTGTTGACTAGTGATAAGGC	Quantitative expression of *MtMOT1.4*
5MTMOT1.5+1260QF	CTTGTGGGGTCTAGTGCTGC	Quantitative expression of *MtMOT1.5*
3MTMOT1.5+1367QR	ACTTGCTCTTGACCTTCATGC	Quantitative expression of *MtMOT1.5*
35MtUBqF	GAACTTGTTGCATGGGTCTTGA	Quantitative expression of *MtUbiquitin carboxyl-terminal hydrolase*
3MtUBqR	CATTAAGTTTGACAAAGAGAAAGAGACAGA	Quantitative expression of *MtUbiquitin carboxyl-terminal hydrolase*
5MTMOT13F	AAAACTGCAGCATTTTATTTTCTTCCATTGGTTC	*MtMOT1.3* cDNA cloning into pDR196
3MTMOT13_534R	AAAACTCGAGAAATTGCCATATTTTTGAAATCCC	*MtMOT1.3* cDNA cloning into pDR196
5MtMOt13-1085pGW	GGGGACAAGTTTGTACAAAAAAGCAGGCTAAAT-GACATATTTGCATAAAATTGG	*MtMOT1.3* promoter cloning into pGWB3 or pGWB13 vectors

5MtMOT13p2000GW	GGGGACAAGTTTGTACAAAAAAGCAGGCTAGTG-CATAGACAACATCAAAAAGC	*MtMOT1.3* promoter cloning into pGWB3 vector
3MtMOT13+1441GW	GGGGACCACTTTGTACAAGAAAGCTGGGTAGAT-GAGGCTTTTCTGATTCCAAAT	*MtMOT1.3* genomic region cloning into pGWB13 or pGWB5 vectors
5MtMOT12+1GW-C	GGGGACAAGTTTGTACAAAAAAGCAGGCTATGG-AAAACCAAAACCCTCC	

Table S2. Occurrence of MOT1 members in the genome of legume and non-legume plants.

Legumes	Members of the MOT1 family
Arachis duranensis	3
Cajanus cajan	3
Cicer arietinum	4
Glycine max	9
Medicago truncatula	5
Phaseolus vulgaris	4
Pisum sativum	4
Vigna angularis	5
Average	4.62
Non-legume plants	
Arabidopsis thaliana	2
Hordeon vulgare	2
Populus trichocarpa	2
Ricinus communis	3
Solanum tuberosum	2
Sorghum bicolor	3
Vitis vinifera	2
Zea mays	2
Average	2.25

Figure S1. Determination of MOT1 gene family expression in nodulated and nitrogen-

fertilized *M. truncatula* plants relative to the to the internal standard gene *Ubiquitin*

carboxyl-terminal hydrolase. Data are the mean ± SD of two independent experiments

with 4 pooled plants. n.d., non-detected.

Figure S2. Determination of *MtMOT1.3* regulation by molybdate relative to the internal

standard gene *Ubiquitin carboxyl-terminal hydrolase*. Data are the mean ± SD of three

independent experiments with 4 pooled plants each.

Figure S3. Growth of *S. cerevisiae* transformed with pDR196 or with pDR196 expressing

MtMOT1.3 under different concentrations of sodium molybdate. (a) 0 μM added sodium

molybdate. (b) 10 μM added sodium molybdate. (c) 30 μM added sodium molybdate.(d)

50 μM added sodium molybdate. (e) 100 μM added sodium molybdate. (f) 1000 μM

added sodium molybdate. Data are the mean ± SD of two independent experiments.

Figure S4. Autofluorescence control for Alexa594 emission. Immunolocalization of

MtMOT1.3-HA in nodule sections from 28 dpi *M. truncatula* plants transformed with

pGWB13::MtMOT1.3-HA. Left nodule was incubated with a mouse anti-HA antibody,

while just PBS was used for the right one. The image is the result of a z-project of 22

optical sections combining a tilescan of 14 areas.

Figure S5. MtMOT1.3 is localized in the plasma membrane. Cross section of a 28 dpi *M.*

truncatula nodule expressing MtMOT1.3-HA and infected with a S. *meliloti* strain

constitutively expressing mCherry (red). DNA is stained with DAPI (blue). (a) Nodule

section incubated with an anti-HA mouse antibody and a anti-mouse Alexa488-

conjugated antibody (green). Left panel differential interface contrast image; top middle

panel, S. meliloti distribution; lower middle panel, DNA distribution; top right panel

MtMOT1.3-HA distribution; lower right panel overlay of DAPI, mCherry, and Alexa488

signals. Bars = 10 μm. (b) Nodule section prepared as in (a) but not adding the anti-HA

mouse antibody. Top left panel, S. meliloti distribution; lower left panel, DNA

distribution; top right panel distribution of Alexa488 signal; lower right panel differential

interface contrast image. Bars = 20 μm.

Figure S6. Immunolocalization of MtMOT1.3-HA in crude protein extracts from 28-dpi

M. truncatula roots and nodules obtained from control and pGWB13::MtMOT1.3-HA

transformed plants.

Figure S7. Transient expression of MtMOT1.3-GFP in *N. benthamiana* leaves. (a) to (c)

Plasma membrane marker-CFP expression. (a) Plasma membrane marker-CFP signal in

the green channel. (b) Plasma membrane marker-CFP in the cyan channel. (c) Overlay of

green and cyan channels and differential interference contrast. Bars = 25 μm. (d) to (f)

MtMOT1.3-GFP expression. (d) MtMOT1.3-GFP signal in the green channel. (e)

MtMOT1.3-GFP signal in the cyan channel. (f) Overlay of green and cyan channels and

differential interference contrast. Bars = 25 μm.

Figure S8. Nitrogenase activity in 28-dpi nodules. Acetylene reduction was measured in

duplicate from two sets of four pooled plants. Data are the mean ± SD.

Figure S9. Nodule development in wild type and mot1.3-1 plants. Logitudinal sections of

wild type (a) and mot1.3-1 (b) nodules stained with toluidine blue. Bars = 100 μm. (c)

Nodule growth in wild type and mot1.3-1 plants. Bars = 100 μm (d) Nodulation kinetics

in wild type and mot1-3 plants. Data are the mean ± SD from ten plants.

Figure S10. *mot1.3-1* mutant aberrant phenotype is restored upon supplementation with

molybdate. (a) Growth of representative plants of wild type and *mot1.3-1* mutant. Bar =

3 cm. (b) Dry weight of shoots and roots. Data are the mean ± SD of at least 6 independent

plants. Asterisk indicates significant differences: *$p < 0.05$, **$p < 0.01$. (c) Number of

nodules per plant. Data are the mean ± SD of at least 6 independent plants. (d) Nitrogenase

activity in 28-dpi nodules. Acetylene reduction was measured in duplicate from two sets

of four pooled plants. Data are the mean ± SD. Asterisk indicates significant differences:

*$p < 0.05$. (e) Nodule length in 28-dpi nodules. Data are the mean ± SD of 15 nodules

from from 6 plants. Asteriks indicates significant differences: *$p < 0.05$.

Figure S11. *mot1.3-1* plants show reduced nitrogenase activity also in hydroponics. (a)

Growth of representative plants of wild type and mot1.3-1 mutant. Bar = 3 cm. (b) Dry

weight of shoots and roots. Data are the mean ± SD of at least 8 independent plants.

Asterisk indicates signifi-cant differences: *$p < 0.05$, **$p < 0.01$. (c) Number of nodules

per plant. Data are the mean ± SD of at least 7 independent plants. (d) Nitrogenase activity

in 28-dpi nodules. Acetylene reduction was measured in duplicate from two sets of four

pooled plants. Data are the mean ± SD. Asterisk indicates significant differences: *$p <$

0.05

Figure S12. Expression of *MtMOT1.1, MtMOT1.2, MtMOT1.4,* and *MtMOT1.5* in

mot1.3-1 nodules relative to the internal standard gene *Ubiquitin carboxyl-terminal*

hydrolase and relativised to the expresion value in wild type plants (100 %). Data are the

mean ± SD of three independent experiments with 4 pooled plants each.

Figure S1

Figure S2

Figure S3

Figure S4

Figure S5

a

b

Figure S6

Figure S7

Figure S8

Figure S9

Figure S10

Figure S11

Figure S12